Title page

Codon Usage Bias Analysis of the *WRKY* Gene Family in *Musa acuminata*

Jiaman Sun[1, 2†], Ji Zhang[2†], Jinzhong Zhang[1], Brett J. Ferguson[3], Andrew Chen[4*]

1. Guangdong Provincial Key Laboratory of Conservation and Precision Utilization of Characteristic

Agricultural Resources in Mountainous Areas, School of Life Science, Jiaying University, Meizhou,

Guangdong, 514015, China.

2. Guangxi Academy of Agricultural Sciences, Nanning 530007, China.

3. Integrative Legume Research Group, School of Agriculture and Food Sustainability, The University

of Queensland, St. Lucia, QLD 4072, Australia

4. School of Agriculture and Food Sustainability, The University of Queensland, St Lucia, Brisbane,

QLD 4072, Australia.

* Corresponding author, a.chen2@uq.edu.au

†These authors contributed equally to the work.

Abstract

Codon usage bias (CUB) is a universal phenomenon across all organisms and species, providing valuable insights into codon usage patterns to understand evolutionary mechanisms. The *WRKY* transcription factor family plays a pivotal role in regulating plant responses to various stresses at both physiological and biochemical levels. Bananas (*Musa* spp.), widely consumed as a fruit or staple food, are an important horticulture crop in the tropical and subtropical regions around the world. A total of 151 *WRKY* transcription factors were identified across the genome of the *Musa acuminata* banana cultivar 'Guijiao 9', which has been shown to be resistant to the Tropical Race 4 of the Fusarium wilt of banana. The codons of these transcription factors, selected based on their expression from RNA-Seq data, were analysed to investigate the evolutionary patterns of WRKY genes in *Musa acuminata*. The average GC content of *MaWRKY* genes was 55.49%, with a GC3 content of 62.13%, indicating a preference for G/C-ending codons. Among the codons, 27 were identified as high frequency, with 22 ending in G or C. The high effective number of codons (ENC) values (34.53–60.79) and low codon adaptation index (CAI) values (0.143–0.3) suggested weak CUB and that was associated with low gene expression levels. ENC-plot, PR2-plot, and neutrality analysis revealed that both natural selection and mutation pressure contributed to the observed CUB, with natural selection being the dominant factor influencing the codon usage of *MaWRKY* genes in *M. acuminata* 'Guijiao 9'. Fourteen optimal codons, all ending in G or C, were identified. This analysis provides a theoretical foundation for further understanding the evolutionary mechanisms of *WRKY* genes and identifying key genes involved in resistance to biotic stress in *Musa*. Analysing the evolution of *WRKY* family genes can offer valuable genetic insights into banana growth and development, supporting the phenotypic selection of banana hybrids in breeding programs.

Keywords: *Musa acuminata*, 'Guijiao 9', *WRKY* transcription factors, codon usage bias, natural selection, mutation pressure, Fusarium wilt of banana, *Fusarium oxysporum* f. sp. *cubense*, Tropical Race 4.

Introduction

A codon is a triad nucleotide sequence that encodes amino acids on messenger RNA. Codons that encode the same amino acid are referred to as synonymous codons. Over the course of evolution, certain species and specific genes tend to favour particular codons for encoding amino acids, leading to synonymous codon usage bias (Kumar et al., 2016). This bias influences gene expression levels, with stronger codon usage bias generally correlating with higher gene expression (Jiang et al., 2008). Low-frequency codons help maintain codon diversity and contribute to gene expression regulation, playing a critical role in species stability and evolution of each species (Liu et al., 2010). Three primary forces shaping codon usage bias have been identified: mutation pressure, natural selection and random genetic drift (Behura and Severson, 2013). Mutation pressure causes deviations in nucleotide composition, while natural selection restricts codon usage bias to optimise protein production efficiency in highly expressed genes (Guan et al., 2018). Additional factors influencing codon usage bias include nucleic acid composition (Mondal et al., 2016), gene expression level (Hambuch and Parsch, 2005), gene length (Wei et al., 2014), and tRNA abundance (Deng et al., 2020).Analysis of codon usage bias is crucial for understanding the evolutionary mechanisms and codon usage patterns of specific plants. Synonymous codon usage bias differs between monocotyledons and dicotyledons. Monocotyledons tend to favour codons ending in G/C, while dicotyledons prefer A/T endings (Murray et al., 1989). GC content also serves as a distinguishing factor between these two groups. Related species often exhibit similar codon usage biases, making it an important tool for inferring evolutionary relationships among different plant species (Ma et al., 2015).

Codon usage bias reflects the origins, evolution, and mutations of species or genes, providing essential insights for gene function analysis, protein expression and protein structure research (Wu et al., 2015; He et al., 2016). In recent years, studies on codon preference in gene families and individual genes across plants (Liu et al., 2020), animals (Uddin et al., 2020), viruses (Deb et al., 2020; Xin et al., 2020) and other organisms has been increasingly widespread. *WRKY* transcription factor family is one of the largest transcription factor families in plants, playing a key role in physiological and biochemical processes such as hormone synthesis, signal transduction, plant growth and development, metabolism regulation and biological responses to both biotic and abiotic stresses (Govardhana and Satyan, 2020; Li et al., 2020). The widespread use of next-generation sequencing technologies for whole genome and transcriptome sequencing, along with the availability of bioinformatics tools have greatly advanced in-depth studies of WRKY transcription factor family genes and their functional roles in plant's ability to adapt to environmental challenges (Wani et al., 2021; Goyal et al., 2023; Liu et al., 2023; Deng et al., 2024).Banana (*Musa.* spp) is an important fruit crop, which supplies essential nutrients to millions of people around the world (Drenth and Kema, 2021). Cultivated bananas originated from natural intra- and interspecific hybridisation between two diploid species, *Musa acuminata (*AA genome) and *Musa balbisiana* (BB genome) (Drenth and Kema, 2021). However, due to various biotic and abiotic stresses, global banana yields have significantly declined in recent years (Siamak and Zheng, 2018; Patel et al., 2019).

One of the major diseases threatening the banana production dominated by 'Cavendish' from the AAA subgroup is the Fusarium wilt of banana, caused by the *Fusarium oxysporum* f. sp. *cubense* tropical race 4 (*Foc* TR4) (Zorrilla-Fontanesi et al., 2020; Roberts et al., 2024). Current efforts have been focused on containment and deterrence through biosecurity measures; biological control agents; integrated disease management strategies; and the development of resistant banana varieties (Swarupa et al., 2014; Ploetz,

2015; Dita et al., 2018; Pegg et al., 2019). Resistant (R) genes, such as Resistance Gene Analog 2 (RGA2),
which was isolated from *Foc TR4*-resistant *Musa acuminata* ssp. *malaccensis*, demonstrated strong
resistance in the field when over-expressed in 'Cavendish' transgenic lines (Dale et al., 2017). Forward
genetic studies have also identified QTLs conferring resistance to *Foc* TR4 and subtropical race 4 (STR4)
(Ahmad et al., 2020; Chen et al., 2023a, 2023b). Candidate R genes, specifically pattern recognition
receptors, were identified.

Several *WRKY* genes, such as *MaWRKY28, MaWRKY71, MaWRKY40 and MaWRKY22,* have been
associated with banana's response to biotic stress, including *Foc* TR4 (Sun et al., 2019; Li et al., 2012).
However, research on the codon usage pattern of WRKY genes in banana has not been reported. In our
previous study, *Musa. acuminata* banana 'Guijiao 9' exhibited strong resistance to *Foc*-TR4, with lower
disease incidence and severity compared to susceptible banana cultivars (Sun et al., 2019). In this study, 151
members of WRKY gene family were identified from *M*. *acuminata* 'Guijiao 9' transcriptome data. Factors
influencing codon preference of *MaWRKY* and potential evolutionary models were determined by analysing
the CUB of *MaWRKY* genes in 'Guijiao 9'. These findings contribute to understanding the function of the
WRKY gene family and provide insights for codon optimisation of *MaWRKY* in the process of biotic stress.

Results

Codon composition of *MaWRKY* genes reveals a G/C bias

The codon composition and ENC values of 151 banana *MaWRKY* family members were statistically
analysed, with detailed codon bias parameters provided in Additional File 2. The ENC values ranged from
34.53 to 60.79, indicating varying levels of codon bias, although the overall CUB was weak. CAI values for
MaWRKY genes ranged from 0.143 to 0.3, suggesting low gene expression levels.

The average content of A3s and T3s was not significantly different from G3s and C3s ($P > 0.05$) (Fig. 1).
However, the combined average content of G3s+C3s diferred significantly from that of A3s+T3s. The
average GC content was 55.49%, ranging from 51.05% to 58.95%, indicating a relatively high GC content
and a preference for G/C-ending codons in *MaWRKY* genes.

Analysis of G/C content at different codon positions showed that GC3s (62.13% on average) were much
higher than GC1s (54.65%) and GC2s (48.24%), suggesting preference for G or C at the third codon
position. GC3s exhibited greater variability (ranging from 54.20% to 69.40%) compared to GC1s (ranging
from 52.24% to 57.93%) and GC2s (ranging from 45.65% to 50.75%). This variability indicates that the first
and second codon positions are relatively stable, while the third position shows greater fluctuation,
potentially contributing to the observed codon usage bias.

Codon Usage Parameter Correlation Analysis

The Pearson Correlation coefficients for various codon usage indices were calculated and are presented in
Table 1 to assess the relationships between factors influencing codon usage. ENC value showed a significant
positive correlation with A3s ($r = 0.726$, $p < 0.05$) and T3s ($r = 0.690$, $p < 0.05$) but was negatively
correlated with G3s ($r = -0.616$, $p < 0.05$), C3s ($r = -0.703$, $p < 0.05$), GC ($r = -0.704$, $p < 0.05$) and GC3s (r

= -0.750, p < 0.05). These findings indicate that the nucleotide composition at the third position of
synonymous codons strongly influences codon usage bias.

Additionally, G3s was positively correlated with C3s (r = 0.615, p < 0.05), GC3s (r = 0.844, p < 0.01) and
GC (r = 0.695, p < 0.01). Conversely, A3s exhibited a negative correlation with G3s (r = -0.801, p < 0.01),
C3s (r = -0.842, p < 0.01), GC3s (r = -0.938, p < 0.01) and GC(r = -0.938, p < 0.01). Similarly, T3s showed
a negative correlation with G3s (r = -0.776, p < 0.05), C3s (r = -0.913, p < 0.01), GC3s (r = -0.959, p < 0.01),
and GC (r = -0.881, p < 0.01). These negative correlations suggest that the frequency of A or T at the third
codon position has little influence on the codon usage bias of *MaWRKY* genes.

Principal Component Analysis

Principal Component Analysis (PCA) was performed on the RSCU values of *MaWRKY* genes sequences.
The contribution of the first 20 factors to the variance in codon usage bias is shown in Fig. 2A. The first four
factors accounted for 58.47% of the total variance, capturing the majority of the differences in codon usage
patterns.
PCA analysis of the 59 codons in *MaWRKY* family members, categorised by their endings (A, T, C or G),
revealed distinct distribution patterns (Fig. 2B). Codons ending in A and T were clustered on the left side of
axis 1, and those ending in G and C were primarily located on the right side. This suggests similar RSCU
usage patterns for codons ending in A or T and for those ending in G or C. The distribution of codons ending
in A and T was denser and showed partial overlap, whereas codons ending in G and C were more dispersed,
with those ending in C being spread across all four quadrants. These findings indicate a natural
evolutionary or long-time domestication bias in banana codon usage toward G and C endings.

Factors affecting codon usage bias in *MaWRKY* genes of *Musa acuminata*

ENC-plot analysis commonly used to assess the effect of mutation pressure on codon usage bias. The ENC
values are plotted against GC3 values (Fig. 3A). The standard curve shows that the relationship between
ENC and GC3s is shaped primarily by mutation pressure rather than selection. The ENC values for
MaWRKY family members generally align with the standard curve in the ENC-plot, indicating that mutation
pressure is a major factor influencing codon usage bias for these genes (Fig. 3A). However, the ENC values
of many *MaWRKY* genes deviate significantly from the standard curve (Fig. 3A), suggesting that mutation
pressure is not the sole factor driving codon bias. Other factors, such as natural selection and gene
expression, may also contribute to the observed codon bias.
154
PR2 analysis was performed to assess the impact of mutation and selection pressure on codon usage by
examining whether there was a mutation imbalance between A/T (U) and C/G. In the PR2 plot, A3 / (A3 +
T3) is plotted as the ordinate and G3 / (G3 + C3) as the abscissa for *MaWRKY* family members to explore
the influence of evolutionary factors (Fig. 3B). The A3 / (A3 + T3) or G3 / (G3 + C3) values for most
MaWRKY genes deviated from 0.5, suggesting that codon bias was additional influenced by factors other
than base mutation of encoding gene amino acids (Fig. 3B). This indicates that additional pressures, such as
natural selection, likely played a role in the evolution of *MaWRKY* genes.
162
To further identify the factors influencing codon usage bias, a neutrality plot analysis was performed by
comparing GC3 (abscissa) and GC12 (ordinate) to examine the role of mutation-selection equilibrium in

codon usage variation. A linear regression line was plotted with GC3 (abscissa) and GC12 (ordinate) values
(Fig. 3C). A regression coefficient (RC) value of less than 0.5 suggests a greater influence of natural
selection, while an RC greater than 0.5 indicates a greater impact of mutation pressure. The RC value of
0.1718 indicates that the natural selection had a more significant effect on the codon usage bias of *MaWRKY*
gene family in banana than mutation pressure (Fig. 3C).
170

Relative synonymous codon usage (RSCU) in *MaWRKY* gene family

To investigate the patterns of synonymous codon usage bias in *MaWRKY* gene family, the RSCU of 59
synonymous codons (excluding the initiation codon, termination codon and tryptophan) was analysed. The
results revealed 27 high-frequency codons (RSCU > 1) in *MaWRKY* family members, including AGG, AGA,
CUC, CUG, UUG, AGC, UCC, UGG, GUG, GUC, ACC, ACG, GGC, GCC, GCU, CCG, CCA, AUC, UGC,
AAG, UUC, AAC, GAG, UAC, CAG, CAC, and GAU, respectively (Fig. 4). Among these, 22 codons
ended in G/C, indicating a preference for G or C at the end of the codons in *MaWRKY* members. The codon
AGG of Arginine (RSCU = 1.800) exhibited the strongest preference, accounting for 30% of the
synonymous codons, while the codon AUC for Isoleucine accounted for 59.0%. These high-frequency
codons show preferential usage, which contributed to the deviation of the ENC value of *MaWRKY* genes
from an ENC value of 61.

The optimal codons of *MaWRKY* Genes

The optimal codons in *MaWRKY* genes of *M. acuminata* 'Guijiao 9' were identified by combining
high-frequency codons (RSCU > 1) and highly expressed codon (ΔRSCU ≥ 0.08). The analysis revealed 26
highly expressed codons with ΔRSCU≥ 0.08 (Table 2) and 27 high-frequency codons with RSCU > 1 (Fig.
4). In total, 14 optimal codons were determined for *MaWRKY* genes in *M. acuminata* 'Guijiao 9', including
CUC, CUG, AUC, GUG, AGC, CCG, ACC, UAC, CAC, CAG, AAC, GAG, UUC and GGC. Notably, all of
these optimal codons end with G or C (Table 2).

Codon usage patterns of WRKY Family Gene members across different plant species

Using the Euclidean distance, the RSCU values of 25 *WRKY* gene family members across different plant
species were clustered with SPSS software v22.0 (Fig. 5). The clustering results of *WRKY* genes in these
plant species were consistent with their phylogenetic tree, as shown in Phytozome v12.1
(https://phytozome.jgi.doe.gov/). This finding suggests a correlation between species classification and
codon usage patterns. It also indicates that the codon usage patterns of *WRKY* genes in different species align
with the evolutionary processes of their respective genomes.
196
A negative correlation was observed between GC3s content and ENC value when comparing the GC3s and
ENC values of WRKY gene family members from different plant species (Fig. 5). The variation in GC3s
content and ENC values differed among different plants categories. For example, in monocotyletons, the
GC3s content ranged from 60.03% to 75.88%, and the ENC value ranged from 42.67 to 52.20, while in
dicotyledons, the GC3s content varied between 37.77% and 52.98%, and the ENC value ranged from 51.85
to 54.26. Plant species that are closely related species showed similar GC3s content and ENC value. 'Guijiao
9' and another *Musa acuminata* representative were clustered adjacently due to their similar ENC and GC3

values, as both belong to the monocotyledon group. The ENC value of 'Guijiao 9' and *Musa acuminata* is
most similar to that of tree species like *Populus trichocarpa*, *Manihot esculenta*, and *Citrus sinensis*, but
significantly different from gramineous species such as *Oryza sativa*, *Zea mays* PH207, and *Setaria viridis.*
207

A heat map analysis of the 59 RSCU values (excluding the initiation codon, termination codon and
tryptophan) for 25 plant species revealed that the codons were divided into two groups: those ending with
G/C and those ending with A/T. Among the 25 plant species, Algae, Pteridophytes and monocotyledons
favoured codons ending in G/C, while dicotyledons preferred those ending in A/T (Fig. 6).

Discussion

During the process of gene expression, the selective use of codons varies from species to species. Different
organisms exhibit different preference for synonymous codons encoding the same amino acid, closely linked
to their genetic characteristics. CUB can influence mRNA stability, transcription, protein translation
accuracy, and protein folding, thereby fine-tuning gene expression.
217

In this study, 151 *MaWRKY* genes were identified in *M. acuminata* using bioinformatics tools for codon
analysis. GC composition has been shown to influence codon and amino acid usage, with the third base of a
codon could directly reflecting codon usage patterns (Chen et al., 2013; Gao et al., 2022). The average GC
content and GC3 content of *MaWRKY* codons were 55.49% and 62.13% respectively, indicating a
preference for G/C-ended codons over A/U-ended codons in *M. acuminata*. This preference was furher
supported by RSCU analysis, which confirmed *MaWRKY* genes favour G/C-ending codons. These results
were consistent with the codon usage patterns observed in other monocotyledons, such as in *Musa basjoo*,
Zea mays and *oryza sativa* (Ma et al., 2015).
226

The ENC value and CAI value provide insights into CUB and gene expression levels. Higher ENC values
suggest lower CUB and gene expression, while higher CAI values indicate better codon adaptiveness and
potentially higher gene expression levels. In this study, the *MaWRKY* genes exhibited high ENC values
(34.53-60.79) and low CAI values (0.143-0.3), indicating weak CUB and low gene expression levels of
these genes in 'Guijiao 9'. This was consistent with studies showing that most *WRKY* family genes display
stress-induced expression patterns (Dong et al., 2003; Kayum et al., 2015). Furthermore, the GC3 content
showed greater variability than GC1 and GC2 with codons ending in G/C exhibiting a broad distribution.
These findings suggest that nucleotide composition plays a significant role in influencing CUB in *MaWRKY*
genes.Mutation pressure and natural selection are recognised as important factors influencing codon usage
bias (Sharp et al., 2010; He et al., 2016). In this study, ENC-Plot analysis showed that most ENC values for
MaWRKY codons did not align with the expected curve, with a few codons exhibiting strong bias near an
ENC value of 35. This finding indicates that mutation pressure is not the primary factor driving codon
preference. Instead, other factors, such as natural selection and gene expression, likely play significant roles.
240

PR2 plot analysis further supported this conclusion, as the A3 / (A3 + T3) and G3 / (G3 + C3) values for
most *MaWRKY* members deviated from 0.5. This deviation suggests that codon bias in *MaWRKY* genes is
influenced more by pressures like natural selection than by mutation. Neutrality plot analysis provided
additional evidence, with a regression coefficient of 0.17, highlighting that natural selection exerts a stronger
influence on codon bias than mutation pressure.

Base on ENC, PR2 and neutrality plot analyses, the CUB of *MaWRKY* genes in *M. acuminata* is influenced

by multiple factors, with natural selection being the dominant force. These findings align with the study by

He et al. (2016), which reported that Ginkgo biloba unique genes involved in environment adaptation

preferentially using G/C-ending codons, with natural selection as the primary driver of codon usage bias.

Interestingly, while monocotyledons like *M. acuminata* favour G/C-ending codons, dicotyledons such as

Helianthus annuus (Gao et al., 2022) and *Brassica napus* (Li et al., 2013) exhibit a preference for A/T

(U)-ending codons. In these species, mutation pressure appears to play a larger role in shaping codon usage

bias (Gao et al., 2022; Li et al., 2013). Thus, the codon usage preferences of *WRKY* gene families vary

significantly among different species, reflecting distinct evolutionary pressures and adaptations. An optimal

codon is defined as one that is over-represented in highly expressed genes and is associated with higher gene

expression levels and efficient translation (Zhou et al., 2009). It was generally believed that GC-rich

organisms favour codons ending with G or C, while AT-rich organisms prefer codons ending with A or T

(Hershberg et al., 2009; Palidwor et al., 2015). In our study, all 14 identified optimal codons of *MaWRKY*

genes ended with G or C (Table 2), aligning with the GC-rich composition of *MaWRKY* genes in *M.*

acuminata. Similar patterns have been observed in other monocotyledon, such as *Oryza sativa* (Liu et al.,

2004) and *Zea mays* (Liu et al., 2010), whose genomes are GC-rich, and whose optimal codons also end

with G or C. This finding has practical implications for introducing point mutations and codon modifications

to enhance the production of specific proteins in host cells. Codon optimisation based on these preferences

can improve gene expression efficiency in biotechnological and agriculture applications.

RSCU values serve as a significant indictor of evolutionary relationships among different organisms and are

widely used in studies of biological evolution in plants. Cluster analysis of RSCU values for *WRKY* genes

across various plant species revealed that the codon usage patterns of *WRKY* gene families align with the

codon usage preferences of their respective genomes during evolution (Fig. 5). *MaWRKY* genes clustered

with other monocotyledon species in a single branch, exhibiting a preference for codons ending with G/C

(Fig. 6). This suggests that the codon usage patterns of *WRKY* family genes are consistent among related

species, reflecting evolutionary.

Conclusions

In the present study, 151 coding sequences of *WRKY* genes from *M. acuminata* 'Guijiao 9' were analysed to

investigate codon usage bias and its potential influencing factors. The results revealed that *MaWRKY* genes

in *M. acuminata* 'Guijiao 9' exhibit a preference for codons ending in G/C. Overall, the codon usage bias

was weak, and the expression levels of these genes were low, indicating high variability in synonymous

codon usage among *MaWRKY* genes. While mutation pressure played a role, natural selection emerged as

the primary factor in shaping the codon usage patterns of *MaWRKY* genes during evolution. The codon

usage patterns of *MaWRKY* genes provide valuable insights into their evolutionary dynamics and offer a

foundation for identifying and developing suitable gene targets with resistance to biotic stress.

Materials and Methods

Screening, Identification and Characterization of Sequence

The complete genome of *M. acuminata* double-haploid 'Pahang' was downloaded in FASTA format from the National Centre for Biotechnology Information (NCBI) banana genome database (https://www.ncbi.nlm.nih.gov/datasets/taxonomy/214687/, GenBank assembly accession: GCA_904845865.1), and was used as the reference genome. The Coding Domain Sequences (CDS) of the *WRKY* gene family were retrieved from the transcriptome datat of *Musa acuminata* banana 'Guijiao 9' (Additional File 1). Corresponding amino acid and nucleotide sequences were obtained through local BLAST. HMMER 3.3 (http://hmmer.org/download.html) was used to identify sequences containing complete WRKY domain using the reference sequence (PF03106) obtained from PFAM (http://pfam.sanger.ac.uk/). The presence of the WRKY domain in these sequences was further confirmed using NCBI's CDD tool (https://www.ncbi.nlm.nih.gov/cdd). ORFfinder was then used to identify CDS from the retained sequences (https://www.ncbi.nlm.nih.gov/orffinder/). Ultimately, a total of 151 *M. acuminata* *WRKY* genes (*MaWRKY)* were identified and included in the subsequent analysis (Additional File 2).

The CDS sequences of the *WRKY* gene family from 24 different species, including Bryophyte, Algae, Pteriophyte, monocotyledon and dicotyledon, were downloaded from the JGI Phytozome12.1 database (https://phytozome.jgi.doe.gov/pz/portal.html) and utilised for Cluster analysis.

Indices of Codon Usage Bias

Following the screening of *MaWRKY* gene CDS sequences, the key indices of codon usage bias were calculated, including the effective number of codons (ENC), codon adaptation index (CAI), relative synonymous codon usage (RSCU), total GC content of each CDS (GC), and GC content at the first, second and third codon positions (GC1, GC2, GC3). Additionally, the nucleotide composition at synonymous third codon positions (A3s, T3s, G3s, C3s) was analysed. These calculations were performed using the CodonW program (version 1.4.2). Correlations between nucleotide contents were assessed using SPSS statistical software (version 23.0), with values exceeding 0.5 and below -0.5 indicating strong positive and strong negative correlations, respectively.

The Effective Number of Codons (ENC) is a key metric for evaluating the degree of codon usage bias. ENC values range from 20, representing extreme bias where only one codon is used for each amino acid, to 61, indicating no bias where all synonymous codons are equally utilised. Higher ENC values suggest weaker codon usage bias and lower expression levels (Sharp and Li, 1986; Wright, 1990; Zhao et al., 2016). Typically, an ENC value above 40 indicates weak CUB (Wright, 1990). The Codon Adaptation Index (CAI) ranges from 0 to 1, reflecting how well a gene's codon usage aligns with those of highly expressed genes (Sharp and Li, 1987). Higher CAI values signify better adaptiveness and higher potential expression efficiency. Relative Synonymous Codon Usage (RSCU) measures the observed frequency of a codon relative to its expected frequency, independent of the amino acid composition of gene product (Wang et al., 2016). An RSCU value of 1 signifies no bias, meaning the codon is used as frequently as expected. Values greater than 1 denote a preference for the codon (positive bias), while values below 1 indicate under utilisation (negative bias) (Zhao et al., 2016).

ENC-plot Analysis

ENC-plot analysis is commonly used to identify factors influencing codon usage bias by plotting ENC values against GC3s values. The standard curve represents the relationship between ENC and GC3s, helping to distinguish whether mutation pressure or natural selection is the dominant factor (Jiang et al., 2008; Liu et al., 2019). When codon usage bias is primarily driven by mutation pressure, the points tend to lie on or just below the standard curve. In contrast, if natural selection and other factors play a larger role, the points will fall below the standard curve (Wright, 1990).

Parity Rule 2 (PR2) Bias Plot Analysis

A PR2 bias plot is created by plotting the AT bias (A3 / (A3+ T3)) on the y-axis and the GC bias (G3 / (G3 + C3)) on the x-axis. The midpoint at 0.5 indicates an equal balance between G = C and A = T, suggesting no significant bias from either mutation or selection pressure (Deb et al., 2020). If the genes are clustered near the centre, it suggests that the base frequencies are relatively balanced, and the codon bias is mainly affected by mutation pressure. Conversely, If the genes are far from the centre, other factors may be affecting the codon usage bias.

Neutrality Plot Analysis

A neutrality plot is used to assess the extent to which codon usage bias is influenced by mutation versus selection in organisms, by comparing GC3 (x-axis) and GC12 (y-axis). In this plot, each gene is represented by a point. If the regression coefficient of the plot approaches 1, the points will show a clear pattern, indicating selection plays a significant role in codon usage bias. If the regression coefficient deviates from 1, it suggests that factors other than mutation pressure are also influencing the codon usage bias (Sueoka, 1988).

Relative Synonymous Codon Usage (RSCU) Cluster Analysis

RSCU cluster analysis was employed to group similar elements. Euclidean distance, a common method for calculating clustering distance, measures the linear correlation between two variables. A smaller clustering distance indicates higher similarity between the clustered genes. The RSCU values of the *MaWRKY* gene family members in 'Guijiao 9' were compared with those from 24 other plant species to reveal the evolutionary relationships and characteristics among them.

Principal Component Analysis (PCA) of RSCU

Principal Component Analysis (PCA) is a multivariate statistical method to examine relationships among multiple variables, and is often applied to analyse trends in synonymous codon usage patterns (Kanaya et al., 1996). In this study, the RSCU values for 59 synonymous codons from *MaWRKY* genes was reduced from 151 dimensions (representing 151 *MaWRKY* genes) to two principal components through dimensionality reduction. Fifty-nine synonymous codons did not include the initiation codon AUG, tryptophan codon UGG, and the termination codons UAA, UAG, and UGA (Zhang et al., 2018).

Optimal Codon Analysis

The optimal codon plays a significant role in improving the speed and accuracy of translation (Duret et al., 1999). To predict the optimal codon for *MaWRKY*, the ΔRSCU method was applied (Qiu et al., 2011). The *MaWRKY* gene sequences were sorted based on their ENC values. The top 10% and the bottom 10% of the

genes, based on ENC value, were selected to create two new databases (high expression and low expression, respectively), and the RSCU values for the codons were calculated. A codon was considered a high-frequency codon if its RSCU value was greater than 1, while a codon was regarded as a high-expression codon if the ΔRSCU value was equal to or greater than 0.08. A codon is defined as optimal if it met both criteria: RSCU> 1 and ΔRSCU≥ 0.08.

Statistical analysis

The CDS sequences of the *MaWRKY* were analysed using CodonW (http://codonw.sourceforge.net/) and EMBOSS online software (http://www.bioinformatics.nl/emboss-explorer/). Codon-related parameters were statistically analysed using Microsoft Office Excel 2016 and MeV software version 4.9.0 (www.tm4.org/mev.html) was used to construct the heat map. ENC-plot analysis, Neutrality plot, and Box diagram were generated using Origin 8.0 and HemI 1.0 (http://hemi.biocuckoo.org/down.php) software. Clustering analysis and principal component analysis（PCA）of RSCU values for *WRKY* genes in different plant species were performed by Statistical Product and Service Solutions (SPSS) version 22.0.

Abbreviations
CUB   Codon usage bias
ENC   Effective number of codons
CAI   Codon adaptation index
RSCU   Relative synonymous codon usage
PCA   Principal component analysis
PR2   Parity Rule 2
GC1, GC2, GC3   The GC content at the first, second and third position, respectively
A3s, T3s, G3s, C3s   The content of each nucleotide of the codon at synonymous third positions

Supplementary Information

Additional file 1. The RNA-Seq data of *Musa acuminata* banana 'Guijiao 9'

Additional file 2. The composition indices values of codon usage in MaWRKY genes

Authors'contributions
Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Writing - original draft, and Writing - review & editing: J.S.; Data Curation, Formal Analysis, Investigation, Methodology, Visualization, Writing - review & editing: J.Z.; Formal Analysis, Investigation, Supervision, and Writing - review & editing: Jinzhong Z; Supervision, and Writing - review & editing: B.J.F.; Formal Analysis, Investigation, Supervision, Writing - original draft, and Writing - review & editing: A.C. All authors have read and approved the final manuscript.

Funding
This work was supported by Guangxi Natural Science Foundation (2021GXNSFAA196014) and Guangxi Science and Technology Planning Project (Guike AA21196005). A.C. was supported by The Bill and Melinda Gates Foundation (Project Grant ID: OPP1093845) through its grant to the International Institute of Tropical Agriculture (IITA) under the project Accelerated Breeding of Better Bananas, grant number IITA

20600.15/0008-8—Phase II as well as Hort Innovation Australia through grant 'BA21000', using the banana

research and development levy and contributions from the Australian Government. Hort Innovation is the

grower-owned, not-for-profit research and development corporation for Australian horticulture.

Availability of data and materials

The genome datasets generated and analysed in the study are available in NCBI,

https://www.ncbi.nlm.nih.gov/datasets/taxonomy/214687/, and supplementary files.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Declaration of Interest Statement

The authors declare no competing interests.

Table 1. Pearson correlation coefficients among parameters influencing codon bias in *MaWRKY* genes of

Musa acuminata banana 'Guijiao 9'. A3s, T3s, G3s and C3s denote the nucleotide content at the third codon

position. GC represents the overall GC content of each coding domain sequence, while GC1, GC2, and GC3

refer to the GC content at the first, second, and third codon positions, respectively. ENC indicates Effective

Number of Codons.

Indices	A3s	T3s	G3s	C3s	GC	GC1	GC2	GC3s	ENC
A3s	1.000								
T3s	.809[**]	1.000							
G3s	-0.801[**]	-0.776[*]	1.000						
C3s	-0.842[**]	-0.913[**]	0.615[*]	1.000					
GC	-0.896[**]	-0.881[**]	0.695[*]	0.840[**]	1.000				
GC1s	-0.479	-0.373	0.246	0.356	0.705[*]	1.000			
GC2s	-0.443	-0.441	0.154	0.370	0.681[*]	0.572	1.000		
GC3s	-0.938[**]	-0.959[**]	0.844[**]	0.929[**]	0.916[**]	0.417	0.417	1.000	
ENC	0.726[*]	0.690[*]	-0.616[*]	-0.703[*]	-0.704[*]	-0.357	-0.317	-0.750[*]	1.000

418   * and ** indicate *p* values < 0.05 and   0.01, respectively.

419   **Table 2.** Identification of optimal codons in MaWRKY Genes of *M. acuminata* 'Guijiao 9'.

| AA | Codon | RSCU | | ΔRSCU | AA | Codon | RSCU | | ΔRSCU |
		High expression	Low expression				High expression	Low expression	
Leu (L)	UUA	0.37	0.40	-0.03	Tyr (Y)	UAU	0.48	0.73	-0.26
	UUG	0.48	1.63	-1.14		UAC*	1.52	1.27	0.26
	CUU*	0.73	0.69	0.04	TER	UAA*	0.38	0.00	0.38
	CUC*	2.13	1.10	1.03		UAG	0.75	2.25	-1.50
	CUA	0.37	0.53	-0.16		UGA*	2.25	0.75	1.50
	CUG*	1.92	1.66	0.26	His (H)	CAU	0.77	1.27	-0.50
Ile (I)	AUU	0.36	0.64	-0.27		CAC*	1.23	0.73	0.50
	AUC*	2.43	1.95	0.48	Gln (Q)	CAA	0.69	1.21	-0.52
	AUA	0.21	0.42	-0.21		CAG*	1.31	0.79	0.52
Met (M)	AUG	1.00	1.00	0.00	Asn (N)	AAU	0.40	0.99	-0.59
Val (V)	GUU	0.56	0.69	-0.14		AAC*	1.60	1.02	0.59
	GUC	1.00	1.24	-0.24	Lys (K)	AAA	0.49	0.50	-0.01
	GUA	0.29	0.47	-0.18		AAG	1.51	1.50	0.01
	GUG*	2.16	1.59	0.57	Asp (D)	GAU	0.79	1.23	-0.44
Ser (S)	UCU	0.67	0.89	-0.22		GAC*	1.21	0.77	0.44
	UCC	1.43	0.64	0.80	Glu (E)	GAA	0.40	0.78	-0.38
	UCA	0.37	0.95	-0.58		GAG*	1.60	1.22	0.38
	AGU*	0.27	0.40	-0.13	Cys (C)	UGU	0.41	0.61	-0.20
	AGC*	2.12	1.90	0.22		UGC	1.44	1.39	0.05
	UCG	1.14	1.22	-0.08	Phe (F)	UUU	0.31	0.68	-0.37
Pro (P)	CCU	1.07	1.36	-0.29		UUC*	1.69	1.32	0.37
	CCC	1.02	0.70	0.32	Trp (W)	UGG	1.00	1.00	0.00
	CCA	0.60	1.34	-0.74	Arg (R)	CGU*	0.71	0.12	0.59
	CCG*	1.31	0.60	0.71		CGC*	1.29	0.30	0.98
Thr (T)	ACU	0.74	0.81	-0.08		CGA*	0.49	0.41	0.08
	ACC*	1.69	0.51	1.18		CGG*	1.11	0.68	0.42
	ACA	0.43	1.53	-1.10		AGA	1.23	2.12	-0.90
	ACG	1.14	1.15	-0.01		AGG	1.19	2.36	-1.18
Ala (A)	GCU	1.05	1.12	-0.07	Gly (G)	GGU*	0.72	0.38	0.34
	GCC*	0.99	0.62	0.37		GGC*	2.05	1.63	0.42
	GCA*	1.00	0.89	0.12		GGA	0.75	1.08	-0.33
	GCG	0.96	1.38	-0.42		GGG	0.63	0.90	-0.28

420   * indicates ΔRSCU ≥ 0.08, the underlined text represents high-frequency codons (RSCU > 1), and the bold
421   text highlights the optimal codon in MaWRKY genes of *M. acuminata* 'Guijiao 9'.

Fig. 1 Nucleic acid composition of *MaWRKY* codons. The Y-axis represents the nucleic acid content, while the X-axis corresponds to parameters related to *MaWRKY* codons. These parameters included the total GC content of each coding domain sequence (CDS), the nucleotide composition at the third codon position (A3s, T3s, G3s, C3s), the GC content at each codon position-GC1, GC2, and GC3 (representing the first, second, and third positions, respectively) for each CDS.

Fig. 2 Principal component analysis (PCA) of *MaWRKY* codon usage frequency. (**A**) The relative and cumulative contributions of the first 20 factors to the total variance in PCA; (**B**) PCA plot of codon RSCU values on codons ending with A, T, G, or C. Green squares represent codons ending with A; red triangles indicate those ending with T, orange stars indicate codons ending with G, and blue circles indicate codons ending with C.

Fig. 3 (A) ENC-plot analysis of *MaWRKY* against GC3s. Points on or near the curve indicate bias caused by mutation

pressure, while points away from the curve suggest influence from natural selection or other factors. The red-dotted line

means the lowest ENC value of *MaWRKY*. **(B)** PR2 plot analysis. The midpoint at 0.5 represents an equal balance

between G = C and A = T, indicating no bias between mutation and selection pressure. Deviations from 0.5 suggest that

codon bias is primarily influenced by factors other than base mutation of gene-encoding amino acids. **(C)** Neutrality plot

analysis of *MaWRKY*. The plot compares GC3 (x-axis) and GC12 (y-axis). An RC value less than 0.5 indicates a

stronger influence of natural selection, whereas an RC greater than 0.5 suggests a greater impact of mutation pressure.

The red straight line represents the fitted curve.

Arg	AGG	1.800	AGA	1.328	CGG	0.923	CGC	0.832	CGA	0.638	CGU	0.480
Leu	CUC	1.752	CUG	1.614	UUG	1.126	CUU	0.804	CUA	0.403	UUA	0.302
Ser	AGC	1.452	UCC	1.279	UCG	1.155	UCU	0.915	UCA	0.674	AGU	0.534
Val	GUG	1.769	GUC	1.128	GUU	0.722	GUA	0.382	- 0	- 0
Thr	ACC	1.463	ACG	1.009	ACA	0.806	ACU	0.723	- 0	- 0
Gly	GGC	1.409	GGA	0.997	GGG	0.878	GGU	0.716	- 0	- 0
Ala	GCC	1.089	GCU	1.031	GCG	0.958	GCA	0.895	- 0	- 0
Pro	CCG	1.075	CCA	1.028	CCU	0.979	CCC	0.918	- 0	- 0
Ile	AUC	1.770	AUU	0.649	AUA	0.581	- 0	- 0	- 0
Cys	UGC	1.487	UGU	0.513	- 0	- 0	- 0	- 0
Lys	AAG	1.433	AAA	0.567	- 0	- 0	- 0	- 0
Phe	UUC	1.425	UUU	0.575	- 0	- 0	- 0	- 0
Asn	AAC	1.335	AAU	0.665	- 0	- 0	- 0	- 0
Glu	GAG	1.303	GAA	0.698	- 0	- 0	- 0	- 0
Tyr	UAC	1.289	UAU	0.711	- 0	- 0	- 0	- 0
Gln	CAG	1.188	CAA	0.812	- 0	- 0	- 0	- 0
His	CAC	1.133	CAU	0.867	- 0	- 0	- 0	- 0
Asp	GAU	1.004	GAC	0.996	- 0	- 0	- 0	- 0

1.8   0.9   0

Fig. 4 Relative synonymous codon usage (RSCU) analysis for each amino acid in the *WRKY* gene family of *Musa acuminata* banana 'Guijiao 9'. The red color bar represents the RSCU value, with deeper shades indicating higher RSCU values

Fig. 5 Cluster Analysis of RSCU values of 25 *WRKY* family members across different plant species, along

with the mean GC3 and ENC values for each corresponding species. Purple represents Bryophytes, orange

represents Algae, green represents Pteridophytes, red represents monocotyledons, and blue represents

dicotyledons.

Fig. 6 Heatmap analysis of the RSCU values for 25 WRKY family members across different plant species.

The color bars represent the RSCU values, with a colour gradient from green to red indicating low to high

RSCU values, respectively. The redder the color, the higher the RSCU value, while the greener the color, the

lower the RSCU value.

References

Ahmad, F., Martawi, N.M., Poerba, Y.S., Hans, S., Henk, K., Gert, H.J., 2020. Genetic mapping of Fusarium

wilt resistance in a wild banana Musa acuminate ssp. malaccensis accession. Theor. Appl. Genet.

133(12): 3409-3418.

Behura, S.K., Severson, D.W., 2013. Codon usage bias: causative factors, quantification methods and

genome-wide patterns: with emphasis on insect genomes. Biol Rev. 88: 49-61.

Chen, A., Sun, J.M., Martin, G., Gray, L.A., Hribová, E., Christelová, P., Yahiaoui, N., Rounsley,

S., Lyons, R., Batley, J., Chen, N., Hamill, S., Rai, S.K., Coin, L., Uwimana, B., D'Hont,

A., Dolezel, J., Edwards, D., Swennen, R., Aitken, E.A.B., 2023a. Identification of a Major

QTL-Controlling Resistance to the Subtropical Race 4 of *Fusarium oxysporum* f. sp. *cubense* in *Musa*

acuminata ssp. *malaccensis*. Pathogens 12, 2: 289.

Chen, A., Sun, J.M., Viljoen, A., Mostert, D., Xie, Y.C., Mangila, L., Bothma, S., Lyons, R., Hřibová, E.,

Christelová, P., 2023b. Genetic Mapping, Candidate Gene Identification and Marker Validation for

Host Plant Resistance to the Race 4 of *Fusarium oxysporum* f. sp. *cubense* using *Musa acuminata* ssp.

malaccensis. Pathogens 12, 6: 820.

Chen, L., Liu, T., Yang, D., Nong, X., Xie, Y., Fu, Y., 2013. Analysis of codon usage patterns in *Taenia*

pisiformis through annotated transcriptome data. Biochem Biophys Res Commun. 430:1344-1348.

Dale, J., James, A., Paul, J.Y., Khanna, H., Smith, M., Peraza-Echeverria, S., Garcia- Bastidas, F., Kema, G.,

Waterhouse, P., Mengersen, K., Harding, R., 2017. Transgenic Cavendish bananas with resistance to

Fusarium wilt tropical race 4. Nat. Commun. 8(1): 1496.

Deb, B., Uddin, A., Chakraborty, S., 2020. Codon usage pattern and its influencing factors in different

genomes of hepadnaviruses. Arch Virol. 165: 557-570.

Deng, Y., Hedayioglu, F., Kalfon, J., Chu, D., Haar, T., 2020. Hidden patterns of codon usage bias across
kingdoms. J R Soc Interface 17: 20190819.

Deng, P.J., Wang, Z., Li, W.Y., Chen X.H., Liu, D.Q., 2024. WRKY11 up-regulated dirigent expression to
enhance lignin/lignans accumulation in Lilium regale Wilson during response to Fusarium wilt. Journal
of Integrative Agriculture, 23(8): 2703-2722.

Dita, M., Barquero, M., Heck, D., Mizubuti, E.S.G., Staver, C.P., 2018. Fusarium Wilt of Banana: Current
Knowledge on Epidemiology and Research Needs Toward Sustainable Disease Management. Front.
Plant Sci. 9:1468.

Dong, J., Chen, C., Chen, Z., 2003. Expression profiles of the Arabidopsis WRKY gene superfamily during
plant defense response. Plant Mol Biol. 51: 21-37.

Drenth, A., Kema, G., 2021. The Vulnerability of Bananas to Globally Emerging Disease Threats.
Phytopathology 111: 2146-2161.

Duret, L., Mouchiroud, D., 1999. Expression pattern and, surprisingly, gene length shape codon usage in
Caenorhabditis, Drosophila, and Arabidopsis. PANS. 96: 4482-4487.

Gao, Y., Lu, Y., Song, Y., Jing, L., 2022. Analysis of codon usage bias of WRKY transcription factors in
Helianthus annuus. BMC Genomic Data 23:46.

Govardhana, M., Satyan, K.B., 2020. In-silico Analysis of Cucumber (*Cucumis sativus* L.) Genome for
WRKY Transcription Factors and cis-acting Elements. Comput Bio Chem. 85: 107212.

Goyal, P., Devi, R., Verma, B., Hussain, S., Arora, P., Tabassum, R., Gupta, S., 2023.
WRKY transcription factors: evolution, regulation, and functional diversity in plants. Protoplasma
260: 331–348.

Guan, D.L., Ma, L.B., Salabat, K.M., Zhan, X.X., Xu, S.Q., Xie, J.Y., 2018. Analysis of codon usage
patterns in *Hirudinaria manillensis* reveals a preference for GC-ending codons caused by dominant
selection constraints. BMC Genomics19: 542.

Hambuch, T.M., Parsch, J., 2005. Patterns of synonymous codon usage in Drosophila melanogaster genes
with sex-biased expression. Genetics 170: 1691-1700.

He, B., Dong, H., Jiang, C., Cao, F., Xu, L.A., 2016. Analysis of codon usage patterns in Ginkgo biloba
reveals codon usage tendency from A/U-ending to G/C-ending. Sci Rep. 6: 35927.

He, B., Dong, H., Jiang, C., Cao, F.L., Tao, S.T., Xu, L.A., 2016. Analysis of codon usage patterns in
Ginkgo biloba reveals codon usage tendency from A/U-ending to G/C ending. Sci. Rep. 6: 35927.

Hershberg, R., Petrov, D.A., 2009. General rules for optimal codon choice. Plos genet. 5: e1000556.

Jiang, Y., Deng, F., Wang, H., Hu, Z., 2008. An extensive analysis on the global codon usage pattern of
baculoviruses. Arch Virol. 153: 2273.

Kanaya, S., Kudo, Y., Nakamura, Y., Ikemura, T., 1996. Detection of genes in Escherichia coli sequences
determined by genome projects and prediction of protein production levels, based on multivariate
diversity in codon usage. Bioinformatics. 12: 213-225.

Kayum, M.A., Jung, H.J., Park, J.I., Ahmed, N.U., Saha, G., Yang, T.J., 2015. Identification and expression
analysis of WRKY family genes under biotic and abiotic stresses in *Brassica rapa*. Mol Genet
Genomics. 290: 79-95.

Kumar, N, Bera, B.C., Greenbaum, B.D., Bhatia, S., Sood, R., Selvaraj, P., Anand, T., Tripathi, B.N.,
Virmani, N., 2016. Revelation of influencing factors in overall codon usage bias of equine influenza
viruses. Plos one 11(4): e0154376.

Li, C.Y., Deng, G.M., Yang, J., Viljoen, A., Jin, Y., Kuang, R.B., Zuo, C.W., Lv, Z.C., Yang, Q.S., Sheng,

O., Wei, Y.R., Hu, C.H., Dong, T.,Yi, G.J., 2012. Transcriptome profiling of resistant and susceptible Cavendish banana roots following inoculation with *Fusarium oxysporum* f. sp *cubense* tropical race 4. BMC Genom. 13: 374.

Li, G.Y., Wang, Z., Zhang, Z.Y., Fang, H.D., Tan, X.L., 2013. The base composition and codon use of the WRKY gene family of the *Brassica napus*. J Biol. 30: 42-45.

Li, X., Tang, Y., Zhou, C., Zhang, L., Lv, J., 2020. A Wheat WRKY Transcription Factor TaWRKY46 Enhances Tolerance to Osmotic Stress in transgenic Arabidopsis Plants. Int J Mol Sci. 21: 1321.

Liu, H., He, R., Zhang, H., Huang, Y., Tian, M., Zhang, J., 2010. Analysis of synonymous codon usage in *Zea mays*. Mol Biol Rep. 37: 677.

Liu, H., Lu, Y., Lan, B., Xu, J., 2020. Codon usage by chloroplast gene is bias in *Hemiptelea davidii*. J Genet. 99: 8.

Liu, J.N., Fang, H.C., Liang, Q., Dong, Y.H., Wang, C.X., Yan, L.P., Ma, X.M., Zhou, R., Lang, X.Y., Gai, S.S., Wang, L.C., Xu, S.Q., Yang, K.Q., Wu, D.J., 2023. Genomic analyses provide insights into the evolution and salinity adaptation of halophyte *Tamarix chinensis*. GigaScience 12: 1–17.

Liu, Q., Feng, Y., Dong, H., Xue, Q., 2004. Synonymous codon usage bias in Oryza sativa. Plant Sci. 167: 101-105.

Liu, S., Qiao, Z., Wang, X., Zeng, H., Li, Y., Cai, N., Chen, Y., 2019. Analysis of codon usage patterns in "Lonicerae Flos" (*Lonicera macranthoides* Hand. -Mazz.) based on transcriptome data. Gene. 705: 127-132.

Ma, Q., Li, C., Wang, J., Wang, Y., Ding, Z., 2015. Analysis of synonymous codon usage in FAD7 genes from different plant species. Genet. Mol Res. 14: 1414-1422.

Mondal, S.K., Kundu, S., Das, R., Roy, S., 2016. Analysis of phylogeny and codon usage bias and relationship of GC content, amino acid composition with expression of the structural nif genes. J Biomol Struct Dyn. 34: 1649-1666.

Murray, E.E., Lotzer, J., Eberle, M., 1989. Codon usage in plant genes. Nucleic Acids Res. 17: 477-498.

Palidwor, G.A., Perkins, T.J., Xia, X.A., 2010. General model of codon bias due to GC mutational bias. PLoS One. 5: e13431.

Patel, P., Yadav, K., Srivastava, A.K., Suprasanna, P., Ganapathi, T.R., 2019. Overexpression of native Musa-miR397 enhances plant biomass without compromising abiotic stress tolerance in banana. Sci Rep. 9: 1-15.

Pegg, K.G., Coates, L.M., O'Neill, W.T., Turner, D.W., 2019. The Epidemiology of Fusarium Wilt of Banana. Front Plant Sci. 10, doi:10.3389/fpls.2019.01395.

Ploetz, R.C., Randy, C., 2015. Fusarium Wilt of Banana. *Phytopathology* 105, 12: 1512-21.

Qiu, S., Bergero, R., Zeng, K., Charlesworth, D., 2011. Patterns of codon usage bias in *Silene latifolia*. Mol Biol Evol. 28: 771-780.

Roberts, J.M., Carvalhais, L.C., O'Dwyer, C., Rincón-Flórez, V.A., Drenth, A., 2024. Diagnostics of Fusarium wilt in banana: Current status and challenges. Plant Pathol 73(4): 760-776.

Sharp, P.M., Emery, L.R., Zeng, K., 2010. Forces that influence the evolution of codon bias. Phil Trans R Soc B. 365: 1203-1212.

Sharp, P.M., Li, W.H., 1986. Codon usage in regulatory genes in *Escherichia coli* does not reflect selection for 'rare'codons. Nucleic acids Res. 14:7737-7749.

Sharp, P.M., Li, W.H., 1987. The codon adaptation index-a measure of directional synonymous codon usage bias, and its potential applications. Nucleic Acids Res. 15: 1281-95.

Siamak, S.B., Zheng, S., 2018. Banana Fusarium wilt (*Fusarium oxysporum* f. sp. *cubense*) control and

resistance, in the context of developing wilt-resistant bananas within sustainable production systems.

Horti Plant J. 4: 208-218.

Sueoka, N., 1988. Directional mutation pressure and neutral molecular evolution. PANS. 85: 2653-2657.

Sun, J., Zhang, J., Fang, H., Peng, L., Wei, S., Li, C., Zheng, S.J., Lu, J., 2019. Comparative transcriptome

analysis reveals resistance-related genes and pathways in *Musa acuminata* banana 'Guijiao 9' in

response to Fusarium wilt. Plant Physiol Bioch. 141: 83-94.

Swarupa, V., Ravishankar, K.V., Rekha, A., 2014. Plant defense response against *Fusarium oxysporum* and

strategies to develop tolerant genotypes in banana. Planta 239, 735–751.

Uddin, A., Mazumder, T.H., Barbhuiya, P.A., Chakraborty, S., 2020. Similarities and dissimilarities of

codon usage in mitochondrial ATP genes among fishes, aves, and mammals. IUBMB Life72: 899-914.

Wang, S.F., Su, M.W., Tseng, S.P., Li, M.C., Tsao, C.H., Huang, S.W., 2016. Analysis of codon usage

preference in hemagglutinin genes of the swine-origin influenza A (H1N1) virus. J Microbio Immuno.

49: 477-86.

Wani, S.H., Anand, S., Singh, B., Bohra, A., Joshi, R., 2021. WRKY transcription factors and plant defense

responses: latest discoveries and future prospects. Plant Cell Rep 40(7):1071-1085

Wei, L., He, J., Jia, X., Qi, Q., Liang, Z., Zheng, H., Ping, Y., Liu, S.Y., Sun, J.C., 2014. Analysis of codon

usage bias of mitochondrial genome in *Bombyx mori* and its relation to evolution. BMC Evol Biol. 14:

262.

Wright, F., 1990. The effective number of codons used in a gene. Gene. 87: 23-29.

Wu, Y., Zhao, D., Tao, J., 2015. Analysis of codon usage patterns in herbaceous Peony (*Paeonia lactiflora*

Pall.) based on transcriptome data. Genes 6: 1125.

Xin, W., Liu, Y., Yang, Y., Sun, T., Niu, L., Ge, J., 2020. Detection, genetic, and codon usage bias analyses

of the VP2 gene of mink bocavirus. Virus Genes 56: 306-315.

Zhang, X., Cai, Y., Zhai, X., Liu, J., Zhao, W., Ji, S., Su, S., Zhou, J.Y., 2018. Comprehensive analysis of

codon usage on rabies virus and other lyssaviruses. Int J Mol Sci. 19: 2397.

Zhao, Y., Zheng, H., Xu, A., Yan, D., Jiang, Z., Qi, Q., Sun, J.C., 2016. Analysis of codon usage bias of

envelope glycoprotein genes in nuclear polyhedrosis virus (NPV) and its relation to evolution. BMC

Genom. 17: 677.

Zhou, T., Weems, M., Wilke, C.O., 2009. Translationally optimal codons associate with structurally

sensitive sites in proteins. Mol Biol Evol. 26: 1571-1580.

Zorrilla-Fontanesi, Y., Pauwels, L., Panis, B., Signorelli, S., Vanderschuren, H., Swennen,

R., 2020. Strategies to revise agrosystems and breeding to control Fusarium wilt of banana. Nat Food

599–604.

(A)

(B)

Axis2(7.95%)

Axis1(21.94%)

A
T
G
C

Arg	AGG 1.800	AGA 1.328	CGG 0.923	CGC 0.832	CGA 0.638	CGU 0.480
Leu	CUC 1.752	CUG 1.614	UUG 1.126	CUU 0.804	CUA 0.403	UUA 0.302
Ser	AGC 1.452	UCC 1.279	UCG 1.155	UCU 0.915	UCA 0.674	AGU 0.534
Val	GUG 1.769	GUC 1.128	GUU 0.722	GUA 0.382	- 0	- 0
Thr	ACC 1.463	ACG 1.009	ACA 0.806	ACU 0.723	- 0	- 0
Gly	GGC 1.409	GGA 0.997	GGG 0.878	GGU 0.716	- 0	- 0
Ala	GCC 1.089	GCU 1.031	GCG 0.958	GCA 0.895	- 0	- 0
Pro	CCG 1.075	CCA 1.028	CCU 0.979	CCC 0.918	- 0	- 0
Ile	AUC 1.770	AUU 0.649	AUA 0.581	- 0	- 0	- 0
Cys	UGC 1.487	UGU 0.513	- 0	- 0	- 0	- 0
Lys	AAG 1.433	AAA 0.567	- 0	- 0	- 0	- 0
Phe	UUC 1.425	UUU 0.575	- 0	- 0	- 0	- 0
Asn	AAC 1.335	AAU 0.665	- 0	- 0	- 0	- 0
Glu	GAG 1.303	GAA 0.698	- 0	- 0	- 0	- 0
Tyr	UAC 1.289	UAU 0.711	- 0	- 0	- 0	- 0
Gln	CAG 1.188	CAA 0.812	- 0	- 0	- 0	- 0
His	CAC 1.133	CAU 0.867	- 0	- 0	- 0	- 0
Asp	GAU 1.004	GAC 0.996	- 0	- 0	- 0	- 0

1.8   0.9   0

GC3s content

ENC