1 **Compensatory evolution following deleterious episodes of GC-biased**

2 **gene conversion in rodents**

4   Marie Riffis[1], Nathanaëlle Saclier[1], Nicolas Galtier[1]

6   [1] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

8   Corresponding authors : Nicolas Galtier : nicolas.galtier@umontpellier.fr, Marie Riffis :

9   marie.riffis@umontpellier.fr

Abstract

GC-biased gene conversion (gBGC) is a widespread evolutionary force associated with meiotic recombination that favours the accumulation of deleterious AT to GC substitutions in proteins, moving them away from their fitness optimum. In many mammals recombination hotspots have a rapid turnover, leading to episodic gBGC, with the accumulation of deleterious mutations stopping when the recombination hotspot dies. Selection is therefore expected to act to repair the damage caused by gBGC episodes through compensatory evolution. However, this process has never been studied or quantified so far. Here, we analysed the nucleotide substitution pattern in coding sequences of a highly diversified group of *Murinae* rodents. Using phylogenetic analyses of about 70,000 coding exons, we identified numerous exon-specific, lineage-specific gBGC episodes, characterised by a clustering of synonymous AT to GC substitutions and by an increasing rate of non-synonymous AT to GC substitutions, many of which are potentially deleterious. Analysing the molecular evolution of the affected exons in downstream lineages, we found evidence for pervasive compensatory evolution after deleterious gBGC episodes. Compensation appears to occur rapidly after the end of the episode, and to be driven by the standing genetic variation rather than new mutations. Our results demonstrate the impact of gBGC on the evolution of amino-acid sequences, and underline the key role of epistasis in protein adaptation. This study contributes to a growing body of literature emphasizing that adaptive mutations, which arise in response to environmental changes, are just one subset of beneficial mutations, alongside mutations resulting from oscillations around the fitness optimum.

Key words : recombination, compensatory mutations, epistasis, *Hydromyini*, mutation load

Introduction

GC-biased gene conversion (gBGC) is a transmission bias associated with recombination that favours G and C alleles over A and T alleles. This evolutionary force, discovered in the 2000s, results from a repair bias in meiotic recombination that leads to non-Mendelian segregation favouring G and C alleles in highly recombining regions (Eyre-Walker 1999; Galtier et al. 2001). gBGC is widespread in the living world and numerous studies have demonstrated its presence in a wide range of organisms (Duret and Galtier 2009; Pessia et al. 2012; Galtier et al. 2018) including yeasts (Mancera et al. 2008; Lesecque et al. 2013), plants (Hämälä and Tiffin 2020), hymenoptera (Wallberg et al. 2015), birds (Bolívar et al. 2016; Smeds et al. 2016; Corcoran et al. 2017; Rousselle et al. 2019) and mammals, among which primates and rodents have been the most thoroughly studied (Galtier et al. 2009; Glémin et al. 2015; Williams et al. 2015; Halldorsson et al. 2016; Li et al. 2019). gBGC has a strong influence on the evolution of patterns of genomic variation. In particular, it drives genomic base composition by increasing the frequency of G and C alleles in regions with a high recombination rate (Pessia et al. 2012; Mugal et al. 2015), as demonstrated by numerous studies in mammals (Meunier and Duret 2004; Duret and Arndt 2008; Duret and Galtier 2009; Romiguier et al. 2010; Clément and Arndt 2013; Pracana et al. 2020). A striking example is the mouse *Fxy* gene, which has experienced a spectacular increase in GC content following its recent translocation into the high-recombining pseudoautosomal region (Galtier and Duret 2007). By increasing the frequency of GC alleles and decreasing the frequency of AT alleles at neutral sites, gBGC mimics the effect of natural selection, with AT to GC mutations having a higher probability of increasing in frequency and eventually reaching fixation, as if they were positively selected. Thus, gBGC affects the site frequency spectrum (SFS), the ratio of non-synonymous over synonymous substitutions (dN/dS) and the estimation of the adaptive rate, and can confound classical tests of positive selection (Galtier and Duret 2007; Berglund et al. 2009; Galtier et al. 2009; Ratnakumar et al. 2010; Bolívar et al. 2016; Corcoran et al. 2017; Rousselle et al. 2019).

Importantly, gBGC also applies to functional sites, where it interferes with both positive and negative selection. In particular, because gBGC tends to increase the frequency of GC alleles regardless of their fitness, it promotes the fixation of deleterious AT to GC mutations that would otherwise be eliminated by purifying selection. gBGC therefore creates an additional mutation load within genomes. The existence of such a gBGC load has been studied theoretically: Duret and Galtier (2009) suggested by simulation that gBGC causes an increase from 4% to 16% of the deleterious substitution rate, while Glémin (2010) modeled how gBGC affects fitness by maintaining recessive deleterious mutations at high frequency in populations. Several studies have also empirically demonstrated the existence of a gBGC

load that affects protein evolution in various organisms. Hämälä and Tiffin (2020) found that in *Arabidopsis thaliana*, protein evolution is strongly influenced by gBGC, which contributes to the accumulation of maladaptive mutations, limits the purging of deleterious mutations, and contributes to a global reduction in the efficiency of genome-scale selection. By analysing proteins with an accelerated rate of evolution, Berglund et al. (2009) and (Galtier et al. 2009) showed in primates that proteins can accumulate bursts of presumably deleterious amino-acid changes due to strong gBGC. In human populations, there is evidence for an increased load of disease-associated AT to GC mutations induced by gBGC (Necşulea et al. 2011; Lachance and Tishkoff 2014). In the mouse, after its translocation to the highly recombining pseudoautosomal region, the *Fxy* gene experienced the quick fixation of many deleterious AT to GC mutations, which would otherwise have been counter-selected (Galtier and Duret 2007).

The recombination rate varies not only among genomic regions, but also potentially in time. If the recombination map is evolutionarily unstable, such that the recombination rate of a particular locus goes up and down, we expect gBGC to be episodic. In many mammals, recombination is concentrated in hotspots, and the location of hotspots is controlled by the PRDM9 protein, which binds to specific DNA motifs (Baudat et al. 2010; Myers et al. 2010; Parvanov et al. 2010; Sandor et al. 2012). The PRDM9 gene and its binding motifs evolve very rapidly, resulting in a rapid turnover of recombination hotspots (Oliver et al. 2009; Myers et al. 2010; Berg et al. 2011; Lesecque et al. 2014; Baker et al. 2015; Latrille et al. 2017). Since gBGC is associated with recombination, this implies that gBGC should proceed in an episodic manner in mammals carrying a functional PRDM9, such as primates and rodents. Suppose that a recombination hotspot appears near a functional element (e.g. an exon). We expect a strong impact of gBGC during the episode, with an accumulation of deleterious AT to GC mutations within the concerned element, and we expect this effect to stop once the recombination hotspot dies and the gBGC episode ends. Consider a coding region affected by an episode of gBGC leading to the fixation of deleterious AT to GC mutations, thus altering the protein function. One might expect that once the recombination hotspot has died and the gene is no longer under strong gBGC, selection may act to repair the damage caused by the gBGC episode through compensatory evolution. By analysing the expected effect of gBGC on the distribution of fitness effect of new mutations, a recent theoretical study indeed suggested that gBGC episodes might considerably alter the evolutionary trajectory of the affected proteins (Joseph 2024). However, no empirical analysis of this process has been conducted so far.

Phylogenetic analyses have been carried out to identify functional sequences having undergone gBGC episodes in specific lineages (Galtier and Duret 2007; Galtier et al. 2009; Galtier 2021). This was achieved by mapping substitutions to the branches of the species

tree, and identifying clusters of AT to GC changes. A similar approach might be taken to

analyse the evolution of these functional sequences in downstream lineages to test whether

post-gBGC episode compensatory evolution is detectable - the expectation would be an

acceleration of the non-synonymous substitution rate after gBGC episodes. This requires a

high enough phylogenetic resolution, such that gBGC-induced and compensatory

substitutions happen in different branches of the tree and can therefore be distinguished from

each other.

Murinae rodents are a particularly suitable group to study these issues, because (1) they have

been shown to experience numerous and pronounced episodes of gBGC (Galtier 2021) ; (2)

in rodents a large proportion of recombination hotspots are determined by the PRDM9 gene

(Joseph et al. 2024) and there is a rapid turnover of these hotspots (Smagulova et al. 2016;

Latrille et al. 2017) ; (3) the house mouse (*Mus musculus*) genome is particularly well known

and reliably annotated; and (4) the recent sequencing of the entire exome of many Murinae

species offer a large sample of proteins with a particularly high phylogenetic resolution

(Roycroft, Achmadi, et al. 2021; Roycroft, MacDonald, et al. 2021).

However, it is important to note that gBGC in mammals and rodents is not always episodic.

This is because the recombination rate variation is not only regulated by PRDM9, and

hotspots can be determined by other means than PRDM9, even when it is present (Joseph et

al. 2024). At fine scale, PRDM9-independent hotspots have been documented (Axelsson et

al. 2012; Auton et al. 2013) and are often localised at the 5' end of genes in CpG islands

(Kawakami et al. 2017; Joseph et al. 2024; Raynaud et al. 2025). On a large scale, variation

in recombination rate can be determined by chromosome size or telomere proximity (Stapley

et al. 2017). These non-PRDM9 determinants of the localisation of recombination hotspots

can create regions where gBGC can persist over time at variable spatial scales, ranging from

Kb to Mb. Such a persistent effect of gBGC has been documented, for example, in the

genomes of gerbils, a group close to Murinae, where the authors found large clusters of about

100 Kb of increasing GC content (Pracana et al. 2020). To study compensation, we should

therefore be careful to target only gBGC events that are localised and short-lived, when it

makes sense to investigate the protein response once the episode has ended. If gBGC is less

precisely located and persistent in time, it is likely to be difficult to distinguish compensatory

evolution from the continuing effect of gBGC.

Here, we used phylogenetic analyses of 71,040 coding exons from 48 species of Murinae

rodents from the Hydromyini tribe to search for a signature of compensatory evolution after

gBGC episodes have damaged coding sequences. Hydromyini is an Australasian taxon that

originated about 10 Mya and is characterised by a particularly high diversification rate

(Roycroft, MacDonald, et al. 2021; Roycroft, Achmadi, et al. 2021). This makes it a particularly

appropriate group to carry out compensation analyses, as they provide us with a high

phylogenetic and temporal resolution. By analysing synonymous substitutions, we first
identified exon-specific, lineage-specific gBGC episodes. We then analysed the molecular
evolution of the affected exons, as well as other exons of the same gene, after the episodes,
focusing on the non-synonymous variation. We uncover ubiquitous compensatory evolution
after gBGC episodes and show that protein re-adaptation in Hydromyini rodents is rapid and
probably driven by the standing genetic variation.
153
154 **Results**
155
156 <u>Overview of the data</u>
157
We analysed whole-exome data from 48 species of *Hydromyini*, a highly diversified group of
Australasian *Murinae*. Using the house mouse genome as a reference for orthology calling,
we generated 71,040 high-quality alignments of coding exons from 15,415 genes. We only
kept the exons found in at least 3 species, the average number of species per exon being 32
(Fig. 1A). The average, minimum, and maximum number of aligned exons per gene were 4.6,
1, and 67, respectively; and 12,158 genes had 3 or more aligned exons (Fig. 1B). The
average exon length, GC12 and GC3, were 241 bp (exons <100pb were discarded), 49.45%
and 56.11%, respectively. We reconstructed the species phylogeny (Fig. 1C) based on the
concatenation of these alignments. The phylogeny was well resolved topologically, with a
bootstrap support of 100% for all internal branches. A high temporal resolution was also
achieved: the tree contained many short branches, with an average branch length of 0.0038
substitutions per site. The distributions of exon length, branch length, GC3 and GC12 are
shown in Supplementary Figure S1A.
We mapped synonymous and non-synonymous AT to GC (Weak to Strong : WS), GC to AT
(SW) and GC-conservative (SSWW) nucleotide substitutions in all branches of the phylogeny,
for each exon (Fig. 2A & B). We report a total of 1,855,929 synonymous substitutions and
1,320,787 non-synonymous substitutions. The counts for each type of substitution are
provided in Supplementary Table S1.
176

177
178 **Figure 1: Overview of the *Hydromyini* dataset analysed in this study. A**: distribution of
179 the number of species per gene alignment (vertical line represents the average) ; **B**:
180 distribution of the number of exons analysed per gene (vertical line represents the average) ;
181 **C**: phylogeny of the 48 *Hydromyini* species, reconstructed from the concatenation of all
182 exons.

Figure 2: Summary of the analyses. A: For each exon, the data consist of a pruned tree derived from the reference phylogeny (Fig. 1C) and a corresponding DNA sequence alignment. **B**: Substitution mapping was performed for each exon, distinguishing each type of substitution. **C**: A gBGC episode (red) in a specific exon and branch was detected by identifying clusters of synonymous WS substitutions that exceed random expectations in both the tree and sequence. Based on synonymous WS and SW substitution counts across the affected branch, its descendants, the targeted exon, and other exons of the same gene, gBGC episodes were annotated as "punctual," "two-branch," or "all exons," depending on their extent. Only "punctual" episodes were retained for compensatory evolution analyses. **D**: To test for compensatory evolution after gBGC episodes, the number of non-synonymous substitutions in descending branches was compared to an expected distribution generated by randomizing substitutions. A significant excess of non-synonymous substitutions indicates compensatory evolution.

<u>gBGC episodes</u>

Detection of the episodes

Our first aim was to identify gBGC episodes by analysing the clustering of synonymous WS
(S-WS) substitutions across branches and exons (Fig. 2C). Specifically, we defined a gBGC
episode when we detected an excess of S-WS substitutions in a specific branch for a specific
exon, compared to a random distribution of substitutions (see Material and Methods). Using
this method, we detected a total of 29,845 gBGC episodes. gBGC episodes occurred in
slightly longer exons and branches than the overall mean (240 bp and 0.0038 substitutions
per site, respectively), with an average exon length of 267 bp and an average branch length of
0.0054 substitutions per site. The GC-content distribution of exons affected by one or several
episodes was indistinguishable from the overall distribution, suggesting a homogeneous
distribution of episodes across genes. The distributions of exon length, branch length, GC3
and GC12 of episodes are available in Supplementary Figure S1B. As expected, we observe
an overrepresentation of S-WS substitutions compared to synonymous SW (S-SW)
substitutions in most branches detected as episodes (Fig. 3).

Figure 3: Weak-to-Strong dS (dS$_{WS}$) and Strong-to-Weak dS (dS$_{SW}$) in branches without
(in grey) and with gBGC episodes (in red). For branches with episodes, data are shown
for all types of episodes ("All episodes") and for episodes affecting a single branch in a
single exon ("Punctual episodes"), which are the subset used in compensation analyses. dS
corresponds to the number of synonymous substitutions divided by the number of third
codon positions, weighted by GC3 (for SW) or by AT3 (for WS). For a better visualization,
we show here only a random sample of our data : 100,000 points for "No episode", and for
episodes a number of points which is proportional to the proportion of the category in all
data.

Overflowing of the episodes and selection of punctual episodes

Our rationale involves analysing gBGC episodes that are localised and short-lived ('PRDM9-

like' episodes), *i.e.* episodes affecting a single branch in a single exon. To check whether the
detected episodes meet this criterion, we looked at the number of S-WS substitutions in the
descending branches and in the other exons of the same genes. We compared the observed
amount with an expected distribution obtained via 1000 randomisations of the substitutions.
We observed an excess of S-WS substitutions and a deficit of S-SW substitutions in the
branches immediately downstream of the episodes (Fig. 4A), as well as in the episode's
branches in the other exons of the same gene where the episodes were detected (Fig. 4B).
This excess of S-WS and deficit of S-SW substitutions is a typical feature of gBGC and
indicates that some of the detected episodes are not punctual, but rather extend into more
than one branch and/or more than one exon.
274

275
Figure 4: Overflowing of the gBGC episodes. A: in the downstream branches in the focus
exon ; **B**: in the focus branches in other exons of the same gene. Vertical lines correspond to
the total observed number of synonymous substitutions summed on all branches and exons
concerned. Density curves correspond to the expected distribution of the total number of
synonymous substitutions summed on all branches and exons concerned. ** : p-value <
0.01. Left panels (red): S-WS substitutions. Right panels (blue): S-SW substitutions.
282
283
We therefore developed an ML method (Supplementary Methods) to classify the gBGC
episodes into 3 categories : punctual (localised, short-lived episodes occurring in a single
exon and a single branch of the tree, the ones we are interested in), two-branch (episodes
occurring in a single exon but affecting the downstream lineages) and all-exons (gene-wide

gBGC) (Fig. 2C). This only concerns episodes having occurred in internal branches of the
tree. 18,730 terminal-branch episodes were therefore discarded at this stage. Single-exon
genes were also excluded at this stage since they do not allow distinguishing the two-branch
and all-exons categories - there were 600 of them. The best-fit parameters modeled our set of
detected episodes as a mixture of punctual, two-branch and all-exons episodes in proportions
0.65, 0.28 and 0.07, respectively, with an equilibrium GC3 of 94.0%, 94.6% and 84.7%,
respectively. We then assigned a category to each episode according to the episode-specific
posterior probabilities calculated by the method (Supplementary Methods, equations 8-10).
364 episodes (3.35%) were considered to belong to neither of these categories per their
Goodness of Fit statistics (GoF < 0.2). 7,572 gBGC episodes were classified as punctual and
deemed suitable for compensatory evolution analysis. Figure 3 shows that the ratio of S-SW
to S-WS substitution rate tends to be higher in episodes annotated as punctual (Fig. 3C)
compared to the whole set of episodes (Fig. 3B).

Non-synonymous evolution in "punctual" episode-carrying branches

There were 6.9 times more non synonymous WS (NS-WS) substitutions in branches carrying
a punctual episode than in branches without episodes. This ratio was much lower for the other
types of substitutions: it was 2.7 and 4.7 for the SW and SSWW types, respectively. The ratio
remained almost unchanged after correction for exon and branch length (Supplementary
Table S1). This confirms our working hypothesis that gBGC episodes are responsible for an
increase in the rate of NS-WS (potentially deleterious) substitutions. 2,159 of the episodes
identified as punctual were associated with at least one NS-WS substitution, while 5,413 had
none.

Compensatory evolution analyses

Our second aim was to investigate whether gBGC episodes are followed by a phase of
compensatory protein evolution (Fig. 2D). To do this, we focused on non-synonymous (NS)
substitutions, and analysed whether they tend to be in excess in the branches downstream of
gBGC episodes. We therefore counted the number of NS substitutions in the two daughter
branches of punctual episode-carrying branches, and compared it with a null distribution
generated by randomising NS substitutions across exon trees. We made this comparison
separately for two types of episodes : (1) those associated with at least one NS-WS
substitution, *i.e.* episodes that have potentially produced deleterious substitutions after which
compensation might happen; and (2) those not associated with any NS-WS substitution, *i.e.*
episodes after which *a priori* no compensation is expected. Here we only considered punctual

episodes occurring in an internal branch of the species tree, for which downstream lineages
can be examined. There were 7,676 such episodes, of which 2,159 were associated with a
NS-WS substitution. The two types of episodes did not seem to differ in terms of gBGC
strength, with an average number of S-WS substitutions per site of 0.014 and 0.015 for
episodes with and without NS substitutions, respectively. We carried out this analysis in the
exons affected by the episodes, but also in the other exons of the same gene.

Signal of compensatory evolution after "punctual" gBGC episodes

In exons affected by punctual episodes, we observe a significant excess of NS substitutions
after episodes with at least one NS-WS substitution, but no such excess after episodes
without any NS-WS substitution (Fig. 5A). This is consistent with the hypothesis that proteins
respond to deleterious gBGC episodes by a phase of compensatory evolution. Interestingly,
this response is largely due to WS substitutions, and also to SSWW substitutions, but not to
SW substitutions (Fig. 5B). As a control, we performed the same analyses with the two other
types of episodes: the "two-branch" episodes (Supplementary Figure S2) and the "all-exons"
episodes (Supplementary Figure S3), and we did not find the same signal of compensation as
after the "punctual episodes". For the two-branch episodes, we observed an excess of NS-WS
substitutions in the descending branches irrespective of the presence of a substitution during
the episode. This suggests that this excess of NS substitutions after "two-branch" episodes is
probably not due to compensation but rather to the overflow of the episodes in the descending
branches. We observed a similar signal for the "all-exons" episodes, although it is less clear,
maybe due to the small sample size. We also reconducted the analysis using more stringent
criteria of topological concordance (see Materials and Methods) between exon and reference
trees, and found that the results were qualitatively unchanged (Supplementary Figure S4).

Figure 5: Compensation after "punctual" gBGC episodes. A: Total number of non-

synonymous substitutions observed in branches following the episodes compared with the
expected distribution (summed on all exons and branches concerned). Distinction is made
between episodes with at least 1 NS-WS substitution (upper panel) and episodes with no
NS-WS substitution (lower panel). **B**: Same thing detailed for each of the 3 types of
substitutions (red: WS, blue: SW and green: SSWW). n.s.: p-value >= 0.05 ; ** : p-value <
0.01.
359
360 *The signal of compensation is only detectable after short branches*
361
We reconducted the above analysis after splitting the data into 5 bins of episode-carrying
branch length (Fig. 6). The results show that when episodes happen in branches longer than
~0.006 substitutions per site, the compensation signal is no longer detectable in the
descending branches. This suggests that compensation occurs quickly after the end of gBGC
episodes, such that the episode and the compensatory phase can both happen in the same
branch if long enough.

total number of NS substitutions

Figure 6: Compensation after "punctual" episodes is only detected after short
carrying-episode branches. The data were separated in 5 bins of episode-carrying branch
lengths: <0.002 (n=628); >=0.002 and < 0.004 (n=778), >=0.004 and <0.006 (n=434),
>=0.006 and <0.008 (n=92) and >=0.008 (n=227). n.s.: p-value >= 0.05 ; *: p-value < 0.05 ;
**: p-value < 0.01.

No detectable excess of substitutions before episodes

An excess of NS substitutions after punctual episodes with NS substitutions (Fig. 5A) is

expected as soon as any process, biological or artefactual, favoring the presence (or
inference) of NS substitutions in connected branches is at work. One such possibility is time-
autocorrelation of substitutions, in the absence of compensatory evolution. Suppose for
instance that the NS substitution rate of an exon varies in time and among clades, perhaps
due to fluctuations of the overall selective pressure. In such a case, branches from clades with
higher substitution rates would carry more substitutions than expected under our
randomisation procedure, which might lead to excessive co-occurrence of substitutions in
connected branches. To make sure the observed pattern indeed reflects a phenomenon of
compensation, we conducted a similar analysis using episode ascending branches instead of
descending branches. This analysis revealed no excess of NS substitutions in branches
immediately upstream of punctual episodes, irrespective of the existence of a NS substitution
within the episode (Supplementary Figure S5). We conclude that the excess of NS
substitution in the downstream branches of an episode indeed reflects an evolutionary
response to episode-associated NS substitutions, rather than a clustering of substitutions
among connected branches.

Compensatory substitutions occur close to deleterious substitutions

We focused on predicted compensation cases where there was exactly one (supposedly
deleterious) NS-WS substitution in the episode branch and exactly one (supposedly
compensatory) NS substitution of any type in the descending branches (n=456 cases). We
tested whether the predicted compensatory substitutions occurred at positions closer to the
predicted deleterious substitution than expected by chance. This was achieved by counting
the number of cases where the two substitutions were at a given distance from each other,
with distances ranging from 1 to 12 codons and divided into 3 bins (Fig. 7). We found that this
happened significantly more often than expected by chance for short distances (from 1 to 4
codons) (p-value < 0.01, Fig. 7), and that the signal decreased as the distance increased,
suggesting that the predicted deleterious and compensatory substitutions are structurally
related.

408
Figure 7 : Predicted compensatory substitutions occurred at sites close to the
predicted deleterious substitutions. The plots represent the total number of compensatory
substitutions that are at a given distance from the deleterious substitutions, for
compensatory cases where there is exactly 1 deleterious NS-WS substitution and exactly 1
compensatory NS substitution (n=456). The distances between the deleterious and the
compensatory substitutions are divided into 3 bins, from upper panel to lower panel : 1-4
codons, 5-8 codons, 9-12 codons. The vertical line corresponds to the observed value and
the distribution corresponds to the expected values based on 10,000 randomisations of the
positions of the compensatory substitutions. **: p-value < 0.01 ; n.s.: p-value >= 0.05.
418
419

Compensation in the other exons during but not after punctual episodes

We also looked for potential compensation in the other exons of the same gene, during the
episode (*i.e.* in the same branch) and after the episode (in the descending branches). We
observed a significant signal of compensation during the episodes (Fig. 8A), but not after (Fig.
8B). We carried out the same analysis using only the other exons adjacent to the affected
ones, which gave a very similar pattern.

Figure 8 : Compensation analyses in other exons of the same gene for "punctual"
episodes. Observed value and expected distribution of the sum of the total number of NS
substitutions of all types in other exons. **A**: during episodes (in the same branches) ; **B**: after
episodes (in the descending branches). Distinction is made between episodes with at least 1
NS-WS substitution (upper panel) and episodes without NS-WS substitution (lower panel).
n.s.: p-value >= 0.05 ; **: p-value < 0.01.

Discussion

Analysing the nucleotide substitution pattern in the exon coding sequences of 48 species of
Hydromyini, a highly diversified group of Murinae rodents, we detected exon-specific, lineage-
specific gBGC episodes and found evidence for compensatory evolution in the descending
branches.

<u>Prominent episodic gBGC in Hydromyini</u>

We found that the nucleotide substitution pattern in *Hydromyini* is characterised by a
clustering of WS substitutions, as found by Galtier (2021) at the Murinae scale, and by a an
overrepresentation of dS_{WS} compared to dS_{SW} in most branches, as reported in the gerbil
(*Gerbillinae*), brown rat (*Rattini*), and house mouse (*Murini*) lineages (Pracana et al. 2020).
We identified >30,000 episodes, *i.e.*, branches with a significant excess of S-WS substitutions
in a particular exon. This is much higher than the few hundreds analyzed by Berglund et al.
(2009) and Galtier et al. (2009) in hominid primates. In *Hydromyini* 32% of exons and 100% of
genes were affected by at least one gBGC episode. These results underscore the prominence

of gBGC in this group, as previously suggested in other taxa of Murinae (Pracana et al. 2020;
Galtier 2021). They confirm that *Hydromyini* are an appropriate group to study episodic gBGC
and its consequences on protein evolution.
We highlight that a small but significant fraction of the detected gBGC episodes, referred to
"all-exons"(about 7%), extend throughout the gene (and perhaps beyond) and persist over
time - in contrast to the transient nature expected in "PRDM9-like" episodes. This suggests
that in *Hydromyini,* large (>10kb) DNA segments undergo prolonged phases of GC-
enrichment, as described by Pracana et al. (2020) in *Gerbillinae*. The mechanism behind this
regional effect remains unclear and warrants further investigation (Brekke et al. 2023). Most of
our detected episodes (around 65%) were classified as "punctual", *i.e.*, were predicted to have
occurred in a single exon and a single branch. These are suitable for studying compensatory
evolution after a phase of intense gBGC. As we expected, gBGC episodes induced an
increased rate of NS-WS substitutions, as previously reported in primates (Berglund et al.
2009; Galtier et al. 2009). Many of these substitutions are potentially deleterious (Glémin
2010), perhaps calling for a compensatory response.
Somewhat surprisingly, gBGC episodes were also accompanied by an increase in the rate of
SW and SSWW substitutions - albeit weaker than the WS one (Supplementary table S1). We
identify at least two non-exclusive explanations for this result. First, recombination is known to
be mutagenic (Halldorsson et al. 2019). gBGC episodes are therefore likely associated with
many new mutations of all types, which might eventually reach fixation and contribute
substitutions along the same branch. Secondly, it should be recalled that genealogies, and
particularly branch lengths, differ among loci due to the variance of the coalescent (Rannala et
al. 2020). This is expected to create a correlation between substitution counts across
categories: exon genealogies containing by chance one or several particularly long branches
are likely to exhibit elevated substitution counts of all kinds in those branches.
479
480   Post-episode compensatory evolution
481
482   Once a gBGC-induced deleterious mutation has occurred, we expect that other mutations
483   occurring at different sites might compensate for the deleterious effect of the first mutation and
484   eventually reach fixation, leading to an increase in the NS substitution rate after the gBGC
485   episode.
486
We report a significant excess of NS substitutions after "punctual" gBGC episodes having
caused a NS-WS substitutions (potentially deleterious episodes), but not after those that have
not (non-deleterious episodes) (Fig. 5A). We suggest that the excess of NS substitutions after
"punctual" deleterious episodes probably reflect compensatory evolution by epistasis in

response to the alteration of the function of the hit protein. Compensation was detectable here
thanks to the high temporal resolution of the Hydromyini tree, which contains many short
branches (Fig. 1C). Indeed, the signal of compensation in the descending branches was only
detectable when the branch carrying the episode was shorter than 0.006 substitutions per site
(Fig. 6). Such short branches are rare in typical species trees. The rapid radiation of
Hydromyini (about 50 species with a ~10 My old common ancestor ; Roycroft, Achmadi, et al.
2021) appears to offer a particularly favorable context for the study of compensatory
evolution.

We performed a number of controls to confirm that the detected excess of NS substitutions
after episodes is a compensatory response to gBGC, rather than an effect of overflowing
gBGC that would continue to "break up" the sequence in the descending branches. First, we
note that the signal of compensatory evolution is clearly detectable only in "punctual"
episodes, not in the "two-branch" and "all-exons" ones (Supplementary Figures S2 & S3). This
does not suggest that gBGC overflowing explains our result, even though our assignment of
episodes into the three categories is probably unperfect. Secondly, if the post-episode excess
of NS substitutions were due to overflowing of gBGC, we would expect to detect it after non-
deleterious episodes as well. Indeed, non-deleterious episodes do not appear weaker than
deleterious ones, as they do not contain less S-WS substitutions on average, even after
correcting for sequence length and GC-content. Moreover, the excess of NS substitutions was
observed not only for the WS substitutions, but also for the GC-conservative SSWW
substitutions, which are not expected to be directly affected by gBGC. Finally, the absence of
an excess of NS substitutions in the ascending branches of deleterious episodes
(Supplementary Figure S5) confirms that the signal we report in Figure 5 reflects an
evolutionary response to gBGC episodes, not an exon-specific clustering of NS substitutions
in specific portions of the tree due to, *e.g.*, a transient increase in the NS substitution rate.

Our analysis uses a common tree topology for all exon alignments, thus assuming that the
consensus Hydromyini phylogeny (Fig. 1C) appropriately depicts the phylogeny of every
analyzed exon - please note that alignments strongly departing from this assumption were
filtered out, see Material and Methods. Still, it is likely that, for some alignments, the true exon
phylogeny differs from the consensus tree, owing to, *e.g.*, incomplete lineage sorting or gene
flow. Topological differences between the true and assumed trees are expected to affect
substitution mapping, and particularly, generate erroneous inferred substitutions, possibly
across connected branches. Note, however, that our analysis is unlikely to be strongly
affected by the problem of false multiple substitutions at the same site - the typical
consequence of exon tree/species tree discordance. Indeed, the episode-associated and

compensatory substitutions we detect occur at distinct sites. It is difficult to conceive how
discordance could result in mismapping a substitution to the episode-carrying branch at site X,
and to a downstream branch at site Y. This does not suggest that exon tree/species tree
discordance is a major confounder, as confirmed by the fact that our results were robust to an
increased stringency in low-discordance exon selection (Supplementary Figure S4).

In the cases where there was exactly one NS-WS (probably deleterious) substitution in the
episode branch and exactly one NS (probably compensatory) substitution in the descending
branches, we showed that the downstream substitution occurred in positions close to the
upstream one more often than expected by chance (Fig. 7). This is an additional argument in
favour of the hypothesis of compensatory evolution in response to gBGC. Indeed, this is
consistent with the report that, in a wide range of taxa covering Virus, Prokaryotes and
Eukaryotes, the distance between experimentally confirmed compensatory and deleterious
mutations in the protein sequence is shorter than expected by chance (Poon and Chao 2005;
Davis et al. 2009). We also detected a more distant response, in other exons of the same
gene (Fig. 8A). This might correspond to compensatory substitutions at sites close to the sites
of the deleterious mutation in the tertiary structure of the protein (Davis et al. 2009; Ivankov et
al. 2014; Storz 2018; Chaurasia and Dutheil 2022). Compensatory evolution in other exons
was detected in the episode-carrying branch but not in downstream branches (Fig. 8B).
Further analyses of this effect should take into account the tertiary structure of the affected
proteins and the predicted effect of the supposedly deleterious and supposedly compensatory
substitutions.

New mutations or standing variation?

Positive selection on beneficial alleles for (re)adaptation can occur either from new beneficial
mutations or from pre-existing (=standing) genetic variation, meaning newly beneficial alleles
that were neutral or deleterious before the environmental change (in the case of adaptation),
or before the change in genetic background (in our case of readaptation) (Barrett and Schluter
2008). Adaptation from standing variation is expected to be faster than adaptation from new
mutations: pre-existing beneficial alleles are already present (no waiting time) and start at a
higher frequency than new beneficial mutations (higher fixation probability). As a result,
adaptation occurs mainly from standing variation when it occurs over short timescales (Barrett
and Schluter 2008; Illingworth et al. 2012). We found that compensatory substitutions are
detectable after episodes occurring in short branches, but not after episodes occurring in long
branches (Fig. 6). A natural explanation of this result is that compensation happens rapidly
after the episode ends (Supplementary Figure S6). According to this interpretation, when the

episode-carrying branch is longer than ~0.006, the compensatory substitutions happen in the
same branch as the deleterious ones, and cannot be detected by our approach. This 0.006
threshold is roughly twice the average heterozygosity of Hydromyini exons (Supplementary
Figure S1), suggesting that the response time is of the order of magnitude of the lifespan of a
polymorphic mutation. Furthermore, we have shown that compensation occurs mainly via WS
substitutions, and to a lesser extent SSWW ones (Fig. 5B). This is intriguing at first sight, but
makes sense if one thinks of compensation from standing variation. Since a gBGC episode
tends to increase the population frequency of polymorphic WS mutations, and decrease the
SW ones, compensatory mutations picked up from the post-episode standing variation are
likely to be of the WS kind, and unlikely to be of the SW kind. If however compensation
happened mainly from new mutations, we would expect all mutation types to respond
similarly. These two observations – fast response, GC-biased compensatory substitutions –
argue in favor of standing genetic variation, not new mutations, being the dominating source
of compensatory beneficial alleles after gBGC episodes in Hydromyini.
The relative contribution of the standing variation versus new mutations to adaptation has
been studied in different ways. Experimental evolution studies, mainly in yeast, have yielded
mixed conclusions, some suggesting that adaptation is mainly driven by standing variation
(Burke et al. 2014), while others suggested that both sources of variation play a role (Ament-
Velásquez et al. 2022). Similar conclusions about the contribution of both the standing
variation and new mutations were drawn from studies that identify the origin of adaptation-
associated mutations by comparing the genome sequences of a presumably adapted lineage
(*e.g.* polar bear) and its presumably non-adapted source population (brown bear) (Samaniego
Castruita et al. 2020; Estandía et al. 2023). Finally, a widely used method, applicable to
diverse organisms, is to analyse population genomic patterns, with the assumption that
adaptation from standing variation leaves a weaker pattern of selective sweep than expected
from new mutations (Barrett and Schluter 2008). These soft selective sweeps have been
investigated using large-scale polymorphism data analyses and found to be pervasive in
various organisms, such as *D. melanogaster* (Garud et al. 2015) or humans, in which standing
variation was suggested to be the dominant mode of adaptation (McManus et al. 2017;
McManus et al. 2017). However, methods to distinguish between soft and hard sweeps are
complex and confounded by neutral demographic processes, making such inferences tricky
(Harris et al. 2018). Here we explore a new way to assess the origin of readaptation after
gBGC in rodent species and find that it seems to be mainly driven by standing variation rather
than new mutations, in line with previous studies having highlighted the important role of
standing genetic variation in adaptation. This approach might easily be extended to other
taxa, gBGC being ubiquitous, and in particular to investigate a potential effect of effective
population size. If mutation limits adaptation (Rousselle et al. 2020), then small-Ne taxa such

as primates might lack the genetic diversity necessary for a rapid response from standing
variation, and instead resort to new mutations.

The above piece of discussion points to a process whereby a (future) compensatory mutation
is already present at polymorphic stage during the episode, increases in frequency as the
episode unfolds, and reaches fixation after the episode has ended. This standing variation-
based scenario raises the possibility that the compensatory mutation reaches fixation not in
just one of the two descending lineages, but in both. If it happened, such a convergent
compensatory process would fail our detection method: because in this scenario the two
descending lineages carry the same state, our method would presumably map a single
substitution in the episode-carrying branch, instead of one substitution in each of the
descending branches. Our approach is therefore conservative with this respect and probably
underestimates the actual prevalence of compensatory evolution, to an extent that needs to
be quantified.

Epistasis vs. back mutations

Compensation is a special case of epistasis, when the fitness of an allele depends on the
allelic state at other loci. The literature on epistasis is rich, with most studies on the topic
agreeing that epistasis is ubiquitous and has a major impact on the long-term evolutionary
trajectories of proteins (Ivankov et al. 2014; Starr and Thornton 2016; Storz 20018). There are
three main ways to study epistasis in the existing literature (reviewed in Starr and Thornton
(2016)). (1) Experimental studies of specific cases (few proteins) where deleterious genotypes
are created and readaptation by compensatory mutations is studied in the lab (Poon et al.
2005; Filteau et al. 2015; Rojas Echenique et al. 2019; Serrano et al. 2021). (2) Search for
coevolving sites across phylogenies, which works well in the case of a long-term pairwise
interaction, but loses power if multiple sites are in interaction. This applies to studies
identifying pairs of sites that tend to undergo substitutions in the same branches (Chaurasia
and Dutheil 2022), as well as studies detecting an effect of substitutions at one site on the
substitution rate at another site (e.g. Neverov et al. 2021). (3) Comparative sequence
analysis, which typically showed that some disease-associated mutations in humans are fixed
in other taxa, indirectly demonstrating the existence of compensatory epistasis (Kern and
Kondrashov 2004; Azevedo et al. 2009). Of note these studies, while agreeing that epistasis
is prevalent, are largely based on a small number of genes with well-characterised functions.
Episodic gBGC, which promotes deleterious alleles throughout the genome (and also
prevents their elimination by purifying selection), allowed the analysis of a large sample of
proteins having undergone functional degradation. Our study thus provides a new and original
perspective on the question, as we have studied at the whole exome level the readaptation of

proteins temporarily displaced from their optimum. By considering gBGC-associated NS
substitutions, we focus on a pool of substitutions presumably enriched in deleterious changes.
This offers the opportunity of analyzing compensatory evolution irrespective of the
identification of pairs of sites with correlated substitution patterns. Indeed, even a residue
experiencing a single gBGC-associated deleterious substitution across the whole tree, which
in turn induces a single compensatory substitution, can contribute to the signal we detect. Our
results confirm that compensation by epistasis appears to be ubiquitous in protein evolution.

Experimental studies of compensation generally create highly deleterious genotypes and
study protein readaptation starting far from the optimum. Poon and Chao (2005), for instance,
introduced missense mutations in the φX174 bacteriophage genome, which reduced fitness
by more than 50% in most mutants. In our case, the re-adapting proteins probably start
relatively close to their optimal state since gBGC is a weak evolutionary force (Lartillot 2013;
Galtier 2021). This enabled us to show that epistasis is pervasive even under weak selection,
a situation that presumably concerns a large number of mutations, and is relevant for
understanding patterns of genomic variation. This may explain why we observed a lower ratio
of compensatory to deleterious mutations (not even 1:1, Fig. 5) than in experimental studies.
For example, Poon and Chao (2005) reported an   average of 11.8 compensatory mutations
for 1 deleterious mutation.

Although our study focuses on epistatic compensation, it should be recalled that re-adaptation
to the optimal state can also be achieved by back-mutations, *i.e* mutations occurring at the
same site as the deleterious mutation and allowing a return to the initial state. Here we were
not able to detect such back-mutations, probably missing a part of the re-adaptation.
Comparing polymorphism and substitution patterns, (Latrille et al. 2024) have recently shown
that back-mutations represent a substantial proportion of beneficial mutations in mammalian
proteins. Moreover, a recent study has theoretically investigated the process of back-mutation
after gBGC (Joseph 2024). Applying plausible amounts of gBGC to empirical fitness
landscapes, this study shows that, by promoting the fixation of deleterious WS mutations and
taking the protein away from its fitness optimum, gBGC significantly alters the distribution of
fitness effects of future mutations, creating opportunities for beneficial back-mutations in
substantial numbers. Back-mutations therefore appear to represent a significant proportion of
the beneficial mutations that enable a return to the lost optimum. To quantify the relative
contribution of compensatory (epistasis) vs. back-mutations in post gBGC episode protein re-
adaptation is an appealing goal. If immediate, back-mutations allow the protein to return to the
optimum in a single step - an efficient re-adaptation process. If however a pre-existing
epistatic compensatory mutation increases in frequency or reaches fixation before the back

mutation happens, then this might be less beneficial, the genetic background having changed.
The dynamic of back-mutation in the context of epistasis has been theoretically investigated
by McCandlish et al. (2016), who showed that the rate of back-mutation is expected to
decrease with time when the site is involved in at least one epistatic interaction, and that
back-mutations tend to occur either almost immediately after the deleterious substitution, or
much later. The interaction between gBGC and selection might offer the opportunity to
empirically test these predictions.

<u>Perspectives on gBGC as a confounding effect in detecting adaptation</u>

Detection of adaptation is a widely studied topic in evolutionary biology and methods for
inferring positive selection are based on the detection of rapid fixation events that result in an
increased substitution rate (McDonald and Kreitman 1991). It has been largely demonstrated
that gBGC can counfound these tests by promoting the fixation of deleterious WS mutations
(Galtier and Duret 2007; Berglund et al. 2009; Galtier et al. 2009; Ratnakumar et al. 2010;
Bolívar et al. 2016; Corcoran et al. 2017; Rousselle et al. 2019). Another previously neglected
confounding effect has recently emerged in the literature, namely the confusion between
beneficial and adaptive mutations (Stolyarova et al. 2020; Joseph 2024; Latrille et al. 2024).
These authors point to the difference between beneficial mutations, which increase fitness,
and adaptive mutations, which are the subset of beneficial mutations responding to an
environmental change. In mammals, it has been quantified that non-adaptive beneficial back-
mutations, which occur after a fitness decrease (*e.g.* due to drift or to gBGC), account for a
large proportion (between 20% and 40%) of beneficial mutations (Latrille et al. 2024). We
showed that, in Hydromyini, episodic gBGC contributes to this effect by moving proteins away
from their optimum, leading to the fixation of compensatory mutations that are beneficial but
not adaptive. Our study therefore demonstrates the influence of gBGC on protein evolution,
and joins this early literature highlighting the non-negligible proportion of non-adaptive
mutations among beneficial mutations, here focusing on compensatory mutations instead of
back-mutations.

Materials and Methods

<u>Data, Data cleaning, Sequence Assembly</u>

Hydromyini whole-exome sequencing raw reads generated by Roycroft, MacDonald, et al.
(2021) and Roycroft, Achmadi, et al. (2021) were downloaded from the NCBI database
(references : PRJNA729818 and PRJNA705792). This dataset is composed of 48 species, of

which 5 are extinct. Reads came from a single individual in 42 species, from 2 individuals in 3
species, and from 3 individuals in 3 species. Raw reads were quality trimmed using
Trimmomatic version 0.39 (Bolger et al. 2014) and pooled for the species with several
individuals. Cleaned reads were assembled using TRINITY version 2.13.2 (Grabherr et al.
2011) to obtain a set of contigs of at least 200 bp for each species.

Orthologous exon call

To identify orthologous coding exon sequences in our assembled contigs, we downloaded
house mouse reference coding exons from the Ensembl database ('Ensembl genes 109' -
'Mouse genes GRCm39') and used a reciprocal best BLAST-hit strategy. First, the reference
exon sequences were blasted against our contig sequences with the blastn function of the
BLAST tool version 2.12.0 (Altschul et al. 1990), requiring >85% sequence identity, an
alignment length >=100bp and covering at least 50% of the reference exon length. The first hit
contigs were trimmed by only keeping the parts included in the BLAST alignments. Finally, a
reciprocal blast of the trimmed contigs against the reference exon set was performed, and we
only kept the contigs hitting the same exon that hit them in the first place.

Alignment and alignment filtering

For each reference exon, orthologous sequences were aligned with the OMM-MACSE
pipeline version 12.01 (Scornavacca et al. 2019), which produces a codon-aware alignment of
nucleotides using MACSE version 2.08 (Ranwez et al. 2018) and applies HmmCleaner
version 1.8 for filtering out dubious parts of the alignment (Di Franco et al. 2019). Alignments
were further filtered in order to exclude suspicious sequences. We mapped raw reads to our
exons with bwa-mem version 0.7.17-r1188 (Li 2013) and calculated the per sequence
average coverage depth and heterozygosity using samtools version 1.19.2 (Li et al. 2009) and
Angsd version 0.930 (Korneliussen et al. 2014), respectively. Sequences with an average
coverage depth below 5X or greater than the species mean plus 4 standard deviations were
excluded. Sequences with an heterozygosity greater than the species mean heterozygosity
plus 3 standard deviations were excluded, as this may be indicative of hidden paralogy
(Roycroft, Achmadi, et al. 2021). We also excluded from the alignments the sequences that
contained stop codons. After these filtering steps, a total of 71,040 exon alignments from
15,415 genes were retained.

Phylogeny

A species tree was reconstructed with IQ-TREE version 2.0.7 (Minh et al. 2020) from the
concatenation of all filtered alignments, using a GTR model and 1000 ultrafast bootstrap
replicates. The tree was rooted with the Ape R package, version 5.8 (Paradis and Schliep
2019) using *Haeromys minahassae* as the outgroup, in agreement with Roycroft, Achmadi, et
al. (2021).
We also created one tree per gene, because we need for our analyses that all exons of a
same gene have the same topology. To do that, for each gene *g*, we first identified the subset
of species for which a sequence of each exon of *g* was available, then we obtained the gene
tree by pruning the consensus phylogeny, only keeping species from this subset (Ape
"prune_tree" function).
760
761 <u>Topological discordance filtering</u>
762
Our analysis assumes that the consensus tree appropriately represents the phylogeny of the
analyzed exons. This might be untrue for some of the exons. In particular, exon alignments
containing paralogous sequences are likely to depart from this assumption. To filter them out,
for each exon alignment, we first reconstructed a *de novo* tree using IQ-TREE, and calculated
the sum of its branch lengths. Then, still using IQ-TREE, we forced the consensus topology,
re-estimated branch lengths, and again calculated their sum. We discarded exon alignments
for which the constrained total tree length (consensus topology) was 1.5 times as long, or
more, as the unconstrained one (free topology). Here we reasoned that an alignment strongly
rejecting the consensus topology should manifest itself by a large number of homoplasic
substitutions when the consensus tree is forced. Other thresholds were tried for this filter,
which did not alter the results (Supplementary Figure S4).
774
775 <u>Substitution mapping</u>
776
We realised a substitution mapping for each exon alignment using the BppSuite (Guéguen et
al. 2013). We used the BppML program version 3.0.0 to optimise gene tree branch lengths
under the YN98 codon model (Yang and Nielsen 1998). Then, we estimated the number of
synonymous and non-synonymous substitutions that have occurred in each branch at each
codon for each alignment for the four different mutational categories (WS, SW, SS, WW) in an
empirical Bayesian framework using the MapNH program version 2 (Romiguier et al. 2012)
with the parameter "map.type = Combination(reg1=DnDs, reg2=SW)".
784
785 <u>Detection of gBGC episodes</u>
786

A signature of episodic gBGC is an aggregation of WS substitutions in time and space. We
used these two criteria to define and detect gBGC episodes in specific branches and specific
exons. We calculated for each exon in each branch of the tree the total number of S-WS
substitutions, and performed permutations to determine if this number is greater than
expected in the absence of gBGC. A permutation was achieved by redistributing the total
number of S-WS substitutions of an exon across the gene tree, the probability for a
substitution to be assigned to a particular branch being proportional to the length of that
branch. This was performed 1000 times to give, for each exon, the expected distribution of the
number of S-WS substitutions in every branch of the tree in the absence of gBGC. An episode
of gBGC was called in a given exon in a given branch when the observed number of S-WS
substitutions in the branch was in the top 1% of the expected distribution.

For exons longer than 1000 bp, we added a criterion of spatial aggregation. We calculated
Moran's I, a metric already used by Galtier (2021) to detect gBGC, to measure the spatial
aggregation of S-WS substitutions across the sequence length. We generated 1000
permutations by redistributing the substitutions occurring in a specific branch along the exon
sequence. This provided us with the expected distribution of Moran's I expected under the
hypothesis of a random spatial distribution of S-WS substitutions. An episode of gBGC was
called when the observed Moran's I was in the top 1% of the expected distribution.

<u>Selection of 'PRDM9-like' episodes</u>

We developed a maximum likelihood (ML) classifier of gBGC episodes in order to distinguish
localised, short-lived episodes from large-scale and/or long-lasting effects of gBGC. Briefly,
we model our set of episodes as a mixture of three categories: punctual (episodes only hitting
one exon in one branch of the tree), two-branch (episodes hitting a single exon but affecting
downstream lineages) and all-exons (gene-wide gBGC). The model parameters were the
proportion of the three categories and the WS/SW balance at synonymous positions in the
branch carrying the episode and its two descending branches. Episodes occurring in a
terminal branch were therefore disregarded. Having fitted the model by maximising the
likelihood, the posterior probability for a given episode to belong to a particular category was
calculated in the empirical Bayesian framework. An episode-specific measure of the goodness
of fit was also calculated by comparing the likelihood of this and a degenerate model. The
details of the method are presented in the Supplementary Methods. We selected episodes for
which the posterior probability of the punctual category was higher than the other two
categories, and the goodness-of-fit p-value was above 0.2.

<u>Analysis of compensatory evolution</u>

Having called gBGC episodes based on synonymous substitutions, we focused on non-
synonymous substitutions in downstream branches to analyse the protein response. We
reasoned that, if there is compensatory evolution, one should observe a surplus of NS
substitutions in the two branches immediately downstream of the branch carrying the episode.
Permutations were performed 1000 times for each exon to redistribute the NS substitutions of
each type (WS, SW, SS, WW) across branches, blocking branches where gBGC episodes
were detected, *i.e.*, the NS substitutions in these branches were not redistributed, and the
redistributed substitutions could not be assigned to these branches. Observed values and
expected distributions were summed across all branches immediately downstream of
episodes in all exons touched by an episode. This allowed us to compare the observed
number of NS substitutions in branches after episodes with its expected distribution in the
absence of compensatory evolution, controlling that no episode is detected in these branches.
We distinguished two types of episodes : (1) those having produced at least one NS-WS
substitution, potentially deleterious, after which compensatory evolution might happen, and (2)
those with no NS-WS substitution, *i.e* after which there is *a priori* no compensation to be
expected. We searched for such compensation in the exons which have endured the gBGC
episodes and also in the other exons of the same gene. We also looked if there is
compensation during the episode in the other exons.

We then focused on the cases of potential compensation where there is exactly 1 (supposedly
deleterious) NS-WS substitution in the episode branch and exactly 1 (supposedly
compensatory) NS substitution of any type in the descending branches. We looked at the
distance (in number of codons) between these two substitutions and assessed whether the
compensatory substitutions are closer to the deleterious substitutions than would be expected
by chance. To do this, the positions of the compensatory substitutions were randomised 1000
times, by drawing an integer between 1 and the number of codons of the considered exon.
We then counted the number of cases where the observed distance between the deleterious
and the compensatory substitutions were less than or equal to 3 codons, and compared it with
the expected distribution of the same counts obtained from the randomised positions.

Data accessibility

All   the   scripts   and   analyzed   data   files   are   available   from
https://github.com/Marie-98/gBGC_comp *and* *10.5281/zenodo.14534842*

Acknowledgments

We wish to thank Khalid Belkhir, Benjamin Penaud, Vincent Ranwez and Iago Bonnici for their advices and help with bioinformatics ; Laurent Duret, Marie Sémon, Thibault Latrille, Julien Joseph, Stéphanie Bedhomme and two anonymous reviewers for insightful feedback and discussions ; and Emily Roycroft for her advice on data filtering.

Funding

This work was not supported by any research grant.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.

Ament-Velásquez SL, Gilchrist C, Rêgo A, Bendixsen DP, Brice C, Grosse-Sommer JM, Rafati N, Stelkens R. 2022. The Dynamics of Adaptation to Stress from Standing Genetic Variation and de novo Mutations. *Mol. Biol. Evol.* 39:msac242.

Auton A, Li YR, Kidd J, Oliveira K, Nadel J, Holloway JK, Hayward JJ, Cohen PE, Greally JM, Wang J, et al. 2013. Genetic Recombination Is Targeted towards Gene Promoter Regions in Dogs. *PLOS Genet.* 9:e1003984.

Axelsson E, Webster MT, Ratnakumar A, Consortium TL, Ponting CP, Lindblad-Toh K. 2012. Death of PRDM9 coincides with stabilization of the recombination landscape in the dog genome. *Genome Res.* 22:51–63.

Azevedo L, Carneiro J, van Asch B, Moleirinho A, Pereira F, Amorim A. 2009. Epistatic interactions modulate the evolution of mammalian mitochondrial respiratory complex components. *BMC Genomics*. 10:266.

Barrett RDH, Schluter D. 2008. Adaptation from standing genetic variation. *Trends Ecol. Evol.* 23:38–44.

Berglund J, Pollard KS, Webster MT. 2009. Hotspots of Biased Nucleotide Substitutions in Human Genes. *PLOS Biol.* 7:e1000026.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinforma. Oxf. Engl.* 30:2114–2120.

Bolívar P, Mugal CF, Nater A, Ellegren H. 2016. Recombination Rate Variation Modulates Gene Sequence Evolution Mainly via GC-Biased Gene Conversion, Not Hill–Robertson Interference, in an Avian System. *Mol. Biol. Evol.* 33:216–227.

Brekke TD, Papadopulos AST, Julià E, Fornas O, Fu B, Yang F, de la Fuente R, Page J, Baril T, Hayward A, et al. 2023. A New Chromosome-Assigned Mongolian Gerbil Genome Allows Characterization of Complete Centromeres and a Fully

Heterochromatic Chromosome. *Mol. Biol. Evol.* 40:msad115.

Burke MK, Liti G, Long AD. 2014. Standing Genetic Variation Drives Repeatable
Experimental Evolution in Outcrossing Populations of Saccharomyces cerevisiae.
Mol. Biol. Evol. 31:3228–3239.

Chaurasia S, Dutheil JY. 2022. The Structural Determinants of Intra-Protein Compensatory
Substitutions. *Mol. Biol. Evol.* 39:msac063.

Clément Y, Arndt PF. 2013. Meiotic recombination strongly influences GC-content evolution
in short regions in the mouse genome. *Mol. Biol. Evol.* 30:2612–2618.

Corcoran P, Gossmann TI, Barton HJ, The Great Tit HapMap Consortium, Slate J, Zeng K.
2017. Determinants of the Efficacy of Natural Selection on Coding and Noncoding
Variability in Two Passerine Species. *Genome Biol. Evol.* 9:2987–3007.

Davis BH, Poon AFY, Whitlock MC. 2009. Compensatory mutations are repeatable and
clustered within proteins. *Proc. R. Soc. B Biol. Sci.* 276:1823–1827.

Di Franco A, Poujol R, Baurain D, Philippe H. 2019. Evaluating the usefulness of alignment
filtering methods to reduce the impact of errors on evolutionary inferences. *BMC*
Evol. Biol. 19:21.

Duret L, Arndt PF. 2008. The Impact of Recombination on Nucleotide Substitutions in the
Human Genome. *PLOS Genet.* 4:e1000071.

Duret L, Galtier N. 2009. Biased Gene Conversion and the Evolution of Mammalian
Genomic Landscapes. *Annu. Rev. Genomics Hum. Genet.* 10:285–311.

Estandía A, Sendell-Price AT, Robertson BC, Clegg SM, unpublished data,
https://www.biorxiv.org/content/10.1101/2023.06.29.546893v2.full.pdf, last accessed
April 2, 2025.

Eyre-Walker A. 1999. Evidence of Selection on Silent Site Base Composition in Mammals:
Potential Implications for the Evolution of Isochores and Junk DNA. *Genetics*.
152:675–683.

Filteau M, Hamel V, Pouliot M, Gagnon-Arsenault I, Dubé AK, Landry CR. 2015.
Evolutionary rescue by compensatory mutations is constrained by genomic and
environmental backgrounds. *Mol. Syst. Biol.* 11:832.

Galtier N. 2021. Fine-scale quantification of GC-biased gene conversion intensity in
mammals. *Peer Community J.* 1.

Galtier N, Duret L. 2007. Adaptation or biased gene conversion? Extending the null
hypothesis of molecular evolution. *Trends Genet.* 23:273–277.

Galtier N, Duret L, Glémin S, Ranwez V. 2009. GC-biased gene conversion promotes the
fixation of deleterious amino acid changes in primates. *Trends Genet.* 25:1–5.

Galtier N, Piganeau G, Mouchiroud D, Duret L. 2001. GC-Content Evolution in Mammalian
Genomes: The Biased Gene Conversion Hypothesis. *Genetics*. 159:907–911.

Galtier N, Roux C, Rousselle M, Romiguier J, Figuet E, Glémin S, Bierne N, Duret L. 2018.
Codon Usage Bias in Animals: Disentangling the Effects of Natural Selection,

938   Effective Population Size, and GC-Biased Gene Conversion. *Mol. Biol. Evol.*
939   35:1092–1103.

940 Garud NR, Messer PW, Buzbas EO, Petrov DA. 2015. Recent Selective Sweeps in North
941   American Drosophila melanogaster Show Signatures of Soft Sweeps. *PLOS Genet.*
942   11:e1005004.

943 Glémin S. 2010. Surprising Fitness Consequences of GC-Biased Gene Conversion: I.
944   Mutation Load and Inbreeding Depression. *Genetics*. 185:939–959.

945 Glémin S, Arndt PF, Messer PW, Petrov D, Galtier N, Duret L. 2015. Quantification of GC-
946   biased gene conversion in the human genome. *Genome Res.* 25:1215–1228.

947 Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
948   Raychowdhury R, Zeng Q, et al. 2011. Trinity: reconstructing a full-length
949   transcriptome without a genome from RNA-Seq data. *Nat. Biotechnol.* 29:644–652.

950 Guéguen L, Gaillard S, Boussau B, Gouy M, Groussin M, Rochette NC, Bigot T, Fournier D,
951   Pouyet F, Cahais V, et al. 2013. Bio++: Efficient Extensible Libraries and Tools for
952   Computational Molecular Evolution. *Mol. Biol. Evol.* 30:1745–1750**.**

953 Halldorsson BV, Hardarson MT, Kehr B, Styrkarsdottir U, Gylfason A, Thorleifsson G, Zink F,
954   Jonasdottir Adalbjorg, Jonasdottir Aslaug, Sulem P, et al. 2016. The rate of meiotic
955   gene conversion varies by sex and age. *Nat. Genet.* 48:1377–1384.

956 Halldorsson BV, Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP,
957   Gunnarsson B, Oddsson A, Halldorsson GH, Zink F, et al. 2019. Characterizing
958   mutagenic effects of recombination through a sequence-level genetic map. *Science*.
959   363:eaau1043.

960 Hämälä T, Tiffin P. 2020. Biased Gene Conversion Constrains Adaptation in Arabidopsis
961   thaliana. *Genetics* [Internet] 215:831–846. Available from:
962   https://doi.org/10.1534/genetics.120.303335

963 Harris RB, Sackman A, Jensen JD. 2018. On the unfounded enthusiasm for soft selective
964   sweeps II: Examining recent evidence from humans, flies, and viruses. *PLOS Genet.*
965   14:e1007859

966 Illingworth CJR, Parts L, Schiffels S, Liti G, Mustonen V. 2012. Quantifying Selection Acting
967   on a Complex Trait Using Allele Frequency Time Series Data. *Mol. Biol. Evol.*
968   29:1187–1197.

969 Ivankov DN, Finkelstein AV, Kondrashov FA. 2014. A structural perspective of compensatory
970   evolution. *Curr. Opin. Struct. Biol.* 26:104–112.

971 Joseph J. 2024. Increased Positive Selection in Highly Recombining Genes Does not
972   Necessarily Reflect an Evolutionary Advantage of Recombination. *Mol. Biol. Evol.*
973   41:msae107**.**

974 Joseph J, Prentout D, Laverré A, Tricou T, Duret L. 2024. High prevalence of PRDM9-
975   independent recombination hotspots in placental mammals. *Proc. Natl. Acad. Sci.*
976   121:e2401973121.

977 Kawakami T, Mugal CF, Suh A, Nater A, Burri R, Smeds L, Ellegren H. 2017. Whole-
978   genome patterns of linkage disequilibrium across flycatcher populations clarify the

causes and consequences of fine-scale recombination rate variation in birds. *Mol. Ecol.* 26:4158–4172.

Kern AD, Kondrashov FA. 2004. Mechanisms and convergence of compensatory evolution in mammalian mitochondrial tRNAs. *Nat. Genet.* 36:1207–1212.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics*. 15:356.

Lachance J, Tishkoff SA. 2014. Biased Gene Conversion Skews Allele Frequencies in Human Populations, Increasing the Disease Burden of Recessive Alleles. *Am. J. Hum. Genet.* 95:408–420.

Lartillot N. 2013. Phylogenetic Patterns of GC-Biased Gene Conversion in Placental Mammals and the Evolutionary Dynamics of Recombination Landscapes. *Mol. Biol. Evol.* 30:489–502.

Latrille T, Duret L, Lartillot N. 2017. The Red Queen model of recombination hot-spot evolution: a theoretical investigation. *Philos. Trans. R. Soc. B Biol. Sci.* 372:20160463.

Latrille T, Joseph J, Hartasánchez DA, Salamin N. 2024. Estimating the proportion of beneficial mutations that are not adaptive in mammals. *PLoS genetics.* 20(12): e1011536.

Lesecque Y, Mouchiroud D, Duret L. 2013. GC-Biased Gene Conversion in Yeast Is Specifically Associated with Crossovers: Molecular Mechanisms and Evolutionary Significance. *Mol. Biol. Evol.* 30:1409–1419.

Li H, unpublished data, https://arxiv.org/abs/1303.3997, last accessed, April 2, 2025.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* 25:2078–2079.

Li R, Bitoun E, Altemose N, Davies RW, Davies B, Myers SR. 2019. A high-resolution map of non-crossover events reveals impacts of genetic diversity on mammalian meiotic recombination. *Nat. Commun.* 10:3900.

Mancera E, Bourgon R, Brozzi A, Huber W, Steinmetz LM. 2008. High-resolution mapping of meiotic crossovers and non-crossovers in yeast. *Nature*. 454:479–485.

McCandlish DM, Shah P, Plotkin JB. 2016. Epistasis and the Dynamics of Reversion in Molecular Evolution. *Genetics*. 203:1335–1351.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the Adh locus in Drosophila. *Nature*. 351:652–654.

McManus KF, Taravella AM, Henn BM, Bustamante CD, Sikora M, Cornejo OE. 2017. Population genetic analysis of the DARC locus (Duffy) reveals adaptation from standing variation associated with malaria resistance in humans. *PLOS Genet.* 13:e1006560**.**

Meunier J, Duret L. 2004. Recombination Drives the Evolution of GC-Content in the Human Genome. *Mol. Biol. Evol.* 21:984–990.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. *Mol. Biol. Evol.* 37:1530–1534.

Mugal CF, Weber CC, Ellegren H. 2015. GC-biased gene conversion links the recombination
landscape and demography to genomic base composition. *BioEssays*. 37:1317–
1326.

Necşulea A, Popa A, Cooper DN, Stenson PD, Mouchiroud D, Gautier C, Duret L. 2011.
Meiotic recombination favors the spreading of deleterious mutations in human
populations. *Hum. Mutat.* 32:198–206.

Neverov AD, Popova AV, Fedonin GG, Cheremukhin EA, Klink GV, Bazykin GA. 2021.
Episodic evolution of coadapted sets of amino acid sites in mitochondrial proteins.
PLOS Genet. 17:e1008711.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*. 35:526–528.

Pessia E, Popa A, Mousset S, Rezvoy C, Duret L, Marais GAB. 2012. Evidence for
Widespread GC-biased Gene Conversion in Eukaryotes. *Genome Biol. Evol.* 4:675–
682.

Poon A, Chao L. 2005. The Rate of Compensatory Mutation in the DNA Bacteriophage
φX174. *Genetics*. 170:989–999.

Poon A, Davis BH, Chao L. 2005. The Coupon Collector and the Suppressor Mutation:
Estimating the Number of Compensatory Mutations by Maximum Likelihood.
Genetics. 170:1323–1332.

Pracana R, Hargreaves AD, Mulley JF. 2020. Runaway GC Evolution in Gerbil Genomes.
Mol. Biol. Evol. 37:2197–2210.

Rannala B, Leache A, Edwards S, Yang Z. 2020. The multispecies coalescent model and
species tree inference. In: Scornavacca C, Delsuc F, Galtier N, editors.
Phylogenetics in the Genomic Era. Self Published. p. 3.3:1-3.3:21.

Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. 2018. MACSE v2: Toolkit for the
Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Mol.*
Biol. Evol. 35:2582–2584.

Ratnakumar A, Mousset S, Glémin S, Berglund J, Galtier N, Duret L, Webster MT. 2010.
Detecting positive selection within genomes: the problem of biased gene conversion.
Philos. Trans. R. Soc. B Biol. Sci. 365:2571–2580.

Raynaud M, Sanna P, Joseph J, Clément J, Imai Y, Lareyre J-J, Laurent A, Galtier N,
Baudat F, Duret L, et al. 2025. PRDM9 drives the location and rapid evolution of
recombination hotspots in salmonid fish. *PLOS Biol.* 23:e3002950.

Rojas Echenique JI, Kryazhimskiy S, Nguyen Ba AN, Desai MM. 2019. Modular epistasis
and the compensatory evolution of gene deletion mutants.Butler G, editor. *PLOS*
Genet. 15:e1007958.

Romiguier J, Figuet E, Galtier N, Douzery EJP, Boussau B, Dutheil JY, Ranwez V. 2012.
Fast and Robust Characterization of Time-Heterogeneous Sequence Evolutionary

Processes Using Substitution Mapping. *PLOS ONE*. 7:e33852.

Romiguier J, Ranwez V, Douzery EJP, Galtier N. 2010. Contrasting GC-content dynamics
across 33 mammalian genomes: Relationship with life-history traits and chromosome
sizes. *Genome Res.* 20:1001–1009.

Rousselle M, Laverré A, Figuet E, Nabholz B, Galtier N. 2019. Influence of Recombination
and GC-biased Gene Conversion on the Adaptive and Nonadaptive Substitution Rate
in Mammals versus Birds. *Mol. Biol. Evol.* 36:458–471.

Rousselle M, Simion P, Tilak M-K, Figuet E, Nabholz B, Galtier N. 2020. Is adaptation limited
by mutation? A timescale-dependent effect of genetic diversity on the adaptive
substitution rate in animals. *PLOS Genet.* 16:e1008668.

Roycroft E, Achmadi A, Callahan CM, Esselstyn JA, Good JM, Moussalli A, Rowe KC. 2021.
Molecular Evolution of Ecological Specialisation: Genomic Insights from the
Diversification of Murine Rodents. *Genome Biol. Evol.* 13:evab103.

Roycroft E, MacDonald AJ, Moritz C, Moussalli A, Portela Miguez R, Rowe KC. 2021.
Museum genomics reveals the rapid decline and extinction of Australian rodents
since European settlement. *Proc. Natl. Acad. Sci.* 118:e2021390118.

Samaniego Castruita JA, Westbury MV, Lorenzen ED. 2020. Analyses of key genes involved
in Arctic adaptation in polar bears suggest selection on both standing variation and
de novo mutations played an important role. *BMC Genomics*. 21:543.

Scornavacca C, Belkhir K, Lopez J, Dernat R, Delsuc F, Douzery EJP, Ranwez V. 2019.
OrthoMaM v10: Scaling-Up Orthologous Coding Sequence and Exon Alignments
with More than One Hundred Mammalian Genomes. *Mol. Biol. Evol.* 36:861–862.

Serrano C, Teixeira CSS, Cooper DN, Carneiro J, Lopes-Marques M, Stenson PD, Amorim
A, Prata MJ, Sousa SF, Azevedo L. 2021. Compensatory epistasis explored by
molecular dynamics simulations. *Hum. Genet.* 140:1329–1342.

Smagulova F, Brick K, Pu Y, Camerini-Otero RD, Petukhova GV. 2016. The evolutionary
turnover of recombination hot spots contributes to speciation in mice. *Genes Dev.*
30:266–280.

Smeds L, Mugal CF, Qvarnström A, Ellegren H. 2016. High-Resolution Mapping of
Crossover and Non-crossover Recombination Events by Whole-Genome Re-
sequencing of an Avian Pedigree. *PLOS Genet.* 12:e1006044.

Stapley J, Feulner PGD, Johnston SE, Santure AW, Smadja CM. 2017. Variation in
recombination frequency and distribution across eukaryotes: patterns and processes.
Philos. Trans. R. Soc. B Biol. Sci. 372:20160455.

Starr TN, Thornton JW. 2016. Epistasis in protein evolution. *Protein Sci.* 25:1204–1218.

Stolyarova AV, Nabieva E, Ptushenko VV, Favorov AV, Popova AV, Neverov AD, Bazykin
GA. 2020. Senescence and entrenchment in evolution of amino acid sites. *Nat.*
Commun. 11:4603.

Storz JF. 2018. Compensatory mutations and epistasis for protein function. *Curr. Opin.*
Struct. Biol. 50:18–25.

Wallberg A, Glémin S, Webster MT. 2015. Extreme Recombination Frequencies Shape
Genome Variation and Evolution in the Honeybee, Apis mellifera. *PLOS Genet.*
11:e1005189.

Williams AL, Genovese G, Dyer T, Altemose N, Truax K, Jun G, Patterson N, Myers SR,
Curran JE, Duggirala R, et al. 2015. Non-crossover gene conversions show strong
GC bias and unexpected clustering in humans.de Massy B, editor. *eLife*. 4:e04637.

Yang Z, Nielsen R. 1998. Synonymous and nonsynonymous rate variation in nuclear genes
of mammals. *J. Mol. Evol.* 46:409–418.