1   **Experimental validation of genome-environment associations in Arabidopsis**
2
3

4   Yuxin Luo[1,2], Claire M. Lorts[1], Erica H. Lawrence-Paul[1], Jesse R. Lasky[1]
5
6   [1]Department of Biology, Pennsylvania State University
7
8   [2]Correspondence

Abstract

Identifying the genetic basis of local adaptation is a key goal in evolutionary biology. Allele frequency clines along environmental gradients, known as genotype-environment associations (GEA), are often used to detect potential loci causing local adaptation but are rarely followed by experimental validation. Here, we tested loci identified in three moisture-related GEA studies on *Arabidopsis*. We studied 42 GEA-identified genes using t-DNA knockout lines under drought and tested effects on flowering time, an adaptive trait, and genotype-by-environment (GxE) interactions on performance and fitness. In total, 16/42 genes had significant effects on traits involved in local adaptation or performance response to environment. We found that *wrky38* mutants had significant GxE effects for fitness; *lsd1* plants had a significant GxE effect for flowering time, while 11 genes showed flowering time effects with no drought interaction. However, most GEA candidates did not exhibit GxE. In the follow-up experiments, *wrky38* caused decreased stomatal conductance and specific leaf area under drought, indicating potentially adaptive drought avoidance. Additionally, GEA identified natural putative LoF variants of *WRKY38* associated with dry environments, as well as alleles associated with variation in *LSD1* expression. While only a few GEA-identified genes were validated for GxE interactions for fitness under drought, we likely overlooked some because experiments might not well represent natural environments and t-DNA insertions might not well represent natural alleles. Nevertheless, GEAs apparently identified some genes contributing to local adaptation. GEA and follow-up experiments are straightforward to implement in model systems and demonstrate prospects for GEA discovery of new local adaptations.

Keywords: Genome-environment associations (GEA); Drought stress; Local adaptation; t-DNA knockout; *WRKY38*; *LSD1*.

INTRODUCTION

A substantial portion of genetic variation within species in ecologically important traits is likely due to local adaptation to the environment (Savolainen et al., 2013; Tigano and Friesen, 2016; Wadgymar et al., 2022). Local adaptation is defined by a genotype-by-environment (GxE) interaction where local genotypes have higher fitness than foreign genotypes (Kawecki & Ebert, 2004). Identifying locally adaptive alleles and traits is of great interest in plant and animal breeding and conservation, where these alleles can be deployed for genetic improvement and prediction (Cortés et al., 2022; Crossa et al., 2017; Lasky et al., 2023; Whiting et al., 2024). Central questions of local adaptation include the effect size and number of involved mutations (Yeaman and Whitlock, 2011), whether locally adapted mutations have tradeoffs among different environments (Lee et al., 2024), whether the same or different mutations are involved in local adaptation to the same environments in different populations (Ralph and Coop, 2015), and which environmental gradients drive locally adaptive genetic variation (Lasky et al., 2012).

Conducting multiple common garden experiments to map populations and genetic loci contributing to fitness tradeoffs has been a gold standard approach (Clausen et al., 1941). However, such experiments are logistically challenging or impossible in many systems. Alternative approaches using geographic patterns in allele frequency have emerged for identifying loci potentially underlying local adaptation. Loci showing substantial allele frequency differences between populations, i.e. certain environment variables, may indicate those driving local adaptation (Coop et al., 2010; Endler, 1973). With the increasing availability of high-throughput genomic data, Genome-Environment Association (GEA) studies have been widely employed to identify putative locally adapted loci (Lasky et al., 2023; Rellstab et al., 2015).

GEAs employ diverse methods to identify loci where allele frequency is closely associated with environmental conditions (Lasky et al., 2023; Rellstab et al., 2015). Methods differ depending on whether they are single-locus or multi-loci (e.g. genome-wide) models (Betancourt et al., 2021; Gehan et al., 2015; Hager et al., 2022; Hancock et al., 2011; Lasky et al., 2012; Lee et al., 2024), where the latter may capture groups of loci covarying with the environment that are potentially relevant for polygenic adaptation (Forester et al., 2018). Additionally, there are methods that control for genome-wide patterns of similarity among populations using random effects (Zoubarev et al., 2012). While these approaches may reduce spurious associations due to population structure-environmental covariation, they can also be too conservative (Alonso-Blanco et al., 2016). Thirdly, some approaches synthesize the evidence from GEA with additional evidence, such as multiple common garden experiments (Capblancq et al., 2023; Lasky et al., 2018; Soudi et al., 2023). While these methods are now standard approaches in ecological genomics, the findings are rarely followed by experimental validation of variation at individual loci (Monroe et al., 2018; Nyine et al., 2025; Tergemina et al., 2025).

To experimentally test the hypothesis that variation at a given locus causes GxE for fitness, approaches that isolate experimental genetic variation to that locus are the most powerful. Creating near-isogenic lines (NILs) segregating at a given locus is ideal for testing GxE effects in a natural genetic background, but it is a time-consuming process. Gene knockout mutants in model species like *Arabidopsis* provide an alternative tool for testing GxE effects of variation at a particular gene. For example, various Arabidopsis knockout lines carrying t-DNA

insertions within genes that disrupt their function provide valuable resources for studying the
effects of individual genes on fitness and other ecologically important traits. Although these
mutants may not fully represent natural allelic variation for local adaptation, using them may be
a good option for bulk experimental screening of candidate loci (Chong and Stinchcombe, 2019;
Monroe et al., 2018). Resequencing of large numbers of natural *Arabidopsis* genotypes has
yielded more detailed information on functional variation (Alonso-Blanco et al., 2016), which can
be linked to loci of interest to make inferences about functional variants generating strong
signals in GEA studies.
Arabidopsis thaliana is a small annual plant native to a wide range across Eurasia and
Africa with diverse climates (Yim et al., 2024). Many studies have demonstrated potential
drought adaptation of *A. thaliana* in quantitative traits (Dittberner et al., 2018), SNPs or genes
(El-Soda et al., 2015; Exposito-Alonso et al., 2018; Hancock et al., 2011), and gene expression
(Lasky et al., 2014). *A. thaliana* employs diverse strategies for drought adaptation such as
drought escape via phenology, drought avoidance via traits reducing water loss, and drought
tolerance via enhancing water acquisition or water-use efficiency (Lovell et al., 2013).
Here we focus on studying loci identified based on 3 different approaches, with the goal
of testing whether identified genes indeed cause genotype-drought effects on fitness or other
aspects of performance. We are also interested in whether the genes identified as drought-
adapted in our experiments have allele frequency clines across the worldwide distribution of *A.*
thaliana. For mutants identified as showing interesting genotype-environment interactions in our
initial drought screen experiment, we followed up by testing their response to an additional,
different drought treatment.
We tested the following hypotheses:
1. Some genes identified through GEA as potentially drought adapted exhibit natural
variation in putative gene function assessed from resequencing data. Such genes would
be especially amenable to using t-DNA knockout lines for testing the effects of natural
variation. Mutations in genes without natural loss-of-function variation can still be
assessed using t-DNA lines, though their relevance to natural variants is less clear.
2. Knockout mutants of these genes display significant genotype-drought interactions
affecting traits (e.g. flowering time) known to be under changing selection in drought,
fitness or performance traits in controlled drought experiments; and
3. These genes show climate-associated allelic variation and genomic signatures of
selection, patterns consistent with adaptation to dry climates.
In our main drought screen, we found that *wrky38* had significant GxE effects on several
fitness-related traits, and *lsd1* had a significant GxE effect on flowering time. Both *WRKY38* and
LSD1 were top candidates associated with moisture in their originated studies (Lasky et al.,
2018, 2012). Additionally, *WRKY38* was also found most strongly associated with winter low
temperatures in the combined common garden study (Lasky et al., 2018), while knockouts of
LSD1 were found to affect chilling sensitivity (Huang et al., 2010) and allelic variation was
associated with the second principal component of climate (moisture related, Lee et al., 2017) in
Arabidopsis. Both genes happen to play a role in the salicylic acid (SA) pathway (Bernacki et al.,
2019; Kim et al., 2008; Szechyńska-Hebda et al., 2016; Wituszynska et al., 2013), primarily
known for microbial defense response but which can be activated by abiotic stress (reviewed in
Miura and Tada, 2014; Wu et al., 2019). Incorporating previous studies and our results, we

decided to use the two mutants to follow up with intermediate drought and freezing experiments,
in order to test their possible adaptation in different drought and cold regimes in terms of their
ecophysiological responses.

METHODS

Selecting genes from published genome-environment
associations

We selected genes to test using five SNP lists from three published GEA studies that employed
three different approaches. The first approach was based on mixed models of genome-wide
associations with climate, where we included random effects to account for genomic similarity
(Kang et al., 2008). For these mixed models, we calculated SNP associations with intra-annual
monthly precipitation variability, growing season precipitation variability, or inter-annual growing
season precipitation variability (i.e. 3 separate environmental GWAS, Lasky et al., 2014). We
took the SNPs from all three climate associations and combined them into one list ranked by *p*-
value for the mixed model association test. The second approach was a multivariate ordination
where we identified SNPs most strongly associated with the first axis, which was associated
with seasonality of temperature and precipitation, or the first axis after partialing out spatial
variables, which was strongly associated with summer moisture (Lasky et al., 2012). We took
SNPs with the strongest absolute loadings on each of these axes. The third approach was
based on integrated information from four common gardens across Europe and also the broader
ecotype panel to identify SNPs that showed the strongest SNP-climate associations and GxE
for fitness favoring home climate alleles in monthly growing season precipitation variability or
aridity index (i.e. annual precipitation/PET; Lasky et al., 2018). In total, we generated 1, 2, and 2
lists of SNPs from the three studies respectively.

From the top SNPs in each list, we only kept SNPs at least 250 kb apart. Starting from
the SNP with the lowest *p*-values, we took up to 3 closest genes within 5kb of the SNP until we
had at least 15 genes for each of the four lists of SNPs, or in the case of the combined climate
association study, until we had at least 25 genes.

This procedure resulted in a list of 90 candidate genes. We chose 44 SALK lines
representing knockouts of 42 genes, including 42 lines with insertions within exons and 2 lines
with insertions within introns and confirmed homozygosity using tDNA Express: Arabidopsis
Gene Mapping Tool (Alonso et al., 2003; Table S1). Among the 44 SALK lines targeting 42
genes, three lines (SALK_048316C, SALK_085852C, and SALK_063969) targeting AT5G65080
(*MAF5*) and AT3G45860 (*CRK4*), although not top candidates, were included in the experiment
because their potential functional significance in drought adaptation (Kim and Sung, 2010;
Ratcliffe et al., 2003; Zhao et al., 2023). As a result, 40 top-candidate genes were validated in
this study.

Functional variation in the screened loci in the 1001 Genomes

We analyzed putative functional variation (hereafter FV) in the 1001 Genomes resequencing data for all genes screened. We used the 1001 Genomes Polymorphism Browser (https://tools.1001genomes.org/polymorph/) to identify potential large-effect mutations in each gene, which were considered as putative loss-of-function (LoF) mutations. If more than one isoform existed for a gene, we recorded the case of the most common isoform, except when no functional variant was found for this isoform, we instead used the most common isoform that had functional variants. We summarized the counts of each type of functional variation and the positions where the variations occurred.

Main drought screen

Detailed experimental settings are provided in Supplementary Methods. Briefly, we programmed the chamber temperature and photoperiod based on that of the *Lip-0* ecotype from southern Poland following Lorts and Lasky (2020) (Table S2). Each tray represented either a drought or well-watered treatment, and seven replicates per mutant were randomly assigned positions for each treatment. Seeds were stratified in DI water for 4-5 days prior to planting.

Pots were maintained at 2.5-5 cm of bottom water before drought treatments, which began 19 days post-planting. Bottom water was removed from drought trays, while well-watered trays retained a constant 2.5 cm of water and received 2.5 ml of top water every other day. After five weeks, all pots were top-watered with a 12 ml miracle grow solution every other day to ensure nutrition.

At maturity, fecundity was assessed when siliques dried and rosette leaves senesced. We recorded inflorescence height and secondary branch numbers were recorded. For reproductive fitness estimates, we measured total silique number and silique length for 6 siliques to calculate average silique length, then calculated the total silique length as the product of silique number and average silique length.

After measurements, aboveground dry biomass was obtained, and rosette tissue was ground for $\delta^{13}C$ and $\delta^{15}N$ isotope analysis. $\delta^{13}C$ represents estimate for water use efficiency (WUE).

Follow-up drought and freezing experiments

Intermediate drought experiment on *wrky38* and *lsd1*

For intermediate drought (ID), plants were grown at 20/14°C (12h/12h) with varying watering frequencies (every 3 days for well-watered, every 6 days for drought). We tracked stomatal conductance (g_{sw}) and Fv/Fm for 10 for four plants per genotype per treatment. Specific leaf area (SLA), relative water content (RWC), and leaf isotope analyses ($\delta^{13}C$, C:N and $\delta^{15}N$) were analyzed using five to six 40-day-old rosettes (see Supplementary Methods for details).

We recorded the flowering time for six replicates per genotype per treatment. As in the main drought screen, we measured the same growth and fitness traits and isotopes.

Overnight freezing experiment on *wrky38* and *lsd1*

Twelve replicates of *Col*, *wrky38*, and *lsd1* were grown at 20/14°C (12h/12h) until Day 30, then acclimated to 10/4℃ for 2 weeks before freezing treatment at 10/-2℃. We surrounded all pots with extra soil for insulation to prevent unnaturally soil freezing. On day 60, we measured diameter as an estimate of growth under freezing. After temperature was restored on Day 64, flowering time, biomass, silique traits, and inflorescence length were recorded as described above.

Statistical analyses

In both the drought experiments, we were primarily interested in whether the knockouts had genotype-specific drought responses, i.e., whether there were genotype-by-drought interactions, as well as whether the knockouts altered average phenotypes under drought, i.e., whether genotype effects were significant. Thus, we applied linear mixed models (LMMs) to test traits we measured between *Col* and each mutant separately, considering genotypes as fixed effects and tray numbers as random effects. To account for false positives arising from multiple LMMs within the dataset, we calculated adjusted *p*-values using a False Discovery Rate (FDR) threshold of 0.05, applying the *p.adjust()* function in R. We summarized how genes, originally identified in previous studies, were selected for knockout experiments and examined whether they exhibited significant genotype effects. Additionally, we calculated the ratio of genes tested with significant genotype effects relative to the total number of genes selected for screening in each original study. We also applied LMMs on the daily g_{sw} and Fv/Fm data measured in the ID experiment to detect if the mutants differed from *Col* in ecophysiology on any of the 10 measuring days.

It was not feasible to control moisture in the freezing experiment as we did in the drought follow-up because soil lost water much more slowly than it did in higher temperatures, so we used *DunnettTest()* function in the R package *DescTools* (Signorell et al., 2023) to conduct Dunnett's Test on traits measured in the freezing experiment, using *Col* as control and test for differences of the two mutants to control separately.

Population genetic variation at *WRKY38*

Because we found some putative LoF for *WRKY38* were relatively common in the 1001 Genomes accessions, we further analysed the geographic and genetic structure of these variants. We used the imputed SNP matrix (hdf5 file) and short indels (vcf file) downloaded from the 1001 Genome Project website (https://1001genomes.org/accessions.html) for the following analyses.

We calculated Tajima's D across chromosome 5 (Chr 5) in 5-kb windows using VCFtools (--TajimaD 5000) to test possible selection signals on *WRKY38* in natural accessions. The most common functional variant for *WRKY38* is a frameshift at position 7495793 on Chr 5 (Frameshift$_{7495793}$; Table S3). To detect linkage disequilibrium and potential haplotype structure (i.e. non-random associations among SNPs within the genomic region) between

Frameshift$_{7495793}$ and adjacent SNPs, we calculated the square of the Spearman's correlation
coefficient (r^2). This calculation was performed between the presence/absence of
Frameshift$_{7495793}$ and the alleles of each SNP within 5 kb upstream and downstream of *WRKY38*
using the *cor()* function in base R.
To characterize the geographic and population structure of natural putative LoF
mutations in 1001 Genomes accessions, we plotted mutations and their genetic clusters by
different shapes and outline colors on a map. We also constructed neighbor-joining (NJ) trees
built with the entire gene sequence of *WRKY38* (including coding and non-coding sequences),
and a genome-wide NJ tree. We referred to the population structure of 1001 Genomes
accessions in Alonso-Blanco et al. (2016), where the authors classified 9 genetic clusters and
one admixed group. For simplicity of data presentation, we combined the 9 groups by 5 major
regions — West Mediterranean (W-Med), North Mediterranean (N-Med), Central and Western
Europe (CW-Euro), North Europe (N-Euro), and Asia. Accessions that were introduced,
classified in the admixed group, or might be potential contaminations (Pisupati et al., 2017) were
not included in the 5 groups.
We used the R package *maps* to visualize the distribution of the common putative LoF
mutations of *WRKY38*. To build the genome-wide tree, we annotated the vcf file using SnpEff
(Cingolani et al., 2012), extracted the synonymous sites, filtered out sites with minor allele
frequency below 0.01 (--maf 0.01) and missing rate over 10% (--max-missing 0.9), and
generated a distance matrix using PLINK (Purcell et al., 2007). Both NJ trees were constructed
using the *nj()* function from package *ape* (Paradis and Schliep, 2019).
We hypothesized that the natural *WRKY38* putative LoF variants could underlie local
adaptation. To identify which climate variables might best explain *WRKY38* functional variation
and be most closely tied to mechanisms of selection, we used LMMs to scan for correlations
across the 19 bioclimatic variables. Worldclim 2 bioclimatic data were retrieved using the
getData() function from the R package raster (Fick and Hijmans, 2017; Hijmans et al., 2015).
Models were implemented using *coxme* and *kinship2* packages to include kinship in the models
as random effects (Sinnwell et al., 2014; Therneau and Therneau, 2015). We focused on testing
the local adaptation of intact *WRKY38*, Frameshift$_{7495793}$, the most common and widespread FV,
and Frameshift$_{7495793\&94}$, which accounted for a 3-bp frameshift that led back to the normal open
reading frame (ORF) thus suspected to be potentially functional. Since mixed models have low
power when causal variants are correlated with the genomic background, we also used Welch's
t-test (*t.test()*) in R to compare the differences in bioclimatic variables for accessions with/out the
intact gene or the FVs.

Variation in *LSD1* expression

Only 5 of the total 1135 accessions had *LSD1* putative LoF mutations. Thus, we suspected that
LoF may not be the major cause of the strong association between climate and the *LSD1*-
targeting SNP in Lasky et al. (2018), and the adaptation might exist due to amino acid changes
or *cis*-regulatory variation impacting expression. Therefore, we scanned across 19 bioclimatic
variables and SNPs (ref/alt alleles) within 10kb around *LSD1* using univariate GEAs with
GEMMA (Zoubarev et al., 2012) for 933 native accessions with known coordinates that were not
potential contaminants (Pisupati et al., 2017). We found elevated correlations with climate at

SNPs 1kb upstream of *LSD1* and the start of the gene sequence, suggesting that the adaptation
may occur at the promoter region. These SNPs may influence gene expression by altering
transcription factor binding motifs (Shastry, 2009; Wang et al., 2005).
Thus, we further asked if these SNPs impact *LSD1* expression using the published 1001
Genomes transcriptome data (http://signal.salk.edu/1001.php; Kawakatsu et al., 2016). Among
728 accessions with published transcriptome data, 1 accession was excluded from the following
analysis according to Pisupati et al. (2017). We compared the expression levels of *LSD1*
between the two alleles of the top climate-associated SNPs in the putative promoter region by
Welch's t-test (*t.test()*) in R. We also used linear models (*lm()*) to find possible associations
between *LSD1* expression and bioclimatic variables. Additionally, we investigated whether
SNPs with elevated correlations to climate are enriched in bioclimatic variables associated with
LSD1 expression.

290 RESULTS

291 Widespread putative functional variation in the experimentally
292 screened genes in the 1001 Genomes accessions

Across all 1135 accessions, 1070 exhibit at least one high-impact functional variant (i.e. putative
LoF) over the 44 experimentally screened genes. Specifically, 1973 putative frameshifts
occurred in 943 genotypes as well as 1116 stop-gained FVs in 842 genotypes, appearing to be
the most prevalent types of putative high-impact variants over the identified genes (Table S3).
Splice donor variants were found in 303 accessions, and splice-acceptor, stop-lost, and start-
lost variants were found in 29, 5, and 4 accessions, respectively.
Putative frameshifts were most common in AT1G19410 (*FBD*), occurring in 639
accessions. Frameshift variants were also common in AT5G22570 (*WRKY38*), detected in 250
accessions. Stop-gained had the most counts in AT5G22900 (*CHX3*), with 492 accessions
having the putative FVs. Splice-donor was also the most prevalent in the *FBD* gene, exhibited
by 220 accessions. The rest of the high-impact effects were rare among the screened loci,
suggesting purifying selection on amino acid sequences. We found 14 accessions with splice-
acceptor variants, 2 accessions with stop-lost, and 2 accessions with start-lost at AT5G10960
(*CAF1I*), AT3G45860 (*CRK4*), and AT5G22570 (*WRKY38*), respectively.
Some mutations in the screened genes had potential dual effects, and such dual effects
were locusspecific. A 1-bp deletion or insertion at the same position in AT1G76090 (*SMT3*)
caused both frameshift and start-loss, resulting in the most common dual effects observed in 95
accessions. Two insertions in AT3G45860 (*CRK4*) could each lead to dual effects of frameshift
and stop-gain, together occurring in 21 accessions. In rare cases, a 1-bp deletion in
AT5G40830 (*ICA*) caused frameshift and stop-lost in 4 accessions, while a 3-bp deletion in
AT5G65050 (*AGL31*/*MAF2*) caused stop-lost and disruptive-inframe-deletion effects in 1
accession.

Prevalent treatment effects, variable genotype effects, and few GxE effects on fitness and phenology in the main drought screen

315
316

Drought treatment effects on phenotypes were strong and largely consistent across different t-
DNA insertion lines (Figure 1). Specifically, the drought effects were significant with FDR=0.05
across all 44 tested lines, causing lower aboveground biomass, smaller inflorescence length
and weight, fewer siliques, shorter average and total silique length, and delayed flowering, with
line CS68738 (*lsd1*) being the only exception with no significant drought effect on flowering time
(Figure 1). The drought effect on rosette weight was also common, significantly reducing rosette
weights in 25 of 44 lines at FDR = 0.05.

Genotype effects varied across the 44 lines tested (Figure 1, Table S4-6). Genotype
effects were most common for flowering time, with 20 having nominally significant effects (α =
0.05) on flowering, primarily accelerating flowering, except for *wrky38*, which exhibited delayed
flowering under drought. Thirteen insertion lines were significant with FDR = 0.05 (12
accelerated flowering and 1 delayed flowering (*wrky38*). Rosette and inflorescence weights
were the traits with the next most insertion lines with genotype effects, 14 lines having nominally
significant effects and 5 significant at FDR = 0.05 that mostly led to increased rosette weight
and reduced inflorescence weight. Genotype effects on other traits measured were less
common, with 5,5,5,3,1 lines having nominally significant effects and 1,0,0,1,1 lines significant
with FDR = 0.05 for silique number, average and total silique length, inflorescence length, and
aboveground biomass, respectively.

We found few genotype-by-treatment interactions caused by the t-DNA insertions.
CS864818, a *WRKY38* (AT5G22570) knockout line, was the only one having a GxE effect that
was significant at FDR = 0.05 on inflorescence length, silique numbers, total silique length,
aboveground biomass, and inflorescence weight (Table 1,S5-S6). Generally, *wrky38* showed
less drought sensitivity than *Col* while being smaller and less fecund in well-watered conditions
except that the mutant had slightly longer inflorescence lengths than *Col* (*wrky38*: 40.1±3.010
cm; *Col*: 36.0±0.713 cm; Figure 2). When considering only genotype effects, the knockout of
WRKY38 caused significantly later flowering, longer inflorescence, more siliques, and greater
rosette weights (Figure 2, Table 1,S4-S6. However, under drought conditions, only flowering
time and inflorescence length were significantly different from *Col*.

Another mutant, *lsd1*, was one of only two having nominally significant genotype-by-
treatment interaction (GxTrt) on flowering time (p = 0.0015; the other line SALK_012432 had p =
0.0479). The mutation significantly accelerated flowering compared to *Col*, and its interaction
with drought exacerbated early flowering (Table 1). Notably, the mutant flowered earlier than
Col under both WW and drought conditions (WW: 58.1±1.06 vs. 58.6±0.972 days; drought:
52.3±3.58 vs. 68.1±1.94 days), and unlike *Col* the effect of drought on flowering time was not
significant (p = 0.3981). Drought had significant effects on fitness-related traits that reduced
silique numbers, average silique length, and total silique length while the GxTrt effects on
fitness were generally not significant (Table 1,S4-S6). lsd1 plants had significantly lower
inflorescence and rosette weights, thus lower aboveground biomass, compared to *Col* (Figure 2,
Table 1,S4-S6).

Both genotypes had increased WUE under drought compared to WW conditions (Figure
2), with treatment effects on WUE significant in LMMs that include *Col* and each mutant (p_{wrky38}

= 9.33e-18, p_{lsd1} = 1.15e-6). The genotype effect of *WRKY38* knockout also significantly
increased its WUE compared to *Col* (p = 8.81e-8), while that of *lsd1* was not significant (p =
0.756; Figure 2). *wrky38* had an increased $\delta^{15}N$ ratio under drought compared to the well-
watered condition, but its responses under both conditions were not substantially different from
Col (p_G = 0.5010, p_{Trt} = 0.5575; Figure 2). *lsd1* mutants, however, had $\delta^{15}N$ being lower under
drought while slightly higher under WW compared to *Col* (p_G = 0.0387, p_{GxTrt} = 4.22e-4),
suggesting a possible nitrogen source shift under drought. C:N ratios had been overall reduced
under drought, probably the result of reduced biomass under drought (Figure 2).

Comparing candidates from different GEA approaches

We compared each approach in identifying genes that influence fitness and performance traits
under drought, potentially indicating their role in drought adaptation. Among the 40 top-
candidate genes from the original studies, 12, 12, 8, 6, and 7 genes were identified through
RDA, partial RDA, univariate associations, the combined common garden approach with aridity
index, and the combined common garden approach with monthly growing season precipitation,
respectively (Table S1). Four genes were identified as top candidates by multiple approaches
(Table S1). Among the knockout lines of these genes, only SALK_051689 (AT5G10960) had
marginally significant genotype effect that accelerated flowering time (p_{adj} = 0.045, Table S5).
RDA and partial RDA proved to be the most effective methods for identifying drought-
adapted genes, with each approach having 6 out of 12 genes exhibiting significant genotype
effects at FDR = 0.05 across fitness and performance traits in the main drought screen. The
combined common garden approach followed, with 2 out of 6 and 2 out of 7 genes identified via
association with aridity index and monthly precipitation showing significant genotype effects at
FDR = 0.05.
In contrast, none of the 8 genes identified by univariate associations showed significant
genotype effects at FDR = 0.05, although 5 genes exhibited significant genotype effects at α =
0.05 before applying false discovery rate corrections.

Stable but less pronounced fitness, phenology, and ecophysiological responses of *wrky38* and *lsd1* under follow-up mild drought

Under the milder follow-up drought (ID), *wrky38* and *lsd1* plants also exhibited similar but less
pronounced responses compared to those observed in the main drought screen. The mild
drought significantly reduced fitness and inflorescence length across both *Col* and the mutants,
as determined by mixed linear models that included *Col* and one mutant (*wrky38* or *lsd1*).
Specifically, the drought significantly reduced fitness and inflorescence length for both *wrky38*
and *lsd1*, while flowering time was not significantly impacted (Figure 3, Table 2,S7). Plants of
wrky38 mutants did not exhibit significant genotype-specific responses to drought compared to
Col across the traits we measured (Table 2). However, the direction of *wrky38* differences was
consistent with the earlier drought screen; e.g., *wrky38* had higher fitness and biomass than*Col*
under drought, but the opposite under well-watered conditions (Figure 3). In contrast, *lsd1*

mutants showed significant genotype effects that reduce inflorescence length and total silique
length and a significant GxE effect for silique number (Table 2). This indicates that *lsd1* mutants
not only have reduced fitness under drought conditions, but the negative impact of drought was
even more pronounced in *lsd1* mutants compared to *Col*.
During the 10 days of ecophysiological tracking, we built LMMs to compare the daily g_{sw}
and Fv/Fm of the two mutants with *Col*. The stomatal conductance (g_{sw}) of all three genotypes
exhibited fluctuations that were associated with the regular watering schedule (Figure S1).
wrky38 had potentially adaptive reduced g_{sw} compared to *Col* and under drought, with genotype
effects significant during the initial days of tracking, when drought-treated plants first
experienced water limitation (9 days after treatment initiation; Table S3). A significant GxE
interaction with drought was also observed (Table S8). The *lsd1* mutant also had reduced g_{sw}
under drought, but the genotype effect was insignificant (Figure S1; Table S8). None of the
Fv/Fm values we measured suggested major oxidative stress (Figure S2), coinciding with
Zivcak et al. (2013), indicating a mild drought. No genotype or treatment pattern was detected
for Fv/Fm.
The drought effect significantly reduced SLA when considering *Col* and *wrky38* but was
not significant in the model including *Col* and *lsd1* (Figure 3, Table 2). Genotype effects on SLA
were significant for both mutants, but *wrky38* had a lower SLA (potentially adaptive under
drought) while *lsd1* had a higher SLA compared to *Col* (Figure 3, Table 2). RWC exhibited little
variation between genotypes and treatments except for *lsd1* plants, which had a significant
genotype effect resulting in lower RWC (Table S7). The $\delta^{13}C$ response to treatment was
generally lower in the follow-up than in the main drought screen, indicating a milder drought in
the follow-up. The treatment effect on WUE was only significant for *wrky38* (p = 0.0029),
whereas the genotype effect on WUE was only significant for *lsd1* (p = 3.7e-4). No significant
GxE effects were detected for SLA, RWC, or $\delta^{13}C$ (Table S7).

422 Reduced vegetative growth or fitness of *wrky38* and *lsd1* under
423 overnight freezing

Treated with overnight freezing for 30 days, both *lsd1* and *wrky38* had significantly smaller
diameters than *Col* (p_{wrky38} = 0.0184, p_{lsd1} = 3.7e-4), representing their reduced vegetative
growth under low temperatures (Figure 4). After returning to warmer temperatures, we found no
variation in flowering time and silique number between *Col* and the two mutants. However, *lsd1*
had a shorter inflorescence length (p = 0.0360), average silique length (p = 0.0023), and total
silique length (p = 0.0218) than *Col*. Both *lsd1* and *wkry38* had smaller biomass than *Col* (p_{wrky38}
= 0.0026, p_{lsd1} = 2.6e-4), which might also be due to reduced vegetative growth. Overall, *lsd1*
plants had reduced growth and fitness compared to *Col* under freezing, consistent with findings
of Huang et al. (2010) that *lsd1* is chilling sensitive. Although putative LoF variants in *WRKY38*
were strongly associated with warm winter temperatures (Lasky et al. 2018), *WRKY38*
knockouts did not exhibit fitness differences compared to *Col*, though their vegetative growth
was inhibited by overnight freezing compared to *Col*.

WRKY38 allelic variation associated with climatic moisture and temperature

To evaluate potential selection on *WRKY38* and its surrounding genomic region, we analyzed SNP diversity patterns and examined allele associations indicating possible selection effects. The Tajima's D of the 5-kb window containing the *WRKY38* gene was above 0 (Tajima's D = 1.429) and is above the average of Chromosome 5 (average Tajima's D = 0.660), but it did not significantly deviate from the distribution of Tajima's D across the chromosome (z-score = 0.659; Figure S3). The overall r^2 between SNPs in 10 kb around *WRKY38* and the most common functional variant (Frameshift$_{7495793}$) also indicated some haplotype structure, with the frameshift having elevated correlations (r^2 >0.4) for SNPs located within 2 kb upstream of *WRKY38* and in the first 500 bp of the gene (also where the frameshift was located, maximum r^2 = 0.5104; Figure S4). However, this was not a dramatically long region of linkage, suggesting no evidence for a recent sweep (cf. the strong correlation observed with chr4:10999188 and other SNPs in a 300 kb window around *LSD1,* Lee et al., 2017). Nevertheless, the putative LoF mutations carried large numbers of alternate SNP alleles across the locus, indicating some divergence and linkage (Figure 5i,S5). The haplotype structure at *WRKY38* can also be seen in the *WRKY38* gene neighbor-joining tree. While the genome-wide NJ tree shows a star-like structure consistent with the recent expansion of Arabidopsis (Lee et al. 2017) with accessions clustered based on geography (Figure 5f), the *WRKY38* tree shows strong divergence between sequences with Frameshift$_{7495793}$ versus putative intact *WRKY38* sequences (Figure 5g,h). This allelic variation is segregated across diverse lineages, including the relicts (Figure 5a,b,c,f). Coincidentally, the rare FVs were all unique to their lineages, suggesting recent origins (Figure 5f; Table S9). In contrast to the widespread Frameshift$_{7495793}$, they might reflect newer local putative LoF mutations.

The gene tree of WRKY38 shows that alleles with the same functional variants exhibit similar haplotype patterns, regardless of their genetic cluster (Figure 5g,h). For example, although most N-Euro accessions are in the non-Frameshift$_{7495793}$ clade, all 12 N-Euro accessions having Frameshift$_{7495793}$ are in the Frameshift$_{7495793}$ clade, overlapping with W-Med, CW-Euro, and Asian accessions carrying this allele. The only accession having Frameshift$_{7495794}$ and the only one with Stop-lost$_{7495609}$ are both within Frameshift$_{7495793}$ clade, while the 6 accessions having Frameshift$_{7496041}$ and 2 accessions having Stop-gained$_{7496443}$, which are grouped at the same tip on the gene tree, only occurred in the non-Frameshift$_{7495793}$ clade.

Inconsistency between the genomewide and *WRKY38* gene tree confirmed *WRKY38* allelic variation independent of population genetic structure. Next, we investigated whether this variation is associated with climatic factors, particularly moisture and drought related variables. We did not find significant correlations at α = 0.05 between the frequency of Frameshift$_{7495793}$ (the most common FV) or Frameshift$_{7495793\&94}$ (the FV that putative restores the normal ORF) and the bioclimatic variables when considering kinship (Table S10), which coincided with the findings of Alonso-Blanco et al. (2016) that almost no SNPs are significantly correlated with climate in environmental GWAS at FDR = 0.05 after considering population structure for 1001 Genomes accessions. Despite this, without accounting for kinship, significant differentiation was detected at α = 0.05 for 11, 12, and 4 bioclimatic variables between accessions with and without intact *WRKY38*, Frameshift$_{7495793}$, and Frameshift$_{7495793\&94}$, respectively (Table S11). Among the

significant associations, precipitation of the warmest quarter (bio18) had the lowest p-values for
both intact *WRKY38* (p = 6.8e-10) and Frameshift$_{7495793}$ (p = 8.8e-9). Specifically, intact
WRKY38 occurred more frequently in regions with wet summers, while Frameshift$_{7495793}$ was
more common in hot and dry regions, suggesting the drought adaptation of *WRKY38* putative
LoF. Frameshift$_{7495793\&94}$ (which has a net in-frame effect) was most significantly correlated with
higher isothermality (bio3; p = 3.7e-4), followed by higher mean temperature of the wettest
quarter (bio8; p = 6.4e-4).

LSD1 allelic variation and expression associated with climatic moisture and temperature

Mixed models (GEMMA) detected strong correlations between SNPs within 10kb around *LSD1*,
particularly for SNPs within 1 kb upstream and the first 500 bp of *LSD1*, and multiple bioclimatic
variables (Figure 6a). These SNPs, located in the promoter region of *LSD1*, may influence gene
expression by altering transcription factor binding motifs (Shastry, 2009; Wang et al., 2005).
Thus, we hypothesized that these SNPs might contribute to local adaptation via changes in
gene expression.
To validate the hypothesis, we picked a top climate-associate SNP (Chr4, position
11003866) from the 1 kb upstream region of *LSD1* and tested for *LSD1* expression differences
between ref/alt alleles. This SNP had the strongest correlation, indicated by the lowest p-value,
with precipitation of the warmest quarter (bio18; wald-p = 2.390e-41; Figure 6b), followed by the
minimum temperature of the driest quarter (bio9; wald-p = 2.140e-36; Figure 6c). We found that
the alternative allele corresponded to significantly lower *LSD1* expression compared to the
reference allele (p = 0.003; Figure 6d), suggesting a connection between lower *LSD1*
expression and hot and dry climates.
Using linear models, we found that *LSD1* expression was significantly greater for
genotypes from wetter summer climates (precipitation of the warmest quarter (p=8.683e-5) and
precipitation of the wettest month (p=0.039)). Across the 10 kb region around *LSD1*, 68 of the
201 SNPs were most strongly associated with bio18 (precipitation of the warmest quarter).
Moreover, when limiting to SNPs within 1kb upstream of *LSD1*, i.e. the region we thought to be
linked with *LSD1*, the ratio became 25 of 35 SNPs. The results indicated that *LSD1* might play a
role in local adaptation to moisture regimes by *cis*-regulatory variation.

DISCUSSION

Local adaptation is a major source of phenotypic variation within species, but traditional
methods of identifying locally adapted genes are logistically challenging. Genotype-environment
associations (GEA) offer an easy-to-implement tool to generate hypotheses about genes
involved in local adaptation, but most published results remain as untested hypotheses. Here
we screened 42 genes identified from genome-wide GEA studies and found at least two had
effects on environmental responses and physiology linked to drought adaptation, suggesting
potential local adaptations.

In our main drought experiment, among 42 potential drought-adapted genes detected
with 3 different GEA approaches, the tDNA knockout of *WRKY38* (detected with RDA, Lasky et
al., 2012) exhibited a significant GxE effect on fitness-related traits, and *lsd1* mutant (gene
detected with combined GEA-common garden synthesis, Lasky et al., 2018) had a significant
GxE effect on flowering time, which could potentially indicate variation in drought escape versus
drought avoidance (Lovell et al., 2013). Coincidentally, these genes were both the top (#1)
candidate genes with the strongest GEA in the studies used to select them. The physiological
effect of *wrky38* identified in the follow-up screening suggests mechanisms of drought
avoidance for *wrky38*, via reduced stomatal conductance and SLA. Furthermore, the high
frequency of putative natural loss-of-function alleles of *WRKY38* in Arabidopsis populations
found in the drier southern part of its range suggests potential local adaptation.

528 Genotype and genotype-by-enviroment effects on phenotypes

Although the screened genes were the ones with the largest effects in the original
studies, the GxE effects for fitness that were significant in the main drought screen were no
longer significant under the milder drought conditions of the follow-up ID experiment. Several
factors could impact the results. First, the milder conditions of the follow-up drought experiment
compared to the main drought screen may have reduced the G×E effect on fitness. Second,
over-simplified laboratory environments could lead to systematic bias in plant fitness (Anderson
et al., 2013). In natural environments, drought conditions are usually coupled with elevated
temperatures, a variable not accounted for in controlled experiments designed to isolate the
effects of a single environmental factor. In the laboratory, when only a single environmental
variable is changed between treatments, a conditional neutrality, that a gene has fitness
advantages under one treatment and is neutral under the other, can be more common
(Anderson et al., 2013). Lastly, the knockout alleles of a gene may not be the mechanism of
local adaptation in nature. Local adaptation may also occur through *cis*-regulatory effects on
gene expression (Hämälä et al., 2020; Lasky et al., 2014) or changes in amino acid sequences.
Therefore, when screening candidate genes for local adaptation, it is important to consider lines
with significant genotype effects alongside those showing significant G×E effects for fitness. As
G×E effects may not always be significant due to the factors discussed above, genotype effects
may still capture environment-associated differentiation.
Besides G×E effects on fitness, G×E effects on flowering time are also noteworthy when
assessing local adaptation to drought stress, as early flowering is a critical drought escape
strategy (Franks, 2011; Kenney et al., 2014). Chong and Stinchcombe (2019) reported
systematic directional effects of flowering time among randomly selected SALK lines. In their
study, 11 out of 27 lines under long-day conditions and 3 out of 27 lines under short-day
conditions exhibited significantly different flowering times compared to *Col*, with all showing
delayed flowering. However, in our main drought screen, 19 out of 20 lines with significant
genotype effects at $\alpha = 0.05$, and 12 out of 13 lines with significant genotype effects at FDR =
0.05, exhibited accelerated flowering (Figure 1, Table S6). Among these, *wrky38* was the only
line that showed delayed flowering (a potential drought avoidance strategy, Ludlow, 1989).
Since the genes we selected were candidates for drought adaptation, the accelerated flowering

observed in these lines may indicate their potential for drought adaptation via escape (Ludlow,
1989).
Though we only observed GxE for fitness in a small number of genes, evidence of
associations between their natural functional variation and moisture-related bioclimate variables
demonstrated the underlying biological bases of potential local adaptation of loci identified by
GEA. Setting up this screen was relatively easy due to the availability of existing tDNA
resources, making the approach an efficient way to identify and screen potential locally adapted
genes in model organisms where such resources are available or relatively easily generated.

566 Comparison of common GEA approaches

Since the advent of GEA, a wide variety of methods have been proposed (reviewed by
Rellstab et al., 2015). We tested candidates from three approaches here: multivariate ordination
(RDA) of genotype and environment (Forester et al., 2018; Lasky et al., 2012), univariate
climate mixed model associations (Lasky et al., 2014), and univariate climate models integrated
with fitness data from multiple common gardens (Lasky et al., 2018). Even considering lines
exhibiting significant genotype effects in addition to GxTrt effects, the number of positives
remains too small to definitely compare these statistical GEA approaches. While Lotterhos
(2023) recently criticized RDA as inappropriate for GEA due to a high false positive rate, the
author used arbitrary and uncalibrated significance thresholds. By contrast, Figure S8 in
Lotterhos (2023) showed that RDA was among the best methods based on the area under the
precision-recall curve (AUC-PR). Our results further support this, showing that RDA and partial
RDA had the highest rate of genes exhibiting significant genotype effects at FDR = 0.05,
followed by the combined common garden approach. In comparison, none of the tested genes
identified through univariate associations showed significant genotype effects at FDR = 0.05.
We propose that using ranked GEA results is more informative than focusing on arbitrary
significance thresholds; as mentioned previously, *WRKY38* and *LSD1* were the top-ranked loci
in their respective studies (RDA and combined GEA-common garden mixed models,
respectively; Lasky et al. 2012, 2018).

585 Potential mechanisms of drought adaptation at *WRKY38* and
586 *LSD1*

The two mutants presented opposite responses under drought. *wrky38* had a significantly
smaller SLA when compared to *Col* and when under drought, indicating how *wrky38* maintained
stable RWC under well-watered and drought conditions. *wrky38* mutants have reduced g_{sw},
SLA, and leaf area (result not shown) under drought in the follow-up experiment and increased
WUE in response to drought during both experiments (Figure 2-3), suggesting a drought-
avoidance strategy of natural *WRKY38* LoF genotypes. In contrast, *lsd1* had a larger SLA than
Col, thus potentially more water loss (Niinemets, 2001), which may result in its reduced RWC.
Instead, *lsd1* mutants showed stable accelerated flowering under drought in our two drought
experiments, indicating a drought-escape strategy.
WRKY38 negatively regulates plant basal pathogen defense by suppressing
PATHOGENESIS-RELATED GENE 1 (*PR1*) expression, a defense gene induced by salicylic

acid (SA) (Kim et al., 2008). Additionally, *WRKY38* is also involved in drought- and cold-related
responses in barley (Marè et al., 2004). Among *Arabidopsis WRKY* members that were also
acting in the SA pathway for plant basal defense responses (Salinas et al., 2024), *AtWRKY54*
and *AtWRKY70* were found to negatively impact osmotic tolerance by reducing stomatal closure
(Li et al., 2013), while *AtWRKY46* was found to regulate light-dependent stomatal opening in
guard cells (Ding et al., 2014). Based on previous studies and our findings, we hypothesize that
WRKY38 inhibits stomatal closure. If so, the lower g_{sw} we found in knockouts could reduce
water loss through the stomata and enhance fitness under drought.
In 1001 Genome accessions, natural *WRKY38* LoF alleles were strongly associated with
lower July moisture compared to intact alleles. Despite this strong association, these putative
LoF alleles were not fixed or entirely absent in any region. Our population genetics analyses
revealed that LoF *WRKY38* exists across the entire distribution region of *A. thaliana* and in
every lineage group (Figure 5a-f). We suspected that the LoF of *WRKY38* might have originated
before the migration across Eurasia of *A. thaliana* and colonized drier regions during this
process. In both drought experiments, *wrky38* generally had lower fitness compared to *Col*
under well-watered conditions. However, *wrky38* maintained relatively stable fitness under
drought stress, while *Col* showed a substantial fitness decrease. The drought insensitivity of
wrky38 may explain how natural LoF *WRKY38* dominates drier regions during migration.
The absence of LSD1 protein in *lsd1* mutants promotes the accumulation of SA
(Bernacki et al., 2021), which adversely affects plant development and reproduction while
accelerating flowering in Arabidopsis (Salinas et al., 2024). These findings explained the
consistent accelerated flowering behavior observed across our two drought experiments.
However, *lsd1* plants differed in their fitness responses to the long-term water deficiency of our
main drought screen and periodical moisture fluctuation in our follow-up experiments (Figure 2-
3). Specifically, *lsd1* mutants had similar fitness to *Col* under well-watered and drought in the
main experiment, but they were less fit than *Col* (i.e. genotype effect was significant) in the
follow-up. Previous laboratory studies have shown that *lsd1* exhibits similar survival and seed
production to the wild type under nonlethal water deficiency. However, under lethal drought
stress that caused complete mortality in wild-type plants, *lsd1* demonstrated a significantly
higher survival rate, albeit with reduced seed production. These findings suggest a trade-off
between survival and fecundity under severe stress in controlled laboratory conditions
(Szechyńska-Hebda et al., 2016; Wituszynska et al., 2013). When grown in the field, *lsd1*
mutants did not exhibit seed yield differences compared to the wild type (Bernacki et al., 2019;
Szechyńska-Hebda et al., 2016; Wituszynska et al., 2013). This could explain our finding that
the *LSD1* gene had lower expression in the hotter and drier regions, as reduced expression
could similarly mitigate fitness trade-offs, enhancing survival in stressful climates.

Freezing responses of *wrky38* and *lsd1*

We conducted the freezing experiment because both genes may also be involved in adaptation
to freezing stress. A SNP within the *WRKY38* coding region (Chr. 5, pos:7496047) was the top
candidate loci associated with temperature of the coldest month in Lasky et al. (2018), and
LSD1 is associated with chilling sensitivity (Huang et al., 2010). Moreover, these two genes are
both in the salicylic acid (SA) pathway that can be activated by cold stress (Miura and Tada,

2014; Wu et al., 2019). *WRKY38* was positively regulated by SA (Kim et al., 2008) while *LSD1*
conditionally regulated SA concentration with the existences of *ENHANCED DISEASE*
SUSCEPTIBILITY1 (*EDS1*) and *PHYTOALEXIN DEFICIENT4* (*PAD4*) (Bernacki et al., 2019;
Szechyńska-Hebda et al., 2016; Wituszynska et al., 2013).
WRKY38 knockouts exhibited similar fitness to *Col* despite reduced vegetative growth
after long-term overnight freezing (Figure 4). Considering that *WRKY38* was originally identified
through RDA (Lasky et al., 2012), this finding suggests a potential role in multivariate
adaptation. The LoF allele might be advantageous in drier and warmer climates, while the
functional allele might be more beneficial in cooler and wetter climates. *LSD1* knockouts had
significantly reduced vegetative growth and fitness under freezing, coincident with the findings
of Huang et al. (2010). Both mutants exhibited some disadvantages under freezing conditions
compared to *Col*, indicating possible links between the function of these genes and adaptation
to cold climates. Although both drought and freezing conditions can limit water availability and
there is overlap in some molecular pathways or responses to each set of conditions (Kim et al.,
2024), here our results suggest the drought-adaptive variants are not cold-adapted.

Conclusion

In this study, we tested the potential role of 44 t-DNA knockout mutants of GEA-identified genes
in adaptation to drought stress using common garden experiments. While most mutants did not
exhibit significant G×E effects for flowering time, performance, or fitness, two knockouts, *wrky38*
and *lsd1*, demonstrated evidence of drought adaptation in both the main drought screen and
follow-up intermediate drought common gardens. Natural variation in the function or expression
of these genes across a moisture gradient further supports the utility of GEA approaches for
generating hypotheses, though further experiments are required to test hypotheses. Our
findings highlight the promise of GEA methods for uncovering novel local adaptations to
environmental stressors.

Acknowledgments

666   We thank Cody Depew for assistance in ordering knockout lines, Amanda Penn for assistance
667   with experiments, and Diana Gamba and Yuxing Xu for assistance with computation. Funding
668   was provided by NIH R35GM138300 to JRL.

References

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., Gadrinab, C., Heller, C., Jeske, A., Koesema, E., Meyers, C.C., Parker, H., Prednis, L., Ansari, Y., Choy, N., Deen, H., Geralt, M., Hazari, N., Hom, E., Karnes, M., Mulholland, C., Ndubaku, R., Schmidt, I., Guzman, P., Aguilar-Henonin, L., Schmid, M., Weigel, D., Carter, D.E., Marchand, T., Risseeuw, E., Brogden, D., Zeko, A., Crosby, W.L., Berry, C.C., Ecker, J.R., 2003. Genome-Wide Insertional Mutagenesis of Arabidopsis thaliana. Science 301, 653–657. https://doi.org/10.1126/science.1086391

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K.M., Cao, J., Chae, E., Dezwaan, T.M., Ding, W., Ecker, J.R., Exposito-Alonso, M., Farlow, A., Fitz, J., Gan, X., Grimm, D.G., Hancock, A.M., Henz, S.R., Holm, S., Horton, M., Jarsulic, M., Kerstetter, R.A., Korte, A., Korte, P., Lanz, C., Lee, C.-R., Meng, D., Michael, T.P., Mott, R., Muliyati, N.W., Nägele, T., Nagler, M., Nizhynska, V., Nordborg, M., Novikova, P.Y., Picó, F.X., Platzer, A., Rabanal, F.A., Rodriguez, A., Rowan, B.A., Salomé, P.A., Schmid, K.J., Schmitz, R.J., Seren, Ü., Sperone, F.G., Sudkamp, M., Svardal, H., Tanzer, M.M., Todd, D., Volchenboum, S.L., Wang, C., Wang, G., Wang, X., Weckwerth, W., Weigel, D., Zhou, X., 2016. 1,135 Genomes Reveal the Global Pattern of Polymorphism in *Arabidopsis thaliana*. Cell 166, 481–491. https://doi.org/10.1016/j.cell.2016.05.063

Anderson, J.T., Lee, C.-R., Rushworth, C.A., Colautti, R.I., Mitchell-Olds, T., 2013. Genetic trade-offs and conditional neutrality contribute to local adaptation. Mol. Ecol. 22, 699–708. https://doi.org/10.1111/j.1365-294X.2012.05522.x

Bernacki, M.J., Czarnocka, W., Rusaczonek, A., Witoń, D., Kęska, S., Czyż, J., Szechyńska-Hebda, M., Karpiński, S., 2019. LSD1-, EDS1- and PAD4-dependent conditional correlation among salicylic acid, hydrogen peroxide, water use efficiency and seed yield in Arabidopsis thaliana. Physiol. Plant. 165, 369–382. https://doi.org/10.1111/ppl.12863

Bernacki, M.J., Rusaczonek, A., Czarnocka, W., Karpiński, S., 2021. Salicylic Acid Accumulation Controlled by LSD1 Is Essential in Triggering Cell Death in Response to Abiotic Stress. Cells 10, 962. https://doi.org/10.3390/cells10040962

Betancourt, N.J., Rajpurohit, S., Durmaz, E., Fabian, D.K., Kapun, M., Flatt, T., Schmidt, P., 2021. Allelic polymorphism at foxo contributes to local adaptation in Drosophila melanogaster. Mol. Ecol. 30, 2817–2830. https://doi.org/10.1111/mec.15939

Capblancq, T., Lachmuth, S., Fitzpatrick, M.C., Keller, S.R., 2023. From common gardens to candidate genes: exploring local adaptation to climate in red spruce. New Phytol. 237, 1590–1605. https://doi.org/10.1111/nph.18465

Chong, V.K., Stinchcombe, J.R., 2019. Evaluating Population Genomic Candidate Genes Underlying Flowering Time in *Arabidopsis thaliana* Using T-DNA Insertion Lines. J. Hered. 110, 445–454. https://doi.org/10.1093/jhered/esz026

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., Ruden, D.M., 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6, 80–92. https://doi.org/10.4161/fly.19695

Clausen, J., Keck, D.D., Hiesey, W.M., 1941. Regional Differentiation in Plant Species. Am. Nat. 75, 231–250.

Coop, G., Witonsky, D., Di Rienzo, A., Pritchard, J.K., 2010. Using Environmental Correlations to Identify Loci Underlying Local Adaptation. Genetics 185, 1411–1423. https://doi.org/10.1534/genetics.110.114819

Cortés, A.J., López-Hernández, F., Blair, M.W., 2022. Genome–Environment Associations, an
Innovative Tool for Studying Heritable Evolutionary Adaptation in Orphan Crops and
Wild Relatives. Front. Genet. 13. https://doi.org/10.3389/fgene.2022.910386
Crossa, J., Pérez-Rodríguez, P., Cuevas, J., Montesinos-López, O., Jarquín, D., Campos, G. de
los, Burgueño, J., González-Camacho, J.M., Pérez-Elizalde, S., Beyene, Y.,
Dreisigacker, S., Singh, R., Zhang, X., Gowda, M., Roorkiwal, M., Rutkoski, J.,
Varshney, R.K., 2017. Genomic Selection in Plant Breeding: Methods, Models, and
Perspectives. Trends Plant Sci. 22, 961–975.
https://doi.org/10.1016/j.tplants.2017.08.011
Ding, Z.J., Yan, J.Y., Xu, X.Y., Yu, D.Q., Li, G.X., Zhang, S.Q., Zheng, S.J., 2014. Transcription
factor WRKY46 regulates osmotic stress responses and stomatal movement
independently in Arabidopsis. Plant J. 79, 13–27. https://doi.org/10.1111/tpj.12538
Dittberner, H., Korte, A., Mettler-Altmann, T., Weber, A.P.M., Monroe, G., de Meaux, J., 2018.
Natural variation in stomata size contributes to the local adaptation of water-use
efficiency in *Arabidopsis thaliana*. Mol. Ecol. 27, 4052–4065.
https://doi.org/10.1111/mec.14838
El-Soda, M., Kruijer, W., Malosetti, M., Koornneef, M., Aarts, M.G.M., 2015. Quantitative trait
loci and candidate genes underlying genotype by environment interaction in the
response of Arabidopsis thaliana to drought. Plant Cell Environ. 38, 585–599.
https://doi.org/10.1111/pce.12418
Endler, J.A., 1973. Gene Flow and Population Differentiation. Science 179, 243–250.
https://doi.org/10.1126/science.179.4070.243
Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H.A., Weigel, D., 2018.
Genomic basis and evolutionary potential for extreme drought adaptation in Arabidopsis
thaliana. Nat. Ecol. Evol. 2, 352–358. https://doi.org/10.1038/s41559-017-0423-0
Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for
global land areas. Int. J. Climatol. 37, 4302–4315. https://doi.org/10.1002/joc.5086
Forester, B.R., Lasky, J.R., Wagner, H.H., Urban, D.L., 2018. Comparing methods for detecting
multilocus adaptation with multivariate genotype–environment associations. Mol. Ecol.
27, 2215–2233. https://doi.org/10.1111/mec.14584
Franks, S.J., 2011. Plasticity and evolution in drought avoidance and escape in the annual plant
Brassica rapa. New Phytol. 190, 249–257. https://doi.org/10.1111/j.1469-
8137.2010.03603.x
Gehan, M.A., Park, S., Gilmour, S.J., An, C., Lee, C.-M., Thomashow, M.F., 2015. Natural
variation in the C-repeat binding factor cold response pathway correlates with local
adaptation of Arabidopsis ecotypes. Plant J. 84, 682–693.
https://doi.org/10.1111/tpj.13027
Hager, E.R., Harringmeyer, O.S., Wooldridge, T.B., Theingi, S., Gable, J.T., McFadden, S.,
Neugeboren, B., Turner, K.M., Jensen, J.D., Hoekstra, H.E., 2022. A chromosomal
inversion contributes to divergence in multiple traits between deer mouse ecotypes.
Science 377, 399–405. https://doi.org/10.1126/science.abg0718
Hämälä, T., Gorton, A.J., Moeller, D.A., Tiffin, P., 2020. Pleiotropy facilitates local adaptation to
distant optima in common ragweed (Ambrosia artemisiifolia). PLOS Genet. 16,
e1008707. https://doi.org/10.1371/journal.pgen.1008707
Hancock, A.M., Brachi, B., Faure, N., Horton, M.W., Jarymowycz, L.B., Sperone, F.G.,
Toomajian, C., Roux, F., Bergelson, J., 2011. Adaptation to Climate Across the
Arabidopsis thaliana Genome. Science 334, 83–86.
https://doi.org/10.1126/science.1209244
Hijmans, R.J., Van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J.A., Lamigueiro,
O.P., Bevan, A., Racine, E.B., Shortridge, A., 2015. Package 'raster.' R Package 734,
473.

Huang, X., Li, Y., Zhang, X., Zuo, J., Yang, S., 2010. The Arabidopsis LSD1 gene plays an important role in the regulation of low temperature-dependent cell death. New Phytol. 187, 301–312. https://doi.org/10.1111/j.1469-8137.2010.03275.x

Kang, H.M., Zaitlen, N.A., Wade, C.M., Kirby, A., Heckerman, D., Daly, M.J., Eskin, E., 2008. Efficient Control of Population Structure in Model Organism Association Mapping. Genetics 178, 1709–1723. https://doi.org/10.1534/genetics.107.080101

Kawakatsu, T., Huang, S.C., Jupe, F., Sasaki, E., Schmitz, R.J., Urich, M.A., Castanon, R., Nery, J.R., Barragan, C., He, Y., 2016. Epigenomic diversity in a global collection of Arabidopsis thaliana accessions. Cell 166, 492–505.

Kenney, A.M., McKay, J.K., Richards, J.H., Juenger, T.E., 2014. Direct and indirect selection on flowering time, water-use efficiency (WUE, δ13C), and WUE plasticity to drought in Arabidopsis thaliana. Ecol. Evol. 4, 4505–4521. https://doi.org/10.1002/ece3.1270

Kim, D.-H., Sung, S., 2010. The Plant Homeo Domain finger protein, VIN3-LIKE 2, is necessary for photoperiod-mediated epigenetic regulation of the floral repressor, MAF5. Proc. Natl. Acad. Sci. 107, 17029–17034. https://doi.org/10.1073/pnas.1010834107

Kim, J.-S., Kidokoro, S., Yamaguchi-Shinozaki, K., Shinozaki, K., 2024. Regulatory networks in plant responses to drought and cold stress. Plant Physiol. 195, 170–189. https://doi.org/10.1093/plphys/kiae105

Kim, K.C., Lai, Z., Fan, B., Chen, Z., 2008. Arabidopsis WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. Plant Cell 20, 2357–2371. https://doi.org/10.1105/tpc.107.055566

Lasky, J.R., Des Marais, D.L., Lowry, D.B., Povolotskaya, I., McKay, J.K., Richards, J.H., Keitt, T.H., Juenger, T.E., 2014. Natural Variation in Abiotic Stress Responsive Gene Expression and Local Adaptation to Climate in Arabidopsis thaliana. Mol. Biol. Evol. 31, 2283–2296. https://doi.org/10.1093/molbev/msu170

Lasky, J.R., Des Marais, D.L., McKAY, J.K., Richards, J.H., Juenger, T.E., Keitt, T.H., 2012. Characterizing genomic variation of Arabidopsis thaliana: the roles of geography and climate. Mol. Ecol. 21, 5512–5529. https://doi.org/10.1111/j.1365-294X.2012.05709.x

Lasky, J.R., Forester, B.R., Reimherr, M., 2018. Coherent synthesis of genomic associations with phenotypes and home environments. Mol. Ecol. Resour. 18, 91–106. https://doi.org/10.1111/1755-0998.12714

Lasky, J.R., Josephs, E.B., Morris, G.P., 2023. Genotype–environment associations to reveal the molecular basis of environmental adaptation. Plant Cell 35, 125–138. https://doi.org/10.1093/plcell/koac267

Lee, C.-R., Svardal, H., Farlow, A., Exposito-Alonso, M., Ding, W., Novikova, P., Alonso-Blanco, C., Weigel, D., Nordborg, M., 2017. On the post-glacial spread of human commensal Arabidopsis thaliana. Nat. Commun. 8, 14458. https://doi.org/10.1038/ncomms14458

Lee, G., Sanderson, B.J., Ellis, T.J., Dilkes, B.P., McKay, J.K., Ågren, J., Oakley, C.G., 2024. A large-effect fitness trade-off across environments is explained by a single mutation affecting cold acclimation. Proc. Natl. Acad. Sci. 121, e2317461121. https://doi.org/10.1073/pnas.2317461121

Li, J., Besseau, S., Törönen, P., Sipari, N., Kollist, H., Holm, L., Palva, E.T., 2013. Defense-related transcription factors WRKY70 and WRKY54 modulate osmotic stress tolerance by regulating stomatal aperture in Arabidopsis. New Phytol. 200, 457–472. https://doi.org/10.1111/nph.12378

Lorts, C.M., Lasky, J.R., 2020. Competition × drought interactions change phenotypic plasticity and the direction of selection on Arabidopsis traits. New Phytol. 227, 1060–1072. https://doi.org/10.1111/nph.16593

Lotterhos, K.E., 2023. The paradox of adaptive trait clines with nonclinal patterns in the underlying genes. Proc. Natl. Acad. Sci. 120, e2220313120. https://doi.org/10.1073/pnas.2220313120

Lovell, J.T., Juenger, T.E., Michaels, S.D., Lasky, J.R., Platt, A., Richards, J.H., Yu, X., Easlon, H.M., Sen, S., McKay, J.K., 2013. Pleiotropy of FRIGIDA enhances the potential for multivariate adaptation. Proc. R. Soc. B Biol. Sci. 280, 20131043. https://doi.org/10.1098/rspb.2013.1043

Ludlow, M.M., 1989. Strategies of response to water stress. In K. H. Kreeb, H. Richter, & T. M. Hinckley (Eds.), Structural and Functional Responses to Environmental Stresses (pp. 269–281). SPB Academic.

Luo, Y.; Lorts, C.M.; Lawrence-Paul, E.; Lasky, J.R. (Forthcoming 2025). Dataset from Experimental validation of genome-environment associations in Arabidopsis [Dataset]. Dryad. https://doi.org/10.5061/dryad.qrfj6q5s4

Marè, C., Mazzucotelli, E., Crosatti, C., Francia, E., Stanca, A. ?michele, Cattivelli, L., 2004. Hv-WRKY38: a new transcription factor involved in cold- and drought-response in barley. Plant Mol. Biol. 55, 399–416. https://doi.org/10.1007/s11103-004-0906-7

Miura, K., Tada, Y., 2014. Regulation of water, salinity, and cold stress responses by salicylic acid. Front. Plant Sci. 5.

Monroe, J.G., Powell, T., Price, N., Mullen, J.L., Howard, A., Evans, K., Lovell, J.T., McKay, J.K., 2018. Drought adaptation in Arabidopsis thaliana by extensive genetic loss-of-function. eLife 7, e41038. https://doi.org/10.7554/eLife.41038

Niinemets, Ü., 2001. Global-Scale Climatic Controls of Leaf Dry Mass Per Area, Density, and Thickness in Trees and Shrubs. Ecology 82, 453–469. https://doi.org/10.1890/0012-9658(2001)082[0453:GSCCOL]2.0.CO;2

Nyine, M., Davidson, D., Adhikari, E., Clinesmith, M., Wang, H., Akhunova, A., Fritz, A., Akhunov, E., 2025. Genomic signals of ecogeographic adaptation in a wild relative are associated with improved wheat performance under drought stress. Genome Biology 26, 35. https://doi.org/10.1186/s13059-025-03500-1

Paradis, E., Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics 35, 526–528.

Pisupati, R., Reichardt, I., Seren, Ü., Korte, P., Nizhynska, V., Kerdaffrec, E., Uzunova, K., Rabanal, F.A., Filiault, D.L., Nordborg, M., 2017. Verification of Arabidopsis stock collections using SNPmatch, a tool for genotyping high-plexed samples. Sci. Data 4, 170184. https://doi.org/10.1038/sdata.2017.184

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., De Bakker, P.I., Daly, M.J., 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. Am. J. Hum. Genet. 81, 559–575.

Ralph, P.L., Coop, G., 2015. Convergent Evolution During Local Adaptation to Patchy Landscapes. PLOS Genet. 11, e1005630. https://doi.org/10.1371/journal.pgen.1005630

Ratcliffe, O.J., Kumimoto, R.W., Wong, B.J., Riechmann, J.L., 2003. Analysis of the Arabidopsis MADS AFFECTING FLOWERING Gene Family: MAF2 Prevents Vernalization by Short Periods of Cold [W]. Plant Cell 15, 1159–1169. https://doi.org/10.1105/tpc.009506

Rellstab, C., Gugerli, F., Eckert, A.J., Hancock, A.M., Holderegger, R., 2015. A practical guide to environmental association analysis in landscape genomics. Mol. Ecol. 24, 4348–4370. https://doi.org/10.1111/mec.13322

Salinas, P., Velozo, S., Herrera-Vásquez, A., 2024. Salicylic acid accumulation: emerging molecular players and novel perspectives on plant development and nutrition. J. Exp. Bot. erae309. https://doi.org/10.1093/jxb/erae309

Savolainen, O., Lascoux, M., Merilä, J., 2013. Ecological genomics of local adaptation. Nat. Rev. Genet. 14, 807–820. https://doi.org/10.1038/nrg3522

Shastry, B.S., 2009. SNPs: Impact on Gene Function and Phenotype, in: Komar, A.A. (Ed.), Single Nucleotide Polymorphisms: Methods and Protocols. Humana Press, Totowa, NJ, pp. 3–22. https://doi.org/10.1007/978-1-60327-411-1_1

Signorell, A., Aho, K., Alfons, A., Anderegg, N., Aragon, T., Arppe, A., Baddeley, A., Barton, K.,
Bolker, B., Borchers, H., 2023. DescTools: tools for descriptive statistics. R package
version 0.99. 26. Compr. R. Arch Netw 289–291.
Sinnwell, J.P., Therneau, T.M., Schaid, D.J., 2014. The kinship2 R package for pedigree data.
Hum. Hered. 78, 91–93.
Soudi, S., Jahani, M., Todesco, M., Owens, G.L., Bercovich, N., Rieseberg, L.H., Yeaman, S.,
2023. Repeatability of adaptation in sunflowers reveals that genomic regions harbouring
inversions also drive adaptation in species lacking an inversion. eLife 12, RP88604.
https://doi.org/10.7554/eLife.88604
Szechyńska-Hebda, M., Czarnocka, W., Hebda, M., Karpiński, S., 2016. PAD4, LSD1 and
EDS1 regulate drought tolerance, plant biomass production, and cell wall properties.
Plant Cell Rep. 35, 527–539. https://doi.org/10.1007/s00299-015-1901-y
Tergemina, E., Ansari, S., Salt, D.E., Hancock, A.M., 2025. Multiple independent MGR5 alleles
contribute to a clinal pattern in leaf magnesium across the distribution of Arabidopsis
thaliana. New Phytol. n/a. https://doi.org/10.1111/nph.70069
Therneau, T.M., Therneau, M.T.M., 2015. Package 'coxme'. R Package Version 2.
Tigano, A., Friesen, V.L., 2016. Genomics of local adaptation with gene flow. Mol. Ecol. 25,
2144–2164. https://doi.org/10.1111/mec.13606
Wadgymar, S.M., DeMarche, M.L., Josephs, E.B., Sheth, S.N., Anderson, J.T., 2022. Local
Adaptation: Causal Agents of Selection and Adaptive Trait Divergence. Annu. Rev. Ecol.
Evol. Syst. 53, 87–111. https://doi.org/10.1146/annurev-ecolsys-012722-035231
Wang, X., Tomso, D.J., Liu, X., Bell, D.A., 2005. Single nucleotide polymorphism in
transcriptional regulatory regions and expression of environmentally responsive genes.
Toxicol. Appl. Pharmacol., Living in a Safe Chemical World. Proceedings of the 10th
International Congress of Toxicology 11-15 July, 2004, Tampere, Finland 207, 84–90.
https://doi.org/10.1016/j.taap.2004.09.024
Whiting, J.R., Booker, T.R., Rougeux, C., Lind, B.M., Singh, P., Lu, M., Huang, K., Whitlock,
M.C., Aitken, S.N., Andrew, R.L., Borevitz, J.O., Bruhl, J.J., Collins, T.L., Fischer, M.C.,
Hodgins, K.A., Holliday, J.A., Ingvarsson, P.K., Janes, J.K., Khandaker, M., Koenig, D.,
Kreiner, J.M., Kremer, A., Lascoux, M., Leroy, T., Milesi, P., Murray, K.D., Pyhäjärvi, T.,
Rellstab, C., Rieseberg, L.H., Roux, F., Stinchcombe, J.R., Telford, I.R.H., Todesco, M.,
Tyrmi, J.S., Wang, B., Weigel, D., Willi, Y., Wright, S.I., Zhou, L., Yeaman, S., 2024. The
genetic architecture of repeated local adaptation to climate in distantly related plants.
Nat. Ecol. Evol. 8, 1933–1947. https://doi.org/10.1038/s41559-024-02514-5
Wijfjes, R. Y., Boesten, R., Becker, F.F. M., Theeuwen, T.P.J.M., Snoek, B.L., Mastoraki, M.,
Verheijen, J.J., Güvencli, N., Denkers, L.-A.M., Koornneef, M., van Eeuwijk, F.A., Smit,
S., de Ridder, D., Aarts, M.G.M., 2024. Allelic variants confer Arabidopsis adaptation to
small regional environmental differences. Plant J. 120, 1662–1681.
https://doi.org/10.1111/tpj.17067
Wituszynska, W., Slesak, I., Vanderauwera, S., Szechynska-Hebda, M., Kornas, A., Van Der
Kelen, K., Mühlenbock, P., Karpinska, B., Mackowski, S., Van Breusegem, F., Karpinski,
S., 2013. Lesion simulating disease1, enhanced disease susceptibility1, and phytoalexin
deficient4 conditionally regulate cellular signaling homeostasis, photosynthesis, water
use efficiency, and seed yield in Arabidopsis. Plant Physiol. 161, 1795–1805.
https://doi.org/10.1104/pp.112.208116
Wu, Z., Han, S., Zhou, H., Tuang, Z.K., Wang, Y., Jin, Y., Shi, H., Yang, W., 2019. Cold stress
activates disease resistance in Arabidopsis thaliana through a salicylic acid dependent
pathway. Plant Cell Environ. 42, 2645–2663. https://doi.org/10.1111/pce.13579
Yeaman, S., Whitlock, M.C., 2011. THE GENETIC ARCHITECTURE OF ADAPTATION
UNDER MIGRATION–SELECTION BALANCE. Evolution 65, 1897–1911.
https://doi.org/10.1111/j.1558-5646.2011.01269.x

Yim, C., Bellis, E.S., DeLeo, V.L., Gamba, D., Muscarella, R., Lasky, J.R., 2024. Climate biogeography of Arabidopsis thaliana: Linking distribution models and individual variation. J. Biogeogr. 51, 560–574. https://doi.org/10.1111/jbi.14737

Zhao, M., Li, M., Huang, M., Liang, C., Chen, D., Hwang, I., Zhang, W., Wang, M., 2023. The cysteine-rich receptor-like kinase CRK4 contributes to the different drought stress response between Columbia and Landsberg erecta. Plant Cell Environ. 46, 3258–3272. https://doi.org/10.1111/pce.14665

Zivcak, M., Brestic, M., Balatova, Z., Drevenakova, P., Olsovska, K., Kalaji, H.M., Yang, X., Allakhverdiev, S.I., 2013. Photosynthetic electron transport and specific photoprotective responses in wheat leaves under drought stress. Photosynth. Res. 117, 529–546. https://doi.org/10.1007/s11120-013-9885-3

Zoubarev, A., Hamer, K.M., Keshav, K.D., McCarthy, E.L., Santos, J.R.C., Van Rossum, T., McDonald, C., Hall, A., Wan, X., Lim, R., Gillis, J., Pavlidis, P., 2012. Gemma: a resource for the reuse, sharing and meta-analysis of expression profiling data. Bioinformatics 28, 2272–2273. https://doi.org/10.1093/bioinformatics/bts430

Captions

Figure 1 Reaction norms (lines) and average trait values (boxes) of a subset of traits measured during the main drought screen across all 44 mutants and *Col*, with *Col*, *lsd1* (CS68738), and *wrky38* (CS864818) highlighted. Each line connects the average trait values of one genotype under well-watered (WW) and drought (D) treatments, with *Col*, *lsd1* (CS68738), and *wrky38* (CS864818) highlighted. Mutants showing significant genotype effects at FDR = 0.05 from linear mixed models (LMMs) comparing *Col* and one mutant are indicated by dashed lines.

Figure 2 A subsetof the traits measured during the main drought screen of *Col*, *lsd1* (CS68738), and *wrky38* (CS864818). Text annotations on the box plots indicate the significance of genotype (G), treatment (E), and genotype-treatment interaction (G×E) effects at α = 0.05, based on LMMs comparing *Col* and one mutant. The text color denotes the mutant genotype. Significance levels are represented as follows: *, $0.01 \leq p < 0.05$; **, $0.001 \leq p < 0.01$; ***, $p < 0.001$.

Figure 3 Traits measured in the follow-up intermediate drought experiment of *Col*, *lsd1* (CS68738), and *wrky38* (CS864818). Text annotations on the box plots indicate the significance of genotype (G), treatment (E), and genotype-treatment interaction (G×E) effects at α = 0.05, based on LMMs comparing *Col* and one mutant genotype. The text color denotes the mutant genotype. Significance levels are represented as follows: *, $0.01 \leq p < 0.05$; **, $0.001 \leq p < 0.01$; ***, $p < 0.001$.

Figure 4 Traits measured in the follow-up freezing experiment of *Col*, *lsd1* (CS68738), and *wrky38* (CS864818). Asterisks indicate significant trait differences between *Col* and each mutant based on Dunnett's test: *, $0.01 \leq p < 0.05$; **, $0.001 \leq p < 0.01$; ***, $p < 0.001$.

Figure 5 Natural variation of *WRKY38* putative LoF in the 1001 Genomes accessions. **a**, distribution of native *Arabidopsis* accessions from 1001 Genomes. Colors indicate ADMIXTURE genetic clusters, originally defined by Alonso-Blanco et al. (2016) and modified in this study. Accessions with clearly assigned genetic clusters are represented by filled icons in their respective cluster colors, while admixed accessions are shown with unfilled icons. **b-e**, distributions of natural *WRKY38* putative functional variation: **b**, distribution of accessions with intact *WRKY38*; **c**, distribution of accessions with Frameshift$_{7495793}$, the most common FV of *WRKY38* ; **d**, distribution of accessions with Frameshift$_{7495793\&94}$, a mutation that restores the

normal open reading frame; **e**, distribution of accessions with other rare FVs of *WRKY38*. **f,**
WRKY38 FVs on the genomewide NJ tree, the relict lineage (Alonso-Blanco et al., 2016) was
marked by an arrow. **g**, *WRKY38* FVs on the NJ gene tree, symbols as in (**f**). **h,**
Frameshift$_{7495793}$ on the NJ gene tree. **i**, numbers of alternative SNPs in accessions with intact
WRKY38 and putative LoF FVs within 10 kb around *WRKY38*. The number of accessions with
each type of functional variation is indicated in parentheses.
Figure 6 Genotype-environment associations and *LSD1* expression. **a**, Scanning the *LSD1*
region to identify potential SNP-environment associations using LMM within 10kb. Each point
represents the smallest p-value from the 19 associations between each SNP and the 19
bioclimate variables. *LSD1* gene region is shown between the dashed lines. These associations
are meant to identify where in the locus variation is most strongly associated with climate. **b**, the
difference in precipitation of the warmest quarter (bio18) between reference and alternative
alleles at a top climate-associated SNP (Chr4, position 11003866). **c**, the difference in the
minimum temperature of the driest quarter (bio9) at an SNP (Chr4, position 11003866). **d**, the
difference in *LSD1* expression at an SNP (Chr4, position 11003866). **b-d**, asterisks indicate
significant trait differences from t-test: *, $0.01 \leq p < 0.05$; **, $0.001 \leq p < 0.05$; ***, $p < 0.001$.
Table 1 Summary statistics from mixed linear models (LMMs) assessing a subset of
performance and fitness traits for *Col*, CS864818 (*wkry38*) and CS68738 (*lsd1*) under well-
watered and drought conditions during the main drought screen. Each LMM independently
analyzes *Col* and one mutant (either *wrky38* or *lsd1*). β represents the estimated effect size, and
subscripts indicate effects attributed to genotype (G), treatment (Trt) or genotype-by-treatment
interaction (GxTrt); *p* represents the *p*-value at α = 0.05, and p_{adj} represents the adjusted *p*-value
at FDR = 0.05. A negative $β_G$ reflects a lower trait value in the mutant compared to *Col*, while a
positive $β_{Trt}$ reflects a lower trait value under drought compared to well-watered conditions.
Table 2 Summary statistics from mixed linear models (LMMs) assessing a subset of
performance and fitness traits for *Col*, CS864818 (*wkry38*) and CS68738 (*lsd1*) under well-
watered and intermediate drought (ID) conditions during the follow-up intermediate drought
experiment. Each LMM independently analyzes *Col* and one mutant (either *wrky38* or *lsd1*). β
represents the estimated effect size, *p* represents the *p*-value at α = 0.05, and subscripts
indicate effects attributed to genotype (G), treatment (Trt) or genotype-by-treatment interaction
(GxTrt). A negative $β_G$ reflects a lower trait value in the mutant compared to *Col*, while a positive
$β_{Trt}$ reflects a lower trait value under ID compared to well-watered conditions.

Author contributions

997   YL, CML, and JRL designed the study, CL and YL performed experiments, YL conducted data
998   analyses, YL, ELP, and JRL interpreted results, YL, CML, and JRL wrote the manuscript, all
999   authors edited and approved the manuscript.

Figures and Tables

1001 **Figure 1**
1002

1003
1004 **Figure 2**
1005

1006
1007 **Figure 3**
1008

1009
1010 **Figure 4**
1011

1012   **Figure 5**
1013

1014
1015   **Figure 6**

1026 **Table 1**

Col-CS864818 (*wkry38*)	Genotype Effect				Treatment Effect				Genotype-by-Treatment		
	β_G	*p*	*p*$_{adj}$		β_{Trt}	*p*	*p*$_{adj}$		β_{GxTrt}	*p*	*p*$_{adj}$
Days to Flower (days)	15.86	**<0.0001**	**0.0002**		-9.57	**<0.0001**	**<0.0001**		-4.76	0.2380	0.8081
Inflorescence Length (cm)	27.33	**<0.0001**	**0.0014**		19.74	**0.0086**	**0.0086**		-23.21	**0.0006**	**0.0274**
Total Silique Length (cm)	15.82	**0.0018**	0.0782		120.03	**0.0009**	**0.0009**		-131.47	**0.0002**	**0.0076**
Biomass (g)	0.08	0.6793	0.9218		0.18	**0.0059**	**0.0059**		-0.14	**0.0004**	**0.0189**

Col-CS68738 (*lsd1*)	Genotype Effect				Treatment Effect				Genotype-by-Treatment		
	β_G	*p*	*p*$_{adj}$		β_{Trt}	*p*	*p*$_{adj}$		β_{GxTrt}	*p*	*p*$_{adj}$
Days to Flower (days)	-15.86	**0.0009**	**0.0082**		-9.57	0.3981	0.3981		15.43	**0.0015**	0.0675
Inflorescence Length (cm)	19.74	0.3586	0.6422		-6.20	**<0.0001**	**<0.0001**		20.14	0.0852	0.9698
Total Silique Length (cm)	17.75	0.2947	0.6362		120.03	**<0.0001**	**<0.0001**		-15.68	0.4049	0.9684
Biomass (g)	-0.13	**<0.0001**	**<0.0001**		0.18	**<0.0001**	**<0.0001**		0.03	0.4058	0.7440

1027 Note: *p*-values less than 0.05 are presented in bold to indicate statistical significance.

1028

1029 **Table 2**

Col-CS864818 (*wkry38*)	β_G	*p*$_G$	β_{Trt}	*p*$_{Trt}$	β_{GxTrt}	*p*$_{GxTrt}$
Days to Flower (days)	0.50	0.1503	-0.33	0.9684	0.93	0.4626
Inflorescence Length (cm)	7.75	0.2416	12.67	**0.0105**	-8.11	0.2551
Silique Number	5.00	0.9632	26.67	**0.0023**	-10.50	0.4705
Total Silique Length (cm)	3.63	0.8752	25.68	**<0.0001**	-9.23	0.2925
Specific Leaf Area (mm^2/mg)	-3.80	**0.0005**	1.19	**0.0313**	1.43	0.4181

Col-CS68738 (*lsd1*)	β_G	*p*$_G$	β_{Trt}	*p*$_{Trt}$	β_{GxTrt}	*p*$_{GxTrt}$
Days to Flower (days)	1.42	0.7466	-0.33	0.1908	-3.63	0.1365
Silique Number	-25.00	0.8293	26.67	**0.0007**	48.51	**0.0081**
Inflorescence Length (cm)	-21.10	**<0.0001**	12.67	**<0.0001**	8.65	0.1897
Total Silique Length (cm)	-18.25	**0.0234**	25.68	**<0.0001**	14.45	0.1378
Specific Leaf Area (mm^2/mg)	3.11	0.0368	1.19	0.4479	1.41	0.6989

1030 Note: *p*-values less than 0.05 are presented in bold to indicate statistical significance.