NF-κB is a Central Regulator of Hypoxia-Induced Gene Expression

Dilem Shakir[1], Michael Batie[1], Chun-Sui Kwok[1], Simon J. Cook[2], Niall S. Kenneth[1], and Sonia Rocha[1]*

[1]Department of Biochemistry, Cell and System Biology. Institute of Systems, Molecular and Integrative Biology, University of Liverpool. Liverpool L697ZB. United Kingdom.
[2]Signalling Programme, The Babraham Institute, Babraham Research Campus, Cambridge CB22 3AT. United Kingdom.

*Corresponding author: Sonia Rocha, srocha@liverpool.ac.uk

Keywords: Hypoxia, NF-kappaB, RNA-seq, transcriptional repression

Abstract

Hypoxia is both a physiological and pathological signal in cells. Changes in gene expression play a critical role in the cellular response to hypoxia, enabling cells to adapt to reduced oxygen availability by regulating metabolism, survival and angiogenesis. These changes are primarily mediated by the HIF family of transcription factors, however other transcription factors such as NF-κB, are also activated by hypoxia. Although NF-κB is known to be activated under hypoxic conditions the extent to which NF-κB contributes to the hypoxic response remains poorly understood. In this study we analysed hypoxia-induced, NF-κB-dependent gene expression, to define the NF-κB-dependent hypoxic signature. Our analysis reveals that a significant proportion of hypoxia-induced gene expression changes are NF-κB dependent, with the majority of genes downregulated by hypoxia requiring NF-κB for their repression. We show that, while the NF-κB-mediated hypoxic response may vary between cell types, there are a core subset of hypoxia-inducible genes that require NF-κB across multiple cell backgrounds. We demonstrate that NF-κB is critical for reactive oxygen species (ROS) generation and regulation of genes involved in oxidative phosphorylation under hypoxic conditions. This work highlights that NF-κB has a central role in the hypoxia response, regulating a broader set of genes than previously thought, providing new understanding of the regulation of gene expression by hypoxia and NF-κB.

Introduction

Hypoxia, or reduction in oxygen homeostasis, is both a physiological and pathological cue (Kaelin & Ratcliffe, 2008; Kenneth & Rocha, 2008; Xie & Simon, 2017). As oxygen is required for viability of many organisms, complex and intricate mechanisms have evolved for cells and organisms to be able to sense hypoxia and respond appropriately. These responses vary from increasing oxygen supply, to reducing oxygen consumption, to eventual adaptation or cell death (Rocha, 2007). At the cellular level, the best understood response occurs via the Hypoxia Inducible Factor (HIF) transcription factor family. HIFs are controlled by a family of 2-oxoglutarate dependent dioxygenases called Prolyl-hydroxylases (PHDs), that modify HIF, making it a target for ubiquitylation via the von Hipple Lindau (VHL) containing E3-ligase complex and destruction by the 26S proteasome (An & Rettig, 2005; Bruegge *et al*, 2007; Fandrey *et al*, 2006). In addition to HIF, other transcription factors are also activated in response to hypoxia, including the NF-κB family of transcription factors (Cummins & Taylor, 2005; Kenneth & Rocha, 2008) NF-κB is best known for its role in innate immunity and its response and control of inflammation, but is also activated in response to physical, physiological and oxidative stresses (Perkins & Gilmore, 2006). The NF-κB transcription factor family is composed of 7 distinct proteins that are encoded by 5 different genes, namely RelA (p65), RelB, cRel, NF-κB1 (p105/p50) and NF-κB2 (p100/p52) (Perkins & Gilmore, 2006). Unlike the other NF-κB subunits, NF-κB1 and NF-κB2 are synthesised as pro-forms, p105 and p100, which are proteolytically processed to their respective active forms, p50 and p52 (Perkins & Gilmore, 2006). Regulation of the NF-κB family occurs at several levels from transcription to post-translational modifications and also cytoplasmic sequestration via their interaction with a family of inhibitors, called Inhibitors of κB, or IκBs (Perkins & Gilmore, 2006). Given its complexity, NF-κB gene targets are numerous and can also be characterised based on their timing of expression, such as early, mid and late genes (Tian *et al*, 2005). Furthermore, NF-κB activity is under several positive and negative feedback loops depending on the activating stimulus, including NF-κB subunits themselves, IκBs, and upstream activators and regulators (Prescott *et al*, 2021). NF-κB is activated by almost all stresses investigated so far but is best characterised following infection with bacteria or virus, or in response to pro-inflammatory cytokines recognised by members of the tumour necrosis factor receptor superfamily (TNFRSF) (Perkins, 2007). Receptor activation initiates a signalling cascade involving the IκB kinase (IKK) complex that phosphorylates IκB, leading to its degradation and the release of NF-κB for nuclear translocation and control of gene transcription

(Adhikari *et al*, 2007). Many more levels of complexity and control exist in the NF-κB pathway, making it one of the most complex transcription factors to study. However, its importance is well established in immunity, response to inflammation, cell death, proliferation and DNA damage responses (Almaden *et al*, 2016; Gerondakis & Siebenlist, 2010; Liu *et al*, 2017; Qin *et al*, 2007; Volcic *et al*, 2012; Xia *et al*, 2018; Xu *et al*, 2015).

NF-κB activation in response to hypoxia has been observed by several studies, including some of our own (Al-Anazi *et al*, 2018; Chandel *et al*, 2000; Culver *et al*, 2010; D'Ignazio *et al*, 2017; D'Ignazio & Rocha, 2016; Koong *et al*, 1994; Melvin *et al*, 2011; Patel *et al*, 2017; Taylor *et al*, 1999). While the exact mechanism of activation slightly differs from study to study, it is well established that hypoxia-dependent activation of NF-κB requires IKK activation, and this has been seen in the fruit fly, mouse and human cells (Bandarra *et al*, 2014; Culver *et al.*, 2010). Additional mechanistic insights include a requirement for a calcium-dependent IKK upstream kinase (Culver *et al.*, 2010), Transforming Growth Factor Activated Kinase (TAK1) (Adhikari *et al.*, 2007; Melvin *et al.*, 2011), input from PHD enzymes (Cummins & Taylor, 2005; Wilson *et al*, 2020) and lack of IκB degradation (Culver *et al.*, 2010).

It is well established that cells have a specific transcriptional response to hypoxia, and analysis of several RNA-sequencing datasets revealed a conserved hypoxia inducible signature (Puente-Santamaria *et al*, 2022). There is also a recognised set of genes that are repressed in response to hypoxia, but the mechanisms controlling this are poorly understood (Batie *et al*, 2018; Cavadas *et al*, 2017). However, whether the NF-κB response to hypoxia is a general feature of most cells, or if there is a common gene expression signature, is not known. Here we combined analysis of publicly available datasets across different cellular backgrounds, with transcriptomics analysis of siRNA-mediated depletion of NF-κB subunits in hypoxia to uncover the contribution of NF-κB to the cellular response. Our analysis shows NF-κB is required for activation of a subset of hypoxia activated genes (i.e., 30%) but strikingly is required for the repression of around 60% of hypoxia repressed genes. Our analysis reveals a core set of hypoxia-induced NF-κB target genes across all cell types, with some cell type specificity. Interestingly, most of the genes were not previously known to be dependent on NF-κB to be regulated in hypoxia, and only a small subset is recognised NF-κB target genes, demonstrating the context-dependent nature of NF-κB. Lastly, we demonstrate that NF-κB is required to maintain reactive oxygen levels necessary for cell signalling.

Methods

Cell culture

Human cervical carcinoma HeLa, lung carcinoma A549, brain glioblastoma U87, breast adenocarcinoma MCF-7 cell lines were obtained from the American Type Culture Collection (ATCC, USA) and IKKα and IKKβ double knockout (DKO) HCT116 cell lines were previously described in (Prescott *et al*, 2022). All of the cell lines were maintained at 5% CO_2 and 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% v/v foetal bovine serum (FBS), 1% L-glutamine (Lonza, UK), and 1% penicillin-streptomycin (Lonza, UK). Cell lines were cultured not more than 30 passages and routinely tested for mycoplasma contamination using MycoAlert Mycoplasma Detection Kit (Lonza, UK).

Treatments and siRNA transfections

Hypoxia treatments were performed by incubating cells in an InVivo300 Hypoxia Workstation (Baker Ruskin, UK) at 1% O_2, 5% CO_2 and 37°C. For stabilisation of NRF2, cells were treated with 100 nM of Bardoxolone methyl **(**CDDO-Me, Sigma) for 16 h, and 2.5 µM of Saha (HDAC inhibitor) for 24 h. Cells were transfected with 27 nM of small interfering RNA (siRNA) oligonucleotides (Eurofins Genomics, Germany) for 48 h using Interferin (Polyplus, France) transfection reagent according to manufacturer's instructions. The following oligonucleotide sequences were used for siRNA knockdowns: Control CAGUCGCGUUUGCGACUGG RelA GCCCUAUCCCUUUACGUCA, RelB GGAUUUGCCGAAUUAACAA, cRel ACAGCUGAAUGAUAUUGAA.

RNA extraction and RNA-seq data analysis

RNA was extracted using the RNeasy Mini Kit (Qiagen, UK) following manufacturer's instructions. RNA-seq was performed on a Novaseq 6000 (Illumina, UK) with paired end 150 bp run type. Sequencing reads were aligned to the human genome assembly version hg38 (GRCh38, Ensemble) using STAR sequence aligner (Dobin *et al*, 2013) to generate coordinate-sorted binary alignment map (bam) files. Read counts for each transcript (GRCh38, Ensembl) were generated using the featureCounts function of Subread (Liao *et al*, 2013). Differential expression analysis was performed using R Bioconductor package DESeq2 (Love *et al*, 2014) with filtering for effect size (>±0.58 $\log_2$ fold change) and statistical significance (FDR <0.05) to determine differentially expressed genes (DEGs). Sample to sample distance comparison heatmaps using rlog transformed read count data were generated using R Bioconductor packages DESeq2 and pheatmap. PCA plot of rlog transformed read count data were generated

using R Bioconductor packages DESeq2 and ggplot2. Volcano plots for display of differential expression analysis were generated using R Bioconductor package EnhancedVolcano. Z-score heatmaps of VST transformed read count data were generated using DESeq2 and pheatmap. RelA, RelB, and cRel dependent and independent hypoxia responsive genes were determined by overlapping hypoxia control siRNA DEGs (compared to normoxia control siRNA) with hypoxia RelA, RelB, and cRel siRNA DEGs (compared to normoxia control siRNA). NF-κB dependent and independent hypoxia responsive genes were determined by overlapping hypoxia control siRNA DEGs (compared to normoxia control siRNA) with the combined list of RelA, RelB, and cRel siRNA DEGs (compared to normoxia control siRNA). Overrepresentation analysis (ORA) was performed using WEB-based Gene SeT AnaLysis Toolkit (WebGestalt) (Liao *et al*, 2019) in overrepresentation analysis (ORA) mode using the Molecular Signatures Database (MSigDB) hallmark gene sets (Liberzon *et al*, 2015; Subramanian *et al*, 2005). Motif enrichment analysis was performed using the findMotifs tool of HOMER (Heinz *et al*, 2010) with gene promoters defined as 400 bps upstream to 100 bps downstream of transcription start sites.

qPCR

RNA was extracted using the RNeasy Mini Kit (Qiagen, UK) following manufacturer's instructions. RNA was converted to cDNA using iscript reverse transcription kit (Bio-Rad) and qPCR analysis of transcript expression changes was performed by running 3 µl of DNA on a QuantStudio 1 qPCR platform (Applied Biosystems) with power-track Sybr green reaction mix (Applied Biosystems) in a final reaction of 15 µl. The following qPCR primers were used: Actin F CTGGGAGTGGGTGGAGGC and R TCAACTGGTCTCAAGTCAGTG, 18S F AAACGGCTACCACATCCAAG and R CGCTCCCAAGATCCAACTAC, CA9 F CTTTGCCAGAGTTGACGAGG and R CAGCAACTGCTCATAGGCAC, RelA F CTGCCGGGATGGCTTCTAT and R CCGCTTCTTCACACACTGGAT, RelB F TCCCAACCAGGATGTCTAGC and R AGCCATGTCCCTTTTCCTCT, cRel F CTGCCTTCTTCAAGCTGGTC and R CGCTTCCATTCCGACTATGT, GADD45B F TCGGCCAAGTTGATGAATG and R TGAGCGTGAAGTGGATTTG, JUNB F GACCAAGAGCGCATCAAA and R TCTTCACCTTGTCCTCCA, NFIX F TGTTGATGACGTGTTCTATCC and R CCAGCTTTCCTGACTTCTTTA, USP28 F GAAGTAGAGGAGTGGGAAGA and R GAAGGCTCTTGTGATGTAGAG, KMT2E F GCCAACTGCCCTACATAAA and R GCTCTTTACCCAGGAAGAATAC, TGFA F GCATGTGTCTGCCATTCT and R GTGATGGCCTGCTTCTTC, VIM F

CCAGCTAACCAACGACAAA and R TCCTCTCTCTGAAGCATCTC, SAP30 F GAGCGCAAGGCATCTTTA and R AATCACCTCCATCATCATCAC, EGLN3 F CTACGTCAAGGAGAGGTCTAA and R CAGATAGTAGATGCAGGTGATG, KLF10 F TGCCTTCGTGTTGAAATCC and R GGGCGCGATTATGCAATTA, BCL3 F CTCTCCATATTGCTGTGGTG and R TGTCTGCCGTAGGTTGT, FTH1 F AGTGCCGTTGTTCAGTTC and R AGACAGCCACACCTTAGT, GCLM F CTGTTCAGTCCTTGGAGTTG and R CTCCCAGTAAGGCTGTAAATG, NQO1 F CCTGGAAGGATGGAAGAAAC and R GAATCCTGCCTGGAAGTTTAG, AIFM1 F CCAGCCACCTTCTTTCTATG and R CCATGTTGTCTCTCACATCC, IDH1 F GCAGTACAAGTCCCAGTTT and R TGAAGCCTCCCTCTGATT, SOD1 F TCGAGCAGAAGGAAAGTAATG and R CCTGCTGTATTATCTCCAAACT, CCND3 F GATTTCCTGGCCTTCATTCT and R GGGTACATGGCAAAGGTATAA, CASP10 F CGTATCAAGGAGAGGAAGAAC and R TGTGGCTCTGTTACCATTAC, LAMTOR2 F CACCCTGCTGCTGAATAA and R CCCAGATGTTACTGGCTATG.

Statistical analysis

For qPCR analysis comparing two conditions, statistical significance was determined via Student's t-test. For qPCR analysis comparing more than two conditions, statistical significance was determined via one-way ANOVA with post-hoc Dunnett's test. For overlap of differentially expressed genes (DEGs) identified by RNA-seq, statistical significance was determined via hypergeometric test using Venn diagram function of an online data analysis tool with its default parameters (Iqbal, 2021). For all other statistical analysis, default settings of the analysis tool were used. In all cases, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Data mining of publicly available datasets

HeLa HIF ChIP-seq dataset (Ortmann *et al*, 2021) (GSE169040) was downloaded from the Gene Expression Omnibus (Edgar *et al*, 2002). MCF-7 (GSE153291), A549 (GSE186370), HCT116 (GSE81513), U87 (GSE78025), and 501-MEL (GSE132624) hypoxia and HUVEC TNF-α (Fowler *et al*, 2022) (GSE201466) RNA-seq datasets were downloaded from the Gene Expression Omnibus (Edgar *et al.*, 2002) and processed using the RNA-seq data analysis pipeline described above. HUVEC hypoxia RNA-seq data was obtained from (Tiana *et al*, 2018). NRF2 target genes were obtained from (Hayes & Dinkova-Kostova, 2014) and (Morgenstern *et al*, 2024). NF-κB target genes were downloaded from the Gilmore laboratory

website https://www.bu.edu/nf-kb/gene-resources/target-genes/ and converted to Ensembl gene annotation format using R Bioconductor package biomaRt (Durinck *et al*, 2005).

Cellular Reactive Oxygen Species detection

Cells were fixed and cellular reactive oxygen species (ROS) were measured using CellROX Green reagent (ThermoFisher) according to the manufacturer's instructions. Cells were imaged using an Epifluorescence microscopy (Zeiss Axio Observer Z.1). Images were quantified using OMERO open microscopy environment (Allan *et al*, 2012). Scale bars = 20µm.

Immunoblots

Cells were lysed in Radio Immunoprecipitation Assay (RIPA) lysis buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 1% v/v NP40, 0.25% w/v Na-deoxycholate, 0.1% w/v SDS, 10 mM NaF, 2 mM Na3VO4, and 1 protease inhibitor tablet (Thermo Fisher Scientific) per 10 mL of lysis buffer). Cells containing the lysis buffer scraped into Eppendorf tubes and incubated on ice for 10 minutes to complete the lysis process. The samples were centrifuged for 15 minutes at 13,000 rpm, 4°C. The supernatant was collected, and protein concentration of cell lysates were determined using Bradford assay (BioRad). Cells treated with hypoxia were harvested in the hypoxia chamber to prevent reoxygenation. 20 µg of protein was prepared in 2× SDS loading buffer (100 mM Tris-HCl pH 6.8, 20% v/v glycerol, 4% w/v SDS, 200 mM DTT and Bromophenol Blue). SDS-PAGE and immunoblots were carried out using standard protocols. The following primary antibodies were used for immunoblotting: RelA (sc-8008, Santa Cruz), RelB (10544, Cell Signaling), cRel (4727, Cell Signaling), HIF-1α (610958, BD Biosciences), β-Actin (66009-1-lg, Proteintech), IDH1 (12332-1-AP, Proteintech), SOD1 (sc-17767, Santa Cruz), Total Oxphos Antibody Cocktail for detecting ATP5A, UQCRC2, and SDHB (ab110411, Abcam), NRF2 (12721, Cell Signaling), BNIP3L (12396, Cell Signaling), Vimentin (5741, Cell Signaling), DYRK1B (5672, Cell Signaling). ImageJ was used for quantification of immunoblots with band intensities normalised to β-Actin.

Model figure was created in BioRender. Rocha, S. (2025) https://BioRender.com/

Results

Transcriptional response profile of hypoxia stimulation is NF-𝜅B dependent.

As the majority of NF-κB target genes have been characterised in the context of inflammatory signalling, we aimed to determine the contribution NF-κB to the transcriptional response to hypoxia within a given cell type. As such, to identify hypoxia-inducible, NF-κB-dependent

gene signatures, an unbiased high-throughput RNA-sequencing (RNA-seq) analysis was performed. HeLa cells were transfected with control siRNA or siRNA to NF-κB subunits, RelA, RelB or cRel and exposed to 21% O_2 (normoxia) or 1% O_2 (hypoxia) for 24 hours (h) prior to profiling for global transcriptomic analysis using RNA-seq (Figure 1A). Principal component analysis of the collected RNA-seq data showed that samples clustered by condition (Sup. Figure S1A). Control samples treated with normoxia clustered together with relatively weak correlation to the rest of the samples, while hypoxia-treated samples grouped close to one another, with the exception of siRelA-treated samples (Sup. Figure S1A). Also, close clustering of replicates within each condition demonstrated their relative similarity to each other. In this analysis, efficiency of the hypoxia stimulation was investigated using DESeq2 normalised counts acquired from RNA-seq analysis showing differential expression of the genes in hypoxia samples, compared to normoxia. As expected, the expression of Carbonic Anhydrase (*CA9*), a core hypoxia upregulated gene, was increased in response to hypoxia (Sup. Figure S1B). Additionally, DESeq2 normalised counts of *RelA*, *RelB*, and *cRel* were reduced by the respective siRNA treatments, demonstrating efficient siRNA depletion (Sup. Figure S1C-E). Successful depletion of NF-κB subunits by siRNA treatment was shown at the protein level by Western blot (Sup. Figure S1F). To investigate the impact of individual NF-κB subunit depletions on hypoxia gene expression changes, volcano plots were generated following differential expression analysis by comparing each condition to normoxia control (Figure 1B-E). This analysis revealed that more than 2,000 genes were differentially expressed at 5% false discovery rate (FDR) and log2 fold change 0.58 in hypoxia control siRNA treated samples compared to normoxia control siRNA treated samples (Figure 1B, Dataset S1). Overall, RelA depletion induced the highest transcriptional changes, followed by RelB or cRel depletions upon hypoxia stimulation (Figure 1C-E, Dataset S1). When we compared the effect of individual NF-κB subunits in hypoxia inducible gene signatures, all three subunits showed a shared proportion of genes between each other, while also having a high portion of genes regulated by RelA, RelB or cRel, exclusively (Figure 1F-G, Dataset S1).

Figure 1. Gene expression changes in response to hypoxia and NF-κB subunit depletions in HeLa cells.
A. RNA-seq (n=2) in HeLa cells cultured at 21% oxygen (normoxia) or exposed to 24 h 1% oxygen (hypoxia), transfected with control siRNA or RelA, RelB or cRel siRNAs. **B-E.** Differential expression analysis volcano plots for 24 h hypoxia control, siRelA, siRelB or sicRel compared to normoxia control. Blue points indicate DEGs. **F-G.** Overlap of siRelA, siRelB and sicRel upregulated genes in hypoxia, compared to normoxia control (**F**) and siRelA, siRelB and sicRel downregulated genes in hypoxia, compared to normoxia control (**G**) in HeLa cells.

To identify NF-κB-dependent, hypoxia-inducible gene signatures we performed integrative analysis, comparing the list of differentially hypoxia-regulated genes with the combined list of NF-κB subunit depleted, differentially hypoxia-regulated genes. Overlap analysis revealed that 35.5% (556 out of 1568) of the hypoxia-upregulated genes (Figure 2A, Sup. Figure S2A), and 62.6% (410 out of 655) of the hypoxia-downregulated genes were NF-κB-dependent (Figure

2B Sup. Figure S2B). Overall, this analysis highlights the importance of NF-κB in the regulation of the transcriptional response to hypoxia. Furthermore, the heatmap generated using the NF-κB dependent hypoxia up- and down-regulated genes showed that individual NF-κB subunit depletions displayed different transcriptional profiles (Figure 2C). We also investigated the number of NF-κB dependent hypoxia-regulated genes shared by the different Rel subunits, revealing that although 41 genes were simultaneous dependent on all NF-κB subunits for hypoxia upregulation, most of the other genes were regulated by individual subunits (Figure 2D, Dataset S2). Similar results were also obtained for repressed genes in hypoxia (Figure 2E, Dataset S2). This analysis revealed that the highest proportion hypoxia regulated genes RelB-dependent followed by cRel and RelA.

To characterise the identified NF-κB-dependent, hypoxia-inducible gene signatures, their association with hallmark gene sets was investigated. Hypoxia, glycolysis, and TNF-α signalling via NF-κB were the top three gene signatures significantly enriched at RelA-dependent 24 h hypoxia up-regulated DEGs. RelB-dependent 24 h hypoxia up-regulated DEGs were enriched in Epithelial–mesenchymal transition (EMT) and myogenesis gene signatures. Finally, EMT, TNF-α signalling via NF-κB, and inflammatory response were the top three gene signatures significantly enriched at cRel-dependent 24 h hypoxia upregulated DEGs (Figure 2F). Oxidative phosphorylation and ROS gene sets were significantly enriched at RelA-, RelB- and cRel-dependent 24 h hypoxia downregulated DEGs (Figure 2G). Motif enrichment analysis showed HIF-1α and HIF-1β motifs were enriched at RelA- and RelB-dependent 24 h hypoxia upregulated DEGs (Sup. Figure S3A), and HIF-2α and HIF-1β motifs were enriched at cRel-dependent 24 h hypoxia upregulated DEGs (Sup. Figure S3A), suggesting a role of HIF subunits in NF-κB controlled gene expression increases in hypoxia. The motif enrichment analysis of each NF-κB subunit-dependent hypoxia downregulated DEGs showed a variety of known NF-κB associated transcription factors (Sup. Figure S3B), indicating distinct transcriptional regulation of RelA, RelB and cRel dependent hypoxia downregulated gene signatures. To further elucidate the correlation of NF-κB controlled hypoxic gene expression changes with HIF binding sites, we analysed publicly available HIF subunit ChIP-seq datasets in HeLa cells exposed to hypoxia (Sup. Figure S4A-C). Similar to motif enrichment analysis, HIF subunit (HIF-1α, HIF-2α and HIF-1β) binding sites were enriched at RelA-, RelB-, and cRel-dependent 24 h hypoxia upregulated DEGs, but not at RelA-, RelB-, and cRel-dependent 24 h hypoxia downregulated DEGs, further indicating HIF involvement in NF-κB dependent hypoxia induced gene activation. We next further investigated if NF-κB subunits had been

found at the promoters of hypoxia regulated genes. To this end, we took advantage of the ChIP-Atlas database (Zou *et al*, 2024). This analysis revealed significant enrichment of RelA (80%) but also RelB (22%) and cRel (18%) in both hypoxia upregulated and downregulated genes (Sup. Figure S5A). However, similar results were obtained when analysing NF-κB subunit occupancy at NF-κB independent hypoxia responsive genes (Sup. Figure S5B), demonstrating enrichment of Rel binding sites from the ChIP atlas are not specific to NF-κB-dependent hypoxia responsive genes. This was expected, as many sites bound by transcription factors are not always functional, and their use varies between stimuli and cell background.

We also investigated if NF-κB-dependent, hypoxia-regulated genes have been previously described as direct NF-κB targets in the Gilmore database (Sup. Figure S5C). Interestingly, only a small number of NF-κB target genes from this database were present, although they are favoured at NF-κB dependent hypoxia activated over repressed genes.

In summary, this analysis showed that NF-κB plays a previously unrecognised and crucial role in the control of gene expression in hypoxia in the cell system analysed.

Figure 2. NF-κB is required for 35% of hypoxia induced genes and 63% of hypoxia repressed genes.
A-B. NF-κB dependence of hypoxia inducible Differential Expressed Genes (DEGs); percentage of NF-κB dependent DEGs are labelled. **C.** Heatmap displaying Z score transformed gene expression levels for NF-κB dependent hypoxia regulated genes. **D-E.** Overlap of RelA, RelB or cRel-dependent hypoxia up- or down-regulated DEGs. **F-G.** Over Representation Analysis (ORA) was performed through WEB-based Gene SeT AnaLysis Toolkit using the Molecular Signatures Database hallmark gene sets, investigating NF-κB dependent hypoxia upregulated DEGs (**F**) or downregulated DEGs (**G**). Dashed line shows statistical significance threshold of FDR 0.05 (-Log$_{10}$FDR 1.3).

NF-κB-dependent, hypoxia-inducible gene expression changes are also observed by qPCR.

To validate the obtained data from the RNA-seq analysis, we selected several genes with high log2 fold change in hypoxia compared to normoxia, and some known HIF and/ or NF-κB target genes amongst the 556 NF-κB-dependent, hypoxia-up-regulated genes (*BCL3*, *KMT2E*, *SAP30*, *TGFA*, *EGLN3*, *USP28*, *VIM*, *NFIX*, *GADD45B*, *JUNB*, *KLF10*, *FTH1*) and 410 NF-κB-dependent, hypoxia-down-regulated genes (*GCLM*, *CCND3*, *NQO1*, *AIFM1*, *SOD1*, *CASP10*, *LAMTOR2*, *IDH1*) to analyse by quantitative polymerase chain reaction (qPCR) (Figure 3A, Sup. Table S1). HeLa cells were transfected with control siRNA or siRNAs to NF-κB subunits, RelA, RelB, or cRel and exposed, or not, to 24 h hypoxia prior to qPCR analysis. In this analysis, efficiency of the hypoxia stimulation was measured using *CA9* (Sup. Figure S6A), as expected expression of this gene was highly upregulated in response to hypoxia. Additionally, RNA levels of *RelA*, *RelB*, and *cRel* were reduced in response to their respective siRNA treatments, demonstrating robust siRNA depletion (Sup. Figure S6B-D). RNA levels of all the selected NF-κB-dependent, hypoxia-upregulated genes were significantly increased in hypoxia control siRNA treated samples compared to normoxia control siRNA treated samples (Figure 3B-D). Next, effect of the NF-κB subunits was tested individually on hypoxia up- and down-regulated gene expression changes. The hypoxia-induced increase in *USP28*, *VIM*, *NFIX*, and *FTH1* was significantly decreased by RelA depletion compared to hypoxia control sample (Figure 3B), whilst *BCL3* showed a slight decrease upon RelA depletion in hypoxia, though this result was not statistically significant (Sup. Figure S6E). *KMT2E* levels were not altered by RelA depletion (Sup. Figure S6E). All the selected RelB-dependent hypoxia-upregulated genes, *USP28*, *VIM*, *NFIX*, *SAP30*, *TGFA*, and *EGLN3*, significantly decreased following RelB depletion in hypoxia compared to hypoxia control sample (Figure 3C). Similarly, the hypoxia induced increase in *USP28*, *FTH1*, and *JUNB* significantly decreased following cRel depletion (Figure 3D). *GADD45B* showed a slight decrease with cRel depletion in hypoxia, however this result was not statistically significant (Sup. Figure S6F) and *KLF10* was not responsive to cRel depletion (Sup. Figure S6F). In summary, 5 out of 6 RelA dependent hypoxia up-regulated genes, 6 out of 6 RelB dependent hypoxia-up-regulated genes, and 4 out of 5 cRel dependent hypoxia up-regulated genes were successfully validated by qPCR (Figure 3B-D). RNA levels of the selected NF-κB-dependent, hypoxia down-regulated genes significantly decreased in hypoxia control siRNA treated samples compared to normoxia control siRNA treated samples (Figure 3E-F). Following siRNA depletion of RelA, *GCLM* and *IDH1* significantly increased expression in hypoxia compared to hypoxia control (Figure 3E). Similarly, RelB depletion followed by hypoxia exposure, increased the mRNA abundance of

GCLM, *IDH1*, *AIFM1*, *CCND3*, and *CASP10*, compared to hypoxia control (Figure 3F). *SOD1* slightly increased with RelB depletion in hypoxia, however this result is not significant (Sup. Figure S6G). None of the selected cRel-dependent, hypoxia-downregulated genes were affected by cRel depletion in hypoxia, except *SOD1*, where cRel depletion gave the opposite effect to the RNA-seq analysis (Sup. Figure S6H). Also, LAMTOR2 was predicted to be both RelB- and cRel-dependent hypoxia regulated gene, however RelB or cRel depletion showed no effect on its mRNA abundance (Sup. Figure S6G-H). In conclusion, 5 out of 7 RelB-dependent, and 2 out of 2 RelA-dependent hypoxia down-regulated genes were successfully validated by qPCR, however none of the selected RNA-seq cRel dependent genes had an effect on reversing the impact of hypoxia stimulation on the gene expressions. This indicates that RelA and RelB have a more dominant role in hypoxia-induced down-regulation of genes, compared to cRel. Interestingly, when we investigated the protein levels of DYRK1B and Vimentin by western blot (Sup. Figure 7A-C), we confirmed that all NF-κBs tested were required for hypoxia-induced Vimentin, while cRel and, to a lesser extent, RelA were required for DYRK1B induction by hypoxia (Sup. Figure 7A-C).

To test if the NF-κB-dependent gene regulation that we observed in hypoxia, was also present under normal oxygen conditions, we performed qPCR experiments following siRNA depletion of the NF-κB subunits in basal oxygen levels (Sup. Figure S8A-H). Firstly, we confirmed efficient depletion of RelA, RelB and cRel by immunoblot analysis (Sup. Figure S8A-C). Of the 4 qPCR-validated RelA-dependent, hypoxia-upregulated genes, only FTH1 was regulated by RelA under normal oxygen conditions (Sup. Figure S8D). On the other hand, none of the RelB-dependent hypoxia-upregulated genes changed their expression levels in normoxia when RelB was depleted (Sup. Figure S8E). Regarding cRel-dependent hypoxia upregulated genes, JUNB and FTH1 displayed cRel-dependent regulation in normoxia (Sup. Figure S8F). Of the NF-κB-dependent hypoxia repressed genes, none of the RelA-dependent genes were controlled by RelA under normoxia and only 1 out of 5 (i.e., GCLM) of the RelB-dependent genes were regulated by RelB depletion in normoxia (Sup. Figure S8G-H). Overall, a majority of RelA- and RelB-dependent hypoxia responsive genes were specifically controlled under low oxygen, and not under basal conditions. Conversely, some of the cRel-dependent, hypoxia-upregulated genes were also altered in basal oxygen levels, indicating a broader mechanism.

Figure 3. Validation of NF-κB dependent hypoxia inducible gene expression changes in HeLa cells.
A. Log₂Fold change of selected NF-κB dependent hypoxia up- and down-regulated DEGs showing altered transcript levels in hypoxia compared to normoxia identified by RNA-seq (n=2). **B-F.** qPCR analysis in HeLa cells cultured at 21% oxygen (normoxia) or 24 h 1% oxygen (hypoxia), transfected with control siRNA or RelA, RelB or cRel siRNAs. RelA- (**B**), RelB- (**C**), and cRel-dependent (**D**) hypoxia upregulated DEGs. RelA- (**E**) and RelB-dependent (**F**) hypoxia downregulated DEGs. Relative mRNA expression levels of the indicated genes were analysed using 18S as a normalising gene. Graphs show mean (n=4) ± SEM, * P < 0.05, ** P < 0.01, *** P < 0.001. Statistical significance was determined via one-way ANOVA with post-hoc Dunnett's test.

NF-κB controlled hypoxia gene signatures in other cell types.

Given that we had uncovered a variety of genes that were dependent on NF-κB for the response to hypoxia in HeLa cells, we wanted to examine if these genes were also hypoxia-regulated in other cell types. To address this question, we analysed hypoxia (24 hours, 1% oxygen) RNA-

sequencing data in human cancer cell lines across different tissues, A549 (lung carcinoma), U87 (glioblastoma), HCT-116 (colorectal carcinoma), MCF-7 (breast adenocarcinoma), 501-mel (melanoma metastasis) and SKNAS (neuroblastoma) (Figure 4A; Sup. Table S2-3; Dataset S3). Following quality control and the analysis pipeline (Sup. Figure S9), volcano plots were created demonstrating the differentially expressed genes (DEGs) that were significantly changed (both upregulated or downregulated) across each cell line in response to hypoxia (Figure 4B). Comparing these hypoxia gene signatures across different cell backgrounds with our HeLa-derived, NF-*κ*B-dependent hypoxia signature revealed a very high overlap in cells such as U87, HCT-116 and MCF-7 and more modest in SKNAS cells, across all NF-*κ*B subunits (Figure 4C; Sup. Figure S10A). We also confirmed that a subset of these genes had been previously described as direct NF-*κ*B target genes in the Gilmore database (Figure 4D; Sup. Figure S10B). Finally, we were able to validate by qPCR analysis the vast majority of the NF-*κ*B dependent control over selected genes in U87, MCF-7 and A549 cells (Figure 5B-L; Sup. Figure S11A-F; Sup. Figure S12A-C; Sup. Figure S13A-Q; Sup. Table S1). Once again, RelB seemed to have the most widespread impact on hypoxia dependent gene changes across all the cell systems investigated (Figure 5B-L). Furthermore, some were also dependent on RelA, and very few were controlled by cRel (Figure 5B-L). Taken together these data indicate that the role of NF-*κ*B in controlling gene expression is common across a variety of cell lines. As a complementary approach we also analysed hypoxia responsive genes using wild type and IKKα/β double knockout HCT116 CRISPR cell lines (Prescott *et al.*, 2022). Six of the 9 genes we identified as NF-*κ*B-dependent, hypoxia upregulated genes in HeLa cells (RNA-seq and validated by qPCR - Sup. Figure S11D), were also hypoxia responsive in HCT116 cells and displayed NF-*κ*B dependence in the HCT116 CRISPR cell lines (Sup. Figure S14A-B). These results, in which NF-*κ*B was inhibited at the level of the IKKs, support our siRNA depletion analysis in a subset of identified NF-*κ*B-dependent, hypoxia-inducible genes, ruling out off target effects of the siRNA approach.

Figure 4. Identification of NF-κB dependent hypoxia inducible gene signature in different cell types.

A. RNA-seq (n=3) in the indicated cell types cultured at 21% oxygen (normoxia) or exposed to 24 h 1% oxygen (hypoxia). **B.** Volcano plots displaying differential expression analysis comparing 24h hypoxia to control from RNA-seq datasets in the indicated cell lines. **C.** Overlap of hypoxia inducible DEGs identified in different cell lines' RNA-seq datasets with NF-κB dependent hypoxia inducible DEGs identified in HeLa RNA-seq experiment following siRNA depletion of individual NF-κB subunits exposed or not to 24 h hypoxia. Percentage of NF-κB dependent hypoxia inducible DEGs are displayed **D.** Overlap of hypoxia inducible DEGs identified in different cell lines' RNA-seq datasets with the Gilmore laboratory's NF-κB target genes. Percentage of NF-κB target genes are displayed.

Figure 5. Analysis of hypoxia inducible NF-κB-dependent genes across various cell backgrounds.
A. Key for qPCR analysis graphs. **B-C.** Hypoxia upregulated NF-κB-dependent genes in U87 cells. **D.** Hypoxia downregulated NF-κB-dependent genes in U87 cells. **E-G.** Hypoxia upregulated NF-κB-dependent genes in

MCF-7 cells. **H-I.** Hypoxia downregulated NF-κB-dependent genes in MCF-7 cells. **J-K.** Hypoxia upregulated NF-κB-dependent genes in A549 cells. **L.** Hypoxia downregulated NF-κB-dependent genes in A549 cells. Relative mRNA expression levels of the indicated genes were analysed using Actin as a normalising gene for U87 and MCF-7 cell lines (**B-I**), and 18S for A549 cells (**J-L**). Graphs show mean (n=3) ± SEM, * P < 0.05, ** P < 0.01, *** P < 0.001. Statistical significance was determined via one-way ANOVA with post-hoc Dunnett's test.

Given that all our analysis had been done in cancer cells, it is possible that role of NF-κB in hypoxia we have uncovered was restricted to cancer cells. To begin to address this aspect, we analysed Human umbilical vein endothelial cells (HUVECs) a non-transformed cell line for which RNA-sequencing data was available in the NCBI Geo database (Tiana *et al.*, 2018). Although, the time of hypoxia exposure was less than the one we had previously used (16 h versus 24 h), these cells still mounted a robust hypoxia response at the transcriptional level (743 genes upregulated and 668 genes downregulated, Sup. Figure S15A). When compared to our HeLa cell data, HUVECs shared 232 upregulated genes but only 82 down regulated genes (Sup. Figure S15B-C). Comparison of the NF-κB dependent hypoxia-inducible gene from HeLa and HUVEC cells indicates that NF-κB is required for 45% of hypoxia repressed genes HUVECs share with HeLa cells, and 28% of hypoxia induced genes (Sup. Figure S15D-E). HUVECs have also previously been analysed for their response to TNF-α, a canonical NF-κB activating stimulus (Fowler *et al.*, 2022). Furthermore, this study also determined the role of RelA in HUVEC responses to TNF-α (Fowler *et al.*, 2022). Using these datasets, we made a series of observations. HUVECs are highly responsive, at a transcriptional level, to TNF-α, with 1640 genes upregulated after 4 h and 1445 genes upregulated after 10 h, 1078 of which were already upregulated after 4 h (Sup. Figure S16A, C, Dataset S4). Similarly, 1687 genes were down regulated after 4 h TNF-α and 1722 genes were downregulated after 10 h TNF-α, with 990 genes shared between the treatment periods (Sup. Figure S16B, D, Dataset S4). Depletion of RelA in this cell system revealed that it was needed for 40% of genes upregulated at 4 h and 49.6% of genes upregulated after 10 h TNF-α (Sup. Figure S16E). Similarly, RelA was needed for 43.2% of TNF-α downregulated genes after 4 h and 40.5% of downregulated genes after 10 h (Sup. Figure S16F). This indicates that RelA has a significant role in the response of HUVEC cells to TNF-α, controlling gene activation and repression. We next compared the RelA-dependent TNF-α regulated genes to our RelA-dependent hypoxia gene signature obtained from HeLa cells. This revealed that 14 of the RelA-dependent TNF-α-induced genes were also observed to be RelA dependent in hypoxia (Sup. Figure S16G). When we analysed the downregulated genes, 15 genes were dependent on RelA for their repression in both Hypoxia and TNF-α (Sup. Figure S16H). Interestingly, 28 genes that require RelA for

repression in TNF-α, require RelA for their induction in hypoxia, while 6 genes that required RelA for TNF-α induction required RelA for hypoxia repression. Taken together this data supports our model, while for some genes, hypoxia works similarly to a canonical inducer of NF-κB, more often hypoxia utilises NF-κB to achieve the opposite result in the control of gene expression, even in non-cancer cell types.

NF-κB is important for hypoxia induced cellular responses.

Given the extent of the NF-κB contribution to the hypoxia transcriptional response (35% of activated and 62% of repressed genes), we next determined if any of the cellular responses to hypoxia were altered in the absence of NF-κB subunits. Pathway analysis of the genes repressed in hypoxia that depend on NF-κB indicated Reactive Oxygen Species (ROS) and Oxidative phosphorylation as two of the main hallmarks altered. To validate this at the cellular level, we measured ROS production using a molecular dye (CellRox); we have previously used this to demonstrate that hypoxia induces ROS generation in these cells (Batie *et al*, 2019) as described in other systems (Alva *et al*, 2024). As expected, hypoxia exposure resulted in a significant increase in ROS (Figure 6A-B). Importantly, depletion of RelA, and to a lesser extent RelB and cRel, reduced the ROS levels generated by hypoxia (Figure 6A-B). As this ROS probe primarily detects superoxide in cells (McBee *et al*, 2017), we investigated genes specifically associated with detoxifying superoxide in cells. One such gene is superoxide dismutase 1, SOD1. Although, we could not demonstrate a statistically significant change in levels of mRNA for SOD1 when NF-κB subunits were depleted in hypoxia (Sup. Figure S6G-H), Western blot analysis confirmed that hypoxia does lead to reduced SOD1 expression (Figure 6C-D), and that RelB, and to lesser extent cRel, relieved this repression when depleted (Figure 6C-D). These results suggest that ROS generation in hypoxia requires NF-κB via a mechanism that partially requires SOD1 reduction. We also investigated if NRF2 is induced in hypoxia as marker of oxidative stress (Sup. Figure S17). NRF2 levels were unchanged in response to 1 and 24 hours of 1% oxygen exposure (Sup. Figure S17A) and only a small number of NRF2 target genes (Hayes & Dinkova-Kostova, 2014; Morgenstern *et al.*, 2024) were upregulated in hypoxia from our RNA-seq analysis (Sup. Figure S17B), thus there is limited evidence for induction of NRF2, at least at the time and level of hypoxia exposure employed here, suggesting the level of ROS induced by hypoxia may be important for signalling but may not activate the NRF2 stress response.

Similarly, we investigated a panel of proteins involved in oxidative phosphorylation (Figure 7, Sup. Figure S18). This panel contained antibodies for proteins such as ATP Synthase F1 subunit alpha (ATP5A); Ubiquinol-cytochrome C reductase core protein 2 (UQCRC2), Succinate Dehydrogenase subunit B (SDHB) and Isocitrate Dehydrogenase 1 (IDH1). Hypoxia led to reductions in several proteins represented in this panel, such as ATP5A, UQCRC2, NDUFB8, COX2 and IDH1 (Figure 7A-B, Sup. Figure S18A-E). Depletion of NF-κB led to reversal of the protein levels to those of normoxia or slightly above for IDH1 (Figure 7A-B, Sup. Figure S18A-E). These data support our RNA-seq analysis, where oxidative phosphorylation was a hallmark that was significantly repressed in hypoxia, dependent on NF-κB presence (Figure 2G). To confirm that hypoxia-dependent repression of protein levels of oxidative phosphorylation pathway components, we also examined HCT116 and their IKKα/IKKβ CRISPR knockout derivatives (Prescott *et al.*, 2022)(Figure 7C). In these cells, where NF-kB signalling is abolished, hypoxia exposure did not result in reduced levels of ATP5A, UQCRC2, COX2 and NDUFB8 (Figure 7C). Only one of the IKKα/IKKβ KO clones reversed SOD1 repression in hypoxia (Figure 7C).

NF-κB was previously shown to actively repress gene expression via HDAC activity (Campbell *et al*, 2004; Elsharkawy *et al*, 2010; Rocha *et al*, 2003). To test if some of NF-κB-repressed genes also require HDAC activity, we treated HeLa cells with an HDAC inhibitor, Saha, alone or in combination with hypoxia. This analysis revealed that while HDAC inhibition reversed hypoxia induced repression of ATP5A, UQCRC2 and COX2 (Figure 7D), it had no effect on hypoxia-induced repression of SOD1 (Sup. Figure S19), causing SOD1 levels to reduce in normoxia. Taken together, these results suggest that NF-κB represses gene expression in hypoxia by mechanisms requiring HDAC activity and additional unknown repressors.

Figure 6. NF-κB is required for ROS generation in hypoxia.
A. Cellular ROS measurements using CellRox staining and immunofluorescence analysis in HeLa cells, cultured at 21% oxygen (control) or 24 h 1% oxygen (hypoxia), transfected with control siRNA or RelA, RelB or cRel siRNAs. Representative images from 3 independent experiments are shown. **B.** Quantification of relative ROS signal in individual cells from 3 biological replicates, represented as a beeswarm with box plot, * P < 0.05, ** P < 0.01, *** P < 0.001 (compared to hypoxia control siRNA treated cells). Statistical significance was determined via one-way ANOVA with post-hoc Dunnett's test. **C.** Immunoblot analysis of the indicated protein in HeLa cells exposed or not to 1% oxygen for 24 h, with siRNA transfection of control, RelA, RelB or cRel. **D.** Signal intensity normalised to actin (mean n=3, ± SEM) from immunoblot analysis in (C). * P < 0.05, ** P < 0.01, *** P < 0.001 (compared to hypoxia control siRNA treated cells). Statistical significance was determined via one-way ANOVA with post-hoc Dunnett's test.

Figure 7. NF-κB is required for expression of oxidative phosphorylation related proteins in hypoxia.
A and B. Immunoblot analysis of the indicated proteins in HeLa cells cultured at 21% oxygen (control) or 24 h 1% oxygen (hypoxia), transfected with control siRNA or RelA, RelB or cRel siRNAs. Representative images

from 3 independent experiments are shown. **C.** HCT-116 wildtype (WT) and IKKα/IKKβ double knockout cells, were cultured at 21% oxygen (control) or 24 h 1% oxygen (hypoxia) prior to cell lysis. Western blot was performed with the indicated antibodies. **D.** Immunoblot analysis of the indicated proteins in HeLa cells cultured at 21% oxygen (control) or 24 h 1% oxygen (hypoxia), with or without 2.5 μM Saha (HDAC inhibitor) for 24 h. Representative images from 3 independent experiments are shown.

Discussion

Here, we have investigated the contribution NF-κB makes to the transcriptional response to hypoxia. We find that NF-κB is required for induction of genes in hypoxia but seems especially important for gene repression under these conditions (Figure 8).

Figure 8. NF-κB is a crucial transcription factor in hypoxia.
Upon exposure to hypoxia, or reduced oxygen availability, NF-κB associates with co-activators to induce gene expression. However, NF-κB can also associate with co-repressors, leading to significant repression of gene expression under hypoxic conditions

NF-κB pathway activation in hypoxia has been well documented across cells, tissues and organisms (Bandarra *et al.*, 2014; Chandel *et al.*, 2000; Cummins & Taylor, 2005; D'Ignazio & Rocha, 2016; Koong *et al.*, 1994; Taylor *et al.*, 1999; Van Welden *et al*, 2017). While the exact mechanism of activation may depend on the system and timing of the investigation, it was implied that NF-κB is part of the cellular response to hypoxia. However, to what extent NF-κB contributes to the transcriptional response to hypoxia was not known. Our understanding of hypoxia is dominated by the HIFs, whose role as a master regulator of the transcriptional response is well established from direct chromatin binding (Batie *et al*, 2022), to a conserved list of genes consistently induced in hypoxia (Puente-Santamaria *et al.*, 2022). Our analysis of depletion of individual Rel subunits reveals that NF-κB is required for 35% of all hypoxia-induced genes but strikingly is required for over 60% of all hypoxia-repressed genes. Interestingly, RelB seems as important as RelA, something previously not appreciated in the response to hypoxia. Very few studies have investigated RelB's role in hypoxia (Patel *et al.*, 2017; Riedl *et al*, 2021). These studies have mostly focused on either gene induction (Patel *et al.*, 2017) or used a hypoxia mimetic as inducer (Riedl *et al.*, 2021). None of these studies investigated RelB-mediated gene regulation in hypoxia, making our study one of the first to use an unbiased approach to analyse the contribution of RelB to the transcriptional profile of cells exposed to hypoxia. Of note, RelA is able to directly regulate the expression of other NF-κB subunits directly (Oeckinghaus & Ghosh, 2009), so the effects of RelA may represent a contributions of other NF-κB subunits as well. Interestingly, NF-κB-dependent hypoxia signatures are more widely regulated across cell types than validated NF-κB targets, suggesting that perhaps NF-κB-dependent genes in hypoxia are different from standard cytokine induced ones. Although analysis of the ChIP-Atlas database, confirmed that 80% of the genes we have identified to be NF-κB dependent have NF-κB subunits bound to them, a similar NF-κB binding percentage was present in genes found to be independent of NF-κB. Further analysis performing ChIP-seq or CUT&RUN in hypoxia for NF-κB subunits would help distinguish between direct and indirect regulation of genes by NF-κB in hypoxia. It is also important to acknowledge that NF-κB is critical to maintain HIF gene expression in different cells and organisms suggesting a wider role for NF-κB in underpinning HIF activity (D'Ignazio & Rocha, 2016; Kenneth *et al*, 2009; Rius *et al*, 2008; van Uden *et al*, 2008; van Uden *et al*, 2011). Interestingly, both transcription factors are evolutionarily conserved and present in the primitive organism *Trichoplax adhaerens* (Loenarz *et al*, 2011; Romanova & Moroz, 2024; Williams & Gilmore, 2020). Although their functional crosstalk has not been investigated in *T.*

adhaerens, it is conserved in *Drosophila melanogaster* (Bandarra *et al.*, 2014; Bandarra *et al*, 2015; van Uden *et al.*, 2011), indicating an ancient association between these two pathways.

In our analysis in HeLa cells, individual depletion of NF-κB subunits only resulted in around 10-20% reduction in HIF-1α mRNA (Sup. Figure S20) and no visible changes to HIF-1α protein levels after 24 hours hypoxia exposure. This supports our original findings that more than one NF-κB subunit is required to control HIF-1α levels (van Uden *et al.*, 2008). Furthermore, it is possible that NF-κBs control over HIF-1α is observed at earlier response times of hypoxia, as after 24h of hypoxia HIF-1α protein levels are already reduced in most cell types. Nevertheless, for genes known to be HIF-dependent, NF-κB's function could be indirect. This is also supported by the analysis of HIF ChIP-seq data where there was a positive correlation between HIF binding and upregulation of genes that are also NF-κB dependent. However, a more thorough investigation of the interplay between HIF and NF-κB, across a range of timepoints in response to hypoxia, is required.

Unlike HIF, NF-κB is able to actively repress genes (Burkhart *et al*, 2005; Campbell *et al.*, 2004; Elsharkawy *et al.*, 2010; Grigoriadis *et al*, 1996; Saha *et al*, 2008) by actively recruiting repressor proteins such as HDACs (Elsharkawy *et al.*, 2010), or by engaging with repressor complexes such as EZH2 directly (Dardis *et al*, 2023) or via nuclear sumoylated-IκBα (Mulero *et al*, 2013). These effects have been observed in response to a variety of stimuli or in developmental stages of differentiation (Burkhart *et al.*, 2005; Campbell *et al.*, 2004; Elsharkawy *et al.*, 2010; Grigoriadis *et al.*, 1996; Mulero *et al.*, 2013; Saha *et al.*, 2008). However, the role NF-κB in gene repression is still under investigated, despite its importance when studied (Campbell *et al.*, 2004; Grigoriadis *et al.*, 1996; Mulero *et al.*, 2013). Our results reveal the extent of NF-κB contribution to the hypoxia transcriptional response, highlights the importance of NF-κB during such a critical developmental and pathological stimulus. Our results suggest that NF-κB may repress gene expression via HDAC activity. However, HDAC inhibition did not reverse hypoxia induced repression for all the proteins we investigated. In the future, it will be important to determine the exact mechanism by which NF-κB mediates its repressive effects in hypoxia as our results indicate that HDACs are only a part of the mechanism.

The hallmark signatures for hypoxia repressed genes controlled by NF-κB were ROS signalling and Oxidative Phosphorylation. Although distinct, these signatures are intrinsically linked in hypoxia (Hamanaka & Chandel, 2010; Solaini *et al*, 2010). Given the reduction in oxygen,

mitochondria are considered one of the sources of ROS due to alteration in the electron transport chain activity (Alva *et al.*, 2024; Kracht *et al*, 2020; Solaini *et al.*, 2010). Furthermore, reducing the expression of genes associated with oxidative phosphorylation in time of oxygen deficiency is a cost-effective mechanism to conserve energy (Kracht *et al.*, 2020). Although technically we are unable to measure oxygen consumption in hypoxia, we could determine that several proteins associated with this function were reduced in hypoxia and that NF-κB depletion reversed this repression. This might be, in the long term, detrimental for the cell, as cell death could be a consequence of prolonged NF-κB depletion in hypoxia.

NF-κB has been previously shown to translocate to mitochondria as well (Ivanova & Perkins, 2019), a process important for ROS generation. Our data supports NF-κB's role in generation of ROS in response to hypoxia. ROS's role in hypoxia is multifaceted (Alva *et al.*, 2024; Hamanaka & Chandel, 2010), and hence it contributes not only to pathology but is also used as a signalling molecule. However, these roles seem to be quite dependent on the extent of ROS generation cell type-specific (Alva *et al.*, 2024).

Our study has several limitations to consider. For this study we only used cancer cells and given NF-κB's pleotropic effects, it is possible that in non-transformed cells, NF-κB has a different role. We analysed a non-transformed cell type, HUVECs, with available transcriptomics in hypoxia (Tiana *et al.*, 2018) and a known NF-κB inducer, TNF-α (Fowler *et al.*, 2022). This analysis also correlated well with our cancer cell analysis, revealing that NF-κB is important for repression of genes in hypoxia but also for some of the hypoxia induced genes. Another limitation of our study is that we only assessed the role of NF-κB in hypoxia and not at basal conditions. Thus, it is possible that some of the genes we have demonstrated to be regulated in hypoxia could also be similarly controlled in normoxic conditions. However, qPCR analysis revealed that this is not the case. Similarly, analysis of HUVECs revealed that only a subset of TNF-α induced/repressed genes are also controlled by RelA under non-stimulated conditions. Another limitation is the use of only one siRNA sequence per NF-κB subunit. Whilst were careful to validate and quantify NF-κB subunit knockdown in each case we also validated our results with additional orthogonal approaches to inhibit NF-κB subunit such as the IKKα/IKKβ CRISPR knockout cells. Nonetheless, it is still possible that some off target effects are present; however, this is the same with any experimental approach. Despite these limitations, our study uncovers a pivotal role for NF-κB in regulating the transcriptional response to hypoxia. Strikingly, we find that NF-κB's primary function under these conditions is to mediate transcriptional repression (Figure 8), a contribution that has been largely underappreciated.

While previous studies have highlighted the importance of this mechanism (Batie *et al.*, 2018; Cavadas *et al.*, 2017), our understanding of how gene repression is orchestrated in hypoxia remains limited. This work takes a critical step toward filling that gap.

Data Availability

The HeLa NF-κB depletion in response to hypoxia RNA-seq dataset is accessible at NCBI GEO: GSE260616. The SKNAS hypoxia stimulation RNA-seq dataset is accessible at NCBI GEO: GSE284326.

Acknowledgements

We would like to acknowledge the members of Rocha and Kenneth labs for helpful discussions and the member of the hypoxia and NF-kappaB community for sharing their RNA-seq data. We thank Dr. Marco Sciacovelli (University of Liverpool) for providing the OxPhos antibody set and Dr. Dom Byrne for valuable reagents. We acknowledge the Liverpool Centre for Cell Imaging (CCI) for provision of imaging equipment, in particular the Epifluorescent microscope funded by MRC grant number MR/K015931/1, OMERO and technical assistance. This work was funded by a Wellcome trust collaborator grant to SR (206293/Z/17/Z) and the University of Liverpool, CSK is supported by a fellowship from the Royal College of Surgeons in England (RCSEng), a Tung Postgraduate fellowship and Alder Hey Children's Charity. NKS is supported by Worldwide Cancer Research [grant number 24-0353].

Competing interests

The authors declare no competing interests.

Author contributions

DS and SR conceptualised and oversaw the project. DS, MB and CSK performed the RNA seq analysis. DS performed most of the experiments and analysed the data associated with it. MB and SR performed some of the validation experiments. SC provided the IKK CRISPR HCT-116 cells. DS, MB, NSK, SC and SR wrote the manuscript. SR obtained the funding.

References:

Adhikari A, Xu M, Chen ZJ (2007) Ubiquitin-mediated activation of TAK1 and IKK. *Oncogene* 26: 3214-3226

Al-Anazi A, Parhar R, Saleh S, Al-Hijailan R, Inglis A, Al-Jufan M, Bazzi M, Hashmi S, Conca W, Collison K *et al* (2018) Data on hypoxia-induced VEGF, leptin and NF-kB p65 expression. *Data Brief* 21: 2395-2397

Allan C, Burel JM, Moore J, Blackburn C, Linkert M, Loynton S, Macdonald D, Moore WJ, Neves C, Patterson A *et al* (2012) OMERO: flexible, model-driven data management for experimental biology. *Nat Methods* 9: 245-253

Almaden JV, Liu YC, Yang E, Otero DC, Birnbaum H, Davis-Turak J, Asagiri M, David M, Goldrath AW, Hoffmann A (2016) B-cell survival and development controlled by the coordination of NF-kappaB family members RelB and cRel. *Blood* 127: 1276-1286

Alva R, Wiebe JE, Stuart JA (2024) Revisiting reactive oxygen species production in hypoxia. *Pflugers Arch* 476: 1423-1444

An J, Rettig MB (2005) Mechanism of von Hippel-Lindau protein-mediated suppression of nuclear factor kappa B activity. *Mol Cell Biol* 25: 7546-7556

Bandarra D, Biddlestone J, Mudie S, Muller HA, Rocha S (2014) Hypoxia activates IKK-NF-kappaB and the immune response in Drosophila melanogaster. *Biosci Rep* 34

Bandarra D, Biddlestone J, Mudie S, Muller HA, Rocha S (2015) HIF-1alpha restricts NF-kappaB-dependent gene expression to control innate immunity signals. *Dis Model Mech* 8: 169-181

Batie M, Del Peso L, Rocha S (2018) Hypoxia and Chromatin: A Focus on Transcriptional Repression Mechanisms. *Biomedicines* 6

Batie M, Frost J, Frost M, Wilson JW, Schofield P, Rocha S (2019) Hypoxia induces rapid changes to histone methylation and reprograms chromatin. *Science* 363: 1222-1226

Batie M, Kenneth NS, Rocha S (2022) Systems approaches to understand oxygen sensing: how multi-omics has driven advances in understanding oxygen-based signalling. *Biochem J* 479: 245-257

Bruegge K, Jelkmann W, Metzen E (2007) Hydroxylation of hypoxia-inducible transcription factors and chemical compounds targeting the HIF-alpha hydroxylases. *Curr Med Chem* 14: 1853-1862

Burkhart BA, Hebbar PB, Trotter KW, Archer TK (2005) Chromatin-dependent E1A activity modulates NF-kappaB RelA-mediated repression of glucocorticoid receptor-dependent transcription. *J Biol Chem* 280: 6349-6358

Campbell KJ, Rocha S, Perkins ND (2004) Active repression of antiapoptotic gene expression by RelA(p65) NF-kappa B. *Mol Cell* 13: 853-865

Cavadas MAS, Cheong A, Taylor CT (2017) The regulation of transcriptional repression in hypoxia. *Exp Cell Res* 356: 173-181

Chandel NS, Trzyna WC, McClintock DS, Schumacker PT (2000) Role of oxidants in NF-kappa B activation and TNF-alpha gene transcription induced by hypoxia and endotoxin. *J Immunol* 165: 1013-1021

Culver C, Sundqvist A, Mudie S, Melvin A, Xirodimas D, Rocha S (2010) Mechanism of hypoxia-induced NF-kappaB. *Mol Cell Biol* 30: 4901-4921

Cummins EP, Taylor CT (2005) Hypoxia-responsive transcription factors. *Pflugers Arch* 450: 363-371

D'Ignazio L, Batie M, Rocha S (2017) Hypoxia and Inflammation in Cancer, Focus on HIF and NF-kappaB. *Biomedicines* 5

D'Ignazio L, Rocha S (2016) Hypoxia Induced NF-kappaB. *Cells* 5

Dardis GJ, Wang J, Simon JM, Wang GG, Baldwin AS (2023) An EZH2-NF-kappaB regulatory axis drives expression of pro-oncogenic gene signatures in triple negative breast cancer. *iScience* 26: 107115

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21

Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W (2005) BioMart and Bioconductor: a powerful link between biological databases and microarray data analysis. *Bioinformatics* 21: 3439-3440

Edgar R, Domrachev M, Lash AE (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* 30: 207-210

Elsharkawy AM, Oakley F, Lin F, Packham G, Mann DA, Mann J (2010) The NF-kappaB p50:p50:HDAC-1 repressor complex orchestrates transcriptional inhibition of multiple pro-inflammatory genes. *J Hepatol* 53: 519-527

Fandrey J, Gorr TA, Gassmann M (2006) Regulating cellular oxygen sensing by hydroxylation. *Cardiovasc Res* 71: 642-651

Fowler JWM, Zhang R, Tao B, Boutagy NE, Sessa WC (2022) Inflammatory stress signaling via NF-kB alters accessible cholesterol to upregulate SREBP2 transcriptional activity in endothelial cells. *Elife* 11

Gerondakis S, Siebenlist U (2010) Roles of the NF-kappaB pathway in lymphocyte development and function. *Cold Spring Harb Perspect Biol* 2: a000182

Grigoriadis G, Zhan Y, Grumont RJ, Metcalf D, Handman E, Cheers C, Gerondakis S (1996) The Rel subunit of NF-kappaB-like transcription factors is a positive and negative regulator of macrophage gene expression: distinct roles for Rel in different macrophage populations. *EMBO J* 15: 7099-7107

Hamanaka RB, Chandel NS (2010) Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes. *Trends Biochem Sci* 35: 505-513

Hayes JD, Dinkova-Kostova AT (2014) The Nrf2 regulatory network provides an interface between redox and intermediary metabolism. *Trends Biochem Sci* 39: 199-218

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* 38: 576-589

Iqbal AD, C.; Metge, F.; Rosskopp, D.; Boucas, J. (2021) Flaski.

Ivanova IG, Perkins ND (2019) Hypoxia induces rapid, STAT3 and ROS dependent, mitochondrial translocation of RelA(p65) and IkappaBalpha. *Biosci Rep* 39

Kaelin WG, Jr., Ratcliffe PJ (2008) Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* 30: 393-402

Kenneth NS, Mudie S, van Uden P, Rocha S (2009) SWI/SNF regulates the cellular response to hypoxia. *J Biol Chem* 284: 4123-4131

Kenneth NS, Rocha S (2008) Regulation of gene expression by hypoxia. *Biochem J* 414: 19-29

Koong AC, Chen EY, Giaccia AJ (1994) Hypoxia causes the activation of nuclear factor kappa B through the phosphorylation of I kappa B alpha on tyrosine residues. *Cancer Res* 54: 1425-1430

Kracht M, Muller-Ladner U, Schmitz ML (2020) Mutual regulation of metabolic processes and proinflammatory NF-kappaB signaling. *J Allergy Clin Immunol* 146: 694-705

Liao Y, Smyth GK, Shi W (2013) The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res* 41: e108

Liao Y, Wang J, Jaehnig EJ, Shi Z, Zhang B (2019) WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res* 47: W199-W205

Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P (2015) The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* 1: 417-425

Liu T, Zhang L, Joo D, Sun SC (2017) NF-kappaB signaling in inflammation. *Signal Transduct Target Ther* 2: 17023-

Loenarz C, Coleman ML, Boleininger A, Schierwater B, Holland PW, Ratcliffe PJ, Schofield CJ (2011) The hypoxia-inducible transcription factor pathway regulates oxygen sensing in the simplest animal, Trichoplax adhaerens. *EMBO Rep* 12: 63-70

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550

McBee ME, Chionh YH, Sharaf ML, Ho P, Cai MW, Dedon PC (2017) Production of Superoxide in Bacteria Is Stress- and Cell State-Dependent: A Gating-Optimized Flow Cytometry Method that Minimizes ROS Measurement Artifacts with Fluorescent Dyes. *Front Microbiol* 8: 459

Melvin A, Mudie S, Rocha S (2011) Further insights into the mechanism of hypoxia-induced NFkappaB. [corrected]. *Cell Cycle* 10: 879-882

Morgenstern C, Lastres-Becker I, Demirdogen BC, Costa VM, Daiber A, Foresti R, Motterlini R, Kalyoncu S, Arioz BI, Genc S *et al* (2024) Biomarkers of NRF2 signalling: Current status and future challenges. *Redox Biol* 72: 103134

Mulero MC, Ferres-Marco D, Islam A, Margalef P, Pecoraro M, Toll A, Drechsel N, Charneco C, Davis S, Bellora N *et al* (2013) Chromatin-bound IkappaBalpha regulates a subset of polycomb target genes in differentiation and cancer. *Cancer Cell* 24: 151-166

Oeckinghaus A, Ghosh S (2009) The NF-kappaB family of transcription factors and its regulation. *Cold Spring Harb Perspect Biol* 1: a000034

Ortmann BM, Burrows N, Lobb IT, Arnaiz E, Wit N, Bailey PSJ, Jordon LH, Lombardi O, Penalver A, McCaffrey J *et al* (2021) The HIF complex recruits the histone methyltransferase SET1B to activate specific hypoxia-inducible genes. *Nat Genet* 53: 1022-1035

Patel H, Zaghloul N, Lin K, Liu SF, Miller EJ, Ahmed M (2017) Hypoxia-induced activation of specific members of the NF-kB family and its relevance to pulmonary vascular remodeling. *Int J Biochem Cell Biol* 92: 141-147

Perkins ND (2007) Integrating cell-signalling pathways with NF-kappaB and IKK function. *Nat Rev Mol Cell Biol* 8: 49-62

Perkins ND, Gilmore TD (2006) Good cop, bad cop: the different faces of NF-kappaB. *Cell Death Differ* 13: 759-772

Prescott JA, Balmanno K, Mitchell JP, Okkenhaug H, Cook SJ (2022) IKKalpha plays a major role in canonical NF-kappaB signalling in colorectal cells. *Biochem J* 479: 305-325

Prescott JA, Mitchell JP, Cook SJ (2021) Inhibitory feedback control of NF-kappaB signalling in health and disease. *Biochem J* 478: 2619-2664

Puente-Santamaria L, Sanchez-Gonzalez L, Pescador N, Martinez-Costa O, Ramos-Ruiz R, Del Peso L (2022) Formal Meta-Analysis of Hypoxic Gene Expression Profiles Reveals a Universal Gene Signature. *Biomedicines* 10

Qin ZH, Tao LY, Chen X (2007) Dual roles of NF-kappaB in cell survival and implications of NF-kappaB inhibitors in neuroprotective therapy. *Acta Pharmacol Sin* 28: 1859-1872

Riedl T, Faure-Dupuy S, Rolland M, Schuehle S, Hizir Z, Calderazzo S, Zhuang X, Wettengel J, Lopez MA, Barnault R *et al* (2021) Hypoxia-Inducible Factor 1 Alpha-Mediated RelB/APOBEC3B Down-regulation Allows Hepatitis B Virus Persistence. *Hepatology* 74: 1766-1781

Rius J, Guma M, Schachtrup C, Akassoglou K, Zinkernagel AS, Nizet V, Johnson RS, Haddad GG, Karin M (2008) NF-kappaB links innate immunity to the hypoxic response through transcriptional regulation of HIF-1alpha. *Nature* 453: 807-811

Rocha S (2007) Gene regulation under low oxygen: holding your breath for transcription. *Trends Biochem Sci* 32: 389-397

Rocha S, Martin AM, Meek DW, Perkins ND (2003) p53 represses cyclin D1 transcription through down regulation of Bcl-3 and inducing increased association of the p52 NF-kappaB subunit with histone deacetylase 1. *Mol Cell Biol* 23: 4713-4727

Romanova DY, Moroz LL (2024) The ancestral architecture of the immune system in simplest animals. *Front Immunol* 15: 1529836

Saha A, Hammond CE, Trojanowska M, Smolka AJ (2008) Helicobacter pylori-induced H,K-ATPase alpha-subunit gene repression is mediated by NF-kappaB p50 homodimer promoter binding. *Am J Physiol Gastrointest Liver Physiol* 294: G795-807

Solaini G, Baracca A, Lenaz G, Sgarbi G (2010) Hypoxia and mitochondrial oxidative metabolism. *Biochim Biophys Acta* 1797: 1171-1177

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES *et al* (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 102: 15545-15550

Taylor CT, Fueki N, Agah A, Hershberg RM, Colgan SP (1999) Critical role of cAMP response element binding protein expression in hypoxia-elicited induction of epithelial tumor necrosis factor-alpha. *J Biol Chem* 274: 19447-19454

Tian B, Nowak DE, Brasier AR (2005) A TNF-induced gene expression program under oscillatory NF-kappaB control. *BMC Genomics* 6: 137

Tiana M, Acosta-Iborra B, Puente-Santamaria L, Hernansanz-Agustin P, Worsley-Hunt R, Masson N, Garcia-Rio F, Mole D, Ratcliffe P, Wasserman WW *et al* (2018) The SIN3A histone deacetylase complex is required for a complete transcriptional response to hypoxia. *Nucleic Acids Res* 46: 120-133

van Uden P, Kenneth NS, Rocha S (2008) Regulation of hypoxia-inducible factor-1alpha by NF-kappaB. *Biochem J* 412: 477-484

van Uden P, Kenneth NS, Webster R, Muller HA, Mudie S, Rocha S (2011) Evolutionary conserved regulation of HIF-1beta by NF-kappaB. *PLoS Genet* 7: e1001285

Van Welden S, Selfridge AC, Hindryckx P (2017) Intestinal hypoxia and hypoxia-induced signalling as therapeutic targets for IBD. *Nat Rev Gastroenterol Hepatol* 14: 596-611

Volcic M, Karl S, Baumann B, Salles D, Daniel P, Fulda S, Wiesmuller L (2012) NF-kappaB regulates DNA double-strand break repair in conjunction with BRCA1-CtIP complexes. *Nucleic Acids Res* 40: 181-195

Williams LM, Gilmore TD (2020) Looking Down on NF-kappaB. *Mol Cell Biol* 40

Wilson JW, Shakir D, Batie M, Frost M, Rocha S (2020) Oxygen-sensing mechanisms in cells. *FEBS J* 287: 3888-3906

Xia ZB, Meng FR, Fang YX, Wu X, Zhang CW, Liu Y, Liu D, Li GQ, Feng FB, Qiu HY (2018) Inhibition of NF-kappaB signaling pathway induces apoptosis and suppresses proliferation and angiogenesis of human fibroblast-like synovial cells in rheumatoid arthritis. *Medicine (Baltimore)* 97: e10920

Xie H, Simon MC (2017) Oxygen availability and metabolic reprogramming in cancer. *J Biol Chem* 292: 16825-16832

Xu RX, Liu RY, Wu CM, Zhao YS, Li Y, Yao YQ, Xu YH (2015) DNA damage-induced NF-kappaB activation in human glioblastoma cells promotes miR-181b expression and cell proliferation. *Cell Physiol Biochem* 35: 913-925

Zou Z, Ohta T, Oki S (2024) ChIP-Atlas 3.0: a data-mining suite to explore chromosome architecture together with large-scale regulome data. *Nucleic Acids Res* 52: W45-W53