Pathogen and pest communities in agroecosystems across climate gradients: Anticipating future challenges in the highland tropics

Romaric A. Mouafo-Tchinda[1,2,3]*, Aaron I. Plex Sulá[1,2,3], Berea A. Etherton[1,2,3], Joshua S. Okonya[4], Gloria Valentine Nakato[5], Yanru Xing[1,2,3], Jacobo Robledo[1,2,3], Ashish Adhikari[1,2,3], Guy Blomme[6], Déo Kantungeko[7], Anastase Nduwayezu[8], Jan F. Kreuze[9], Jürgen Kroschel[9,10], James P. Legg[11], and Karen A. Garrett[1,2,3]*

[1] Plant Pathology Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL, USA

[2] Global Food Systems Institute, University of Florida, Gainesville, FL, USA

[3] Emerging Pathogens Institute, University of Florida, Gainesville, FL, USA

[4] Association for Strengthening Agricultural Research in Eastern and Central Africa, Entebbe, Uganda

[5] International Institute of Tropical Agriculture, Arusha, Tanzania

[6] Bioversity International, Addis Ababa, Ethiopia

[7] International Institute of Tropical Agriculture, Bujumbura, Burundi

[8] Rwanda Agriculture Development and Animal Resources Board (RAB), Ruhengeri, Rwanda

[9] International Potato Center (CIP), Lima, Peru

[10] Hans-Ruthenberg-Institute for Tropical Agricultural Sciences, University of Hohenheim, Stuttgart, Germany

[11] International Institute of Tropical Agriculture, Dar es Salaam, Tanzania

* Corresponding authors

Emails: rmouafotchinda@ufl.edu*, karengarrett@ufl.edu*

ABSTRACT

CONTEXT: Tropical cropping systems must adapt to the current and future geographic distribution of pathogen and pest communities. An important research gap is how climate change may shift the distribution of pathogens and pests in tropical lowlands and highlands.

OBJECTIVE: We evaluated the current geographic risk of 27 pathogens and pests in the production of four food security crops (banana, cassava, potato, and sweetpotato) in the Great Lakes region of Africa, and the potential future risk under climate change. Models for each pathogen and pest indicate the potential for changes in geographic distribution, with model fit indicating the potential for decision support systems to facilitate management.

METHODS: First, cropland connectivity analysis identified locations likely important in the spread of crop-specific pathogens and pests, such as locations in Rwanda and Burundi. Second, we surveyed the 27 economically important pathogens and pests in Rwanda and Burundi, mapping the distribution of each across climate gradients and quantifying patterns of association. Third, we used machine learning to develop models of each species as a function of environmental variables, including host landscape variables. We also evaluated the increase in temperature across altitudes under future climate change scenarios in this region.

RESULTS AND CONCLUSIONS: Among the ten machine-learning algorithms evaluated, random forests and support vector machines generally performed best for predicting severity and infestation. Host landscape variables were useful predictors for some species. Based on climate matching, 44% of the pathogens and pests could become more common with warmer temperatures at higher altitudes, while 17% may become less common.

SIGNIFICANCE: This study indicates adaptation priorities for crop health in a region with multiple challenges to agricultural sustainability. The models developed here also indicate which species may have more potential and relevance for future development

54   of pathogen and pest forecasts.

56   **Keywords:** Climate matching | crop pathogens and pests | cropland connectivity | food
57   security | machine learning | vegetatively propagated crops

1. Introduction

1.1. Impact of climate and crop landscape structure on plant pathogens and pests

Agroecosystems in the tropics, such as the Great Lakes region of Africa, are often challenged by climate variability and societal dynamics including demographic pressures, socio-political instability, and resource-related conflict (Ogot and Niane, 1984; Ong'ayo, 2018). Climate change is likely to affect agricultural productivity and exacerbate the proliferation of some pathogens and pests in this region (Ebregt *et al.*, 2005; Ferdu and Solbreck, 2010; Okonya and Kroschel, 2013; Gallego‑Tévar *et al.*, 2024). Ecosystems can be reshaped by new temperature and precipitation patterns, driving pathogens and pests to new locations (Sparks *et al.*, 2014; Cilas *et al.*, 2016; Garrett *et al.*, 2022a; Garrett *et al.*, 2022b). The interactions among climate heterogeneity, the geographic structure of agroecosystems, and pathogens and pests in tropical regions such as the Great Lakes region represent a major knowledge gap. Research has examined how climate affects the dynamics of specific pathogens and pests (Kroschel *et al.*, 2014; Adhikari *et al.*, 2015; Blomme *et al.*, 2020). However, field studies that allow for simultaneous comparisons of many different pathogens and pests along climate gradients are rare, and little is known about how regional factors, such as the geographic structure of agroecoystems in the Great Lakes region, exacerbate risks. It is an open question how pathogen and pest communities are distributed across altitudes in this region, what variables are predictive of pathogen and pest community structure, and how their distribution may shift under climate change.

Climate matching studies identify which locations have similar climate conditions, such that they may be equally suitable for a species; ecological communities may move following changes in where suitable climate conditions are found in the future (Colwell *et al.*, 2008; Singh *et al.*, 2023; Chevalier *et al.*, 2024). For example, pathogen and pest communities now at hotter, lower altitudes may move to what are currently cooler, higher altitudes if there is warming due to climate change. The dynamics of pathogens and pests in response to climate have been widely studied (Hodkinson, 2005; Pautasso *et al.*, 2012; Juroszek *et al.*, 2022; Mishra *et al.*, 2024). Higher altitudes are commonly

associated with lower species richness, disease severity, and pest damage (Hodkinson,

2005; Blomme *et al.*, 2020). Since altitude is typically negatively correlated with

temperature (Poveda *et al.*, 2012), altitude can be a useful proxy when evaluating the

effect of temperature on plant disease and pest damage. For example, previous studies

in South America and East Africa predicted that rising temperatures due to climate

change will modify the geographic distribution of pathogens and pests, increasing plant

disease severity or pest damage at higher altitudes (Poveda *et al.*, 2012; Blomme *et al.*,

2020). New climate conditions favorable to certain pathogens and pests are likely to

occur at higher altitudes, disrupting ecological balances and exposing higher-altitude

plants and ecosystems to new risks.

The geographic distribution of crop production can also influence geographic

patterns of disease severity and pest damage. Crop landscape structure has been well

documented to influence crop damage from diseases and pests (Thies and Tscharntke,

1999; Poveda *et al.*, 2012). Analysis of the potential invasion network for pathogens and

pests in crop landscapes, such as cropland connectivity analysis, has been used to

identify candidate priority locations for surveillance and management of host-specific

pathogens and pests (Xing *et al.*, 2020; Andersen Onofre *et al.*, 2021; Buddenhagen *et*

al., 2022). An analysis of cropland geographic structure indicates the geographic

distribution of potential spread of pathogens and pests. Higher host plant availability

generally increases the local risk of host-specific pathogen and pest establishment,

while greater host connectivity can also increase the risk of spread and establishment.

1.2. Disaster plant pathology: Vulnerabilities and solutions in the Great Lakes

region

Many families living in the Great Lakes region rely on food security crops such as

banana and plantain, cassava, potato, and sweetpotato. However, many households

remain food insecure because of declining agricultural productivity, driven by frequent

disasters in the region, including conflicts, health emergencies, and natural disasters

that displace populations and disrupt agriculture (Ochieng *et al.*, 2014; Ochieng *et al.*,

2015; Ong'ayo, 2018). These challenges are compounded by the cascading effects of

high prevalence of plant pests and diseases, low agricultural input use, and loss of

harvest and assets (Kroschel *et al.*, 2014; McEwan, 2016; Almekinders *et al.*, 2019).

The Rwandese genocide in 1994, the civil war in Burundi from 1993 to 2005 and also in

2015, and the persistent conflicts in Eastern DR Congo are particularly important

humanitarian disasters in the Great Lakes region, which have left an indelible mark

(Ong'ayo, 2018). Health emergencies and natural disasters, such as severe droughts in

Ethiopia, Kenya, and Uganda (2015-2016) and devastating floods in Rwanda, Uganda,

and DR Congo (2020), have severely restricted agricultural practices such as effective

surveillance and management of fields, causing considerable damage to infrastructure

and cropland (Ochieng *et al.*, 2014; Ochieng *et al.*, 2015).

Disaster plant pathology, integrating concepts from plant pathology and disaster

science, addresses the risks and potential impact of plant diseases before, during, and

after disaster events that threaten agricultural systems and ecosystems (Etherton *et al.*,

2024; Mouafo-Tchinda *et al.*, 2024). It focuses on how disturbances modify host-

pathogen dynamics and increase damage and the spread of pathogens, and on the

development of strategies for better managing plant health. A combination of global

and national research efforts has been adopted to manage the impacts that these

disasters, pathogens, and pests may have on agroecological systems (RTB, 2013;

Science Group Project on Accelerated Varietal Improvement and Seed Systems in

Africa, 2024). For example, disease control programs which encourage farmers to

recognize and destroy diseased plants, and disease-resistant crop breeding programs

are currently underway across the Great Lakes region.

When models of pathogen and pest responses to environmental variables are

available, decision-support tools can be developed to predict where specific disease-

and pest-resistant varieties will be needed in the long-run, and in the short-run whether

likely economic injury levels motivate management actions by growers (Kaundal *et al.*,

2006; Carisse and Fall, 2021; Fenu and Malloci, 2021). Pest and disease forecasting

can reduce production costs, pesticide use, and environmental pollution, protecting the

health of farmers and consumers. Machine learning can be used to analyze high

volume datasets, and improve the predictive power, accuracy, and efficiency for

diseases and pests (Kaundal et al., 2006; Garrett et al., 2022a). Implementing decision
support systems in low-resource countries such as those in the Great Lakes region is
challenging; in addition to the often-limited availability of high-resolution local weather
data as input to models, field observations and georeferenced datasets for the many
important pathogen and pest species are often unavailable for constructing predictive
models.

1.3. Objectives

In this study, we evaluated the geographic distribution and risk factors for 27
pathogens and pests in the production of four food security crops — banana (including
plantain), cassava, potato, and sweetpotato — in the Great Lakes region of Africa,
including the potential future risk under climate change. The first objective was to
analyze the geographic risk of host-specific pathogen and pest spread based on host
landscape connectivity for these crops. The second objective was to characterize the
seasonal and geographic occurrence and associations of 27 economically important
pathogens and pests affecting these crops, through field surveys conducted in two key
locations in the region, Rwanda and Burundi. Part of this dataset has been used to
characterize the prevalence of potato viruses and vectors (Okonya *et al.*, 2021) and the
occurrence of banana pests and diseases across altitudes (Nakato *et al.*, 2023). The
third objective was to predict pathogen and pest severity/infestation based on climate
variables and host landscape structure using machine learning, and to assess which
pathogens and pests are likely to experience altitudinal range shifts under projected
changes in temperature. The field surveys were implemented along environmental
gradients to compare lower- and higher-altitude pathogen and pest communities, where
lower-altitude communities have the potential to move toward higher altitudes in the
future. The simultaneous study of these 27 pathogens and pests allows a unique direct
comparison of their associations with environmental predictors. These analyses can
also support decision-making by (i) national agricultural agencies that plan geographic
surveillance strategies for economically important pathogens and pests, (ii) local
agricultural research communities and international partners as they prioritize efforts to
improve current and future mitigation responses to pathogens and pests, and (iii) farmer

communities that need to be prepared for changes in pathogen and pest dynamics due
to natural or human-driven disasters, including changing climates.

2. Materials and methods

2.1. Landscape structures influencing pathogen and pest risk

2.1.1. Identifying locations with high cropland connectivity

The spatial pattern of host availability is a key ecological trait that influences the
spread of host-specific pathogens and pests. We analyzed the connectivity of host
landscapes in the Great Lakes region to map the potential establishment and regional
spread of host-specific pathogens and pests, to identify candidate priority locations for
spatially targeted surveillance in four food security crops — banana (including plantain),
cassava, potato, and sweetpotato. This cropland connectivity analysis assesses how
crop production is connected in a landscape, based on host availability and physical
proximity, in a gravity model (Jongejans *et al.*, 2015). Cropland connectivity analysis
identifies potential habitat networks through which pests and pathogens can spread, a
first approximation to epidemic or invasion networks, which can be refined as more data
become available (Margosian *et al.*, 2009; Xing *et al.*, 2020; Andersen Onofre *et al.*,
2021).

For each crop, we used publicly available estimates of the cropland harvested
area (or cropland density) across sub-Saharan Africa in 2017 (International Food Policy
Research Institute, 2020) as a proxy for host availability to host-specific pathogens and
pests. Maps of cropland density, aggregated at a spatial resolution of 10-min (or grid
cells of image 9.27 km × 9.27 km), were used to evaluate the potential epidemic or
invasion network using a gravity model (Xing *et al.*, 2020). Here the gravity model
incorporated two pathogen or pest dispersal kernels: $c_i c_j d_{ij}^{-\beta}$ for the inverse power law
model, and $c_i c_j e^{-\gamma d_{ij}}$ for the negative exponential model. In each dispersal model, d_{ij} is
the Vincenty ellipsoid distance between a pair of cropland nodes (locations) i and j, c_i is
the cropland density for node i, and β and γ are dispersal parameters describing the

relative likelihood of pathogen or pest movement across d_{ij} (Xing *et al.*, 2020; Andersen

Onofre *et al.*, 2021).

We calculated a cropland connectivity risk index (CCRI) (Xing *et al.*, 2020) for

each location with crop production in the region as a measure of its likely importance in

epidemic or invasion networks. The CCRI was the weighted mean of four network

metrics (node centralities), evaluated using the geohabnet package v2.0.0 (Keshav *et*

al., 2024) in R v4.4.0 (R Core Team, 2024), averaged across the results from a

sensitivity analyses for a range of dispersal parameters, as described by Mouafo-

Tchinda *et al.* (2024). Two CCRI formulations were used to capture key facets of node

connectivity. The first CCRI considered ("CCRI-betweenness") was the mean of

betweenness centrality weighted by 1/2, node strength weighted by 1/6, eigenvector

centrality weighted by 1/6, and the sum of nearest neighbors' degrees weighted by 1/6.

The second, "CCRI-closeness", considered 1/2 closeness centrality instead of

betweenness centrality, with the same weights for the other metrics. We emphasized

betweenness or closeness centrality to account for, respectively, the importance of a

node as a bridge between parts of the network and the relative facility with which a

location can be reached from all other locations in the network.

Habitat connectivity analysis using the geohabnet package (Keshav *et al.*, 2024)

is part of the R2M toolbox for rapid risk assessment supporting mitigation of pathogens

and pests (garrettlab.com/r2m). Other examples of applications of habitat connectivity

and R2M tools, in addition to those discussed elsewhere in this paper, illustrate how

these tools can be used in an integrated analysis (Andersen Onofre *et al.*, 2019;

Garrett, 2021; Etherton *et al.*, 2023; Mouafo-Tchinda *et al.*, 2024; Etherton *et al.*, 2025).

2.1.2. Associations among altitude, crop landscape structure, and climate

We characterized associations among altitude, crop landscape structure, and

climate in the Great Lakes region. The map of altitudes was retrieved from the Shuttle

Radar Topography Mission (SRTM) database, using the geodata package v0.5-8

(Hijmans *et al.*, 2023) in R. We also retrieved the 19 bioclimatic variables for the region

from Worldclim database v2.1 (Fick and Hijmans, 2017) using the geodata package

version 0.5-8 (Hijmans *et al.*, 2023) and the terra package v1.8-0 (Hijmans, 2024) in R.

For crop landscape structure variables, we used the raster maps of cropland density

and connectivity (CCRI-closeness) generated in section 2.1.1, using the terra package

(Hijmans *et al.*, 2022). We assessed the associations between altitude, bioclimatic

variables, and cropland structure by calculating the Pearson correlations between these

variables using the metan package v1.18.0 (Olivoto and Lúcio, 2020) in R.

2.2. Pathogen and pest communities

2.2.1. Field sampling and spatiotemporal structure of pathogen and pest communities

For each crop species, we identified fields for sampling along four survey

transects in Rwanda and Burundi, each transect representing an altitude gradient. In

Burundi, the field surveys included cropland around Lake Tanganyika in the Rusizi

watershed, representing the Bubanza, Bujumbura Rural, Cibitoke, and Muramvya

provinces (Fig. 1C). In Rwanda, the field surveys were in major cropland around Lake

Kivu in the Ruhengeri watershed, which includes part of the Northern and Western

provinces (Fig. 1C). The survey locations represented major crop landscapes in both

countries, encompassing a mixture of cropping systems and a range in altitude from

900 to 2800 m.a.s.l. (Fig. 1a-b) (Kroschel *et al.*, 2014). Fields for sampling were

selected at regular intervals along the transects regardless of whether diseases or pest

damage were apparent. The number of fields sampled in Burundi and Rwanda was 189

and 103 for banana, 78 and 116 for potato, and 107 and 104 for sweetpotato,

respectively. For cassava, sampling was only carried out in 50 fields in Burundi, with the

second Burundi survey not conducted due to civil unrest, and Rwanda was not sampled

because few farmers grew cassava in the Rwanda study area.

Sampling was conducted in each watershed during two cropping seasons, dry

and rainy. In both countries, two rainy seasons occur: one from March to May and the

other from September to December (WorldData.info, 2024). The dry seasons occur

from June to August and January to February (WorldData.info, 2024). For each field

sampled, we evaluated crop-specific pest or disease intensity (Table 2) based on

prevalence in the region, disease severity, infestation rate and intensity, and/or the

number of pests, using methods as described by Nakato *et al.* (2023), Okonya *et al.*

(2021), and Legg *et al.* (2009). A total of 27 economically important pathogens and

pests in the region were included in this study, including 6 for banana, 6 for cassava, 7

for potato and 8 for sweetpotato (Table 2). Here prevalence refers to the presence or

absence of a pest or disease in a field, while disease severity, pest infestation intensity

or infestation rate is the percentage of leaves or plants damaged by a disease or pest.

For infestation rate, infestation intensity, and severity, twenty plants per field, selected in

a systematic pattern, were assessed for visual symptoms of damage by pests and

diseases. The percentage of plants infested was noted as the infestation rate. The

percentage of symptomatic leaves per plant was noted as infestation intensity for pests

and severity for diseases. For damage by some virus diseases and pests such as

cassava mosaic disease (CMD) and cassava green mite (CGM), crop damage was

scored from 1 (representing no symptoms) to 5 (the most severe) (Sseruwagi *et al.*,

2004; Legg *et al.*, 2009; Okonya *et al.*, 2021). The disease or pest damage score

measures the severity or infestation intensity in a field, based on visual assessments

and a predefined scale. Thirty plants were randomly sampled at regular intervals along

the season and damage was assessed by scoring each plant on a scale of 1 to 5 as

described by Legg *et al.* (2009).

For banana, fields were surveyed in Burundi in March 2015 (during the rainy

season) and in July 2016 (dry season), while fields were surveyed in Rwanda in July

2015 (dry season) and in November 2016 (rainy season). Cassava fields were surveyed

only once in Burundi in March 2015 (rainy season), for reasons described above. For

Fig. 1. Location of fields of banana, cassava, potato, and sweetpotato where pathogen and pest surveys were carried out in Africa (A), in the Great Lakes region (B), in transects across altitude gradients, in Rwanda and Burundi (C). White lines in (C) indicate the province boundaries of Rwanda and yellow lines indicate the province boundaries of Burundi. Crosses and circles represent locations surveyed during the dry and rainy season, respectively. The complete surveillance effort included 412 crop fields.

potato, field sampling was conducted in Burundi in July 2016 (dry season) and

December 2017 (rainy season), and in Rwanda in June-July 2015 (dry season) and

November-December 2017 (rainy season). For sweetpotato, the field surveys took

place in Burundi in November-December 2015 (rainy season) and August 2016 (dry

season), and in Rwanda in July 2015 (dry season) and November 2017 (rainy season).

The irregular dates for these crop-field surveys were due in part to the civil unrest

prevailing in the region at the time the study was conducted.

2.2.2. Quantifying patterns of field-scale association of crop pathogens and pests

We evaluated the patterns of association of pathogens and pests based on

prevalence and severity/infestation using non-metric multidimensional scaling (NMDS)

(Kruskal, 1964) and heatmaps (Pryke *et al.*, 2007) for visualization. We initially

calculated five measures of dissimilarity (Bray, Manhattan, Jaccard, Chao, and robust-

Aitchison) in the NMDS analysis using the ggvegan package v0.1.999 (Simpson, 2020)

in R. We selected the Bray distance in the final analysis due to its highest goodness of

fit. A heatmap was prepared using the metan package v1.18.0 (Olivoto and Lúcio, 2020)

in R to map the association of pathogens and pests based on prevalence and

severity/infestation rates.

2.3. Prediction of pathogen and pest damage and geographic range shifts

2.3.1. Machine learning to predict pathogen and pest levels

Weather, climate, and geographic variables were used to predict severity or

infestation rates/intensity, comparing machine learning algorithms. We retrieved

weather variables (monthly, quarterly, and annual temperature, precipitation, and

relative humidity) from the NASAPOWER database using the nasapower package

v4.0.10 (Sparks, 2018) in R. The climate variables included the 19 bioclimatic variables,

solar radiation (kJ m-2 day-1), wind speed (m s-1), and water vapor pressure (kPa)

described, along with the altitude variable, in section 2.1.2. The CCRIs and cropland

density were assembled as described in section 2.1.1.

Data were preprocessed to exclude collinearity between variables. A stratified

split was then used to divide the data into training and test sets in an 8:2 ratio. We then

trained ten machine-learning algorithms: logistic regression (GLM and GLMNET),

support vector machine (SVM, with linear and radial basis function), neural network

(NN), k-nearest neighbors (KNN), stochastic gradient boosting (GBM, generalized

boosted modeling), bagged trees, classification and regression trees (CART), and

random forest (RF). We compared the performance of these algorithms to predict the

severity or infestation of each pathogen and pest using the caret package v6.0-94

(Kuhn, 2008) in R. We selected SVM (with radial basis function) and RF for further

analyses based on their having the highest performance in terms of MAE (mean

absolute error), RMSE (root mean squared error) and R-squared (Kaundal *et al.*, 2006).

324   In the next stage using the SVM and RF models, we executed a 10-fold cross-

validation with ten repetitions to prevent overfitting the training model and used a grid

search for hyperparameter tuning. The tuning process was driven by trainControl, with

RMSE-based model evaluation to prioritize accuracy. Hyperparameter tuning for SVM

included C (regularization parameter) and Gamma (kernel coefficient), both with a

tuneLength of 10. Hyperparameter tuning for RF included mtry (the number of

predictors randomly selected at each division) with a tuneLength of 10. We trained the

model using the train function in the caret package and estimated the performance

parameters MAE, RMSE, and R-squared. We used the resulting model to predict

severity or infestation rate/intensity for the test data and estimated the performance

parameters (RMSE and R-squared).

2.3.2. Pathogen and pest communities across altitudinal gradients

336   To study the geographic range of pathogen and pest communities associated

with the four crops in the sample transects in Rwanda and Burundi, we evaluated how

pathogen and pest levels (prevalence, disease severity or infestation rate/intensity)

changed with altitude across the sampling locations. We also evaluated the pathogen

and pest richness (number of pathogens and pests present from those included in the

sampling) in communities during the dry and rainy seasons, across the altitude

gradients. The associations between prevalence, infestation/severity and richness of

pathogens and pests across altitude gradients were evaluated based on Pearson's

correlation.

2.3.3. Climate change: impact on regional temperature patterns

346   We assessed potential climate shifts along the altitude gradients in the Great

Lakes region, using current and future temperature data from WorldClim database v2.1

(Fick and Hijmans, 2017). We used future climatic variables from three global circulation

models (GCMs: GISS-E2-1-G, MIROC6 and CNRM-CM6-1), two Shared

Socioeconomic Pathways (SSP1-2.6 and SSP5-8.5), and four future time-periods:

2021-2040, 2041-2060, 2061-2080, and 2081-2100. We selected these GCM modeling

groups because of their equilibrium climate sensitivity (ECS) (Hausfather, 2019). ECS

indicates the severity of future warming impacts. CNRM-CM6-1 assumes a high value

of ECS, 4.9 C, while GISS-E2-1-G and MIROC6 assumes low values of ECS, 2.7 and

2.6 respectively (Hausfather, 2019). The two SSPs used in this study represent different

emission pathways by the year 2100: sustainability (SSP1-2.6) and fossil-fuel

development (SSP5-8.5) emission scenarios featured by the radiation forcing of 2.6 and

8.5 W/m², respectively (Pinnegar *et al.*, 2021). We calculated the average annual

temperature across the three selected GCMs for each future time-period and SSP

combination. We used the resulting density plots to assess potential shifts in the

distribution of annual mean temperature at low (<1500 m.a.s.l.) and high (>1500

m.a.s.l.) altitudes, using the ggplot2 package v3.5.0 (Wickham, 2016) in R.

3. Results

3.1. Landscape structures influencing pathogen and pest risk

3.1.1. Identifying locations with high cropland connectivity

The analysis of cropland connectivity in the Great Lakes region (i) quantifies how

the geographic distribution of crop production could influence the potential spread of

crop-specific pathogens and pests and (ii) identifies candidate priority locations for

pathogen and pest surveillance for each food security crop (Fig. 2). These maps of

cropland connectivity highlight locations in the region that are likely important for

understanding and managing the risk of crop-specific pathogens and pests.

In these maps of cropland connectivity, a higher CCRI-betweenness (Fig. 2A) or CCRI-

closeness indicates the potential to play an important role in the establishment and

spread of crop-specific pathogens and pests affecting these crops. Locations with high

CCRI are candidate priorities for surveillance and mitigation programs. Although CCRI-

betweenness and CCRI-closeness may not yield identical CCRI values, they tended to

highlight the same potential locations for prioritization (Fig. 2A). For banana, locations in

Rwanda, Burundi, the DR Congo, Uganda, Tanzania, Ethiopia, Malawi, and

Mozambique have a high CCRI (Fig. 2). Locations in the southern, eastern, and

northern provinces of Rwanda have the highest CCRI-betweenness (Fig. 2A and 2B).

For cassava, several locations across the region have a high CCRI. In Burundi,

locations in the Cibitoke, Bubanza, Kirundo, Ngozi, Bujumbura Rural and Gitega

provinces have the highest CCRI (Fig. 2A and B). For potato, locations in Rwanda,

Burundi, the DR Congo, Uganda, Tanzania, Ethiopia, and Malawi have high CCRI-

betweenness (Fig. 2). These maps also show that locations in the northern, western,

and southern provinces of Rwanda have the highest CCRI-betweenness (Fig. 2A and

B). For sweetpotato, locations in Rwanda, Burundi, Uganda, Tanzania, Malawi, and

Ethiopia have a high CCRI (Fig. 2). These maps indicate the important role that Rwanda

and Burundi play for understanding and mitigating the risks associated with the spread

of pathogens and pests of these crops.

Maps of cropland connectivity (Fig. 2B) also illustrate the potential roles of the

locations we selected for field sampling crop pathogens and pests in the Ruhengeri

watershed in Rwanda and in the Rusizi watershed in Burundi. The connectivity of these

landscapes varies from crop to crop. Some sampled locations had high cropland

connectivity (e.g., cassava and sweetpotato in Burundi transects and potato in Rwanda

transects) compared to other locations. Overall, the surveyed locations around Lake

Kivu in Rwanda and Lake Tanganyika in Burundi, as well as other locations in the two

countries, are likely to be important candidate locations for surveillance for the

establishment and spread of banana, cassava, potato, and sweetpotato pathogens and

pests in both countries, based on cropland connectivity (Fig. 2B).

Fig. 2. Maps of a cropland connectivity risk index (CCRI-betweenness) for four food security crops in (A) the Great Lakes region of Africa and (B) in a focused geographical extent in Rwanda and Burundi. The CCRI indicates how important a location is likely to be for potential spread of a host-specific pathogen or pest, based on the network of host availability. The CCRI is a weighted mean of four network metrics, including betweenness centrality in the results shown here. In (A), dark blue lines indicate national boundaries. In (B), blue lines indicate the provinces of Rwanda and green lines indicate the provinces of Burundi.

3.1.2. Associations among altitude, landscape structure and climate

The associations among (a) altitude, (b) crop landscape structure, and (c) climate

variables such as temperature and precipitation, were analyzed for the Great Lakes

region. Higher temperatures were generally associated with lower altitudes. There was

a strong negative correlation between altitude and most temperature-related bioclimatic

variables (Tem, Tmin, TColQ, TDriQ, TWetQ, TWarQ, Tmax, TColdQ). In contrast,

altitude was weakly positively correlated with some precipitation-related bioclimatic

variables such as PrecS, PWetQ, and PWetM, and negatively correlated with others, including Prec, PDriQ, PWarQ, and PColQ (Fig. 3). In this region, higher cropland density and cropland connectivity of banana, cassava, potato, and sweetpotato (Fig. 3) were at higher altitudes (except cassava density). Cropland connectivity of these crop species had a stronger positive correlation with altitude compared to cropland density. Cropland density and cropland connectivity for each of these four crops were negatively correlated with most temperature-related bioclimatic variables and positively correlated with most precipitation-related bioclimatic variables (Fig. 3).

Fig. 3 Associations between altitude, bioclimatic variables, and crop landscape structure (crop density and cropland connectivity) in the Great Lakes region of Africa, based on Pearson's correlation. Strong positive correlations are in dark blue, while strong negative correlations are in dark red. White boxes without labels indicate no association.

3.2. Pathogen and pest communities

3.2.1. Spatiotemporal structure of pathogen and pest communities

420 Most of the targeted pathogens and pests were found in Burundi and Rwanda in
421 both crop-growing seasons (Fig. 4). Sweetpotato mealybugs, sweetpotato butterfly
422 (*Acraea acerata*) and sweetpotato aphids were observed only in Burundi (Fig. 4). The
423 observed prevalence of each pest or disease often varied by country, crop, and season.

Fig. 4. Seasonal distribution of observed pathogens and pests in smallholder production of banana, potato and sweetpotato in Burundi and Rwanda. (Cassava data are not represented here). The length of the bars represents the prevalence for both seasons (where a total of 2 indicates presence in every field in both seasons). Abbreviations: Banana bunchy top virus (BBTV), *Cosmopolites sordidus* (CS), *Ralstonia solanacearum* (RS), sweetpotato virus (SPV), and Xanthomonas wilt of banana (XW).

424 In both countries and seasons, the highest prevalence was observed for *Ralstonia*
425 *solanacearum* and virus symptoms in potato, virus symptoms in sweetpotato, and
426 *Cosmopolites sordidus* in banana.

3.2.2. Associations among pathogens and pests

The NMDS analysis identified a limited set of pathogens/pests whose levels were

associated, although some were common enough that they frequently co-occurred with

others (Fig. 5). A few pathogens and pests were associated in terms of prevalence and

severity/infestation in the crop fields. Among banana diseases/pests, Fusarium wilt,

Xanthomonas wilt and plant parasitic nematodes were associated in terms of both

prevalence and severity (Fig. 5). In cassava, all pathogens and pests except *B. tabaci*

were associated in terms of prevalence. However, *B. tabaci* was slightly associated with

CMV in terms of infestation, which is expected as *B. tabaci* vectors CMV (Fig. 5). In

potato, viruses and *R. solanacearum* were associated in terms of both prevalence and

severity/infestation (Fig. 5). In sweetpotato fields, *B. tabaci* and sweetpotato aphids

were associated, having similar infestation levels across fields.

Fig. 5. Relationships among economically important pathogens and pests and altitude in four major crops in Burundi and Rwanda, using non-metric multidimensional scaling (NMDS) based on Bray's distance dissimilarities of disease severity or infestation intensity. Each dot represents a surveyed crop field. A line (or vector) indicates whether (line direction) and how strongly (line length) the pathogen or pest is associated with others. Nearly parallel lines in the same quadrant indicate pathogens and pests often co-occur in a crop. (Cassava data was collected only in Burundi for one season). Abbreviations: *Acraea acerata* (AA), *Alternaria bataticola* (AB), Banana bunchy top virus (BBTV), *Bemisia tabaci* (cassava, potato or sweetpotato whitefly; BT), cassava bacterial blight (CBB), Cassava brown streak virus (CBSV), cassava green mite (CGM), cassava mealybug (CM), Cassava mosaic virus (CMV), *Cylas brunneus* (CB), *Cylas puncticollis* (CP), *Cosmopolites sordidus* (CS), Fusarium wilt of banana (FW), leaf miner fly (LMF), *Myzus persicae* (MP), *Phytophthora infestans* (PI), plant parasitic nematodes (PPN), *Pentalonia nigronervosa* (PN), *Phthorimaea operculella* (PO), potato viruses (PV), *Ralstonia solanacearum* (RS), *Synanthedon* sp. (clearwing moth; Ssp), sweetpotato mealybug (SPM), sweetpotato virus (SPV), sweetpotato weevils (SPW), and Xanthomonas wilt of banana (XW).

Analyses of associations between pairs of pathogens/pests provide a more

detailed perspective on which syndromes are common. In banana, some pairs like *C.*

sordidus and *P. nigronervosa*, plant parasitic nematodes and Fusarium wilt, plant

parasitic nematodes and *P. nigronervosa*, *P. nigronervosa* and Xanthomonas wilt, and

Fusarium wilt and Xanthomonas wilt were positively correlated ($p < 0.05$) in terms of

both prevalence and severity/infestation, while other pairs were only positively

correlated ($p < 0.05$) in terms of prevalence: *C. sordidus* and plant parasitic nematodes,

plant parasitic nematodes and Xanthomonas wilt, plant parasitic nematodes and BBTV,

P. nigronervosa and BBTV, and BBTV and Xanthomonas wilt (Fig. 6).

In cassava, CBSV and mealybug were positively correlated ($r = 0.69$, $p < 0.001$

in terms of prevalence but negatively correlated ($r = -0.44$, $p < 0.01$) in terms of

severity/infestation (Fig. 6). Only *B. tabaci* and CMV were positively correlated in terms

of severity/infestation ($r = 0.64$, $p < 0.001$) while other pathogens and pests were

positively correlated in terms of prevalence ($p < 0.05$). In potato, viruses and *R.*

solanacearum was positively correlated in terms of prevalence ($r = 0.33$, $p < 0.001$) and

severity ($r = 0.21$, $p < 0.01$), while *P. infestans* and *M. persicae*, and leaf miner fly and

M. persicae were negatively correlated ($p < 0.05$) in terms of prevalence and

severity/infestation, and leaf miner fly and *R. solanacearum* were positively correlated (p

< 0.05) in terms of prevalence but negatively correlated ($p < 0.05$) in terms of

severity/infestation (Fig. 6). In sweetpotato, *C. brunneus* and *C. puncticollis* were

positively correlated in terms of prevalence ($r = 0.58$, $p < 0.001$) and infestation ($r = $

0.18, $p < 0.01$), while *A. bataticola* and weevils (*Cylas* spp.) were negatively correlated

in terms of prevalence ($r = -0.16$, $p < 0.05$) but positively correlated in terms of

severity/infestation ($r = 0.18$, $p < 0.05$) (Fig. 6). *A. bataticola* and mealybug, *A. bataticola*

and aphids, *A. bataticola* and *B. tabaci*, *A. bataticola* and *Acraea acerata*, and aphids

and *Synanthedon* sp. were negatively correlated in terms of prevalence (Fig. 6).

Fig. 6. Association of pathogens and pests in banana, cassava, potato, and sweetpotato in smallholder farms in Burundi and Rwanda, based on Pearson's correlation of prevalence. (Cassava data was collected only in Burundi for one season).

Abbreviations: *Acraea acerata* (AA), *Alternaria bataticola* (AB), Banana bunchy top virus (BBTV), *Bemisia tabaci* (cassava, potato or sweetpotato whitefly; BT), cassava bacterial blight (CBB), Cassava brown streak virus (CBSV), cassava green mite (CGM), cassava mealybug (CM), Cassava mosaic virus (CMV), *Cylas brunneus* (CB), *Cylas puncticollis* (CP), *Cosmopolites sordidus* (CS), Fusarium wilt of banana (FW), leaf miner fly (LMF), *Myzus persicae* (MP), *Phytophthora infestans* (PI), plant parasitic nematodes (PPN), *Pentalonia nigronervosa* (PN), *Phthorimaea operculella* (PO), potato viruses (PV), *Ralstonia solanacearum* (RS), *Synanthedon* sp. (clearwing moth; Ssp), sweetpotato mealybug (SPM), sweetpotato virus (SPV), sweetpotato weevil (SPW), and Xanthomonas wilt of banana (XW).

3.3. Prediction of pathogen and pest damage and geographic range shifts

3.3.1. Machine learning to predict individual pathogen and pest levels

467
468

We compared machine learning algorithms in terms of their ability to predict
severity and infestation in the field samples at monthly, quarterly, and annual intervals
(Table 1). SVM and RF had the highest model performance for some pathogens/pests.
For the SVM model, the performance characteristics across all pathogens and pests
were 3-16% MAE for training, 7-20% and 4-26% RMSE, and 0.20-0.72 and 0.09-0.72
R-squared for training and testing, respectively, compared to the RF model with 3-17%
MAE for training, 8-21% and 5-23% RMSE, and 0.23-0.72 and 0.12-0.73 R-squared for
training and testing, respectively (Table 1). Among the 27 pathogens and pests tested,
the three with the highest predictive performance metrics per crop were reported in
Table 1.

The predictive performance was good for certain pathogens and pests, but poor
for others (Table 1). As an example, SVM and RF performed well in predicting the
monthly, quarterly, and annual infestation intensity of leaf miner fly in this region,
explaining 69-71% of the variability in the data (R-squared) with low errors. The RMSE
was 17-18% during training and 16-17% during prediction, demonstrating that both
models generalized well to unseen data. In addition, the MAE of 11-13% indicated a low
average error in training. The tight alignment between training and prediction metrics
showed the robustness, stability and reliability of these models to predict leaf miner fly
infestation intensity.

Pathogens and pests differed in the importance of predictor variables.
Contemporary weather variables, such as monthly temperature, precipitation, and
relative humidity (extracted through NASAPOWER), generally had high contribution
scores. Cropland structure variables also played significant roles, with contribution
scores reaching 100 for cropland density, 81 for cropland connectivity, and 69 for
altitude. For example, the key variables for predicting leaf miner fly infestation intensity

(monthly) using SVM were monthly temperature, precipitation, and relative humidity,

precipitation in the driest month, cropland connectivity, cropland density and annual

temperature range (i.e. difference between the maximum temperature of warmest

month and the minimum temperature of coldest month).

Table 1. Support vector machine (SVM) and random forest (RF) performance for

predicting pathogen and pest levels in smallholder farms in Burundi and Rwanda, for

selected species with higher model performance. The response variable was severity

for the pathogens and infestation intensity for pests, except for infestation rate for

sweetpotato aphids.

| | | | SVM | | | | | | Random Forest | | | | |
| | | | Training | | | Prediction | | | Training | | | Prediction | |
Crop	Pathogen/pest	Timeframe	%RMSE	R-squared	%MAE	%RMSE	R-squared	%RMSE	R-squared	%MAE	%RMSE	R-squared
Banana	*P. nigronervosa*	Monthly	18	0.63	13	21	0.56	16	0.72	12	18	0.66
		Quarterly	18	0.63	13	21	0.56	16	0.72	12	19	0.64
		Annual	19	0.63	14	21	0.55	16	0.71	11	19	0.63
	Banana bunchy top virus	Monthly	7	0.22	3	4	0.10	8	0.25	4	5	0.12
		Quarterly	7	0.22	3	4	0.11	8	0.23	4	5	0.13
		Annual	7	0.20	3	4	0.09	8	0.24	4	5	0.13
	Plant parasitic nematodes	Monthly	11	0.42	6	15	0.29	10	0.42	6	14	0.31
		Quarterly	11	0.40	6	15	0.25	11	0.37	7	14	0.32
		Annual	11	0.39	6	15	0.23	11	0.37	7	14	0.29
Potato	Leaf miner fly	Monthly	18	0.69	13	16	0.71	18	0.70	12	16	0.71
		Quarterly	18	0.69	13	16	0.71	18	0.71	11	16	0.69
		Annual	18	0.70	13	16	0.72	17	0.72	11	17	0.69
	Ralstonia solanacearum	Monthly	14	0.30	10	14	0.13	13	0.40	10	15	0.12
		Quarterly	15	0.28	10	14	0.13	13	0.41	10	15	0.12
		Annual	15	0.28	11	14	0.16	12	0.39	10	15	0.13
	Potato viruses	Monthly	20	0.31	16	21	0.34	21	0.28	17	21	0.38
		Quarterly	20	0.32	16	21	0.33	20	0.31	16	21	0.38
		Annual	20	0.31	16	21	0.34	21	0.30	17	21	0.37
Sweet potato	*Alternaria bataticola*	Monthly	15	0.35	9	10	0.42	14	0.39	9	10	0.39
		Quarterly	14	0.41	9	9	0.43	13	0.45	8	9	0.42
		Annual	15	0.35	8	9	0.47	14	0.38	9	9	0.43
	Sweetpotato weevils (*Cylas* spp.)	Monthly	19	0.44	10	25	0.26	18	0.38	10	22	0.44
		Quarterly	19	0.42	10	26	0.21	19	0.38	11	22	0.41
		Annual	20	0.41	10	25	0.27	19	0.36	11	23	0.40
	Sweetpotato aphids	Monthly	14	0.60	6	9	0.60	10	0.53	5	7	0.73
		Quarterly	14	0.60	6	9	0.58	10	0.58	4	8	0.65
		Annual	14	0.58	6	9	0.58	9	0.54	4	9	0.53

3.3.2. Pathogen and pest communities across altitudinal gradients

For each crop, pathogen and pest communities differed at low versus high altitudes (Table 2). Although there was not strong evidence for differences across the altitudinal gradients for 5 out of 27 pathogens and pests (correlations with p > 0.05 in Table 1), likely due in some cases to high variability across samples, there was evidence for differences in some pathogens and pests that distinguished the pathogen/pest communities (correlations in Table 2 with lower p-values). For instance, crop production at higher altitudes (>1,500 m.a.s.l) had more *Myzus persicae*, *Phthorimaea operculella*, and cassava green mites. At lower altitudes, crop fields had more Fusarium wilt and Xanthomonas wilt of bananas, *P. infestans*, *A. acerata* and *Bemisia tabaci* (Table 2). The distribution of these pathogens and pests across altitudes was supported in the analysis combining observations from Burundi and Rwanda; overall, for banana, potato, and sweetpotato, there was evidence that 44% of the targeted pathogens and pests were more common at low altitudes, and that 17% were more common at higher altitudes.

When analyzing data for Rwanda and Burundi separately, differences across altitudes for pathogens and pests were specific to the measure being used (e.g., disease intensity; Table 2) or the country. For example, potato fields at high altitudes had a higher prevalence of potato leaf miner fly, while there was not evidence for a difference in the infestation intensity of this pest across the altitudinal gradient. Similarly, there was stronger evidence that altitude was negatively correlated with the prevalence and severity of banana Fusarium wilt in Burundi (p < 0.001) compared to Rwanda (p = 0.069 for severity and p = 0.85 for prevalence). At the community level, potato fields in Rwanda and cassava fields in Burundi had higher pathogen and pest richness at higher altitudes, while richer pathogen and pest communities were found in banana fields at lower altitudes.

Table 2. Correlation between altitude and pathogen and pest prevalence, severity, and infestation in Burundi and Rwanda (PRV: prevalence, SEV: severity, Rate: infestation rate, INF: infestation intensity, NUM: number of pests).

Crop	Disease/ Pathogen/ Pest	Variables	Correlation in Burundi		Correlation in Rwanda		Correlation in both countries combined	
			r	P-value	r	P-value	r	P-value
Banana	Xanthomonas wilt	**PRV**	-0.08	0.28	-0.07	0.49	**-0.13**	**0.022**
		SEV	-0.02	0.76	-0.04	0.7	**-0.13**	**0.028**
	Fusarium wilt	**PRV**	**-0.37**	**<0.001**	-0.02	0.85	**-0.27**	**<0.001**
		SEV	**-0.33**	**<0.001**	0.18	0.069	**-0.34**	**<0.001**
	Banana bunchy top virus	**PRV**	-0.13	0.065	0	-	**-0.49**	**<0.001**
		SEV	-0.01	0.9	0	-	**-0.22**	**<0.001**
	Plant parasitic nematodes	PRV	-0.01	0.90	0.05	0.65	-0.11	0.065
		INF	-0.13	0.07	-0.03	0.78	-0.02	0.72
	Banana weevil (*Cosmopolites sordidus*)	**PRV**	0.03	0.700	-0.14	0.16	**0.18**	**0.003**
		INF	-0.07	0.37	**-0.38**	**<0.001**	**-0.13**	**0.034**
	Banana aphid (*Pentalonia nigronervosa*)	**PRV**	0.03	0.68	-0.08	0.43	**-0.14**	**0.018**
		INF	0.09	0.24	0.09	0.37	**-0.25**	**<0.001**
		Richness	**-0.16**	**0.028**	0.80	0.41	**-0.30**	**<0.001**
Potato	Leafminer fly	**PRV**	0.16	0.16	**0.26**	**0.006**	0.09	0.22
		INF	0.17	0.13	0.01	0.93	0.04	0.62
		Rate	0.14	0.21	0.13	0.16	0.01	0.88
	Potato aphid (*Myzus persicae*)	**PRV**	0.02	0.86	**0.24**	**0.011**	**0.23**	**0.001**
		Rate	0.07	0.54	**0.25**	**0.008**	**0.21**	**0.004**
	Potato whitefly (*Bemisia tabaci*)	**PRV**	**-0.57**	**<0.001**	0.04	0.7	**-0.17**	**0.02**
		NUM	**-0.38**	**<0.001**	0.08	0.39	0.05	0.45
	Potato tuber moth (*Phthorimaea operculella*)	**PRV**	**0.34**	**0.003**	0.03	0.75	**0.16**	**0.028**
		Rate	**0.36**	**0.001**	-0.05	0.58	0.15	0.03
	Ralstonia solanacearum	PRV	-0.09	0.41	0.10	0.31	0.04	0.61
		SEV	0.04	0.73	-0.08	0.41	-0.03	0.66
	Potato viruses	PRV	0	-	0.10	0.3	0.10	0.16
		SEV	0.04	0.71	**-0.21**	**0.025**	-0.61	0.4
	Phytophthora infestans	**PRV**	-0.20	0.082	0.07	0.46	**-0.18**	**0.014**
		SEV	-0.21	0.068	-0.06	0.53	**-0.14**	**0.044**
		Richness	-0.15	0.18	**0.26**	**0.005**	0.1	0.18
Sweet Potato	*Alternaria bataticola*	**PRV**	**0.58**	**<0.001**	0.05	0.61	**0.54**	**<0.001**
		SEV	**0.48**	**<0.001**	0.04	0.65	**0.32**	**<0.001**
	Sweetpotato viruses	PRV	0.01	0.98	0	-	0.03	0.63
		SEV	0.06	0.51	-0.15	0.14	-0.08	0.24
	Sweetpotato whitefly (*Bemisia tabaci*)	**PRV**	-0.11	0.26	0	-	**-0.22**	**0.001**
		NUM	-0.05	0.61	0	-	-0.11	0.13
	Sweetpotato butterfly (*A.*	**PRV**	**-0.19**	**0.045**	0	-	**-0.32**	**<0.001**
		Rate	-0.10	0.29	0	-	**-0.19**	**0.005**

Crop	Pest/Disease	Metric						
	acerata)	NUM	**-0.19**	**0.045**	0	-	**-0.22**	**0.001**
	Sweetpotato	PRV	-0.15	0.13	0.05	0.63	**-0.17**	**0.016**
	weevils (*Cylas*	INF	0.02	0.81	0.08	0.43	**0.17**	**0.013**
	spp.)	Rate	-0.03	0.76	-0.05	0.62	**-0.17**	**0.013**
	Cylas puncticollis	PRV	-0.08	0.43	0.04	0.69	-0.08	0.24
		NUM	**-0.23**	**0.016**	-0.06	0.54	-0.04	0.61
	C. brunneus	PRV	0.08	0.39	0.06	0.58	0.02	0.8
		NUM	-0.01	0.99	**-0.20**	**0.042**	-0.02	0.75
	Synanthedon sp.	PRV	**-0.28**	**0.004**	**0.26**	**0.007**	0.01	0.91
	Sweetpotato	PRV	**-0.20**	**0.036**	0	-	**-0.21**	**0.002**
	mealybug							
	Sweetpotato	PRV	**0.24**	**0.013**	0	-	0.01	0.94
	aphids	Rate	**0.45**	**<0.001**	0	-	**0.22**	**0.002**
		NUM	**0.35**	**<0.001**	0	-	**0.17**	**0.01**
		Richness	0.05	0.6	0.10	0.34	-0.02	0.8
Cassava	Cassava whitefly	PRV	**-0.80**	**<0.001**	-	-	-	-
	(*Bemisia tabaci*)	NUM	**-0.32**	**0.032**	-	-	-	-
	Cassava green	PRV	**0.37**	**0.012**	-	-	-	-
	mite	INF	0.06	0.68	-	-	-	-
	Cassava mealybug	PRV	0.29	0.057	-	-	-	-
		INF	0.27	0.072	-	-	-	-
	Cassava mosaic	PRV	**0.31**	**0.037**	-	-	-	-
	virus	SEV	**-0.52**	**<0.001**	-	-	-	-
	Cassava brown	PRV	0.17	0.25	-	-	-	-
	streak virus	SEV	**-0.47**	**0.002**	-	-	-	-
	Cassava bacterial	PRV	0.12	0.43	-	-	-	-
	blight	SEV	0.25	0.093	-	-	-	-
		Richness	**0.78**	**<0.001**	-	-	-	-

3.3.3. Climate change and impact on regional temperature patterns

The frequency distribution of annual temperatures across altitudes in the Great

Lakes region illustrates the transformative impact of climate change (Fig. 7). A

progressive shift in temperature distribution is expected, highlighting how currently

cooler climates at higher altitudes are likely to experience much higher temperatures

(Fig. 7). A general warming trend is projected at both high and low altitudes. There is a

greater temperature increase in the SSP5-8.5 scenario than in the SSP1-2.6 scenario.

Currently, the mean annual temperature in the region ranges between 17-30°C at low

altitudes, with the mode around 24°C, and between 10-22°C at higher altitudes, with the

mode around 17°C. As climate change progresses, and depending on socio-economic

pathways, this temperature landscape is likely to change substantially. Under the SSP1-

2.6 scenario, by 2090 (2081-2100), temperatures are expected to range between 20-

32°C at low altitudes, with the mode around 26°C, and between 10-25°C at high

altitudes, with mode 20°C. The high-emission scenario (SSP5-8.5) predicts that

temperatures will range between 22-36°C at low altitudes, with the mode around 28°C,

and between 12-30°C at higher altitudes, with the mode around 25°C.

These projections suggest future climate conditions at higher altitudes could

become more favorable for the 44% of pathogens and pests we evaluated that currently

are more common at lower altitudes (section 3.3.2). Likewise, pathogens and pests

currently more common at higher altitudes (17% of those evaluated in section 3.3.2)

may become less common as the locations warm in the future. These percentages are

based on banana, potato, and sweetpotato, the three crops for which we had the most

complete data sets for analysis in predictive models based on field data collected to

allow direct comparisons across pathogens and pests.

Fig. 7. Shift in annual temperature range at lower and higher elevations in the Great Lake region of Africa. Density is the probability density or likelihood of temperature rates occurring at locations in the Great Lakes region of Africa. TP: time period of the projections (while the 'Current' reference remains the same). SSP: Shared Socioeconomic Pathway, sustainability (SSP1-2.6) and fossil-fueled development (SSP5-8.5) emission scenarios. Here, high altitude refers to locations above 1500 m.a.s.l. and low altitude those below 1500 m.a.s.l.

4. Discussion

This study identified environmental predictors for a set of important pathogens and pests in tropical agroecosystems key to food security. We identified candidate surveillance locations based on cropland (host) connectivity, a starting point for understanding geographic spread and establishment of host-specific pathogens and pests. Based on the current geographic and seasonal distribution of 27 economically important pathogens and pests across climate gradients, we evaluated future climate scenarios. The 44% of the pathogens and pests currently more abundant at lower altitudes could become more common at higher altitudes under climate change, while the 17% more abundant at higher altitudes could become less common overall in the future. Identifying differences in pathogen and pest communities and their predictors contributes to a more complete understanding of agroecosystems, which can be translated to inform management strategies, as we discuss below.

4.1. Pathogen and pest risk across landscape structures

The cropland connectivity analysis provides a first approximation to the geographic distribution of pathogen and pest risk, based on host availability. This analysis identified candidate priority locations for surveillance and mitigation of the potential spread of host-specific pathogens and pests based on networks of host availability (Fig. 2). Analysis of cropland connectivity provides a baseline for quantifying invasion risks for pathogens and pests at a national or global scale (Andersen Onofre *et al*., 2021; Mouafo-Tchinda *et al*., 2024; Etherton *et al*., 2025). In the Great Lakes region, locations in Rwanda and Burundi had a high CCRI for all four food-security crops, highlighting their vulnerability to pathogen and pest spread in the absence of phytosanitary measures. Maps of cropland connectivity can inform monitoring of outbreaks of introduced or emerging diseases and pests by national plant protection organizations (NPPOs). Ongoing improvements to understanding other components of pathogen and pest risk – such as the effects of climate change, local trade, genetic resistance deployment, and other management adoption – could be integrated in more complete habitat connectivity analyses as additional resources become available

591   (Keshav *et al.*, 2024).

592   **4.2. Pathogen and pest communities**

593   Analysis of the on-farm distribution of 27 pathogens and pests highlights how
these communities change along climate gradients, in terms of seasonal abundance
(Fig. 4), community structure (Fig. 5), and interspecific pathogen/pest associations (Fig.
6). Previous studies have typically addressed how individual tropical pathogens or pests
are distributed along climate gradients (Ebregt *et al.*, 2005; Okonya and Kroschel, 2013;
Nyang'au *et al.*, 2021); a community perspective provides a more realistic scenario for
understanding regional pathogen and pest management (Savary *et al.*, 2000; Were *et
al.*, 2013; Makiola *et al.*, 2021). The study addresses this knowledge gap by identifying
which pathogens/pests are most prevalent across seasons, which pathogen/pest
occurrences are associated, and how climate gradients are likely to influence disease
severity and pest infestation. Evaluating the interactions and cumulative effects of
multiple pathogens/pests in specific production environments can increase our
understanding of injury profiles (Savary *et al.*, 2000), the cumulative damage to plants
caused by the set of biotic stressors including insects, pathogens and weeds.

607   Our findings highlight injury profiles for food security crops across climate
gradients. Some associations are already well-known; as expected, BBTV and its vector
P. nigronervosa were positively correlated (Hu *et al.*, 1996; Blomme *et al.*, 2020). BBTV
has been present in Africa for decades, its spread continues to have an impact on new
regions, and its management remains a major challenge (Rybicki, 2015; Ngatat *et al.*,
2024; Ocimati *et al.*, 2024). We also found strong associations between certain
pathogens and pests; Fusarium wilt, Xanthomonas wilt and plant parasitic nematodes
were associated in banana, viruses and *R. solanacearum* in potato, and *B. tabaci* and
aphids in sweetpotato. These pathogen/pest associations underscore the need for
integrated management strategies that address the spread and combined impacts of
multiple pathogens and pests on plants (Mouafo-Tchinda *et al.*, 2022; Alcalá Briseño *et
al.*, 2023).

4.3. Climate-based predictions of pathogen and pest levels

In the Great Lakes region, temperature generally decreased with increasing
altitude (Fig. 3), where higher temperatures were associated with an increase in levels
of some pathogens and pests and a decrease in others (Table 2). Whiteflies were more
common at low altitudes for cassava, potato and sweetpotato. Aphids were more
common at low altitudes for banana, but more common at high altitudes for potato. The
observed differences in the altitudinal distribution of these pests could be attributed to
their species composition, host specificity, and the influence of species-specific
ecological adaptations (Aleuy and Kutz, 2020; McCulloch and Waters, 2023). Aphid
communities included different species, with *P. nigronervosa* more frequent in low-
altitude plants, and *M. persicae* more frequent in high-altitude plants (Malumphy *et al.*,
2017; Wells and Clark, 2019; Wieczorek *et al.*, 2019; Aleuy and Kutz, 2020).

Weather variables are well known to be useful for predicting disease severity
(Kaundal *et al.*, 2006; Carisse and Fall, 2021; Fenu and Malloci, 2021). Cropland
geography variables (cropland density, connectivity), altitude, and climate variables
(annual mean temperature, temperature annual range, and annual precipitation) were
key for most of the pathogens and pests we evaluated. Model performance depends on
data availability, the predictor variables, and the biological features of pathogens and
pests (Xiao *et al.*, 2018; Fenu and Malloci, 2021). The data from Rwanda and Burundi
supported prediction of some pathogens, while others, like BBTV, would require more
study. The models effectively predicted leaf miner fly infestation intensity, accounting for
69-71% of the variability with low training and prediction errors. These models could
serve as a prototype for predictive models of leaf miner fly infestation intensity in this
region (Table 1).

Temperature influences the life cycle, reproduction, and migration patterns of
pathogens and pests (Awmack *et al.*, 1997; Yamamura and Kiritani, 1998; Mouafo-
Tchinda *et al.*, 2021). In response to regional warming, certain pathogens and pests
may become more common at specific altitudes (Poveda *et al.*, 2012; Datta *et al.*, 2017;
Blomme *et al.*, 2020), which will require adaptation of pathogen and pest management
to future conditions. Several studies suggest that climate change shifts some species
ranges to higher altitudes (Bale *et al.*, 2002; Colwell *et al.*, 2008; Bebber *et al.*, 2013),

and the new results from this study clarify the specific effects on individual pathogens

and pests in Rwanda and Burundi for key food security crops.

Pathogen/pest richness in banana decreased with altitude, while it increased in

cassava. In addition, the severity of certain pathogens/diseases such as BBTV, CMV, *P.*

infestans and Fusarium wilt decreased with altitude, while the severity or infestation of

A. bataticola and aphids in sweetpotato increased. In general, we found evidence that

44% of the pathogens and pests currently more common at low altitude in banana,

potato, and sweetpotato could shift to higher altitudes under climate change, based on

climate matching. Specifically, those expected to become more common at higher

altitudes are Xanthomonas wilt, Fusarium wilt, BBTV, and *P. nigronervosa*, in banana;

sweetpotato mealybug, *A. acerata*, sweetpotato weevils in sweetpotato; and *B. tabaci* in

potato and sweetpotato. The 17% of pathogens and pests currently more common at

high altitudes could become less common. Cropland connectivity was positively

correlated with altitude and was a key predictor of severity/infestation of some

pathogens and pests. This suggests that food-security crops at higher altitudes may

experience substantial flows of pathogens and pests, amplifying the risk of damage.

Customizing risk assessments for these specific pest/pathogen communities across

altitudes could make management strategies better adapted to the challenges posed by

climate change for each crop in the region.

4.4. Translating agroecosystem understanding of pathogens and pests to support

current and future agriculture

Historically, global and national research and extension efforts have been

employed to manage diseases and pests in the Great Lakes region (RTB, 2013;

Kroschel *et al.*, 2014). Integrated management systems combining cultural control,

surveillance, awareness creation and involving policy makers and farmers have been

adopted (Legg *et al.*, 2006; Gotor *et al.*, 2022). There has been support for active

management programs where clean planting material is provided to farmers to

encourage them to destroy diseased plants. Efforts have been made to create

awareness and empower farmers to recognize and destroy diseased plants. A

challenge for these efforts is limited resources to support ongoing, sustainable plans by

governments and national programs, especially after project lifespans are over. The

results of this new study can help to inform prioritization and efficient use of available

resources. Currently, for many food security crops, efforts are underway to improve

seed health, obtain resistance to key pathogens and pests, and adapt these improved

technologies and techniques for use in humanitarian settings (RTB, 2013; Andrade-

Piedra *et al.*, 2020; Andrade-Piedra *et al.*, 2023).

Pathogen and pest risk analysis can help to ensure the economic, social, and

environmental sustainability of crop production (Kaundal *et al.*, 2006; Fall *et al.*, 2016;

Carisse and Fall, 2021). Image analysis apps such as the PlantVillage Nuru app and the

Tumaini app can help smallholder farmers in the region diagnose plant damage caused

by pests and diseases, and offer national plant protection organizations and extension

staff a tool for surveillance and disease mapping (Ramcharan *et al.*, 2019; Selvaraj *et

al.*, 2019; Kreuze *et al.*, 2022). Predicting the timing of pathogen and pest threats would

enable farmers to adjust management strategies in the short run and would support

prioritization in research and breeding programs in the long run.

Ongoing improvements in predicting pathogen and pest impacts depend on

weather data availability, which is often limited in low-income and disaster-prone

countries. Satellite platforms like NASA POWER offer a useful alternative (Sparks,

2018), providing near-real-time weather data and allowing researchers in resource-

limited settings to develop useful models without the need for extensive ground

observations. Another challenge is detection of transient pests such as aphids, which

move frequently and rapidly across fields (Loxdale, 2018). Innovative sampling

techniques, such as suction traps and environmental DNA (eDNA) analysis, could be

helpful for monitoring pest populations more accurately (Taberlet *et al.*, 2012; Poppinga

et al., 2016). However, implementing these advanced approaches in low-income and

disaster-prone areas such as the Great Lakes region remains a challenge due to limited

resources and technical capacity, requiring scalable and cost-effective solutions.

In disaster-prone regions, humanitarian organizations and government agencies
must prioritize key areas where abandonment or poor management of staple crops is
particularly important to regional spread of pathogens and pests (Etherton *et al.*, 2024;
Mouafo-Tchinda *et al.*, 2024). This study identifies candidate priority locations for
mitigation of the establishment and spread of pathogens and pests, characterizes
pathogen and pest associations (injury profiles), and evaluates potential altitudinal shifts
under the influence of climate change. Future research can build on these results by
validating and refining the models for each pathogen/pest, and developing tools that
growers and policymakers can easily implement. Ongoing improvement can increase
the adaptability and effectiveness of plant health management strategies, a no-regrets
adaptation strategy as agricultural systems face increasing challenges due to climate
change.

Acknowledgements

We appreciate support from the CGIAR Seed Equal Research Initiative, the
CGIAR Roots Tubers and Bananas Research Program, USAID Bureau of Humanitarian
Assistance (BHA) award number 720BHA22IO00136, the OneCGIAR Initiative on Plant
Health, and the CGIAR Trust Fund (www.cgiar.org/funders/); we thank all donors and
organizations which globally support the work of CGIAR through their contributions to
the CGIAR Trust Fund. We also appreciate support from USDA Animal and Plant
Health Inspection Service (APHIS) Cooperative Agreements AP21PPQS&T00C195 and
AP22PPQS&T00C133. The opinions expressed in this article are those of the authors
and do not necessarily reflect the view of USAID BHA or USDA APHIS.

Competing interests

730   The authors declare that they have no known competing financial interests or
731 personal relationships that could have appeared to influence the work reported in this
732 paper.

References

Adhikari, U., Nejadhashemi, A.P., Woznicki, S.A., 2015. Climate change and eastern
Africa: a review of impact on major crops. Food and Energy Security 4, 110-132.
Alcalá Briseño, R.I., Batuman, O., Brawner, J., Cuellar, W.J., Delaquis, E., Etherton, B.A.,
French-Monar, R.D., Kreuze, J.F., Navarrete, I., Ogero, K., et al. 2023. Translating
virome analyses to support biosecurity, on-farm management, and crop breeding.
Frontiers in Plant Science 14, 1056603.
Aleuy, O.A., Kutz, S., 2020. Adaptations, life-history traits and ecological mechanisms of
parasites to survive extremes and environmental unpredictability in the face of climate
change. International Journal for Parasitology: Parasites and Wildlife 12, 308-317.
Almekinders, C.J., Walsh, S., Jacobsen, K.S., Andrade-Piedra, J.L., McEwan, M.A., de
Haan, S., Kumar, L., Staver, C., 2019. Why interventions in the seed systems of roots,
tubers and bananas crops do not reach their full potential. Food Secur. 11, 23-42.
Andersen Onofre, K.F., Buddenhagen, C.E., Rachkara, P., Gibson, R., Kalule, S., Phillips,
D., Garrett, K.A., 2019. Modeling epidemics in seed systems and landscapes to guide
management strategies: The case of sweet potato in northern Uganda. Phytopathology
109, 1519-1532.
Andersen Onofre, K.F., Forbes, G.A., Andrade-Piedra, J.L., Buddenhagen, C.E., Fulton,
J.C., Gatto, M., Khidesheli, Z., Mdivani, R., Xing, Y., Garrett, K.A., 2021. An integrated
seed health strategy and phytosanitary risk assessment: Potato in the Republic of
Georgia. Agr Syst 191, 103144.
Andrade-Piedra, J., Almekinders, C.J., McEwan, M., Kilwinger, F., Mayanja, S., Mulugo, L.,
Delaquis, E., Garrett, K., Omondi, B.A., Rajendran, S., 2020. User guide to the toolbox
for working with root, tuber and banana seed systems. Lima, Peru. CGIAR Research
Program on Roots, Tubers and Bananas. RTB User Guide.
Andrade-Piedra, J., Dontsop, P., Dunia, D., Kulakow, P., Harahagazwe, D., Mouafo-
Tchinda, R., Omondi, A., Ogero, K., Rajendran, S., McEwan, M., 2023.
Tools4SeedSystems: Working towards resilience through root, tuber and banana crops
in humanitarian settings. Capacity needs assessment for root, tuber and banana seed
interventions in humanitarian settings: Cameroon and DRC. Technical Report. 65.
Awmack, C., Harrington, R., Leather, S., 1997. Host plant effects on the performance of
the aphid *Aulacorthum solani* (Kalt.)(Homoptera: Aphididae) at ambient and elevated
CO_2. Global Change Biology 3, 545-549.
Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K.,
Butterfield, J., Buse, A., Coulson, J.C., Farrar, J., 2002. Herbivory in global climate
change research: direct effects of rising temperature on insect herbivores. Global
Change Biology 8, 1-16.
Bebber, D.P., Ramotowski, M.A.T., Gurr, S.J., 2013. Crop pests and pathogens move
polewards in a warming world. Nature Climate Change 3, 985-988.
Blomme, G., Ocimati, W., Amato, S., Felde, A.z., Kamira, M., Bumba, M., Bahati, L., Amini,
D., Ntamwira, J., 2020. Banana pest risk assessment along banana trade axes running
from low to high altitude sites, in the Eastern DR Congo and in Burundi. African Journal
of Agricultural Research 16(9), 1253-1269.
Buddenhagen, C.E., Xing, Y., Andrade-Piedra, J.L., Forbes, G.A., Kromann, P., Navarrete,
I., Thomas-Sharma, S., Choudhury, R.A., Andersen Onofre, K.F., Schulte-Geldermann,

E., Etherton, B.A., Plex Sula, A.I., Garrett, K.A., 2022. Where to invest project efforts for
greater benefit: A framework for management performance mapping with examples for
potato seed health. Phytopathology 112, 1431-1443.
Carisse, O., Fall, M.L., 2021. Decision trees to forecast risks of strawberry powdery mildew
caused by *Podosphaera aphanis*. Agriculture 11, 29.
Chevalier, M., Broennimann, O., Guisan, A., 2024. Climate change may reveal currently
unavailable parts of species' ecological niches. Nature Ecology & Evolution 8, 1298–
1310.
Cilas, C., Goebel, F.-R., Babin, R., Avelino, J., 2016. Tropical crop pests and diseases in a
climate change setting—a few examples. In: Torquebiau, E. (Ed.), Climate change and
agriculture worldwide, Springer, Dordrecht, pp. 73-82.
Colwell, R.K., Brehm, G., Cardelús, C.L., Gilman, A.C., Longino, J.T., 2008. Global
warming, elevational range shifts, and lowland biotic attrition in the wet tropics. Science
322, 258-261.
Datta, A., Kühn, I., Ahmad, M., Michalski, S., Auge, H., 2017. Processes affecting
altitudinal distribution of invasive *Ageratina adenophora* in western Himalaya: The role
of local adaptation and the importance of different life-cycle stages. PLOS One 12,
e0187708.
Ebregt, E., Struik, P., Odongo, B., Abidin, P., 2005. Pest damage in sweet potato,
groundnut and maize in north-eastern Uganda with special reference to damage by
millipedes (Diplopoda). NJAS-Wageningen Journal of Life Sciences 53, 49-69.
Etherton, B., Plex Sula, A., Mouafo-Tchinda, R., Kakuhenzire, R., Kassaye, H., Asfaw, F.,
Kosmakos, V., McCoy, R., Xing, Y., Yao, J., 2025. Translating Ethiopian potato seed
networks: Identifying strategic intervention points for managing bacterial wilt and other
diseases. Agr Syst 222, 104167.
Etherton, B.A., Choudhury, R., Alcalá-Briseño, R., Xing, Y., Sulá, A.P., Carrillo, D.,
Wasielewski, J., Stelinski, L., Grogan, K., Ballen, F., 2023. Are avocados toast? A
framework to analyze decision-making for emerging epidemics, applied to laurel wilt.
Agr Syst 206, 103615.
Etherton, B.A., Choudhury, R.A., Alcalá Briseño, R.I., Mouafo-Tchinda, R.A., Plex Sulá,
A.I., Choudhury, M., Adhikari, A., Lei, S.L., Kraisitudomsook, N., Buritica, J.R., 2024.
Disaster plant pathology: Smart solutions for threats to global plant health from natural
and human-driven disasters. Phytopathology 114, 855-868.
Fall, M.L., Van der Heyden, H., Carisse, O., 2016. A quantitative dynamic simulation of
Bremia lactucae airborne conidia concentration above a lettuce canopy. PLoS One 11,
e0144573.
Fenu, G., Malloci, F.M., 2021. Forecasting plant and crop disease: an explorative study on
current algorithms. Big Data and Cognitive Computing 5, 2.
Ferdu, A., Solbreck, C., 2010. Oviposition preference and larval performance of the sweet
potato butterfly *Acraea acerata* on *Ipomoea* species in Ethiopia. Agricultural and Forest
Entomology 12, 161-168.
Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1‑km spatial resolution climate surfaces
for global land areas. International Journal of Climatology 37, 4302-4315.
Gallego‑Tévar, B., Gil‑Martínez, M., Perea, A., Pérez‑Ramos, I.M., Gómez-Aparicio, L.,
2024. Interactive Effects of Climate Change and Pathogens on Plant Performance: A
Global Meta‑Analysis. Global Change Biology 30, e17535.

Garrett, K., Bebber, D., Etherton, B., Gold, K., Sulá, A.P., Selvaraj, M.G., 2022a. Climate
change effects on pathogen emergence: artificial intelligence to translate big data for
mitigation. Annual Review of Phytopathology 60, 357-378

Garrett, K.A., 2021. Impact network analysis and the INA R package: Decision support for
regional management interventions. Methods in Ecology and Evolution 12, 1634-1647.

Garrett, K.A., Thomas-Sharma, S., Forbes, G.A., Hernandez Nopsa, J.F., Plex Sulá, A.I.,
2022b. Climate change and plant pathogen invasions. In: Ziska, L. (Ed.), Invasive
Species and Global Climate Change. CABI, pp. 22-49.

Gotor, E., Di Cori, V., Pagnani, T., Kikulwe, E., Kozicka, M., Caracciolo, F., 2022. Public
and private investments for banana Xanthomonas Wilt control in Uganda: the economic
feasibility for smallholder farmers. African Journal of Science, Technology, Innovation
and Development 14, 135-146.

Hausfather, Z., 2019. CMIP6: the next generation of climate models explained. Carbon
Brief.

Hijmans, R., Barbosa, M., Ghosh, A., Mandel, A., 2023. geodata: Download geographic
data. R package version 0.5.

Hijmans, R.J., 2024. terra: Spatial Data Analysis. R package version 1.8-0.

Hijmans, R.J., Bivand, R., Forner, K., Ooms, J., Pebesma, E., Sumner, M.D., 2022.
Package 'terra'. Maintainer: Vienna, Austria.

Hodkinson, I.D., 2005. Terrestrial insects along elevation gradients: species and
community responses to altitude. Biological reviews 80, 489-513.

Hu, J.S., Wang, M., Sether, D., Xie, W., Leonhardt, K.W., 1996. Use of polymerase chain
reaction (PCR) to study transmission of banana bunchy top virus by the banana aphid
(*Pentalonia nigronervosa*). Annals of Applied Biology 128, 55-64.

International Food Policy Research Institute, M., 2020. Spatially-disaggregated crop
production statistics data in Africa south of the Sahara for 2017. Harvard Dataverse.

Jongejans, E., Skarpaas, O., Ferrari, M.J., Long, E.S., Dauer, J.T., Schwarz, C.M.,
Rauschert, E.S.J., Jabbour, R., Mortensen, D.A., Isard, S.A., Lieb, D.A., Sezen, Z.,
Hulting, A.G., Shea, K., 2015. A unifying gravity framework for dispersal. Theoretical
Ecology 8, 207-223.

Juroszek, P., Bartsch, L., Fontaine, J.F., Racca, P., Kleinhenz, B., 2022. Summary of the
worldwide available crop disease risk simulation studies that were driven by climate
change scenarios and published during the past 20 years. Plant Pathology 71, 1815–
1838.

Kaundal, R., Kapoor, A.S., Raghava, G.P., 2006. Machine learning techniques in disease
forecasting: a case study on rice blast prediction. BMC bioinformatics 7, 1-16.

Keshav, K., Plex-Sula, A., Garrett, K.A., 2024. geohabnet: Geographical Risk Analysis
Based on Habitat Connectivity. R Package, Version 2.0.0. CRAN.

Kreuze, J., Adewopo, J., Selvaraj, M., Mwanzia, L., Kumar, P.L., Cuellar, W.J., Legg, J.P.,
Hughes, D.P., Blomme, G., 2022. Innovative digital technologies to monitor and control
pest and disease threats in root, tuber, and banana (RT&B) cropping systems: Progress
and prospects. Root, Tuber and Banana Food System Innovations: Value Creation for
Inclusive Outcomes, pp. 261-288.

Kroschel, J., Okonyaet, J., Juarez, H., Forbes, G., Kreuze, J., Beed, F., Blomme, G., Legg,
J., 2014. Management of critical pests and diseases in RTB crops under changing

climates, through risk assessment, surveillance and modeling. RTB Workshop Reports;
CGIAR Research Program on Roots, Tubers and Bananas (RTB): Lima, Peru, p. 20.
Kruskal, J.B., 1964. Nonmetric multidimensional scaling: a numerical method.
Psychometrika 29, 115-129.
Kuhn, M., 2008. Building predictive models in R using the caret package. Journal of
Statistical Software 28, 1-26.
Legg, J., Bouwmeester, H., Bigirimana, S., Tata-Hangy, W., Obiero, H., Gashaka, G.,
Mkamilo, G., Ndyetabula, I., Jeremiah, S., Alicai, T., 2009. Cassava Disease
Surveillance Surveys 2009. Mapping Report, Great Lakes Cassava Initiative. p. 46.
Legg, J.P., Owor, B., Sseruwagi, P., Ndunguru, J., 2006. Cassava mosaic virus disease in
East and Central Africa: epidemiology and management of a regional pandemic.
Advances in Virus Research 67, 355-418.
Loxdale, H.D., 2018. Aspects, including pitfalls, of temporal sampling of flying insects, with
special reference to aphids. Insects 9, 153.
Makiola, A., Holdaway, R.J., Wood, J.R., Orwin, K.H., Glare, T.R., Dickie, I.A., 2021.
Environmental and plant community drivers of plant pathogen composition and
richness. The New Phytologist 233, 496-504.
Malumphy, C., Eyre, D., Anderson, H., 2017. Tobacco, sweet potato or silver leaf whitefly:
Bemisia tabaci. Department for Environment Food and Rural Affairs, United Kingdom.
Hal, p. 6.
Margosian, M.L., Garrett, K.A., Hutchinson, J.S., With, K.A., 2009. Connectivity of the
American agricultural landscape: assessing the national risk of crop pest and disease
spread. BioScience 59, 141-151.
McCulloch, G.A., Waters, J.M., 2023. Rapid adaptation in a fast‑changing world: Emerging
insights from insect genomics. Global Change Biology 29, 943-954.
McEwan, M., 2016. Sweetpotato seed systems in sub-saharan Africa: A literature review to
contribute to the preparation of conceptual frameworks to guide practical interventions
for root, tuber and banana seed systems. RTB Working Paper. CGIAR Research
Program on Roots, Tubers and Bananas (RTB), Apartado 1558, Lima 12, Peru, p. 45.
Mishra, K., Subbanna, A., Rajashekara, H., Paschapur, A.U., Jeevan, B., Singh, A.K.,
Maharana, C., 2024. Perceptions on Disease and Pest Status of Major Cultivated Crops
in Indian Himalayas Under Changing Climate. Adapting to Climate Change in
Agriculture-Theories and Practices: Approaches for Adapting to Climate Change in
Agriculture in India, pp. 121-141.
Mouafo-Tchinda, R.A., Beaulieu, C., Fall, M.L., Carisse, O., 2021. Effect of temperature on
aggressiveness of *Plasmopara viticola* f. sp. *aestivalis* and *P. viticola* f. sp. *riparia* from
eastern Canada. Canadian Journal of Plant Pathology 43, 73-87.
Mouafo-Tchinda, R.A., Etherton, B.A., Plex Sulá, A.I., Andrade-Piedra, J., Ogero, K.,
Omondi, B.A., McEwan, M.A., Tene Tayo, P.M., Harahagazwe, D., Cherinet, M., 2024.
Pathogen and pest risks to vegetatively propagated crops in humanitarian contexts:
Towards a national plant health risk analysis for Cameroon and Ethiopia. bioRxiv,
2024.2002.2012.580019.
Mouafo-Tchinda, R.A., Fall, M.L., Beaulieu, C., Carisse, O., 2022. Competition Between
Plasmopara viticola Clade *riparia* and Clade *aestivalis*: A Race to Lead Grape Downy
Mildew Epidemics. Plant Disease 106, 2866-2875.

Nakato, G.V., Okonya, J.S., Kantungeko, D., Ocimati, W., Mahuku, G., Legg, J.P.,
Blomme, G., 2023. Influence of altitude as a proxy for temperature on key *Musa* pests
and diseases in watershed areas of Burundi and Rwanda. Heliyon 9.
Ngatat, S., Hanna, R., Doumtsop Fotio, A.R., Lienou, J.A., Nanga Nanga, S., Fotso Kuate,
A., Osundahunsi, B., Fiaboe, K.K., Ndemba, B., Dossa, G.S., 2024. Distribution and
diversity of emergent Banana bunchy top virus infecting banana and plantain in
Cameroon, Central Africa. Journal of Phytopathology 172, e13279.
Nyang'au, D., Atandi, J., Cortada, L., Nchore, S., Mwangi, M., Coyne, D., 2021. Diversity
of nematodes on banana (*Musa* spp.) in Kenya linked to altitude and with a focus on the
pathogenicity of *Pratylenchus goodeyi*. Nematology 24, 137-147.
Ochieng, J., Knerr, B., Owuor, G., Ouma, E., 2015. Agricultural commercialization and
household food security: The case of smallholders in Great Lakes Region of Central
Africa. International Association of Agricultural Economists, Milan, Italy 212588, p. 30.
Ochieng, J., Ouma, E., Birachi, E., 2014. Gender participation and decision making in crop
management in Great Lakes Region of Central Africa. Gender, Technology and
Development 18, 341-362.
Ocimati, W., Ogwal, G., Tazuba, A.F., Kubiriba, J., Tugume, J., Erima, R., Okurut, W.,
Mahuku, G., Kutunga, D., Blomme, G., 2024. Mapping the vulnerability of banana
production landscapes in Uganda to banana bunchy top disease. Frontiers in Agronomy
6, 1401478.
Ogot, B.A., Niane, D.T., 1984. The Great Lakes Region. General history of Africa IV: Africa
from the twelfth to the sixteenth century, 498-524.
Okonya, J.S., Gamarra, H., Nduwayezu, A., Bararyenya, A., Kroschel, J., Kreuze, J., 2021.
Serological survey and metagenomic discovery of potato viruses in Rwanda and
Burundi reveals absence of PVY in Burundi and first report of TRV in potatoes in sub-
Saharan Africa. Virus Research 302, 198487.
Okonya, J.S., Kroschel, J., 2013. Incidence, abundance and damage by the sweet potato
butterfly (*Acraea acerata* Hew.) and the African sweet potato weevils (*Cylas* spp.)
across an altitude gradient in Kabale district, Uganda. Int J AgriScience 3, 814-824.
Olivoto, T., Lúcio, A.D.C., 2020. metan: An R package for multi‑environment trial analysis.
Methods in Ecology and Evolution 11, 783-789.
Ong'ayo, A.O., 2018. Displacement and cross-border mobility in the Great Lakes region-
re-thinking underlying factors and implications for regional management of migration.
Africa Insight 48, 62-85.
Pautasso, M., Döring, T.F., Garbelotto, M., Pellis, L., Jeger, M.J., 2012. Impacts of climate
change on plant diseases—opinions and trends. European Journal of Plant Pathology
133, 295-313.
Pinnegar, J.K., Hamon, K.G., Kreiss, C.M., Tabeau, A., Rybicki, S., Papathanasopoulou,
E., Engelhard, G.H., Eddy, T.D., Peck, M.A., 2021. Future socio-political scenarios for
aquatic resources in Europe: A common framework based on shared-socioeconomic-
pathways (SSPs). Frontiers in Marine Science 7, 568219.
Poppinga, S., Weisskopf, C., Westermeier, A.S., Masselter, T., Speck, T., 2016. Fastest
predators in the plant kingdom: functional morphology and biomechanics of suction
traps found in the largest genus of carnivorous plants. AoB Plants 8, plv140.

Poveda, K., Martínez, E., Kersch‑Becker, M.F., Bonilla, M.A., Tscharntke, T., 2012.
Landscape simplification and altitude affect biodiversity, herbivory and Andean potato
yield. Journal of Applied Ecology 49, 513-522.
Pryke, A., Mostaghim, S., Nazemi, A., 2007. Heatmap visualization of population based
multi objective algorithms. Evolutionary Multi-Criterion Optimization: 4th International
Conference, EMO 2007, Matsushima, Japan, March 5-8, 2007. Proceedings 4.
Springer, pp. 361-375.
R Core Team, R., 2024. R: A language and environment for statistical computing. R
Foundation for Statistical Computing.
Ramcharan, A., McCloskey, P., Baranowski, K., Mbilinyi, N., Mrisho, L., Ndalahwa, M.,
Legg, J., Hughes, D.P., 2019. A mobile-based deep learning model for cassava disease
diagnosis. Frontiers in plant science 10, 272.
RTB, C., 2013. Workshop on tools for managing seed degeneration in RTB crops (5-8
February 2013, Arusha, Tanzania). CRP-RTB, p. 23.
Rybicki, E.P., 2015. A Top Ten list for economically important plant viruses. Archives of
virology 160, 17-20.
Savary, S., Willocquet, L., Elazegui, F.A., Teng, P.S., Van Du, P., Zhu, D., Tang, Q.,
Huang, S., Lin, X., Singh, H., 2000. Rice pest constraints in tropical Asia:
characterization of injury profiles in relation to production situations. Plant Disease 84,
341-356.
Science Group Project on Accelerated Varietal Improvement and Seed Systems in Africa,
2024. CGIAR project on Accelerated Varietal Improvement and Seed Systems in Africa:
Annual Technical Report 2023.
Selvaraj, M.G., Vergara, A., Ruiz, H., Safari, N., Elayabalan, S., Ocimati, W., Blomme, G.,
2019. AI-powered banana diseases and pest detection. Plant Methods 15, 1-11.
Simpson, G., 2020. ggvegan; ggplot2-based plots for vegan package. version 0.1. 0. In:
https://github.com/gavinsimpson/ggvegan; consulted: May.
Singh, B.K., Delgado-Baquerizo, M., Egidi, E., Guirado, E., Leach, J.E., Liu, H., Trivedi, P.,
2023. Climate change impacts on plant pathogens, food security and paths forward.
Nature Reviews Microbiology 21, 640–656.
Sparks, A.H., 2018. nasapower: a NASA POWER global meteorology, surface solar
energy and climatology data client for R. Journal of Open Source Software 3, 1035.
Sparks, A.H., Forbes, G.A., Hijmans, R.J., Garrett, K.A., 2014. Climate change may have
limited effect on global risk of potato late blight. Global Change Biology 20, 3621-3631.
Sseruwagi, P., Sserubombwe, W., Legg, J.P., Ndunguru, J., Thresh, J., 2004. Methods of
surveying the incidence and severity of cassava mosaic disease and whitefly vector
populations on cassava in Africa: a review. Virus Research 100, 129-142.
Taberlet, P., Coissac, E., Hajibabaei, M., Rieseberg, L.H., 2012. Environmental DNA.
Molecular Ecology 21.
Thies, C., Tscharntke, T., 1999. Landscape structure and biological control in
agroecosystems. Science 285, 893-895.
Wells, K., Clark, N.J., 2019. Host specificity in variable environments. Trends in
Parasitology 35, 452-465.
Were, H., Kabira, J., Kinyua, Z., Olubayo, F., Karinga, J., Aura, J., Lees, A., Cowan, G.,
Torrance, L., 2013. Occurrence and distribution of potato pests and diseases in Kenya.
Potato Research 56, 325-342.

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New
York. ISBN 978-3-319-24277-4.
Wieczorek, K., Fulcher, T.K., Chłond, D., 2019. The composition of the aphid fauna
(Insecta, Hemiptera) of the Royal Botanic Gardens, Kew. Scientific Reports 9, 10000.
WorldData.info, 2024. Climate comparison between Rwanda and Burundi based on a long-
term analysis of the last 20 years.
Xiao, Q., Li, W., Chen, P., Wang, B., 2018. Prediction of crop pests and diseases in cotton
by long short term memory network. Intelligent Computing Theories and Application:
14th International Conference, ICIC 2018, Wuhan, China, August 15-18, 2018,
Proceedings, Part II 14. Springer, pp. 11-16.
Xing, Y., Hernandez Nopsa, J.F., Andersen, K.F., Andrade-Piedra, J.L., Beed, F.D.,
Blomme, G., Carvajal-Yepes, M., Coyne, D.L., Cuellar, W.J., Forbes, G.A., Kreuze, J.F.,
Kroschel, J., Kumar, P.L., Legg, J.P., Parker, M., Schulte-Geldermann, E., Sharma, K.,
Garrett, K.A., 2020. Global cropland connectivity: A risk factor for invasion and
saturation by emerging pathogens and pests. BioScience 70, 744-758.
Yamamura, K., Kiritani, K., 1998. A simple method to estimate the potential increase in the
number of generations under global warming in temperate zones. Applied Entomology
and Zoology 33, 289-298.
1022