# Pathogen and pest communities in food security crops across climate

# gradients: Anticipating future challenges in the Great Lakes region of Africa

Romaric A. Mouafo-Tchinda[1,2,3]*, Aaron I. Plex Sulá[1,2,3], Berea A. Etherton[1,2,3], Joshua S. Okonya[4],

Gloria Valentine Nakato[5], Yanru Xing[1,2,3], Jacobo Robledo Buritica[1,2,3], Ashish Adhikari[1,2,3], Guy

Blomme[6], Déo Kantungeko[7], Anastase Nduwayezu[8], Jan Kreuze[9], Jürgen Kroschel[9,10], James

Legg[11], and Karen A. Garrett[1,2,3]*

[1] Plant Pathology Department, University of Florida, Gainesville, FL, USA
[2] Global Food Systems Institute, University of Florida, Gainesville, FL, USA
[3] Emerging Pathogens Institute, University of Florida, Gainesville, FL, USA
[4] Association for Strengthening Agricultural Research in Eastern and Central Africa, Entebbe,
Uganda
[5] International Institute of Tropical Agriculture, Arusha, Tanzania
[6] Bioversity International, Addis Ababa, Ethiopia
[7] International Institute of Tropical Agriculture, Bujumbura, Burundi
[8] Rwanda Agriculture Development and Animal Resources Board (RAB), Ruhengeri, Rwanda
[9] International Potato Center (CIP), Lima, Peru
[10] Hans-Ruthenberg-Institute for Tropical Agricultural Sciences, University of Hohenheim,
Stuttgart, Germany
[11] International Institute of Tropical Agriculture, Dar es Salaam, Tanzania
20
21
22 * Corresponding authors
23

Emails: rmouafotchinda@ufl.edu*, karengarrett@ufl.edu*

**Keywords:** Climate matching | crop pathogens and pests | cropland connectivity | food security

| machine learning |vegetatively propagated crops

Abstract

Natural and human-driven disasters are a significant challenge to the sustainable production of

food security crops in the Great Lakes region of Africa. A relevant research gap is the effect of

climate change on the distribution of pathogens and pests of food security crops in this region.

We evaluated the current geographic risk of pathogens and pests in the production of four food

security crops — banana, cassava, potato, and sweetpotato — in the region, and the potential

future risk under climate change. First, cropland connectivity analysis identified locations likely

important in the spread and establishment of crop-specific pathogens and pests for each crop,

with locations in Rwanda and Burundi emerging as important across crops. Second, we surveyed

27 economically important pathogens and pests in Rwanda and Burundi, mapping their

distribution across landscapes and quantifying patterns of association. Cropland density,

cropland connectivity, altitude, and contemporary and long-term temperature and precipitation

were strongly associated with disease severity and pest infestation. Among ten machine-learning

algorithms evaluated, random forests and support vector machines generally performed best for

predicting severity and infestation. Third, an increase in temperature across altitudes is

projected under future climate change scenarios in this region. We found evidence that 44% of

the pathogen and pest species we studied in banana, potato, and sweetpotato could become

more common with warmer temperatures at higher altitudes, while 17% may become less

common. Ongoing development of pathogen and pest forecasts can guide strategic surveillance,

mitigation and adaptation to future epidemics and pest invasions in high-risk crop locations

under climate change.

# 1 | INTRODUCTION

## 1.1 | Impact of climate and crop landscape structure on pathogen and pest damage

The Great Lakes region of Africa faces recurrent disasters, exacerbated by climate variability, fragile ecosystems, and societal dynamics including demographic pressures, socio-political instability, and resource-related conflicts (Ong'ayo, 2018, Ogot & Niane, 1984). The geographic structure of cropland in this region, including crop density, cropland connectivity, and crop diversity, is influenced by heterogeneous agricultural practices, creating local habitats that can support the spread of pathogens and pests. Suitable climate factors such as rainfall, temperature, and relative humidity are essential to achieving optimal plant growth and maximum crop yields. However, climate change is likely to affect agricultural productivity and exacerbate the proliferation of some pathogens and pests in this region (Okonya & Kroschel, 2013, Ebregt *et al.*, 2005, Ferdu & Solbreck, 2010, Gallego–Tévar *et al.*, 2024). It reshapes ecosystems by altering temperature and precipitation patterns, driving pathogens and pests into new locations (Cilas *et al.*, 2016, Garrett *et al.*, 2022a, Garrett *et al.*, 2022b, Sparks *et al.*, 2014). The interactions among climate variability, the geographic structure of cropland, and the biology of pathogens and pests in this region remain poorly understood and represent a major knowledge gap. Although research has examined how climate change affects the dynamics of specific pathogens and pests (Adhikari *et al.*, 2015, Blomme *et al.*, 2020, Kroschel *et al.*, 2014), little is known about how regional factors—such as the geographic structure of cropland of key food security crops in the Great Lakes region—exacerbate these risks, including the potential for altitudinal shifts of pathogen and pest communities under climate change.

Climate matching identifies which locations have similar climate conditions, which may be

equally suitable for a species; ecological communities may move following changes in where

suitable climate conditions are found in the future (Colwell *et al.*, 2008, Chevalier *et al.*, 2024,

Singh *et al.*, 2023). For example, pathogen and pest communities now at hotter, lower altitudes

may move to what are currently cooler, higher altitudes if there is warming due to climate

change. The dynamics of pathogens and pests across climate gradients have been widely studied

(Hodkinson, 2005, Mishra *et al.*, 2024, Juroszek *et al.*, 2022, Pautasso *et al.*, 2012). Higher

altitudes are commonly associated with lower species richness, disease severity, and pest

damage (Blomme et al., 2020, Hodkinson, 2005). Since altitude is typically negatively correlated

with temperature (Poveda *et al.*, 2012), altitude can be a useful proxy when evaluating the effect

of temperature on crop disease and pest damage. For example, previous studies in South

America and East Africa predicted that rising temperatures due to climate change will modify the

geographic distribution of pathogens and pests, increasing crop disease severity or pest damage

at higher altitudes (Blomme et al., 2020, Poveda et al., 2012). New climate conditions favorable

to certain pathogens and pests are likely to occur at higher altitudes, disrupting ecological

balances and exposing high-altitude crops and ecosystems to new risks.

The geographic distribution of crop production can also influence geographic patterns of disease

severity and pest damage. Crop landscape structure has been well documented to influence crop

damage from diseases and pests (Poveda et al., 2012, Thies & Tscharntke, 1999). Analysis of the

network structure of crop landscapes, such as cropland connectivity analysis, has been used to

identify candidate priority locations for surveillance and management of crop-specific pathogens

and pests (Xing *et al.*, 2020, Andersen Onofre *et al.*, 2021, Buddenhagen *et al.*, 2022). A general

assessment of cropland geographic structure helps us understand the potential introduction,

establishment, and spread of pathogens and pests. Higher crop density increases the local risk of

crop-specific pathogen and pest establishment, while greater cropland connectivity also

increases the risk of spread and establishment.

**1.2 | Disaster plant pathology: Vulnerabilities and solutions in the Great Lakes region**

Many families living in the Great Lakes region rely on food security crops such as banana and

plantain, cassava, potato, and sweetpotato. However, many households remain food insecure

because of declining agricultural productivity, driven by frequent disasters in the region,

including conflicts, health emergencies, and natural disasters that displace populations and

disrupt agriculture (Ochieng *et al.*, 2015, Ochieng *et al.*, 2014, Ong'ayo, 2018). These challenges

are compounded by the cascading effects of high prevalence of pests and diseases, low

agricultural input use, loss of harvest and assets, dysfunctional or non-existent markets, high

food prices, and socioeconomic conflicts (McEwan, 2016, Almekinders *et al.*, 2019, Kroschel et

al., 2014). The Rwandese genocide in 1994, the civil war in Burundi from 1993 to 2005 and also

in 2015, and the persistent conflicts in Eastern DR Congo are particularly important humanitarian

disasters in the Great Lakes region, which have left an indelible mark (Ong'ayo, 2018). Health

emergencies and natural disasters, such as severe droughts in Ethiopia, Kenya, and Uganda

(2015-2016) and devastating floods in Rwanda, Uganda, and DR Congo (2020), have severely

restricted agricultural practices such as effective surveillance and management of fields, causing

considerable damage to infrastructure and cropland (Ochieng et al., 2015, Ochieng et al., 2014).

Disaster plant pathology, integrating concepts from plant pathology and disaster science,

addresses the risks and potential impact of plant diseases before, during, and after disaster

events that threaten agricultural systems and ecosystems (Etherton *et al.*, 2024, Mouafo-

Tchinda *et al.*, 2024). It focuses on disturbances that modify host-pathogen dynamics and

increase damage and the spread of pathogens, and the development of strategies for better

managing plant health.  A combination of global and national research efforts has been adopted

to manage the impacts that these disasters, pathogens, and pests may have on agroecological

systems (RTB, 2013, Science Group Project on Accelerated Varietal Improvement and Seed

Systems in Africa, 2024). For example, disease control programs which encourage farmers to

recognize and destroy diseased plants, and disease-resistant crop breeding programs are

currently underway across the Great Lakes region.

When models of pathogen and pest responses to environmental variables are available,

decision-support tools can be developed to predict where specific disease- and pest-resistant

varieties will be needed in the long-run, and in the short-run whether likely economic injury

levels motivate management actions by growers (Fenu & Malloci, 2021, Kaundal *et al.*, 2006,

Carisse & Fall, 2021). Pest and disease forecasting can reduce production costs, pesticide use,

and environmental pollution, protecting the health of farmers and consumers. Machine learning

can analyze high volume datasets, and improve the predictive power, accuracy, and efficiency

for diseases and pests (Garrett et al., 2022a, Kaundal et al., 2006). Implementing decision

support systems in low-resource countries such as those in the Great Lakes region is challenging;

in addition to the often-limited availability of high-resolution local weather data as input to

models, field observations and georeferenced datasets for the many important pathogen and

pest species are often unavailable for constructing predictive models.

## 1.3 | Objectives

In this study, we evaluated the geographic risk of pathogens and pests in the production of four food security crops — banana (including plantain), cassava, potato, and sweetpotato — in the Great Lakes region of Africa, including the potential future risk under climate change. The first objective was to analyze the geographic risk of crop-specific pathogen and pest spread based on crop landscape connectivity for these crops. The second objective was to characterize the seasonal and geographic occurrence and associations of 27 economically important pathogens and pests affecting these crops, through field surveys conducted in two key locations in the region, Rwanda and Burundi. Part of this dataset has been used to characterize the prevalence of potato viruses and vectors (Okonya *et al.*, 2021) and the occurrence of banana pests and diseases across altitudes (Nakato *et al.*, 2023).  The third objective was to predict pathogen and pest severity/infestation based on climate variables and crop landscape structure using machine learning, and to assess which pathogens and pests are likely to experience altitudinal range shifts under projected changes in temperature. The field surveys were implemented along environmental gradients to compare low- and high-altitude pathogen and pest communities, where lower-altitude communities have the potential to move toward higher altitudes in the future. These risk assessments can support decision-making by (i) national agricultural agencies that plan geographic surveillance strategies for economically important pathogens and pests, (ii) local agricultural research communities and international partners as they prioritize efforts to improve current and future mitigation responses to pathogens and pests, and (iii) farmer communities that need to be prepared for changes in pathogen and pest dynamics due to natural or human-driven disasters or changing climates.

2 | METHODS

2.1 | Landscape structures influencing pathogen and pest risk

2.1.1 | Identifying locations with high cropland connectivity

The spatial pattern of host availability is a key ecological trait that influences the spread of crop-

specific pathogens and pests. We analyzed the connectivity of crop landscapes in the Great Lakes

region to map the potential establishment and regional spread of crop-specific pathogens and

pests, to identify candidate priority locations for spatially targeted surveillance in four food

security crops — banana (including plantain), cassava, potato, and sweetpotato. This cropland

connectivity analysis assesses how crop production is connected in a landscape, based on host

availability and physical proximity in a gravity model (Jongejans *et al.*, 2015). Cropland

connectivity analysis identifies potential habitat networks through which pests and pathogens

can spread between croplands, a first approximation to epidemic or invasion networks, which

can be improved as more data become available (Margosian *et al.*, 2009, Xing et al., 2020,

Andersen Onofre et al., 2021).

For each crop, we used publicly available estimates of the cropland harvested area (or cropland

density) across sub-Saharan Africa in 2017 (International Food Policy Research Institute, 2020) as

a proxy for host availability to crop-specific pathogens and pests. Maps of cropland density,

aggregated at a spatial resolution of 10-min (or grid cells of image 9.27 km × 9.27 km), were used

to evaluate the potential epidemic or invasion network using a gravity model (Xing et al., 2020).

Here the gravity model incorporated two pathogen or pest dispersal kernels: $c_i c_j d_{ij}^{-\beta}$ for the

inverse power law model, and $c_i c_j e^{-\gamma d_{ij}}$ for the negative exponential model. In each dispersal

model, $d_{ij}$ is the Vincenty ellipsoid distance between a pair of cropland nodes (locations) $i$ and $j$,

$c_i$ is the cropland density for node $i$, and $\beta$ and $\gamma$ are dispersal parameters describing the

relative likelihood of pathogen or pest movement across $d_{ij}$ (Xing et al., 2020, Andersen Onofre

et al., 2021).

We calculated a cropland connectivity risk index (CCRI) (Xing et al., 2020) for each location with

crop production in the region as a measure of its likely importance in epidemic or invasion

networks. The CCRI was the weighted mean of four network metrics (node centralities),

evaluated using the geohabnet package v2.0.0 (Keshav *et al.*, 2024) in R v4.4.0 (R Core Team,

2024), averaged across the results from a sensitivity analyses for a range of dispersal

parameters, as described by Mouafo-Tchinda et al. (2024). Two CCRI formulations were used to

capture key facets of node connectivity. The first CCRI considered ("CCRI-betweenness") was the

mean of betweenness centrality weighted by 1/2, node strength weighted by 1/6, eigenvector

centrality weighted by 1/6, and the sum of nearest neighbors' degrees weighted by 1/6. The

second, "CCRI-closeness", considered 1/2 closeness centrality instead of betweenness centrality,

with the same weights for the other metrics. We emphasized betweenness or closeness

centrality to account for, respectively, the importance of a node as a bridge between parts of the

network and the relative facility with which a location can be reached from all other locations in

the network.

Habitat connectivity analysis using the geohabnet package (Keshav et al., 2024) is part of the

R2M toolbox for rapid risk assessment supporting mitigation of pathogens and pests

([garrettlab.com/r2m](garrettlab.com/r2m)). Other examples of applications of habitat connectivity and R2M tools, in

addition to those discussed elsewhere in this paper, illustrate how these tools can be used in an

integrated analysis (Mouafo-Tchinda et al., 2024, Andersen Onofre *et al.*, 2019, Etherton *et al.*,

2025, Etherton *et al.*, 2023, Garrett, 2021).

### 2.1.2 | Associations among altitude, crop landscape structure, and climate

We characterized associations among altitude, crop landscape structure, and climate in the

Great Lakes region. The map of altitudes was retrieved from the Shuttle Radar Topography

Mission (SRTM) database, using the geodata package v0.5-8 (Hijmans *et al.*, 2023) in R. We also

retrieved the 19 bioclimatic variables for the region from Worldclim database v2.1 (Fick &

Hijmans, 2017) using the geodata package version 0.5-8 (Hijmans et al., 2023) and the terra

package v1.8-0 (Hijmans, 2024) in R. For crop landscape structure variables, we used the raster

maps of cropland density and connectivity (CCRI-closeness) generated in section 2.1.1, using the

terra package (Hijmans *et al.*, 2022). We assessed the associations between altitude, bioclimatic

variables, and cropland structure by calculating the Pearson correlations between these

variables using the metan package v1.18.0 (Olivoto & Lúcio, 2020) in R.

### 2.2 | Pathogen and pest communities

### 2.2.1 | Field sampling and spatiotemporal structure of pathogen and pest communities

For each crop species, we identified fields for sampling along four survey transects in Rwanda

and Burundi, each transect representing an altitude gradient. In Burundi, the field surveys

included cropland around Lake Tanganyika in the Rusizi watershed, representing the Bubanza,

Bujumbura Rural, Cibitoke, and Muramvya provinces (Figure 1C). In Rwanda, the field surveys

were in major cropland around Lake Kivu in the Ruhengeri watershed, which includes part of the

Northern and Western provinces (Figure 1C). The survey locations represented major crop

landscapes in both countries, encompassing a mixture of cropping systems and a range in

altitude from 900 to 2800 m.a.s.l. (Figure 1a-b) (Kroschel et al., 2014). Fields for sampling were

selected at regular intervals along the transects regardless of whether diseases or pest damage

were apparent. The number of fields sampled in Burundi and Rwanda was 189 and 103 for

banana, 78 and 116 for potato, and 107 and 104 for sweetpotato, respectively. For cassava,

sampling was only carried out in 50 fields in Burundi, with the second Burundi survey not

conducted due to civil unrest, and Rwanda was not sampled because few farmers grew cassava

in the Rwanda study area.

Sampling was conducted in each watershed during two cropping seasons, dry and rainy. In both

countries, two rainy seasons occur: one from March to May and the other from September to

December (WorldData.info, 2024). The dry seasons occur from June to August and January to

February (WorldData.info, 2024). For each field sampled, we evaluated crop-specific pest or

disease intensity (Table 2) based on prevalence in the region, disease severity, infestation rate

and intensity, and/or the number of pests, using methods as described by Nakato et al. (2023),

Okonya et al. (2021), and Legg *et al.* (2009). A total of 27 economically important pathogens and

pests in the region were included in this study, including 6 for banana, 6 for cassava, 7 for potato

and 8 for sweetpotato (Table 2). Here prevalence refers to the presence or absence of a pest or

disease in a field, while disease severity, pest infestation intensity or infestation rate is the

percentage of leaves or plants damaged by a disease or pest. For infestation rate, infestation

intensity, and severity, twenty plants per field, selected in a systematic pattern, were assessed

for visual symptoms of damage by pests and diseases. The percentage of plants infested was

noted as the infestation rate. The percentage of symptomatic leaves per plant was noted as

infestation intensity for pests and severity for diseases. For damage by some virus diseases and

pests such as cassava mosaic disease (CMD) and cassava green mite (CGM), crop damage was

scored from 1 (representing no symptoms) to 5 (the most severe) (Legg et al., 2009, Okonya et

al., 2021, Sseruwagi *et al.*, 2004). The disease or pest damage score measures the severity or

infestation intensity in a field, based on visual assessments and a predefined scale. Thirty plants

were randomly sampled at regular intervals along the season and damage was assessed by

scoring each plant on a scale of 1 to 5 as described by Legg et al. (2009).

For banana, fields were surveyed in Burundi in March 2015 (during the rainy season) and in July

2016 (dry season), while fields were surveyed in Rwanda in July 2015 (dry season) and in

November 2016 (rainy season). Cassava fields were surveyed only once in Burundi in March 2015

FIGURE 1| Location of fields of banana, cassava, potato, and sweetpotato where pathogen and pest surveys were carried out in Africa (A), in the Great Lakes region (B), in transects across altitude gradients, in Rwanda and Burundi (C). White lines in (C) indicate the province boundaries of Rwanda and yellow lines indicate the province boundaries of Burundi. Crosses and circles represent locations surveyed during the dry and rainy season, respectively. The complete surveillance effort included 412 crop fields.

(rainy season), for reasons described above. For potato, field sampling was conducted in Burundi

in July 2016 (dry season) and December 2017 (rainy season), and in Rwanda in June-July 2015

(dry season) and November-December 2017 (rainy season). For sweetpotato, the field surveys

took place in Burundi in November-December 2015 (rainy season) and August 2016 (dry season),

and in Rwanda in July 2015 (dry season) and November 2017 (rainy season).  The irregular dates

for these crop-field surveys were due in part to the civil unrest prevailing in the region at the

time the study was conducted.

2.2.2 | Quantifying patterns of field-scale association of crop pathogens and pests

We evaluated the patterns of association of pathogens and pests based on prevalence and

severity/infestation using non-metric multidimensional scaling (NMDS) (Kruskal, 1964) and

heatmaps (Pryke *et al.*, 2007) for visualization. We initially calculated five measures of

dissimilarity (Bray, Manhattan, Jaccard, Chao, and robust-Aitchison) in the NMDS analysis using

the ggvegan package v0.1.999 (Simpson, 2020) in R. We selected the Bray distance in the final

analysis due to its highest goodness of fit. A heatmap was prepared using the metan package

v1.18.0 (Olivoto & Lúcio, 2020) in R to map the association of pathogens and pests based on

prevalence and severity/infestation rates.

2.3 | Prediction of pathogen and pest damage and geographic range shifts

2.3.1 | Machine learning to predict pathogen and pest levels

Weather, climate, and geographic variables were used to predict severity or infestation

rates/intensity, comparing machine learning algorithms. We retrieved weather variables

(monthly, quarterly, and annual temperature, precipitation, and relative humidity) from the

NASAPOWER database using the nasapower package v4.0.10 (Sparks, 2018) in R. The climate

variables included the 19 bioclimatic variables, solar radiation (kJ m-2 day-1), wind speed (m s-1),

and water vapor pressure (kPa) described, along with the altitude variable, in section 2.1.2. The

CCRIs and cropland density were assembled as described in section 2.1.1.

Data were preprocessed to exclude collinearity between variables. A stratified split was then

used to divide the data into training and test sets in an 8:2 ratio. We then trained ten machine-

learning algorithms: logistic regression (GLM and GLMNET), support vector machine (SVM, with

linear and radial basis function), neural network (NN), k-nearest neighbors (KNN), stochastic

gradient boosting (GBM, generalized boosted modeling), bagged trees, classification and

regression trees (CART), and random forest (RF). We compared the performance of these

algorithms to predict the severity or infestation of each pathogen and pest using the caret

package v6.0-94 (Kuhn, 2008) in R. We selected SVM (with radial basis function) and RF for

further analyses based on their having the highest performance in terms of MAE (mean absolute

error), RMSE (root mean squared error) and R-squared (Kaundal et al., 2006).

In the next stage using the SVM and RF models, we executed a 10-fold cross-validation with ten

repetitions to prevent overfitting the training model and used a grid search for hyperparameter

tuning. The tuning process was driven by trainControl, with RMSE-based model evaluation to

prioritize accuracy. Hyperparameter tuning for SVM included C (regularization parameter) and

Gamma (kernel coefficient), both with a tuneLength of 10. Hyperparameter tuning for RF

included mtry (the number of predictors randomly selected at each division) with a tuneLength

of 10. We trained the model using the train function in the caret package and estimated the

performance parameters MAE, RMSE, and R-squared. We used the resulting model to predict

severity or infestation rate/intensity for the test data and estimated the performance

parameters (RMSE and R-squared).

### 2.3.2 | Pathogen and pest communities across altitudinal gradients

To study the geographic range of pathogen and pest communities associated with the four crops

in the sample transects in Rwanda and Burundi, we evaluated how pathogen and pest levels

(prevalence, disease severity or infestation rate/intensity) changed with altitude across the

sampling locations. We also evaluated the pathogen and pest richness (number of pathogens

and pests present from those included in the sampling) in communities during the dry and rainy

seasons, across the altitude gradients. The associations between prevalence, infestation/severity

and richness of pathogens and pests across altitude gradients were evaluated based on

Pearson's correlation.

**2.3.3 | Climate change: impact on regional temperature patterns**

We assessed potential climate shifts along the altitude gradients in the Great Lakes region, using

current and future temperature data from WorldClim database v2.1 (Fick & Hijmans, 2017). We

used future climatic variables from three global circulation models (GCMs: GISS-E2-1-G, MIROC6

and CNRM-CM6-1), two Shared Socioeconomic Pathways (SSP1-2.6 and SSP5-8.5), and four

future time-periods: 2021-2040, 2041-2060, 2061-2080, and 2081-2100. We selected these

GCM modeling groups because of their equilibrium climate sensitivity (ECS) (Hausfather, 2019).

ECS indicates the severity of future warming impacts. CNRM-CM6-1 assumes a high value of ECS,

4.9 C, while GISS-E2-1-G and MIROC6 assumes low values of ECS, 2.7 and 2.6 respectively

(Hausfather, 2019). The two SSPs used in this study represent different emission pathways by

the year 2100: sustainability (SSP1-2.6) and fossil-fuel development (SSP5-8.5) emission

scenarios featured by the radiation forcing of 2.6 and 8.5 W/m², respectively (Pinnegar *et al.*,

2021). We calculated the average annual temperature across the three selected GCMs for each

future time-period and SSP combination. We used the resulting density plots to assess potential

shifts in the distribution of annual mean temperature at low (<1500 m.a.s.l.) and high (>1500

m.a.s.l.) altitudes, using the ggplot2 package v3.5.0 (Wickham, 2016) in R.

3 | RESULTS

3.1 | Landscape structures influencing pathogen and pest risk

3.1.1 | Identifying locations with high cropland connectivity

The analysis of cropland connectivity in the Great Lakes region (i) indicates quantitatively how

the geographic distribution of crop production could facilitate the potential spread of crop-

specific pathogens and pests and (ii) identifies candidate priority locations for pathogen and pest

surveillance for each food security crop (Figure 2). These maps of cropland connectivity highlight

locations in the region that are likely important for understanding and managing the risk of crop-

specific pathogens and pests.

In these maps of cropland connectivity, a higher CCRI-betweenness (Figure 2A) or CCRI-closeness

indicates the potential to play an important role in the establishment and spread of crop-specific

pathogens and pests affecting these crops, locations with high CCRI are candidate priorities for

surveillance and mitigation programs. Although CCRI-betweenness and CCRI-closeness may not

yield identical CCRI values, they highlight the same potential locations for prioritization (Figure

2A). The overall values of CCRI-closeness were higher than those of CCRI-betweenness, which

may reflect the greater influence of proximity-based connectivity compared to pathway-based

connectivity in the network. Combining these metrics could provide complementary insights into

prioritizing key locations within the network. For banana, locations in Rwanda, Burundi, the DR

Congo, Uganda, Tanzania, Ethiopia, Malawi, and Mozambique have a high CCRI (Figure 2).

Locations in the southern, eastern, and northern provinces of Rwanda have the highest CCRI-

betweenness (Figure 2A and 2B). For cassava, several locations across the region have a high

CCRI. In Burundi, locations in the Cibitoke, Bubanza, Kirundo, Ngozi, Bujumbura Rural and Gitega

provinces have the highest CCRI (Figure 2A and B). For potato, locations in Rwanda, Burundi, the

DR Congo, Uganda, Tanzania, Ethiopia, and Malawi have high CCRI-betweenness (Figure 2).

These maps also show that locations in the northern, western, and southern provinces of

Rwanda have the highest CCRI-betweenness (Figure 2A and B). For sweetpotato, locations in

Rwanda, Burundi, Uganda, Tanzania, Malawi, and Ethiopia have a high CCRI (Figure 2). These

maps indicate the important role that Rwanda and Burundi play for understanding and

mitigating the risks associated with the spread of pathogens and pests of these crops.

Maps of cropland connectivity (Figure 2B) also illustrate the potential roles of the locations we

selected for field sampling crop pathogens and pests in the Ruhengeri watershed in Rwanda and

in the Rusizi watershed in Burundi. The connectivity of these landscapes varies from crop to

crop. Some sampled locations had high cropland connectivity (e.g., cassava and sweetpotato in

Burundi transects and potato in Rwanda transects) compared to other locations. Overall, the

surveyed locations around Lake Kivu in Rwanda and Lake Tanganyika in Burundi, as well as other

locations in the two countries, are likely to be important candidate locations for surveillance for

the establishment and spread of banana, cassava, potato, and sweetpotato pathogens and pests

in both countries, based on cropland connectivity (Figure 2B).

FIGURE 2 | Maps of a cropland connectivity risk index (CCRI-betweenness) for four food security crops in (A) the Great Lakes region of Africa and (B) in a focused geographical extent in Rwanda and Burundi. The CCRI indicates how important a location is likely to be for potential spread of a host-specific pathogen or pest, based on the network of host availability. The CCRI is a weighted mean of four network metrics, including betweenness centrality in the results shown here. In (A), dark blue lines indicate national boundaries. In (B), blue lines indicate the provinces of Rwanda and green lines indicate the provinces of Burundi.

3.1.2 | Associations among altitude, landscape structure and climate

The associations among (a) altitude, (b) crop landscape structure, and (c) climate variables such

as temperature and precipitation, were analyzed for the Great Lakes region. Higher

temperatures were generally associated with lower altitudes. There was a strong negative

correlation between altitude and most temperature-related bioclimatic variables (Tem, Tmin,

TColQ, TDriQ, TWetQ, TWarQ, Tmax, TColdQ). In contrast, altitude was weakly positively

correlated with some precipitation-related bioclimatic variables such as PrecS, PWetQ, and

PWetM, and negatively correlated with others, including Prec, PDriQ, PWarQ, and PColQ (Figure

3). In this region, higher cropland density and cropland connectivity of banana, cassava, potato,

and sweetpotato (Figure 3) were at higher altitudes (except cassava density). Cropland

connectivity of these crops showed a stronger positive correlation with altitude compared to

cropland density. Cropland density and cropland connectivity for each of these four crops were

negatively correlated with most temperature-related bioclimatic variables and positively

correlated with most precipitation-related bioclimatic variables (Figure 3).

**FIGURE 3 |** Associations between altitude, bioclimatic variables, and crop landscape structure (crop density and cropland connectivity) in the Great Lakes region of Africa, based on Pearson's correlation. Strong positive correlations are in dark blue, while strong negative correlations are in dark red. White boxes without labels indicate no association.

379    3.2 | Pathogen and pest communities

380    3.2.1 | Spatiotemporal structure of pathogen and pest communities

381    Most of the targeted pathogens and pests were found in Burundi and Rwanda in both crop-

382    growing seasons (Figure 4). Sweetpotato mealybugs, sweetpotato butterfly (*Acraea acerata*) and

383    sweetpotato aphids were observed only in Burundi (Figure 4). The observed prevalence of each

FIGURE 4 | Seasonal distribution of observed pathogens and pests in smallholder production of banana, potato and sweetpotato in Burundi and Rwanda. (Cassava data are not represented here). The length of the bars represents the prevalence for both seasons (where a total of 2 indicates presence in every field in both seasons). Abbreviations: Banana bunchy top virus (BBTV), *Cosmopolites sordidus* (CS), *Ralstonia solanacearum* (RS), sweetpotato virus (SPV), and Xanthomonas wilt of banana (XW).

pest or disease often varied by country, crop, and season. In both countries and seasons, the

highest prevalence was observed for *Ralstonia solanacearum* and virus symptoms in potato,

virus symptoms in sweetpotato, and *Cosmopolites sordidus* in banana.

3.2.2 | Associations among pathogens and pests

The NMDS analysis indicated few pathogens/pests whose levels are associated, although some

were common enough that they frequently co-occurred with others (Figure 5). A few pathogens

and pests were associated in terms of prevalence and severity/infestation in the crop fields.

Among banana diseases/pests, Fusarium wilt, Xanthomonas wilt and plant parasitic nematodes

were associated in terms of both prevalence and severity (Figure 5). In cassava, all pathogens

and pests except *B. tabaci* were associated in terms of prevalence. However, *B. tabaci* was

slightly associated with CMV in terms of infestation, which is expected as *B. tabaci* vectors CMV

(Figure 5). In potato, viruses and *R. solanacearum* were associated in terms of both prevalence

and severity/infestation (Figure 5). In sweetpotato fields, *B. tabaci* and sweetpotato aphids were

associated having similar infestation levels across fields.

FIGURE 5 | Relationships among economically important pathogens and pests and altitude in four major crops in Burundi and Rwanda, using non-metric multidimensional scaling (NMDS) based on Bray's distance dissimilarities of disease severity or infestation intensity. Each dot represents a surveyed crop field. A line (or vector) indicates whether (line direction) and how strongly (line length) the pathogen or pest is associated with others. Nearly parallel lines in the same quadrant indicate pathogens and pests often co-occur in a crop. (Cassava data was collected only in Burundi for one season). Abbreviations: *Acraea acerate* (AA), *Alternaria bataticola* (AB), Banana bunchy top virus (BBTV), *Bemisia tabaci* (cassava, potato or sweetpotato whitefly; BT), cassava bacterial blight (CBB), Cassava brown streak virus (CBSV), cassava green mite (CGM), cassava mealybug (CM), Cassava mosaic virus (CMV), *Cylas brunneus* (CB), *Cylas puncticollis* (CP), *Cosmopolites sordidus* (CS), Fusarium wilt of banana (FW), leaf miner fly (LMF), *Myzus persicae* (MP), *Phytophthora infestans* (PI), plant parasitic nematodes (PPN), *Pentalonia nigronervosa* (PN), *Phthorimaea operculella* (PO), potato viruses (PV), *Ralstonia solanacearum* (RS), *Synanthedon* sp. (clearwing moth; Ssp), sweetpotato mealybug (SPM), sweetpotato virus (SPV), sweetpotato weevils ( SPW ), and Xanthomonas wilt of banana (XW).

Analyses of associations between pairs of pathogens/pests provide a more detailed perspective

on which syndromes are important. In banana, some pairs like *C. sordidus* and *P. nigronervosa*,

plant parasitic nematodes and Fusarium wilt, plant parasitic nematodes and *P. nigronervosa*, *P.*

*nigronervosa* and Xanthomonas wilt, and Fusarium wilt and Xanthomonas wilt were positively

correlated ($p < 0.05$) in terms of both prevalence and severity/infestation, while other pairs were

only positively correlated ($p < 0.05$) in terms of prevalence: *C. sordidus* and plant parasitic

nematodes, plant parasitic nematodes and Xanthomonas wilt, plant parasitic nematodes and

BBTV, *P. nigronervosa* and BBTV, and BBTV and Xanthomonas wilt (Figure 6).

In cassava, CBSV and mealybug were positively correlated ($r = 0.69$, $p < 0.001$ in terms of

prevalence but negatively correlated ($r = -0.44$, $p < 0.01$) in terms of severity/infestation (Figure

6). Only *B. tabaci* and CMV were positively correlated in terms of severity/infestation ($r = 0.64$, p

$< 0.001$) while other pathogens and pests were positively correlated in terms of prevalence ($p <$

$0.05$). In potato, viruses and *R. solanacearum* was positively correlated in terms of prevalence (r

$= 0.33$, $p < 0.001$) and severity ($r = 0.21$, $p < 0.01$), while *P. infestans* and *M. persicae*, and leaf

miner fly and *M. persicae* were negatively correlated ($p < 0.05$) in terms of prevalence and

severity/infestation, and leaf miner fly and *R. solanacearum* were positively correlated ($p < 0.05$)

in terms of prevalence but negatively correlated ($p < 0.05$) in terms of severity/infestation

(Figure 6). In sweetpotato, *C. brunneus* and *C. puncticollis* were positively correlated in terms of

prevalence ($r = 0.58$, $p < 0.001$)  and infestation ($r = 0.18$, $p < 0.01$), while *A. bataticola* and

weevils (*Cylas* spp.) were negatively correlated in terms of prevalence ($r = -0.16$, $p < 0.05$) but

positively correlated in terms of severity/infestation ($r = 0.18$, $p < 0.05$) (Figure 6). *A. bataticola*

and mealybug, *A. bataticola* and aphids, *A. bataticola* and *B. tabaci*, *A. bataticola* and *Acraea*

*acerate*, and aphids and *Synanthedon* sp. were negatively correlated in terms of prevalence

FIGURE 6 | Association of pathogens and pests in banana, cassava, potato, and sweetpotato in smallholder farms in Burundi and Rwanda, based on Pearson's correlation of prevalence. (Cassava data was collected only in Burundi for one season).

Abbreviations: *Acraea acerate* (AA), *Alternaria bataticola* (AB), Banana bunchy top virus (BBTV), *Bemisia tabaci* (cassava, potato or sweetpotato whitefly; BT), cassava bacterial blight (CBB), Cassava brown streak virus (CBSV), cassava green mite (CGM), cassava mealybug (CM), Cassava mosaic virus (CMV), *Cylas brunneus* (CB), *Cylas puncticollis* (CP), *Cosmopolites sordidus* (CS), Fusarium wilt of banana (FW), leaf miner fly (LMF), *Myzus persicae* (MP), *Phytophthora infestans* (PI), plant parasitic nematodes (PPN), *Pentalonia nigronervosa* (PN), *Phthorimaea operculella* (PO), potato viruses (PV), *Ralstonia solanacearum* (RS), *Synanthedon* sp. (clearwing moth; Ssp), sweetpotato mealybug (SPM), sweetpotato virus (SPV), sweetpotato weevil ( SPW ), and Xanthomonas wilt of banana (XW).

## 3.3 | Prediction of pathogen and pest damage and geographic range shifts

### 3.3.1 | Machine learning to predict pathogen and pest levels

We compared machine learning algorithms in terms of their ability to predict severity and infestation in the field samples at monthly, quarterly, and annual intervals (Table 1). SVM and RF had the highest model performance for some pathogens/pests. For the SVM model, the performance characteristics across all pathogens and pests were 3-16% MAE for training, 7-20% and 4-26% RMSE, and 0.20-0.72 and 0.09-0.72 R-squared for training and testing, respectively, compared to the RF model with 3-17% MAE for training, 8-21% and 5-23% RMSE, and 0.23-0.72 and 0.12-0.73 R-squared for training and testing, respectively (Table 1). Among the 27 pathogens and pests tested, the three with the highest predictive performance metrics per crop were reported in Table 1.

The predictive performance was good for certain pathogens and pests, but poor for others (Table 1). As an example, SVM and RF performed well in predicting the monthly, quarterly, and annual infestation intensity of leaf miner fly in this region, explaining 69-71% of the variability in the data (R-squared) with low errors. The RMSE was 17-18% during training and 16-17% during prediction, demonstrating that both models generalized well to unseen data. In addition, the MAE of 11-13% indicated a low average error in training. The tight alignment between training and prediction metrics showed the robustness, stability and reliability of these models to predict leaf miner fly infestation intensity.

Pathogens and pests differed in the importance of predictor variables. Contemporary weather

variables such as monthly temperature, precipitation, and relative humidity (extracted through

NASAPOWER), generally had high contribution scores. Cropland structure variables also played

significant roles, with contribution scores reaching 100 for cropland density, 81 for cropland

connectivity, and 69 for altitude. For example, the key variables for predicting leaf miner fly

infestation intensity (monthly) using SVM were monthly temperature, precipitation, and relative

humidity, precipitation in the driest month, cropland connectivity, cropland density and annual

temperature range (i.e. difference between the maximum temperature of warmest month and

the minimum temperature of coldest month).

**TABLE 1 |** Support vector machine (SVM) and random forest (RF) performance for predicting
pathogen and pest levels in smallholder farms in Burundi and Rwanda, for selected species with
higher model performance. The response variable was severity for the pathogens and infestation
intensity for pests except for infestation rate for sweetpotato aphids.

| | | | SVM | | | | | Random Forest | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | Training | | | Prediction | | Training | | | Prediction | |
| Crop | Pathogen/pest | Timeframe | %RMSE | R-squared | %MAE | %RMSE | R-squared | %RMSE | R-squared | %MAE | %RMSE | R-squared |
| Banana | *P. nigronervosa* | Monthly | 18 | 0.63 | 13 | 21 | 0.56 | 16 | 0.72 | 12 | 18 | 0.66 |
| | | Quarterly | 18 | 0.63 | 13 | 21 | 0.56 | 16 | 0.72 | 12 | 19 | 0.64 |
| | | Annual | 19 | 0.63 | 14 | 21 | 0.55 | 16 | 0.71 | 11 | 19 | 0.63 |
| | Banana bunchy top virus | Monthly | 7 | 0.22 | 3 | 4 | 0.10 | 8 | 0.25 | 4 | 5 | 0.12 |
| | | Quarterly | 7 | 0.22 | 3 | 4 | 0.11 | 8 | 0.23 | 4 | 5 | 0.13 |
| | | Annual | 7 | 0.20 | 3 | 4 | 0.09 | 8 | 0.24 | 4 | 5 | 0.13 |
| | Plant parasitic nematodes | Monthly | 11 | 0.42 | 6 | 15 | 0.29 | 10 | 0.42 | 6 | 14 | 0.31 |
| | | Quarterly | 11 | 0.40 | 6 | 15 | 0.25 | 11 | 0.37 | 7 | 14 | 0.32 |
| | | Annual | 11 | 0.39 | 6 | 15 | 0.23 | 11 | 0.37 | 7 | 14 | 0.29 |
| Potato | Leaf miner fly | Monthly | 18 | 0.69 | 13 | 16 | 0.71 | 18 | 0.70 | 12 | 16 | 0.71 |
| | | Quarterly | 18 | 0.69 | 13 | 16 | 0.71 | 18 | 0.71 | 11 | 16 | 0.69 |
| | | Annual | 18 | 0.70 | 13 | 16 | 0.72 | 17 | 0.72 | 11 | 17 | 0.69 |
| | *Ralstonia solanacearum* | Monthly | 14 | 0.30 | 10 | 14 | 0.13 | 13 | 0.40 | 10 | 15 | 0.12 |
| | | Quarterly | 15 | 0.28 | 10 | 14 | 0.13 | 13 | 0.41 | 10 | 15 | 0.12 |
| | | Annual | 15 | 0.28 | 11 | 14 | 0.16 | 12 | 0.39 | 10 | 15 | 0.13 |
| | Potato viruses | Monthly | 20 | 0.31 | 16 | 21 | 0.34 | 21 | 0.28 | 17 | 21 | 0.38 |
| | | Quarterly | 20 | 0.32 | 16 | 21 | 0.33 | 20 | 0.31 | 16 | 21 | 0.38 |
| | | Annual | 20 | 0.31 | 16 | 21 | 0.34 | 21 | 0.30 | 17 | 21 | 0.37 |
| Sweet | *Alternaria* | Monthly | 15 | 0.35 | 9 | 10 | 0.42 | 14 | 0.39 | 9 | 10 | 0.39 |

| | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| potato | *bataticola* | Quarterly | 14 | 0.41 | 9 | 9 | 0.43 | 13 | 0.45 | 8 | 9 | 0.42 |
| | | Annual | 15 | 0.35 | 8 | 9 | 0.47 | 14 | 0.38 | 9 | 9 | 0.43 |
| | Sweetpotato weevils (*Cylas* spp.) | Monthly | 19 | 0.44 | 10 | 25 | 0.26 | 18 | 0.38 | 10 | 22 | 0.44 |
| | | Quarterly | 19 | 0.42 | 10 | 26 | 0.21 | 19 | 0.38 | 11 | 22 | 0.41 |
| | | Annual | 20 | 0.41 | 10 | 25 | 0.27 | 19 | 0.36 | 11 | 23 | 0.40 |
| | Sweetpotato aphids | Monthly | 14 | 0.60 | 6 | 9 | 0.60 | 10 | 0.53 | 5 | 7 | 0.73 |
| | | Quarterly | 14 | 0.60 | 6 | 9 | 0.58 | 10 | 0.58 | 4 | 8 | 0.65 |
| | | Annual | 14 | 0.58 | 6 | 9 | 0.58 | 9 | 0.54 | 4 | 9 | 0.53 |

### 3.3.2 | Pathogen and pest communities across altitudinal gradients

For each crop, pathogen and pest communities differed at low versus high altitudes (Table 2).

Although there was not strong evidence for differences across the altitudinal gradients for 5 out

of 27 pathogens and pests (correlations with p > 0.05 in Table 1), likely due in some cases to high

variability across samples, there was evidence for differences in some pathogens and pests that

distinguished the pathogen/pest communities (correlations in Table 2 with lower p-values). For

instance, crop production at higher altitudes (>1,500 m.a.s.l) had more *Myzus persicae*,

*Phthorimaea operculella*, and cassava green mites. At lower altitudes, crop fields had more

Fusarium wilt and Xanthomonas wilt of bananas, *P. infestans*, *A. acerate* and *Bemisia tabaci*

(Table 2). The distribution of these pathogens and pests across altitudes was supported in the

analysis combining observations from Burundi and Rwanda; overall, for banana, potato, and

sweetpotato, there was evidence that 44% of the targeted pathogens and pests were more

common at low altitudes, and that 17% were more common at higher altitudes.

When analyzing data for Rwanda and Burundi separately, differences across altitudes for

pathogens and pests were specific to the measure being used (e.g., disease intensity; Table 2) or

the country. For example, potato fields at high altitudes had a higher prevalence of potato leaf

miner fly, while there was not evidence for a difference in the infestation intensity of this pest

across the altitudinal gradient. Similarly, there was stronger evidence that altitude was

negatively correlated with the prevalence and severity of banana Fusarium wilt in Burundi (p <

0.001) compared to Rwanda (p = 0.069 for severity and p = 0.85 for prevalence). At the

community level, potato fields in Rwanda and cassava fields in Burundi had higher pathogen and

pest richness at higher altitudes, while richer pathogen and pest communities were found in

banana fields at lower altitudes.

TABLE 2 | Correlation between altitude and pathogen and pest prevalence, severity, and
infestation in Burundi and Rwanda (PRV: prevalence, SEV: severity, Rate: infestation rate, INF:
infestation intensity, NUM: number of pests).
485

| Crop | Disease/Pathogen/Pest | Variables | Correlation in Burundi | | Correlation in Rwanda | | Correlation in both countries combined | |
|---|---|---|---|---|---|---|---|---|
| | | | r | P-value | r | P-value | r | P-value |
| Banana | Xanthomonas wilt | PRV | -0.08 | 0.28 | -0.07 | 0.49 | -0.13 | 0.022 |
| | | SEV | -0.02 | 0.76 | -0.04 | 0.7 | -0.13 | 0.028 |
| | Fusarium wilt | PRV | -0.37 | <0.001 | -0.02 | 0.85 | -0.27 | <0.001 |
| | | SEV | -0.33 | <0.001 | 0.18 | 0.069 | -0.34 | <0.001 |
| | Banana bunchy top virus | PRV | -0.13 | 0.065 | 0 | - | -0.49 | <0.001 |
| | | SEV | -0.01 | 0.9 | 0 | - | -0.22 | <0.001 |
| | Plant parasitic nematodes | PRV | -0.01 | 0.90 | 0.05 | 0.65 | -0.11 | 0.065 |
| | | INF | -0.13 | 0.07 | -0.03 | 0.78 | -0.02 | 0.72 |
| | Banana weevil (Cosmopolites sordidus) | PRV | 0.03 | 0.700 | -0.14 | 0.16 | 0.18 | 0.003 |
| | | INF | -0.07 | 0.37 | -0.38 | <0.001 | -0.13 | 0.034 |
| | Banana aphid (Pentalonia nigronervosa) | PRV | 0.03 | 0.68 | -0.08 | 0.43 | -0.14 | 0.018 |
| | | INF | 0.09 | 0.24 | 0.09 | 0.37 | -0.25 | <0.001 |
| | | Richness | -0.16 | 0.028 | 0.80 | 0.41 | -0.30 | <0.001 |
| Potato | Leafminer fly | PRV | 0.16 | 0.16 | 0.26 | 0.006 | 0.09 | 0.22 |
| | | INF | 0.17 | 0.13 | 0.01 | 0.93 | 0.04 | 0.62 |
| | | Rate | 0.14 | 0.21 | 0.13 | 0.16 | 0.01 | 0.88 |
| | Potato aphid (Myzus persicae) | PRV | 0.02 | 0.86 | 0.24 | 0.011 | 0.23 | 0.001 |
| | | Rate | 0.07 | 0.54 | 0.25 | 0.008 | 0.21 | 0.004 |
| | Potato whitefly (Bemisia tabaci) | PRV | -0.57 | <0.001 | 0.04 | 0.7 | -0.17 | 0.02 |
| | | NUM | -0.38 | <0.001 | 0.08 | 0.39 | 0.05 | 0.45 |
| | Potato tuber moth (Phthorimaea operculella) | PRV | 0.34 | 0.003 | 0.03 | 0.75 | 0.16 | 0.028 |
| | | Rate | 0.36 | 0.001 | -0.05 | 0.58 | 0.15 | 0.03 |
| | Ralstonia solanacearum | PRV | -0.09 | 0.41 | 0.10 | 0.31 | 0.04 | 0.61 |
| | | SEV | 0.04 | 0.73 | -0.08 | 0.41 | -0.03 | 0.66 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Potato viruses | PRV | 0 | - | 0.10 | 0.3 | 0.10 | 0.16 |
| | | SEV | 0.04 | 0.71 | **-0.21** | **0.025** | -0.61 | 0.4 |
| | *Phytophthora* | PRV | -0.20 | 0.082 | 0.07 | 0.46 | **-0.18** | **0.014** |
| | *infestans* | SEV | -0.21 | 0.068 | -0.06 | 0.53 | -0.14 | **0.044** |
| | | **Richness** | -0.15 | 0.18 | **0.26** | **0.005** | 0.1 | 0.18 |
| Sweet Potato | *Alternaria bataticola* | **PRV** | **0.58** | **<0.001** | 0.05 | 0.61 | **0.54** | **<0.001** |
| | | **SEV** | **0.48** | **<0.001** | 0.04 | 0.65 | **0.32** | **<0.001** |
| | Sweetpotato viruses | PRV | 0.01 | 0.98 | 0 | - | 0.03 | 0.63 |
| | | SEV | 0.06 | 0.51 | -0.15 | 0.14 | -0.08 | 0.24 |
| | Sweetpotato whitefly (*Bemisia tabaci*) | PRV | -0.11 | 0.26 | 0 | - | **-0.22** | **0.001** |
| | | NUM | -0.05 | 0.61 | 0 | - | -0.11 | 0.13 |
| | Sweetpotato butterfly *(A. acerate)* | **PRV** | **-0.19** | **0.045** | 0 | - | **-0.32** | **<0.001** |
| | | Rate | -0.10 | 0.29 | 0 | - | **-0.19** | **0.005** |
| | | **NUM** | **-0.19** | **0.045** | 0 | - | **-0.22** | **0.001** |
| | Sweetpotato weevils (*Cylas* spp.) | PRV | -0.15 | 0.13 | 0.05 | 0.63 | **-0.17** | **0.016** |
| | | INF | 0.02 | 0.81 | 0.08 | 0.43 | **0.17** | **0.013** |
| | | Rate | -0.03 | 0.76 | -0.05 | 0.62 | **-0.17** | **0.013** |
| | *Cylas puncticollis* | PRV | -0.08 | 0.43 | 0.04 | 0.69 | -0.08 | 0.24 |
| | | **NUM** | **-0.23** | **0.016** | -0.06 | 0.54 | -0.04 | 0.61 |
| | *C. brunneus* | PRV | 0.08 | 0.39 | 0.06 | 0.58 | 0.02 | 0.8 |
| | | NUM | -0.01 | 0.99 | **-0.20** | **0.042** | -0.02 | 0.75 |
| | *Synanthedon* sp. | **PRV** | **-0.28** | **0.004** | **0.26** | **0.007** | 0.01 | 0.91 |
| | Sweetpotato mealybug | **PRV** | **-0.20** | **0.036** | 0 | - | **-0.21** | **0.002** |
| | Sweetpotato aphids | **PRV** | **0.24** | **0.013** | 0 | - | 0.01 | 0.94 |
| | | **Rate** | **0.45** | **<0.001** | 0 | - | **0.22** | **0.002** |
| | | **NUM** | **0.35** | **<0.001** | 0 | - | **0.17** | **0.01** |
| | | Richness | 0.05 | 0.6 | 0.10 | 0.34 | -0.02 | 0.8 |
| Cassava | Cassava whitefly (*Bemisia tabaci*) | **PRV** | **-0.80** | **<0.001** | - | - | - | - |
| | | **NUM** | **-0.32** | **0.032** | - | - | - | - |
| | Cassava green mite | **PRV** | **0.37** | **0.012** | - | - | - | - |
| | | INF | 0.06 | 0.68 | - | - | - | - |
| | Cassava mealybug | PRV | 0.29 | 0.057 | - | - | - | - |
| | | INF | 0.27 | 0.072 | - | - | - | - |
| | Cassava mosaic virus | **PRV** | **0.31** | **0.037** | - | - | - | - |
| | | **SEV** | **-0.52** | **<0.001** | - | - | - | - |
| | Cassava brown streak virus | PRV | 0.17 | 0.25 | - | - | - | - |
| | | **SEV** | **-0.47** | **0.002** | - | - | - | - |
| | Cassava bacterial blight | PRV | 0.12 | 0.43 | - | - | - | - |
| | | SEV | 0.25 | 0.093 | - | - | - | - |
| | | **Richness** | **0.78** | **<0.001** | - | - | - | - |

### 3.3.3 | Climate change and impact on regional temperature patterns

The frequency distribution of annual temperatures across altitudes in the Great Lakes region

illustrates the transformative impact of climate change (Figure 7). A progressive shift in

temperature distribution is expected, highlighting how currently cooler climates at higher

altitudes are likely to experience much higher temperatures (Figure 7). A general warming trend

is projected at both high and low altitudes. There is a greater temperature increase in the SSP5-

8.5 scenario than in the SSP1-2.6 scenario. Currently, the mean annual temperature in the region

ranges between 17-30°C at low altitudes, with the mode around 24°C, and between 10-22°C at

higher altitudes, with the mode around 17°C. As climate change progresses, and depending on

socio-economic pathways, this temperature landscape is likely to change substantially. Under

the SSP1-2.6 scenario, by 2090 (2081-2100), temperatures are expected to range between 20-

32°C at low altitudes, with the mode around 26°C, and between 10-25°C at high altitudes, with

mode 20°C. The high-emission scenario (SSP5-8.5) predicts that temperatures will range

between 22-36°C at low altitudes, with the mode around 28°C, and between 12-30°C at higher

altitudes, with the mode around 25°C.

These projections suggest future climate conditions at higher altitudes could become more

favorable for the 44% of pathogens and pests we evaluated that currently are more common at

lower altitudes (section 3.3.2). Likewise, pathogens and pests currently more common at higher

altitudes (17% of those evaluated in section 3.3.2) may become less common as the locations

warm in the future. These percentages are based on banana, potato, and sweetpotato, the three

crops for which we had the most complete data sets for analysis in predictive models based on

field data collected to allow direct comparisons across pathogens and pests.

FIGURE 7 Shift in annual temperature range at lower and higher elevations in the Great Lake region of Africa. Density is the probability density or likelihood of temperature rates occurring at locations in the Great Lakes region of Africa. TP: time period of the projections (while the 'Current' reference remains the same). SSP: Shared Socioeconomic Pathway, sustainability (SSP1-2.6) and fossil-fueled development (SSP5-8.5) emission scenarios. Here, high altitude refers to locations above 1500 m.a.s.l. and low altitude those below 1500 m.a.s.l.

## 511    4 | DISCUSSION

This study identified candidate surveillance locations based on cropland connectivity, a starting

point in understanding geographic spread and establishment of crop-specific pathogens and

pests in four key food security crops in the Great Lakes region. We also quantified the current

geographic and seasonal distribution of 27 economically important pathogens and pests across

climate gradients. We evaluated future climate scenarios, with evidence that 44% of the

pathogens and pests would become more common at higher altitudes while 17% would become

less common.

### 4.1 | Pathogen and pest risk across landscape structures

The cropland connectivity analysis provides a first approximation to the potential geographic

distribution of pathogen and pest risk. This analysis helped us identify candidate priority

locations for general surveillance and mitigation of the potential spread of crop-specific

pathogens and pests based on networks of host availability (Figure 2). Analysis of cropland

connectivity provides a baseline for quantifying invasion risks for pathogens and pests at a

national or global scale (Andersen Onofre et al., 2021, Etherton et al., 2025, Mouafo-Tchinda et

al., 2024). In the Great Lakes region, locations in Rwanda and Burundi had a high CCRI for all four

food-security crops, highlighting their vulnerability to pathogen and pest spread in the absence

of phytosanitary measures. Maps of cropland connectivity are particularly relevant for

monitoring outbreaks of introduced or emerging diseases and pests, as tools for national plant

protection organizations (NPPOs).

Ongoing improvements to understanding other components of pathogen and pest risk regionally

– such as the effects of climate change, local trade, and genetic resistance deployment – could

be integrated in more complete habitat connectivity analyses as additional resources become

available (Keshav et al., 2024). The disaster plant pathology framework emphasizes the

importance of developing and implementing disease management strategies before, during, and

after disasters (Etherton et al., 2024). Human-driven crises like the ongoing Boko-Haram and

separatist conflicts in Cameroon played a key role in human migration and informal movement

of cassava and sweetpotato planting material between neighboring villages (Mouafo-Tchinda et

al. (2024). The recurrence of crises in the Great Lakes region can increase pathogen and pest risk

nationally or internationally, especially in production regions with high cropland connectivity.

These threats motivate strengthening monitoring of informal seed movement in highly

connected cropland.

## 4.2 | Pathogen and pest communities

Our analysis of the on-farm distribution of 27 pathogens and pests highlights how these

communities change along climate gradients, in terms of seasonal abundance (Figure 4),

community structure (Figure 5), and interspecific pathogen/pest associations (Figure 6). Previous

studies have typically addressed how individual pathogens or pests are distributed along climate

gradients (Ebregt et al., 2005, Okonya & Kroschel, 2013, Nyang'au *et al.*, 2021), yet a community

perspective provides a more realistic scenario for understanding regional pest management

(Makiola *et al.*, 2021, Savary *et al.*, 2000, Were *et al.*, 2013). Our study addresses this knowledge

gap by identifying which pathogens/pests are most prevalent across seasons, which pathogen

and pest occurrences are associated, and how climate gradients are likely to influence disease

severity and pest infestation. Evaluating these interactions and cumulative effects of multiple

pathogens/pests in specific production environments can increase our understanding of injury

profiles (Savary et al., 2000), the cumulative damage to crops caused by the set of biotic

stressors including insects, pathogens and weeds.

Our findings highlight injury profiles for food security crops across climate gradients. Some

associations are already well-known; as expected, BBTV and its vector *P. nigronervosa* were

positively correlated (Blomme et al., 2020, Hu *et al.*, 1996). BBTV has been present in Africa for

decades, its spread continues to have an impact on new regions, and its management remains a

major challenge (Ngatat *et al.*, 2024, Rybicki, 2015, Ocimati *et al.*, 2024). We also found strong

associations between certain pathogens and pests; Fusarium wilt, Xanthomonas wilt and plant

parasitic nematodes were associated in banana, viruses and *R. solanacearum* in potato, and *B.*

*tabaci* and aphids in sweetpotato. These pathogen/pest associations underscore the need for

integrated management strategies that address the spread and combined impacts of multiple

pathogens and pests on crops (Alcalá Briseño *et al.*, 2023, Mouafo-Tchinda *et al.*, 2022).

**4.3 | Climate-based predictions of pathogen and pest levels**

In the Great Lakes region, temperatures decreased with increasing altitude (Figure 3), where

higher temperatures were associated with an increase in the prevalence and severity of some

pathogens and pests and a decrease in others (Table 2). Whiteflies were more common at low

altitudes for cassava, potato and sweetpotato. Aphids were more common at low altitudes for

banana, but more common at high altitudes for potato. The observed differences in the

altitudinal distribution of these pests could be attributed to their species composition, host

specificity, and the influence of species-specific ecological adaptations (McCulloch & Waters,

2023, Aleuy & Kutz, 2020). Aphid communities included different species, with *P. nigronervosa*

more frequent in low-altitude crops, and *M. persicae* more frequent in high-altitude crops (Aleuy

& Kutz, 2020, Wells & Clark, 2019, Malumphy *et al.*, 2017, Wieczorek *et al.*, 2019).

The distribution of host availability across a landscape can influence pathogen/pest prevalence,

severity/infestation, and richness (Etherton et al., 2025, Buddenhagen et al., 2022, Poveda et al.,

2012). In this study, the availability of banana, potato, and sweetpotato increased with altitude

while cassava decreased slightly (Figure 3), which may contribute to differences observed in the

pathogen and pest communities of these crop species. As a crop species becomes more

common, insect and microbial communities may become dominated by specialist species

infesting that crop (Poveda et al., 2012).

Weather variables are well known to be useful for predicting disease severity (Fenu & Malloci,

2021, Kaundal et al., 2006, Carisse & Fall, 2021). Cropland geography variables (cropland density,

connectivity), altitude, and climate variables (annual mean temperature, temperature annual

range, and annual precipitation) were key for most of the pathogens and pests we evaluated.

The performance of machine learning algorithms depends on data availability, the predictor

variables, and the biological features of pathogens and pests (Xiao *et al.*, 2018, Fenu & Malloci,

2021). The data for Rwanda and Burundi supported prediction of some pathogens, while others,

like BBTV, would require more study. The models effectively predicted leaf miner fly infestation

intensity, accounting for 69-71% of the variability with low training and predicting errors. These

models could serve as a baseline for predictive models of leaf miner fly infestation intensity in

this region (Table 1).

Temperature influences the life cycle, reproduction, and migration patterns of pathogens and

pests (Mouafo-Tchinda *et al.*, 2021, Yamamura & Kiritani, 1998, Awmack *et al.*, 1997). In

response to regional warming, certain pathogens and pests may become more common at

specific altitudes (Datta *et al.*, 2017, Poveda et al., 2012, Blomme et al., 2020), which will require

adaptation of pathogen and pest management to future conditions. Several studies suggest that

climate change shifts species ranges to higher altitudes (Colwell et al., 2008, Bebber *et al.*, 2013,

Bale *et al.*, 2002), and the new results from this study clarify the specific effects on individual

pathogens and pests in Rwanda and Burundi for key food security crops.

Pathogen/pest richness in banana decreased with altitude, while it increased in cassava. In

addition, the severity of certain pathogens/diseases such as BBTV, CMV, *P. infestans* and

Fusarium wilt decreased with altitude, while the severity or infestation of *A. bataticola* and

aphids in sweetpotato increased. In general, we found evidence that 44% of the pathogens and

pests currently more common at low altitude in banana, potato, and sweetpotato could shift to

higher altitudes under climate change, based on climate matching. Specifically, those expected

to become more common at higher altitudes are Xanthomonas wilt, Fusarium wilt, BBTV, and *P.*

*nigronervosa*, in banana; sweetpotato mealybug, *A. acerate*, sweetpotato weevils in

sweetpotato; and *B. tabaci* in potato and sweetpotato. The 17% currently more common at high

altitudes could become less common. Cropland connectivity was positively correlated with

altitude and was a key predictor of severity/infestation of some pathogens and pests. This

suggests that food-security crops at higher altitudes may experience substantial flows of

pathogens and pests, amplifying the risk of damage. Customizing risk assessments for these

specific pest/pathogen communities across altitudes could make management strategies better

adapted to the challenges posed by climate change for each crop in the region.

## 4.4 | Translating pathogen and pest risk to support current and future agriculture

Historically, global and national research and extension efforts have been employed to manage

diseases and pests in the Great Lakes region (RTB, 2013, Kroschel et al., 2014). Integrated

management systems combining cultural control, surveillance, awareness creation and involving

policy makers and farmers have been adopted (Gotor *et al.*, 2022, Legg *et al.*, 2006). There has

been support for active management programs where clean planting material is provided to

farmers to encourage them to destroy diseased plants. Efforts have been made to create

awareness and empower farmers to recognize and destroy diseased plants. A challenge for these

efforts is availability of resources to support ongoing, sustainable plans by governments and

national programs, especially after project lifespans are over. The results of this new study can

help to inform prioritization and efficient use of available resources. Currently, for many food

security crops, efforts are underway to improve seed health, obtain resistance to key pathogens

and pests, and adapt these improved technologies and techniques for use in humanitarian

settings (Andrade-Piedra *et al.*, 2020, Andrade-Piedra *et al.*, 2023, RTB, 2013).

Pathogen and pest risk analysis can help to ensure the economic, social, and environmental

sustainability of crop production (Carisse & Fall, 2021, Fall *et al.*, 2016, Kaundal et al., 2006).

Image analysis apps such as the PlantVillage Nuru app and the Tumaini app can help smallholder

farmers in the region diagnose crop damage caused by pests and diseases, and offer national

plant protection organizations and extension staff a tool for surveillance and disease mapping

(Ramcharan *et al.*, 2019, Kreuze *et al.*, 2022, Selvaraj *et al.*, 2019). Predicting the timing of

pathogen and pest threats would enable farmers to adjust management strategies in the short

run and would support prioritization in research and breeding programs in the long run.

Ongoing improvements in predicting pathogen and pest impacts depend on weather data

availability, which is often limited in low-income and disaster-prone countries. Satellite platforms

like NASA POWER offer a useful alternative (Sparks, 2018), providing near-real-time weather

data and allowing researchers in resource-limited settings to develop useful models without the

need for extensive ground observations. Another challenge is detection of transient pests such

as aphids, which move frequently and rapidly across fields (Loxdale, 2018). Addressing this

requires the use of innovative sampling techniques, such as suction traps and environmental

DNA (eDNA) analysis, to monitor pest populations more accurately (Poppinga *et al.*, 2016,

Taberlet *et al.*, 2012). However, implementing these advanced approaches in low-income and

disaster-prone areas such as the Great Lakes region remains a challenge due to limited resources

and technical capacity, requiring scalable and cost-effective solutions.

In disaster-prone regions, humanitarian organizations and government agencies must prioritize

key areas where abandonment or poor management of staple crops is particularly important to

regional spread of pathogens and pests (Etherton et al., 2024, Mouafo-Tchinda et al., 2024). This

study identifies candidate priority locations for mitigation of the establishment and spread of

pathogens and pests, characterizes pathogen and pest associations (injury profiles), and

evaluates potential altitudinal shifts under the influence of climate change. Future research can

build on these results by validating and refining the models for each pathogen/pest, and

developing tools that growers and policymakers can easily implement. Iterative improvement

can increase the adaptability and effectiveness of crop health management strategies, a no-

regrets adaptation strategy as agricultural systems face increasing challenges due to climate

change.

Acknowledgements

We appreciate support from the CGIAR Seed Equal Research Initiative, the CGIAR Roots Tubers

and Bananas Research Program, USAID Bureau of Humanitarian Assistance (BHA) award number

720BHA22IO00136, the OneCGIAR Initiative on Plant Health, and the CGIAR Trust Fund

([www.cgiar.org/funders/](www.cgiar.org/funders/)); we thank all donors and organizations which globally support the

work of CGIAR through their contributions to the CGIAR Trust Fund. We also appreciate support

from USDA Animal and Plant Health Inspection Service (APHIS) Cooperative Agreements

AP21PPQS&T00C195 and AP22PPQS&T00C133. The opinions expressed in this article are those

of the authors and do not necessarily reflect the view of the USAID BHA or USDA APHIS.

References

Adhikari, U., Nejadhashemi, A. P. and Woznicki, S. A. (2015) Climate change and eastern Africa:
a review of impact on major crops. *Food and Energy Security,* **4,** 110-132.
Alcalá Briseño, R. I., Batuman, O., Brawner, J., Cuellar, W. J., Delaquis, E., Etherton, B. A.*, et al.*
(2023) Translating virome analyses to support biosecurity, on-farm management, and
crop breeding. *Frontiers in Plant Science,* **14,** 1056603.
Aleuy, O. A. and Kutz, S. (2020) Adaptations, life-history traits and ecological mechanisms of
parasites to survive extremes and environmental unpredictability in the face of climate
change. *International Journal for Parasitology: Parasites and Wildlife,* **12,** 308-317.
Almekinders, C. J., Walsh, S., Jacobsen, K. S., Andrade-Piedra, J. L., McEwan, M. A., de Haan, S.*,*
*et al.* (2019) Why interventions in the seed systems of roots, tubers and bananas crops
do not reach their full potential. *Food Security,* **11,** 23-42.
Andersen Onofre, K. F., Buddenhagen, C. E., Rachkara, P., Gibson, R., Kalule, S., Phillips, D.*, et al.*
(2019) Modeling epidemics in seed systems and landscapes to guide management
strategies: The case of sweet potato in northern Uganda. *Phytopathology,* **109,** 1519-
1532.

Andersen Onofre, K. F., Forbes, G. A., Andrade-Piedra, J. L., Buddenhagen, C. E., Fulton, J. C., Gatto, M.*, et al.* (2021) An integrated seed health strategy and phytosanitary risk assessment: Potato in the Republic of Georgia. *Agricultural Systems,* **191,** 103144.

Andrade-Piedra, J., Almekinders, C. J., McEwan, M., Kilwinger, F., Mayanja, S., Mulugo, L.*, et al.* (2020) User guide to the toolbox for working with root, tuber and banana seed systems. Lima, Peru. CGIAR Research Program on Roots, Tubers and Bananas. *RTB User Guide*.

Andrade-Piedra, J., Dontsop, P., Dunia, D., Kulakow, P., Harahagazwe, D., Mouafo-Tchinda, R.*, et al.* (2023) Tools4SeedSystems: Working towards resilience through root, tuber and banana crops in humanitarian settings. Capacity needs assessment for root, tuber and banana seed interventions in humanitarian settings: Cameroon and DRC. Technical Report. 65.

Awmack, C., Harrington, R. and Leather, S. (1997) Host plant effects on the performance of the aphid *Aulacorthum solani* (Kalt.)(Homoptera: Aphididae) at ambient and elevated CO2. *Global Change Biology,* **3,** 545-549.

Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, T. M., Brown, V. K.*, et al.* (2002) Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology,* **8,** 1-16.

Bebber, D. P., Ramotowski, M. A. T. and Gurr, S. J. (2013) Crop pests and pathogens move polewards in a warming world. *Nature Climate Change,* **3,** 985-988.

Blomme, G., Ocimati, W., Amato, S., Felde, A. z., Kamira, M., Bumba, M.*, et al.* (2020) Banana pest risk assessment along banana trade axes running from low to high altitude sites, in the Eastern DR Congo and in Burundi. *African Journal of Agricultural Research,* **16(9),** 1253-1269.

Buddenhagen, C. E., Xing, Y., Andrade-Piedra, J. L., Forbes, G. A., Kromann, P., Navarrete, I.*, et al.* (2022) Where to invest project efforts for greater benefit: A framework for management performance mapping with examples for potato seed health. *Phytopathology,* **112,** 1431-1443.

Carisse, O. and Fall, M. L. (2021) Decision trees to forecast risks of strawberry powdery mildew caused by *Podosphaera aphanis*. *Agriculture,* **11,** 29.

Chevalier, M., Broennimann, O. and Guisan, A. (2024) Climate change may reveal currently unavailable parts of species' ecological niches. *Nature Ecology & Evolution,* **8,** 1298–1310.

Cilas, C., Goebel, F.-R., Babin, R. and Avelino, J. (2016) Tropical crop pests and diseases in a climate change setting—a few examples. In: *Climate change and agriculture worldwide.* (Torquebiau, E., ed.). Springer, Dordrecht, pp. 73-82.

Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C. and Longino, J. T. (2008) Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science,* **322,** 258-261.

Datta, A., Kühn, I., Ahmad, M., Michalski, S. and Auge, H. (2017) Processes affecting altitudinal distribution of invasive *Ageratina adenophora* in western Himalaya: The role of local adaptation and the importance of different life-cycle stages. *PloS One,* **12,** e0187708.

Ebregt, E., Struik, P., Odongo, B. and Abidin, P. (2005) Pest damage in sweet potato, groundnut and maize in north-eastern Uganda with special reference to damage by millipedes (Diplopoda). *NJAS-Wageningen Journal of Life Sciences,* **53,** 49-69.

Etherton, B., Plex Sula, A., Mouafo-Tchinda, R., Kakuhenzire, R., Kassaye, H., Asfaw, F*., et al.*
(2025) Translating Ethiopian potato seed networks: Identifying strategic intervention
points for managing bacterial wilt and other diseases. *Agr Syst,* **222,** 104167.
Etherton, B. A., Choudhury, R., Alcalá-Briseño, R., Xing, Y., Sulá, A. P., Carrillo, D*., et al.* (2023)
Are avocados toast? A framework to analyze decision-making for emerging epidemics,
applied to laurel wilt. *Agr Syst,* **206,** 103615.
Etherton, B. A., Choudhury, R. A., Alcalá Briseño, R. I., Mouafo-Tchinda, R. A., Plex Sulá, A. I.,
Choudhury, M*., et al.* (2024) Disaster plant pathology: Smart solutions for threats to
global plant health from natural and human-driven disasters. *Phytopathology,* **114,** 855-
868.
Fall, M. L., Van der Heyden, H. and Carisse, O. (2016) A quantitative dynamic simulation of
Bremia lactucae airborne conidia concentration above a lettuce canopy. *PLoS One,* **11,**
e0144573.
Fenu, G. and Malloci, F. M. (2021) Forecasting plant and crop disease: an explorative study on
current algorithms. *Big Data and Cognitive Computing,* **5,** 2.
Ferdu, A. and Solbreck, C. (2010) Oviposition preference and larval performance of the sweet
potato butterfly *Acraea acerata* on *Ipomoea* species in Ethiopia. *Agricultural and Forest*
*Entomology,* **12,** 161-168.
Fick, S. E. and Hijmans, R. J. (2017) WorldClim 2: new 1‑km spatial resolution climate surfaces
for global land areas. *International Journal of Climatology,* **37,** 4302-4315.
Gallego‑Tévar, B., Gil‑Martínez, M., Perea, A., Pérez‑Ramos, I. M. and Gómez‑Aparicio, L.
(2024) Interactive Effects of Climate Change and Pathogens on Plant Performance: A
Global Meta‑Analysis. *Global Change Biology,* **30,** e17535.
Garrett, K. A., Bebber, D., Etherton, B. A., Gold, K., Plex Sulá, A. I. and Selvaraj, M. G. (2022a)
Climate change effects on pathogen emergence: artificial intelligence to translate big
data for mitigation. *Annual Review of Phytopathology,* **60,** 357-378
Garrett, K. A. (2021) Impact network analysis and the INA R package: Decision support for
regional management interventions. *Methods in Ecology and Evolution,* **12,** 1634-1647.
Garrett, K. A., Thomas-Sharma, S., Forbes, G. A., Hernandez Nopsa, J. F. and Plex Sulá, A. I.
(2022b) Climate change and plant pathogen invasions. In: *Invasive Species and Global*
*Climate Change.* (Ziska, L., ed.). CABI, pp. 22-49.
Gotor, E., Di Cori, V., Pagnani, T., Kikulwe, E., Kozicka, M. and Caracciolo, F. (2022) Public and
private investments for banana Xanthomonas Wilt control in Uganda: the economic
feasibility for smallholder farmers. *African Journal of Science, Technology, Innovation*
*and Development,* **14,** 135-146.
Hausfather, Z. (2019) CMIP6: the next generation of climate models explained. *Carbon Brief*.
Hijmans, R., Barbosa, M., Ghosh, A. and Mandel, A. (2023) geodata: Download geographic data.
R package version 0.5.
Hijmans, R. J. (2024) terra: Spatial Data Analysis. R package version 1.8-0.
Hijmans, R. J., Bivand, R., Forner, K., Ooms, J., Pebesma, E. and Sumner, M. D. (2022) Package
'terra'. *Maintainer: Vienna, Austria*.
Hodkinson, I. D. (2005) Terrestrial insects along elevation gradients: species and community
responses to altitude. *Biological reviews,* **80,** 489-513.

Hu, J. S., Wang, M., Sether, D., Xie, W. and Leonhardt, K. W. (1996) Use of polymerase chain
reaction (PCR) to study transmission of banana bunchy top virus by the banana aphid
(*Pentalonia nigronervosa*). *Annals of Applied Biology,* **128,** 55-64.
International Food Policy Research Institute, M. (2020) Spatially-disaggregated crop production
statistics data in Africa south of the Sahara for 2017. In: *Harvard Dataverse.*
Jongejans, E., Skarpaas, O., Ferrari, M. J., Long, E. S., Dauer, J. T., Schwarz, C. M*., et al.* (2015) A
unifying gravity framework for dispersal. *Theoretical Ecology,* **8,** 207-223.
Juroszek, P., Bartsch, L., Fontaine, J. F., Racca, P. and Kleinhenz, B. (2022) Summary of the
worldwide available crop disease risk simulation studies that were driven by climate
change scenarios and published during the past 20 years. *Plant Pathology,* **71,** 1815–
1838.
Kaundal, R., Kapoor, A. S. and Raghava, G. P. (2006) Machine learning techniques in disease
forecasting: a case study on rice blast prediction. *BMC bioinformatics,* **7,** 1-16.
Keshav, K., Plex-Sula, A. and Garrett, K. A. (2024) geohabnet: Geographical Risk Analysis Based
on Habitat Connectivity. R Package, Version 2.0.0. . CRAN.
Kreuze, J., Adewopo, J., Selvaraj, M., Mwanzia, L., Kumar, P. L., Cuellar, W. J*., et al.* (2022)
Innovative digital technologies to monitor and control pest and disease threats in root,
tuber, and banana (RT&B) cropping systems: Progress and prospects. In: *Root, Tuber*
*and Banana Food System Innovations: Value Creation for Inclusive Outcomes.* pp. 261-
288.
Kroschel, J., Okonyaet, J., Juarez, H., Forbes, G., Kreuze, J., Beed, F*., et al.* (2014) Management
of critical pests and diseases in RTB crops under changing climates, through risk
assessment, surveillance and modeling. In: *RTB Workshop Reports; CGIAR Research*
*Program on Roots, Tubers and Bananas (RTB): Lima, Peru.* pp. 20.
Kruskal, J. B. (1964) Nonmetric multidimensional scaling: a numerical method. *Psychometrika,*
**29,** 115-129.
Kuhn, M. (2008) Building predictive models in R using the caret package. *Journal of Statistical*
*Software,* **28,** 1-26.
Legg, J., Bouwmeester, H., Bigirimana, S., Tata-Hangy, W., Obiero, H., Gashaka, G*., et al.* (2009)
Cassava Disease Surveillance Surveys 2009. Mapping Report, Great Lakes Cassava
Initiative. pp. 46.
Legg, J. P., Owor, B., Sseruwagi, P. and Ndunguru, J. (2006) Cassava mosaic virus disease in East
and Central Africa: epidemiology and management of a regional pandemic. *Advances in*
*Virus Research,* **67,** 355-418.
Loxdale, H. D. (2018) Aspects, including pitfalls, of temporal sampling of flying insects, with
special reference to aphids. *Insects,* **9,** 153.
Makiola, A., Holdaway, R. J., Wood, J. R., Orwin, K. H., Glare, T. R. and Dickie, I. A. (2021)
Environmental and plant community drivers of plant pathogen composition and
richness. *New Phytologist,* **233,** 496-504.
Malumphy, C., Eyre, D. and Anderson, H. (2017) Tobacco, sweet potato or silver leaf whitefly:
*Bemisia tabaci*. In: *Department for Environment Food and Rural Affairs, Unites Kingdom.*
*Hal.* pp. 6.

Margosian, M. L., Garrett, K. A., Hutchinson, J. S. and With, K. A. (2009) Connectivity of the
American agricultural landscape: assessing the national risk of crop pest and disease
spread. *BioScience,* **59,** 141-151.
McCulloch, G. A. and Waters, J. M. (2023) Rapid adaptation in a fast‑changing world: Emerging
insights from insect genomics. *Global Change Biology,* **29,** 943-954.
McEwan, M. (2016) Sweetpotato seed systems in sub-saharan Africa: A literature review to
contribute to the preparation of conceptual frameworks to guide practical interventions
for root, tuber and banana seed systems. RTB Working Paper. Apartado 1558, Lima 12,
Peru: CGIAR Research Program on Roots, Tubers and Bananas (RTB), pp. 45.
Mishra, K., Subbanna, A., Rajashekara, H., Paschapur, A. U., Jeevan, B., Singh, A. K.*, et al.* (2024)
Perceptions on Disease and Pest Status of Major Cultivated Crops in Indian Himalayas
Under Changing Climate. In: *Adapting to Climate Change in Agriculture-Theories and*
*Practices: Approaches for Adapting to Climate Change in Agriculture in India.* pp. 121-
141.
Mouafo-Tchinda, R. A., Beaulieu, C., Fall, M. L. and Carisse, O. (2021) Effect of temperature on
aggressiveness of *Plasmopara viticola* f. sp. *aestivalis* and *P. viticola* f. sp. *riparia* from
eastern Canada. *Canadian Journal of Plant Pathology,* **43,** 73-87.
Mouafo-Tchinda, R. A., Etherton, B. A., Plex Sulá, A. I., Andrade-Piedra, J., Ogero, K., Omondi, B.
A.*, et al.* (2024) Pathogen and pest risks to vegetatively propagated crops in
humanitarian contexts: Towards a national plant health risk analysis for Cameroon and
Ethiopia. *bioRxiv***,** 2024.02.12.580019.
Mouafo-Tchinda, R. A., Fall, M. L., Beaulieu, C. and Carisse, O. (2022) Competition Between
*Plasmopara viticola* Clade *riparia* and Clade *aestivalis*: A Race to Lead Grape Downy
Mildew Epidemics. *Plant Disease,* **106,** 2866-2875.
Nakato, G. V., Okonya, J. S., Kantungeko, D., Ocimati, W., Mahuku, G., Legg, J. P.*, et al.* (2023)
Influence of altitude as a proxy for temperature on key Musa pests and diseases in
watershed areas of Burundi and Rwanda. *Heliyon,* **9**.
Ngatat, S., Hanna, R., Doumtsop Fotio, A. R., Lienou, J. A., Nanga Nanga, S., Fotso Kuate, A.*, et*
*al.* (2024) Distribution and diversity of emergent Banana bunchy top virus infecting
banana and plantain in Cameroon, Central Africa. *Journal of Phytopathology,* **172,**
e13279.
Nyang'au, D., Atandi, J., Cortada, L., Nchore, S., Mwangi, M. and Coyne, D. (2021) Diversity of
nematodes on banana (*Musa* spp.) in Kenya linked to altitude and with a focus on the
pathogenicity of *Pratylenchus goodeyi*. *Nematology,* **24,** 137-147.
Ochieng, J., Knerr, B., Owuor, G. and Ouma, E. (2015) Agricultural commercialization and
household food security: The case of smallholders in Great Lakes Region of Central
Africa. In: *International Association of Agricultural Economists.* Milan, Italy 212588, pp.
30.
Ochieng, J., Ouma, E. and Birachi, E. (2014) Gender participation and decision making in crop
management in Great Lakes Region of Central Africa. *Gender, Technology and*
*Development,* **18,** 341-362.
Ocimati, W., Ogwal, G., Tazuba, A. F., Kubiriba, J., Tugume, J., Erima, R.*, et al.* (2024) Mapping
the vulnerability of banana production landscapes in Uganda to banana bunchy top
disease. *Frontiers in Agronomy,* **6,** 1401478.

Ogot, B. A. and Niane, D. T. (1984) The Great Lakes region. *General history of Africa IV: Africa*
*from the twelfth to the sixteenth century*, 498-524.
Okonya, J. S., Gamarra, H., Nduwayezu, A., Bararyenya, A., Kroschel, J. and Kreuze, J. (2021)
Serological survey and metagenomic discovery of potato viruses in Rwanda and Burundi
reveals absence of PVY in Burundi and first report of TRV in potatoes in sub-Saharan
Africa. *Virus Research, 302,* 198487.
Okonya, J. S. and Kroschel, J. (2013) Incidence, abundance and damage by the sweet potato
butterfly (*Acraea acerata* Hew.) and the African sweet potato weevils (*Cylas* spp.) across
an altitude gradient in Kabale district, Uganda. *Int J AgriScience, 3,* 814-824.
Olivoto, T. and Lúcio, A. D. C. (2020) metan: An R package for multi‑environment trial analysis.
*Methods in Ecology and Evolution, 11,* 783-789.
Ong'ayo, A. O. (2018) Displacement and cross-border mobility in the Great Lakes region-re-
thinking underlying factors and implications for regional management of migration.
*Africa Insight, 48,* 62-85.
Pautasso, M., Döring, T. F., Garbelotto, M., Pellis, L. and Jeger, M. J. (2012) Impacts of climate
change on plant diseases—opinions and trends. *European Journal of Plant Pathology,*
*133,* 295-313.
Pinnegar, J. K., Hamon, K. G., Kreiss, C. M., Tabeau, A., Rybicki, S., Papathanasopoulou, E*., et al.*
(2021) Future socio-political scenarios for aquatic resources in Europe: A common
framework based on shared-socioeconomic-pathways (SSPs). *Frontiers in Marine*
*Science, 7,* 568219.
Poppinga, S., Weisskopf, C., Westermeier, A. S., Masselter, T. and Speck, T. (2016) Fastest
predators in the plant kingdom: functional morphology and biomechanics of suction
traps found in the largest genus of carnivorous plants. *AoB Plants, 8,* plv140.
Poveda, K., Martínez, E., Kersch‑Becker, M. F., Bonilla, M. A. and Tscharntke, T. (2012)
Landscape simplification and altitude affect biodiversity, herbivory and Andean potato
yield. *Journal of Applied Ecology, 49,* 513-522.
Pryke, A., Mostaghim, S. and Nazemi, A. (2007) Heatmap visualization of population based multi
objective algorithms. In: *Evolutionary Multi-Criterion Optimization: 4th International*
*Conference, EMO 2007, Matsushima, Japan, March 5-8, 2007. Proceedings 4.* Springer,
pp. 361-375.
R Core Team, R. (2024) R: A language and environment for statistical computing. R Foundation
for Statistical Computing.
Ramcharan, A., McCloskey, P., Baranowski, K., Mbilinyi, N., Mrisho, L., Ndalahwa, M*., et al.*
(2019) A mobile-based deep learning model for cassava disease diagnosis. *Frontiers in*
*Plant Science, 10,* 272.
RTB, C. (2013) Workshop on tools for managing seed degeneration in RTB crops (5-8 February
2013, Arusha, Tanzania). CRP-RTB, pp. 23.
Rybicki, E. P. (2015) A Top Ten list for economically important plant viruses. *Archives of*
*Virology, 160,* 17-20.
Savary, S., Willocquet, L., Elazegui, F. A., Teng, P. S., Van Du, P., Zhu, D*., et al.* (2000) Rice pest
constraints in tropical Asia: characterization of injury profiles in relation to production
situations. *Plant Disease, 84,* 341-356.

Science Group Project on Accelerated Varietal Improvement and Seed Systems in Africa (2024)
CGIAR project on Accelerated Varietal Improvement and Seed Systems in Africa: Annual
Technical Report 2023.

Selvaraj, M. G., Vergara, A., Ruiz, H., Safari, N., Elayabalan, S., Ocimati, W.*, et al.* (2019) AI-
powered banana diseases and pest detection. *Plant Methods,* **15,** 1-11.

Simpson, G. (2020) ggvegan; ggplot2-based plots for vegan package. version 0.1. 0. *In:*
[*https://github*](https://github). *com/gavinsimpson/ggvegan; consulted: May*.

Singh, B. K., Delgado-Baquerizo, M., Egidi, E., Guirado, E., Leach, J. E., Liu, H.*, et al.* (2023)
Climate change impacts on plant pathogens, food security and paths forward. *Nature*
*Reviews Microbiology,* **21,** 640–656.

Sparks, A. H. (2018) nasapower: a NASA POWER global meteorology, surface solar energy and
climatology data client for R. *Journal of Open Source Software,* **3,** 1035.

Sparks, A. H., Forbes, G. A., Hijmans, R. J. and Garrett, K. A. (2014) Climate change may have
limited effect on global risk of potato late blight. *Global Change Biology,* **20,** 3621-3631.

Sseruwagi, P., Sserubombwe, W., Legg, J. P., Ndunguru, J. and Thresh, J. (2004) Methods of
surveying the incidence and severity of cassava mosaic disease and whitefly vector
populations on cassava in Africa: a review. *Virus Research,* **100,** 129-142.

Taberlet, P., Coissac, E., Hajibabaei, M. and Rieseberg, L. H. (2012) Environmental DNA.
*Molecular Ecology,* **21**.

Thies, C. and Tscharntke, T. (1999) Landscape structure and biological control in
agroecosystems. *Science,* **285,** 893-895.

Wells, K. and Clark, N. J. (2019) Host specificity in variable environments. *Trends in Parasitology,*
**35,** 452-465.

Were, H., Kabira, J., Kinyua, Z., Olubayo, F., Karinga, J., Aura, J.*, et al.* (2013) Occurrence and
distribution of potato pests and diseases in Kenya. *Potato Research,* **56,** 325-342.

Wickham, H. (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.
ISBN 978-3-319-24277-4.

Wieczorek, K., Fulcher, T. K. and Chłond, D. (2019) The composition of the aphid fauna (Insecta,
Hemiptera) of the Royal Botanic Gardens, Kew. *Scientific Reports,* **9,** 10000.

WorldData.info (2024) Climate comparison between Rwanda and Burundi based on a long-term
analysis of the last 20 years.

Xiao, Q., Li, W., Chen, P. and Wang, B. (2018) Prediction of crop pests and diseases in cotton by
long short term memory network. In: *Intelligent Computing Theories and Application:*
*14th International Conference, ICIC 2018, Wuhan, China, August 15-18, 2018,*
*Proceedings, Part II 14.* Springer, pp. 11-16.

Xing, Y., Hernandez Nopsa, J. F., Andersen, K. F., Andrade-Piedra, J. L., Beed, F. D., Blomme, G.*,*
*et al.* (2020) Global cropland connectivity: A risk factor for invasion and saturation by
emerging pathogens and pests. *BioScience,* **70,** 744-758.

Yamamura, K. and Kiritani, K. (1998) A simple method to estimate the potential increase in the
number of generations under global warming in temperate zones. *Applied Entomology*
*and Zoology,* **33,** 289-298.