C12ORF57: a novel principal regulator of synaptic AMPA currents and excitatory neuronal homeostasis

Ruiji Jiang[1]**, Malek Chouchane[1], Ching Moey[2], Jens Bunt[2], Jiang Li[1], Fowzan Alkuraya[4], Eman Alobeid[5], Linda Richards[2,5,] Erik Ullian[6], Elliott Sherr[1]**

1. University of California San Francisco Department of Neurology, San Francisco, CA United States
2. The University of Queensland, Queensland Brain Institute, Brisbane, Australia
3. King Faisal Specialist Hospital and Research Center, Department of Genetics, Riyadh, Saudi Arabia
4. Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia
5. Washington University in Saint Louis, Department of Neuroscience, St Louis, MO United States
6. University of California San Francisco, Department of Ophthalmology, San Francisco, CA, United States

.

Abstract

Biallelic recurrent loss of function mutations in *C12ORF57*, a novel open reading frame, underlie Temtamy syndrome (TS)—a neurodevelopmental disorder characterized by dysgenesis of the corpus callosum, epilepsy, and severe intellectual disability. To study the heretofore unknown function of this gene, we used a knockout (KO) mouse model of its murine ortholog, *Grcc10*. *Grcc10* KO mice exhibit the characteristic phenotypic features seen in human TS patients,

including increased epileptiform activity. Consistent with this propensity for seizures, hippocampal neurons in these mice show significantly increased AMPA receptor expression levels and higher amplitude of miniature excitatory postsynaptic currents (mEPSCs). We also find that GRCC10/C12ORF57 modulates the activity of calcium/calmodulin dependent kinase 4 (CAMK4) and regulates the expression of CREB and ARC, which are involved in synaptic scaling of AMPA receptors. Through multiple lines of inquiry, we establish that C12ORF57/GRCC10 plays a central modulatory role in synaptic homeostasis and uncover a novel mechanism for neuronal excitatory tuning.

Keywords: Temtamy syndrome, C12ORF57, epilepsy, ASD, CAMK4, CREB, ARC, neuronal homeostasis, AMPA receptors

Introduction

C12ORF57 is a three exon gene encoding a highly conserved 126 amino acid, 13 kD protein that is abundantly expressed in human fetal and postnatal brain [1,2]. Recent linkage and sequencing analyses have identified a recurrent autosomal recessive mutation in the start codon (c.1A>G, p.1M>V) of *C12ORF57* as the cause of Temtamy syndrome (TS), an autism and intellectual disability (ID) disorder characterized by ocular coloboma, dysgenesis of the corpus callosum and medically intractable seizures [1–5]. This syndrome was initially reported in Middle Eastern populations but subsequently also noted in other countries [3,6–8]. The above mentioned mutation is observed throughout the Middle East and is the single most common cause of recessive ID and developmental delay in Saudi Arabia, accounting for greater than 1% of intellectual disability with genetic cause in these cohorts [2,3]. Despite the widespread nature of this mutation among affected populations and evolutionary conservation suggesting an important biological function, currently

there is little known about its role in normal neuronal development and homeostasis. This is because C12ORF57 lacks homology to any known vertebrate protein, making it difficult to predict its function (**Supplemental Figure 1**) [2]. To discover the function of this novel gene, we generated and analyzed the consequences of homozygous knockout model of its murine ortholog, *Grcc10,* with which it shares 99.2% amino acid identity (**Supplemental Figure 1**).

We show that *Grcc10* knockout (KO) mice recapitulate many of the phenotypes found in human TS patients, notably epilepsy and corpus callosum anomalies, making it a useful model in which to study the function of this gene. Using biochemical (including yeast-2-hybrid interactions) and electrophysiological assays, we examine the potential mechanisms by which C12ORF57/GRCC10 regulates synaptic excitability. We analyze its interaction with the neuronal homeostasis protein, calcium calmodulin dependent kinase IV (CAMK4), as well as its regulation of CREB, Fos, and ARC, which are implicated in the synaptic scaling of AMPA receptors [9]. Taken together, our data show for the first time, that C12ORF57 is a central regulator of synaptic homeostasis, and also reveal a novel mechanism of neuronal excitatory tuning.

Results

Grcc10 loss of function in mice recapitulates Temtamy syndrome phenotypes

When examined by RT-PCR, the *C12ORF57* transcript was not detected in primary fibroblasts from patients homozygous for the most common c.1 A>G variant , suggesting that their phenotype was consistent with a complete loss of function (LoF) (**Figure 1a**) [2,3]. To model this homozygous inactivation of *C12ORF57* in human disease*,* we generated a transgenic mouse line (*Grcc10<tm1.1(KOMP)Vlcg>*) carrying a beta galactosidase-containing cassette that constitutively disrupts the *Grcc10* gene (*Grcc10* KO). *Grcc10* +/+ (WT), +/- (Het) and -/- (KO)

littermates were born with the expected 1:2:1 Mendelian ratios from heterozygote pairs (X=.014,

dF=2, p=.99, Chi-Square, n=154) (**Figure 1b**). *Grcc10 -/-* pups had high postweaning mortality

compared to WT and Het littermates as early as postnatal day 3 (P3) (X=.014, df=2, p=.00013,

Chi-Square, n=91), with the majority of animals dying before P2 (**Figure 1b**). *Grcc10* KO mice

that survived until weaning age showed significantly diminished physical growth compared to

their WT and Het littermates (11.71±0.11g vs. 4.15±0.99g, respectively, p<.00001 t=33.07,

df=8.18, N=5, Welch's T-Test), suggesting that *Grcc10* is also crucial for health and survival

(**Figure 1c**).

Since our previous work associating *C12ORF57* with ID, ASD, and epilepsy, and its high

expression in human fetal brain tissue strongly suggests a crucial role in normal brain development,

we first investigated *Grcc10* expression in early mouse brain development using *in situ*

hybridization on late embryonic days (E)15 and 18 (E15, E18) and P0 and P10 .[1,3] We observed

that *Grcc10* is broadly expressed in many brain regions at all ages examined. At E15, it is

specifically localized to the cortical plate, glial wedge, and the dorsal ventricular zone. At E18, we

observed higher expression in the cortex, hippocampus, and thalamus that was maintained until

P10 (**Figure 1d**). Interrogation of Allen Brain Atlas cell and region-specific RNA-seq also shows

global expression across multiple brain areas and neuronal cell types (**Supplemental Figure 2**).

The widespread expression of *Grcc10* in the developing cortex suggests that *Grcc10* may play an

early and central role in cortical development. Since previous studies showed that disrupted

cortical lamination is linked to ASD, ID and epilepsy we examined whether this was the case in

Grcc10 KO mice by performing immunohistochemistry on P0 murine cortical slices using

previously validated upper and lower-layer markers *Satb2* and *Ctip2* respectively[10–12]. We

observed no difference in the number of Satb2 WT (1403.00±89.09 cells/HPF) versus KO brains

(1498.20$\pm$40.14, p=.36, t=.95, df=10, Welch's tailed T-test) or Ctip2 neurons in WT

(610.43$\pm$31.00) versus KO brains (633.17$\pm$41.03, p=0.76, t=0.32, df=11, Welch's T-test) or the

pattern of lamination of WT and *Grcc10* KO cortices, suggesting that *Grcc10* loss of function does

not impair gross cortical architecture or layering (**Supplemental Figure 3**).

Our *Grcc10* -/- mice share a shortened corpus callosum structure similar to what is found in

patients with Temtamy syndrome. In P0 KO mice, corpus callosum length was shorter

(792.81$\pm$39.15 μm, p=.017, F=6.41, df=2. N=6, ANOVA) when compared to their WT

(984.53$\pm$12.23μm, p=.023, df=9, Tukey post-hoc test), and heterozygous littermates

(931.89$\pm$47.73μm) thought not statistically significantly so (p=.092, q=3.391, DF=9, Tukey post-

hoc test). The corpus callosum cross sectional area in these P0 KO mice was also reduced

(154859.22$\pm$8441.27μm^2) (p=.0058 ,ANOVA, F=9.65, df=2 N=6) compared to WT

(199602.27$\pm$8702.73μm^2) (p=.012,q=3.546, df=9, N=3, Tukey post-hoc test) and Het mice

(200644.93$\pm$6764.12μm^2, p=.010 q=3.628, DF=2, N=3, Tukey post-hoc test). Rostral to caudal

cortical length between KO (1625.01$\pm$30.49μm) and WT (1706.25 $\pm$56.16 μm,) or Het

(1709.39$\pm$63.84 μm p=.29, ANOVA, F=1.337, df=2, N=6) was not significantly different (**Figure**

1e).

Patients harboring *C12ORF*57 homozygous or biallelic mutant variants have a high rate of epilepsy

that is refractory in many cases [3,4]. We have observed *Grcc10* KO mice displaying spontaneous

seizure activity (**Supplemental Video 1**), but their early mortality and small size prevented us

from conducting EEG monitoring. To assess seizure susceptibility, we tested whether these mice

are more susceptible to pharmacologically induced seizures. We performed intra-peritoneal kainic

acid injection in P21 *Grcc10* KO, Het and WT mice, and scored their behaviors indicative of

seizures based on the Racine scale of 1-5 (**Figure 1f**) [13–16]. We observed that all animals presented with seizures following kainic acid, but not carrier injections. *Grcc10* KO mice reached clinically severe stages (=>5) significantly faster (9.44±1.33 min,p<.0001, 2-way ANVOA F=46.34, F=764.66 N=4) than *Grcc10* WT (36.33±2.84 min, p<.0001, N=6, Tukey post-hoc test) or Het littermates (20.74±1.82 min, N=5 p<.0001, Tukey post-hoc test) (**Figure 1g**). These data suggest that *Grcc10* KO mice have higher underlying susceptibility to induced seizures, analogous to the increased rate of epilepsy seen in human Temtamy syndrome patients.

Grcc10 loss of function causes an increase in AMPA mediated mEPSCs

We next asked what synaptic changes may mediate the severe seizure activity seen in our mouse model. We hypothesized that we would be able to see increased neuronal excitability through AMPA (α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) mediated currents based on substantial evidence from many animal models that shows that these receptors are key mediators of seizure spread and epileptogenesis [17,18]. We measured electrophysiological parameters in *Grcc10* KO neurons by first assessing spontaneous AMPA mediated miniature excitatory postsynaptic currents (mEPSC) using whole cell recordings in isolated excitatory hippocampal neurons from E18 *Grcc10* WT, and KO mice on day *in vitro* (DIV) 14-18. We observed that AMPA-mediated mEPSCs recorded from *Grcc10* KO neurons have a significantly higher mean amplitude than mEPSCs recorded from *Grcc10* WT neurons (35.81±0.90 pA vs .15.49±0.29 pA, respectively, p<.0001, F=367, df=2, N=5 neurons per genotype, ANOVA with Tukey post-hoc test) (**Figure 2a-c**). These data support the hypothesis that *Grcc10* loss of function causes an increase in AMPA-mediated mEPSC amplitudes. In contrast, NMDA amplitude and frequency were not significantly different between WT and KO neurons (13.84±3.93 pA vs. 12.92±2.92 pA,

p=0.20, t=1.27, df=301.4, N=5, Welch's T-test) (**Figure 2d**). GABA current amplitude was also

not significantly different between WT and KO neurons (13.78 ± 3.32pA vs. 15.56 ± 8.37 pA,

(p=.16, t=1.41, df=137, N=5 Welch's T-test) (**Figure 2e**). To confirm that the loss of

C12ORF57/GRCC10 was a necessary and sufficient cause of the increase in AMPA mEPSC

activity in these cultured neurons, we sought to rescue the *Grcc10* KO neurons using lentiviral

vector containing *C12ORF57-P2A-GFP* construct under *Syn2* translational control with a *GFP*

only lentivirus serving as a control (**Supplemental Figure 4**). We transfected E18 *Grcc10* KO

neurons on DIV 7 with both the construct and a *GFP* only vector and compared their

electrophysiology profiles on DIV14-18 with WT lentiviral infected *GFP* controls. *Grcc10* KO

neurons transfected with *C12ORF57* had decreased mEPSC amplitude compared to *GFP*-

transfected KO neurons (18.27 ± 0.47 pA vs. 38.93 ± 0.56 pA, respectively, p<.0001) and had no

statistically significant difference in mEPSC amplitude compared to *GFP* transfected WT neurons

(15.50 ± 0.32 pA ; p=.57, t=0.17, df=137.2 ANOVA, N=4) (**Figure 2f**). Frequency of AMPA

mEPSC also decreased in *C12ORF57*-transfected compared to *GFP*-transfected KO neurons

(5.46 ± 0.61 events/sec vs. 22.46 ± 0.47 events/sec, respectively, p<.0001, ANOVA with Tukey

post-hoc test, F=85.3, df=2, N=5). *C12ORF57*-transfection rescued the phenotype in KO neurons

(19.06 ± 0.81 events/sec, P<.0001, ANOVA Tukey post-hoc test, N=5), which were now

comparable to WT neurons (5.21 ± 1.31 events/sec p>.999, ANOVA with Tukey post-hoc, df=2,

N=5) (**Figure 2g**). These data suggest that C12ORF57 specifically modulates post-synaptic

AMPA current and is sufficient to rescue the excessively increased excitability phenotype seen in

Grcc10 KO neurons and is necessary for normal neuronal electrophysiology.

Loss of *Grcc10/C12ORF57* leads to an increase in surface AMPA receptor localization and

impairs excitatory homeostatic synaptic down-scaling after blocking GABA receptors.

Given our findings of increased AMPA current in *Grcc10-/-* neurons we hypothesized that loss of

Grcc10/C12ORF57 would result in increased GluA2 surface expression. AMPA receptors are

post-synaptic heterotetrameric assemblies of two homologous subunits (GluA1-4), with

compositions containing two GluA2 subunits being the most predominant in hippocampal

pyramidal neurons and mediating the majority of fast excitatory synaptic transmission in the brain.

We stained for membrane bound GluA2 subunits on isolated DIV 17 *Grcc10* WT an KO)

pyramidal hippocampal neurons and quantified the fluorescence intensity of GluA2 positive

puncta per 50 μm of neuronal processes. We observed an increase in the number of GluA2 puncta

(83.72±23.10%,(p=.0024, t=3.62, df=15.32, N=10 per genotype Welch's T-test) and an increase

in puncta intensity (102.43±28.66%, p=.0008, t=3.572 df=51.84, N=19 WT, N=35 KO, Welch's

T-Test) in *Grcc10* KO neurons compared to WT neurons (**Figure 3a**).

We also sought to determine if *Grcc10* KO neurons would be less able to downregulate AMPA

current in response to increased excitation, which would be predicted with decreased CAMK4

activity in the KO mice. To do this, we measured surface AMPA receptor expression after inducing

a chronic increase in postsynaptic activity by treating neurons with bicuculline (a GABA receptor

antagonist) for 48 hours and then measuring GluA2 surface expression [19]. We observed that, as

expected, bicuculline treatment caused a significant decrease in surface GluA2 expression in WT

neurons (-907.00±293.20 puncta/HPF, p=.0041, t=3.09, df=32, untreated N=19, treated N=15

,Welch's T-test). In contrast, bicuculline treatment had no significant effect on GluA2 in *Grcc10*

KO neurons (-255.0±616.9 puncta/HPF, p=.68, T=.41, df=53, N=19 untreated, N=15 bicuculline

treated, Welch's T-Test) (**Figure 3b**). We also observed an increase in dendritic spine density in

in situ Grcc10 KO cortical neurons (1.26+/-0.53 spines/μm) compared to WT cortical neurons

0.91+/-0.22 spines/μm, p=.0023, t=3.185, df=57, N=33, Welch's T-Test) (Figure 3C), though

spine width was not statistically different between the two neuron types (33.53+/-18.57 vs 34.53+/-

19.75, p=.83, t=.198, df=57, N=33, Welch's T-test). These findings suggest that GRCC10 loss of

function increases dendritic spine density and GluA2 surface expression and impairs homeostatic

synaptic down-scaling of GluA2 and results in higher levels of AMPA expression at synapses both

at baseline and under conditions of acute increases in neuronal activity. This data further supports

our observation of increased AMPA current in *Grcc10-/-* mouse neurons described in the previous

section.

C12ORF57/Grcc10 loss of function causes a decrease in the expression of pCREB, cFOS and

ARC

Because the loss of C12ORF57 disrupts synaptic AMPA downscaling and current, we sought to

determine if there were changes to the negative regulation of AMPA synapses involving CREB

and ARC signaling. This is because phosphorylated CREB (p-CREB) regulation of the immediate

early gene, *Arc*, is implicated in the synaptic downscaling of AMPA receptors and the regulation

of surface GluA2 expression in response to neuronal excitation [9,20–28]. To investigate if this

pathway is altered in our KO mice, we stained primary hippocampal *Grcc10* WT and KO neurons

isolated at E18 at 20 days *in vitro* using an antibody that binds specifically to the Ser133 p-CREB

and ARC and measured fluorescence intensity. We observed that KO neurons express a

significantly lower level of Ser133-pCREB (-29.80±8.55%, (p=.0008, t=3.66, df=38, N=20,

Welch's T-Test) and ARC (-65.79±9.18%, p<.0001, t=7.17, df=38 N=24, Welch's T-Test)

compared to WT neurons (**Figure 4a-c**). These data were also supported by western blot findings

of decreased pCREB in whole brain lysates of KO mice compared to WT (-31.26±12.47%,

(p=.024, t=2.5, df=14.72, N=5, Welch's T-Test) and decreased intermediate early genes ARC (-

48.03$\pm$8.44%, p=.0036, t=5.69, df=4.36, N=5 Welch's T-Test) and cFOS (-49.94%$\pm$10.45,

=.0043, t=4.78, df-5.288, N=5, Welch's T-Test) (**Figure 4d-f**). These results suggest that *Grcc10*

loss of function leads to a decrease in CREB phosphorylation and decreased expression of ARC

and cFOS.

To further study the effect of *C12ORF57*/*Grcc10* loss of function on CRE-dependent

transcriptional activation, we performed a dual luciferase assay as described by Bellis et al. [29]. We

transfected both *C12ORF57* WT and KO HEK293T cells with one construct containing *Renilla*

luciferase as a reporter of CRE promoter activity and another one expressing firefly luciferase

under a constitutive promoter as a control. We observed that KO cells had a significant decrease

in normalized CRE-driven *Renilla* luminescence compared to WT cells (-25.40$\pm$3.78%, ANOVA,

F=20.74, p<.0001, Tukey post-hoc test, p=.019, N=3 per experimental condition). Moreover,

transfection of *C12ORF57*-/- cells with C12ORF57 restored CRE driven luciferase signal to

wildtype levels (+27.56$\pm$17.10%, p=0.12, Tukey post-hoc test) (**Figure 4g**). These data show that

loss of C12ORF57 reduces CRE-dependent transcriptional activation, potentially impacting the

negative regulatory elements upstream of AMPA, CREB and ARC.

C12ORF57 enhances CAMK4 activity through interactions with its autoinhibitory domain

The lack of significant homology of GRCC10 to any known protein impaired our ability to predict

its cellular mechanisms *a priori*. To bypass this limitation, we analyzed the GRCC10 protein

interactome using two independent unbiased Yeast 2-hybrid screens utilizing mouse brain libraries

[30]. Both unbiased genome-wide screens resulted in a single positive clone that encoded the calcium

calmodulin associated kinase IV (CAMK4). We conducted direct co-immunoprecipitation of

GRCC10 from whole WT and KO mouse brain lysate and confirmed its interaction with CAMK4

in WT lysate which was abolished in KO lysate (**Figure 5a**). Discovery of C12ORF57's interaction

with CAMK4 is particularly informative because CAMK4 is a known upstream regulator of both

CREB and ARC and plays a role in synaptic downscaling [9,23,31].

Given the close association of C12ORF57/GRCC10 with CAMK4, we next investigated whether

it plays a role in modulating CAMK4 activity. In prior studies, phosphorylated CAMK4 has been

shown to be largely confined to the nucleus[32,33]. This localization for CAMK4 was shown to be

mediated by phosphorylation at amino acid (AA) Thr200 in humans and via Thr196 in mice

[26,27,34,35]. We measured the expression of CAMK4 in *Grcc10* WT versus KO brains by western

blot and observed that differences between total CAMK4 were not significant between WT and

KO brains (-0.74 $\pm$0.73%, p=.34, t=1.01, df=6.97, N=6 Welch's T-Test). However, we did see a

shift in CAMK4 distribution within the nucleus of KO brains compared to WT brains with

significantly less pCAMK4 (-66.66$\pm$6.83%, p=.0037, F=18.33, df=3, N=5, ANOVA with Tukey)

and total CAMK4 (-48.1$\pm$6.73%, p=.002 F=14.89, N=5, df=3, ANOVA with Tukey post-hoc test).

We saw the opposite shift in cytoplasmic fraction, where KO brains showed an increase in total

CAMK4 (+31.6$\pm$20.69%, p=.0035 2-way ANOVA, F=5.50, DF=2, N=6) compared to WT, and a

similar, though not statistically significant, decrease in pCAMK4 (+31.87$\pm$19.51, p=.1848, N=6,

Tukey post-hoc test) (**Figure 5b-g**). These results suggest that loss of C12ORF57 disrupts the

balance of nuclear versus cytoplasmic CAMK4, with increased nuclear phosphorylated CAMK4

in WT brains when compared to KO brains.

To further uncover how C12ORF57/GRCC10 interacts with CAMK4 and because we observed

changes in CAMK4's phosphorylation and localization with C12ORF57 loss, we investigated the

roles of various regulatory domains of CAMK4. We created two modified forms of CAMK4

constructs lacking distinct regulatory domains: one which lacks the autoinhibitory domain (AA

305-321) (NAI), which binds CAMK4-negative regulatory proteins phosphatase 1 and 2 (PP1 and

PP2), and the other lacking the calmodulin binding domain (AA 322-341) (NCB) (**Figure 5h**)[36].

We co-transfected these constructs in HEK293T cells with a flag-tagged version of C12ORF57

and performed a co-immunoprecipitation assay with CAMK4. We observed that the deletion of

the autoinhibitory domain significantly reduced the ability of CAMK4 to bind to C12ORF57-

FLAG (-65.76±6.76%, p<.0001, N=8, ANOVA with Tukey post-hoc test) when compared to wild

type CAMK4, whereas deletion of the calmodulin binding domain resulted in no significant

change in binding (-6.13±6.81%, p=.66, ANOVA with Tukey post-hoc test, N=8) (**Figure 5h-j**).

This suggests that the CAMK4 autoinhibitory domain plays a significant role in its interaction with

C12ORF57.

The association of C12ORF57 with CAMK4's autoinhibitory domain that interacts with protein

phosphatases, along with the decrease in pCAMK4 in KO brain nuclei, led us to hypothesize that

C12ORF57/GRCC10 could disrupt the activity of phosphatases that modulate CAMK4 activity.

To investigate this, we measured *in vitro* CAMK4 kinase activity in the presence or absence of

C12ORF57 and/or serine/threonine-protein phosphatase alpha catalytic subunit 1 (PP1CA) using

an *in vitro* kinase enzyme system that enables the quantification of ATP consumption by CAMK4

via luminescence [31]. We found that the addition of PP1CA significantly decreases CAMK4 *in vitro*

activity by 39.32% (-981.00±75.95 Relative luminesce units, RLU, p=<.0001, ANOVA with

Tukey's post-hoc test), which is expected given its negative regulatory activity *in vivo* (**Figure**

5k). We also found that adding C12ORF57 does not affect CAMK4 kinase activity compared to

CAMK4 alone (-66.33±90.83 RLU, p=85, F=448.3, df=6, N=3,p=.95, ANOVA with Tukey's

post-hoc test). The presence of both C12ORF57 and PP1CA caused CAMK4 activity to decrease

significantly than just PP1CA alone (-681.00±67.39 RLU p>.0001, ANOVA with Tukey's post-

hoc test) (**Figure 5k**). This, in conjunction with our previous data, suggests that C12ORF57

enhances CAMK4 activity by reducing efficacy of phosphatase inhibition by binding to the

autoinhibitory domain of CAMK4 and increasing the relative amount of active phosphorylated

CAMK4.

Overexpression of CAMK4 is sufficient to rescue Grcc10 excitability phenotype.

Because we hypothesize that CAMK4 is downstream of C12ORF57/GRCC10, we tested whether

overexpression of CAMK4 would be sufficient to rescue the increased AMPA mEPSC phenotype

in *Grcc10-/-* neurons. We performed whole cell recordings on isolated excitatory hippocampal

neurons from embryonic day 18, *Grcc10* WT, and KO mice on DIV 14-18 that were transfected

with a lentivirus carrying a mouse *Camk4-Pp2A-Gfp* fusion under *CMV* promoter control versus a

GFP vector as a negative control and a *C12ORF57* positive control. *Grcc10* KO neurons

transfected with *CAMK4* had AMPA amplitudes statistically significantly lower (p<.0001,

F=240.8, dF, 3,N=6, ANOVA) than vector-infected KO neurons (12.54±2.29 pA vs. 46.11± 20.70

pA, respectively, p<.0001, Tukey post-hoc test) and equal to WT neurons (12.30±3.52 pA p=.99,

N=6, Tukey post-hoc test) and *C12ORF57*-transfected KO neurons (12.55±3.684pA, p=.9987,

N=6, Tukey post-hoc test) (**Figure 5l**). These data suggest that CAMK4 is indeed downstream of

C12ORF57 and the overexpression of CAMK4 is sufficient to overcome the loss of C12ORF57

and rescue the AMPA amplitude phenotype.

Discussion

Our *Grcc10* knockout mice model similar brain phenotypes (seizures, shortened corpus

callosum) to those observed in human Temtamy syndrome, caused by mutations in the closely

related and poorly characterized human ortholog, *C12ORF57*. Due to the lack of recognized

protein motifs or domains in C12ORF57, our *Grcc10* knockout mice provide a novel window to

elucidate the precise molecular function of these genes.

We showed that loss of *Grcc10* increased AMPA GluA2 in response to chemical stimulus. We

also showed that loss of *Grcc10* decreased known excitatory synaptic regulators CREB, Arc and

Fos, and that at baseline *Grcc10* -/- neurons express greater levels of AMPA and have increased

mEPSCs. This suggests that C12ORF57 is capable both of regulating the baseline excitatory

potential of neurons as well as the neurons dynamic response to overexcitation, suggesting it acts

as a neuronal excitatory "rheostat" whose loss leads to an unexpected and dramatic increase in

AMPA current and signaling. While the full mechanism and scope of this rheostat function will

require further study to elucidate, we believe CAMK4, the binding partner we discovered

provides plausible mechanistic insight.

The fine tuning of AMPA current would require a protein regulator, which is itself tightly and

dynamically regulated. The activation of CAMK4 is a transient and tightly regulated event

requiring both Ca^{2+}/calmodulin binding and phosphorylation of the kinase at residue Thr200 by

an upstream CaMK kinase (CaMKK) which traffics it into the nucleus [31]. Expression of either

dominant negative or constitutively active forms of CAMK4, is able to scale up or down

excitatory synaptic strength, respectively, and disrupts the cell's ability to dynamically respond

to stimuli [31,37,38]. As expected, a disruption in CAMK4 activity led to a destabilization of

neuronal homeostasis in our *in vitro* murine hippocampal excitatory neuronal model. Loss of

Grcc10 increased AMPA mEPSCs and GluA2 expression and the inability to downregulate

excitatory GluA2 in response to chemical stimulus, as observed with CAMK4 dominant negative

mutations. Further, similar to the overexpression of the dominant negative version of CAMK4,

we observed a decrease in the ratio of nuclear pCAMK4/CAMK4 in our mouse model. Our *in*

vitro data also supports this observation with the loss of C12ORF57 decreasing downstream

CAMK4 signaling through pCREB and ARC. CREB acts upstream of the immediate-early gene

ARC which is responsible for downscaling excitatory synapses in response to neuronal excitation

[26–28]. CREB and ARC signaling has been shown to increase endocytosis of surface GluA2

expression in response to increased calcium flux from neuronal excitation [9,20]. Prior published

evidence showed that ARC is involved in synaptic scaling by interacting with endophilin to

enhance endocytosis of AMPA receptors, which is crucial for neuronal excitatory/inhibitory

homeostasis [20–25]. CAMK4 is known to regulate the activity of CREB by phosphorylating its

Ser133 residue [32,39,40]. Because sequence analysis of human and murine CAMK4 shows identical

calmodulin and autoinhibitory domains, as well as 98.3% amino acid identity between these

orthologs, it is likely that our biochemical findings in mice would generalize to humans.

Previous studies showed that the autoinhibitory domain of CAMK4 stably associates with protein

serine/threonine phosphatases 1 and 2A (PP1; PP2A) to negatively regulate its kinase activity[31,41].

These interactions modulate CAMK4's role in homeostatic excitatory synaptic scaling, a

mechanism by which neurons adapt their synaptic strength through upregulation or

downregulation of surface expression of AMPA receptors [22,37,41]. We noted that C12ORF57 binds

CAMK4 and the presence of the autoinhibitory phosphatase binding domain was important in

maintaining this interaction. Further, we also found that C12ORF57 antagonized the PP1CA-

mediated decrease in CAMK4 activity *in vitro*. These results give a further plausible mechanism

by which C12ORF57 could regulate excitatory synaptic strength through modulation of CAMK4

activity by slowing its dephosphorylation.

Based on our observations, we propose a model where C12ORF57 protects CAMK4 from

dephosphorylation and bolsters its homeostatic role in reining in excitatory synaptic up-scaling in

excitatory neurons (**Figure 6**). In this model, the loss of C12ORF57 leads to a decrease in the

proportion of active, phosphorylated, CAMK4, reducing negative feedback of excitatory synaptic

scaling and increased AMPA expression at synapses. We suggest that these events lead to higher

baseline neuronal excitability and increased seizures seen in both TS patients and our murine

knockout model. Because sequence analysis of human and murine CAMK4 shows identical

calmodulin and autoinhibitory domains, as well as 98.3% amino acid identity between these

orthologs, it is likely that our biochemical findings in mice would generalize to humans. It should

be noted that this is not the first time CAMK4 activity is implicated in epilepsy in humans;

disruptions involving a regulatory miRNA and CAMK4 point mutations have been reported in

certain cases [42–44].

An important question to address in our model is why previous CAMK4 knockout mice did not

present with a similar seizure phenotype [45]. It is possible that loss of C12ORF57 in excitatory

neurons and a subsequent decrease in the pCAMK4/CAMK4 ratio creates a functional CAMK4

knockdown and that cells do not tolerate knockdown as well as a knockout or complete loss of

CAMK4 function. Previous work has shown that CAMK4 is sensitive to changes in both

subcellular localization as well as relative levels in relation to other CaM kinases which bolsters

this hypothesis[32,46,47]. In neuronal subsets such as excitatory neurons which have *high* levels of

endogenous CAMK4, the levels of other CaMKs such as CAMKI are the principle regulators of

neuronal firing, while in neurons with *low* endogenous CAMK4 levels, CAMK4 is the primary

regulator of neuronal firing [48]. For these reasons we propose the data suggests that it is the relative

levels of phosphorylated CAMK4 which cause disruptions in neuronal excitatory homeostasis

which are not captured in previous complete knockout experiments.

We acknowledge that our model is likely simplified, and that further study is required to elucidate

details of how C12ORF57 affects the kinetics, localization, and phosphorylation of various CaM

kinases. Also, while we saw no increased NMDA or GABA activity in *Grcc10* KO neurons, we

cannot fully determine whether this is due to changes in presynaptic release probability versus

purely increased postsynaptic excitation/detection, specifically its effect on AMPA current

frequency. Further work is necessary to determine what, if any, effect C12ORF57 has on

presynaptic transmission. Additionally, the high perinatal mortality makes it difficult to study

certain aspects of the knockout phenotype and limits broader behavioral characterization. The

development of a conditional knockout would be helpful to further studying the phenotypic

features found in human patients, such as optic coloboma, motor delay, social impairment, and

also epilepsy, since complete null mice cannot tolerate EEG placement and monitoring. Another

area for future study is that while most mutations currently discovered in human patients involve

a loss of function of the start codon, there are several missense, nonsense and splice site mutations

reported in compound heterozygotes such as p.L51Q mutations[4,8]. Patients with non-initiator

methionine mutations have similar phenotypes and severity (epilepsy, agenesis of the corpus

callosum and optic coloboma), suggesting that these mutations also result in loss of function of the

protein [1,2,4]. Further research into specific missense mutations may provide additional insight into

key amino acid residues and domains within the C12ORF57 protein.

In conclusion, we have found that the novel *Grcc10* knockout mouse shares many phenotypic

features with Temtamy syndrome patients, making it a suitable model in which to study the

function of its human ortholog, *C12ORF57*. We have shown through multiple lines of evidence in

murine, human cellular, and *in vitro* models that – a) C12ORF57 binds CAMK4, b) C12ORF57

loss decreases CAMK4 phosphorylation and increases mEPSC activity, and c) C12ORF57 loss

decreases downstream signaling through the synaptic scaling pathway of CREB, ARC, and GluA2.

These findings provide the first evidence for the molecular function of C12ORF57 in neuronal

development and homeostasis, as well as a possible explanation for the pathogenesis behind some

of the phenotypic features of Temtamy syndrome. Further studies, perhaps using conditional

knockout models, are needed to shed further light on additional functions of this protein and tease

out its precise role in synaptic homeostasis. Such studies have the potential to give us crucial

insight into how C12ORF57 regulates neuronal excitatory synaptic scaling and protects neurons

against excessive excitability.

Materials and Methods

Animal maintenance and generation of G*rcc10* mouse lines

All procedures of animal maintenance and experiments were in accordance with the regulations of

and approved by IRB at the University of California San Francisco and in compliance with relevant

federal and local laws. All animal breeding and procedures performed in the Richards lab were

approved by The University of Queensland Animal Ethics committee and performed in accordance

with the Australian Code of Practice for the Care and Use of Animals for Scientific Purposes.

Generation of the mouse *Grcc10<tm1.1(KOMP)Vlcg>* line by the Sherr lab was previously

described[49,50] .

Embryos and postnatal pups were bred from wildtype CD1 time-mated dams at The University of

Queensland. The detection of vaginal plug in the following morning after mating was designated

as embryonic day zero (E0). Pups are at postnatal day (P0) on day of birth. Tissue collection and

fixing was as previously described (9), then post-fixed with 4% paraformaldehyde (ProSciTech)

for 2 to 4 days and stored in 1x Dulbecco's phosphate buffered saline (PBS, Lonza) with 0.2%

sodium azide (Sigma Aldrich).

Primary Fibroblast Derivation and RT-PCR

Primary fibroblasts were derived from 10mm skin punch biopsies of C12ORF57 patient as

previously described by Kisiel et al.[51] All fibroblasts were below passage number 3. RNA was

extracted using PureLink RNA mini system (Thermo-fischer 12183018A). cDNA was generated

from RNA samples using superscript IV one Step RT-PCR system (ThermoFischer, 12594025).

RT PCR was run using the following primers: C12ORF57 F:5'-

GGATAACGCCTGCAACGACATG-3'   R: 5'-CTTCGTAGGACTTGACCAAGCG-3';

GAPDH: F: 5'-GTCTCCTCTGACTTCAACAGCG-3'; R: 5'-

ACCACCCTGTTGCTGTAGCCAA-3'

Kainic Acid Treatment

417   Postnatal day 21 genotyped mice received 20 mg/kg intraperitoneal kainic acid (Sigma-Aldrich

418   K0250) solution dissolved in saline or saline vehicle solution using sterile technique. An observer

419   blinded to mouse genotype recorded time to each Racine stage.

C12ORF57 Antibody generation

C12ORF57 peptide was generated using a full-length cDNA of human C12ORF57 conjugated to

N-terminal GST (NCBI9606) produced in BL21(DE3) competent *E.coli.* and purified against a

glutathione agarose (GE Lifescience 17075601). Rabbit polyclonal antibody was produced using

Thermo Fisher Custom Polyclonal Antibody production.

Antibody showed specific bands on western blot at 1:200 dilution in both CMV-C12ORF57-

FLAG at expected 17 kD weight and in Grcc10 WT mouse brains and not in Grcc10 KO brains

(Supplemental Figure 5).

Cellular Fractionation

429 Whole mouse brains were Dounce homogenized in fractionation buffer (20 mM HEPES, 10 mM

430 KCL, 2mM MgCL2, 1 mM EDTA, 1 mM EGTA, 1mM DTT) followed by passage through a 27-

431 gauge needle. The nuclear fraction was separated by centrifugation at 720 xg for 5 min.

432 Cytoplasmic fraction was separated at 10,000 xg for 5 min.

Western Blot

434 Tissue was homogenized with a Dounce homogenizer in RIPA buffer, and lysates were run on a

435 8-20% gradient SDS page. The blots were blocked with Odyssey Li-Cor blocking buffer (Licor).

436 Primary antibodies were followed by an IR fluorescent-conjugated goat anti-mouse and anti-rabbit

437 IgG (1:10 0000, Licor)/ Quantification and IF intensity analysis was done using Odyssey LiCor

438 Image Studio Software v.5.2.5

Antibodies

The following Primary antibodies were used: CAMK4: Invitrogen 8C5B8 (Mouse, MA5-

17038,1:1000), pCAMK4 Th200 (Rabbit Invitrogen PA5-37504, 1:1000), pCREB: (Mouse CST

87G3 #9197, 1:1000), CREB (Rabbit, CST,48H2, 1:1000), GFP (Chicken, A10626,

Invitrogen1:1000), Arc (Proteintech, Rabbit, 16290-1-AP, 1:1000WB), S6 Ribosomal protein,

(Rabbit, 5G10, 2217, Cell Signaling Technology) AMPA (6C4, Invitrogen, 32-0300), and CTCF

(CST, 28995).

The following secondary antibodies were used: IRDye680RD Goat anti-Rabbit IgG (H+L)

1:10,000 (Li-cor, 926-68071), IRDye680 Donkey Anti-Chicken (Li-cor, 926-68075), and

IRDye800 Goat anti-mouse IgG (Li-Cor, 926-32210).

Co-immunoprecipitation

Mouse brains were homogenized in NP40 lysis buffer (1% NP40, 240 mM NaCl, 5 mM EDTA 50

mM Tris ph7.4). 1µg of rabbit anti-C12ORF57 antibody was bound to 2µl of agarose resin and

incubated in 1.5 µg of total lysate overnight and co-immunoprecipitated using Pierce Co-

Immunoprecipitation kit (Thermo Fisher, 26159). Protein was dissociated using low pH Elution

buffer (Thermo Fisher, 21104)

Comparative Pull-down experiments

HEK293 cells were co-transfected with C12ORF57-DYKDDDK and CAMK4 constructs in a

pCMV backbone. Cells were lysed with 1% NP40 lysis buffer in 1x PBS.

1 ug of rabbit Anti-DYKDDDK primary was bound to 20 µl of agarose resin (Pierce, 26159) and

incubated in 1.5 ug of total protein overnight. Western blotting was performed as described above

and bands were quantified using an Odyssey Licor system as described above.

Hematoxylin and eosin staining

Hematoxylin and eosin (H&E) staining using standard H&E protocols was performed on P0

Grcc10 knock-out mouse brains. Dissected brain tissues were sectioned on vibratome (Leica) at

50µm thickness in sagittal and coronal orientation.

In situ hybridization

In situ hybridization was performed as described in Moldrich et al, 2010[52]. Two sets of Grcc10

riboprobes were generated in-house using the primers corresponding to those used in the Allen

Mouse Brain Atlas (2015 Allen Institute for Brain Science http://mouse.brain-

map.org/experiment/show/386251, Forward primer 1: 5' ACTCAAGCACCGAGTGGC 3',

Reverse primer 1: 5' CCTCTGGAACACAAGGGC 3') (Forward primer2: 5'

GCTTCCTTTCTGCCTCCTCT 3', Reverse primer 2: 5' CACAGCAGCAGCACACATAC 3').

and primers designed with Primer3Plus software. These primers were used to amplify a 459 or 519

base pair fragment, respectively, from mouse cortex cDNA. These fragments were purified and

cloned into pGEM-T Vector System (Promega). The plasmids were then linearized (NotI or SacII

restriction enzyme, New England BioLabs), purified (PCR Clean up Kit, Qiagen), transcribed (Sp6

or T7 RNA Polymerase, New England BioLabs) and digoxigenin-labelled (DIG RNA Labelling

Mix, Roche) or fluorescein-labelled (Fluorescein RNA Labelling Mix, Roche) to generate the

riboprobes.

***In Vitro* CAMK4 assay**

CAMK4 kinase activity was measured using ADP Glo System (Promega, V6930). CAMK4,

calmodulin, and reaction buffer were supplied by CAMK4 *in vitro* Kinase system (Promega,

V2951). C12ORF57-FLAG was transfected into HEK293 cells and purified from whole cell lysate

through immunoprecipitation on Anti-DYKDDDK Affinity resin (Genescript, L00432). CAMK4,

C12ORF57, and purified recombinant PP1CA (Origene, TA808819) were mixed in equimolar

amounts with 25 µM ATP and reaction luminescence was measured on a Tecan Spark for an

integration time of 1 sec.

Primary Hippocampal Neuronal Culture and Transfection

Embryos were removed from pregnant dams at embryonic day 18 (E18). Quick genotyping of the

mice was performed by PCR analysis of DNA, extracted from the tail biopsies. Tails were digested

in KAPA Express Extract Kit and then template DNA was added to PCR mix containing: 1X

KAPA2G Robust HotStart ReadyMix, and *Grcc10* primers (Forward*: 3'-*

GCGGACCTCTCTGGGATG-5' Reverse: 3'-GCAGCACACATACGAGAACC-5')

After genotyping, hippocampi were dissected in ice-cold Hanks' balanced salt solution (HBSS;

Gibco) and were incubated in trypsin/EDTA (Gibco BRL) at 37 °C for 25 min and subsequently

transferred to trypsin inhibitor (Worthington) at room temperature for 5 min. After trituration in

DMEM/10% FBS solution, cells were counted and plated on poly-L-lysine (Sigma) and Laminin

(Gibco) pre-coated cover glass (Harvard Apparatus) at a density of 40.000 cells per 10 mm cloning

rings. Cells were maintained in Neurobasal medium (Gibco) containing 1% heat-inactivated FBS,

1x B27-supplement (Gibico), 1X Glutamax (Gibco), 15 mM NaCl. Medium was partially

exchanged for fresh medium every 7 days. On days 3-5 (days in vitro, DIV), mitotic inhibitor 5-

fluoro-2′-deoxyuridine (FUDR, 10 μM) was added to control glial growth.

For the transfected experiments, on day 7, *Grcc10+/+* neurons were infected by custom lentiviral

particles produced from CW307508 (Origene) with C12ORF57-P2A-GFP under control of a

synapsin 2 promoter (Catalog No: CW307508V) at a concentration of 1:100 GFP under *Syn2*

(RC218378LF) or CAMK4-P2A-GFP (Origene Ca MR207505L4V).

In vitro patch clamp recordings

Patch clamp measurements were done at DIV 14-17. Before recordings, Neurobasal medium was

replaced by HEPES ACSF, containing (in mM): NaCl, 140; KCl, 2.5; HEPES, 10; MgCl2, 2;

$CaCl_2$, 2; and dextrose, 10. pH was adjusted to 7.3 with NaOH. mOsm/kg was set to 300. Cultures

were visualized under upright microscopy (Olympus, BX51WI) and cells with pyramidal

morphology were identified by ×40 water-immersion objective (Olympus) and IR-2000 (DAGE-

MTI) camera GFP was identified with 488 nm LED pulse. Whole-cell patch clamp recordings

were done from the soma of the pyramidal hippocampal neurons using Axopatch 200B amplifier

(Molecular Devices, USA). Cell capacitance and access resistance were monitored throughout the

experiments. Signals were acquired with a DigiData-1440A digitizer controlled by pCLAMP-11

software (Molecular Devices, USA). Signals were sampled at 10 kHz and low-pass filtered at

2 kHz. Borosilicate glass pipettes (4–5 MΩ) were pulled with a horizontal puller (P-1000, Sutter

Instrument).

Miniature excitatory currents (mEPSCs) were isolated from the GABAergic transmission by

applying 50 μM $GABA_A$ blocker picrotoxin in ACSF. To capture mEPSCs, voltage-gated Na+

channels and NMDA receptor-mediated currents were blocked with 1 μM TTX and 50 μM D-

AP5, respectively. Patch pipettes were filled with Cs-based intracellular solution (in mM):

$CsMeSO_3$, 135; CsCl, 6; HEPES, 10; phosphocreatine, 10; EGTA, 0.6; Mg-ATP, 3; Na-GTP, 0.5;

QX314-Br, 2. pH was set to 7.35 by CsOH and osmolarity to 305 - 310 mOsm/kg. mEPSCs were

measured from the holding membrane potential of -70 mV. NMDA receptor-mediated currents

were recorded using the same external solution, without magnesium and D-AP5, and with the

addition of NBQX to block AMPA currents (10 μM, Tocris). GABA receptor-mediated currents

were recorded using external solution without picrotoxin, but with D-AP5, TTX and NBQX. Data

analyses were done in Easy Electrophysiology software (Easy Electrophysiology Inc) utilizing a

template modeling process for mEPSC detection.

C12ORF57 HEK293 Knockout

536 sgRNA were generated with the sequence 3'-GATAACGCCTGCAACGACAT-5' and inserted

537 into pCAS_Guide-EF1a-GFP vectors (Origene). HEK293 17T cells were transfected with

538 Lipofectamine 3000 (ThermoFisher, L3000001) and transfection efficiency was determined via

539 immunofluorescence. Lines were picked and maintained with DMEM+10% FBS containing

540 puromycin at 1 ug/ml.

Golgi Cox Staining and dendritic spine density

543 P18 mouse brains were fixed in formalin and frozen sectioned. Golgi-Cox staining was performed

544 on these neurons utilizing the FD rapid GoligStain Kit (FD Neurotechnologies Inc PK401A).

545 Neurons were chosen at random from slices of mouse cortex and imaged on a Nikon T1 inverted

546 microscope. Dendritic spines were counted and analyzed using ImageJ software plug in Dendritic

547 spine counter.

Neuronal Surface Staining

Live cultured KO and WT neurons were incubated with a primary antibody directed against the

extracellular domain of GluA2. Incubation was conducted at room temperature to slow down

endocytosis of the antibody as shown previously [20,37] After incubation with the primary antibody,

cultures were washed and fixed for standard immunostaining. GluA2 positive puncta were imaged

using confocal microscopy. Their number and fluorescence intensity were quantified over 50 μm

of neuronal processes using ImageJ software.

Bicuculline Inhibition

Surface AMPA receptor expression after induction of a chronic increase in postsynaptic activity

was assessed by measuring GluR2 surface expression before and after treatment of neurons with

bicuculline for 48 hours, as described previously[20,49] .

Statistical analysis

Comparisons between Genotypes:

Analysis of Variance (ANOVA): T-test or Mann Whitney Mann Whitney or ANOVA were

appropriate was conducted to assess the overall differences between genotypes. When possible, all

measurements were done by blinded observers. All T-test were two tailed.

Post hoc Analysis: In case of significant differences identified by ANOVA, post hoc tests (Tukey's

Honestly Significant Difference test, Sidak's multiple comparison test) were performed to

determine specific pairwise differences between groups.

Error bars represent ±standard error of the mean (SEM) in all graphs.

Statistical software

All statistical analyses were performed using Prism version 10.0.0. The significance level (α) was

set at 0.05 for statistical tests. Statistical differences are indicated in the figures and tables using

the following symbols: $*P \leq .05, **P \leq .01, ***P \leq .001, ****P < .0001$.

References

1. Li, J. *et al.* Agenesis of the corpus callosum, optic coloboma, intractable seizures,

craniofacial and skeletal dysmorphisms: an autosomal recessive disorder similar to Temtamy

syndrome. *Am. J. Med. Genet. A.* **143A**, 1900–1905 (2007).

2. Akizu, N. *et al.* Whole-exome sequencing identifies mutated c12orf57 in recessive corpus

callosum hypoplasia. *Am J Hum Genet* **92**, 392–400 (2013).

3. Alrakaf, L. *et al.* Further delineation of Temtamy syndrome of corpus callosum and ocular

abnormalities. *Am J Med Genet A* **176**, 715–721 (2018).

4. Zahrani, F., Aldahmesh, M. A., Alshammari, M. J., Al-Hazzaa, S. A. F. & Alkuraya, F. S.

Mutations in c12orf57 cause a syndromic form of colobomatous microphthalmia. *Am. J.*

Hum. Genet. **92**, 387–391 (2013).

5. Temtamy, S. A. *et al.* New autosomal recessive multiple congenital abnormalities/mental

retardation syndrome with craniofacial dysmorphism absent corpus callosum, iris colobomas

and connective tissue dysplasia. *Clin. Dysmorphol.* **5**, 231–240 (1996).

6. Alfiya, F. *et al.* C12orf57 pathogenic variants: a unique cause of developmental

encephalopathy in a south Indian child. *J. Genet.* **101**, 30 (2022).

7. Wang, Y. *et al.* Temtamy syndrome caused by a new C12orf57 variant in a Chinese boy,

including pedigree analysis and literature review. *Exp. Ther. Med.* **19**, 327–332 (2020).

8. Platzer, K. *et al.* Exome sequencing identifies compound heterozygous mutations in

C12orf57 in two siblings with severe intellectual disability, hypoplasia of the corpus

callosum, chorioretinal coloboma, and intractable seizures. *Am. J. Med. Genet. A.* **164A**,

1976–1980 (2014).

9. Turrigiano, G. G. The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135**,

422–435 (2008).

10. Guerrini, R. & Dobyns, W. B. Malformations of cortical development: clinical features and

genetic causes. *Lancet Neurol.* **13**, 710–726 (2014).

11. Franco, S. J. *et al.* Fate-restricted neural progenitors in the mammalian cerebral cortex.

Science **337**, 746–749 (2012).

12. Molyneaux, B. J., Arlotta, P., Menezes, J. R. L. & Macklis, J. D. Neuronal subtype

specification in the cerebral cortex. *Nat. Rev. Neurosci.* **8**, 427–437 (2007).

13. Bertoglio, D. *et al.* Kainic Acid-Induced Post-Status Epilepticus Models of Temporal Lobe

Epilepsy with Diverging Seizure Phenotype and Neuropathology. *Front. Neurol.* **8**, 588

(2017).

14. Tripathi, P. P. *et al.* Increased susceptibility to kainic acid-induced seizures in Engrailed-2

knockout mice. *Neuroscience* **159**, 842–849 (2009).

15. Kokate, T. G., Cohen, A. L., Karp, E. & Rogawski, M. A. Neuroactive steroids protect

against pilocarpine- and kainic acid-induced limbic seizures and status epilepticus in mice.

Neuropharmacology **35**, 1049–1056 (1996).

16. Racine, R. J. Modification of seizure activity by electrical stimulation. II. Motor seizure.

Electroencephalogr Clin Neurophysiol **32**, 281–94 (1972).

17. Anggono, V. & Huganir, R. L. Regulation of AMPA receptor trafficking and synaptic

plasticity. *Curr. Opin. Neurobiol.* **22**, 461–469 (2012).

18. Rogawski, M. A. & Donevan, S. D. AMPA receptors in epilepsy and as targets for

antiepileptic drugs. *Adv. Neurol.* **79**, 947–963 (1999).

19. Schanzenbächer, C. T., Langer, J. D. & Schuman, E. M. Time- and polarity-dependent

proteomic changes associated with homeostatic scaling at central synapses. *eLife* **7**, e33322

(2018).

20. Joseph, A. & Turrigiano, G. G. All for One But Not One for All: Excitatory Synaptic Scaling and Intrinsic Excitability Are Coregulated by CaMKIV, Whereas Inhibitory Synaptic Scaling Is Under Independent Control. *J Neurosci* **37**, 6778–6785 (2017).

21. Rial Verde, E. M., Lee-Osbourne, J., Worley, P. F., Malinow, R. & Cline, H. T. Increased expression of the immediate-early gene arc/arg3.1 reduces AMPA receptor-mediated synaptic transmission. *Neuron* **52**, 461–474 (2006).

22. Shepherd, J. D. *et al.* Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. *Neuron* **52**, 475–484 (2006).

23. Kessels, H. W. & Malinow, R. Synaptic AMPA receptor plasticity and behavior. *Neuron* **61**, 340–350 (2009).

24. Noel, J. *et al.* Surface expression of AMPA receptors in hippocampal neurons is regulated by an NSF-dependent mechanism. *Neuron* **23**, 365–376 (1999).

25. Shepherd, J. D. & Huganir, R. L. The cell biology of synaptic plasticity: AMPA receptor trafficking. *Annu. Rev. Cell Dev. Biol.* **23**, 613–643 (2007).

26. Bito, H., Deisseroth, K. & Tsien, R. W. CREB phosphorylation and dephosphorylation: a Ca(2+)- and stimulus duration-dependent switch for hippocampal gene expression. *Cell* **87**, 1203–14 (1996).

27. Soderling, T. R. The Ca-calmodulin-dependent protein kinase cascade. *Trends Biochem. Sci.* **24**, 232–236 (1999).

28. Nonaka, M. *et al.* Towards a better understanding of cognitive behaviors regulated by gene expression downstream of activity-dependent transcription factors. *Neurobiol. Learn. Mem.* **115**, 21–29 (2014).

29. Bellis, A. D. *et al.* Cellular arrays for large-scale analysis of transcription factor activity.

Biotechnol. Bioeng. **108**, 395–403 (2011).

30. Parrish, J. R., Gulyas, K. D. & Finley, R. L. Yeast two-hybrid contributions to interactome

mapping. *Curr. Opin. Biotechnol.* **17**, 387–393 (2006).

31. Anderson, K. A., Noeldner, P. K., Reece, K., Wadzinski, B. E. & Means, A. R. Regulation

and function of the calcium/calmodulin-dependent protein kinase IV/protein serine/threonine

phosphatase 2A signaling complex. *J. Biol. Chem.* **279**, 31708–31716 (2004).

32. Enslen, H. *et al.* Characterization of Ca2+/calmodulin-dependent protein kinase IV. Role in

transcriptional regulation. *J. Biol. Chem.* **269**, 15520–15527 (1994).

33. Impey, S. *et al.* Phosphorylation of CBP mediates transcriptional activation by neural

activity and CaM kinase IV. *Neuron* **34**, 235–44 (2002).

34. Nakamura, Y. *et al.* Immunohistochemical localization of Ca(2+)/calmodulin-dependent

protein kinase kinase beta in the rat central nervous system. *Neurosci. Res.* **39**, 175–188

(2001).

35. Tokumitsu, H. *et al.* Activation mechanisms for Ca2+/calmodulin-dependent protein kinase

IV. Identification of a brain CaM-kinase IV kinase. *J Biol Chem* **269**, 28640–7 (1994).

36. Naz, H., Islam, A., Ahmad, F. & Hassan, M. I. Calcium/calmodulin-dependent protein

kinase IV: A multifunctional enzyme and potential therapeutic target. *Prog. Biophys. Mol.*

Biol. **121**, 54–65 (2016).

37. Ibata, K., Sun, Q. & Turrigiano, G. G. Rapid synaptic scaling induced by changes in

postsynaptic firing. *Neuron* **57**, 819–826 (2008).

38. Goold, C. P. & Nicoll, R. A. Single-cell optogenetic excitation drives homeostatic synaptic

depression. *Neuron* **68**, 512–28 (2010).

39. Matthews, R. P. *et al.* Calcium/calmodulin-dependent protein kinase types II and IV

differentially regulate CREB-dependent gene expression. *Mol Cell Biol* **14**, 6107–16 (1994).

40. Sun, P., Enslen, H., Myung, P. S. & Maurer, R. A. Differential activation of CREB by

Ca2+/calmodulin-dependent protein kinases type II and type IV involves phosphorylation of

a site that negatively regulates activity. *Genes Dev.* **8**, 2527–2539 (1994).

41. Kasahara, J., Fukunaga, K. & Miyamoto, E. Differential effects of a calcineurin inhibitor on

glutamate-induced phosphorylation of Ca2+/calmodulin-dependent protein kinases in

cultured rat hippocampal neurons. *J. Biol. Chem.* **274**, 9061–9067 (1999).

42. Sun, J. *et al.* Identification of serum miRNAs differentially expressed in human epilepsy at

seizure onset and post-seizure. *Mol. Med. Rep.* **14**, 5318–5324 (2016).

43. Wang, Y. & Li, Z. Association of UGT2B7 and CaMK4 with response of valproic acid in

Chinese children with epilepsy. *Therapie* **75**, 261–270 (2020).

44. Zech, M. *et al.* A unique de novo gain-of-function variant in CAMK4 associated with

intellectual disability and hyperkinetic movement disorder. *Cold Spring Harb Mol Case Stud*

4, (2018).

45. Takao, K. *et al.* Comprehensive behavioral analysis of calcium/calmodulin-dependent

protein kinase IV knockout mice. *PloS One* **5**, e9460 (2010).

46. Wayman, G. A. *et al.* Regulation of axonal extension and growth cone motility by

calmodulin-dependent protein kinase I. *J. Neurosci. Off. J. Soc. Neurosci.* **24**, 3786–3794

(2004).

47. Wayman, G. A., Tokumitsu, H., Davare, M. A. & Soderling, T. R. Analysis of CaM-kinase

signaling in cells. *Cell Calcium* **50**, 1–8 (2011).

48. Cohen, S. M. *et al.* Excitation-Transcription Coupling in Parvalbumin-Positive Interneurons

Employs a Novel CaM Kinase-Dependent Pathway Distinct from Excitatory Neurons.

Neuron **90**, 292–307 (2016).

49. Skarnes, W. C. *et al.* A conditional knockout resource for the genome-wide study of mouse

gene function. *Nature* **474**, 337–42 (2011).

50. Okazaki, Y. *et al.* Analysis of the mouse transcriptome based on functional annotation of

60,770 full-length cDNAs. *Nature* **420**, 563–73 (2002).

51. *Skin Tissue Engineering: Methods and Protocols*. (Humana, New York, NY, 2019).

doi:10.1007/978-1-4939-9473-1.

52. Moldrich, R. X. *et al.* Molecular regulation of the developing commissural plate. *J. Comp.*

Neurol. **518**, 3645–3661 (2010).

Acknowledgments

Thanks to the Linda Richards Lab for their assistance with *in situ* hybrydization lab. Theywere

supported by an NHMRC Principal Research Fellowship (GNT1120615).

We thank the staff of the University of Queensland Biological Resources (UQBR) animal facility

and the QBI Advanced Microscopy and Histology Facilities, for their expertise and assistance in

this project.

Microscopy support was provided by the Center for Advanced Microscopy and the Nikon Imaging

Center at UCSF

Suling Wang for providing the illustrations in Figure 6 and Figure 1 Panel 1D.

We would like to thank the multiple families and patients who provided samples and participated

in our research.

Author Contributions

RJ: drafting/revision of the manuscript for content, Data for found in figures 1-6.

MC: drafting/revision of the manuscript for content, data found in figures 2-4 including

electrophysiology

CM: Contributed data on H&E, immunohistochemistry and *in situ* hybridization.

JB: Data on gene expression

JL: drafting/revision of the manuscript for content, Data found in supplemental figure 1

LR: supervision of laboratory work at University of Queensland, editing manuscript.

EU: drafting/revision of the manuscript for content, supervision of laboratory work at UCSF

ES: drafting/revision of the manuscript for content, supervision of laboratory work

EA and FA: Diagnosed case and established the fibroblast line for human patient

Ethics declarations

Competing interests

The authors declare no competing interests.

Figure 1 *Grcc10-/- Mice Recapitulate the Temtamy Syndrome Phenotype* **a)** RT PCR of C12ORF57 from fibroblasts derived from a human patient with homozygous c.1 A>G mutations in C12ORF57 see loss of transcript when compared to wild type controls. **b)** Ratios of surviving

Grcc10 genotype mice are shown at ages P0, P3, and P21 (N=154, p=0.99, Chi-Square). **c)** Mean

body mass (g) are shown for ages P0, P5, P9, P13, and P21 (N=5 for KO mice N=14 for WT and

heterozygotes, p>.0001 ANOVA) **d)** *In situ* hybridization of C12ORF57 shows diffuse widespread

expression of *Grcc10/C12ORF57* throughout all brain regions throughout embryonic

development. **e)** Representative images of midsagittal slices from *Grcc10+/+, Grcc10+/-* and

Grcc10+/+ mouse brains along with graphs showing mean of cortical length (p=.1127, ANOVA

with Tukey post-hoc test), corpus callosum length (p=.017, ANOVA with Tukey post-hoc test)

and corpus callosum cross sectional area cross-sectional area (p=.0058, ANOVA with Tukey's

post hock). N=6 for KO, N=3 for WT, N=3 for Het. **f)** Pictographic of Racine stages G) Average

Racine stage (x-axis) of mice in time post kainic acid injection (y-axis) for *Grcc10+/+* (blue, N=6),

Grcc10+/-(pink, N=7), and *Grcc10-/-*(green, N=4) (p=.00162, ANOVA with post hoc Tukey).

For all Error bars represent SEM. Statistical differences are indicated in the figures using the

following symbols $*P \leq .05, **P \leq .01, ***P \leq .001, ****P < .0001$.

Figure 2 *Grcc10 -/- neurons have increased mEPSCs amplitude and frequency* **a)** Representative

AMPA mEPSC tracings of *Grcc10* +/+ (WT) neurons (top), *Grcc10* +/- (Het) neurons (middle)

and *Grcc10* -/- neurons (bottom). **b)** Graph of cumulative frequency (x-axis) of mEPSC against

amplitude in DIV 17 pyramidal neurons. The mean curve of each group is bolded.   **c)** Mean AMPA

761 mEPSC amplitudes (pA) (p<.0001, N=5 for each group, ANOVA) **d)** Mean NMDA mEPSC

762 amplitudes (pA). (p=.20 N=5 Welch's T-test) Error bars represent SEM. **e)** Mean GABA mEPSC

763 amplitude (p=.16, N=5 for each group, Welch's T-test) **f and g)** Upper panels: Cumulative

764 frequency of events plotted against event amplitude (left panel) and the time between events

765 (events/second) (right panel) in DIV17 pyramidal neurons including C12ORF57 transfected rescue

766 (purple) and GFP Control (green). The combined mean of group is represented by corresponding

767 bolded line. Lower panels: Mean amplitude of AMPA mEPSC (left panel) and mean events/second

768 of AMPA mEPSC (right panel). For (**f**) and (**g**), N=5 for each group, ANOVA with Tukey post-

769 hoc test). For all graphs, error bars represent SEMs. Statistical differences are indicated in the

770 figures using the following symbols $*P \leq .05$, $**P \leq .01$, $***P \leq .001$, $****P < .0001$.

Figure 3 *Loss of Grcc10/C12ORF57 disrupts normal GluA2 expression in primary murine neurons* **a)** (i) Representative images of surface staining of MAP (white, upper panels) and GluA2 (red, lower panels) in *Grcc10 +/+* and *Grcc10-/-* neurons. Insets 2x magnification. Scale bar 20

μm. Mean GluA2 fluorescent puncta/100μm (ii) (p=.0024, N=10 both genotypes Welch's T- test),

and mean puncta intensity (ii) (p=.0008, N=19 WT, N=35 KO, Welch's T-Test) Error bars

represent SEM. **b)** (i) Sample images of DIV 17 hippocampal primary neurons stained for GluA2

(red) and MAP2 (white) on both *Grcc10* WT and *Grcc10* KO neurons with and without bicuculline

treatment. Scale bar 20 μm. Insets 10x magnification (ii) GluA2 puncta/100 μm on with and

without bicuculine, with mean represented as horizontal line (p=.0041, N=19 WT Ctl, N=15

WT+bicuculline treated Welch's T-test. Plot of GluA2 puncta/100μm on KO neurons (p=.68,

T=.41, df=53, N=20 KO Ctl, N=35 KO+bicuculline treated. Welch's T-Test). Scale bar 20 μm. **c)**

(i) Representative Golgi-Cox-stained dendrites from *Grcc10* WT and KO cortical neurons (scale

bar 5μm). (ii) Plot of spine density (p=.0023, N=33 per group, Welch's T-Test) and (iii) spine

width (p=.83, N=33 per group, Welch's T-Test). Mean represented by horizontal line. Statistical

differences are indicated in the figures using the following symbols $*P \leq .05$, $**P \leq .01$, $***P \leq$

$.001$, $****P < .0001$.

Figure 4 *Loss of C12ORF57 decreases levels of downstream CAMK4 targets* **a)** IHC staining of

primary hippocampal neurons from *Grcc10* (upper panels) and *Grcc10* KO (lower panels) mice

for phospho-CREB (green, left), ARC (red, center) with overlay of MAP-2 Yellow and DAPI

(blue, right). Scale bar 5μm **b and c)** Fluorescent quantification of ARC (**b**) (p=.0008, N=20 for

each genotype, Welch's T-Test) and **p-CREB (c)** (p<.0001, N=24, Welch's T-Test). Normalized means are shown. **d) Left:** Western blot of whole brain lysates of pCREB (top) CREB (middle) and GAPDH controls (bottom). **Right**: Mean ratio of CREB/pCREB fluorescence relative to GAPDH (p=.024, N=10, Welch's T-Test). Normalized to WT Mean. **e) Left:** Western blot of whole brain lysates from *Grcc10* WT and *Grcc10* KO mice with anti-ARC (top) and anti-actin control (bottom) antibodies. Right: Mean ratios of ARC band intensity to actin normalized to mean WT intensity (p=.0036, N=5 for each genotype, Welch's T-Test). **f)** Left: Western blot of whole brain lysates from *Grcc10* KO and WT mice with anti-cFos (top) and anti-actin control (bottom) antibodies. Right: Mean ratio of cFos band intensity to actin normalized to WT mean (p=.0043, N=5 for each genotype) **g)** Mean relative luminescent signal from Renilla luciferase assay on HEK293 cell lysates normalized to WT control (N=6 for each genotype, Welch's T-Test.). For all graphs, error bars represent SEM. Statistical differences are indicated in the figures using the following symbols *P ≤ .05, **P ≤ .01, ***P ≤ .001, ****P < .0001.

Figure 5 *C12ORF57 binds CAMK4 and modulates its phosphorylation and activity* **a)** Western

blot showing co-immunoprecipitation of CAMK4 with C12ORF57 from *Grcc10+/+* whole brain

lysate (left) with *Grcc10-/-* brain negative control (right) showing no GRCC10 expression or

CAMK4 pulldown **b)** Representative CAMK4 and GAPDH western blot bands from brain whole

cell lysates of *Grcc10 +/+* and *Grcc10-/-* mice **c)** Mean relative CAMK4 western blot band

intensity normalized to WT (p=.34, N=6 for each genotype Welch's T-Test) Error bars represent SEM. **d)** Representative CAMK4, pCAMK4 and lamin (control) bands on western blot from nuclear fraction of *Grcc10* WT and KO whole brain lysates **e)** Relative CAMK4, pCAMK4and Lamin band intensity normalized to *Grcc10 WT* levels. (p=.001, N=6 for each genotype, 2-way ANOVA with Sidak's multiple comparison test). **f)** Representative CAMK4, pCAMK4 and GAPDH (control) bands on western blot from cytoplasmic fractions of whole brain g) Relative CAMK4 pCAMK4 and Lamin band intensity normalized to *Grcc10* WT levels (p=.016, N=6 for each genotype, 2-way ANOVA, Sidak's multiple comparison test) in KO compared to WT. **h)** Schematic of WT CAMK4 (top) with kinase domain (blue), calmodulin binding domain (purple) and autoinhibitory/regulatory domain (yellow). Construct Δ(322-341) which lacks the autoinhibitory domain (NAI, middle) and construct Δ(305-321) which lacks the calmodulin binding domain (NCB, bottom) **i)** Representative western blot of co-IP of CAMK4 constructs with C12ORF57-FLAG **j)** Relative CAMK4 band intensity between WT, NCB (p=.24, N=8 replicates for each construct, ANOVA) and NAI (p<.0001, N=8 replicates for each construct, ANOVA) CAMK4 constructs.. **k)** Relative luminescence from *in vitro* CAMK4 kinase activity assay for C12ORF57, CAMK4, CAMK4+C12ORF57, CAMK4+PP1CA, CAMK4+PP1CA+C12ORF57, no CAMK4, and no ATP (N=3 per condition, ANOVA). **l)** Left: Graph of cumulative frequency (x-axis) of mEPSC against amplitude (pA) in DIV 17 pyramidal neurons including CAMK4 transfected control (green) KO (pink) WT (black), and C12ORF57 transfected (purple). The mean curve of each group is bolded. Right: Mean amplitude of WT, KO, CAMK4 transfected (KO+CAMK4) and C12OR57 transfected (KO+C12ORF57) (p<.0001, N=6 for each group, ANOVA). For all graphs, error bars represent SEM. Statistical differences are indicated in the figures using the following symbols *$P \leq .05$, **$P \leq .01$, ***$P \leq .001$, ****$P < .0001$.

Figure 6 *Proposed Model of C12ORF57 Function*

Proposed normal function of C12ORF57 (top panel). Under normal conditions C12ORF57 binds

CAMK4 and blocks dephosphorylation leading to increased active CAMK4 and activation of ARC

and down regulation of AMPA. Loss of C12ORF57 (bottom), where in response to calcium ion

influx during depolarization, more CAMK4 is dephosphorylated, and less is able to translocate to

841   the nucleus, decreasing downstream pCREB transcription and ARC transcription resulting in

842   unopposed long-term potentiation, increased GluA2 on cell surface and more AMPA current.