1   Intestinal RICT-1 regulates the larval germline progenitor pool via the vitellogenin VIT-3
2   in *C. elegans*
3

4   Anke Kloock,
5   E. Jane Albert Hubbard*
6
7   Department of Cell Biology,
8   NYU Grossman School of Medicine
9   New York, NY 10016
10
11
12
13   *Corresponding author

ABSTRACT

Populations of proliferating cells such as stem cells and tumors are often nutrient responsive. Highly conserved signaling pathways communicate information about the surrounding environmental, organismal, and cellular nutrient conditions. One such pathway is the Target of Rapamycin (TOR) pathway. The TOR kinase exists in two complexes, TOR complex 1 (TORC1) and TOR complex 2 (TORC2). TORC1 has been researched extensively and its regulation, particularly by amino acids, is well characterized. TORC1 activity promotes both stem cell fate and proliferation in the *Caenorhabditis elegans* hermaphrodite germline stem cell system to facilitate expansion of the larval germline Progenitor Zone (PZ) pool in response to nutrients. By contrast, a role for TORC2 in germline development has not been investigated. Here, we show that RICT-1, the sole ortholog of the TORC2-specific component RICTOR, also promotes expansion of the larval PZ, acting largely through SGK-1. Further, unlike the germline-autonomous role for TORC1 components, intestinal *rict-1* is both necessary and sufficient for full germline PZ pool establishment. Furthermore, neither DAF-2/IIS nor DAF-7/TGF-ß pathways mediate the effects of RICT-1. Rather, intestinal RICT-1 likely acts via vitellogenins, intestinally produced yolk proteins previously characterized for provisioning the adult germ line, but not previously characterized for a role in larval germ line development. By comparative RNA-seq on staged L4 larvae, we found vitellogenin genes among highly differentially abundant mRNAs. Genetic analysis supports a role for *vit-3* in germline development in a linear pathway with *rict-1*. Our results establish the *C. elegans* germ line as a fruitful model for investigating TORC2 and its connection to stem cells and lipid biology.

Keywords: TORC2, Stem Cells, Rictor, Vitellogenins

INTRODUCTION

Highly conserved signaling pathways govern cellular responses to nutrients and metabolites. These responses are required for the proper regulation of pools of proliferating cells such as stem cells and tumor cells (Puca et al., 2022; Tu et al., 2023).

TOR is a highly conserved serine/threonine kinase that is conserved from yeast to mammals and integrates nutritional signals to regulate cellular responses (reviewed in (Liu and Sabatini, 2020)). TOR acts in two complexes bearing key subunits Raptor in TORC1 and Rictor in TORC2. Both complexes respond to nutritional signals and regulate many nutrient-dependent aspects of stem cell biology (reviewed in (Russell et al., 2011; Saba et al., 2021)). Albeit differently, both complexes respond to amino acid availability and impact lipid metabolism (reviewed in (Caron et al., 2015; Szwed et al., 2021; Zoncu et al., 2011)) and homeostasis (Kim and Chen, 2004). TORC2 furthermore promotes *de novo* fatty acid and lipid synthesis in mice (Guri et al., 2017) and worms (Jones et al., 2009; Soukas et al., 2009). In *C. elegans,* our prior work established that TORC1 components and/or downstream effectors affect germline stem cell proliferation and differentiation germ line-autonomously and in response to amino acid provision (Korta et al., 2012; Roy et al., 2018; Roy et al., 2016).

To address the functions of TORC2 distinct from those of TORC1, Rictor mutants have been characterized in rodents, worms, and flies. In all three systems, reducing Rictor activity results in animals that are smaller and develop slower (Guertin et al., 2006; Hietakangas and Cohen, 2007; Jones et al., 2009; Soukas et al., 2009). Mutants in rodents and worms display high fat accumulation (Frei et al., 2023; Jones et al., 2009; Soukas et al., 2009; Yen et al., 2010), suggesting Rictor facilitates trafficking of lipids from the intestine to other organs (Jones et al., 2009; Soukas et al., 2009). Adult *C. elegans* hermaphrodites bearing reduction-of-function mutations in *rict-1*, the sole Rictor ortholog(Jones et al. 2009), exhibit reduced intestinal expression of the vitellogenin gene *vit-3* (Dowen et al., 2016), and produce fewer offspring (Jones et al., 2009; Soukas et al., 2009). While many cellular mechanisms regulate brood size, one is the number of cells in the early adult progenitor zone (PZ) (Agarwal et al., 2018; Kocsisova et al., 2019), the pool of stem and progenitor cells in the distal part of the hermaphrodite gonad (reviewed in(Hubbard and Schedl, 2019)).

Our results show that *rict-1* mutant adults have a reduced brood size and half the normal pool of cells in the PZ. As in other phenotypic contexts, *rict-1* acts via *sgk-1*, but does not act via downstream IIS or TGFß pathways that are implicated in *rict-1* control of dauer entry (O'Donnell et al., 2018) and in larval PZ expansion (Dalfo et al., 2012; Michaelson et al., 2010; Pekar et al., 2017). We show that intestinal RICT-1 is sufficient and necessary for full establishment of the adult PZ. Our results also indicate that during the L4 larval stage, *rict-1* regulates expression of several members of the conserved ApolipoproteinB-like vitellogenins (Perez and Lehner, 2019)*,* and that *vit-3* contributes to expansion of the progenitor pool, likely downstream of *rict-1*.

RESULTS

RICT-1 is required for normal germline development

To explore effects of TORC2 on the germ line, we examined the effects of reduced *rict-1.* We assessed phenotypes in two available *rict-1* mutants (*ft7* in the third exon and *mg360* in the 11th exon (Jones et al., 2009; Soukas et al., 2009)) and in 6 additional alleles that we generated using CRISPR-Cas9 genome editing. Among three different isoforms (isoform A and B encode identical proteins), *ft7* only affects isoforms A+B, while *mg360* affects all three isoforms (Figure 1A). We generated four CRISPR mutants with Stop-In cassettes (Wang et al., 2018) inserted in the regions of each of the SNPs: *na112* and *na113* in the third exon, and *na114* and *na115* in the 11th exon. We generated two additional mutants: one that deletes 241bp of the 5'UTR and 29bp of the first exon (*na116*), and one that deletes 520bp in total across the first three exons (*na119*). We used this strategy so as not to interfere with the PQN-32 locus that resides within the intron between exons 10 and 11. Strains bearing all these mutations, as well as others used in this study are listed in Table S1.

To gauge the effects of *rict-1* on larval accumulation of the stem/progenitor pool, we counted the number of nuclei per distal gonad arm in the distal progenitor zone (PZ) at the adult molt. We chose to assess the progenitor pool at the molt as this represents a developmental timepoint independent of growth rate and at the start of the early adult steady-state (reviewed in(Hubbard and Schedl, 2019)). All eight *rict-1* mutants displayed a similar significant reduction in the number of PZ nuclei, only reaching half of the wild-type level at the L4 to adult molt (average 107 ± 15.4 across all eight alleles versus 202 ± 20.4 in the wild type) (Figure 1B). We further determined the mitotic index and found that, in contrast to *rsks-1* S6 kinase mutants (Korta et al., 2012), *rict-1* mutants display the same mitotic index as wild-type worms (p=0.4) (Figure 1C). Representative images of the wild-type PZ and the *rict-1(na113)* mutant are shown in Figures 1D and 1E, respectively. We do not see differences when comparing anterior versus posterior position gonad arms (Figure S1B). Given that all *rict-1* mutants behaved similarly, our subsequent studies focused on the *na113* allele.

The brood size in mated worms is correlated with the number of PZ nuclei (Agarwal et al., 2018; Kocsisova et al., 2019). To determine whether the PZ pool reduction we observed in the *rict-1* mutants is associated with reproductive defects, we assessed live broods and, consistent with previous results (Jones et al., 2009; Soukas et al., 2009), we observed a significantly smaller number of hatched offspring for both *rict-1* alleles tested (199 ± 9.21 for *rict-1(ft7)* and 200 ± 3.82 for *rict-1(na113)*) compared with the wild type (312.8 ± 6.77)(Figure 1F,G). We also observed that *rict-1* mutants display an extended reproductive period of ~9 days compared with ~6 days in the wild type (8.95 ± 0.41 days for *na113*, 9.2 ± 0.49 for *ft7*, and 6.35 ± 1.39 for the wild type; Figure 1G,H). Since hermaphrodite brood size is sperm limited (Hodgkin and Barnes, 1991), and the correlation with the PZ can only be assessed in the presence of replete sperm, we assessed live broods in mated worms. We found that, unlike the wild type, where male sperm provision significantly increased offspring number, the number of hatched offspring is not increased in mated *rict-1* mutants (Figure S1A). Furthermore, *rict-1* mutants appear more susceptible to damage inflicted by mating, as the mated hermaphrodite *rict-1* mutant

128 survival was negatively impacted by mating, compared with unmated *rict-1* mutants (data
129 not shown). Despite the extended reproductive period, the overall sum of hatched brood
130 of *rict-1* mutants is still reduced in comparison to wild type (Figure 1F). We conclude that
131 RICT-1 promotes normal germline development, and we focused on the PZ pool as this
132 is a primary early aspect of germline development.
133
134
135 **SGK-1 acts downstream of RICT-1 to establish the early adult PZ pool**
136
We next assessed known pathway components that downstream of RICT-1 and
RICTOR. Canonically, Rictor phosphorylates SGK (reviewed in (Blackwell et al., 2019;
Jang et al., 2022)).
In *C. elegans*, SGK-1 acts genetically downstream of Rictor to regulate fat storage,
body size, developmental rate (Jones et al., 2009; Soukas et al., 2009), lifespan (Soukas
et al., 2009), mitochondrial homeostasis(Gatsi et al., 2014), autophagy (Aspernig et al.,
2019), *skn-1* embryonic phenotypes (Ruf et al., 2013), and the transcription of *vit-3*
(Dowen et al., 2016). To assayed for non-additive genetic interactions of *rict-1* and *sgk-1*
in PZ establishment, we compared PZ counts in the null single and double mutants. We
found that the number of PZ nuclei is similar in *rict-1* or *sgk-1* single mutants (108.76 $\pm$
3.17 and 104.54 $\pm$ 2.15 for *rict-1(na113)* and *sgk-1(ok538)*, respectively) and in the *rict-*
1; sgk-1 double mutant (94.74 $\pm$ 2.15). These results suggesting that, similar to other *rict-*
1 phenotypes, *rict-1* and *sgk-1* are likely acting in a linear relationship for this phenotype
(Figure 2). While PZ counts in the *rict-1*;*sgk-1* double mutant do not show an additive
phenotype in comparison to each single mutant alone, the number of PZ nuclei in the
double mutant is significantly lower than in the *rict-1* mutant alone (p<0.05). This result
suggests that *sgk-1* may have additional minor effects on this phenotype independent of
rict-1 (Figure 2).
The *sgk-1* gain of function allele (*ft15*) (Jones et al., 2009) suppresses some *rict-1*
mutant phenotypes (such as associated learning (Sakai et al., 2017), fat storage, body
size, and developmental delay (Jones et al., 2009)). For both autophagy and *skn-1*
interactions, *sgk-1(gf)* suppression of *rict-1* mutant phenotypes is partial (Aspernig et al.,
2019; Ruf et al., 2013).To determine whether the *sgk-1(ft15) gain-of-function* mutant
suppresses *rict-1,* either entirely or partially, we examined *sgk-1(ft15)* alone and in
combination with *rict-1(na113).* We found that *sgk-1(ft15)* alone displays a PZ count
similar to the wild type (178.74 $\pm$ 5.1 versus 189.76 $\pm$ 4.06). As expected, if *sgk-1* were
acting downstream of *rict-1,* we found a significant, though partial, suppression of the *rict-*
1 PZ phenotype (to 123.06 $\pm$ 3.48 PZ nuclei). Taking the loss- and gain-of-function results
together, we conclude that SGK-1 likely acts downstream of RICT-1, even though the two
may have other minor independent outputs that affect the PZ phenotype.
167
Intestinal RICT-1 contributes to PZ pool establishment
In several different phenotypic contexts, *rict-1* acts from the intestine. These
contexts include dauer entry and foraging behavior (O'Donnell et al., 2018), associative
learning (Sakai et al., 2017), fat accumulation (Jones et al., 2009; Soukas et al., 2009),
and food seeking (McLachlan et al., 2022). Single cell RNA-seq analyses show that *rict-*
1 mRNA is widely expressed but is most highly abundant in the intestine, (from (Cao et

al., 2017; Hutter and Suh, 2016), displayed in Figure S2A). We therefore wondered
whether intestinal RICT-1 might influence the germline PZ, even though TORC1 and
downstream components did not appear to affect the germ line from the soma (Korta et
al., 2012; Roy et al., 2018).
To determine whether intestinal *rict-1(+)* is sufficient to promote accumulation of
the larval PZ pool, we expressed *rict-1(+)* from an intestinal promoter, *ges-1,* on an
extrachromosomal array in the *rict-1(na113)* background. We found that the array-bearing
siblings contain a significantly greater number of PZ nuclei (142.74 $\pm$ 3.94) than their non-
array-bearing siblings (94.97 $\pm$ 3.24). In addition, there is no difference between the PZ
counts of non-array bearing siblings that have maternal but not zygotic *rict-1* with
homozygous *rict-1(na113)* that have neither maternal nor zygotic *rict-1* (Tukey
comparison p=0.533). This result supports the conclusion that *rict-1* does not regulate the
establishment of the PZ through maternal effects (Figure 3A).
To assess whether PZ pool establishment requires intestinal RICT-1, we depleted
RICT-1 in the intestine using tissue-specific auxin-mediated protein degradation (Zhang
et al., 2015). We first generated an sAID::RICT-1 fusion (*rict-1(na117)*) by CRISPR-Cas9
genome editing, and determined that it does not impact protein function, as the number
of PZ nuclei is not significantly different from that of wild-type worms (Figure S2B). We
then crossed it into a strain in which TIR1 is driven by the intestine-specific promoter *ges-*
1p. We observed that in the presence of auxin the number of PZ nuclei is significantly
reduced (153.26 $\pm$ 3.23 with auxin vs. 189.71 $\pm$ 3.21 without auxin). Relative to the vehicle
(EtOH) treatment, auxin did not affect the PZ counts of the other tested strains tested
(wild type, *rict-1(na113)*, *rict-1(na117*[sAID::RICT-1]) or *ieSi61*[*ges-1p*::TIR1] alone),
indicating that the lower number of PZ nuclei is due to auxin induced degradation and not
auxin induced stress (Figure 3B).
Taken together, our sufficiency and necessity tests suggest that intestinal RICT-1
activity is a major contributor to the accumulation of the larval PZ pool as assayed at the
adult molt. Interestingly, intestinally expressed *rict-1* does not entirely rescue to wild-type
levels, nor does intestinal degradation of RICT-1 lower the number of PZ nuclei to the
level of the *rict-1(na113)* (Figure 3B). These results indicate that RICT-1 acts in additional
tissues as well. Single cell RNA-seq analysis (Figure S2A, data from (Cao et al., 2017;
Ghaddar et al., 2023; Hutter and Suh, 2016)) indicates that in addition to the intestine,
larval *rict-1* is expressed in other tissues, including the germ line. Given our prior results
indicating a role for TOR in the germ line (Korta et al., 2012) we tested RICT-1
degradation using *ieSi38*[*sun-1p*::TIR1] and found that it also contributes, though less
than the intestine (Figure S2C). We conclude that the intestine is the primary contributing
source of RICT-1 important for establishment of the early adult germline PZ, with
additional partial contribution from the germ line.
212
213 **Neither DAF-7/TGF-ß nor DAF-2/IIS pathways mediate the effects of *rict-1* on the**
214 **germ line**
215
In the phenotypic context of dauer entry, intestinal *rict-1* signals via neuronal DAF-
28 Insulin and DAF-7 TGF-ß from ASI and ASJ (O'Donnell et al., 2018). This finding was
of particular interest to us since we previously characterized roles for both the Insulin and
TGF-ß pathways in establishing the PZ pool. In both pathways, these same neurons are

implicated: *ins-3* is expressed in ASI and ASJ (Ghaddar et al., 2023; Michaelson et al.,
2010) and was implicated upstream of DAF-2 for PZ phenotypes, and ASI was implicated
by cell ablation for the DAF-7 pathway regulation of the germ line stem cell niche (Dalfo
et al., 2012). In both cases, however, the tissue requirements for the receptors and/or
downstream effectors were different, and effectors acted independently of neurons and
therefore different from their roles in dauer entry (Dalfo et al., 2012; Michaelson et al.,
2010; Pekar et al., 2017). Moreover, in our prior studies, the PZ defect seen upon reduced
daf-2 or *daf-7* was fully suppressible by loss of *daf-16* or *daf-3*, respectively.

If the relevant IIS and/or TGFß pathways mediate the effects of *rict-1*, we would
expect *daf-16* and/or *daf-3* to suppress the *rict-1* mutant PZ defect. To test this
hypothesis, we built double and triple mutants of *rict-1(na113)* with *daf-16(mu86)* and *daf-*
3(e1376) and counted the PZ pool. We found that neither *daf-16(mu86)* nor *daf-3(e1376)*
alone strongly suppressed the PZ defect in the *rict-1* mutant. The PZ of the *daf-16(mu86);*
rict-1(na113) averaged 119.09 ± 2.91 (compared with 97.57 ± 2.73 for *rict-1(na113)*
alone. While this small difference is statistically significant, we note that *daf-16(mu86)* has
a greater (though not statistically significant) average number of PZ nuclei than the wild
type (Figure 4). Most important, the triple mutant *daf-16(mu86); rict-1(na113); daf-*
3(e1376) averages the same number of PZ nuclei (107.27 ± 2.89) as either double mutant
or the *rict-1(na113)* mutant alone (Figure 4). We conclude from these results that neither
daf-16 nor *daf-3* are negatively regulated downstream of RICT-1 as it impacts PZ
accumulation.
241
242 **Transcripts of four vitellogenin genes are depleted in *rict-1* relative to wild type at**
243 **the mid-L4**
244
To determine how *rict-1* may be influencing the larval germ line, we performed
comparative bulk RNA-sequencing of tightly synchronized mid-L4 stage *rict-1* mutants
and wild-type worms in biological triplicate. We detected 14,153 protein coding transcripts
(see Methods). Out of these, 154 are differentially regulated (adjusted p < 0.05 and log2
fold change >2) between the wild type and the *rict-1(na113)* mutant (110 less and 44
more abundant in the *rict-1(na113)* mutant than in the wild type). Applying a more
stringent adjusted p value cut off of p<0.001 (log2 fold change >2) we detect 149
differentially regulated transcripts, out of which 106 are more and 43 are less abundant
in the *rict-1(na113)* mutant than in the wild type. (Figure 5). Our results indicate that
genotype accounts for strongest variance, and that the replicates identify similar sets of
differentially abundant transcripts (Figure 5A, B). Complete transcript counts and statistics
are in Table S3.

Of particular interest were genes expressed in the intestine, and, among these,
vitellogenin genes caught our attention for several reasons*.* The *C. elegans* genome
encodes six vitellogenins, members of a family of conserved yolk proteins (Perez and
Lehner, 2019) best described for their adult function in provisioning oocytes (Grant and
Hirsh, 1999; Kimble and Sharrock, 1983). However, no function in larval stages or the
larval germ line is yet described. We found that four of the six vitellogenins (*vit-1, -3, -4,*
and *-5*) are significantly lower in mid-L4 larval *rict-1* mutants relative to wild type (p<0.001,
log2 fold change < -2) (Figure 5C,D)
265

266
VIT-3 promotes germline progenitor pool establishment
268
Among the vitellogenin genes we detected in our comparative RNA-seq analysis,
rict-1 was previously found to indirectly regulate *vit-3* transcription in the adult intestine
(Dowen, 2019; Dowen et al., 2016). We therefore tested whether *vit-3* could, like *rict-1,*
contribute to accumulation of the larval PZ pool. To this end, we counted the number of
PZ nuclei in worms bearing a loss-of-function *vit-3* mutation. If VIT-3 were important for
PZ establishment we would expect that the PZ pool would be depleted relative to wild
type. We observed that the *vit-3(ok2348)* null mutant contains significantly fewer PZ
nuclei (130.45 $\pm$ 2.3) than the wild type (202.93 $\pm$ 5.35) (Figure 6). If *rict-1* and *vit-3* were
acting in a linear genetic pathway, we would expect that loss of both *rict-1* and *vit-3* would
not further deplete the PZ pool beyond either single mutant alone. We found that the
double *rict-1; vit-3* mutant has the same number of PZ nuclei as the *rict-1(na113)* mutant
alone (108.7 $\pm$ 3.4 vs 99.5 $\pm$ 3, respectively). Together, these results reveal an unexpected
role for *vit-3* in the larval accumulation of germline progenitors, likely downstream of *rict-*
1.
283

284 **DISCUSSION**
285
We generated seven new alleles of *rict-1*, including six similarly behaving alleles
encoding small deletions or early stop codons, and one encoding a fully functional AID-
tagged RICT-1 protein. Using these and prior existing SNP alleles, we show that *rict-1* is
required for several aspects of reproduction, including timing of progeny production. We
focused on a larval requirement for *rict-1* in germline development: the accumulation of
the germline progenitor cells that establishes the early adult steady-state germline
progenitor pool. We determined that the *rict-1* mutant PZ pool reaches only half the
normal number of cells at the adult molt. We observed a similar phenotype upon loss of
sgk-1, a canonical effector downstream of *rict-1*, with no additional defect in the double
mutant and partial but significant suppression of *rict-1* by an *sgk-1(gf)* allele. We also
discovered a prominent role for intestinal *rict-1* in larval establishment of the PZ that is
both sufficient and necessary for establishment of the full complement of wild-type PZ
cells. Despite other phenotypic contexts where *rict-1* depends on pathways previously
implicated in PZ pool regulation, we find no obvious genetic interaction with the insulin or
DAF-7/TGFß pathways. Finally, we identify four vitellogenin genes among transcripts that
are highly reduced in *rict-1* L4 larvae relative to wild type, and demonstrate a role for one
of them, *vit-3*, in promoting the progenitor pool, likely downstream of *rict-1.*
Using multiple mutant alleles of *rict-1*, we confirmed low broods reported by others
(Jones et al., 2009; Soukas et al., 2009), and we also observe an extended timeline of
progeny production and no increase of progeny production upon replete sperm provision.
In mated worms, brood size correlates with the number of cells in the progenitor zone
(PZ) (Agarwal et al., 2018; Kocsisova et al., 2019). Multiple *rict-1* mutants affecting
different areas of the protein all displayed similarly reduced numbers of PZ cells (~half of
the wild type number). These alleles alter coding sequences near the N-terminus (*na116*
and *na119*), the third exon (*ft7*, *na112*, and *na113*) and the 11[th] exon (*mg360*, *na114*, and

na115). The N-terminus and third exon mutants disrupt isoform A and B (Figure 1C), but
nevertheless display the same PZ phenotype as those mutants that affect all three
isoforms. This suggests that the C isoform is unlikely to play a role in the establishment
of the PZ. We also found that the mitotic index of PZ cells in *rict-1* mutants did not differ
from the wild type (Figure 1D), indicating that *rict-1* is unlikely affecting the establishment
of the PZ by markedly slowing cell cycle progression. Exactly how loss of *rict-1* affects
the PZ pool at the cellular level, and whether it affects germline stem cell fate is of future
interest.

Our results suggest that, as in other phenotypic contexts, RICT-1 acts upstream
of SGK-1 to regulate the PZ. Similar to reports for *rict-1* phenotypes affecting fat storage,
body size, developmental rate (Jones et al., 2009; Soukas et al., 2009), lifespan (Soukas
et al., 2009), mitochondrial homeostasis (Gatsi et al., 2014), autophagy (Aspernig et al.,
2019), and *skn-1* in embryonic development (Ruf et al., 2013), we see no additive effect
of a *rict-1; sgk-1* double mutant over either single mutant. We also tested a *sgk-1 (gf)*
allele (Jones et al., 2009), that suppresses some *rict-1* phenotypes entirely (such as those
associated with learning (Sakai et al., 2017), fat storage, body size, and developmental
delay (Jones et al., 2009). Significant, though partial, suppression of *rict-1* mutant
phenotypes were also reported for with phenotypes associated with autophagy (Aspernig
et al., 2019) and *skn-1* interactions (Ruf et al., 2013). As in the latter cases, we observed
significant, though partial, suppression of the *rict-1* mutant PZ defect. These results
suggest that *rict-1* may have additional downstream outputs onto relevant effectors for its
role in establishing the PZ pool.

Our results show that intestinal RICT-1 is both sufficient and necessary for full
establishment of the PZ pool. RICT-1 intestinal sufficiency has been demonstrated for
other phenotypes including entry to the dauer stage, foraging behavior (O'Donnell et al.,
2018), associative learning (Sakai et al., 2017), fat accumulation (Jones et al., 2009;
Soukas et al., 2009), and food seeking (McLachlan et al., 2022)). Tagging RICT-1 at its
endogenous locus with AID, we used the auxin degradation system to also assay the
requirement for intestinally produced *rict-1* in an otherwise wild-type background (Figure
3B). Because auxin exposure prolongs lifespan (Loose and Ghazi, 2021) and promotes
resistance to endoplasmic reticulum stress (Bhoi et al., 2021), we used a minimal
concentration of auxin (1mM) that did not enhance the PZ defect in the *rict-1* mutant nor
in other control strains. Our results indicate that the intestinal rescue of *rict-1(+)* in the *rict-*
1(na113) background does not rescue the number of PZ nuclei to the wild-type level, and
that the intestinal degradation does also not reduce the number PZ nuclei from the wild-
type level to the *rict-1(na113)* level. Results from scRNA-seq of larvae (Cao et al., 2017;
Hutter and Suh, 2016) indicate that *rict-1* expression is broad but is highest in the
intestine, followed by the germ line (Figure S2A), where we also see a partial requirement
(Figure S2C). Whether the intestine and germ line are the only required tissues, and how
the germline-autonomous role interacts with TORC1 function are areas for further
investigation.

In the phenotypic context of dauer entry, RICT-1 acts via both the DAF-28/Insulin
and the DAF-7/TGF-ß in the ASI and ASJ neurons (O'Donnell et al., 2018). However, we
observed that neither the insulin nor TGFß signaling pathways, both of which are
implicated in larval PZ pool expansion (Dalfo et al., 2012; Michaelson et al., 2010; Pekar
et al., 2017), act downstream of RICT-1 to promote PZ pool expansion.

The results of our RNA-seq and genetic analysis suggest that *rict-1* in L4 larvae
transcriptionally regulates at least four of the six vitellogenin genes, and that a *vit-3* mutant
has a PZ defect similar to but not as severe as loss of *rict-1*. No function has been
previously ascribed for vitellogenins in post embryonic development nor in the germ line
prior to adult oocyte provision (reviewed in (Perez and Lehner, 2019)). RICT-1 indirectly
promotes high levels of *vit-3* transcription in the adult intestine (Dowen et al., 2016). In
this scenario, adult transcriptional activation of *vit-3* occurs in response to heterochronic
cues and is controlled by CEH-60, a protein that can also act as a transcriptional repressor
of stress response genes, suggesting a tradeoff between larval stress response and adult
reproduction (Dowen, 2019; Dowen et al., 2016).
Nevertheless, our results point to a possibly different role for vitellogenins, prior to
oocyte provisioning. In combination with the high fat phenotype of the *rict-1* mutant (Jones
et al., 2009; Soukas et al., 2009; Yen et al., 2010) we speculate that RICT-1 regulates the
establishment of the PZ via intestine-germline transport of lipids or other molecules
associated with vitellogenin-containing lipid particles. Vitellogenins are secreted from the
adult intestine into the body cavity (Kimble and Sharrock, 1983), pass through gonad
sheath pores (Hall et al., 1999), and are taken up by developing oocytes via receptor-
mediated endocytosis (Grant and Hirsh, 1999). Molecules associated with vitellogenins
could act as a nutrient source for post-embryonic survival (Perez and Lehner, 2019). The
expression of all six vitellogenins is negligible during the first three larval stages in
comparison to the rapidly increasing expression during the fourth larval stage and
adulthood (Cao et al., 2017; Ghaddar et al., 2023; Hutter and Suh, 2016), a time-frame
that correlates with the rapid expansion of the PZ pool (Killian and Hubbard, 2005). We
note that although loss of *vit-3* causes a mutant phenotype is somewhat less severe than
rict-1, this result suggests that other *vits* from our RNA-seq analysis may contribute to PZ
expansion.
What might vitellogenins provide to the larval germ line? One possibility is that they
facilitate transport of a critical nutrient lipids, such as cholesterol. Another possibility is
that they provide limiting membrane lipids required for the demands of germ cell division.
Yet another possibility is that they associate with as-yet unknown moieties such as
hormones to direct germline development. The oocyte receptor for vitellogenins, RME-2,
is not detectable in the distal gonad during larval stages (Grant and Hirsh, 1999; Salinas
et al., 2007). However, a function for RME-2 as a high affinity receptor prior to oogenesis
has not been formally ruled out. Alternatively, a different receptor may fulfill this role in
larvae. Additional studies are required to determine how RICT-1 regulates larval
vitellogenins and how vitellogenins participate in the establishment of the PZ pool.
In conclusion, we present the first evidence that the TORC2 component RICT-1
contributes to establishment of the full germline stem/progenitor pool. Our results suggest
a model whereby intestinal RICT-1 promotes the expression of vitellogenins that support
germline expansion during larval stages. Due to the high conservation of TORC2
components, vitellogenins, and their receptors, our results may have general implications
for how Rictor and TORC2 regulate germline development in other organisms and, more
generally, how they may support pools of proliferating stem or progenitor cells in other
organisms and organ contexts. Moreover, since vitellogenin-like molecules, such as
ApoB, are implicated in heart disease and other aspects of lipid metabolism, the *C.*

elegans larval germ line may provide in roads to the identification of additional associated
moieties and receptors.

ACKNOWLEDGMENTS

We thank Nicole Hall for help and advice with the RNA extraction protocol and analysis, Amy Webster for feedback on the RNA-seq analysis, Sophie Wasel and Mia Sinks for technical support, and Michael Cammer for microscopy support. We are also grateful for feedback and support from Jeremy Nance and Niels Ringstad, and members of their labs. The Microscopy and Genome Technology cores are partially supported by the Cancer Center Support Grant P30CA016087 at the Laura and Isaac Perlmutter Cancer Center. The computational requirements for this work were supported in part by the NYU Langone High Performance Computing (HPC) Core's resources and personnel. Thanks to Michael O'Donnell for providing strains; some strains were also provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported by NYSTEM Institutional Training Grant Contract #C322560GG and NIH R35GM134876 to E.J.A.H.

MATERIALS AND METHODS

Worm maintenance and husbandry

C. elegans nematodes were grown on NGM solid media at 20°C and fed *Escherichia coli* strain OP50 (Stiernagle, 2006). All strains generated and used in this study are listed with full genotypes in Table S1. Some strains were provided by the CGC. All strains were based on Bristol N2 (RRID:WBSTRAIN:WBStrain00000001). Many experiments required access to information on WormBase (Sternberg et al., 2024).

Determination of progenitor zone nuclei counts and mitotic index

Mid-L4 worms (L4.5; (Mok et al., 2015)) were picked from mixed stage plates and transferred to new plates. 6h later all early adult worms bearing a vulval slit were picked into 1ml of M9+Tween within a 1.5ml of low-retention microfuge tube (02-681-320, Fisherbrand). Worms were then spun down on a table-top centrifuge for 10 seconds, before the supernatant was carefully removed with a pipette under the dissecting scope, to avoid loss of worms. Another 500µl of M9+Tween was added, followed by a centrifugation and supernatant removal. 50µl of 100%EtOH were added to each sample, and worms were allowed to settle at the bottom for ~10min. After 10min as much EtOH was removed as possible without disturbing the worms. Finally, ~5µl of Vectashield+DAPI (H-1000, Vector Laboratories) was added, and worms were gently pipetted onto a 4% agar pad on a glass slide. Slides were kept in the cold and dark and were imaged within 2 weeks after staining. Imaging was performed on Nikon W1 spinning disk confocal microscope with 60x oil immersion magnification with z-stacks of 0.5µm step size. Only one gonad arm was counted per individual worm. The collected files were counted with ImageJ(Arganda-Carreras et al. 2024) to manually count the number of PZ nuclei. Mitotic

444 index presented as a percentage of the number of mitotic figures (identified as
445 characteristic metaphase and anaphase figures) divided by the total number of PZ nuclei.

Brood Size Determination

447 L4 worms were picked from mixed stage plates. Single hermaphrodites were added to
448 plates and, for the mated treatment, were joined by 3 wild-type males, which were
449 replaced every two days. All worms were transferred every day to new plates, and
450 hatched offspring were counted 2 days later.

CRISPR-Cas9 genome editing

CRISPR/Cas9 genome editing was performed using pre-incubated Cas9
(Berkeley)::(crRNA +tracrRNA) (IDT) ribonucleoprotein, and screening was facilitated by
co-CRISPR detection using the *dpy-10(cn64)* mutation as described (Paix et al., 2016).
DNA repair templates contained ~25–35 bps of homology on each arm as ssDNA oligos
(<150 bps) (IDT). The guide RNAs and oligos are listed in Table S4. For Stop-In mutants
(*na112, na113, na114*, and *na115*) the inserted stop cassette led to an early stop in all
frames (Wang et al., 2018) .

RNA extraction for bulk RNA-seq

Worms were bleached and synchronized as previously described (Stiernagle, 2006).
Synchronized L1 larvae were added to 15 10cm NGM plates (each inoculated with 1ml
of OP50) at different times, so that all strains would reach the mid-L4 stage at the same
time (as previously empirically determined). Worms were washed off the plates with
RNAse-free water (10977015, Invitrogen) and collected atop pluristrainers with 40μl mesh
size (#43-10040-40 from Pluristrainer.com) and washed three times with 5ml of RNAse
free H_2O. Pluristrainers were turned over, and 1ml of RNAse-free water was used to wash
the worms off the mesh and collect them in RNAse-free 2ml Eppendorf microfuge tubes
(E0030123620, Eppendorf). Each 1ml of packed worms was resuspended in 4ml of Trizol.
The worm+Trizol mix was vortexed and then carefully dropped into liquid nitrogen and
then thawed at 37ºC. After three freeze-thaw cycles, samples were thawed at 37ºC, and
2ml of Trizol+2ml of Chloroform/packed ml of worms was added and worms and shaken
by hand. Samples were aliquoted into 2ml RNAse-free tubes and incubated at room
temperature for three minutes. Samples were then spun at 12,000g at 4ºC for 15 min.
The aqueous layer was removed into new RNAse-free tubes, and an equal volume of
70% ethanol was added and inverted to mix. This solution was transferred to a spin
column (RNeasy Qiagen kit # 74104) and was spun down at 21k rpf for 30sec. The
column was washed once with buffer RW1 and centrifuged for 15 seconds at 21k rpf
DNase I was added directly to each RNeasy column membrane and was left to incubate
at room temperature for 15 minutes. Each column was washed with Qiagen RW1 and
RPE buffers, before the sample was eluted from the spin column in RNase-free water to
be collected in a fresh 2ml RNase-free tube. RNA quality was initially assessed with a
nanodrop (NanoDrop 8000 UV-Vis Spectrophotometer). When samples were submitted
to the NYU genome technology core. The automated RNA-seq library preparation and
polyA selection was carried out before samples were analyzed with the NovaSeq X.

RNA-seq analysis

Version WBCel235 of the *C. elegans* genome was used to map sequencing reads. Seq-N-Slide was used to align the sequencing results and determine those gene transcripts that are differentially detected between the mutant and wild-type strains (Dolgalev). Transcript counts were restricted to protein coding genes (19,983 genes). Prior to further analysis this count data was further restricted to include only genes with counts-per-million >1 in at least two out of the six libraries (14,153 genes). Principal component analysis (PCA) was performed on log2 normalized CPM values on all genes included in the differential gene expression analysis (14,153 genes), using the stats package in R (version 2024.04.2+764) (RStudioTeam, 2020). Lists of genes with differentially abundant transcripts are in Table S3, along with summary statistics. The entire dataset will be publicly available upon publication at NCBI Gene Expression Omnibus.

Statistical analyses

Plots in Figures 1-6 and Figures S1-S2 were generated using R with minor modifications using Adobe Illustrator. Statistical analysis was conducted in R (version 2024.04.2+764) (RStudioTeam, 2020). All PZ counts and the offspring data were analyzed with a Generalized Mixed Effect Model (Nelder and Wedderburn, 1972), taking the replicate as a random factor into account. The Mitotic Index data was analyzed using a Linear Model (Nelder and Wedderburn, 1972), and the Reproductive Period data was analyzed with a Linear Mixed Effects Model (Nelder and Wedderburn, 1972).

REFERENCES CITED

Agarwal, I., Farnow, C., Jiang, J., Kim, K. S., Leet, D. E., Solomon, R. Z., Hale, V. A. and
Goutte, C. (2018). HOP-1 Presenilin Deficiency Causes a Late-Onset Notch Signaling
Phenotype That Affects Adult Germline Function in Caenorhabditis elegans. *Genetics*
208, 745-762.

Aspernig, H., Heimbucher, T., Qi, W., Gangurde, D., Curic, S., Yan, Y., Donner von
Gromoff, E., Baumeister, R. and Thien, A. (2019). Mitochondrial Perturbations Couple
mTORC2 to Autophagy in C. elegans. *Cell Rep* **29**, 1399-1409 e1395.

Bhoi, A., Palladino, F. and Fabrizio, P. (2021). Auxin confers protection against ER stress in
Caenorhabditis elegans. *Biol Open* **10**.

Blackwell, T. K., Sewell, A. K., Wu, Z. and Han, M. (2019). TOR Signaling in Caenorhabditis
elegans Development, Metabolism, and Aging. *Genetics* **213**, 329-360.

Cao, J., Packer, J. S., Ramani, V., Cusanovich, D. A., Huynh, C., Daza, R., Qiu, X., Lee, C.,
Furlan, S. N., Steemers, F. J., et al. (2017). Comprehensive single-cell transcriptional
profiling of a multicellular organism. *Science* **357**, 661-667.

Caron, A., Richard, D. and Laplante, M. (2015). The Roles of mTOR Complexes in Lipid
Metabolism. *Annu Rev Nutr* **35**, 321-348.

Dalfo, D., Michaelson, D. and Hubbard, E. J. (2012). Sensory regulation of the C. elegans
germline through TGF-beta-dependent signaling in the niche. *Curr Biol* **22**, 712-719.

Dolgalev, I.

Dowen, R. H. (2019). CEH-60/PBX and UNC-62/MEIS Coordinate a Metabolic Switch that
Supports Reproduction in C. elegans. *Dev Cell* **49**, 235-250 e237.

Dowen, R. H., Breen, P. C., Tullius, T., Conery, A. L. and Ruvkun, G. (2016). A microRNA
program in the C. elegans hypodermis couples to intestinal mTORC2/PQM-1 signaling to
modulate fat transport. *Genes Dev* **30**, 1515-1528.

Frei, I. C., Weissenberger, D., Muller, C., Hall, M. N. and Shimobayashi, M. (2023). Hepatic
mTORC2 compensates for loss of adipose mTORC2 in mediating energy storage and
glucose homeostasis. *Am J Physiol Endocrinol Metab* **324**, E589-E598.

Gatsi, R., Schulze, B., Rodriguez-Palero, M. J., Hernando-Rodriguez, B., Baumeister, R.
and Artal-Sanz, M. (2014). Prohibitin-mediated lifespan and mitochondrial stress
implicate SGK-1, insulin/IGF and mTORC2 in C. elegans. *PLoS One* **9**, e107671.

Ghaddar, A., Armingol, E., Huynh, C., Gevirtzman, L., Lewis, N. E., Waterston, R. and
O'Rourke, E. J. (2023). Whole-body gene expression atlas of an adult metazoan. *Sci Adv*
9, eadg0506.

Grant, B. and Hirsh, D. (1999). Receptor-mediated endocytosis in the Caenorhabditis elegans
oocyte. *Mol Biol Cell* **10**, 4311-4326.

Guertin, D. A., Stevens, D. M., Thoreen, C. C., Burds, A. A., Kalaany, N. Y., Moffat, J.,
Brown, M., Fitzgerald, K. J. and Sabatini, D. M. (2006). Ablation in mice of the
mTORC components raptor, rictor, or mLST8 reveals that mTORC2 is required for
signaling to Akt-FOXO and PKCalpha, but not S6K1. *Dev Cell* **11**, 859-871.

Guri, Y., Colombi, M., Dazert, E., Hindupur, S. K., Roszik, J., Moes, S., Jenoe, P., Heim,
M. H., Riezman, I., Riezman, H., et al. (2017). mTORC2 Promotes Tumorigenesis via
Lipid Synthesis. *Cancer Cell* **32**, 807-823 e812.

Hall, D. H., Winfrey, V. P., Blaeuer, G., Hoffman, L. H., Furuta, T., Rose, K. L., Hobert, O.
and Greenstein, D. (1999). Ultrastructural features of the adult hermaphrodite gonad of

Caenorhabditis elegans: relations between the germ line and soma. *Dev Biol* **212**, 101-
123.

Hietakangas, V. and Cohen, S. M. (2007). Re-evaluating AKT regulation: role of TOR
complex 2 in tissue growth. *Genes Dev* **21**, 632-637.

Hodgkin, J. and Barnes, T. M. (1991). More is not better: brood size and population growth in
a self-fertilizing nematode. *Proc Biol Sci* **246**, 19-24.

Hodgkin, J. and Doniach, T. (1997). Natural variation and copulatory plug formation in
Caenorhabditis elegans. *Genetics* **146**, 149-164.

Hubbard, E. J. A. and Schedl, T. (2019). Biology of the Caenorhabditis elegans Germline
Stem Cell System. *Genetics* **213**, 1145-1188.

Hutter, H. and Suh, J. (2016). GExplore 1.4: An expanded web interface for queries on
Caenorhabditis elegans protein and gene function. *Worm* **5**, e1234659.

Jang, H., Park, Y. and Jang, J. (2022). Serum and glucocorticoid-regulated kinase 1: Structure,
biological functions, and its inhibitors. *Front Pharmacol* **13**, 1036844.

Jones, K. T., Greer, E. R., Pearce, D. and Ashrafi, K. (2009). Rictor/TORC2 regulates
Caenorhabditis elegans fat storage, body size, and development through sgk-1. *PLoS Biol*
7, e60.

Killian, D. J. and Hubbard, E. J. (2005). Caenorhabditis elegans germline patterning requires
coordinated development of the somatic gonadal sheath and the germ line. *Dev Biol* **279**,
322-335.

Kim, J. E. and Chen, J. (2004). regulation of peroxisome proliferator-activated receptor-gamma
activity by mammalian target of rapamycin and amino acids in adipogenesis. *Diabetes*
53, 2748-2756.

Kimble, J. and Sharrock, W. J. (1983). Tissue-specific synthesis of yolk proteins in
Caenorhabditis elegans. *Dev Biol* **96**, 189-196.

Kocsisova, Z., Kornfeld, K. and Schedl, T. (2019). Rapid population-wide declines in stem cell
number and activity during reproductive aging in C. elegans. *Development* **146**.

Korta, D. Z., Tuck, S. and Hubbard, E. J. (2012). S6K links cell fate, cell cycle and nutrient
response in C. elegans germline stem/progenitor cells. *Development* **139**, 859-870.

Lin, K., Dorman, J. B., Rodan, A. and Kenyon, C. (1997). daf-16: An HNF-3/forkhead family
member that can function to double the life-span of Caenorhabditis elegans. *Science* **278**,
1319-1322.

Liu, G. Y. and Sabatini, D. M. (2020). mTOR at the nexus of nutrition, growth, ageing and
disease. *Nat Rev Mol Cell Biol* **21**, 183-203.

Loose, J. A. and Ghazi, A. (2021). Auxin treatment increases lifespan in Caenorhabditis
elegans. *Biol Open* **10**.

McLachlan, I. G., Kramer, T. S., Dua, M., DiLoreto, E. M., Gomes, M. A., Dag, U.,
Srinivasan, J. and Flavell, S. W. (2022). Diverse states and stimuli tune olfactory
receptor expression levels to modulate food-seeking behavior. *Elife* **11**.

Meissner, B., Boll, M., Daniel, H. and Baumeister, R. (2004). Deletion of the intestinal
peptide transporter affects insulin and TOR signaling in Caenorhabditis elegans. *J Biol*
Chem **279**, 36739-36745.

Michaelson, D., Korta, D. Z., Capua, Y. and Hubbard, E. J. (2010). Insulin signaling
promotes germline proliferation in C. elegans. *Development* **137**, 671-680.

Mok, D. Z., Sternberg, P. W. and Inoue, T. (2015). Morphologically defined sub-stages of C.
elegans vulval development in the fourth larval stage. *BMC Dev Biol* **15**, 26.

Nelder, J. A. and Wedderburn, R. W. (1972). Generalized Linear Models. *J R Stat Soc Ser a-*
600   *G* **135**, 370-+.

601   **O'Donnell, M. P., Chao, P. H., Kammenga, J. E. and Sengupta, P.** (2018). Rictor/TORC2
602   mediates gut-to-brain signaling in the regulation of phenotypic plasticity in C. elegans.
603   *PLoS Genet* **14**, e1007213.

604   **Paix, A., Schmidt, H. and Seydoux, G.** (2016). Cas9-assisted recombineering in C. elegans:
605   genome editing using in vivo assembly of linear DNAs. *Nucleic Acids Res* **44**, e128.

606   **Pekar, O., Ow, M. C., Hui, K. Y., Noyes, M. B., Hall, S. E. and Hubbard, E. J. A.** (2017).
607   Linking the environment, DAF-7/TGFbeta signaling and LAG-2/DSL ligand expression
608   in the germline stem cell niche. *Development* **144**, 2896-2906.

609   **Perez, M. F. and Lehner, B.** (2019). Vitellogenins - Yolk Gene Function and Regulation in
610   Caenorhabditis elegans. *Front Physiol* **10**, 1067.

611   **Puca, F., Fedele, M., Rasio, D. and Battista, S.** (2022). Role of Diet in Stem and Cancer Stem
612   Cells. *Int J Mol Sci* **23**.

613   **Riddle, D. L., Swanson, M. M. and Albert, P. S.** (1981). Interacting genes in nematode dauer
614   larva formation. *Nature* **290**, 668-671.

615   **Robinson, J. T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G.**
616   **and Mesirov, J. P.** (2011). Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26.

617   **Roy, D., Kahler, D. J., Yun, C. and Hubbard, E. J. A.** (2018). Functional Interactions
618   Between rsks-1/S6K, glp-1/Notch, and Regulators of Caenorhabditis elegans Fertility and
619   Germline Stem Cell Maintenance. *G3 (Bethesda)* **8**, 3293-3309.

620   **Roy, D., Michaelson, D., Hochman, T., Santella, A., Bao, Z., Goldberg, J. D. and Hubbard,**
621   **E. J. A.** (2016). Cell cycle features of C. elegans germline stem/progenitor cells vary
622   temporally and spatially. *Dev Biol* **409**, 261-271.

623   **RStudioTeam** (2020). Integrated Development for R. Boston MA: PBC

624   **Ruf, V., Holzem, C., Peyman, T., Walz, G., Blackwell, T. K. and Neumann-Haefelin, E.**
625   (2013). TORC2 signaling antagonizes SKN-1 to induce C. elegans mesendodermal
626   embryonic development. *Dev Biol* **384**, 214-227.

627   **Russell, R. C., Fang, C. and Guan, K. L.** (2011). An emerging role for TOR signaling in
628   mammalian tissue and stem cell physiology. *Development* **138**, 3343-3356.

629   **Saba, J. A., Liakath-Ali, K., Green, R. and Watt, F. M.** (2021). Translational control of stem
630   cell function. *Nat Rev Mol Cell Biol* **22**, 671-690.

631   **Sakai, N., Ohno, H., Tomioka, M. and Iino, Y.** (2017). The intestinal TORC2 signaling
632   pathway contributes to associative learning in Caenorhabditis elegans. *PLoS One* **12**,
633   e0177900.

634   **Salinas, L. S., Maldonado, E., Macias-Silva, M., Blackwell, T. K. and Navarro, R. E.** (2007).
635   The DEAD box RNA helicase VBH-1 is required for germ cell function in C. elegans.
636   *Genesis* **45**, 533-546.

637   **Soukas, A. A., Kane, E. A., Carr, C. E., Melo, J. A. and Ruvkun, G.** (2009). Rictor/TORC2
638   regulates fat metabolism, feeding, growth, and life span in Caenorhabditis elegans. *Genes*
639   *Dev* **23**, 496-511.

640   **Sternberg, P. W., Van Auken, K., Wang, Q., Wright, A., Yook, K., Zarowiecki, M.,**
641   **Arnaboldi, V., Becerra, A., Brown, S., Cain, S., et al.** (2024). WormBase 2024: status
642   and transitioning to Alliance infrastructure. *Genetics* **227**.

Stiernagle, T. (2006). Maintenance of C. elegans. *WormBook*, 1-11.

Szwed, A., Kim, E. and Jacinto, E. (2021). Regulation and metabolic functions of mTORC1
and mTORC2. *Physiol Rev* **101**, 1371-1426.
Tu, W. B., Christofk, H. R. and Plath, K. (2023). Nutrient regulation of development and cell
fate decisions. *Development* **150**.
Wang, H., Park, H., Liu, J. and Sternberg, P. W. (2018). An Efficient Genome Editing
Strategy To Generate Putative Null Mutants in Caenorhabditis elegans Using
CRISPR/Cas9. *G3 (Bethesda)* **8**, 3607-3616.
Yen, K., Le, T. T., Bansal, A., Narasimhan, S. D., Cheng, J. X. and Tissenbaum, H. A.
(2010). A comparative study of fat storage quantitation in nematode Caenorhabditis
elegans using label and label-free methods. *PLoS One* **5**.
Zhang, L., Ward, J. D., Cheng, Z. and Dernburg, A. F. (2015). The auxin-inducible
degradation (AID) system enables versatile conditional protein depletion in C. elegans.
Development **142**, 4374-4384.
Zoncu, R., Efeyan, A. and Sabatini, D. M. (2011). mTOR: from growth signal integration to
cancer, diabetes and ageing. *Nat Rev Mol Cell Biol* **12**, 21-35.
659
660
661

662
663 **FIGURE LEGENDS**
664

Figure 1: *rict-1* regulates reproduction and the germline progenitor pool. (A) Overview of
the eight different *rict-1* mutants used in this study (two pre-existing SNP mutations in
magenta, four Stop-in mutants in black, and two knock-out mutants in blue). Scale bar
indicates 1kb. Only alleles *mg360*, *na114*, and *na115* affect all isoforms. (B) Number of
PZ nuclei for wild type and eight tested *rict-1* mutants. (C) Mitotic index for wild type and
five tested *rict-1* mutants in comparison to wild type. (D, E) Micrographs of representative
wild type (D) and *rict-1(na113)* mutant (E) progenitor zones. (F) Average of hatched
offspring per adult over a lifetime. (G) Hatched offspring per adult over time; the thick lines
indicate the average per strain, while the thinner lines indicate single worms. Error bars
indicate standard error of the mean. (H) Reproductive period in days. (B, C, F, H) Black
circles indicate the means across independent replicate experiments (shown in different
colors), while the smallest circles indicate gonad arms (n = number of gonad arms
scored). The violin displays the overall distribution of data points. Error bars indicate the
standard error of the mean. Statistical tests are summarized in Table S2. The asterisks
indicate *** p<0.001, ** p<0.01, * p<0.05 of a Linear Mixed Effects Model with a Tukey
Post Hoc Test. Panel B, F, and H were analyzed with a Linear Mixed Model, panel C was
analyzed with a Linear Model, and panel G was analyzed with a Generalized Linear
Model, all followed up with a Tukey Post Hoc test where appropriate.
683

Figure 2: RICT-1 and SGK-1 are likely acting in a linear pathway. Black circles indicate
the means across independent replicate experiments (shown in different colors), while
the smallest circles indicate individual gonad arms (n = number of gonad arms scored).
The violin indicates the overall distribution of data points. Error bars indicate the standard
error of the mean. Statistical tests are summarized in Table S2. The asterisks indicate ***
p<0.001, ** p<0.01, * p<0.05 of a Linear Mixed Effects Model with a Tukey Post Hoc Test.
690

Figure 3: Intestinal RICT-1 is sufficient and necessary for full establishment of the PZ. (A) The intestinal array bearing siblings of *rict-1* (*oyEx713[intestinal rict-1(+)]*) display a significantly higher number of PZ nuclei than their non-array bearing siblings. The non-array bearing siblings are also not significantly different from the non-transgenic *rict-1(na113)* null mutants. (B) In the presence of auxin the Intestinal TIR1 facilitates the degradation of the sAID::RICT-1 (*rict-1(na117)*), while the wildtype, the *rict-1(na113)*, the sAID::RICT-1, and the intestinal TIR1 (*ieSi61*) do not display any sensitivity to auxin. Black circles indicate the means across independent replicate experiments (shown in different colors), while the smallest circles indicate individual gonad arms (n = number of gonad arms scored). The violin indicates the overall distribution of data points. Error bars indicate the standard error of the mean. Statistical tests are summarized in Table S2. The asterisks indicate *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$ of a Linear Mixed Effects Model with a Tukey Post Hoc Test.

Figure 4: Neither the Insulin pathway nor the DAF-7/TGF-ß pathway mediate the effects of RICT-1 on the germ line. Black circles indicate the means across different replicate experiments (shown in different colors), while the smallest circles indicate individual worms (also the number at the bottom of each graph). The violin displays the overall distribution of data points. Error bars indicate the standard error of the mean. Statistical test results are summarized in Table S2. The asterisks indicate *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$ of a Generalized Linear Model with a Tukey Post Hoc Test.

Figure 5: Significant differences in *rict-1* mutant versus wild-type transcript abundance in L4 larvae include four vitellogenin genes. (A) Principal Component Analysis of the two strains, in which each circle indicates one library, color coded by the two different strains. (B) A heatmap in which each line indicates one differentially regulated gene (log2 fold change >2 and adjusted p value < 0.001) between the *rict-1(na113)* mutant and the wild type. Darker colors indicate a stronger differential, where blue colors indicate lower and red higher abundance. (C) A volcano plot displaying genes in (B) as circles in both the *rict-1(na113)* mutant and the wild-type: not significantly regulated (grey), down-regulated in the *rict-1(na113)* mutant (red) or up-regulated in the *rict-1(na113)* mutant (blue) relative to the wild type. Vitellogenin genes are labeled and marked in larger black circles. The dashed line indicates the adjusted p-value threshold of <0.001. (D) The transcription pattern of the 4 differentially regulated vitellogenin genes is displayed in the Integrative Genome Viewer (Robinson et al., 2011). In each case, the top 3 rows are the wild type, and the lower 3 rows are the *rict-1(na113)* mutant; the transcript abundance scale is shown for each gene on the left side of the bottom row.

Figure 6: *rict-1* and *vit-3* are likely acting in a linear pathway. Black circles indicate the means across different replicate experiments (shown in different colors), while the smallest circles indicate individual gonad arms (n = number of gonad arms scored). The violin displays the overall distribution of data points. Error bars indicate the standard error of the mean. Statistical tests are summarized in Table S2. The asterisks indicate *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$ of a Generalized Linear Model with a Tukey Post Hoc Test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6