Genetic Note

A high-quality reference genome and comparative genomics of the widely-farmed banded

cricket (*Gryllodes sigillatus*) identifies selective breeding targets

Shangzhe Zhang[1], Kristin R. Duffield[2,3], Bert Foquet[4,5], Jose L. Ramirez[3], Ben M. Sadd[4], Scott

K. Sakaluk[4], John Hunt[6], Nathan W. Bailey[1,*]

[1]School of Biology, University of St Andrews, St Andrews, Fife KY16 9TH, UK

[2]USDA-ARS, Geospatial and Environmental Epidemiology Research Unit, Mississippi State,

MS, USA

[3]USDA-ARS, National Center for Agricultural Utilization Research, Crop BioProtection

Research Unit, Peoria, IL, USA

[4]School of Biological Sciences, Illinois State University, Normal, IL 61790, USA

[5]McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History,

University of Florida, Gainesville, FL, USA

[6]School of Science, Western Sydney University, Hawkesbury Campus, Penrith NSW 2751,

Australia

* Correspondence to Nathan W. Bailey: nwb3@st-andrews.ac.uk

Keywords: banded cricket | cricket farming | genome-assisted breeding | *Gryllodes sigillatus* |

insect protein

Abstract

Farmed insects have gained attention as an alternative, sustainable source of protein with a lower carbon footprint than traditional livestock. We present a high-quality reference genome for one of the most commonly farmed insects, the banded cricket *Gryllodes sigillatus*. In addition to its agricultural importance, *G. sigillatus* is also a model in behavioural and evolutionary ecology research on reproduction and mating systems. We report comparative genomic analyses that clarify the banded cricket's evolutionary history, identify gene family expansions and contractions unique to this lineage, associate these with agriculturally important traits, and identify targets for genome-assisted breeding efforts. The high-quality *G. sigillatus* genome assembly plus accompanying comparative genomic analyses serve as foundational resources for both applied and basic research on insect farming and behavioural biology, enabling researchers to pinpoint trait-associated genetic variants, unravel functional pathways governing those phenotypes, and accelerate selective breeding efforts to increase the efficacy of large-scale insect farming operations.

Introduction

Global demand for sustainable protein sources has driven interest in alternative agriculture. Such efforts include insect farming, which can meet nutritional needs while significantly reducing environmental impacts (Grabowski et al., 2022; van Huis & Oonincx, 2017). Insects are known for their favourable feed conversion efficiency. They require significantly less land, water, and energy resources compared to traditional livestock, exploit diverse dietary proteins (Morales-Ramos et al., 2020; Kasdorf et al., 2025) and produce fewer greenhouse gases (Lange & Nakamura, 2023). These advantages make them an environmentally friendly option for addressing food security challenges (Huis et al., 2021; van Huis & Oonincx, 2017). Crickets (suborder Ensifera) have been farmed and consumed by humans for millennia, and over 60 species are known to be consumed across 49 countries (Magara et al., 2021). Among these, the banded cricket (*Gryllodes sigillatus*, Figure 1a) has gained considerable attention due to its rapid growth rate, high reproductive capacity, and palatability, making it a preferred species for large-scale farming operations (Kong et al., 2024; Magara et al., 2021).

Despite the growing commercial significance of *G. sigillatus*, publicly available genetic resources for this species are limited. This scarcity of genetic information constrains the potential for selective breeding programs that could enhance desirable traits such as nutritional

value, growth rate, and reproductive efficiency. Recent research on *G. sigillatus* has identified

key environmental factors that influence such traits, including temperature, food stress (Rapkin

et al., 2018; Kasdorf et al., 2025; Muzzatti et al., 2024; Muzzatti et al., 2025a), mating strategies

(Sakaluk et al., 2019) and pests (Muzzatti et al., 2025b). However, genome-based selection has

become central to modern breeding in a wide variety of agricultural species, resulting in rapid

and efficient improvement of specific traits (Johnsson, 2023). Beyond their economic

importance, field crickets – including *G. sigillatus* - have long been a study organism in

evolutionary and behavioural ecology (Bateman & MacFadyen, 1999; Champagnon & Cueva

del Castillo, 2008; Sakaluk et al., 2002; Sakaluk et al., 2019). Males of this polygynandrous

species produce a costly nuptial gift – the spermatophylax – which they transfer to females

upon mating, making *G. sigillatus* a prominent research model for examining mating system

evolution, mate choice, and sexual selection (Ivy & Sakaluk, 2005; Sakaluk, 1984, 1991). A

high-quality genome assembly serves as a foundational resource for such efforts, enabling

researchers to pinpoint trait-associated genetic variants, unravel functional pathways governing

those phenotypes, and accelerate both selective breeding efforts and fundamental research in

behavioural and evolutionary biology (Sinha et al., 2021).

In this study, we produced a high-quality, annotated, chromosome-level reference genome for

G. sigillatus by integrating long-read PacBio HiFi data with high-throughput chromosome

conformation capture (Hi-C) data. Using comparative genomic approaches, we refined the

phylogenetic placement and evolutionary history of *G. sigillatus*. We identified candidate gene

families linked to economically significant traits that underwent changes during the divergence

of *G. sigillatus* from sister taxa and illuminate gene family expansions and contractions

relevant to the unique biology of this species. This research provides a foundational resource

for the genetic analysis and farming of *G. sigillatus*, offering valuable insights into its genomic

features and agriculturally important traits, as well as for basic research on the evolution of

mating systems and reproductive behaviour.

Methods

Sampling and DNA sequencing

For whole genome sequencing, an individual *Gryllodes sigillatus* female was taken from an

inbred laboratory stock line descended from approximately 500 adults collected at Las Cruces,

New Mexico, in 2001 (Ivy & Sakaluk, 2005). We extracted high molecular weight genomic

DNA from liquid nitrogen flash-frozen cricket legs. Following an initial wash with $1\times$ PBS

buffer, DNA was extracted using the Macherey-Nagel™ NucleoBond™ High Molecular
Weight DNA Kit. DNA quality and concentration were assessed with both Nanodrop and Qubit.
Sequencing library preparation and whole genome sequencing were performed using the
PacBio SMRT® system by BGI Genomics, resulting in 96.25 Gb of HiFi reads with a coverage
depth of approximately 48×. Additionally, flash-frozen muscle tissue was used to prepare and
sequence a Hi-C library, also by BGI Genomics, resulting in 151.48 Gb of ca. 72× Hi-C reads
(Table S1).
98
99 **Bulk RNA-seq**
100 RNA-seq data used to facilitate genome annotation was generated using two different methods.
101 This was done so as to include reads from as wide a range of tissues as possible to increase the
102 completeness of the genome annotation. We used data from whole bodies and accessory glands
103 of males, plus whole bodies, ovaries, and heads of females (Table S2).
104
Whole body RNA-seq data was generated from laboratory stock crickets as follows. These
were descended from 500 adult *G. sigillatus* collected in Las Cruces, New Mexico (USA) in
2001 (Ivy & Sakaluk, 2005). Rearing followed previously described methods (Duffield et al.,
2021, 2022). Approximately 500 crickets were housed in ventilated, 55 L plastic storage bins
packed with egg carton. Crickets were given *ad libitum* food (approximately equal parts Purina
Cat Chow Complete Cat Food and Teklad Global 18% Protein Extruded Rodent Diet pellets)
and water (glass vials plugged with moist cotton). Colonies were housed in an environmental
chamber at 32 °C on a 16:8 hour light:dark cycle. Experimental crickets were frozen and stored
at -80 °C at 7 days post-eclosion and shipped on dry ice to LC Sciences (Houston, TX) for
further processing. Total RNA was extracted using Trizol (Invitrogen, USA) following the
manufacturer's procedure. Total RNA quality and quantity were analysed using a Bioanalyzer
2100 and RNA 6000 Nano LabChip Kit (Agilent, USA), with samples obtaining RIN
number >7.0. Then, mRNA was purified in two rounds using Dynabeads Oligo (dT) (Thermo
Fisher, USA) and subsequently fragmented with a Magnesium RNA Fragmentation Module
(New England Biolabs, USA) at 94℃ for 5-7min. The cleaved RNA fragments were reverse-
transcribed to cDNA using SuperScript™ II Reverse Transcriptase (Invitrogen, USA) and then
used to synthesize U-labelled second-stranded DNAs with *E. coli* DNA polymerase I (New
England Biolabs, USA), RNase H (New England Biolabs, USA), and dUTP Solution (Thermo
Fisher, USA). Dual-index adapters were ligated to the fragments, and size selection was

performed with AMPureXP beads. After the heat-labile UDG enzyme (New England Biolabs,

USA) treatment of the U-labelled second-stranded DNAs, the ligated products were amplified

with PCR by under the following conditions: initial denaturation at 95℃ for 3 min; 8 cycles of

denaturation at 98℃ for 15 sec, annealing at 60℃ for 15 sec, and extension at 72℃ for 30 sec;

and then final extension at 72℃ for 5 min. The average insert size for the final cDNA libraries

was 300±50 bp. Sequencing was performed using 2×150bp paired-end sequencing (PE150) on

an Illumina Novaseq™ 6000 following the vendor's recommended protocol (Table S2).

RNA-seq data for male accessory glands, female ovaries, and female heads were generated

from a lab-maintained line (Burns-Dunn et al., 2024; McKermitt et al., 2024), that originated

from the same crickets collected in Las Cruces, 2001 (Ivy & Sakaluk, 2005). Approximately

300 adults were housed in ventilated 62.5 L plastic storage bins packed with egg carton.

Nymphs were maintained in 6 L bins. Crickets were given *ad libitum* food (approximately

equal parts Purina Cat Chow Complete Cat Food and Teklad Global 18% Protein Extruded

Rodent Diet pellets) and water (glass vials plugged with moist cotton). Colonies were

maintained at a constant temperature (30 ± 2 °C) and under a reversed photoperiod (14:10 hr

light:dark cycle). Seven-day old crickets were mated, and two hours later they were stunned on

ice for tissue dissections. Each tissue was immediately snap frozen in liquid nitrogen and stored

-80 °C until further processing. RNA extractions were performed as described in Foquet et al.

(2023). RNA was extracted using a Trizol (Thermo Fisher Scientific) and 1-bromo-3-

chloropropane (BCP, Acros Organics) phase separation using standard procedures, followed

by a DNAse treatment using a TURBO DNA-free kit (Thermo Fisher Scientific). RNA

qualities and concentration were measured on a MultiSkan GO microplate spectrophotometer

with a μDrop adapter plate (ThermoScientific). Only RNA with final 260/230 and 260/280

values higher than two were used for RNA sequencing. RNA samples were sent to Novogene

for library preparation, sequencing, and read pre-processing. Sample quality was assessed with

a Bioanalyzer 2100 (Agilent, USA), with samples obtaining RIN number >5.0 used for

sequencing. RNA was processed with an Illumina TruSeq Stranded mRNA Kit with polyA

enrichment, and sequencing of paired end 150bp reads on a NovaSeq were performed by

Novogene.

Genome assembly

We used HiFiasm v0.19.5-r587 (Cheng et al., 2021), with default settings to generate a draft assembly of *G. sigillatus*. The primary assembly was retained for subsequent refinement and analysis. To eliminate duplicates within the contig-level assembly, Purge_Dups v1.2.6 (Guan et al., 2020) was employed. This identified redundant contigs by mapping the original HiFi reads back to the assembly, removing repetitive or duplicated sequences that could inflate assembly size and misrepresent genomic content. To remove mitochondrial sequences from the draft assembly, MitoHiFi v3.2.1 (Uliano-Silva et al., 2023) was used. This tool identified and removed contigs associated with the mitochondrial genome by comparing them against the published *G. sigillatus* mitochondrial sequence (MW365703.1, Yang et al., 2021), ensuring that only nuclear genomic sequences remained in the primary assembly.

Hi-C data were then incorporated to scaffold the contigs and achieve chromosome-level assembly following Arima Hi-C mapping guidelines (https://github.com/ArimaGenomics/mapping_pipeline). Briefly, this involved mapping with BWA-MEM2 v2.0pre2 (Vasimuddin et al., 2019) and removing PCR duplicates using Picard v3.1.0 (https://broadinstitute.github.io/picard). All Hi-C reads were mapped to the contig-level assembly, providing spatial genomic interaction information. YahS v1.1a-r3 (Zhou et al., 2023) was used to scaffold contigs based on these mapping results, integrating long-range contact information to order and orient contigs accurately. To further refine scaffolding, we used Juicer tools (https://github.com/aidenlab/Juicebox) and Juicebox v3.0.0 (https://github.com/aidenlab/JuicerTools) to visualise the Hi-C contact heatmap and then manually inspect and correct misassembled regions.

Once the final scaffolds were generated, BlobTools2 (Challis et al., 2020) was used to screen for potential contaminant sequences originating from other species. This step identified 23 scaffolds that likely represented contamination, and these were removed to produce the final, cleaned scaffold-level genome assembly. This comprehensive, iterative workflow maximised the likelihood of obtaining an accurate, high-quality chromosome-level *G. sigillatus* genome assembly suitable for further genomic analyses and annotation.

Genome quality assessment

Genome assembly statistics, including total length, N50, and GC content, were calculated using SeqKit v2.4.0 (Shen et al., 2016) to provide a comprehensive overview of the assembly quality.

189 To assess the completeness of the assembled genome, BUSCO v5.4.6 (Manni et al., 2021) was
190 employed, aligning the nucleotide sequences against the Arthropoda dataset (Odb10). This
191 analysis enabled the evaluation of the assembly's coverage and accuracy based on the presence
192 of conserved single-copy orthologs representative of the Arthropoda lineage.
193
194 To evaluate the integrity and accuracy of the genome assembly, we employed the k-mer-based
195 method Merqury v1.3 (Rhie et al., 2020). This approach involves mapping k-mers derived from
196 raw HiFi reads to the contig-level assembly, enabling a detailed assessment of assembly quality.
197 Specifically, Merqury calculates the Quality Value (QV) score, which serves as a metric for
198 the overall accuracy of the assembly and estimates the error rate at the level of individual bases.
199
For evaluating the completeness of the protein-coding gene annotation (see below), BUSCO
was again utilised with the Insecta dataset in 'transcriptome' mode, which is designed to
analyse transcriptome assemblies. This allowed for a thorough assessment of gene prediction
quality, ensuring that the annotated protein-coding genes were representative and complete. By
using both genome and transcriptome completeness assessments, we ensured a robust
evaluation of the assembly's accuracy and the integrity of its functional annotations.
206
Genome annotation
To identify repetitive elements in the genome sequence, three sequential methods were
employed to ensure thorough coverage. First, Tandem Repeats Finder v4.09 (TRF; Benson,
1999) was used to detect simple sequence repeats. The repetitive regions identified by TRF
were masked with 'N' in the genome sequence, effectively obscuring these regions for
downstream analyses. Next, transposable elements (TEs) within the genome were identified
using RepeatMasker v4.1.2-p1 (Tarailo-Graovac & Chen, 2009), with the previously masked
genome sequence serving as input. During this step, the genomic sequences were aligned
against the Repbase database v20181026 (Bao et al., 2015) to specifically search for TEs
homologous to the Arthropoda lineage. The genome was again subjected to hard-masking
based on the identified TEs. Lastly, RepeatModeler2 (Flynn et al., 2020) was applied to conduct
a self-alignment of the genomic sequences to uncover any additional repetitive elements. The
results from RepeatModeler2 were incorporated into a custom repeat database, and
RepeatMasker was run once more to identify repeats based on alignment hits. All repetitive
elements identified through this comprehensive protocol were soft-masked in subsequent
analyses, preserving their sequence while distinguishing them for further study.

For gene prediction, we integrated both homologous protein sequences and RNA-seq data.

Protein sequences from *D. melanogaster* (BDGP6.46, Ensembl; Harrison et al., 2024), *G.*

bimaculatus (Ylla et al., 2021), *T. occipitalis* (Kataoka et al., 2020) and *T. oceanicus* v2 (Zhang

et al., 2024), and reviewed orthologs from Arthropoda were obtained from SwissProt to guide

the annotation process. Additionally, RNA-seq data from multiple tissues and both sexes were

utilised to facilitate transcriptome-based annotation. Prior to gene prediction, raw RNA-seq

reads were quality-trimmed using Fastp v0.23.2 (Chen et al., 2018) and subsequently mapped

to the reference genome with HISAT2 v2.2.0 (Kim et al., 2015), ensuring accurate alignment

for transcript assembly.

To construct the final gene set, we employed a combined annotation strategy that integrated

results from three distinct gene prediction approaches. First, BRAKER3 v3.0 (Gabriel et al.,

2024) was used for *ab initio* gene prediction, leveraging both protein and transcript evidence

to train and predict gene structures. Second, Miniprot v0.10-r225 (H. Li, 2023) was employed

to map homologous protein sequences from closely related species and Arthropoda orthologs

to the reference genome, providing potential coordinates and open reading frame (ORF)

structures of protein-coding genes. Third, RNA-seq data were assembled using two different

tools, StringTie v2.2.1 (Shumate et al., 2022) and Trinity v2.14.0 (Grabherr et al., 2011), to

capture transcript diversity. The Trinity-assembled transcriptome was filtered to retain

sequences longer than 1 Kb, and redundancy was reduced using CD-HIT v4.8.1 (W. Li &

Godzik, 2006). The PASA pipeline v2.5.3 (https://github.com/PASApipeline/PASApipeline)

was then utilised to predict gene structures by mapping these transcripts to the reference

genome. For the transcriptome assembled by StringTie, potential ORFs were identified using

TransDecoder v5.7.1 (https://github.com/TransDecoder/TransDecoder), and coordinates were

mapped to the reference genome. Finally, the EvidenceModeler v2.1.0 (Haas et al., 2008) was

employed to integrate all prediction results into a unified gene set, while the PASA pipeline

refined the final gene models by aligning the transcriptome to predicted gene structures. This

comprehensive approach resulted in a high-confidence, annotated gene set for the target

genome.

Functional annotation

The functions of protein-coding genes were annotated by mapping the isoform sequences

against multiple publicly available functional databases. For Non-redundant (Nr; Sayers et al.,

2022) and Swiss-Prot/TrEMBL (Boeckmann et al., 2003), protein sequences were mapped to

an Insecta-specific subset using DIAMOND blastp v2.0.14.152 (Buchfink et al., 2021),

enabling accurate identification of protein functions within this taxonomic group. To identify

and annotate functional domains within the protein sequences, InterProScan (Paysan-Lafosse

et al., 2023) was employed, which integrates multiple signature databases for a thorough

domain analysis. Additionally, EggNOG mapper (Cantalapiedra et al., 2021) was utilised to

conduct EggNOG (Huerta-Cepas et al., 2019) analysis, from which Gene Ontology (GO) terms

were extracted to provide insights into the biological processes, molecular functions, and

cellular components associated with each protein. Finally, KEGG pathway (Kanehisa et al.,

2016) annotations were acquired using the KAAS online server (Moriya et al., 2007), allowing

for the identification of biochemical pathways and the functional context of the proteins within

broader metabolic networks.

Ortholog group identification and phylogenetic analysis

Orthologous protein sequences from six other genomes, including *Anabrus simplex* (NCBI

RefSeq accession: GCF_040414725.1 by United States Department of Agriculture), *Gryllus*

longicercus (Szrajer et al., 2024), *Gryllus bimaculatus* (Ylla et al., 2021), *Teleogryllus*

oceanicus (Zhang et al., 2024), *Teleogryllus occipitalis* (Kataoka et al., 2020) and *Schistocerca*

americana (iqSchAmer2.1, https://www.dnazoo.org/assemblies/schistocerca_americana),

along with our *G. sigillatus* proteins, were collected for ortholog group identification using

OrthoFinder v2.2.5 (Emms & Kelly, 2019). To construct a robust phylogenetic framework, we

focused on single-copy orthologs, which were used to build a maximum-likelihood

phylogenetic tree, providing insights into the evolutionary relationships among these species.

To estimate divergence times, we incorporated reference divergence times from fossil evidence

obtained from the TimeTree database (Kumar et al., 2022), ensuring that our analysis was

anchored to well-established evolutionary benchmarks. The Bayesian inference tool

MCMCTree, part of the PAML v4.10.7 package (Reis & Yang, 2011), was then employed to

estimate divergence times across all nodes within the phylogeny using only four-fold sites

within all single-copy gene sequences.

Gene family analysis

Using the ultrametric tree generated by MCMCTree, we analysed the expanded and contracted

gene families across all nodes of the phylogeny with CAFE5 v1.1 (Mendes et al., 2021), a

software designed for modelling gene family evolution. To optimise the analysis, we tested a

range of discrete gamma rate (k) values from 2 to 6, which account for the variation in
evolutionary rates across gene families. The k=4 that yielded the highest maximum likelihood
was selected for the final model, ensuring an accurate fit to the data. To identify gene families
that experienced significant changes specifically in *G. sigillatus*, we conducted a Likelihood
Ratio Test (LRT), which enabled us to distinguish gene families that had undergone statistically
significant expansions or contractions.
297

298 **Statistical analysis and visualisation**
299 Statistical analyses were performed using R v4.0.2. Diagrams were generated using ggplot2 in
300 the tidyverse package (Wickham et al., 2019). The genomic circos plot was generated using
301 shinyCircos v2.0 (Wang et al., 2023).
302

Results and Discussion
Chromosome-level genome assembly of *G. sigillatus*
We generated a high-quality, chromosome level annotated reference genome for *G. sigillatus*
(Figure 1b; Table 1; Table S3). The completed assembly has a total size of 2.17 Gb and
comprises 11 pseudochromosomes (Figure 1c; Table S4), consistent with the known karyotype
of *G. sigillatus* (You et al., 2007). Based on the latter karyotype study, the longest scaffold in
the assembly was marked as chromosome X and autosomes were named by order of length.
The assembly has a scaffold N50 of 200.35 Mb and a maximum scaffold length of 420.78 Mb
(Figure 1b, Table S3), and k-mer based quality estimation yielded a QV score of 61.21 and per-
base error rate of 7.56×10^{-7} (Table S4), indicating a highly contiguous and complete genome
assembly. To further assess the completeness and quality of the assembly, we performed
Benchmarking Universal Single-Copy Orthologs (BUSCO) analysis using the insecta_odb10,
which yielded a BUSCO recovery score of 99.49% (Table S5).
316

Genome annotation revealed that repetitive sequences accounted for 48.22% of the genome,
including DNA transposons (16.22%), long interspersed nuclear elements (LINEs, 24.13%),
short interspersed nuclear elements (SINEs, 1.81%), and long terminal repeat (LTR)
retrotransposons (0.07%) (Figure 1b; Table S6). This repeat content is comparable to that
observed in related species (Szrajer et al., 2024; Ylla et al., 2021; Zhang et al., 2024),
suggesting relatively conserved genomic architecture and evolutionary patterns of transposable
elements.
324

We predicted a total of 16,179 protein-coding genes using a combination of *ab initio* prediction
and evidence-based methods, incorporating RNA-seq data (Figure 1b; Table S2) and protein
homology. Functional annotation was successful for 84.56% of the total predicted genes, using
alignment against several public databases including the Non-redundant Protein Sequence
Database (Nr), TrEMBL, Gene Ontology (GO), EggNOG, and the Kyoto Encyclopedia of
Genes and Genomes (KEGG).
331
332 **Table 1. Summary of *Gryllodes sigillatus* genome assembly and annotation.**

Genome assembly	*G. sigillatus*
Total length (bp)	2,173,690,933
Contig N50 (bp)	9,022,296
Scaffold N50 (bp)	200,356,516
Longest scaffold (bp)	420,784,874
Hi-C anchoring rate (%)	98.86
BUSCO (%)	99.49
Repeat content (%)	48.22
Gene number	16,179

Figure 1. *Gryllodes sigillatus* and its genome characteristics. (a) A female *G. sigillatus*. **(b)** Features of the *G. sigillatus* genome. Outer track (grey) illustrates the 11 chromosomes, with the largest identified as the putative X chromosome on the basis of a published karyotype study (You et al., 2007). Genome position is indicated by the outer numbers. The inner purple track shows GC content, and the inner pink and green tracks show genome-wide gene and repeat density, respectively. **(c)** Hi-C contact map of the genome assembly indicates high contiguity and coverage. **Photo credit**: NW Bailey.

Genome evolution and phylogenetic analysis

Our study provides the first reference genome within the *Gryllodes* genus. We also performed a phylogenetic analysis that broadly supports the known evolutionary history of *Gryllodes* (Song et al., 2020). A total of six Ensiferan and one Caeliferan genome (the latter as an outgroup) were selected to identify 15,123 orthogroups of which 4,157 comprised single-copy orthologs (Figure S1). A maximum-likelihood species tree was constructed using concatenated alignments of these orthologs, confirming previously-reconstructed evolutionary relationships (Song et al., 2020), validating the inferred speciation timeline, and positioning *Gryllodes* within its evolutionary context (Figure 2a). Divergence time estimates indicate that *Gryllodes* diverged from other *Gryllidae* species approximately 141-199 million years ago (Mya).

Analysis of gene family evolution across orthopterans revealed 377 expanded and 783 contracted gene families in *G. sigillatus* (Figure 2a). Focusing on gene families that exhibited significant changes, we identified 28 that were notably expanded in *G. sigillatus* relative to other orthopteran species (Likelihood Ratio Test, $P < 0.01$), consisting of 146 protein-coding genes in *G. sigillatus* (Table S7). Many of these expanded families are linked to histone-related functions. Histones are central to nucleosome formation, which packages DNA within the nucleus and regulates gene expression through chromatin remodelling. In insects, histone modifications influence development (Glastad et al., 2019), stress responses (Gupta & Nair, 2024), and genome stability (Glastad et al., 2019), and the expansion of histone-related gene families in *G. sigillatus* may be associated with an enhanced capacity for chromatin remodelling and genome organisation, consistent with adaptations that enable *G. sigillatus* to be resilient in diverse ecological niches.

Other expanded gene families are associated with metabolism and energy production, including those encoding succinyl-CoA ligase (SCL), NADH dehydrogenase (complex I subunit 1), and trypsin. Succinyl-CoA ligase, a key enzyme in the tricarboxylic acid (TCA) cycle, catalyses the conversion of succinyl-CoA to succinate, producing ATP or GTP through substrate-level phosphorylation. An expansion in SCL genes may enhance TCA cycle efficiency, boosting energy production to meet increased metabolic demands for activities such as locomotion, growth, and reproduction; NADH dehydrogenase, a mitochondrial complex I component, initiates the electron transport chain by transferring electrons from NADH to ubiquinone, driving ATP synthesis via oxidative phosphorylation (Garcia et al., 2017). The expansion of genes encoding NADH dehydrogenase subunits could increase electron transport efficiency,

improving ATP generation. Energy metabolism pathways and their evolution in other systems
have been linked to the immunity-metabolism interface (Dolezal et al., 2019), detoxification
of and resistance to xenobiotics (Gospodaryov et al., 2020), biological invasion (Wang et al.,
2013), dispersal and flight (Yang et al., 2014; Treidel et al., 2023). While flight performance
is less relevant in *G. sigillatus*, flight muscles are used for male reproductive signaling –
acoustic advertisement song – which has been shown to be energetically constrained (Houslay
et al., 2017). Overall, metabolic enhancement may further provide *G. sigillatus* with greater
resilience to environmental fluctuations in energy availability, supporting sustained activity
and survival across diverse conditions. Concentrating on these gene families during selective
breeding efforts, for example through marker-assisted selection, could facilitate genetic
improvements in nutritional requirements, energy efficiency, growth rates, and reproductive
success.

The expansion of the trypsin gene family suggests adaptations in digestive processes. Trypsins
are serine proteases involved in protein digestion in the insect midgut, hydrolysing peptide
bonds to aid nutrient absorption (Meriño-Cabrera et al., 2022; Muhlia-Almazán et al., 2008).
An increased repertoire of trypsin genes may allow *G. sigillatus* to more effectively exploit
diverse dietary proteins, supporting rapid growth and high reproductive output. Trypsins are
also implicated in immune responses, such as activating the prophenoloxidase system involved
in melanisation and pathogen defence (González-Santoyo & Córdoba-Aguilar, 2012; Kanost
& Gorman, 2008); thus, their expansion may also enhance immune capabilities in *G. sigillatus*.
Determining whether such genes involving immunity are associated with increased resistance
to pests and pathogens in a farming setting can help identify strains with specific haplotypes
that strengthen immunity to pests, reduce mortality rates, and minimise reliance on chemical
treatments. An accurate assessment of genetic variation at such loci across the global range of
farmed cricket species is a priority for future work, as the evolutionary potential for responses
to artificial selection are currently unknown, and natural variation is likely to be considerably
higher than that available in farmed stock populations.

Finally, the expansion of the cytochrome P450 (CYP) gene family, known for its roles in
detoxification and the immune response, was observed (Feyereisen, 2012). Cytochrome P450
enzymes are involved in the metabolism of endogenous substrates and detoxification of
xenobiotics (Nauen et al., 2022), including plant secondary metabolites and synthetic
insecticides (Dermauw et al., 2020). Additionally, CYP enzymes contribute to the synthesis

and degradation of hormones such as ecdysteroids and juvenile hormones, which regulate

development and reproduction. CYP expansion suggests that *G. sigillatus* may possess an

enhanced ability to detoxify a broad range of harmful compounds, offering a selective

advantage in chemically diverse environments and in energy conversion from agricultural by-

products, thus contributing to more sustainable and economical cricket farming.

These findings highlight potentially distinctive genetics underlying adaptive traits in *G.*

sigillatus, shedding light on its evolutionary trajectory within Orthoptera. The expansion of

gene families related to chromatin remodelling, metabolism, digestion, and detoxification

suggests that *G. sigillatus* has evolved genetic adaptations enhancing its physiological

functions and ecological resilience, which is consistent with its global, cosmopolitan

distribution in tropical regions and introduction into many non-native habitats. These

adaptations likely support rapid responses to environmental stresses, optimise energy use, and

strengthen defences against pathogens and toxins. Genomic resources will help to devise

strategies for selectively breeding *G. sigillatus* for optimal reproductive traits in farming

applications, as well as for unpicking the evolutionary causes and consequences of nuptial gift-

giving, multiple mating, and mating system evolution in fundamental behavioural ecology

research.

Figure 2. Evolution of gene family expansion and contraction in selected orthopterans. (a) Phylogenetic tree illustrates the divergence times and gene family evolution across seven Orthoptera species. Branch lengths represent evolutionary distances, with upper and lower divergence time estimates (in units of 100 million years ago, Mya) labelled along each branch. Numbers in red and green indicate the number of significantly expanded (+) and contracted (−) gene families along each lineage, respectively. **(b)** Heatmap illustrating counts of gene families that are significantly expanded in *G. sigillatus* (*ii*), with darker shades representing higher counts. Gene families marked with an asterisk (*****) represent those that failed to match a verified family name across Insecta but have homologous sequences in human gene families, indicating possible highly conserved functions. **Photo credits:** *Anabrus simplex*, *Gryllodes sigillatus*, *Gryllus bimaculatus* and *Teleogryllus oceanicus* by NW Bailey; *Gryllus longicercus* by Jeff Hollenbeck, modified by Shangzhe Zhang, licensed under CC BY-ND-NC 1.0; *Teleogryllus occipitalis* by Takaaki Hattori, modified by Shangzhe Zhang, licensed under CC-BY 4.0; *Schistocerca americana* by Brandon Woo with permission granted.

Conclusion

There are over 6,000 species of crickets in the infraorder Gryllidea, over 60 of which are known to be consumed and, in some cases, farmed by humans (Cigliano et al., 2018; Magara et al., 2021). Generating a high-quality chromosome-scale assembly of one of the most globally widespread and commonly-farmed species, *Gryllodes sigillatus*, is a key step in promoting the adoption of genome-informed farming improvements for insects. The widespread farming of *G. sigillatus* is likely driven by their worldwide, cosmopolitan distribution and ease of collection (Weissman et al., 2012). However, there are many cricket species with farming potential, and the resources developed here will facilitate their study and use more broadly (Dossey et al., 2023; Magara et al., 2019, 2021). Recently, genomic resources, such as genome assemblies, transcriptome sequences, have increasingly deployed to improve strategies for farming crickets (Kataoka et al., 2022; Nakamura et al., 2022). Comparative and phylogenetic approaches continue to advance our understanding of the evolutionary history of crickets and allied taxa (Gray et al., 2020; Song et al., 2020; Dong et al., 2024), and genomic approaches in these species benefit from a longstanding history of major contributions to fundamental research in evolutionary and behavioural biology. Recent work, for example, has shed light on the genomics of speciation (Xu & Shaw, 2021; Yusuf et al., 2024), adaptation in the wild (Pascoal et al., 2020; Zhang et al., 2021; Zhang et al., 2024; Rayner et al. 2024), evolutionary developmental biology (Barnett et al., 2019; Ohde et al., 2022), and sperm competition, mate choice, and sexual selection (Simmons et al., 2013; Xu & Shaw, 2019). In addition, our findings provide valuable resources for precisely identifying genetic markers using approaches such as genome-wide association study (GWAS) and marker-assisted selection, and for guiding the genomic editing of breeding traits through RNA*i* or CRISPR-Cas9. These technologies have been successfully applied in other farmed crickets, including *Acheta domesticus* (Dossey et al., 2023) and *Gryllus bimaculatus* (Watanabe et al., 2017), and could be similarly implemented in *G. sigillatus*. Future research efforts will benefit not only from exploring the potential for *G. sigillatus* in greater depth using these genome-informed tools for applied and basic research, but also developing genomic resources across a broader taxonomic sampling of commonly studied Gryllids.

Supplementary information

Supplementary Materials include Figs. S1, Table S1 to S7, Data table S1.

Data availability

All genomic data (long-reads sequencing data, and Hi-C sequencing data) have been deposited at European Nucleotide Archive (ENA) under PRJEB84028. Transcriptome data have been deposited at PRJEB85996 and PRJNA1204288. Sequences of the genome assembly is publicly available at ENA under accession GCA_965111905. Annotation files of the genome assembly has been deposited at Zenodo as DOI: 10.5281/zenodo.14617131.

Acknowledgements

The authors gratefully acknowledge funding support from the Australian Research Council to JH (DP180101708), the UK Natural Environment Research Council to NWB (NE/W001616/1), the China Scholarship Council to SZ (202106180022), and the United States National Science Foundation to SKS, BMS, and JH (IOS 16–54028). This research was also supported in part by the U.S. Department of Agriculture, Agricultural Research Service (Project Number: 6064-32000-001-000D to KRD and 5010-22410-023-00-D to JLR). We thank the UK Biotechnology and Biological Sciences Research Council (BB/S019669/1 and BB/X019683/1) and James Hutton Institute for access to the UK's Crop Diversity Bioinformatics High Performance Computing Cluster, with particularly acknowledgement of Ian Milne's inexhaustible patience as sysadmin. Lianne Baker assisted with sample shipping and Tanya Sneddon with DNA extractions. The authors are grateful for constructive feedback on the manuscript from Xin Du. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

Conflict of interest statement

The authors declare no conflict of interest.

Author contributions

Conceived the study: JH, NWB. Designed experiments: all authors. Secured funding: JH, BMS, JLR, KRD, SKS, NWB. Performed experimental work: KRD, BF, SZ. Analysed data: KRD, BF, SZ. Led manuscript writing: SZ, NWB. Contributed to writing: JH, BF, SKS, BMS, JLR, KRD.

References

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA*, *6*(1), 11. https://doi.org/10.1186/s13100-015-0041-9

Barnett, A. A., Nakamura, T., Extavour, C. G. (2019). Hox genes limit germ cell formation in the short germ insect *Gryllus bimaculatus*. *Proceedings of the National Academy of Sciences, U.S.A.* *116* (33), 16430-16435. https://doi.org/10.1073/pnas.1816024116

Bateman, P. W., & MacFadyen, D. N. (1999). Mate Guarding in the Cricket Gryllodes sigillatus: Influence of Multiple Potential Partners. *Ethology*, *105*(11), 949–957. https://doi.org/10.1046/j.1439-0310.1999.00484.x

Benson, G. (1999). Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research*, *27*(2), 573–580. https://doi.org/10.1093/nar/27.2.573

Blankers, T., Oh, K. P., Bombarely, A., Shaw, K. L. (2018) The genomic architecture of a rapid island radiation: recombination rate variation, chromosome structure, and genome assembly of the Hawaiian cricket *Laupala*. *Genetics* *209*(4), 1329-1344. https://doi.org/10.1534/genetics.118.300894

Boeckmann, B., Bairoch, A., Apweiler, R., Blatter, M.-C., Estreicher, A., Gasteiger, E., Martin, M. J., Michoud, K., O'Donovan, C., Phan, I., Pilbout, S., & Schneider, M. (2003). The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Research*, *31*(1), 365–370. https://doi.org/10.1093/nar/gkg095

Buchfink, B., Reuter, K., & Drost, H.-G. (2021). Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nature Methods*, *18*(4), 366–368. https://doi.org/10.1038/s41592-021-01101-x

Burns-Dunn, S., Mortys, T., House, C. M., Mitchell, C., Duffield, K. R., Foquet, B., Sadd, B. M., Sakaluk, S. K., & Hunt, J. (2024). Sexually antagonistic coevolution of the male nuptial gift and female feeding behaviour in decorated crickets. *Proceedings of the Royal Society B: Biological Sciences*, *291*(2026), 20240804. https://doi.org/10.1098/rspb.2024.0804

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Molecular Biology and Evolution*, *38*(12), 5825–5829. https://doi.org/10.1093/molbev/msab293

541   Challis, R., Richards, E., Rajan, J., Cochrane, G., & Blaxter, M. (2020). BlobToolKit –
542   Interactive Quality Assessment of Genome Assemblies. *G3 Genes|Genomes|Genetics*,
543   *10*(4), 1361–1374. https://doi.org/10.1534/g3.119.400908

544   Champagnon, J., & Cueva del Castillo, R. (2008). Female Mate Choice, Calling Song and
545   Genetic Variance in the Cricket, *Gryllodes sigillatus*. *Ethology*, *114*(3), 223–230.
546   https://doi.org/10.1111/j.1439-0310.2007.01470.x

547   Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ
548   preprocessor. *Bioinformatics*, *34*(17), i884–i890.
549   https://doi.org/10.1093/bioinformatics/bty560

550   Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de
551   novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2),
552   Article 2. https://doi.org/10.1038/s41592-020-01056-5

553   Cigliano, M., Braun, H., Eades, D., & Otte, D. (2018). *"Infraorder Gryllidea". Orthoptera*
554   *Species File. Version 5.0/5.0*. Facultad de Ciencias Naturales y Museo, Universidad
555   Nacional de La Plata (UNLP).

556   Dermauw, W., Van Leeuwen, T., & Feyereisen, R. (2020). Diversity and evolution of the P450
557   family in arthropods. *Insect Biochemistry and Molecular Biology*, *127*, 103490.
558   https://doi.org/10.1016/j.ibmb.2020.103490

559   Dolezal, T., Krejcova, G., Bajgar, A., Nedbalova, P., & Strasser, P. (2019). Molecular
560   regulations of metabolism during immune response in insects. *Insect Biochemistry and*
561   *Molecular Biology*, *109*, 31–42. https://doi.org/10.1016/j.ibmb.2019.04.005

562   Dong, J., Liu, Y., Tan, M. K., Wahab, R. A., Nattier, R., Chifflet-Belle, P., & Robillard, T.
563   (2024). Museomics allows comparative analyses of mitochondrial genomes in the
564   family Gryllidae (Insecta, Orthoptera) and confirms its phylogenetic relationships.
565   *PeerJ*, *12*, e17734. https://doi.org/10.7717/peerj.17734

566   Dossey, A. T., Oppert, B., Chu, F.-C., Lorenzen, M. D., Scheffler, B., Simpson, S., Koren, S.,
567   Johnston, J. S., Kataoka, K., & Ide, K. (2023). Genome and Genetic Engineering of the
568   House Cricket (Acheta domesticus): A Resource for Sustainable Agriculture.
569   *Biomolecules*, *13*(4), Article 4. https://doi.org/10.3390/biom13040589

570   Duffield, K. R., Foquet, B., Stasko, J. A., Hunt, J., Sadd, B. M., Sakaluk, S. K., & Ramirez, J.
571   L. (2022). Induction of Multiple Immune Signaling Pathways in *Gryllodes sigillatus*
572   Crickets during Overt Viral Infections. *Viruses*, *14*(12), Article 12.
573   https://doi.org/10.3390/v14122712

Duffield, K. R., Hunt, J., Sadd, B. M., Sakaluk, S. K., Oppert, B., Rosario, K., Behle, R. W., &
Ramirez, J. L. (2021). Active and Covert Infections of Cricket Iridovirus and Acheta
domesticus Densovirus in Reared *Gryllodes sigillatus* Crickets. *Frontiers in*
Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.780796

Emms, D. M., & Kelly, S. (2019). OrthoFinder: Phylogenetic orthology inference for
comparative genomics. *Genome Biology*, *20*(1), 238. https://doi.org/10.1186/s13059-
019-1832-y

Feyereisen, R. (2012). 8—Insect CYP Genes and P450 Enzymes. In L. I. Gilbert (Ed.), *Insect*
Molecular Biology and Biochemistry (pp. 236–316). Academic Press.
https://doi.org/10.1016/B978-0-12-384747-8.10008-X

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.
(2020). RepeatModeler2 for automated genomic discovery of transposable element
families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.
https://doi.org/10.1073/pnas.1921046117

Foquet, B., Rapkin, J., Sharma, M. D., Sadd, B. M., Sakaluk, S. K., & Hunt, J. (2023).
Transcriptomic responses of females to consumption of nuptial food gifts as a potential
mediator of sexual conflict in decorated crickets. *Journal of Evolutionary Biology*,
36(1), 183–194. https://doi.org/10.1111/jeb.14114

Gabriel, L., Brůna, T., Hoff, K. J., Ebel, M., Lomsadze, A., Borodovsky, M., & Stanke, M.
(2024). BRAKER3: Fully automated genome annotation using RNA-seq and protein
evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Research*, *34*(5),
769–777. https://doi.org/10.1101/gr.278090.123

Garcia, C. J., Khajeh, J., Coulanges, E., Chen, E. I., & Owusu-Ansah, E. (2017). Regulation of
Mitochondrial Complex I Biogenesis in *Drosophila* Flight Muscles. *Cell Reports*, *20*(1),
264–278. https://doi.org/10.1016/j.celrep.2017.06.015

Glastad, K. M., Hunt, B. G., & Goodisman, M. A. D. (2019). Epigenetics in Insects: Genome
Regulation and the Generation of Phenotypic Diversity. *Annual Review of Entomology*,
64, 185–203. https://doi.org/10.1146/annurev-ento-011118-111914

González-Santoyo, I., & Córdoba-Aguilar, A. (2012). Phenoloxidase: A key component of the
insect immune system. *Entomologia Experimentalis et Applicata*, *142*(1), 1–16.
https://doi.org/10.1111/j.1570-7458.2011.01187.x

Gospodaryov, D. V., Strilbytska, O. M., Semaniuk, U. V., Perkhulyn, N. V., Rovenko, B. M.,
Yurkevych, I. S., Barata, A. G., Dick, T. P., Lushchak, O. V., & Jacobs, H. T. (2020).
Alternative NADH dehydrogenase extends lifespan and increases resistance to

xenobiotics in *Drosophila*. *Biogerontology*, *21*(2), 155–171.
https://doi.org/10.1007/s10522-019-09849-8

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis,
X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke,
A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev,
A. (2011). Trinity: Reconstructing a full-length transcriptome without a genome from
RNA-Seq data. *Nature Biotechnology*, *29*(7), 644–652.
https://doi.org/10.1038/nbt.1883

Grabowski, N. T., Abdulmawjood, A., Acheuk, F., Barragán Fonseca, K., Chhay, T., Costa
Neto, E. M., Ferri, M., Franco Olivas, J., González Aguilar, D. G., Keo, S.,
Lertpatarakomol, R., Miech, P., Piofczyk, T., Proscia, F., Mitchaothai, J., Guerfali, M.
M., Sayed, W., Tchibozo, S., & Plötz, M. (2022). Review: Insects—A Source of Safe
and Sustainable Food?— "Jein" (Yes and No). *Frontiers in Sustainable Food Systems*,
5. https://doi.org/10.3389/fsufs.2021.701797

Gray, D. A., Weissman, D. B., Cole, J. A., Lemmon, E. M., & Lemmon, A. R. (2020).
Multilocus phylogeny of Gryllus field crickets (Orthoptera: Gryllidae: Gryllinae)
utilizing anchored hybrid enrichment. *Zootaxa*, *4750*(3), Article 3.
https://doi.org/10.11646/zootaxa.4750.3.2

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying
and removing haplotypic duplication in primary genome assemblies. *Bioinformatics*,
36(9), 2896–2898. https://doi.org/10.1093/bioinformatics/btaa025

Gupta, A., & Nair, S. (2024). Epigenetic processes in insect adaptation to environmental stress.
Current Opinion in Insect Science, 101294. https://doi.org/10.1016/j.cois.2024.101294

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell, C. R.,
& Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using
EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biology*,
9(1), R7. https://doi.org/10.1186/gb-2008-9-1-r7

Harrison, P. W., Amode, M. R., Austine-Orimoloye, O., Azov, A. G., Barba, M., Barnes, I.,
Becker, A., Bennett, R., Berry, A., Bhai, J., Bhurji, S. K., Boddu, S., Branco Lins, P.
R., Brooks, L., Ramaraju, S. B., Campbell, L. I., Martinez, M. C., Charkhchi, M.,
Chougule, K., … Yates, A. D. (2024). Ensembl 2024. *Nucleic Acids Research*, *52*(D1),
D891–D899. https://doi.org/10.1093/nar/gkad1049

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H.,
Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., & Bork, P. (2019).

eggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology
resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, *47*(D1),
D309–D314. https://doi.org/10.1093/nar/gky1085

Huis, A. van, Rumpold, B., Maya, C., & Roos, N. (2021). Nutritional Qualities and
Enhancement of Edible Insects. *Annual Review of Nutrition*, *41*(Volume 41, 2021),
551–576. https://doi.org/10.1146/annurev-nutr-041520-010856

Ivy, T. M., & Sakaluk, S. K. (2005). Polyandry Promotes Enhanced Offspring Survival in
Decorated Crickets. *Evolution*, *59*(1), 152–159. https://doi.org/10.1111/j.0014-
3820.2005.tb00902.x

Johnsson, M. (2023). Genomics in animal breeding from the perspectives of matrices and
molecules. *Hereditas*, *160*(1), 20. https://doi.org/10.1186/s41065-023-00285-w

Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., & Tanabe, M. (2016). KEGG as a
reference resource for gene and protein annotation. *Nucleic Acids Research*, *44*(D1),
D457–D462. https://doi.org/10.1093/nar/gkv1070

Kanost, M. R., & Gorman, M. J. (2008). 4—PHENOLOXIDASES IN INSECT IMMUNITY.
In N. E. Beckage (Ed.), *Insect Immunology* (pp. 69–96). Academic Press.
https://doi.org/10.1016/B978-012373976-6.50006-9

Kasdorf, S. Y., Muzzatti, M. J., Haider, F., Bertram, S. M., & MacMillan, H. A. (2025).
Brewery waste as a sustainable protein source for the banded cricket (Gryllodes
sigillatus). *Journal of Insects as Food and Feed*. https://doi.org/10.1163/23524588-
00001368

Kataoka, K., Minei, R., Ide, K., Ogura, A., Takeyama, H., Takeda, M., Suzuki, T., Yura, K., &
Asahi, T. (2020). The Draft Genome Dataset of the Asian Cricket *Teleogryllus*
occipitalis for Molecular Research Toward Entomophagy. *Frontiers in Genetics*, *11*.
https://doi.org/10.3389/fgene.2020.00470

Kataoka, K., Togawa, Y., Sanno, R., Asahi, T., & Yura, K. (2022). Dissecting cricket genomes
for the advancement of entomology and entomophagy. *Biophysical Reviews*, *14*(1), 75–
97. https://doi.org/10.1007/s12551-021-00924-4

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low
memory requirements. *Nature Methods*, *12*(4), Article 4.
https://doi.org/10.1038/nmeth.3317

Kong, J. D., Ritchie, M. W., Vadboncoeur, É., MacMillan, H. A., & Bertram, S. M. (2024).
Growth, development, and life history of a mass-reared edible insect, *Gryllodes*

sigillatus (Orthoptera: Gryllidae) (p. 2024.10.08.617250). *bioRxiv*.

https://doi.org/10.1101/2024.10.08.617250

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher, G.,

& Hedges, S. B. (2022). TimeTree 5: An Expanded Resource for Species Divergence

Times. *Molecular Biology and Evolution*, *39*(8), msac174.

https://doi.org/10.1093/molbev/msac174

Houslay, T. M., Houslay, K. F., Rapkin, J., Hunt, J., & Bussière, L. F. (2017). Mating

opportunities and energetic constraints drive variation in age-dependent sexual

signalling. *Functional Ecology*, *31*(3), 728–741. https://doi.org/10.1111/1365-

2435.12766

Lange, K. W., & Nakamura, Y. (2023). Potential contribution of edible insects to sustainable

consumption and production. *Frontiers in Sustainability*, *4*.

https://doi.org/10.3389/frsus.2023.1112950

Li, H. (2023). Protein-to-genome alignment with miniprot. *Bioinformatics*, *39*(1), btad014.

https://doi.org/10.1093/bioinformatics/btad014

Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics (Oxford, England)*, *22*(13), 1658–1659.

https://doi.org/10.1093/bioinformatics/btl158

Magara, H. J. O., Niassy, S., Ayieko, M. A., Mukundamago, M., Egonyu, J. P., Tanga, C. M.,

Kimathi, E. K., Ongere, J. O., Fiaboe, K. K. M., Hugel, S., Orinda, M. A., Roos, N., &

Ekesi, S. (2021). Edible Crickets (Orthoptera) Around the World: Distribution,

Nutritional Value, and Other Benefits—A Review. *Frontiers in Nutrition*, *7*.

https://doi.org/10.3389/fnut.2020.537915

Magara, H. J. O., Tanga, C. M., Ayieko, M. A., Hugel, S., Mohamed, S. A., Khamis, F. M.,

Salifu, D., Niassy, S., Sevgan, S., Fiaboe, K. K. M., Roos, N., & Ekesi, S. (2019).

Performance of Newly Described Native Edible Cricket *Scapsipedus icipe* (Orthoptera:

Gryllidae) on Various Diets of Relevance for Farming. *Journal of Economic*

Entomology, *112*(2), 653–664. https://doi.org/10.1093/jee/toy397

Manni, M., Berkeley, M. R., Seppey, M., & Zdobnov, E. M. (2021). BUSCO: Assessing

Genomic Data Quality and Beyond. *Current Protocols*, *1*(12), e323.

https://doi.org/10.1002/cpz1.323

McKermitt, J. T., Foquet, B., Kuna, W., Hunt, J., Sadd, B. M., & Sakaluk, S. K. (2024).

Experimental evolution under varying sex ratio and behavioral plasticity in response to

perceived competitive environment independently affect calling effort in male crickets.

Evolution, *78*(3), 453–462. https://doi.org/10.1093/evolut/qpad224

Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2021). CAFE 5 models variation

in evolutionary rates among gene families. *Bioinformatics*, *36*(22–23), 5516–5518.

https://doi.org/10.1093/bioinformatics/btaa1022

Meriño-Cabrera, Y. B., Oliveira, M. G. de A., Meriño-Cabrera, Y. B., & Oliveira, M. G. de A.

(2022). Trypsins: Structural Characterization and Inhibition Focus in Insects. In

Hydrolases. IntechOpen. https://doi.org/10.5772/intechopen.102632

Morales-Ramos, J. A., Rojas, M. G., Dossey, A. T., & Berhow, M. (2020). Self-selection of

food ingredients and agricultural by-products by the house cricket, *Acheta domesticus*

(Orthoptera: Gryllidae): A holistic approach to develop optimized diets. *PLOS ONE*,

15(1), e0227400. https://doi.org/10.1371/journal.pone.0227400

Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C., & Kanehisa, M. (2007). KAAS: An

automatic genome annotation and pathway reconstruction server. *Nucleic Acids*

Research, *35*(suppl_2), W182–W185. https://doi.org/10.1093/nar/gkm321

Muhlia-Almazán, A., Sánchez-Paz, A., & García-Carreño, F. L. (2008). Invertebrate trypsins:

A review. *Journal of Comparative Physiology B*, *178*(6), 655–672.

https://doi.org/10.1007/s00360-008-0263-y

Muzzatti, M. J., Kong, J. D., McColville, E. R., Brzezinski, H., Stabile, C. C., MacMillan, H.

A., & Bertram, S. M. (2025a). Diet particle size influences tropical house cricket life

history. *Journal of Insects as Food and Feed.* https://doi.org/10.1163/23524588-

00001365

Muzzatti, M. J., Ritchie, M. W., Bess, E. C., Bertram, S. M., & MacMillan, H. A. (2025b).

Farmed crickets (Orthoptera: Gryllidae) raised with dermestids (Coleoptera:

Dermestidae) suffer from reduced and delayed growth, but not enough to explain

reports of dramatic yield loss. *Journal of Economic Entomology*, 118(1), 160–171.

https://doi.org/10.1093/jee/toae208

Nakamura, T., Ylla, G., & Extavour, C. G. (2022). Genomics and genome editing techniques

of crickets, an emerging model insect for biology and food science. *Current Opinion in*

Insect Science, *50*, 100881. https://doi.org/10.1016/j.cois.2022.100881

Nauen, R., Bass, C., Feyereisen, R., & Vontas, J. (2022). The Role of Cytochrome P450s in

Insect Toxicology and Resistance. *Annual Review of Entomology*, *67*(Volume 67, 2022),

105–124. https://doi.org/10.1146/annurev-ento-070621-061328

Ohde, T., Mito, T., & Niimi, T. (2022). A hemimetabolous wing development suggests the wing origin from lateral tergum of a wingless ancestor. *Nature Communications*, *13*(1), 979. https://doi.org/10.1038/s41467-022-28624-x

Pascoal, S., Risse, J. E., Zhang, X., Blaxter, M., Cezard, T., Challis, R. J., Gharbi, K., Hunt, J., Kumar, S., Langan, E., Liu, X., Rayner, J. G., Ritchie, M. G., Snoek, B. L., Trivedi, U., Bailey, N. W. (2020). Field cricket genome reveals the footprint of recent, abrupt adaptation in the wild. *Evolution Letters 4*(1), 19-33. https://doi.org/10.1002/evl3.148

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B. L., Salazar, G. A., Bileschi, M. L., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D. H., Letunić, I., Marchler-Bauer, A., Mi, H., Natale, D. A., Orengo, C. A., Pandurangan, A. P., Rivoire, C., … Bateman, A. (2023). InterPro in 2022. *Nucleic Acids Research*, *51*(D1), D418–D427. https://doi.org/10.1093/nar/gkac993

Rapkin, J., Jensen, K., Archer, C. R., House, C. M., Sakaluk, S. K., Castillo, E. del, & Hunt, J. (2018). The Geometry of Nutrient Space–Based Life-History Trade-Offs: Sex-Specific Effects of Macronutrient Intake on the Trade-Off between Encapsulation Ability and Reproductive Effort in Decorated Crickets. *The American Naturalist*, *191*(4), 452–474. https://doi.org/10.1086/696147

Rayner, J. G., Eichenberger, F., Bainbridge, J. V. A., Zhang, S., Zhang, X., Yusuf, L. H. Balenger, S., Gaggiotti, O. E., Bailey, N. W. (2024). Competing adaptations maintain nonadaptive variation in a wild cricket population. *Proceedings of the National Academy of Sciences, U.S.A. 121*(32), e2317879121. https://doi.org/10.1073/pnas.2317879121

Reis, M. dos, & Yang, Z. (2011). Approximate Likelihood Calculation on a Phylogeny for Bayesian Estimation of Divergence Times. *Molecular Biology and Evolution*, *28*(7), 2161–2172. https://doi.org/10.1093/molbev/msr045

Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology*, *21*(1), 245. https://doi.org/10.1186/s13059-020-02134-9

Sakaluk, S. K. (1984). Male Crickets Feed Females to Ensure Complete Sperm Transfer. *Science*, *223*(4636), 609–610. https://doi.org/10.1126/science.223.4636.609

Sakaluk, S. K. (1991). Post-copulatory mate guarding in decorated crickets. *Animal Behaviour*, *41*(2), 207–216. https://doi.org/10.1016/S0003-3472(05)80472-5

Sakaluk, S. K., Duffield, K. R., Rapkin, J., Sadd, B. M., & Hunt, J. (2019). The troublesome gift: The spermatophylax as a purveyor of sexual conflict and coercion in crickets. In

M. Naguib, L. Barrett, S. D. Healy, J. Podos, L. W. Simmons, & M. Zuk (Eds.),

Advances in the Study of Behavior. Volume 51 (pp. 1–30). Academic Press.

https://doi.org/10.1016/bs.asb.2018.12.001

Sakaluk, S. K., Schaus, J. M., Eggert, A., Snedden, W. A., & Brady, P. L. (2002).

POLYANDRY AND FITNESS OF OFFSPRING REARED UNDER VARYING

NUTRITIONAL STRESS IN DECORATED CRICKETS. *Evolution*, *56*(10), 1999–

2007. https://doi.org/10.1111/j.0014-3820.2002.tb00126.x

Sayers, E. W., Bolton, E. E., Brister, J. R., Canese, K., Chan, J., Comeau, D. C., Connor, R.,

Funk, K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop,

S., Lu, Z., Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., … Sherry, S. T.

(2022). Database resources of the national center for biotechnology information.

Nucleic Acids Research, *50*(D1), D20–D26. https://doi.org/10.1093/nar/gkab1112

Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: A Cross-Platform and Ultrafast Toolkit for

FASTA/Q File Manipulation. *PLOS ONE*, *11*(10), e0163962.

https://doi.org/10.1371/journal.pone.0163962

Shumate, A., Wong, B., Pertea, G., & Pertea, M. (2022). Improved transcriptome assembly

using a hybrid of long and short reads with StringTie. *PLOS Computational Biology*,

18(6), e1009730. https://doi.org/10.1371/journal.pcbi.1009730

Simmons, L. W., Tan, Y.-F., & Millar, A. H. (2013). Sperm and seminal fluid proteomes of

the field cricket eleogryllus oceanicus: Identification of novel proteins transferred to

females at mating. *Insect Molecular Biology*, *22*(1), 115–130.

https://doi.org/10.1111/imb.12007

Sinha, P., Singh, V. K., Bohra, A., Kumar, A., Reif, J. C., & Varshney, R. K. (2021). Genomics

and breeding innovations for enhancing genetic gain for climate resilience and nutrition

traits. *Theoretical and Applied Genetics*, *134*(6), 1829–1843.

https://doi.org/10.1007/s00122-021-03847-6

Song, H., Béthoux, O., Shin, S., Donath, A., Letsch, H., Liu, S., McKenna, D. D., Meng, G.,

Misof, B., Podsiadlowski, L., Zhou, X., Wipfler, B., & Simon, S. (2020). Phylogenomic

analysis sheds light on the evolutionary pathways towards acoustic communication in

Orthoptera. *Nature Communications*, *11*(1), 4939. https://doi.org/10.1038/s41467-020-

18739-4

Szrajer, S., Gray, D., & Ylla, G. (2024). The genome assembly and annotation of the cricket

Gryllus longicercus. *Scientific Data*, *11*(1), 708. https://doi.org/10.1038/s41597-024-

03554-z

Tarailo-Graovac, M., & Chen, N. (2009). Using RepeatMasker to identify repetitive elements
in genomic sequences. *Current Protocols in Bioinformatics, Chapter 4*, 4.10.1-4.10.14.
https://doi.org/10.1002/0471250953.bi0410s25

Treidel, L. A., Goswami, P., & Williams, C. M. (2023). Changes in mitochondrial function
parallel life history transitions between flight and reproduction in wing polymorphic
field crickets. *American Journal of Physiology-Regulatory, Integrative and
Comparative Physiology*, *324*(6), R735–R746.
https://doi.org/10.1152/ajpregu.00191.2022

Uliano-Silva, M., Ferreira, J. G. R. N., Krasheninnikova, K., Blaxter, M., Mieszkowska, N.,
Hall, N., Holland, P., Durbin, R., Richards, T., Kersey, P., Hollingsworth, P., Wilson,
W., Twyford, A., Gaya, E., Lawniczak, M., Lewis, O., Broad, G., Martin, F., Hart,
M., … Darwin Tree of Life Consortium. (2023). MitoHiFi: A python pipeline for
mitochondrial genome assembly from PacBio high fidelity reads. *BMC Bioinformatics*,
24(1), 288. https://doi.org/10.1186/s12859-023-05385-y

van Huis, A., & Oonincx, D. G. A. B. (2017). The environmental sustainability of insects as
food and feed. A review. *Agronomy for Sustainable Development*, *37*(5), 43.
https://doi.org/10.1007/s13593-017-0452-8

Vasimuddin, Md., Misra, S., Li, H., & Aluru, S. (2019). Efficient Architecture-Aware
Acceleration of BWA-MEM for Multicore Systems. *2019 IEEE International Parallel
and Distributed Processing Symposium (IPDPS)*, 314–324.
https://doi.org/10.1109/IPDPS.2019.00041

Wang, Y., Jia, L., Tian, G., Dong, Y., Zhang, X., Zhou, Z., Luo, X., Li, Y., & Yao, W. (2023).
shinyCircos-V2.0: Leveraging the creation of Circos plot with enhanced usability and
advanced features. *iMeta*, *2*(2), e109. https://doi.org/10.1002/imt2.109

Wang, Y.-L., Wang, Y.-J., Luan, J.-B., Yan, G.-H., Liu, S.-S., & Wang, X.-W. (2013). Analysis
of the transcriptional differences between indigenous and invasive whiteflies reveals
possible mechanisms of whitefly invasion. *PLoS ONE*, *8*(5), e62176.
https://doi.org/10.1371/journal.pone.0062176

Watanabe, T., Noji, S., & Mito, T. (2017). Genome editing in the cricket, *Gryllus bimaculatus*.
Methods in Molecular Biology, 1630, 219–233. https://doi.org/10.1007/978-1-4939-
7128-2_18

Weissman, D. B., Gray, D. A., Pham, H. T., & Tijssen, P. (2012). Billions and billions sold:
Pet-feeder crickets (Orthoptera: Gryllidae), commercial cricket farms, an epizootic

densovirus, and government regulations make for a potential disaster. *Zootaxa*, *3504*(1),

Article 1. https://doi.org/10.11646/zootaxa.3504.1.3

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund,

G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S.

M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019).

Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686.

https://doi.org/10.21105/joss.01686

Xu, M. & Shaw, K. L. (2019). Genetic coupling of signal and preference facilitates sexual

isolation during rapid speciation. *Proceedings of the Royal Society of London, B 286*

(1913), 20191607. https://doi.org/10.1098/rspb.2019.1607

Xu, M., & Shaw, K. L. (2021). Extensive Linkage and Genetic Coupling of Song and

Preference Loci Underlying Rapid Speciation in Laupala Crickets. *Journal of Heredity*,

112(2), 204–213. https://doi.org/10.1093/jhered/esab001

Yang, J., Dong, H., He, M., & Gao, J. (2021). Mitochondrial genome characterization of

Gryllodes sigillatus (Orthoptera: Gryllidae) and its phylogenetic implications.

Mitochondrial DNA Part B, *6*(3), 1056–1058.

https://doi.org/10.1080/23802359.2021.1899078

Yang, Y., Xu, S., Xu, J., Guo, Y., & Yang, G. (2014). Adaptive Evolution of Mitochondrial

Energy Metabolism Genes Associated with Increased Energy Demand in Flying Insects.

PLOS ONE, *9*(6), e99120. https://doi.org/10.1371/journal.pone.0099120

Ylla, G., Nakamura, T., Itoh, T., Kajitani, R., Toyoda, A., Tomonari, S., Bando, T., Ishimaru,

Y., Watanabe, T., Fuketa, M., Matsuoka, Y., Barnett, A. A., Noji, S., Mito, T., &

Extavour, C. G. (2021). Insights into the genomic evolution of insects from cricket

genomes. *Communications Biology*, *4*(1), Article 1. https://doi.org/10.1038/s42003-

021-02197-9

You, J.-X., Li, N., Ren, B.-Z., Yuan, H.-B., & Shang, L.-N. (2007). The study on karyotypes

of five Grylloidea species (Orthoptera: Grylloidea) in Northeast China. *Zootaxa*,

1611(1), Article 1. https://doi.org/10.11646/zootaxa.1611.1.5

Yusuf, L. H., Pascoal, S., Moran, P. A., & Bailey, N. W. (2024). Testing the genomic overlap

between intraspecific mating traits and interspecific mating barriers. *Evolution Letters*,

8(6), 902–915. https://doi.org/10.1093/evlett/qrae042

Zhang, X., Rayner, J. G., Blaxter, M., & Bailey, N. W. (2021) Rapid parallel adaptation despite

gene flow in silent crickets. *Nature Communications 12*(1), 50.

https://doi.org/10.1038/s41467-020-20263-4.

Zhang, X., Blaxter, M., Wood, J. M. D., Tracey, A., McCarthy, S., Thorpe, P., Rayner, J. G.,

Zhang, S., Sikkink, K. L., Balenger, S. L., & Bailey, N. W. (2024). Temporal genomics

in Hawaiian crickets reveals compensatory intragenomic coadaptation during adaptive

evolution. *Nature Communications*, *15*(1), 5001. https://doi.org/10.1038/s41467-024-

49344-4

Zhou, C., McCarthy, S. A., & Durbin, R. (2023). YaHS: Yet another Hi-C scaffolding tool.

Bioinformatics, *39*(1), btac808. https://doi.org/10.1093/bioinformatics/btac808

outer

GC content
Gene density

inner

Repeats density