RNAseq sheds light on "who is doing what" in the coral *Porites lutea*

Kshitij Tandon[1,2,3*], Juntong Hu[1], Francesco Ricci[4,5], Linda Louise Blackall[1], Mónica Medina[6],
Michael Kühl[7], Heroen Verbruggen[1,2,8]

[1]School of BioSciences, The University of Melbourne, Melbourne, VIC 3010, Australia
[2]Melbourne Integrative Genomics, The University of Melbourne, Melbourne, VIC 3010,
Australia
[3]Department of Microbiology and Immunology at the Peter Doherty Institute of Infection and
Immunity, The University of Melbourne, Melbourne, VIC 3010, Australia
[4]Department of Microbiology, Biomedicine Discovery Institute, Monash University, Clayton,
VIC 3800, Australia
[5]Securing Antarctica's Environmental Future, Monash University, Clayton, VIC 3800,
Australia
[6]Department of Biology, Pennsylvania State University, University Park, PA 16802, USA
[7]Marine Biological Section, Department of Biology, University of Copenhagen, DK-3000
Helsingør, Denmark
[8]CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório
Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

*Corresponding author: Kshitij Tandon (k.tandon@unimelb.edu.au)

Abstract

Global decline of coral reefs due to climate change necessitates nature-based protection
strategies for these crucial ecosystems. Developing such strategies requires a thorough
understanding of the complex roles and interactions occurring within the coral holobiont. Using
RNAseq, we investigated the active microbiome of healthy stony coral *Porites lutea*, focusing
on the coral tissue, the green endolithic algal layer (*Ostreobiu*m layer), and the deeper coral
skeleton. We identified distinct, metabolically active communities within these compartments
and highlight substantial metabolic redundancy across carbon, nitrogen, and sulphur
pathways. Our study provides first transcriptomic evidence of *Ostreobium's* ability to transfer
fixed carbon to other holobiont members and the coral host. Additionally, we highlight critical
roles of diverse coral holobiont members in nutrient cycling and maintaining homeostasis
through scavenging of reactive oxygen and nitrogen species. This study provides novel
molecular-level understanding of the functional roles played by diverse coral holobiont
members in their respective compartments and underscores that corals harbour distinct
microbiomes with wide-ranging functions.

Introduction

Coral reefs owe their success to the symbiosis between reef-building, stony corals (Scleractinia) (*1–3*) and a large diversity of closely associated bacteria, archaea, viruses, and microeukaryotes, collectively termed the coral holobiont (*3–6*). These symbionts *sensu lato* encompass organisms spanning the full spectrum of host-microbe interactions, from mutualistic partners that enhance coral fitness to opportunistic pathogens, as well as commensals and other microorganisms whose ecological roles may shift along this continuum depending on environmental conditions. While the most well-characterized symbiosis in corals involves photosynthetic, endosymbiotic microalgae in the family Symbiodiniaceae residing in the gastrodermal tissues of the animal host (*7–9*), a compendium of prokaryotes associated with corals has identified 39 bacterial and 2 archaeal phyla (*10*). This diverse microbial assemblage is not only distinct between different compartments within the coral (*11*), but fine-scale mapping of the coral skeleton has revealed a stratified community across its depth (*12–14*). In recent years, other microeukaryotes besides Symbiodiniaceae, including endolithic algae of the genus *Ostreobium*, fungi, and apicomplexans have also garnered attention (*15, 16*).

The diversity and composition of coral holobiont members have been investigated through metabarcoding of ribosomal marker genes (16S rRNA, 18S rRNA, ITS2) (*17–19*) (*20, 21*), providing invaluable insights into the diverse and structured symbiont composition *sensu lato* within coral compartments (*12, 13, 22, 23*). Combined culture-based and molecular approaches have expanded our understanding of the functions of coral holobiont members in maintaining coral health and resilience (*24–28*). Symbiodiniaceae, in particular, play a crucial role by supplying corals with photosynthates, meeting a significant portion of their energy demand (*29*). Coral-associated bacteria and archaea have been implicated in a myriad of functions essential for sustaining coral health and resilience, including stress alleviation through scavenging of reactive oxygen and nitrogen species (ROS and RNS), protection against pathogens and nutrient cycling (*24–26, 28, 30–32*). Coral skeleton-dwelling endolithic green algae of the genus *Ostreobium,* which form conspicuous green bands beneath the coral tissue, have also been suggested to support corals by providing photosynthates during coral bleaching (*33–35*), but we lack a detailed understanding of the metabolic interaction between the coral animal and *Ostreobium* (*15*).

Most of our present knowledge on the role of microbiota for coral function and health comes from amplicon and metagenomic sequencing studies, with potential functions inferred using correlative, bioinformatic analyses. Despite these advances and the extensive information now available on the gene-encoded functional potential of microbiome and microeukaryotic

members through genomic and metagenomic studies, a significant knowledge gap persists regarding actual microbial metabolic activity and its spatial organisation within the coral holobiont. Genomic data alone cannot predict the actual contributions of specific functions by symbionts *in situ*, as genes encoding these functions may be variably expressed or not expressed at all. RNAseq enables the analysis of the pool of actively expressed genes in an environmental sample, highlighting the ecophysiologically relevant processes occurring at the time of sampling (*36*). Therefore, taking a transcriptomics view can provide insights into which community members perform which functional roles.

Obtaining a molecular level understanding of the functional roles of holobiont members is crucial especially given the rapid decline of coral reefs worldwide due the adverse effects of climate change (*37*). Addressing this knowledge gap will also be beneficial for developing effective nature-based interventions to protect coral reefs, including improved assisted evolution strategies and development of coral probiotics (*2*, *38*). Both approaches rely on a comprehensive understanding of the roles played by diverse microbial symbionts in maintaining coral health.

In this study, we sought to answer the fundamental question of 'who is doing what' in a healthy coral holobiont across different compartments. Using RNAseq, we identified the active symbiont community of the coral *Porites lutea*, characterizing their specific activities and locations within the coral, from the tissue to the endolithic algal layer (Ostreobium-layer) and deeper coral skeleton. We focused on key nutrient cycling pathways, including carbon fixation (C), photosynthesis (P), nitrogen (N) and sulfur (S) metabolism, sugar export, and the scavenging of reactive oxygen and nitrogen species (ROS and RNS).

Methods:

Coral sampling and colony fragmentation Fragments from six healthy-looking, *Porites lutea* colonies, spaced at least 15 m apart, were collected at low tide (1-2 m depth) from the research zone of the Heron Island reef flat, southern Great Barrier Reef (23°44′S, 151°91′E), in October 2022. Four fragments (from colonies we numbered 1, 3, 4 and 5) were collected during daytime, while two (colonies 6 and 7) were collected five-six hours after sunset. All collected colonies exhibited a visually discernible *Ostreobium-layer* in the coral skeleton . 10-50 mm below the coral tissue layer (Fig. 1A). The fragments were collected using a sterile hammer and chisel and were placed in separate sterile zip-loc polyethylene bags in seawater and flash frozen in liquid nitrogen immediately upon collection. A 5-mm thick slice was cut from the frozen samples, perpendicular to the vertical growth axis, using a diamond saw with a continuous flow of sterile filtered (0.2 μm) seawater. Later, pliers were used to obtain

113 fragments from three distinct layers, i.e., the coral tissue, the green *Ostreobium*-layer, and the
114 tan fraction of the skeleton below the green band (termed the skeleton layer). During this
115 process, the material was kept frozen by repeated immersion in liquid nitrogen. Each layer
116 was kept frozen at -80ºC, transported to the University of Melbourne on dry ice, and processed
117 immediately upon arrival.
118

Total RNA extraction
For RNA extraction, samples of individual layers were ground to a fine powder in liquid nitrogen
using pre-cooled mortars and pestles. Approximately 200 mg of powdered tissue samples and
between 300-500 mg of powdered *Ostreobium-layer* and skeleton layer were used for RNA
extraction with 1 mL of pre-cooled TRIzol reagent (Invitrogen, USA). Total RNA was extracted
following the manufacturer's protocol with some modifications (**see supplementary file 1 for**
details). Quantification of total RNA was conducted using the Qubit RNA HS Assay kit
(Thermo Fisher Scientific, USA).
127

RNA sequencing and data-preprocessing
We did not perform any ribosomal RNA depletion due to the dual aim of sequencing mRNA
and rRNA from both eukaryotes and prokaryotes simultaneously and profiling the
transcriptionally active microbial and microeukaryotic community composition using 16S and
18S rRNA abundance (see below). Total RNA libraries were prepared for 36 samples (6
colonies x 3 layers x 2 replicates per layer) and sequenced on the Illumina NovaSeq platform
by Azenta (Suzhou, China). Sequencing yielded approximately 23 million reads per sample
with 2x150 bp reads.
136

Raw reads were quality checked and trimmed using FastQC v0.11.9
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc) and Trimmomatic v0.39(*39*),
respectively. Trimmomatic parameters were as follows: *TruSeq3-PE.fa:2:30:5 HEADCROP:5*
SLIDINGWINDOW:4:20 MINLEN:50. Quality-filtered reads were first used to identify and filter
out rRNA reads using SortMeRNA v4.3.6(*40*) with Silva v138.1 LSU and SSU (*SILVA138.1*
LSU and SSU NR99 database, last accessed Feb 10, 2023)(*41*). Reads identified as rRNA
were taxonomically classified using PhyloFlash v3.4(*42*) (with *almost-everything*), which
rapidly reconstructs the SSU rRNAs (NTUs, nearest taxonomic units) and taxonomically
annotates them using Silva 138.1 NR99 SSU database. Reads identified as rRNA were
removed from the raw-read set, and non-rRNA reads were mapped to the *Porites lutea*
genome (accessed from www.reefgenomics.org) using bowtie2 v2.3.4.1(*43*) and samtools
1.17(*44*) to identify and remove host-related reads.

High-quality, non-rRNA and non-host reads were co-assembled using Trinity v2.15.1(*45*) with
parameters *--min_contig_length 250 and --no_salmon*. Taxonomic annotation of assembled
transcripts was obtained using the lowest common ancestor approach by performing *blastx*
(46) searches against the NR database (last accessed April 2023) with diamond
v2.1.6.160(*47*), using an *e-value cutoff of 1e-5* and processed through MEGAN6 community
edition v6.24.20, using the *megan-map-Feb2022-ue.db* and meganising the *daa* file using
daa2info (*48*). This process was used to identify and remove any transcripts annotated as
Scleractinia and other Eumetazoa. The remaining transcripts were quantified using RSEM
(RNA-Seq by Expectation and Maximisation)(*49*), implemented in Trinity through the utility
script *align_and_estimate_abundance*.*pl*. This approach generated both gene and isoform
expression profiles. Isoform-level expression data was used for the analysis. Expression
values per transcript were averaged across replicates for each layer (tissue, *Ostreobium*-layer
and skeleton).
162
Community composition analysis
Taxonomic classification of NTUs assembled from rRNA reads was conducted to profile the
abundance of transcriptionally active community members. NTUs classified as "Chloroplast"
or "Mitochondria" were removed. Abundance values were converted to relative abundance for
compositional analysis using barplots and log-transformed for diversity analysis using PCoA.
These analyses were performed using R packages phyloseq v1.42.0(*50*), metagMisc v0.5.0
(https://github.com/vmikk/metagMisc) and ggplot2 v34.3 (*51*). We used the quantified and
taxonomically annotated transcripts assembled from mRNA reads to profile the community
composition encoding for different functional guilds based on the transcriptome data and
compared them to rRNA based analysis using the same approach.
173
Functional analysis
To ascertain functional roles, we first filtered transcripts using Trimmed Mean of the M (TMM)
values of 0.1 and presence in at least 2 samples to identify truly expressed transcripts. This
approach filters out i) any transcripts from potential residual DNA or very low expressed
transcripts, ii) transcripts originating from spurious mapping of reads. Transcripts deemed truly
expressed were then grouped in a compartment-specific manner to obtain tissue, *Ostreobium*-
layer and skeleton-layer specific transcriptome. These compartment specific transcripts were
processed   through   Trinotate   v4.0.0(*52*)   and   Transdecoder   v4
(https://github.com/TransDecoder/TransDecoder) to identify the longest open reading frames
and predict the proteins from transcripts ≥50 amino acids. Predicted proteins were functionally
annotated using a combination of EggNOG mapper v2.1.12(*53*) with database eggnogDB

v5.0.2(*54*) with *e-value cutoff 1e-5*, ghostkoala (*55*) with database
c_family_euk+genus_prok+viruses parameters and Interproscan v5.63-95.0(*56*) with *e value*
cutoff 1e-5 to obtain a comprehensive understanding of functional orthologous groups
(EggNOG and COG), KEGG pathways, including Kegg Ontology, KOs and protein families
(pfam). Metabolic markers for carbon-fixation (C), photosynthesis (P), nitrogen (N) and sulfur
(S) metabolism were identified from a previous study (*57*). Additionally, we focussed on genes
involved in sugar export and scavenging of ROS and RNS. Expression profiles of transcripts
functionally annotated with marker KOs or pfams were plotted using heatmaps using
ComplexHeatmap(*58*) in R.
194
195 **Results and Discussion**
196
197 **Corals harbor compartment-specific active symbiont community**
198
One of our primary objectives was to profile the transcriptionally active community within the
coral holobiont and compare these findings with the community composition obtained from
rRNA analysis. To achieve this, we examined the total and active community composition
across the coral tissue, the endolithic *Ostreobium-layer*, and the underlying coral skeleton,
and compared it with their rRNA based community composition (**Fig 1A)**. For total community
composition, we taxonomically analysed 28,717 rRNA NTUs (6,010 from tissue layer, 15,942
from *Ostreobium* layer, 26,712 from skeleton layer) belonging to Archaea (2.92% - 840
sequences), Eukaryota (12.68% - 3,644 sequences), and Bacteria (84.38% - 24,233
sequences) (**Fig 1B**). This observation offers essential insights into the diversity and
composition of the coral holobiont. Unlike traditional amplification-based metabarcoding
analyses, which are often biased (*59*) and fail to provide accurate quantitative distinctions
between prokaryotes and eukaryotes, using rRNAs assembled from total RNA delivers a more
precise and comprehensive overview of the community composition. Eukaryotes (Amorphea,
SAR, Archaeplastida) were the dominant groups in the coral tissue (**Supplementary Fig S1A**)
and the *Ostreobium layer* (**Supplementary Fig S1B**), while the deeper skeleton was
dominated by Bacteria (Proteobacteria, Planctomyceota, Desulfobacteria, and Cyanobacteria)
(**Supplementary Fig S1C**). Our study shows that bacterial diversity is highest in the skeleton,
where the majority of bacterial NTU sequences were assembled, confirming previous reports
of greater bacterial richness in the skeleton compared to tissue compartments (*13*, *19*).
218
After initially evaluating NTUs based on rRNAs, we extended our analysis to the remaining
RNA sequence annotations. To profile active microbial community composition, we used a
curated set of 371,193 taxonomically annotated transcripts, selected from a total of 780,431
assembled sequences. After excluding 111,367 transcripts belonging to Scleractinia,

Chordata, and Eumetazoa, 259,826 (47.56%) transcripts remained. These transcripts
represent Eukaryotes (52.48% - 136,366), Bacteria (44.99% - 116,900), Archaea (1.97% -
5,131), and viruses (0.54% - 1,429) (**Fig 1B**) and revealed a high relative abundance of
alveolates, the eukaryotic group containing Apicomplexa and dinoflagellates, including
Symbiodiniaceae in coral tissue (**Supplementary Fig S2A)**. Green algae, Streptophyta and
Chlorophyta, dominated the *Ostreobium layer* and deeper skeleton. (**Supplementary Fig S2B**
and C). The high relative abundance of dinoflagellate (Symbiodiniaceae) expressed
transcripts in coral tissue is unsurprising, as they are the most dominant microeukaryotes in
the coral tissue, meeting most of the coral's energy demands (*60*). However, we identified
dinoflagellate and Apicomplexa transcripts in all compartments. Apicomplexans have been
found in healthy coral colonies across the geographic distribution and taxonomic diversity of
corals, but their role is yet to be characterized (*16*, *61*). Bacteria and Fungi were present in all
compartments, with bacteria as dominant members of the skeletal microbiome
(**Supplementary Figure S2C)**. Proteobacteria, Cyanobacteria and Actinobacteria were the
dominant bacterial phyla in all coral compartments. Members of the phylum Proteobacteria
are some of the most widespread coral microbiome members (*10*, *17*, *62*). Cyanobacteria
have also previously been identified as the dominant prokaryote group in the *Ostreobium layer*
and deeper skeleton of *P. lutea* from the Red Sea (*63*). Obtaining annotation for less than 50%
of the total assembled transcripts highlights the immense diversity of uncharacterised
symbionts associated with *P. lutea*, despite it being one of the best investigated coral species
with several metagenomic and physiological studies (*24*, *25*, *64*, *65*). These findings align with
earlier studies using the metatranscriptomics approach to profile transcriptionally active
microbiomes in other ecologically diverse systems, including soil (*66–68*), sponge (*69*) and
ocean microbiomes (*57*, *70*).

Fig. 1. Taxonomic and functional overview based on rRNA and transcriptome data A) illustration of a coral fragment depicting the three compartments: Tissue, Ostreobium-layer and Skeleton. B) Bar plots showing overall community composition based on rRNA and transcriptome data. C) UpSet plot illustrating shared and unique transcripts across different coral compartments. D) Venn diagram showing counts and percentage of transcripts which are deemed expressed, expressed and taxonomically annotated, annotated at KO and at OG levels.

Functional annotation of transcripts remains a bottleneck

Analysis of active metabolic pathways in a healthy coral holobiont revealed 261,884 truly expressed transcripts through TMM filtering. These transcripts were distributed across the three investigated compartments, with the highest overlap observed between the *Ostreobium layer* and deeper skeleton (89,682 shared transcripts). Additionally, a substantial number of uniquely expressed transcripts were identified in the deeper skeleton-layer (48,966), while the coral tissue exhibited the lowest count of uniquely expressed transcripts (**Fig. 1C**). Recent metabarcoding and metagenomic studies have highlighted the skeleton as the most diverse compartment of corals (*13*, *19*, *63*). In line with these observations, our coral skeleton samples

revealed 178,580 uniquely predicted proteins from the expressed transcripts. However, only 30.64% (54,731) could be annotated using KO ids and 29.25% (52,252) using OGs (**Fig 1D**). The extraordinary uniqueness of skeletal proteins underscores the added value of sampling genomically underexplored environments, such as the coral skeleton. The low functional annotation observed in our dataset is consistent with a recent study on gene expression changes and community turnover of global oceans, which similarly found less than 25% of the transcripts assigned to KOs (*57*). Annotations based on COG revealed a notable proportion of proteins encoded by transcripts involved in signal transduction, intracellular trafficking, secretion, vesicular transport, post-translational modification, protein turnover, and chaperones (**Supplementary Figure S3)**. While most transcripts remained unannotated, the presence of these functional categories suggests potential microbial involvement in cellular communication and protein homeostasis within the coral holobiont. Although our data do not directly demonstrate nutrient or metabolite exchange, these functional signals are consistent with prior hypotheses that such microbial activities may support coral resilience, particularly in oligotrophic environments. Further targeted studies, including isotopic tracing or compartment-specific transcriptomics, will be needed to directly test these functional roles.

Distinct photosymbionts drive carbon fixation in coral compartments

To reveal the functional roles of transcriptionally active holobiont members in healthy *P. lutea* coral, we identified key microbial processes, such as photosynthesis, carbon fixation, sugar export, and nitrogen and sulphur cycling. Our analysis of pathway-specific metabolic markers highlighted the taxonomic affiliations of transcripts and their expression profiles, shedding light on the active contributions of each microbial group to these essential processes. Among the most highly expressed genes were those associated with photosynthetic carbon fixation, including photosystem I (PsaA, 59 transcripts), photosystem II (PsbA, 67 transcripts) and the Calvin–Benson cycle enzyme, RuBisCO subunits (ribulose-1,5-bisphosphate carboxylase/oxygenase, *rbc*L (79) and *rbc*S (18) genes) (**Fig 2**). While rbcL can be expressed by chemotrophs (*71*), our taxonomic annotations did not identify chemotrophic affiliations, suggesting that carbon fixation in this context is predominantly photoautotrophic. These genes were actively transcribed by Symbiodiniaceae, stramenopiles, Bryopsidales (including the endolithic green alga *Ostreobium*), Cyanobacteria, Gammaproteobacteria, and unclassified Viridiplantae and Chlorophyta in different coral compartments. Symbiodiniaceae dominated expression in the tissue, Cyanobacteria were active throughout all compartments, and *Ostreobium* and other Ulvophyceae were the main contributors in the deeper skeleton layers (**Supplementary data file 2**). The co-expression of *rbc*L/*rbc*S with *psb*A and *psa*A supports the role of chlorophototrophy as the dominant carbon fixation strategy in the coral holobiont (*72–74*).

In the skeleton, we also detected transcripts indicative of alternative carbon fixation pathways.

The expression of *cox*L, *cox*M, and *cox*S, affiliated with Cyanobacteria and Proteobacteria,

suggests potential activity of carbon monoxide dehydrogenase (CODH), enabling CO

oxidation and energy generation through chemosynthesis. Additionally, transcripts for *acs* and

*acs*B, encoding the acetyl-CoA synthase complex of the Wood–Ljungdahl pathway, were

expressed in the skeleton and affiliated with anaerobic bacteria. This pathway enables fixation

of CO_2 and CO into acetyl-CoA, a central metabolite in microbial energy and biosynthetic

pathways (*63, 75*). These transcriptional signals reflect the metabolic versatility of skeleton-

associated microbial communities and suggest adaptation to low-oxygen, possibly CO-rich

microenvironments (*14, 76*). Although we did not directly measure substrate availability or

fluxes, the coexistence of phototrophic and anaerobic carbon fixation pathways implies that

microbial communities play diverse roles in internal carbon cycling. These findings highlight

the functional potential of coral-associated microbes to persist in spatially heterogeneous,

nutrient-poor reef environments; however, further experimental work is needed to determine

the direct impacts of these processes on coral physiology or holobiont productivity.

Besides *psa*A and *psb*A, we detected expression of several other photosynthetic marker

genes across all three coral holobiont compartments. These included, *petC* , which encodes

the Rieske iron–sulfur protein of the cytochrome b6f complex that links Photosystem II to

Photosystem I, and *petE* ncoding plastocyanin, which facilitates electron transfer between the

cytochrome b6f complex and Photosystem I. The expression of these genes supports the

widespread presence of oxygenic phototrophs throughout the coral tissue, *Ostreobium* layer,

and skeleton. transcripts for cyanobacteria-specific antenna proteins—including *apcA, apcF,*

cpcA, cpeA and *cpeT*—were most abundant in the Ostreobium layer and deeper skeleton,

suggesting a greater prevalence and transcriptional activity of Cyanobacteria in these

compartments compared to the tissue (**Fig 2**). This pattern is consistent with taxonomic

assignments based on both rRNA (**Supplementary Fig S1**) and transcriptome levels

(Supplementary Fig S2) and aligns with previous findings showing an enrichment of

cyanobacterial and eukaryotic phototrophs in the skeletal layers of *P. lutea* These antenna

and photosynthetic genes, many of which are involved in the Calvin–Benson cycle, collectively

accounted for more than 50% of the total carbon fixation gene transcripts in the skeleton. We

also observed expression of *pufM*, a marker gene for anoxygenic photosynthesis that encodes

a core component of the bacterial photosynthetic reaction centre. Expression of *pufM* was

confined to the skeleton, suggesting the presence of obligate anaerobes, such as green sulfur

and purple non-sulfur bacteria. This observation is in agreement with hyperspectral pigment

imaging, which revealed pigment signatures consistent with anoxygenic phototrophs in the
skeletal layers of *P. lutea* (*14*).

The observed compartment-specific expression patterns of photosynthetic genes reflect
microbial adaptations to distinct light regimes within the coral holobiont and provide insight
into the metabolic plasticity of associated phototrophs. The elevated expression of
cyanobacterial antenna protein genes (*apcA*, *apcF*, *cpcA*, *cpeA*, *cpeT*) in the Ostreobium
layer and deeper skeleton suggests that Cyanobacteria inhabiting these low-light regions may
utilize alternative chlorophyll pigments, such as chlorophyll *d* and *f*, to absorb far-red light that
penetrates coral tissue. This strategy likely enhances photosynthetic efficiency under reduced
irradiance and aligns with previous reports of far-red light utilization by endolithic
Cyanobacteria and *Ostreobium* (*77–79*).

Within the same skeletal microhabitat, the green alga Ostreobium appears to employ
complementary adaptations, including the use of chlorophyll b and an expanded suite of light-
harvesting complex (LHC) proteins in both photosystems (*15*). Genomic features such as
Lhca1 gene duplication and the loss of photoprotective genes further reflect specialization to
persist under light-limited conditions (80). In contrast, the tissue compartment, characterized
by higher light availability, showed dominant transcriptional activity from Symbiodiniaceae and
other phototrophs known to rely on chlorophyll *a* and *c*, consistent with adaptation to higher
irradiance (81). The distinct expression of photosynthetic genes and pigment-related functions
across coral compartments emphasizes the niche differentiation and photoadaptive strategies
among coral-associated phototrophs. These spatial patterns of gene expression illustrate the
functional diversity of the holobiont and suggest mechanisms by which microbial communities
could contribute to energy capture across microhabitats with steep light gradients.

Not only Symbiodiniaceae provide fixed carbon to the coral holobiont
To identify members of the coral holobiont involved in the export of photosynthetically fixed
carbon compounds (photosynthates), we focused on genes encoding SWEET ("Sugars Will
Eventually be Exported Transporter") proteins. SWEETs are bidirectional sugar uniporters that
facilitate the diffusion of sugars such as glucose along concentration gradients across
biological membranes (*82, 83* We identified transcripts encoding SWEET proteins, specifically
associated with Pfam domains PF04193 and PF03083, across all three major compartments
of the coral holobiont (**Fig 2**). Taxonomic annotation of these transcripts revealed that sugar
exporters were predominantly eukaryotic, including Symbiodiniaceae, the endolithic
alga *Ostreobium*, and unclassified green algae within the Chlorophyta (**Supplementary data**
file 2). These organisms share a common capacity for oxygenic photosynthesis via the Calvin

cycle, in which glyceraldehyde-3-phosphate (G3P), serves as a precursor for sugar

biosynthesis, including glucose. Glucose is likely the principal carbohydrate exported via

SWEET proteins (*84*), suggesting that these eukaryotic phototrophs contribute substantially

to the pool of photosynthates available within the coral holobiont.

Interestingly, despite previous studies reporting sugar transporters in coral-associated

bacteria (*26*), we did not detect transcripts encoding semiSWEET proteins, the prokaryotic

homologs of SWEETs. This absence may reflect a biological partitioning of function; wherein

eukaryotic symbionts serve as primary carbohydrate exporters. Alternatively, it could be a

result of detection limits or the divergence of bacterial sugar transporters from characterized

semiSWEETs. Regardless, our findings highlight a potentially central role for eukaryotes in

sugar-based metabolic exchange among holobiont members.

Notably, we identified SWEET gene expression in *Ostreobium*, providing the first genomic and

transcriptomic evidence that this skeleton-dwelling alga has the capacity to export sugars.

These sugars may diffuse into the coral skeleton, potentially serving as an energy source for

both the coral host and other microbial symbionts. This finding provides molecular support for

prior radiolabeling studies using 14C and 13C-labeled photoassimilates , which demonstrated

that fixed carbon from endolithic algae can be translocated into coral host lipids under both

healthy and bleached conditions (*34*, *35*). The ability of *Ostreobium* to contribute sugars under

thermal stress, when Symbiodiniaceae populations are diminished, suggests a compensatory

role in mitigating coral energy deficits (*85*). This metabolic capacity may be especially

important during bleaching events, offering a survival advantage by partially offsetting the loss

of primary photosynthate sources.

In addition to phototrophs, we also detected the expression of SWEET-encoding transcripts

in coral-associated fungi. This is the first direct molecular evidence of fungal participation in

sugar transport within the coral holobiont. While the functional implications remain to be fully

resolved, fungi are increasingly recognized for their roles in shaping coral-associated

microbiomes and reef-scale biogeochemical cycling through nutrient processing and chemical

interactions (*86*). Their involvement in fixed carbon exchange may reflect metabolic cross-

feeding, carbon recycling, or interactions with other holobiont members, broadening our

understanding of fungal contributions to holobiont function.

The identification of multiple eukaryotic lineages capable of translocating sugars underscores

the complex and cooperative nature of carbon cycling within the coral holobiont. These sugar-

exporting symbionts likely play critical roles in sustaining coral energy homeostasis. However,

this capability may also pose risks: elevated extracellular sugar concentrations could
unintentionally fuel the proliferation of opportunistic or pathogenic bacteria, potentially
destabilize the microbial community and compromise coral health. Thus, while eukaryotic
sugar exporters contribute to coral resilience, their activity may also influence disease
susceptibility, highlighting the need for further investigation into the ecological and evolutionary
consequences of carbon transfer dynamics in the coral holobiont.
419

420
421
Fig. 2. Carbon Fixation, Photosynthesis and Sugar Export Across Coral Compartments.
Heatmap displaying the expression of marker genes for carbon fixation photosynthesis, and
SWEET protein-mediated sugar export across various coral compartments. Expression values
for both were averaged for the count of transcripts and log2(expression + 1) transformed.

426
427
Skeleton dwelling microbes mediate nitrogen metabolism

Our analysis revealed active nitrogen metabolism within the skeleton of *P. lutea*, highlighting
the role of skeleton-associated microbes in sustaining nitrogen cycling within the coral
holobiont. Expression of canonical nitrogen fixation marker genes nifK, nifD and nifH was
detected in both the *Ostreobium* layer and the deeper skeleton, suggesting active nitrogen
fixation in these predominantly anoxic microenvironments (**Fig. 3A**). Nitrogen cycling is
essential in coral reef ecosystems, where nutrient scarcity demands tight microbial recycling

to support coral growth and productivity (*31*). While nitrogen-cycling microbes are widely
documented in coral-associated microbiomes (*31*, *87*), our study identified Cyanobacteria as
the predominant taxonomically resolved nitrogen fixers in *P. lutea*. This contrasts with earlier
reports emphasizing heterotrophic nitrogen-fixing bacteria (*88–90*) and instead aligns with
recent genome-resolved metagenomic studies showing low diversity of nitrogen-fixing taxa in
this species (*24*, *25*). It is important to note that a substantial proportion of transcripts remain
unclassified, and additional nitrogen-fixing taxa may be present but undetected. Nonetheless,
the predominance of cyanobacterial transcripts underscores their likely importance in
supporting nitrogen-limited growth of Symbiodiniaceae (*9*, *91*), and highlights their key role in
the coral holobiont's nitrogen economy.

We also identified nitrification, the aerobic microbial process converting ammonia to
hydroxylamine and subsequently to nitrite and nitrate, to be ubiquitous across all coral
compartments (**Fig 3A**). This process is vital for nitrogen turnover and ammonium regulation
, reducing toxicity, and maintaining nitrogen turnover within the holobiont, with implications for
both coral host and symbiont physiology (*31*). Transcripts encoding nitrification-related genes
were primarily assigned to Thaumarchaeota, Gammaproteobacteria, and Chloroflexi,
particularly in the deeper skeleton (**Fig 3B and Fig 5**). These taxa likely contribute to ammonia
oxidation, although unclassified transcripts and incomplete database coverage suggest that
additional nitrifying microbes may be involved.

In contrast, denitrification was spatially restricted to the *Ostreobium* layer and deeper skeleton
compartments, consistent with oxygen-limited conditions in these compartments. We detected
expression of key denitrification marker genes (*napA, nirS, norB, and nosZ*) in both eukaryotic
phototrophs (Symbiodiniaceae, Ostreobium) and bacterial symbionts, including
Rhodospirillales, Chlorobi, Proteobacteria, and Acidobacteria (**Fig. 3A**; **supplementary data
file 3**). The subsequent steps in denitrification, conversion of nitrite to nitric oxide, nitrous
oxide, and nitrogen during denitrification were most commonly attributed to Thaumarchaeota,
and Proteobacteria (α- and γ-), and Bacteroidota. Importantly, we also detected fungal
(Ascomycota) transcripts encoding NIT-6 gene, involved in dissimilatory nitrate reduction to
ammonium (DNRA), providing evidence for fungal contributions to nitrogen recycling within
the holobiont.

The interplay between nitrification, denitrification, and DNRA, particularly within the skeleton,
appears crucial for maintaining nitrogen homeostasis. DNRA may dominate under low-oxygen
conditions (e.g., at night), recycling nitrate into ammonium when aerobic nitrification is less
active. This ensures stable ammonium availability, potentially supporting coral growth. In

contrast, coupling of nitrification and denitrification may help regulate nitrogen levels under
nutrient-rich conditions, preventing excessive proliferation of Symbiodiniaceae and
maintaining a balanced holobiont metabolism.
476
Overall, our results expand current understanding of nitrogen cycling in coral holobionts by
resolving the microbial taxa actively involved and mapping their compartment-specific
activities. The coral skeleton emerges as a metabolically active zone for both aerobic and
anaerobic nitrogen transformations, with stratified microbial communities performing
complementary functions. These findings underscore the importance of skeleton-associated
microbes in holobiont nutrient dynamics and reef ecosystem functioning.
483
Sulphur metabolising microbes are widespread across coral compartments
Sulfur, abundant in seawater primarily as inorganic sulfate, is assimilated by the coral
host, Symbiodiniaceae, and various members of the coral microbiome into essential organic
compounds, making it a critical element for holobiont growth and function (*92–94*). In this
study, we identified expression of transcripts encoding sulfur cycling marker genes across all
three coral compartments (Fig. 3B), underscoring the distributed nature of sulfur metabolism
within the holobiont. Notably, transcript abundance for most sulfur-related genes was highest
in the skeleton, indicating a prominent role for the endolithic microbiome in coral sulfur cycling.
492
We detected expression of anaerobic dissimilatory sulfate reduction (DSR) marker genes,
aprA and aprB, exclusively in the skeleton compartment. Taxonomic annotation identified
these transcripts as originating from canonical sulfate-reducing bacteria (SRB),
including Deltaproteobacteria, as well as anoxygenic phototrophic Chlorobi (Fig. 3B).
Although Chlorobi carry the aprAB genes, they typically participate in sulfur oxidation rather
than sulfate reduction (95, 96), and their presence here suggests diverse sulfur redox
processes within the skeleton. This observation supports prior metabarcoding and
metagenomic studies, which reported abundant SRBs and assembled their genomes from the
skeleton of *P. lutea* and *I. palifera* (24). The co-detection of SRBs and Chlorobi within the
same compartment aligns with previous reports of anoxic microniches in coral skeletons (*14*,
24, *27*, *76*) and suggests the presence of syntrophic interactions, wherein SRB-produced
sulfide serves as an electron donor for phototrophic Chlorobi. Such metabolic coupling has
been observed in microbial mats, stratified lake waters, and coral skeletons (*27*, *97–99*). The
simultaneous expression of aprAB genes by both groups in this study provides molecular
evidence for this syntrophy within the coral holobiont.
508
We also identified a complex and spatially distributed expression pattern of **assimilatory**

sulfate reduction (ASR) genes. Transcripts for cysC, cysI, cysJ, and SUOX were detected
across all compartments, whereas cysD expression was restricted to the deeper skeleton (Fig.
3B). The gene encoding cysK, which links assimilatory sulfate reduction to L-cysteine
biosynthesis, was consistently expressed in all compartments. These findings suggest
widespread ASR activity throughout the holobiont, supporting sulfur assimilation and amino
acid biosynthesis. Taxonomic analysis of ASR gene transcripts revealed a diverse array
of eukaryotic and prokaryotic symbionts. These included microeukaryotes such
as Symbiodiniaceae, *Ostreobium*, and other Chlorophyta, as well as bacteria
including Cyanobacteria, Thaumarchaeota, and Proteobacteria (Fig. 3B). Notably, all ASR
transcripts were exclusively assigned to algae and bacteria, consistent with the known
distribution of this pathway among plants, algae, and microbes(*100*).
521

Together, these results provide new insights into the compartmentalized and microbially
mediated sulfur cycling processes within *P. lutea*. The coral skeleton emerges as a key site
for both dissimilatory and assimilatory sulfur transformations, driven by specialized microbial
consortia. The detection of potential syntrophic sulfur interactions between SRBs and Chlorobi
adds a new layer of complexity to our understanding of the coral skeleton as a metabolically
active niche, reinforcing its contribution to holobiont nutrient cycling and resilience.
528

Fig 3. Transcriptome- based expression profiles of Nitrogen and Sulfur metabolism across coral compartments. A) Heatmap showing the expression profiles of marker genes involved in nitrogen metabolism across coral compartments, along with a bar plot indicating the count of transcripts. The nitrogen cycle is illustrated, highlighting subpathways and contributing community members. B) Heatmap showing the expression profiles of marker genes involved in sulphur metabolism across coral compartments, along with a bar plot

536 indicating the count of transcripts. The sulphur metabolism pathway is illustrated, highlighting
537 subpathways and contributing community members. Expression values for both were
538 averaged and log2(expression + 1) transformed.
539
540

541
542 **Figure 5.** Overview of the transcriptionally dominant taxa and their functional contribution in
543 different compartments of the coral identified in this study.
544
545

Microeukaryotes are the major scavengers of ROS and RNS in *P. lutea*

Coral reefs are facing unprecedented threats due to rising sea surface temperatures, resulting

in more frequent and prolonged coral bleaching events (*37*). Most theories on the onset of

coral bleaching suggest that the overproduction of reactive oxygen species (ROS) and

reactive nitrogen species (RNS) is a key trigger, leading to cellular damage and stress under

adverse environmental conditions (*101*). Rapid scavenging of ROS and RNS is thus crucial in

maintaining homeostasis and protecting the coral holobiont in stressed and healthy corals. We

observed widespread expression of seven protein families with functions related to ROS or

RNS scavenging across the tissue, *Ostreobium* layer, and deeper skeleton compartments.

Specifically, transcripts encoding for genes involved in RNS scavenging, including nitric oxide

reductases (PF00175: *hmp*; PF00115 and PF00034: *nor*BC) and peroxynitrite to nitrate

reduction (PF10417: *ahp*C) were expressed in all compartments (**Fig 4**). A single transcript

encoding nitrous oxide reductase (PF18764: *nos*Z) was identified and expressed exclusively

in deeper skeleton samples. This enzyme catalyses the last step of the dissimilatory metabolic

pathway of denitrifying bacteria, which were abundant in the deeper skeleton. Similar

observations were made for ROS scavenging expression profiles. We identified 48, 43, and

41 transcripts encoding the general ROS scavenger Glutaredoxin (PF00462), as well as

hydrogen peroxide scavengers Peroxiredoxins (PF00578) and peroxidases (PF00141). These

transcripts were expressed in all coral compartments along with transcripts encoding

superoxide dismutases (PF00081 and PF02777: SOD Mn and PF00080:SOD cu-zn) (**Fig 4**).

The majority of ROS and RNS scavenging transcripts were of micro-eukaryotic origin (76%),

including Symbiodiniaceae, *Ostreobium*, Chlorophyta, and Fungi, while others were from

archaeal and bacterial sources (24%), encompassing Thaumarchaeota, Proteobacteria

(Rhodospirillales, Xanthomonadales), Cyanobacteria, Chlorobi, Chloroflexi and Bacteroidetes

(*Candidatus* Amoebophilus) (**Supplementary data file 4**).

Scavenging reactive oxygen (ROS) and nitrogen species (RNS) is a key function of bacterial

candidates for coral probiotics, as emphasised in recent studies identifying potential probiotic

bacteria for corals (*2, 26, 102–104*). However, the identification of the majority of ROS and

RNS scavenging transcripts in micro-eukaryotic inhabitants of the coral holobiont rather than

in bacteria, raises the question of which organisms are the main players in scavenging ROS

and RNS under thermal stress. This finding suggests that current strategies for designing coral

probiotics, which primarily focus on bacterial functions, should be refined through additional

molecular and experimental validation to provide functional insights. These insights could

guide the assembly of a more comprehensive consortium of beneficial symbionts for corals.

Fig. 4. Transcriptome-based expression profiles of Reactive Oxygen Species (ROS) and Reactive Nitrogen Species (RNS) marker genes across coral compartments. Heatmap displaying the expression of different protein families, count of transcripts for each family involved in ROS and RNS scavenging across coral compartments. Expression values for both were averaged and log2(expression + 1) transformed.

Conclusion

This study provides novel insights into the transcriptionally active communities within the coral holobiont, with a focus on their functional roles across distinct ecological compartments: coral tissue, the endolithic *Ostreobium* layer, and the deeper skeleton layer. Our analysis revealed a nuanced stratification of microbial activity, with microeukaryotes dominating key metabolic processes in the coral tissue and *Ostreobium* layer, while prokaryotes emerge as the primary active symbionts in the deeper skeleton (**Fig 5**). These findings underscore the profound influence of physicochemical gradients—including light penetration, nutrient availability, and carbon distribution—in sculpting both community composition and functional specialization within the holobiont. Functional analysis highlighted the critical contributions of *Symbiodiniaceae, Ostreobium*, and Cyanobacteria to photosynthate production, with *Symbiodiniaceae* predominantly occupying the coral tissue, while *Ostreobium* and Cyanobacteria play pivotal roles in the endolithic *Ostreobium*-layer (**Fig 5**). These symbionts, evolutionarily adapted to varying light conditions and utilizing distinct photosynthetic pigments, demonstrated unique nutrient cycling capabilities. Notably, in addition to *Symbiodiniaceae*, we identified *Ostreobium* and fungi as the only taxa capable of exporting sugars via SWEET proteins, providing novel insights into nutrient exchange mechanisms within the holobiont. Analysis of nitrogen metabolic pathways revealed Cyanobacteria as the exclusive nitrogen fixers, with other nitrogen and sulphur metabolic processes likely distributed across holobiont members, suggesting a robust functional redundancy that potentially enhances ecosystem resilience (**Fig 5**). Our analysis also identified microeukaryotes as the primary scavengers of ROS and RNS across all compartments, playing a crucial role in maintaining holobiont health, particularly under thermal stress conditions (**Fig 5**). These findings offer a comprehensive view of the metabolic activities within the coral microbiome, elucidating the intricate functional roles of its diverse members. While this research significantly advances our understanding of coral holobiont functions, it simultaneously highlights the complexity of these systems and the necessity for more targeted research focussed on the microenvironmental niches of the coral holobiont. Future investigations should employ advanced sampling techniques and high-resolution analytical methods to more precisely map the functional dynamics between coral hosts, their symbionts, and the surrounding microenvironment (*11*, *105*, *106*). Ultimately, the insights gained from this

study are invaluable for advancing coral reef conservation efforts. By identifying functionally relevant symbiotic members, we can better inform the design of coral probiotics and other strategies aimed at enhancing coral resilience in the face of climate change and other environmental stressors.

Data and code availability

Raw reads generated in this study have been submitted to NCBI SRA under the Bioproject PRJNA1207708. All codes and files required to regenerate the figures have been made available via github.

Acknowledgement

Authors would like to acknowledge staff at the Heron Island Research Station for their assistance during sample collection. All coral samples were collected under Permit G22/47504.1 issued to KT and FR by Great Barrier Reef Marine Park Authority, Australia. Authors would also like to acknowledge reviewers of the manuscript for their suggestions.

Funding

This work was funded through the Australian Research Council grant DP200101613 (to H.V., L.L.B., M.M., and M.K.), the Faculty of Science (University of Melbourne, to H.V.). K.T. acknowledges support from the University of Melbourne Early Career Research Grant. M.K. acknowledges support from the Gordon and Betty Moore Foundation through grant no. GBMF9206 (https://doi.org/10.37807/GBMF9206). M.M. acknowledges support from NOAA CRCP: National Oceanic and Atmospheric Administration Coral Reef Conservation Program NA19NOS4820132. H.V. is supported by the Fundação para a Ciência e a Tecnologia (CEECIND:2023.06155). Computational analysis of this work was supported by the University of Melbourne's Research Computing Services and the Petascale Campus Initiative.

References

1. C. Jaspers, S. Fraune, A. E. Arnold, D. J. Miller, T. C. G. Bosch, C. R. Voolstra, Consortium of Australian Academy of Science Boden Research Conference Participants, Resolving structure and function of metaorganisms through a holistic framework combining reductionist and integrative approaches. *Zoology* **133**, 81–87 (2019).

2. R. S. Peixoto, M. Sweet, H. D. M. Villela, P. Cardoso, T. Thomas, C. R. Voolstra, L. Høj, D. G. Bourne, Coral Probiotics: Premise, Promise, Prospects. *Annu Rev Anim Biosci* **9**, 265–288 (2021).

3. M. J. H. van Oppen, L. L. Blackall, Coral microbiome dynamics, functions and design in a changing world. *Nat. Rev. Microbiol.* **17**, 557–567 (2019).

4.   E. Rosenberg, O. Koren, L. Reshef, R. Efrony, I. Zilber-Rosenberg, The role of
microorganisms in coral health, disease and evolution. *Nat. Rev. Microbiol.* **5**, 355–362
(2007).

5.   F. Rohwer, V. Seguritan, F. Azam, N. Knowlton, Diversity and distribution of coral-
associated bacteria. *Mar. Ecol. Prog. Ser.* **243**, 1–10 (2002).

6.   R. V. Thurber, J. P. Payet, A. R. Thurber, A. M. S. Correa, Virus-host interactions and
their roles in coral reef health and disease. *Nat. Rev. Microbiol.* **15**, 205–216 (2017).

7.   T. C. LaJeunesse, J. E. Parkinson, P. W. Gabrielson, H. J. Jeong, J. D. Reimer, C. R.
Voolstra, S. R. Santos, Systematic Revision of Symbiodiniaceae Highlights the Antiquity
and Diversity of Coral Endosymbionts. *Curr. Biol.* **28**, 2570–2580.e6 (2018).

8.   L. Muscatine, The role of symbiotic algae in carbon and energy flux in reef corals.
Ecosystems of the world **25**, 75–87 (1990).

9.   L. A. Morris, C. R. Voolstra, K. M. Quigley, D. G. Bourne, L. K. Bay, Nutrient Availability
and Metabolism Affect the Stability of Coral-Symbiodiniaceae Symbioses. *Trends*
Microbiol. **27**, 678–689 (2019).

10.   M. J. Huggett, A. Apprill, Coral microbiome database: Integration of sequences reveals
high diversity and relatedness of coral-associated microbes. *Environ. Microbiol. Rep.*
11, 372–385 (2019).

11.   E. Bollati, D. J. Hughes, D. J. Suggett, J.-B. Raina, M. Kühl, Microscale sampling of the
coral gastrovascular cavity reveals a gut-like microbial community. *anim microbiome* **6**
(2024).

12.   M. J. Sweet, A. Croquer, J. C. Bythell, Bacterial assemblages differ between
compartments within the coral holobiont. *Coral Reefs* **30**, 39–52 (2011).

13.   V. R. Marcelino, M. J. van Oppen, H. Verbruggen, Highly structured prokaryote
communities exist within the skeleton of coral colonies. *ISME J.* **12**, 300–303 (2018).

14.   F. Ricci, K. Tandon, M. Moßhammer, E. H.-J. Cho, L. L. Blackall, M. Kühl, H.
Verbruggen, Fine-scale mapping of physicochemical and microbial landscapes of the
coral skeleton. *Environ. Microbiol.* **25**, 1505–1521 (2023).

15.   K. Tandon, M. M. Pasella, C. Iha, F. Ricci, J. Hu, C. J. O'Kelly, M. Medina, M. Kühl, H.
Verbruggen, Every refuge has its price: Ostreobium as a model for understanding how
algae can live in rock and stay in business. *Semin. Cell Dev. Biol.* **134**, 27–36 (2023).

16.   A. M. Bonacolta, B. A. Weiler, T. Porta-Fitó, M. Sweet, P. Keeling, J. del Campo,
Beyond the Symbiodiniaceae: diversity and role of microeukaryotic coral symbionts.
Coral Reefs **42**, 567–577 (2023).

17.   M. McCauley, T. L. Goulet, C. R. Jackson, S. Loesgen, Systematic review of cnidarian
microbiomes reveals insights into the structure, specificity, and fidelity of marine
associations. *Nat. Commun.* **14**, 4899 (2023).

18.   R. Rowan, D. A. Powers, Ribosomal RNA sequences and the diversity of symbiotic
dinoflagellates (zooxanthellae). *Proc. Natl. Acad. Sci. U. S. A.* **89**, 3639–3643 (1992).

19.   F. J. Pollock, R. McMinds, S. Smith, D. G. Bourne, B. L. Willis, M. Medina, R. V.
Thurber, J. R. Zaneveld, Coral-associated bacteria demonstrate phylosymbiosis and
cophylogeny. *Nat. Commun.* **9**, 4921 (2018).

20. L. Fujise, D. J. Suggett, M. Stat, T. Kahlke, M. Bunce, S. G. Gardner, S. Goyen, S.
Woodcock, P. J. Ralph, J. R. Seymour, N. Siboni, M. R. Nitschke, Unlocking the
phylogenetic diversity, primary habitats, and abundances of free-living Symbiodiniaceae
on a coral reef. *Mol. Ecol.* **30**, 343–360 (2021).

21. L. K. Lee, C. P. Leaw, L. C. Lee, Z. F. Lim, K. S. Hii, A. A. Chan, H. Gu, P. T. Lim,
Molecular diversity and assemblages of coral symbionts (Symbiodiniaceae) in diverse
scleractinian coral species. *Mar. Environ. Res.* **179**, 105706 (2022).

22. V. R. Marcelino, K. M. Morrow, M. J. H. van Oppen, D. G. Bourne, H. Verbruggen,
Diversity and stability of coral endolithic microbial communities at a naturally high pCO2
reef. *Mol. Ecol.* **26**, 5344–5357 (2017).

23. F. Ricci, A. Fordyce, W. Leggat, L. L. Blackall, T. Ainsworth, H. Verbruggen, Multiple
techniques point to oxygenic phototrophs dominating the Isopora palifera skeletal
microbiome. *Coral Reefs* **40**, 275–282 (2021).

24. K. Tandon, F. Ricci, J. Costa, M. Medina, M. Kühl, L. L. Blackall, H. Verbruggen,
Genomic view of the diversity and functional role of archaea and bacteria in the skeleton
of the reef-building corals Porites lutea and Isopora palifera. *Gigascience* **12** (2022).

25. S. J. Robbins, C. M. Singleton, C. X. Chan, L. F. Messer, A. U. Geers, H. Ying, A.
Baker, S. C. Bell, K. M. Morrow, M. A. Ragan, D. J. Miller, S. Forêt, ReFuGe2020
Consortium, C. R. Voolstra, G. W. Tyson, D. G. Bourne, A genomic view of the reef-
building coral Porites lutea and its microbial symbionts. *Nat Microbiol* **4**, 2090–2100
(2019).

26. T. Doering, K. Tandon, S. H. Topa, S. J. Pidot, L. L. Blackall, M. J. H. van Oppen,
Genomic exploration of coral-associated bacteria: identifying probiotic candidates to
increase coral bleaching resilience in Galaxea fascicularis. *Microbiome* **11**, 185 (2023).

27. S.-H. Yang, K. Tandon, C.-Y. Lu, N. Wada, C.-J. Shih, S. S.-Y. Hsiao, W.-N. Jane, T.-C.
Lee, C.-M. Yang, C.-T. Liu, V. Denis, Y.-T. Wu, L.-T. Wang, L. Huang, D.-C. Lee, Y.-W.
Wu, H. Yamashiro, S.-L. Tang, Metagenomic, phylogenetic, and functional
characterization of predominant endolithic green sulfur bacteria in the coral Isopora
palifera. *Microbiome* **7**, 3 (2019).

28. M. Sweet, H. Villela, T. Keller-Costa, R. Costa, S. Romano, D. G. Bourne, A. Cárdenas,
M. J. Huggett, A. H. Kerwin, F. Kuek, M. Medina, J. L. Meyer, M. Müller, F. J. Pollock,
M. S. Rappé, M. Sere, K. H. Sharp, C. R. Voolstra, N. Zaccardi, M. Ziegler, R. Peixoto,
Insights into the Cultured Bacterial Fraction of Corals. *mSystems* **6**, e0124920 (2021).

29. P. G. Falkowski, Z. Dubinsky, L. Muscatine, J. W. Porter, Light and the Bioenergetics of
a Symbiotic Coral. *Bioscience* **34**, 705–709 (1984).

30. K. Tandon, C.-Y. Lu, P.-W. Chiang, N. Wada, S.-H. Yang, Y.-F. Chan, P.-Y. Chen, H.-Y.
Chang, Y.-J. Chiou, M.-S. Chou, W.-M. Chen, S.-L. Tang, Comparative genomics:
Dominant coral-bacterium Endozoicomonas acroporae metabolizes
dimethylsulfoniopropionate (DMSP). *ISME J.* **14**, 1290–1303 (2020).

31. N. Rädecker, C. Pogoreutz, C. R. Voolstra, J. Wiedenmann, C. Wild, Nitrogen cycling in
corals: the key to understanding holobiont functioning? *Trends Microbiol.* **23**, 490–497
(2015).

32. M. J. Neave, C. T. Michell, A. Apprill, C. R. Voolstra, Endozoicomonas genomes reveal
functional adaptation and plasticity in bacterial strains symbiotically associated with

745   diverse marine hosts. *Sci. Rep.* **7**, 40579 (2017).

746 33. D. Schlichter, H. Kampmann, S. Conrady, Trophic potential and photoecology of
747    endolithic algae living within coral skeletons. *Mar. Ecol.* **18**, 299–317 (1997).

748 34. M. Fine, Y. Loya, Endolithic algae: an alternative source of photoassimilates during
749    coral bleaching. *Proceedings of the Royal Society of London. Series B: Biological*
750    *Sciences* **269**, 1205–1210 (2002).

751 35. L. Sangsawang, B. E. Casareto, H. Ohba, H. M. Vu, A. Meekaew, T. Suzuki, T. Yeemin,
752    Y. Suzuki, 13C and 15N assimilation and organic matter translocation by the endolithic
753    community in the massive coral Porites lutea. *R Soc Open Sci* **4**, 171201 (2017).

754 36. E. A. Ottesen, C. R. Young, S. M. Gifford, J. M. Eppley, R. Marin 3rd, S. C. Schuster, C.
755    A. Scholin, E. F. DeLong, Ocean microbes. Multispecies diel transcriptional oscillations
756    in open ocean heterotrophic bacterial assemblages. *Science* **345**, 207–212 (2014).

757 37. T. P. Hughes, J. T. Kerry, S. R. Connolly, J. G. Álvarez-Romero, C. M. Eakin, S. F.
758    Heron, M. A. Gonzalez, J. Moneghetti, Emergent properties in the responses of tropical
759    corals to recurrent climate extremes. *Curr. Biol.* **31**, 5393–5399.e3 (2021).

760 38. M. J. H. van Oppen, J. K. Oliver, H. M. Putnam, R. D. Gates, Building coral reef
761    resilience through assisted evolution. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 2307–2313
762    (2015).

763 39. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
764    sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

765 40. E. Kopylova, L. Noé, H. Touzet, SortMeRNA: fast and accurate filtering of ribosomal
766    RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

767 41. P. Yilmaz, L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J.
768    Peplies, W. Ludwig, F. O. Glöckner, The SILVA and "All-species Living Tree Project
769    (LTP)" taxonomic frameworks. *Nucleic Acids Res.* **42**, D643–D648 (2013).

770 42. H. R. Gruber-Vodicka, B. K. B. Seah, E. Pruesse, phyloFlash: Rapid Small-Subunit
771    rRNA Profiling and Targeted Assembly from Metagenomes. *mSystems* **5** (2020).

772 43. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
773    **9**, 357–359 (2012).

774 44. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis,
775    R. Durbin, 1000 Genome Project Data Processing Subgroup, The Sequence
776    Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

777 45. M. G. Grabherr, B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X.
778    Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A.
779    Gnirke, N. Rhind, F. di Palma, B. W. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman,
780    A. Regev, Full-length transcriptome assembly from RNA-Seq data without a reference
781    genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

782 46. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
783    search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

784 47. B. Buchfink, K. Reuter, H.-G. Drost, Sensitive protein alignments at tree-of-life scale
785    using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

48. D. H. Huson, S. Beier, I. Flade, A. Górska, M. El-Hadidi, S. Mitra, H.-J. Ruscheweyh, R.
Tappu, MEGAN Community Edition - Interactive Exploration and Analysis of Large-
Scale Microbiome Sequencing Data. *PLoS Comput. Biol.* **12**, e1004957 (2016).

49. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

50. P. J. McMurdie, S. Holmes, phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

51. H. Wickham, ggplot2. *Wiley Interdiscip. Rev. Comput. Stat.* **3**, 180–185 (2011).

52. D. M. Bryant, K. Johnson, T. DiTommaso, T. Tickle, M. B. Couger, D. Payzin-Dogru, T.
J. Lee, N. D. Leigh, T.-H. Kuo, F. G. Davis, J. Bateman, S. Bryant, A. R. Guzikowski, S.
L. Tsai, S. Coyne, W. W. Ye, R. M. Freeman Jr, L. Peshkin, C. J. Tabin, A. Regev, B. J.
Haas, J. L. Whited, A Tissue-Mapped Axolotl De Novo Transcriptome Enables
Identification of Limb Regeneration Factors. *Cell Rep.* **18**, 762–776 (2017).

53. C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas,
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

54. J. Huerta-Cepas, D. Szklarczyk, D. Heller, A. Hernández-Plaza, S. K. Forslund, H.
Cook, D. R. Mende, I. Letunic, T. Rattei, L. J. Jensen, C. von Mering, P. Bork, eggNOG
5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314 (2019).

55. M. Kanehisa, Y. Sato, K. Morishima, BlastKOALA and GhostKOALA: KEGG Tools for
Functional Characterization of Genome and Metagenome Sequences. *J. Mol. Biol.* **428**,
726–731 (2016).

56. P. Jones, D. Binns, H.-Y. Chang, M. Fraser, W. Li, C. McAnulla, H. McWilliam, J.
Maslen, A. Mitchell, G. Nuka, S. Pesseat, A. F. Quinn, A. Sangrador-Vegas, M.
Scheremetjew, S.-Y. Yong, R. Lopez, S. Hunter, InterProScan 5: genome-scale protein
function classification. *Bioinformatics* **30**, 1236–1240 (2014).

57. G. Salazar, L. Paoli, A. Alberti, J. Huerta-Cepas, H.-J. Ruscheweyh, M. Cuenca, C. M.
Field, L. P. Coelho, C. Cruaud, S. Engelen, A. C. Gregory, K. Labadie, C. Marec, E.
Pelletier, M. Royo-Llonch, S. Roux, P. Sánchez, H. Uehara, A. A. Zayed, G. Zeller, M.
Carmichael, C. Dimier, J. Ferland, S. Kandels, M. Picheral, S. Pisarev, J. Poulain, Tara
Oceans Coordinators, S. G. Acinas, M. Babin, P. Bork, C. Bowler, C. de Vargas, L.
Guidi, P. Hingamp, D. Iudicone, L. Karp-Boss, E. Karsenti, H. Ogata, S. Pesant, S.
Speich, M. B. Sullivan, P. Wincker, S. Sunagawa, Gene Expression Changes and
Community Turnover Differentially Shape the Global Ocean Metatranscriptome. *Cell*
179, 1068–1083.e21 (2019).

58. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

59. L. M. van der Loos, R. Nijland, Biases in bulk: DNA metabarcoding of marine
communities and the methodology involved. *Mol. Ecol.* **30**, 3270–3288 (2021).

60. S. K. Davy, D. Allemand, V. M. Weis, Cell biology of cnidarian-dinoflagellate symbiosis.
Microbiol. Mol. Biol. Rev. **76**, 229–261 (2012).

61. W. K. Kwong, J. Del Campo, V. Mathur, M. J. A. Vermeij, P. J. Keeling, A widespread

coral-infecting apicomplexan with chlorophyll biosynthesis genes. *Nature* **568**, 103–107
(2019).

62. A. Oren, G. M. Garrity, Valid publication of the names of forty-two phyla of prokaryotes.
Int. J. Syst. Evol. Microbiol. **71** (2021).

63. A. Cárdenas, J.-B. Raina, C. Pogoreutz, N. Rädecker, J. Bougoure, P. Guagliardo, M.
Pernice, C. R. Voolstra, Greater functional diversity and redundancy of coral endolithic
microbiomes align with lower coral bleaching susceptibility. *ISME J.* **16**, 2406–2420
(2022).

64. J. R. Guest, J. Low, K. Tun, B. Wilson, C. Ng, D. Raingeard, K. E. Ulstrup, J. T. I. Tanzil,
P. A. Todd, T. C. Toh, D. McDougald, L. M. Chou, P. D. Steinberg, Coral community
response to bleaching on a highly disturbed reef. *Sci. Rep.* **6**, 20717 (2016).

65. P. Buerger, G. M. Schmidt, M. Wall, C. Held, C. Richter, Temperature tolerance of the
coral Porites lutea exposed to simulated large amplitude internal waves (LAIW). *J. Exp.*
Mar. Bio. Ecol. **471**, 232–239 (2015).

66. J. Peng, C.-E. Wegner, Q. Bei, P. Liu, W. Liesack, Metatranscriptomics reveals a
differential temperature effect on the structural and functional organization of the
anaerobic food web in rice field soil. *Microbiome* **6**, 169 (2018).

67. N. Vannier, F. Mesny, F. Getzke, G. Chesneau, L. Dethier, J. Ordon, T. Thiergart, S.
Hacquard, Genome-resolved metatranscriptomics reveals conserved root colonization
determinants in a synthetic microbiota. *Nat. Commun.* **14**, 8274 (2023).

68. S. R. Law, A. R. Serrano, Y. Daguerre, J. Sundh, A. N. Schneider, Z. R. Stangl, D.
Castro, M. Grabherr, T. Näsholm, N. R. Street, V. Hurry, Metatranscriptomics captures
dynamic shifts in mycorrhizal coordination in boreal forests. *Proc. Natl. Acad. Sci. U. S.*
A. **119**, e2118852119 (2022).

69. S. Campana, A. Riesgo, E. Jongepier, J. Fuss, G. Muyzer, J. M. de Goeij, Meta-
transcriptomic comparison of two sponge holobionts feeding on coral- and macroalgal-
dissolved organic matter. *BMC Genomics* **23**, 674 (2022).

70. A. Schoenle, M. Hohlfeld, K. Hermanns, F. Mahé, C. de Vargas, F. Nitsche, H. Arndt,
High and specific diversity of protists in the deep-sea basins dominated by diplonemids,
kinetoplastids, ciliates and foraminiferans. *Commun Biol* **4**, 501 (2021).

71. S. L. Pichard, L. Campbell, J. H. Paul, Diversity of the ribulose bisphosphate
carboxylase/oxygenase form I gene (rbcL) in natural phytoplankton communities. *Appl.*
Environ. Microbiol. **63**, 3600–3606 (1997).

72. B. K. Swan, M. Martinez-Garcia, C. M. Preston, A. Sczyrba, T. Woyke, D. Lamy, T.
Reinthaler, N. J. Poulton, E. D. P. Masland, M. L. Gomez, M. E. Sieracki, E. F. DeLong,
G. J. Herndl, R. Stepanauskas, Potential for chemolithoautotrophy among ubiquitous
bacteria lineages in the dark ocean. *Science* **333**, 1296–1300 (2011).

73. J. A. Raven, Contributions of anoxygenic and oxygenic phototrophy and
chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquat. Microb.*
Ecol. **56**, 177–192 (2009).

74. F. Ricci, C. Greening, Chemosynthesis: a neglected foundation of marine ecology and
biogeochemistry: (Trends in Microbiology, Published online January 30, 2024). *Trends*
Microbiol. **32**, 723 (2024).

75. M. P. Lesser, M. Sabrina Pankey, M. Slattery, K. J. Macartney, D. J. Gochfeld,
Microbiome diversity and metabolic capacity determines the trophic ecology of the
holobiont in Caribbean sponges. *ISME Commun.* **2**, 112 (2022).

76. M. Kühl, G. Holst, A. W. D. Larkum, P. J. Ralph, IMAGING OF OXYGEN DYNAMICS
WITHIN THE ENDOLITHIC ALGAL COMMUNITY OF THE MASSIVE CORAL
PORITES LOBATA(1). *J. Phycol.* **44**, 541–550 (2008).

77. R. Samsonoff Wa Maccoll, Biliproteins and phycobilisomes from cyanobacteria and red
algae at the extreme habitats. *Arch* **176**, 400–405 (2001).

78. A. de los Ríos, M. Grube, L. G. Sancho, C. Ascaso, Ultrastructural and genetic
characteristics of endolithic cyanobacterial biofilms colonizing Antarctic granite rocks:
Antarctic endolithic cyanobacterial biofilms. *FEMS Microbiol. Ecol.* **59**, 386–395 (2007).

79. S. H. Magnusson, M. Fine, M. Kühl, Light microclimate of endolithic phototrophs in the
scleractinian corals Montipora monasteriata and Porites cylindrica. *Mar. Ecol. Prog. Ser.*
332, 119–128 (2007).

80. C. Iha, K. E. Dougan, J. A. Varela, V. Avila, C. J. Jackson, K. A. Bogaert, Y. Chen, L. M.
Judd, R. Wick, K. E. Holt, M. M. Pasella, F. Ricci, S. I. Repetti, M. Medina, V. R.
Marcelino, C. X. Chan, H. Verbruggen, Genomic adaptations to an endolithic lifestyle in
the coral-associated alga Ostreobium. *Curr. Biol.* **31**, 1393–1402.e5 (2021).

81. M. Lichtenberg, A. W. D. Larkum, M. Kühl, Photosynthetic Acclimation of Symbiodinium
in hospite Depends on Vertical Position in the Tissue of the Scleractinian Coral
Montastrea curta. *Front. Microbiol.* **7**, 230 (2016).

82. O. Hoegh-Guldberg, P. J. Mumby, A. J. Hooten, R. S. Steneck, P. Greenfield, E.
Gomez, C. D. Harvell, P. F. Sale, A. J. Edwards, K. Caldeira, N. Knowlton, C. M. Eakin,
R. Iglesias-Prieto, N. Muthiga, R. H. Bradbury, A. Dubi, M. E. Hatziolos, Coral reefs
under rapid climate change and ocean acidification. *Science* **318**, 1737–1742 (2007).

83. N. Rädecker, C. Pogoreutz, H. M. Gegner, A. Cárdenas, F. Roth, J. Bougoure, P.
Guagliardo, C. Wild, M. Pernice, J.-B. Raina, A. Meibom, C. R. Voolstra, Heat stress
destabilizes symbiotic nutrient cycling in corals. *Proc. Natl. Acad. Sci. U. S. A.* **118**
(2021).

84. J. Ji, L. Yang, Z. Fang, Y. Zhang, M. Zhuang, H. Lv, Y. Wang, Plant SWEET Family of
Sugar Transporters: Structure, Evolution and Biological Functions. *Biomolecules* **12**
(2022).

85. O. Hoegh-Guldberg, Climate change, coral bleaching and the future of the world's coral
reefs. *Mar. Freshwater Res.* **50**, 839–866 (1999).

86. A. Roik, M. Reverter, C. Pogoreutz, A roadmap to understanding diversity and function
of coral reef-associated fungi. *FEMS Microbiol. Rev.* **46** (2022).

87. A. R. Mohamed, M. A. Ochsenkühn, A. M. Kazlak, A. Moustafa, S. A. Amin, The coral
microbiome: towards an understanding of the molecular mechanisms of coral-microbiota
interactions. *FEMS Microbiol. Rev.* **47** (2023).

88. N. D. Olson, T. D. Ainsworth, R. D. Gates, M. Takabayashi, Diazotrophic bacteria
associated with Hawaiian Montipora corals: Diversity and abundance in correlation with
symbiotic dinoflagellates. *J. Exp. Mar. Bio. Ecol.* **371**, 140–146 (2009).

89. K. A. Lema, B. L. Willis, D. G. Bourne, Amplicon pyrosequencing reveals spatial and temporal consistency in diazotroph assemblages of the Acropora millepora microbiome. *Environ. Microbiol.* **16**, 3345–3359 (2014).

90. K. A. Lema, D. G. Bourne, B. L. Willis, Onset and establishment of diazotrophs and other bacterial associates in the early life history stages of the coral Acropora millepora. *Mol. Ecol.* **23**, 4682–4695 (2014).

91. C. Cook, C. D'Elia, Are natural populations of zooxanthellae ever nutrient-limited? *Symbiosis* **4**, 199–211 (1987).

92. J. Li, Q. Yang, J. Dong, M. Sweet, Y. Zhang, C. Liu, Y. Zhang, X. Tang, W. Zhang, S. Zhang, Microbiome Engineering: A Promising Approach to Improve Coral Health. *Proc. Est. Acad. Sci. Eng.* **28**, 105–116 (2023).

93. M. A. Moran, B. P. Durham, Sulfur metabolites in the pelagic ocean. *Nat. Rev. Microbiol.* **17**, 665–678 (2019).

94. T. Higuchi, K. Tanaka, K. Shirai, I. Yuyama, T. Mezaki, N. Takahata, Y. Sano, Sulfur assimilation in corals with aposymbiotic and symbiotic zooxanthellae. *Environ. Microbiol. Rep.* **13**, 98–103 (2021).

95. B. Meyer, J. Kuever, Homology modeling of dissimilatory APS reductases (AprBA) of sulfur-oxidizing and sulfate-reducing prokaryotes. *PLoS One* **3**, e1514 (2008).

96. T. Watanabe, H. Kojima, M. Fukui, Identity of major sulfur-cycle prokaryotes in freshwater lake ecosystems revealed by a comprehensive phylogenetic study of the dissimilatory adenylylsulfate reductase. *Sci. Rep.* **6**, 36262 (2016).

97. J. F. Imhoff, Biology of green sulfur bacteria, *eLS* (2014). https://doi.org/10.1002/9780470015902.a0000458.pub2.

98. C. Ng, M. Z. DeMaere, T. J. Williams, F. M. Lauro, M. Raftery, J. A. E. Gibson, C. Andrews-Pfannkoch, M. Lewis, J. M. Hoffman, T. Thomas, R. Cavicchioli, Metaproteogenomic analysis of a dominant green sulfur bacterium from Ace Lake, Antarctica. *ISME J.* **4**, 1002–1019 (2010).

99. Y.-H. Chen, S.-H. Yang, K. Tandon, C.-Y. Lu, H.-J. Chen, C.-J. Shih, S.-L. Tang, Potential syntrophic relationship between coral-associated Prosthecochloris and its companion sulfate-reducing bacterium unveiled by genomic analysis. *Microb Genom* **7** (2021).

100. I. Kushkevych, J. Cejnar, J. Treml, D. Dordević, P. Kollar, M. Vítězová, Recent Advances in Metabolic Pathways of Sulfate Reduction in Intestinal Bacteria. *Cells* **9** (2020).

101. J. Helgoe, S. K. Davy, V. M. Weis, M. Rodriguez-Lanetty, Triggers, cascades, and endpoints: connecting the dots of coral bleaching mechanisms. *Biol. Rev. Camb. Philos. Soc.* **99**, 715–752 (2024).

102. A. M. Dungan, D. Bulach, H. Lin, M. J. H. van Oppen, L. L. Blackall, Development of a free radical scavenging bacterial consortium to mitigate oxidative stress in cnidarians. *Microb. Biotechnol.* **14**, 2025–2040 (2021).

103. N. Delgadillo-Ordoñez, N. Garcias-Bonet, I. Raimundo, F. C. García, H. Villela, E. O. Osman, E. P. Santoro, J. Curdia, J. G. D. Rosado, P. Cardoso, A. Alsaggaf, A. Barno,

C. P. Antony, C. Bocanegra, M. L. Berumen, C. R. Voolstra, F. Benzoni, S. Carvalho, R.
S. Peixoto, Probiotics reshape the coral microbiome in situ without detectable off-target
effects in the surrounding environment. *Commun Biol* **7**, 434 (2024).

104.   C. R. Voolstra, J.-B. Raina, M. Dörr, A. Cárdenas, C. Pogoreutz, C. B. Silveira, A. R.
Mohamed, D. G. Bourne, H. Luo, S. A. Amin, R. S. Peixoto, The coral microbiome in
sickness, in health and in a changing world. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-
024-01015-3 (2024).

105.   M. J. H. van Oppen, J.-B. Raina, Coral holobiont research needs spatial analyses at
the microbial scale. *Environ. Microbiol.* **25**, 179–183 (2023).

106.   D. J. Hughes, J.-B. Raina, D. A. Nielsen, D. J. Suggett, M. Kühl, Disentangling
compartment functions in sessile marine invertebrates. *Trends Ecol. Evol.* **37**, 740–748
(2022).

968