1 **Metatranscriptomics sheds light on "who is doing what" in the coral *Porites lutea***
2

3 Kshitij Tandon[1,2,3*], Juntong Hu[1], Francesco Ricci[4,5], Linda Louise Blackall[1], Mónica Medina[6],
4 Michael Kühl[7], Heroen Verbruggen[1,2,8]
5
[1]School of BioSciences, The University of Melbourne, Melbourne, VIC 3010, Australia
[2]Melbourne Integrative Genomics, The University of Melbourne, Melbourne, VIC 3010,
Australia
[3]Department of Microbiology and Immunology at the Peter Doherty Institute of Infection and
Immunity, The University of Melbourne, Melbourne, VIC 3010, Australia
[4]Department of Microbiology, Biomedicine Discovery Institute, Monash University, Clayton,
VIC 3800, Australia
[5]Securing Antarctica's Environmental Future, Monash University, Clayton, VIC 3800,
Australia
[6]Department of Biology, Pennsylvania State University, University Park, PA 16802, USA
[7]Marine Biological Section, Department of Biology, University of Copenhagen, DK-3000
Helsingør, Denmark
[8]CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório
Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal
20
21 *Corresponding author: Kshitij Tandon (k.tandon@unimelb.edu.au)
22

Abstract

Global decline of coral reefs due to climate change calls for effective, nature-based
strategies to protect these crucial ecosystems. Developing such strategies requires a
thorough understanding of the complex roles and interactions of key inhabitants within coral
holobionts. Using a metatranscriptomics approach, we investigated the transcriptionally
active microbiome of the healthy stony coral *Porites lutea*, focusing on the coral tissue, the
green endolithic algal layer (*Ostreobiu*m layer), and the deeper coral skeleton. We show
distinct, metabolically active communities within these compartments and highlight
substantial metabolic redundancy across carbon, nitrogen, and sulphur pathways. Our study
provides the first transcriptomic evidence of the ability of *Ostreobium* to transfer fixed carbon
to other members of the holobiont and the coral host. Additionally, we identified critical roles
of diverse coral holobiont members in nutrient cycling and maintaining homeostasis through
scavenging of reactive oxygen and nitrogen species. This study provides novel molecular-
level understanding of the functional roles played by diverse coral holobiont members in their
respective compartments and underscores that corals harbour several distinct microbiomes
with a wide array of functions.

Introduction

Coral reefs owe their success to the symbiosis between reef-building, stony corals (Scleractinia) (*1–3*) and a large diversity of closely associated bacteria, archaea, viruses, and microeukaryotes, collectively termed the coral holobiont (*3–6*). These symbionts *sensu lato* encompass organisms spanning the full spectrum of host-microbe interactions, from mutualistic partners that enhance coral fitness to opportunistic pathogens, as well as commensals and other microorganisms whose ecological roles may shift along this continuum depending on environmental conditions. While the most well-characterized symbiosis in corals involves photosynthetic, endosymbiotic microalgae in the family Symbiodiniaceae residing in the gastrodermal tissues of the animal host (*7–9*), a compendium of prokaryotes associated with corals has identified 39 bacterial and 2 archaeal phyla (*10*). This diverse microbial assemblage is not only distinct between different compartments within the coral (*11*), but fine-scale mapping of the coral skeleton has revealed a stratified community across its depth (*12–14*). In recent years, other microeukaryotes besides Symbiodiniaceae, including endolithic algae of the genus *Ostreobium*, fungi, and apicomplexans have also garnered attention (*15, 16*).

The diversity and composition of coral holobiont members have been investigated through metabarcoding of ribosomal marker genes (16S rRNA, 18S rRNA, ITS2) (*17–19*) (*20, 21*),providing invaluable insights into the diverse and structured symbiont composition *sensu lato* within coral compartments (*12, 13, 22, 23*). Combined culture-based and molecular approaches have expanded our understanding of the functions of coral holobiont members in maintaining coral health and resilience (*24–28*). Symbiodiniaceae, in particular, play a crucial role by supplying corals with photosynthates, meeting a significant portion of their energy demand (*29*). Coral-associated bacteria and archaea have been implicated in a myriad of functions essential for sustaining coral health and resilience, including stress alleviation through scavenging of reactive oxygen and nitrogen species (ROS and RNS), protection against pathogens and nutrient cycling (*24–26, 28, 30–32*). Coral skeleton-dwelling endolithic green algae of the genus *Ostreobium,* which form conspicuous green bands beneath the coral tissue, have also been suggested to support corals by providing photosynthates during coral bleaching (*33–35*), but we lack a detailed understanding of the metabolic interaction between the coral animal and *Ostreobium* (*15*).

Most of our present knowledge on the role of microbiota for coral function and health comes from amplicon and metagenomic sequencing studies, with potential functions inferred using correlative, bioinformatic analyses. Despite these advances and the extensive information now available on the gene-encoded functional potential of microbiome and microeukaryotic members through genomic and metagenomic studies, a significant knowledge gap persists

regarding actual microbial metabolic activity and its spatial organisation within the coral
holobiont. Genomic data alone cannot predict the actual contributions of specific functions by
symbionts *in situ*, as genes encoding these functions may be variably expressed or not
expressed at all. Metatranscriptomics enables the analysis of the pool of actively expressed
genes in an environmental sample, highlighting the ecophysiologically relevant processes
occurring at the time of sampling (*36*). Therefore, metatranscriptomics can provide insights
into which community members perform which functional roles.
83
Obtaining a molecular level understanding of the functional roles of holobiont members is
crucial especially given the rapid decline of coral reefs worldwide due the adverse effects of
climate change (*37*). Addressing this knowledge gap will also be beneficial for developing
effective nature-based interventions to protect coral reefs, including improved assisted
evolution strategies and development of coral probiotics (*2*, *38*). Both of these approaches
rely on a comprehensive understanding of the roles played by diverse microbial symbionts in
maintaining coral health.
In this study, we aimed to address the fundamental question of "who is doing what" in a
healthy coral holobiont across different compartments. Using metatranscriptomics, we aimed
to identify the active symbiont community of the coral *Porites lutea*, characterising their
specific activities and where these are carried out in the coral, from the coral tissue to the
endolithic algal layer (hereafter *Ostreobium*-layer) and the deeper coral skeleton. Our focus
was on key nutrient cycling pathways, including carbon-fixation (C), photosynthesis (P),
nitrogen (N) and sulphur (S) metabolism, sugar-export, and the scavenging of reactive
oxygen and nitrogen species (ROS and RNS).
99
100   **Methods:**
101
Coral sampling and colony fragmentation Fragments from six healthy-looking, *Porites*
lutea colonies, spaced at least 15 m apart, were collected at low tide (1-2 m depth) from the
research zone of the Heron Island reef flat, southern Great Barrier Reef (23°44′S, 151°91′E),
in October 2022. Four fragments (from colonies we numbered 1, 3, 4 and 5) were collected
during daytime, while two (colonies 6 and 7) were collected five-six hours after sunset. All
collected colonies exhibited a visually discernible *Ostreobium-layer* in the coral skeleton . 10-
50 mm below the coral tissue layer (Fig. 1A). The fragments were collected using a sterile
hammer and chisel and were placed in separate sterile zip-loc polyethylene bags in
seawater and flash frozen in liquid nitrogen immediately upon collection. A 5-mm thick slice
was cut from the frozen samples, perpendicular to the vertical growth axis, using a diamond
saw with a continuous flow of sterile filtered (0.2 µm) seawater. Later, pliers were used to
obtain fragments from three distinct layers, i.e., the coral tissue, the green *Ostreobium*-layer,

and the tan fraction of the skeleton below the green band (termed the skeleton layer). During
this process, the material was kept frozen by repeated immersion in liquid nitrogen. Each
layer was kept frozen at -80ºC, transported to the University of Melbourne on dry ice, and
processed immediately upon arrival.
118

Total RNA extraction

119
For RNA extraction, samples of individual layers were ground to a fine powder in liquid
nitrogen using pre-cooled mortars and pestles. Approximately 200 mg of powdered tissue
samples and between 300-500 mg of powdered *Ostreobium-layer* and skeleton layer were
used for RNA extraction with 1 mL of pre-cooled TRIzol reagent (Invitrogen, USA). Total
RNA was extracted following the manufacturer's protocol with some modifications (**see**
supplementary file 1 for details). Quantification of total RNA was conducted using the
Qubit RNA HS Assay kit (Thermo Fisher Scientific, USA).
127

Metatranscriptome sequencing and data-preprocessing

128
We did not perform any ribosomal RNA depletion due to the dual aim of sequencing mRNA
and rRNA from both eukaryotes and prokaryotes simultaneously, and profiling the
transcriptionally active microbial and microeukaryotic community composition using 16S and
18S rRNA abundance (see below). Total RNA libraries were prepared for 36 samples (6
colonies x 3 layers x 2 replicates per layer) and sequenced on the Illumina NovaSeq
platform by Azenta (Suzhou, China). Sequencing yielded approximately 23 million reads per
sample with 2x150 bp reads.
Raw reads were quality checked and trimmed using FastQC v0.11.9
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc) and Trimmomatic v0.39(*39*),
respectively. Trimmomatic parameters were as follows: *TruSeq3-PE.fa:2:30:5*
HEADCROP:5 SLIDINGWINDOW:4:20 MINLEN:50. Quality-filtered reads were first used to
identify and filter out rRNA reads using SortMeRNA v4.3.6(*40*) with Silva v138.1 LSU and
SSU (*SILVA138.1 LSU and SSU NR99 database, last accessed Feb 10, 2023*)(*41*). Reads
identified as rRNA were taxonomically classified using PhyloFlash v3.4(*42*) (with *almost-*
everything), which rapidly reconstructs the SSU rRNAs (NTUs, taxonomic units) and
taxonomically annotates them using Silva 138.1 NR99 SSU database. Reads identified as
rRNA were removed from the raw-read set, and non-rRNA reads were mapped to the
Porites lutea genome (accessed from www.reefgenomics.org) using bowtie2 v2.3.4.1(*43*)
and samtools 1.17(*44*) to identify and remove host-related reads.
High-quality, non-rRNA and non-host reads were co-assembled using Trinity v2.15.1(*45*)
with parameters *--min_contig_length 250 and --no_salmon*. Taxonomic annotation of
assembled transcripts was obtained using the lowest common ancestor approach by

performing *blastx (46)* searches against the NR database (last accessed April 2023) with
diamond v2.1.6.160(*47*), using an *e-value cutoff of 1e-5* and processed through MEGAN6
community edition v6.24.20, using the *megan-map-Feb2022-ue.db* and meganising the *daa*
file using *daa2info* (*48*). This process was used to identify and remove any transcripts
annotated as Scleractinia and other Eumetazoa. The remaining transcripts were quantified
using RSEM (RNA-Seq by Expectation and Maximisation)(*49*), implemented in Trinity
through the utility script *align_and_estimate_abundance.pl*. This approach generated both
gene and isoform expression profiles. Isoform-level expression data was used for the
analysis. Expression values per transcript were averaged across replicates for each layer
(tissue, *Ostreobium*-layer and skeleton).
161
Community composition analysis
Taxonomic classification of NTUs assembled from rRNA reads was conducted to profile the
abundance of transcriptionally active community members. NTUs classified as "Chloroplast"
or "Mitochondria" were removed. Abundance values were converted to relative abundance
for compositional analysis using barplots and log-transformed for diversity analysis using
PCoA. These analyses were performed using R packages phyloseq v1.42.0(*50*), metagMisc
v0.5.0 (https://github.com/vmikk/metagMisc) and ggplot2 v34.3 (*51*). We used the quantified
and taxonomically annotated transcripts assembled from mRNA reads to profile the
community composition encoding for different functional guilds based on the
metatranscriptome data and compared them to rRNA based analysis using the same
approach.
173
Functional analysis
To ascertain functional roles, we first filtered transcripts using Trimmed Mean of the M
(TMM) values of 0.1 and presence in at least 2 samples to identify truly expressed
transcripts. This approach filters out i) any transcripts from potential residual DNA or very
low expressed transcripts, ii) transcripts originating from spurious mapping of reads.
Transcripts deemed truly expressed were then grouped in a compartment-specific manner to
obtain tissue, *Ostreobium*-layer and skeleton-layer specific metatranscriptome. These
compartment specific transcripts were processed through Trinotate v4.0.0(*52*) and
Transdecoder v4 (https://github.com/TransDecoder/TransDecoder) to identify the longest
open reading frames and predict the proteins from transcripts ≥50 amino acids. Predicted
proteins were functionally annotated using a combination of EggNOG mapper v2.1.12(*53*)
with database eggnogDB v5.0.2(*54*) with *e-value cutoff 1e-5*, ghostkoala (*55*) with database
c_family_euk+genus_prok+viruses parameters and Interproscan v5.63-95.0(*56*) with *e value*

cutoff 1e-5 to obtain a comprehensive understanding of functional orthologous groups
(EggNOG and COG), KEGG pathways, including Kegg Ontology, KOs and protein families
(pfam). Metabolic markers for carbon-fixation (C), photosynthesis (P), nitrogen (N) and sulfur
(S) metabolism were identified from a previous study (*57*). Additionally, we focussed on
genes involved in sugar export and scavenging of ROS and RNS. Expression profiles of
transcripts functionally annotated with marker KOs or pfams were plotted using heatmaps
using ComplexHeatmap(*58*) in R.
194
195 **Results and Discussion**
196
197 **Corals harbor compartment-specific active symbiont community**
198
One of our primary objectives was to profile the transcriptionally active community within the
coral holobiont and compare these findings with the community composition obtained from
rRNA analysis. To achieve this, we examined the total and active community composition
across the coral tissue, the endolithic *Ostreobium-layer*, and the underlying coral skeleton,
and compared it with their rRNA based community composition (**Fig 1A**). For total
community composition, we taxonomically analysed 28,717 rRNA NTUs (6,010 from tissue
layer, 15,942 from *Ostreobium* layer, 26,712 from skeleton layer) belonging to Archaea
(2.92% - 840 sequences), Eukaryota (12.68% - 3,644 sequences), and Bacteria (84.38% -
24,233 sequences) (**Fig 1B**). This observation offers essential insights into the diversity and
composition of the coral holobiont. Unlike traditional amplification-based metabarcoding
analyses, which are often biased (*59*) and fail to provide accurate quantitative distinctions
between prokaryotes and eukaryotes, using rRNAs assembled from total RNA delivers a
more precise and comprehensive overview of the community composition. Eukaryotes
(Amorphea, SAR, Archaeplastida) were the dominant groups in the coral tissue
(**Supplementary Fig S1A**) and the *Ostreobium layer* (**Supplementary Fig S1B**), while the
deeper skeleton was dominated by Bacteria (Proteobacteria, Planctomyceota,
Desulfobacteria, and Cyanobacteria) (**Supplementary Fig S1C**). Our study shows that
bacterial diversity is highest in the skeleton, where the majority of bacterial NTU sequences
were assembled, confirming previous reports of greater bacterial richness in the skeleton
compared   to   tissue   compartments   (*13*,   *19*).
219
The active community composition was profiled using a curated set of 371,193 taxonomically
annotated transcripts out of a total of 780,431 assembled. After excluding 111,367
transcripts belonging to Scleractinia, Chordata, and Eumetazoa, 259,826 (47.56%)
transcripts remained. These transcripts represent Eukaryotes (52.48% - 136,366), Bacteria
(44.99% - 116,900), Archaea (1.97% - 5,131), and viruses (0.54% - 1,429) (**Fig 1B**) and

revealed a high relative abundance of alveolates, the eukaryotic group containing
Apicomplexa and dinoflagellates, including Symbiodiniaceae in coral tissue (**Supplementary**
Fig S2A). Green algae, Streptophyta and Chlorophyta, dominated the *Ostreobium layer* and
deeper skeleton. (**Supplementary Fig S2B and C**). The high relative abundance of
dinoflagellate (Symbiodiniaceae) expressed transcripts in coral tissue is unsurprising, as
they are the most dominant microeukaryotes in the coral tissue, meeting most of the coral's
energy demands (*60*). However, we identified dinoflagellate and Apicomplexa transcripts in
all compartments. Apicomplexans have been found in healthy coral colonies across the
geographic distribution and taxonomic diversity of corals, but their role is yet to be
characterized (*16*, *61*). Bacteria and Fungi were present in all compartments, with bacteria
as dominant members of the skeletal microbiome (**Supplementary Figure S2C)**.
Proteobacteria, Cyanobacteria and Actinobacteria were the dominant bacterial phyla in all
coral compartments. Members of the phylum Proteobacteria are some of the most
widespread coral microbiome members (*10*, *17*, *62*). Cyanobacteria have also previously
been identified as the dominant prokaryote group in the *Ostreobium layer* and deeper
skeleton of *P. lutea* from the Red Sea (*63*). Obtaining annotation for less than 50% of the
total assembled transcripts highlights the immense diversity of uncharacterised symbionts
associated with *P. lutea,* despite it being one of the best investigated coral species with
several metagenomic and physiological studies (*24*, *25*, *64*, *65*). These findings align with
earlier studies using the metatranscriptomics approach to profile transcriptionally active
microbiomes in other ecologically diverse systems, including soil (*66–68*), sponge (*69*) and
ocean microbiomes (*57*, *70*).

247
Fig. 1. Taxonomic and functional overview based on rRNA and transcriptome data A)
illustration of a coral fragment depicting the three compartments: Tissue, Ostreobium-layer
and Skeleton. B) Bar plots showing overall community composition based on rRNA and
transcriptome data. C) UpSet plot illustrating shared and unique transcripts across different
coral compartments. D) Venn diagram showing counts and percentage of transcripts which
are deemed expressed, expressed and taxonomically annotated, annotated at KO and at
OG levels.

255
256

Functional annotation of transcripts remains a bottleneck

Analysis of active metabolic pathways in a healthy coral holobiont revealed 261,884 truly
expressed transcripts through TMM filtering. These transcripts were distributed across the
three investigated compartments, with the highest overlap observed between the
Ostreobium layer and deeper skeleton (89,682 shared transcripts). Additionally, a substantial
number of uniquely expressed transcripts were identified in the deeper skeleton-layer
(48,966), while the coral tissue exhibited the lowest count of uniquely expressed transcripts
(**Fig. 1C**). Recent metabarcoding and metagenomic studies have highlighted the skeleton as
the most diverse compartment of corals (*13*, *19*, *63*). In line with these observations, our

coral skeleton samples revealed 178,580 uniquely predicted proteins from the expressed

transcripts. However, only 30.64% (54,731) could be annotated using KO ids and 29.25%

(52,252) using OGs (**Fig 1D**). The extraordinary uniqueness of skeletal proteins

underscores the added value of sampling genomically underexplored environments, such as

the coral skeleton. The low functional annotation observed in our dataset is consistent with a

recent study on gene expression changes and community turnover of global oceans, which

similarly found less than 25% of the transcripts assigned to KOs (*57*). Annotations based on

COG revealed a notable proportion of proteins encoded by transcripts involved in signal

transduction, intracellular trafficking, secretion, vesicular transport, post-translational

modification, protein turnover, and chaperones (**Supplementary Figure S3**). These findings

suggest intricate, interconnected metabolic interactions within coral compartments,

facilitating the exchange of nutrients and metabolites. Such processes are likely crucial for

coral health and stability, especially in nutrient-poor waters, and provide evidence of the

dynamic roles microbial communities play in maintaining coral homeostasis through their

contributions to essential nutrient cycles within the coral holobiont. However, to fully

elucidate these nutritions exchanges, further experimental validation is required to identify

metabolites being exchanged and the potential underlying mechanisms.

Distinct photosymbionts drive carbon fixation in coral compartments

To reveal the functional roles of transcriptionally active holobiont members in healthy *P.*

lutea coral, we identified key microbial processes, such as photosynthesis, carbon fixation,

sugar export, and nitrogen and sulphur cycling. Our analysis of pathway-specific metabolic

markers highlighted the taxonomic affiliations of transcripts and their expression profiles,

shedding light on the active contributions of each microbial group to these essential

processes. The genes with the highest number of transcripts included those encoding

photosynthetic markers for Photosystem I (PsaA, 59 transcripts) and Photosystem II (PsbA,

67 transcripts) and for subunits of RuBisCO(ribulose-1,5-bisphosphate

carboxylase/oxygenase, *rbc*L (79) and *rbc*S (18) genes), the key enzyme in carbon-fixation

via the Calvin-Benson cycle (**Fig 2**). Notably, while chemotrophs can also have a

transcriptionally active *rbc*L gene (*71*), our study did not taxonomically annotate any

chemotrophs, so the expression of these genes primarily reflects activities of phototrophs.

These genes were actively transcribed by Symbiodiniaceae, stramenopiles, Bryopsidales

(including the endolithic green alga *Ostreobium*), Cyanobacteria, Gammaproteobacteria, and

unclassified Viridiplantae and Chlorophyta in different coral compartments. Symbiodiniaceae

were the dominant contributors in coral tissue, while Cyanobacteria were active across all

coral compartments. In the *Ostreobium*-layer and deeper skeleton layer, *Ostreobium* and

other Ulvophyceae were the key contributors (**Supplementary data file 2**). The expression

levels of *rbc*L and *rbc*S correlated with those of *psb*A and *psa*A, indicating that carbon

fixation is primarily driven by chlorophototrophs rather than chemotrophs, which derive

energy from chemical sources (*72–74*). We identified the expression of coxS, coxM, and

coxL transcripts in microbes from coral skeletons, primarily affiliated with Cyanobacteria and

Proteobacteria, indicating the presence of carbon monoxide dehydrogenase (CODH)

activity. This activity enables CO oxidation to CO□, supporting energy generation through

chemosynthesis. These findings suggest that these microbes contribute to carbon cycling

and primary productivity in nutrient-poor, low-light environments. Their metabolic flexibility

highlights a potentially critical role in supporting coral reef ecosystems through alternative

energy pathways. Additionally, the expression of acs and acsB in the skeleton layers

indicates the involvement of bacteria in carbon fixation via the Wood–Ljungdahl pathway,

converting CO□ and CO into acetyl-CoA, a key intermediate in various metabolic processes

(*63, 75*). The expression patterns of these genes underscore the metabolic diversity within

the Ostreobium and skeleton-associated microbial communities, reflecting adaptation to

specific environmental conditions, such as low O_2 (*14, 76*) and the availability of CO,

although the latter warrants further experimental validation. The conversion of both inorganic

and organic carbon into energy may contribute to nutrient cycling, promoting microbial

persistence and growth, and ultimately supporting the overall health and productivity of the

coral   holobiont.

Besides *psa*A and *psb*A, we observed expression of other photosynthetic marker genes,

including those for the photosynthetic reaction centre, *petC* (encoding the Rieske protein, a

part of the cytochrome b6f complex that connects Photosystem II to Photosystem I) and

petE (encoding plastocyanin, which transfers electron between the cytochrome b6f complex

and Photosystem II), across all three holobiont compartments. This indicates the presence of

oxygenic phototrophs throughout the different layers of corals. Notably, genes encoding for

cyanobacteria-specific antenna proteins (*apcA, apcF, cpcA, cpeA* and *cpeT*), were highly

expressed in the *Ostreobium-layer* and deeper skeleton compared to the tissue, suggesting

a higher prevalence and activity of Cyanobacteria in these compartments (**Fig 2**). This

finding aligns with the taxonomic profiling at the rRNA (Supplementary Fig S1) and

metatranscriptome levels (Supplementary Fig S2) and corroborates earlier reports showing a

prevalence of genes with eukaryotic and cyanobacterial origins within the skeletal layers of

P. lutea (63). These genes are involved in the Calvin-Benson cycle and collectively

represent >50% of the relative contribution to the total carbon fixation genes in the skeletal

layers of *P. lutea* (*63*). Additionally, the expression of *pufM,* a marker for anoxygenic

photosynthesis that encodes a protein component of the bacterial photosynthetic reaction

centre, was restricted to the coral skeleton, indicating the presence of obligate anaerobes,
such as green sulphur and purple non-sulfur bacteria, in line with hyperspectral imaging of
pigment distributions over coral skeleton cross-sections in *P. lutea* (*14*).
343

The observed differential expression patterns in the coral compartments reflect adaptations
to varying light conditions and provide new insights into the metabolic flexibility of coral-
associated microbes. The higher expression of cyanobacterial antenna proteins in the
Ostreobium-layer and deeper skeleton, where light is limited, suggests that Cyanobacteria in
these regions utilise alternative chlorophyll pigments, such as chlorophyll d and f, to absorb
far-red light that penetrates through the coral tissue. This adaptation enables efficient
photosynthesis under low-light conditions, supporting the role of Cyanobacteria in energy
generation in these deeper layers. These findings extend earlier reports of the use of far-red
absorbing pigments by Cyanobacteria and Ostreobium, which thrive in endolithic habitats
with limited light (*77–79*). In the Ostreobium layer, the green alga *Ostreobium* also employs
similar strategies, utilising chlorophyll b and enriching its repertoire of light-harvesting
complex proteins in both photosystems to adapt to low-light conditions (*15*). This includes
changes such as the loss of photoprotection genes and duplication of the *Lhca1* gene *(80)*.
In contrast, the tissue compartment, exposed to higher light levels, harbours
Symbiodiniaceae and other phototrophs adapted to higher irradiance, utilising chlorophyll a
and c (*81*). The expression of specific photosynthetic genes across these compartments
underscores the sophisticated strategies employed by coral-associated phototrophs,
including *Ostreobium*, to thrive in their respective microhabitats. These findings highlight the
metabolic diversity within the holobiont and contribute to our understanding of coral
resilience in fluctuating environmental conditions.
364

Not only Symbiodiniaceae provide fixed carbon to the coral holobiont
To identify members of the coral holobiont responsible for exporting photosynthetically fixed
carbon compounds (photosynthates), thereby potentially contributing to coral energy
demands, we focused on genes encoding for "sugars will eventually be exported
transporters" (SWEET) proteins. SWEET proteins act as bidirectional uniporters/facilitators,
enabling the diffusion of sugars across cell membranes along concentration gradients (*82*,
83). We identified the expression of transcripts encoding SWEET proteins (pfams: PF04193
and PF03083) across all three compartments in the coral holobiont (**Fig 2**). Taxonomic
annotation of these transcripts identified eukaryotes, including Symbiodiniaceae,
Ostreobium, and other unclassified green algae (Chlorophyta), as the primary sugar
exporters (**Supplementary data file 2**). All of these organisms perform the Calvin cycle,
from which the intermediate glyceraldehyde-3-phosphate (G3P) serves as a precursor for

production of sugars including glucose, which is likely the main carbohydrate transported by
SWEET proteins (*84*). Interestingly, we did not detect any bacterial transcripts expressing
semiSWEET proteins—the bacterial equivalents of SWEET proteins—despite recent studies
identifying sugar transporters in coral-associated bacteria (*26*). This suggests a potential
specialisation within the coral holobiont, where eukaryotes may play a predominant role in
carbohydrate-based energy exchange between symbionts.
383

We identified the expression of SWEET protein genes in *Ostreobium*, providing the first
genomic and transcriptomic evidence of sugar export capabilities in skeleton-dwelling algae.
This discovery suggests that *Ostreobium* can actively transport sugars outside its cells,
making them potentially available for uptake by the coral host and other members of the
coral microbiome. These results highlight a possible alternative energy source for corals,
particularly under stress conditions when the loss of *Symbiodiniaceae* compromises the
availability of photosynthates such as glucose.
391

We identified the expression of SWEET protein genes in *Ostreobium*, providing the first
genomic and transcriptomic evidence of sugar export by skeleton-dwelling algae. This
suggests that *Ostreobium* actively transports sugars into the coral skeleton, potentially
making them available to the coral host and other members of the microbiome. These
findings highlight a potential alternative carbon source for corals, particularly under stress
conditions when *Symbiodiniaceae* are expelled, limiting the availability of photosynthates
(*85*). Previous studies using 14C and 13C-labeled photoassimilates have demonstrated
carbon translocation from endolithic algae into coral host lipids under both healthy and
bleached conditions (*34, 35*). Our findings provide molecular evidence that *Ostreobium*
contributes to this process, underscoring its role in supporting coral energy demands during
stress or bleaching events. This capacity for sugar export may represent a critical
mechanism for maintaining coral resilience by mitigating energy starvation in the absence of
Symbiodiniaceae.
405

Coral-associated fungi are increasingly recognized for their diverse roles, including
influencing reef-scale biogeochemical cycles and shaping coral-associated and
environmental microbiomes through chemical interactions (*86*). In this study, we identified
expression of fungal transcripts encoding SWEET proteins, providing direct molecular
evidence of fungi actively participating in sugar transport in the coral holobiont. This finding
expands our understanding of their role in carbon cycling by demonstrating their contribution
to fixed carbon transfer within the coral holobiont.
413

414 The identification of diverse eukaryotes capable of translocating sugars underscores their
415 importance as a member of the coral probiotic. However, this capability may also pose risks,
416 as sugar-exporting organisms could inadvertently support the proliferation of opportunistic
417 or pathogenic bacteria, thereby potentially harming the coral. Therefore, further research is
418 essential to elucidate the impacts of these sugar-translocating eukaryotes on coral health.
419

420
421
**Fig. 2. Carbon Fixation, Photosynthesis and Sugar Export Across Coral
Compartments.** Heatmap displaying the expression of marker genes for carbon fixation
photosynthesis, and SWEET protein-mediated sugar export across various coral
compartments. Expression values for both were averaged for the count of transcripts and
log2(expression + 1) transformed.

Skeleton   dwelling   microbes   mediate   nitrogen   metabolism

Our analysis identified the expression of nitrogen-fixation marker genes (nifK, nifD, nifH) in
both the Ostreobium layer and the deeper skeleton of *P. lutea*, indicating active nitrogen
fixation in these anoxic environments (Fig. 3A). Nitrogen cycling within the coral holobiont is
indispensable to reef ecology, enabling corals to thrive in nutrient-deficient waters through
efficient nitrogen assimilation and recycling (*31*). While nitrogen-cycling microbes are
consistently detected in coral-associated microbiota, (*31*, *87*) our study highlights

cyanobacteria as the primary nitrogen fixers in *P. lutea*. This finding contrasts earlier reports that heterotrophic bacteria dominate nitrogen fixation in corals (*88–90*) , aligning instead with recent genome-resolved metagenomic studies showing low diversity among nitrogen-fixing bacteria in this species (*24, 25*). The predominance of cyanobacteria in nitrogen fixation emphasises their critical role in the nitrogen economy of the coral holobiont, particularly in supporting nitrogen-limited growth of Symbiodiniaceae (*9, 91*). These results advance our understanding of the microbial interactions underpinning nutrient dynamics and overall reef health.

We also identified nitrification, the aerobic microbial process converting ammonia to hydroxylamine and subsequently to nitrite and nitrate, to be ubiquitous across all coral compartments (**Fig 3A**). This process is crucial for the coral holobiont as it reduces internal ammonium levels, thereby regulating the growth of Symbiodiniaceae and retaining dissolved inorganic nitrogen in an oxidised state, which is essential for preventing the accumulation of toxic ammonia and facilitating the availability of nitrogen for other metabolic processes (*31*). We identified active expression of the nitrification pathway by Thaumarchaeota, Gammaproteobacteria, and Chloroflexi (**Fig 3B and Fig 5**), particularly in the deeper skeleton, enhancing our understanding of the specific organisms involved and their roles in this process.

In contrast, denitrification was confined to the *Ostreobium* layer and deeper skeleton compartments, where we detected expression of the marker genes *napA, nirS, norB, and nosZ* (**Fig. 3A**). This localised expression indicates that denitrification, a key anaerobic process, primarily occurs in low-oxygen zones, particularly in deeper skeleton areas, where conditions are conducive to the reduction of nitrate to nitrogen gases. Nitrate reduction to nitrite via *NR* and *napA* was observed in both eukaryotic (Symbiodiniaceae and Ostreobium) and bacterial symbionts (including Rhodospirillales, Chlorobi, Proteobacteria, and Acidobacteria) (**Fig. 3A**; supplementary data file 3). The subsequent conversion of nitrite to nitric oxide, nitrous oxide, and nitrogen during denitrification was predominantly carried out by Thaumarchaeota, and Proteobacteria (α- and γ-), and Bacteroidota. Importantly, we identified fungal (Ascomycota) transcripts encoding the NIT-6 gene, responsible for reducing nitrite to ammonium via dissimilatory nitrate reduction to ammonium (DNRA), highlighting the role of fungi in nitrogen cycling in coral.

The interplay between nitrification and denitrification, particularly in the skeleton, is crucial for maintaining nitrogen homeostasis within the coral holobiont. The DNRA process, active under low-oxygen conditions (such as at night), converts nitrate back to ammonium, which

may be essential when nitrification, dependent on oxygen, is less active. This coupling
between DNRA and nitrification ensures stable ammonium levels, which could promote coral
growth. On the other hand, coupling denitrification and nitrification may help regulate
nitrogen availability in nutrient-rich conditions, preventing excessive growth of
Symbiodiniaceae and maintaining a balance within the coral ecosystem.

Our results significantly expand current understanding of nitrogen cycling within coral
holobionts by identifying which microbial taxa are actively involved in these processes and
how different compartments, including the coral skeleton, contribute to nitrogen dynamics.
The skeleton, in particular, plays a critical role in both nitrification and denitrification, with
different microbes in at different depths of the skeleton performing complementary functions
that ensure nitrogen homeostasis within the coral host.

Sulphur metabolising microbes are widespread across coral compartments
Sulphur is readily available in seawater as inorganic sulphate and is assimilated by the coral
host, Symbiodiniaceae, and diverse members of the coral microbiome into organic
compounds, thus playing an essential role in the growth of coral holobionts (*92–94*). We
identified expression of transcripts encoding for marker genes involved in sulphur cycling in
all coral compartments (**Fig 3B**). The majority of sulphur cycling marker genes exhibited
higher expression levels in the skeleton, suggesting a predominant role of the endolithic
microbiome. Anaerobic dissimilatory sulphate reduction marker genes (*apr*A and *apr*B) were
exclusively expressed in the coral skeleton, with taxonomic analysis identifying them as
known sulphate-reducing bacteria (SRB), specifically Deltaproteobacteria and anoxygenic
phototrophic Chlorobi (**Fig 3B**). It is important to note that while Chlorobi possess *apr*A and
*apr*B genes, they are primarily involved in sulphur oxidation rather than sulphate reduction
(*95, 96*). Metabarcoding studies have predicted abundant sulphate reducers in the coral
skeleton, and recent metagenomic studies have assembled genomes of SRBs from the
skeleton of *P. lutea* and *I. palifera (24)*. The coral skeleton not only harbours SRBs but also
anoxygenic phototrophs such as Chlorobi, supporting the presence of anoxic microniches in
the coral skeleton (*14, 24, 27, 76*). SRB and Chlorobi have also been shown to form a
syntrophic relationship, where sulphide produced by SRB is used as an electron donor for
photosynthetic Chlorobi in microbial mats, deeper lake waters and coral skeleton (*27, 97–*
99). Our identification of transcripts from SRBs and Chlorobi within the same compartment of
the holobiont provides molecular evidence for such a syntrophic relationship in the coral
skeleton.
507
We observed a complex expression profile of assimilatory sulphate reduction marker genes

(*cys*C, *cys*I, *cys*J, and SUOX), which were expressed in all three compartments; with the

exception of *cys*D, which was exclusively expressed in the deep skeleton layer (**Fig 3B**).

Additionally, the gene encoding the enzyme *cys*K, crucial for linking assimilatory sulphate

reduction with the biosynthesis of the essential amino acid L-Cysteine, was consistently

expressed across all compartments (**Fig 3B**). Taxonomic analysis of transcripts encoding

these ASR marker genes identified a diverse range of symbiont members, including

microeukaryotes such as Symbiodiniaceae, *Ostreobium* and other Chlorophyta, as well as

prokaryotic groups Cyanobacteria, Thaumarchaeota, and Proteobacteria (**Fig 3B**). The

identification of transcripts encoding ASR exclusively annotated to algae and bacteria aligns

with the understanding that this pathway is restricted to plants, algae and microbes(*100*).

520
Fig 3. Transcriptome- based expression profiles of Nitrogen and Sulfur metabolism
across coral compartments. A) Heatmap showing the expression profiles of marker genes
involved in nitrogen metabolism across coral compartments, along with a bar plot indicating
the count of transcripts. The nitrogen cycle is illustrated, highlighting subpathways and
contributing community members. B) Heatmap showing the expression profiles of marker
genes involved in sulphur metabolism across coral compartments, along with a bar plot

527 indicating the count of transcripts. The sulphur metabolism pathway is illustrated, highlighting
528 subpathways and contributing community members. Expression values for both were
529 averaged and log2(expression + 1) transformed.
530
531

532
533 **Figure 5.** Overview of the transcriptionally dominant taxa and their functional contribution in
534 different compartments of the coral identified in this study.

535
536
Microeukaryotes are the major scavengers of ROS and RNS in *P. lutea*
Coral reefs are facing unprecedented threats due to rising sea surface temperatures,
resulting in more frequent and prolonged coral bleaching events (*37*). Most theories on the
onset of coral bleaching suggest that the overproduction of reactive oxygen species (ROS)
and reactive nitrogen species (RNS) is a key trigger, leading to cellular damage and stress
under adverse environmental conditions (*101*). Rapid scavenging of ROS and RNS is thus
crucial in maintaining homeostasis and protecting the coral holobiont in stressed and healthy
corals. We observed widespread expression of seven protein families with functions related
to ROS or RNS scavenging across the tissue, *Ostreobium* layer, and deeper skeleton
compartments. Specifically, transcripts encoding for genes involved in RNS scavenging,
including nitric oxide reductases (PF00175: *hmp*; PF00115 and PF00034: *nor*BC) and
peroxynitrite to nitrate reduction (PF10417: *ahp*C) were expressed in all compartments (**Fig
4**). A single transcript encoding nitrous oxide reductase (PF18764: *nos*Z) was identified, and
expressed exclusively in deeper skeleton samples. This enzyme catalyses the last step of
the dissimilatory metabolic pathway of denitrifying bacteria, which were abundant in the
deeper skeleton. Similar observations were made for ROS scavenging expression profiles.

We identified 48, 43, and 41 transcripts encoding the general ROS scavenger Glutaredoxin
(PF00462), as well as hydrogen peroxide scavengers Peroxiredoxins (PF00578) and
peroxidases (PF00141). These transcripts were expressed in all coral compartments along
with transcripts encoding superoxide dismutases (PF00081 and PF02777: SOD Mn and
PF00080:SOD cu-zn) (**Fig 4**). The majority of ROS and RNS scavenging transcripts were of
micro-eukaryotic origin (76%)**,** including Symbiodiniaceae, *Ostreobium*, Chlorophyta, and
Fungi, while others were from archaeal and bacterial sources (24%), encompassing
Thaumarchaeota, Proteobacteria (Rhodospirillales, Xanthomonadales), Cyanobacteria,
Chlorobi, Chloroflexi and Bacteroidetes (*Candidatus* Amoebophilus) (**Supplementary data**
file   **4**).

Scavenging reactive oxygen (ROS) and nitrogen species (RNS) is a key function of bacterial
candidates for coral probiotics, as emphasised in recent studies identifying potential probiotic
bacteria for corals (*2, 26, 102–104*). However, the identification of the majority of ROS and
RNS scavenging transcripts in micro-eukaryotic inhabitants of the coral holobiont rather than
in bacteria, raises the question of which organisms are the main players in scavenging ROS
and RNS under thermal stress. This finding suggests that current strategies for designing
coral probiotics, which primarily focus on bacterial functions, should be refined through
additional molecular and experimental validation to provide functional insights. These
insights could guide the assembly of a more comprehensive consortium of beneficial
symbionts for corals.

Fig. 4. Transcriptome-based expression profiles of Reactive Oxygen Species (ROS) and Reactive Nitrogen Species (RNS) marker genes across coral compartments. Heatmap displaying the expression of different protein families, count of transcripts for each family involved in ROS and RNS scavenging across coral compartments. Expression values for both were averaged and log2(expression + 1) transformed.

Conclusion

This study provides novel insights into the transcriptionally active communities within the coral holobiont, with a focus on their functional roles across distinct ecological compartments: coral tissue, the endolithic *Ostreobium* layer, and the deeper skeleton layer. Our analysis revealed a nuanced stratification of microbial activity, with microeukaryotes dominating key metabolic processes in the coral tissue and *Ostreobium* layer, while prokaryotes emerge as the primary active symbionts in the deeper skeleton (**Fig 5**). These findings underscore the profound influence of physicochemical gradients—including light penetration, nutrient availability, and carbon distribution—in sculpting both community composition and functional specialization within the holobiont. Functional analysis highlighted the critical contributions of *Symbiodiniaceae, Ostreobium*, and Cyanobacteria to photosynthate production, with *Symbiodiniaceae* predominantly occupying the coral tissue, while *Ostreobium* and Cyanobacteria play pivotal roles in the endolithic *Ostreobium*-layer (**Fig 5**). These symbionts, evolutionarily adapted to varying light conditions and utilizing distinct photosynthetic pigments, demonstrated unique nutrient cycling capabilities. Notably, in addition to *Symbiodiniaceae*, we identified *Ostreobium* and fungi as the only taxa capable of exporting sugars via SWEET proteins, providing novel insights into nutrient exchange mechanisms within the holobiont. Analysis of nitrogen metabolic pathways revealed Cyanobacteria as the exclusive nitrogen fixers, with other nitrogen and sulphur metabolic

processes likely distributed across holobiont members, suggesting a robust functional
redundancy that potentially enhances ecosystem resilience (**Fig 5**). Our analysis also
identified microeukaryotes as the primary scavengers of ROS and RNS across all
compartments, playing a crucial role in maintaining holobiont health, particularly under
thermal stress conditions (**Fig 5**). These findings offer a comprehensive view of the
metabolic activities within the coral microbiome, elucidating the intricate functional roles of its
diverse members. While this research significantly advances our understanding of coral
holobiont functions, it simultaneously highlights the complexity of these systems and the
necessity for more targeted research focussed on the microenvironmental niches of the coral
holobiont. Future investigations should employ advanced sampling techniques and high-
resolution analytical methods to more precisely map the functional dynamics between coral
hosts, their symbionts, and the surrounding microenvironment (*11*, *105*, *106*). Ultimately, the
insights gained from this study are invaluable for advancing coral reef conservation efforts.
By identifying functionally relevant symbiotic members, we can better inform the design of
coral probiotics and other strategies aimed at enhancing coral resilience in the face of
climate change and other environmental stressors.
619

Data and code availability

Raw reads generated in this study have been submitted to NCBI SRA under the Bioproject

PRJNA1207708. All codes and files required to regenerate the figures have been made

available via github.

Acknowledgement

Authors would like to acknowledge staff at the Heron Island Research Station for their

assistance during sample collection. All coral samples were collected under Permit

G22/47504.1 issued to KT and FR by Great Barrier Reef Marine Park Authority, Australia.

Authors would also like to acknowledge reviewers of the manuscript for their suggestions.

Funding

This work was funded through the Australian Research Council grant DP200101613 (to

H.V., L.L.B., M.M., and M.K.), the Faculty of Science (University of Melbourne, to H.V.). K.T.

acknowledges support from the University of Melbourne Early Career Research Grant. M.K.

acknowledges support from the Gordon and Betty Moore Foundation through grant no.

GBMF9206 (https://doi.org/10.37807/GBMF9206). M.M. acknowledges support from NOAA

CRCP: National Oceanic and Atmospheric Administration Coral Reef Conservation Program

NA19NOS4820132. H.V. is supported by the Fundação para a Ciência e a Tecnologia

(CEECIND:2023.06155). Computational analysis of this work was supported by the

University of Melbourne's Research Computing Services and the Petascale Campus

Initiative.

References

1. C. Jaspers, S. Fraune, A. E. Arnold, D. J. Miller, T. C. G. Bosch, C. R. Voolstra,
Consortium of Australian Academy of Science Boden Research Conference
Participants, Resolving structure and function of metaorganisms through a holistic
framework combining reductionist and integrative approaches. *Zoology* **133**, 81–87
(2019).

2. R. S. Peixoto, M. Sweet, H. D. M. Villela, P. Cardoso, T. Thomas, C. R. Voolstra, L. Høj,
D. G. Bourne, Coral Probiotics: Premise, Promise, Prospects. *Annu Rev Anim Biosci* **9**,
265–288 (2021).

3. M. J. H. van Oppen, L. L. Blackall, Coral microbiome dynamics, functions and design in
a changing world. *Nat. Rev. Microbiol.* **17**, 557–567 (2019).

4. E. Rosenberg, O. Koren, L. Reshef, R. Efrony, I. Zilber-Rosenberg, The role of
microorganisms in coral health, disease and evolution. *Nat. Rev. Microbiol.* **5**, 355–362
(2007).

5. F. Rohwer, V. Seguritan, F. Azam, N. Knowlton, Diversity and distribution of coral-
associated bacteria. *Mar. Ecol. Prog. Ser.* **243**, 1–10 (2002).

6. R. V. Thurber, J. P. Payet, A. R. Thurber, A. M. S. Correa, Virus-host interactions and
their roles in coral reef health and disease. *Nat. Rev. Microbiol.* **15**, 205–216 (2017).

7. T. C. LaJeunesse, J. E. Parkinson, P. W. Gabrielson, H. J. Jeong, J. D. Reimer, C. R.
Voolstra, S. R. Santos, Systematic Revision of Symbiodiniaceae Highlights the Antiquity
and Diversity of Coral Endosymbionts. *Curr. Biol.* **28**, 2570–2580.e6 (2018).

8. L. Muscatine, The role of symbiotic algae in carbon and energy flux in reef corals.
Ecosystems of the world **25**, 75–87 (1990).

9. L. A. Morris, C. R. Voolstra, K. M. Quigley, D. G. Bourne, L. K. Bay, Nutrient Availability
and Metabolism Affect the Stability of Coral-Symbiodiniaceae Symbioses. *Trends*
Microbiol. **27**, 678–689 (2019).

10. M. J. Huggett, A. Apprill, Coral microbiome database: Integration of sequences reveals
high diversity and relatedness of coral-associated microbes. *Environ. Microbiol. Rep.*
11, 372–385 (2019).

11. E. Bollati, D. J. Hughes, D. J. Suggett, J.-B. Raina, M. Kühl, Microscale sampling of the
coral gastrovascular cavity reveals a gut-like microbial community. *anim microbiome* **6**
(2024).

12. M. J. Sweet, A. Croquer, J. C. Bythell, Bacterial assemblages differ between
compartments within the coral holobiont. *Coral Reefs* **30**, 39–52 (2011).

13. V. R. Marcelino, M. J. van Oppen, H. Verbruggen, Highly structured prokaryote
communities exist within the skeleton of coral colonies. *ISME J.* **12**, 300–303 (2018).

14. F. Ricci, K. Tandon, M. Moßhammer, E. H.-J. Cho, L. L. Blackall, M. Kühl, H.
Verbruggen, Fine-scale mapping of physicochemical and microbial landscapes of the
coral skeleton. *Environ. Microbiol.* **25**, 1505–1521 (2023).

15. K. Tandon, M. M. Pasella, C. Iha, F. Ricci, J. Hu, C. J. O'Kelly, M. Medina, M. Kühl, H.
Verbruggen, Every refuge has its price: Ostreobium as a model for understanding how
algae can live in rock and stay in business. *Semin. Cell Dev. Biol.* **134**, 27–36 (2023).

16. A. M. Bonacolta, B. A. Weiler, T. Porta-Fitó, M. Sweet, P. Keeling, J. del Campo,

Beyond the Symbiodiniaceae: diversity and role of microeukaryotic coral symbionts.
Coral Reefs **42**, 567–577 (2023).

17. M. McCauley, T. L. Goulet, C. R. Jackson, S. Loesgen, Systematic review of cnidarian
microbiomes reveals insights into the structure, specificity, and fidelity of marine
associations. *Nat. Commun.* **14**, 4899 (2023).

18. R. Rowan, D. A. Powers, Ribosomal RNA sequences and the diversity of symbiotic
dinoflagellates (zooxanthellae). *Proc. Natl. Acad. Sci. U. S. A.* **89**, 3639–3643 (1992).

19. F. J. Pollock, R. McMinds, S. Smith, D. G. Bourne, B. L. Willis, M. Medina, R. V.
Thurber, J. R. Zaneveld, Coral-associated bacteria demonstrate phylosymbiosis and
cophylogeny. *Nat. Commun.* **9**, 4921 (2018).

20. L. Fujise, D. J. Suggett, M. Stat, T. Kahlke, M. Bunce, S. G. Gardner, S. Goyen, S.
Woodcock, P. J. Ralph, J. R. Seymour, N. Siboni, M. R. Nitschke, Unlocking the
phylogenetic diversity, primary habitats, and abundances of free-living Symbiodiniaceae
on a coral reef. *Mol. Ecol.* **30**, 343–360 (2021).

21. L. K. Lee, C. P. Leaw, L. C. Lee, Z. F. Lim, K. S. Hii, A. A. Chan, H. Gu, P. T. Lim,
Molecular diversity and assemblages of coral symbionts (Symbiodiniaceae) in diverse
scleractinian coral species. *Mar. Environ. Res.* **179**, 105706 (2022).

22. V. R. Marcelino, K. M. Morrow, M. J. H. van Oppen, D. G. Bourne, H. Verbruggen,
Diversity and stability of coral endolithic microbial communities at a naturally high pCO_2
reef. *Mol. Ecol.* **26**, 5344–5357 (2017).

23. F. Ricci, A. Fordyce, W. Leggat, L. L. Blackall, T. Ainsworth, H. Verbruggen, Multiple
techniques point to oxygenic phototrophs dominating the Isopora palifera skeletal
microbiome. *Coral Reefs* **40**, 275–282 (2021).

24. K. Tandon, F. Ricci, J. Costa, M. Medina, M. Kühl, L. L. Blackall, H. Verbruggen,
Genomic view of the diversity and functional role of archaea and bacteria in the skeleton
of the reef-building corals Porites lutea and Isopora palifera. *Gigascience* **12** (2022).

25. S. J. Robbins, C. M. Singleton, C. X. Chan, L. F. Messer, A. U. Geers, H. Ying, A.
Baker, S. C. Bell, K. M. Morrow, M. A. Ragan, D. J. Miller, S. Forêt, ReFuGe2020
Consortium, C. R. Voolstra, G. W. Tyson, D. G. Bourne, A genomic view of the reef-
building coral Porites lutea and its microbial symbionts. *Nat Microbiol* **4**, 2090–2100
(2019).

26. T. Doering, K. Tandon, S. H. Topa, S. J. Pidot, L. L. Blackall, M. J. H. van Oppen,
Genomic exploration of coral-associated bacteria: identifying probiotic candidates to
increase coral bleaching resilience in Galaxea fascicularis. *Microbiome* **11**, 185 (2023).

27. S.-H. Yang, K. Tandon, C.-Y. Lu, N. Wada, C.-J. Shih, S. S.-Y. Hsiao, W.-N. Jane, T.-C.
Lee, C.-M. Yang, C.-T. Liu, V. Denis, Y.-T. Wu, L.-T. Wang, L. Huang, D.-C. Lee, Y.-W.
Wu, H. Yamashiro, S.-L. Tang, Metagenomic, phylogenetic, and functional
characterization of predominant endolithic green sulfur bacteria in the coral Isopora
palifera. *Microbiome* **7**, 3 (2019).

28. M. Sweet, H. Villela, T. Keller-Costa, R. Costa, S. Romano, D. G. Bourne, A. Cárdenas,
M. J. Huggett, A. H. Kerwin, F. Kuek, M. Medina, J. L. Meyer, M. Müller, F. J. Pollock,
M. S. Rappé, M. Sere, K. H. Sharp, C. R. Voolstra, N. Zaccardi, M. Ziegler, R. Peixoto,
Insights into the Cultured Bacterial Fraction of Corals. *mSystems* **6**, e0124920 (2021).

29. P. G. Falkowski, Z. Dubinsky, L. Muscatine, J. W. Porter, Light and the Bioenergetics of
a Symbiotic Coral. *Bioscience* **34**, 705–709 (1984).

30. K. Tandon, C.-Y. Lu, P.-W. Chiang, N. Wada, S.-H. Yang, Y.-F. Chan, P.-Y. Chen, H.-Y.
Chang, Y.-J. Chiou, M.-S. Chou, W.-M. Chen, S.-L. Tang, Comparative genomics:
Dominant coral-bacterium Endozoicomonas acroporae metabolizes
dimethylsulfoniopropionate (DMSP). *ISME J.* **14**, 1290–1303 (2020).

31. N. Rädecker, C. Pogoreutz, C. R. Voolstra, J. Wiedenmann, C. Wild, Nitrogen cycling in
corals: the key to understanding holobiont functioning? *Trends Microbiol.* **23**, 490–497
(2015).

32. M. J. Neave, C. T. Michell, A. Apprill, C. R. Voolstra, Endozoicomonas genomes reveal
functional adaptation and plasticity in bacterial strains symbiotically associated with
diverse marine hosts. *Sci. Rep.* **7**, 40579 (2017).

33. D. Schlichter, H. Kampmann, S. Conrady, Trophic potential and photoecology of
endolithic algae living within coral skeletons. *Mar. Ecol.* **18**, 299–317 (1997).

34. M. Fine, Y. Loya, Endolithic algae: an alternative source of photoassimilates during
coral bleaching. *Proceedings of the Royal Society of London. Series B: Biological
Sciences* **269**, 1205–1210 (2002).

35. L. Sangsawang, B. E. Casareto, H. Ohba, H. M. Vu, A. Meekaew, T. Suzuki, T. Yeemin,
Y. Suzuki, 13C and 15N assimilation and organic matter translocation by the endolithic
community in the massive coral Porites lutea. *R Soc Open Sci* **4**, 171201 (2017).

36. E. A. Ottesen, C. R. Young, S. M. Gifford, J. M. Eppley, R. Marin 3rd, S. C. Schuster, C.
A. Scholin, E. F. DeLong, Ocean microbes. Multispecies diel transcriptional oscillations
in open ocean heterotrophic bacterial assemblages. *Science* **345**, 207–212 (2014).

37. T. P. Hughes, J. T. Kerry, S. R. Connolly, J. G. Álvarez-Romero, C. M. Eakin, S. F.
Heron, M. A. Gonzalez, J. Moneghetti, Emergent properties in the responses of tropical
corals to recurrent climate extremes. *Curr. Biol.* **31**, 5393–5399.e3 (2021).

38. M. J. H. van Oppen, J. K. Oliver, H. M. Putnam, R. D. Gates, Building coral reef
resilience through assisted evolution. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 2307–2313
(2015).

39. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

40. E. Kopylova, L. Noé, H. Touzet, SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

41. P. Yilmaz, L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J.
Peplies, W. Ludwig, F. O. Glöckner, The SILVA and "All-species Living Tree Project
(LTP)" taxonomic frameworks. *Nucleic Acids Res.* **42**, D643–D648 (2013).

42. H. R. Gruber-Vodicka, B. K. B. Seah, E. Pruesse, phyloFlash: Rapid Small-Subunit
rRNA Profiling and Targeted Assembly from Metagenomes. *mSystems* **5** (2020).

43. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

44. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis,
R. Durbin, 1000 Genome Project Data Processing Subgroup, The Sequence

Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

45.   M. G. Grabherr, B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X.
Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A.
Gnirke, N. Rhind, F. di Palma, B. W. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman,
A. Regev, Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

46.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

47.   B. Buchfink, K. Reuter, H.-G. Drost, Sensitive protein alignments at tree-of-life scale
using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

48.   D. H. Huson, S. Beier, I. Flade, A. Górska, M. El-Hadidi, S. Mitra, H.-J. Ruscheweyh, R.
Tappu, MEGAN Community Edition - Interactive Exploration and Analysis of Large-
Scale Microbiome Sequencing Data. *PLoS Comput. Biol.* **12**, e1004957 (2016).

49.   B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

50.   P. J. McMurdie, S. Holmes, phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

51.   H. Wickham, ggplot2. *Wiley Interdiscip. Rev. Comput. Stat.* **3**, 180–185 (2011).

52.   D. M. Bryant, K. Johnson, T. DiTommaso, T. Tickle, M. B. Couger, D. Payzin-Dogru, T.
J. Lee, N. D. Leigh, T.-H. Kuo, F. G. Davis, J. Bateman, S. Bryant, A. R. Guzikowski, S.
L. Tsai, S. Coyne, W. W. Ye, R. M. Freeman Jr, L. Peshkin, C. J. Tabin, A. Regev, B. J.
Haas, J. L. Whited, A Tissue-Mapped Axolotl De Novo Transcriptome Enables
Identification of Limb Regeneration Factors. *Cell Rep.* **18**, 762–776 (2017).

53.   C. P. Cantalapiedra, A. Hernández-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas,
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

54.   J. Huerta-Cepas, D. Szklarczyk, D. Heller, A. Hernández-Plaza, S. K. Forslund, H.
Cook, D. R. Mende, I. Letunic, T. Rattei, L. J. Jensen, C. von Mering, P. Bork, eggNOG
5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314 (2019).

55.   M. Kanehisa, Y. Sato, K. Morishima, BlastKOALA and GhostKOALA: KEGG Tools for
Functional Characterization of Genome and Metagenome Sequences. *J. Mol. Biol.* **428**,
726–731 (2016).

56.   P. Jones, D. Binns, H.-Y. Chang, M. Fraser, W. Li, C. McAnulla, H. McWilliam, J.
Maslen, A. Mitchell, G. Nuka, S. Pesseat, A. F. Quinn, A. Sangrador-Vegas, M.
Scheremetjew, S.-Y. Yong, R. Lopez, S. Hunter, InterProScan 5: genome-scale protein
function classification. *Bioinformatics* **30**, 1236–1240 (2014).

57.   G. Salazar, L. Paoli, A. Alberti, J. Huerta-Cepas, H.-J. Ruscheweyh, M. Cuenca, C. M.
Field, L. P. Coelho, C. Cruaud, S. Engelen, A. C. Gregory, K. Labadie, C. Marec, E.
Pelletier, M. Royo-Llonch, S. Roux, P. Sánchez, H. Uehara, A. A. Zayed, G. Zeller, M.
Carmichael, C. Dimier, J. Ferland, S. Kandels, M. Picheral, S. Pisarev, J. Poulain, Tara
Oceans Coordinators, S. G. Acinas, M. Babin, P. Bork, C. Bowler, C. de Vargas, L.
Guidi, P. Hingamp, D. Iudicone, L. Karp-Boss, E. Karsenti, H. Ogata, S. Pesant, S.

Speich, M. B. Sullivan, P. Wincker, S. Sunagawa, Gene Expression Changes and
Community Turnover Differentially Shape the Global Ocean Metatranscriptome. *Cell*
179, 1068–1083.e21 (2019).

58. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

59. L. M. van der Loos, R. Nijland, Biases in bulk: DNA metabarcoding of marine
communities and the methodology involved. *Mol. Ecol.* **30**, 3270–3288 (2021).

60. S. K. Davy, D. Allemand, V. M. Weis, Cell biology of cnidarian-dinoflagellate symbiosis.
Microbiol. Mol. Biol. Rev. **76**, 229–261 (2012).

61. W. K. Kwong, J. Del Campo, V. Mathur, M. J. A. Vermeij, P. J. Keeling, A widespread
coral-infecting apicomplexan with chlorophyll biosynthesis genes. *Nature* **568**, 103–107
(2019).

62. A. Oren, G. M. Garrity, Valid publication of the names of forty-two phyla of prokaryotes.
Int. J. Syst. Evol. Microbiol. **71** (2021).

63. A. Cárdenas, J.-B. Raina, C. Pogoreutz, N. Rädecker, J. Bougoure, P. Guagliardo, M.
Pernice, C. R. Voolstra, Greater functional diversity and redundancy of coral endolithic
microbiomes align with lower coral bleaching susceptibility. *ISME J.* **16**, 2406–2420
(2022).

64. J. R. Guest, J. Low, K. Tun, B. Wilson, C. Ng, D. Raingeard, K. E. Ulstrup, J. T. I. Tanzil,
P. A. Todd, T. C. Toh, D. McDougald, L. M. Chou, P. D. Steinberg, Coral community
response to bleaching on a highly disturbed reef. *Sci. Rep.* **6**, 20717 (2016).

65. P. Buerger, G. M. Schmidt, M. Wall, C. Held, C. Richter, Temperature tolerance of the
coral Porites lutea exposed to simulated large amplitude internal waves (LAIW). *J. Exp.*
Mar. Bio. Ecol. **471**, 232–239 (2015).

66. J. Peng, C.-E. Wegner, Q. Bei, P. Liu, W. Liesack, Metatranscriptomics reveals a
differential temperature effect on the structural and functional organization of the
anaerobic food web in rice field soil. *Microbiome* **6**, 169 (2018).

67. N. Vannier, F. Mesny, F. Getzke, G. Chesneau, L. Dethier, J. Ordon, T. Thiergart, S.
Hacquard, Genome-resolved metatranscriptomics reveals conserved root colonization
determinants in a synthetic microbiota. *Nat. Commun.* **14**, 8274 (2023).

68. S. R. Law, A. R. Serrano, Y. Daguerre, J. Sundh, A. N. Schneider, Z. R. Stangl, D.
Castro, M. Grabherr, T. Näsholm, N. R. Street, V. Hurry, Metatranscriptomics captures
dynamic shifts in mycorrhizal coordination in boreal forests. *Proc. Natl. Acad. Sci. U. S.*
A. **119**, e2118852119 (2022).

69. S. Campana, A. Riesgo, E. Jongepier, J. Fuss, G. Muyzer, J. M. de Goeij, Meta-
transcriptomic comparison of two sponge holobionts feeding on coral- and macroalgal-
dissolved organic matter. *BMC Genomics* **23**, 674 (2022).

70. A. Schoenle, M. Hohlfeld, K. Hermanns, F. Mahé, C. de Vargas, F. Nitsche, H. Arndt,
High and specific diversity of protists in the deep-sea basins dominated by diplonemids,
kinetoplastids, ciliates and foraminiferans. *Commun Biol* **4**, 501 (2021).

71. S. L. Pichard, L. Campbell, J. H. Paul, Diversity of the ribulose bisphosphate
carboxylase/oxygenase form I gene (rbcL) in natural phytoplankton communities. *Appl.*

Environ. Microbiol. **63**, 3600–3606 (1997).

72. B. K. Swan, M. Martinez-Garcia, C. M. Preston, A. Sczyrba, T. Woyke, D. Lamy, T. Reinthaler, N. J. Poulton, E. D. P. Masland, M. L. Gomez, M. E. Sieracki, E. F. DeLong, G. J. Herndl, R. Stepanauskas, Potential for chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. *Science* **333**, 1296–1300 (2011).

73. J. A. Raven, Contributions of anoxygenic and oxygenic phototrophy and chemolithotrophy to carbon and oxygen fluxes in aquatic environments. *Aquat. Microb. Ecol.* **56**, 177–192 (2009).

74. F. Ricci, C. Greening, Chemosynthesis: a neglected foundation of marine ecology and biogeochemistry: (Trends in Microbiology, Published online January 30, 2024). *Trends Microbiol.* **32**, 723 (2024).

75. M. P. Lesser, M. Sabrina Pankey, M. Slattery, K. J. Macartney, D. J. Gochfeld, Microbiome diversity and metabolic capacity determines the trophic ecology of the holobiont in Caribbean sponges. *ISME Commun.* **2**, 112 (2022).

76. M. Kühl, G. Holst, A. W. D. Larkum, P. J. Ralph, IMAGING OF OXYGEN DYNAMICS WITHIN THE ENDOLITHIC ALGAL COMMUNITY OF THE MASSIVE CORAL PORITES LOBATA(1). *J. Phycol.* **44**, 541–550 (2008).

77. R. Samsonoff Wa Maccoll, Biliproteins and phycobilisomes from cyanobacteria and red algae at the extreme habitats. *Arch* **176**, 400–405 (2001).

78. A. de los Ríos, M. Grube, L. G. Sancho, C. Ascaso, Ultrastructural and genetic characteristics of endolithic cyanobacterial biofilms colonizing Antarctic granite rocks: Antarctic endolithic cyanobacterial biofilms. *FEMS Microbiol. Ecol.* **59**, 386–395 (2007).

79. S. H. Magnusson, M. Fine, M. Kühl, Light microclimate of endolithic phototrophs in the scleractinian corals Montipora monasteriata and Porites cylindrica. *Mar. Ecol. Prog. Ser.* **332**, 119–128 (2007).

80. C. Iha, K. E. Dougan, J. A. Varela, V. Avila, C. J. Jackson, K. A. Bogaert, Y. Chen, L. M. Judd, R. Wick, K. E. Holt, M. M. Pasella, F. Ricci, S. I. Repetti, M. Medina, V. R. Marcelino, C. X. Chan, H. Verbruggen, Genomic adaptations to an endolithic lifestyle in the coral-associated alga Ostreobium. *Curr. Biol.* **31**, 1393–1402.e5 (2021).

81. M. Lichtenberg, A. W. D. Larkum, M. Kühl, Photosynthetic Acclimation of Symbiodinium in hospite Depends on Vertical Position in the Tissue of the Scleractinian Coral Montastrea curta. *Front. Microbiol.* **7**, 230 (2016).

82. O. Hoegh-Guldberg, P. J. Mumby, A. J. Hooten, R. S. Steneck, P. Greenfield, E. Gomez, C. D. Harvell, P. F. Sale, A. J. Edwards, K. Caldeira, N. Knowlton, C. M. Eakin, R. Iglesias-Prieto, N. Muthiga, R. H. Bradbury, A. Dubi, M. E. Hatziolos, Coral reefs under rapid climate change and ocean acidification. *Science* **318**, 1737–1742 (2007).

83. N. Rädecker, C. Pogoreutz, H. M. Gegner, A. Cárdenas, F. Roth, J. Bougoure, P. Guagliardo, C. Wild, M. Pernice, J.-B. Raina, A. Meibom, C. R. Voolstra, Heat stress destabilizes symbiotic nutrient cycling in corals. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

84. J. Ji, L. Yang, Z. Fang, Y. Zhang, M. Zhuang, H. Lv, Y. Wang, Plant SWEET Family of Sugar Transporters: Structure, Evolution and Biological Functions. *Biomolecules* **12** (2022).

85. O. Hoegh-Guldberg, Climate change, coral bleaching and the future of the world's coral
reefs. *Mar. Freshwater Res.* **50**, 839–866 (1999).

86. A. Roik, M. Reverter, C. Pogoreutz, A roadmap to understanding diversity and function
of coral reef-associated fungi. *FEMS Microbiol. Rev.* **46** (2022).

87. A. R. Mohamed, M. A. Ochsenkühn, A. M. Kazlak, A. Moustafa, S. A. Amin, The coral
microbiome: towards an understanding of the molecular mechanisms of coral-microbiota
interactions. *FEMS Microbiol. Rev.* **47** (2023).

88. N. D. Olson, T. D. Ainsworth, R. D. Gates, M. Takabayashi, Diazotrophic bacteria
associated with Hawaiian Montipora corals: Diversity and abundance in correlation with
symbiotic dinoflagellates. *J. Exp. Mar. Bio. Ecol.* **371**, 140–146 (2009).

89. K. A. Lema, B. L. Willis, D. G. Bourne, Amplicon pyrosequencing reveals spatial and
temporal consistency in diazotroph assemblages of the Acropora millepora microbiome.
Environ. Microbiol. **16**, 3345–3359 (2014).

90. K. A. Lema, D. G. Bourne, B. L. Willis, Onset and establishment of diazotrophs and
other bacterial associates in the early life history stages of the coral Acropora millepora.
Mol. Ecol. **23**, 4682–4695 (2014).

91. C. Cook, C. D'Elia, Are natural populations of zooxanthellae ever nutrient-limited?
Symbiosis **4**, 199–211 (1987).

92. J. Li, Q. Yang, J. Dong, M. Sweet, Y. Zhang, C. Liu, Y. Zhang, X. Tang, W. Zhang, S.
Zhang, Microbiome Engineering: A Promising Approach to Improve Coral Health. *Proc.*
Est. Acad. Sci. Eng. **28**, 105–116 (2023).

93. M. A. Moran, B. P. Durham, Sulfur metabolites in the pelagic ocean. *Nat. Rev. Microbiol.*
17, 665–678 (2019).

94. T. Higuchi, K. Tanaka, K. Shirai, I. Yuyama, T. Mezaki, N. Takahata, Y. Sano, Sulfur
assimilation in corals with aposymbiotic and symbiotic zooxanthellae. *Environ. Microbiol.*
Rep. **13**, 98–103 (2021).

95. B. Meyer, J. Kuever, Homology modeling of dissimilatory APS reductases (AprBA) of
sulfur-oxidizing and sulfate-reducing prokaryotes. *PLoS One* **3**, e1514 (2008).

96. T. Watanabe, H. Kojima, M. Fukui, Identity of major sulfur-cycle prokaryotes in
freshwater lake ecosystems revealed by a comprehensive phylogenetic study of the
dissimilatory adenylylsulfate reductase. *Sci. Rep.* **6**, 36262 (2016).

97. J. F. Imhoff, Biology of green sulfur bacteria, *eLS* (2014).
https://doi.org/10.1002/9780470015902.a0000458.pub2.

98. C. Ng, M. Z. DeMaere, T. J. Williams, F. M. Lauro, M. Raftery, J. A. E. Gibson, C.
Andrews-Pfannkoch, M. Lewis, J. M. Hoffman, T. Thomas, R. Cavicchioli,
Metaproteogenomic analysis of a dominant green sulfur bacterium from Ace Lake,
Antarctica. *ISME J.* **4**, 1002–1019 (2010).

99. Y.-H. Chen, S.-H. Yang, K. Tandon, C.-Y. Lu, H.-J. Chen, C.-J. Shih, S.-L. Tang,
Potential syntrophic relationship between coral-associated Prosthecochloris and its
companion sulfate-reducing bacterium unveiled by genomic analysis. *Microb Genom* **7**
(2021).

100.   I. Kushkevych, J. Cejnar, J. Treml, D. Dordević, P. Kollar, M. Vítězová, Recent

Advances in Metabolic Pathways of Sulfate Reduction in Intestinal Bacteria. *Cells* **9**
(2020).

101.   J. Helgoe, S. K. Davy, V. M. Weis, M. Rodriguez-Lanetty, Triggers, cascades, and
endpoints: connecting the dots of coral bleaching mechanisms. *Biol. Rev. Camb. Philos.*
Soc. **99**, 715–752 (2024).

102.   A. M. Dungan, D. Bulach, H. Lin, M. J. H. van Oppen, L. L. Blackall, Development of
a free radical scavenging bacterial consortium to mitigate oxidative stress in cnidarians.
Microb. Biotechnol. **14**, 2025–2040 (2021).

103.   N. Delgadillo-Ordoñez, N. Garcias-Bonet, I. Raimundo, F. C. García, H. Villela, E. O.
Osman, E. P. Santoro, J. Curdia, J. G. D. Rosado, P. Cardoso, A. Alsaggaf, A. Barno,
C. P. Antony, C. Bocanegra, M. L. Berumen, C. R. Voolstra, F. Benzoni, S. Carvalho, R.
S. Peixoto, Probiotics reshape the coral microbiome in situ without detectable off-target
effects in the surrounding environment. *Commun Biol* **7**, 434 (2024).

104.   C. R. Voolstra, J.-B. Raina, M. Dörr, A. Cárdenas, C. Pogoreutz, C. B. Silveira, A. R.
Mohamed, D. G. Bourne, H. Luo, S. A. Amin, R. S. Peixoto, The coral microbiome in
sickness, in health and in a changing world. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-
024-01015-3 (2024).

105.   M. J. H. van Oppen, J.-B. Raina, Coral holobiont research needs spatial analyses at
the microbial scale. *Environ. Microbiol.* **25**, 179–183 (2023).

106.   D. J. Hughes, J.-B. Raina, D. A. Nielsen, D. J. Suggett, M. Kühl, Disentangling
compartment functions in sessile marine invertebrates. *Trends Ecol. Evol.* **37**, 740–748
(2022).

963