1

Aggregation of HAPLN2, a component of the perinodal extracellular matrix, is a hallmark of physiological brain aging in mice

Ayaka Watanabe[1][¶], Shoshiro Hirayama[1][¶], Itsuki Kominato[1], Sybille Marchese[2], Pietro

Esposito[2], Vanya Metodieva[2], Hiroshi Kameda[3], Terunori Sano[4], Sho Takatori[5], Masaki

Takao[4], Masato Koike[3], Juan Alberto Varela[6], Taisuke Tomita[5], Shigeo Murata[1]*

[1] Laboratory of Protein Metabolism, Graduate School of Pharmaceutical Sciences, the
University of Tokyo, Tokyo, Japan

[2] School of Biology, University of St Andrews, St Andrews, United Kingdom

[3] Department of Cell Biology and Neuroscience, Juntendo University Graduate School of
Medicine, Tokyo, Japan

[4] Department of Clinical Laboratory, National Centre of Neurology and Psychiatry,
Tokyo, Japan

[5] Laboratory of Neuropathology and Neuroscience, Graduate School of Pharmaceutical
Sciences, the University of Tokyo, Tokyo, Japan

[6] School of Physics and Astronomy, University of St Andrews, St Andrews, United
Kingdom

* Corresponding author

Email: smurata@g.ecc.u-tokyo.ac.jp (SMurata)

¶These authors contributed equally to this work.

Abstract

Protein aggregation is a hallmark of neurodegenerative diseases and is also observed in the brains of elderly individuals without such conditions, suggesting that aging drives the accumulation of protein aggregates. However, the comprehensive understanding of age-dependent protein aggregates involved in brain aging remains unclear. Here, we investigated proteins that become sarkosyl-insoluble with age and identified hyaluronan and proteoglycan link protein 2 (HAPLN2), a hyaluronic acid-binding protein of the extracellular matrix at the nodes of Ranvier, as an age-dependent aggregating protein in mouse brains. Elevated hyaluronic acid levels and impaired microglial function reduced the clearance of HAPLN2, leading to its accumulation. HAPLN2 oligomers induced microglial inflammatory responses both *in vitro* and *in vivo*. Furthermore, age-associated HAPLN2 aggregation was also observed in the human cerebellum. These findings suggest that HAPLN2 aggregation results from age-related decline in brain homeostasis and may exacerbate the brain environment by activating microglia. This study provides new insights into the mechanisms underlying cerebellar aging and highlights the role of HAPLN2 in age-associated changes in the brain.

Author Summary

To identify age-dependent protein aggregates unrelated to disease, we analyzed the sarkosyl-insoluble proteome of young and aged mouse brains. We discovered that hyaluronan and proteoglycan link protein 2 (HAPLN2), a hyaluronic acid-binding protein existing in the nodal extracellular matrix, accumulated with age. Age-dependently accumulated HAPLN2 formed large protein aggregates that cannot be solubilized by the anionic detergent sarkosyl or by hyaluronidase digestion. In addition, HAPLN2 formed irregularly shaped puncta that were mislocalized from the nodes of Ranvier in the cerebellar white matter of not only aged mice but also aged human brains. Oligomers of full-length HAPLN2 specifically induced microglial activation *in vitro* and *in vivo*. Our findings suggest that the accumulation of HAPLN2 aggregates is a new hallmark of brain aging and a possible factor contributing to brain inflammation.

Introduction

Protein aggregation is associated with a wide range of physiological declines, including neurodegenerative diseases, with amyloid β (Aβ) and tau being prominent examples. These proteins aggregate to cause neurodegenerative diseases such as Alzheimer's disease, disrupting neuronal function and ultimately causing cell death. Intracellular aggregates interfere with essential cellular processes such as protein folding, trafficking, and degradation, resulting in the accumulation of misfolded proteins and cellular stress [1]. This disruption impairs crucial neuronal functions, including transcription [2], mitochondrial function [3], and stress granule dynamics [4, 5]. In neurodegenerative diseases, the accumulation of protein aggregates is also observed in the extracellular space. These aggregates activate microglia, triggering the release of pro-inflammatory cytokines and neurotoxic factors, leading to chronic inflammation detrimental to neuronal health [6].

The impairment of clearance systems in both intracellular and extracellular spaces is thought to be responsible for the accumulation of aberrant protein aggregates. The main intracellular pathways for clearing abnormal proteins are the ubiquitin-proteasome system [7] and the autophagy-lysosome system [8, 9]. Additionally, extracellular mechanisms such as phagocytosis and secretion of proteases by microglia [10, 11], and the glymphatic system contribute to the clearance of protein aggregates from the brain [12]. However, aging is associated with a decline in the capacity of these mechanisms to maintain proteostasis, making organisms more susceptible to protein aggregation. This decline is attributed to factors such as reduced efficiency of protein degradation pathways, decreased expression of chaperone proteins, impaired cellular stress responses, and microglial senescence [13, 14].

Interestingly, aggregates of wild-type Aβ, tau, and TMEM106B have been observed in the brains of non-demented elderly individuals [15–17]. This suggests that protein aggregation is not merely a consequence of neurodegenerative diseases but also an inherent aspect of the aging process itself, raising the possibility that age-dependent protein aggregates contribute to the broader physiological decline during aging.

Indeed, the detrimental effects of age-related protein aggregation have been demonstrated

in model organisms. In *Caenorhabditis elegans*, the accumulation of both intracellular

and extracellular protein aggregates is associated with reduced lifespan, regardless of

disease state. In contrast, removing these aggregates extend lifespan. For example,

reduced daf-2 function suppresses protein aggregation and extends lifespan, whereas

deficiency in daf-16 or hsf-1 induces protein aggregation and shortens lifespan [18–22].

Furthermore, analysis of RIPA buffer-insoluble and sodium deoxycholate-insoluble

proteins by mass spectrometry has revealed that protein aggregation occurs extensively

during physiological aging in the absence of disease [22, 23]. These findings suggest that

the clearance of age-related protein aggregates in *C. elegans* is crucial for maintaining

individual health.

An extracellular protein quality control mechanism exists that disposes of extracellular

waste in *C. elegans*. Coelomocytes remove extracellular debris through non-specific

phagocytosis [24]. Moreover, overexpression of extracellular chaperones in the

pseudocoelom suppresses the accumulation of aggregates and extends lifespan [25].

These findings highlight the importance of extracellular protein quality control and

emphasize the essential role to maintain proteostasis in the extracellular space to ensure

healthy aging.

In contrast to these studies about *C. elegans*, less is known about how age-dependent

protein aggregates contribute to the aging process in mammals. To address this gap, we

employed proteomic analyses to identify proteins that aggregate and accumulate in the

mouse brain with age. Our investigation identified hyaluronan and proteoglycan link

protein 2 (HAPLN2), a perinodal extracellular matrix constituent protein, as an age-

dependent protein aggregate. We found that HAPLN2 accumulates in the cerebellar white

matter and is mislocalized from its typical location at the nodes of Ranvier in aged mouse

brains. Further investigation using recombinant protein revealed that HAPLN2 forms

aggregates under acidic conditions and in the presence of hyaluronic acid, mimicking

age-related changes in the brain microenvironment. Importantly, we also observed age-

dependent HAPLN2 aggregation in human brain sections. These findings suggest that

HAPLN2 aggregation may be a conserved hallmark of brain aging across mammalian

species. Our study highlights the importance of investigating age-related protein aggregation beyond the context of specific neurodegenerative diseases and suggests a potential role for HAPLN2 aggregates in brain aging, possibly by inducing microglial activation and neuroinflammation.

Results

Proteomic analyses of aged mouse brains identified proteins that accumulate in the sarkosyl-insoluble fraction with age

To identify proteins that aggregate in an age-dependent manner, we enriched detergent-insoluble proteins from the whole brains of young (2-month-old) and aged (24-month-old) mice using 1% N-lauryl-sarcosine (sarkosyl) fractionation (Fig 1A). Sarkosyl is a well-established detergent that solubilizes natively folded proteins while enriching protein aggregates associated with neurodegenerative diseases in its insoluble P2 fraction (Fig 1A) [26–31]. Proteins in the sarkosyl-insoluble P2 fraction were subjected to label-free quantitative proteomics. We identified 31 proteins that increased more than 5-fold in the brains of aged mice compared to young mice as candidates for age-dependent protein aggregates (S1 Table).

Age-related accumulation of proteins associated with neurodegenerative diseases is attributed to their inherent resistance to degradation due to aggregation and age-related declines in clearance mechanisms [32, 33]. To identify proteins that accumulate with age, we performed semi-quantitative mass spectrometry analysis using tandem mass tag (TMT) labeling. Whole brains from young and aged mice were lysed with phase transfer surfactant (PTS) buffer (0.5% deoxycholic acid and 0.33% sarkosyl), which increases the solubility of hydrophobic proteins and enhances their trypsin digestion [34]. The PTS-soluble supernatant, containing most of the brain proteome (Fig S1), was subjected to TMT-based proteomic analysis (Fig 1B). We identified 45 proteins that significantly increased by more than 1.2-fold in the aged mouse brains compared to young mouse

brains (S2 Table). Finally, we narrowed down nine proteins showing an age-dependent

increase in both proteomic analyses (Fig 1C).

To elucidate the characteristics of these proteins enriched in both proteomic analyses, we

performed gene ontology analysis (Fig 1D). The term "hyaluronic acid binding" was

significantly enriched, encompassing aggrecan core protein (ACAN), hyaluronan and

proteoglycan link protein (HAPLN) 1, HAPLN2, and versican core protein (VCAN),

which were previously reported to increase with age in rodent brains [35–37]. Therefore,

we focused on these proteins as candidates for proteins that aggregate and accumulate

with age.

Figure 1. Mass Spectrometry Analyses Revealed Sarkosyl-Insoluble Proteins.

Accumulated in Aged Mouse Brains. (A) Workflow (left) illustrating the fractionation of

young (2-month-old) and aged (24-month-old) mouse brains to obtain sarkosyl-insoluble

protein aggregates. A volcano plot (right) represents the results of label-free quantitative

proteomic analysis. The cut-off was set at a false discovery rate (FDR) of 0.01. The

colored region in the volcano plot highlights proteins with a fold change > 5.0 and P-

value < 0.01, as listed in S1 Table. n = 4. Specific proteins are highlighted: ACAN

(green), HAPLN1 (orange), HAPLN2 (red), and VCAN (blue). (B) Tandem mass tag

(TMT)-based mass spectrometry analysis of phase transfer surfactant (PTS)-soluble

supernatant from young (2-month-old) and aged (24-month-old) mouse brains. The cut-

off was set at FDR = 0.01. The colored regions in the volcano plots highlight proteins

with a fold change > 1.2 and P-value < 0.05, as listed in S2 Table. n = 4. (C) A list of

proteins significantly increased in both proteomic analyses described in (A) and (B). (D)

Bar chart showing gene ontology (GO) terms in the molecular function category enriched

among the proteins identified in (C). GO enrichment analysis was performed using the

Database for Annotation, Visualization, and Integrated Discovery (DAVID). Proteins

included in the enriched categories are indicated on the right side of the chart.

HAPLN1 and HAPLN2 formed salkosyl-insoluble protein assemblies in aged mouse brains

To validate the proteomic findings, we performed immunoblotting on each fraction prepared as shown in Figure 1A. Consistent with the proteomic analyses, marked age-dependent increases in ACAN, HAPLN1, and HAPLN2 were observed in the sarkosyl-insoluble P2 fraction, while we were unable to assess VCAN due to a lack of detectable antibodies (Figs 2A and 2B). As expected from previous studies, ubiquitinated proteins also increased in the P2 fraction with age (Figs 2A and 2B), confirming the successful enrichment of protein aggregates [35]. These results also demonstrated the sarkosyl-insolubility of ACAN, HAPLN1, and HAPLN2 in aged mouse brains.

Characteristics of protein aggregates include not only detergent-insolubility but also the formation of high-molecular-weight assemblies trapped on nitrocellulose membranes and aberrant localization as large deposits *in vivo* [36, 37]. Therefore, we performed a series of experiments to determine whether ACAN, HAPLN1, and HAPLN2 are aggregated in aged mouse brains.

First, to investigate whether ACAN, HAPLN1, and HAPLN2 form sarkosyl-insoluble large protein assemblies and are trapped on a nitrocellulose membrane, we performed a filter trap assay using the S1 fraction. The proteins that passed through the nitrocellulose membranes were trapped by the PVDF membrane layered underneath. HAPLN1 and HAPLN2 trapped on the nitrocellulose membrane showed a marked increase with age. In contrast, the amount of ACAN trapped on nitrocellulose membranes did not differ between young (3-month-old) and aged (21-month-old) mice (Fig 2C). The amounts of ACAN, HAPLN1, and HAPLN2 captured in the PVDF membrane were all increased in the S1 fractions of aged mice. These results suggest that HAPLN1 and HAPLN2 formed sarkosyl-insoluble large protein complexes with age, whereas ACAN did not.

ACAN, HAPLN1, and HAPLN2 are known to form the extracellular matrix through non-covalent association with hyaluronic acid. A previous study showed that ACAN extracted from the aged mouse brain was enriched in the Triton X-100-insoluble fraction but was solubilized by degrading high-molecular-weight hyaluronic acid. [38]. Therefore, we next

investigated whether treatment with hyaluronidase solubilizes ACAN, HAPLN1, and HAPLN2 from the sarkosyl-insoluble P2 fraction. We treated the sarkosyl-insoluble P2 fraction with hyaluronidase, followed by ultracentrifugation to yield the supernatant hyaluronidase-extractable S3 and the pellet hyaluronidase-resistant P3 fraction (Fig 2D). ACAN was recovered in the S3 fraction after hyaluronidase treatment (Fig 2E). In contrast, the majority of HAPLN1 and HAPLN2 remained in the P3 fraction, with only a portion shifting to the S3 fraction (Fig 2E). We also performed a filter trap assay on the S3 and P3 fractions. HAPLN1 and HAPLN2 in the P3 fraction were captured on a nitrocellulose membrane, whereas ACAN in the P3 fraction was not captured on a nitrocellulose membrane (Fig 2F). These results suggest that HAPLN1 and HAPLN2 form high molecular weight protein assemblies that remain insoluble in sarkosyl despite hyaluronic acid degradation. We focused on HAPLN2 for further investigation because HAPLN2 showed the highest fold change of the nine proteins in both proteomic analyses (Fig 1C).

Figure 2. Extracellular Matrix Component Proteins Accumulate in the Sarkosyl-Insoluble Fraction of Mouse Brains with Age. (A) Immunoblot analyses of ACAN, HAPLN1, HAPLN2, and ubiquitin in the S1, P1, S2, and P2 fractions, as outlined in Figure 1A, from mouse brains. n = 2. (B) Densitometric quantitation of (A). The expression levels of the indicated proteins were normalized to α-tubulin in the S1 fractions. Error bars represent mean ± S.D. *P*-values by Student's *t*-test. n = 4. (C) Filter trap assay and immunoblot analysis of the S1 fraction from young (3-month-old) and aged (21-month-old) mouse brains. n = 2. (D) Diagram illustrating hyaluronidase treatment of the sarkosyl-insoluble P2 fraction. The P2 fraction was treated with hyaluronidase and centrifuged at $100{,}000 \times g$. (E) Immunoblot analyses of ACAN, HAPLN1, and HAPLN2 in the hyaluronidase-extractable S3 and hyaluronidase-resistant P3 fractions from 27-month-old mouse brains. n = 3. (F) Filter trap assay of the fractions described in (E). n = 3.

HAPLN2 was deposited in the extracellular space in the cerebellar white matter of aged mice

To investigate whether HAPLN2 forms aberrant deposits in aged mouse brains, we examined the distribution and morphological structure of HAPLN2 in mouse brains. Previous studies have shown that HAPLN2 is expressed in the cerebellar white matter of young adult mice [39]. Immunoblot analysis confirmed that the HAPLN2 protein was predominantly detected in the cerebellum of 24-month-old mice, whereas it was only faintly detected in the cerebellum of 4-month-old mice (Fig 3A). Immunohistochemical analysis revealed that the expression of HAPLN2 increased with age in the cerebellar white matter, which is mainly composed of myelinated axons expressing myelin basic protein (MBP) and is surrounded by granular layers stained with 4',6-diamidino-2-phenylindole (DAPI) (Fig 3B). These results demonstrate that HAPLN2 accumulates in the cerebellar white matter of aged mice.

We next examined its detailed structure in the cerebellar white matter of aged mice. Immunohistochemistry analysis showed significantly enlarged puncta of HAPLN2 in the aged mouse brains compared to young mouse brains (Figs 3C and 3D). We then investigated whether the HAPLN2 deposits are localized to the intracellular or extracellular space. Hyaluronic acid is a major component of the extracellular matrix, which is visualized by staining with hyaluronic acid binding protein (HABP). In young mouse samples, HABP staining was weak, and colocalization of HAPLN2 and HABP was unclear. In contrast, in aged mouse brains, both HABP and HAPLN2 staining showed significantly enlarged puncta, and colocalization of HAPLN2 and HABP was clearly observed (Figs 3E–G, and S2A). These results indicate that HAPLN2 accumulates as large deposits with hyaluronic acid in the extracellular space of the cerebellar white matter of aged mice.

To determine which cells in the cerebellum express HAPLN2 and whether the accumulation of HAPLN2 is due to increased mRNA expression or not, we referred to the results of a previous single-cell transcriptome analysis. This data revealed that *Hapln2* mRNA expression was specifically detected in oligodendrocytes and that mRNA levels

remained constant between 2–3 months and 21–22 months of age (Fig 3H) [40]. These

results suggest that the age-dependent accumulation of HAPLN2 protein is not due to the

induction of mRNA expression.

Since the single-cell RNA-seq result indicate that HAPLN2 is expressed in

oligodendrocytes, we performed co-staining of HAPLN2 and the oligodendrocyte marker

MBP [41]. However, the HAPLN2 deposits did not colocalize with oligodendrocytes (Fig

3I). We also examined co-staining of HAPLN2 with ubiquitin or p62, since protein

aggregates that forms intracellular aggregation and then are deposited extracellularly, as

is typically observed with tau, often colocalize with these proteins [42]. However,

HAPLN2 did not colocalize with ubiquitin and p62 (Figs S2B and S2C). These results

suggest that HAPLN2 is secreted from oligodendrocytes and then accumulated in the

extracellular space.

We next investigated whether enlarged HAPLN2 puncta retain their physiological

function. HAPLN2 is normally localized to the perinodal extracellular matrix, which

surrounds the nodes of Ranvier and contains hyaluronic acid [43]. Therefore, we

examined the colocalization between HAPLN2 and contactin-associated protein (Caspr),

which localizes to the nodes of Ranvier. The colocalization of HAPLN2 with Caspr

decreased with age (Figs 3J and 3K). In addition, Mander's overlap coefficient (MOC)

M1, which indexes the proportion of HAPLN2 colocalized with Caspr, decreased with

age (Fig 3L), suggesting that HAPLN2 was abnormally localized outside the nodes of

Ranvier in aged mice. In contrast, MOC M2, an index of the proportion of Caspr

colocalized with HAPLN2, did not change with age (Fig 3M), suggesting that most of

Caspr colocalizes with functional HAPLN2 and that the nodes of Ranvier are not

dysregulated in aged mice. Additionally, the area of HAPLN2 mislocalized from Caspr

was larger than HAPLN2 colocalized with Caspr in 24-month-old mice (Fig 3N). This

further supports that enlarged HAPLN2 puncta have lost their function.

Figure 3. HAPLN2 Forms Aggregate-Like Structures in the Cerebellar White

Matter of Aged Mice. (A) Immunoblot analysis of the S1 fraction from the cerebrum

and cerebellum of mice. n = 2. (B) Representative images of fluorescence

immunohistochemistry of the mouse cerebellum showing HAPLN2, myelin basic protein

(MBP, a marker for myelinated axons), and 4',6-diamidino-2-phenylindole (DAPI). n = 3.

(C) Magnified fluorescence immunohistochemistry images for HAPLN2 corresponding

to the boxed areas in the left panels. n = 3. (D) Quantitation of the area of each HAPLN2-

positive punctum in (C). (E) Representative images of fluorescence

immunohistochemistry for HAPLN2 and biotinylated hyaluronic acid-binding protein

(HABP), a probe for hyaluronic acid. n = 3. (F) Quantitation of the area of each HABP-

positive punctum. (G) Pearson's correlation coefficients for colocalization between

HAPLN2 and HABP. (H) Single-cell transcriptomics data from Ximerakis et al., *Nat*

Neurosci (2019). TPM: transcripts per million. ASC: astrocyte; MG: microglia; NEUR:

mature neuron; OLG: oligodendrocyte. Error bars indicate maximum values, boxes

indicate the first and third quartiles, and solid and dashed lines represent the median and

mean, respectively. (I) Representative images of fluorescence immunohistochemistry for

HAPLN2 and MBP as an oligodendrocyte marker in the cerebellar white matter. n = 3.

(J) Representative images of fluorescence immunohistochemistry for HAPLN2 and

Caspr, a marker for the nodes of Ranvier. n = 3. (K–M) Colocalization analysis between

HAPLN2 and Caspr using Pearson's correlation coefficient (K) and Mander's overlap

coefficients (MOC) M1 (L) and M2 (M). MOC M1 represents the proportion of HAPLN2

overlapping Caspr, while MOC M2 represents the reverse. (N) Quantitation of the area of

each HAPLN2 punctum, categorized by whether it merges with Caspr in 24-month-old

mice. All coefficient calculations were performed using ImageJ (version 1.53t) with the

JACoP plugin. Scale bars: 1 mm (B and C (left)) and 10 μm (C (right); E; I; J). Error bars

represent mean ± S.D. *P*-values by Student's *t*-test.

Deposition of HAPLN2 aggregates starts in middle age in mouse brains

Research in humans and rodents has highlighted middle age as a period marked by the

onset of changes associated with brain aging [44]. To investigate whenHAPLN2 starts to

accumulate, we examined the age-related progression of HAPLN2 accumulation in the

mouse brain. We performed sarkosyl fractionation followed by immunoblot analysis

using young (3- and 6-month-old), middle-aged (12-month-old), and aged (18-month-old)

mice. HAPLN2 began to accumulate in the S1 and P2 fractions from 12-month-old mice

and was further enriched in 18-month-old mice (Fig 4A). Correspondingly, the puncta of

HAPLN2 became larger in 12- and 18-month-old mice, consistent with the immunoblot

analysis (Figs 4B and 4C). These results indicate that HAPLN2 accumulates gradually

from middle age to old age, further confirming the age-dependent accumulation of

HAPLN2 aggregates.

Figure 4. Time-Course Analysis of Age-Dependent HAPLN2 Accumulation in

Mouse Brains. (A) Immunoblot analyses of the S1 and P2 fractions from the mouse

cerebellum at different ages. n = 3. (B) Representative images of fluorescence

immunohistochemistry of the mouse cerebellum showing HAPLN2 and DAPI staining. n

= 3. (C) Quantitation of the area of each HAPLN2-positive structure shown in (B). Error

bars represent mean ± S.D. *P*-values by one-way ANOVA followed by Tukey's *post hoc*

test.

Low pH, high NaCl concentration, and hyaluronic acid promoted HAPLN2

aggregation

To investigate the factors affecting HAPLN2 aggregation, we explored the aggregation

propensity of recombinant HAPLN2 protein. Various factors have been reported to

influence the formation of protein aggregates, including pH and NaCl concentration. For

instance, the TAR DNA-binding protein of 43 kDa (TDP-43) aggregates at high pH and

NaCl concentration [45], while fused in sarcoma (FUS) aggregates at low concentrations

of RNA [46] and in the presence of dextran [47, 48] and NaCl [49]. Therefore, we

incubated recombinant HAPLN2 protein in buffers with varying pH and NaCl

concentration, and in the presence or absence of hyaluronic acid. The formation of

amorphous aggregates of HAPLN2 was promoted at low pH, high NaCl concentration,

and in the presence of hyaluronic acid (Figs 5A, 5B, S3A, and S3B). Quantitative area

measurements revealed that recombinant HAPLN2 formed amorphous aggregates larger

than 10 μm^2 at pH 6.0, while it formed small oligomer-like aggregates less than 10 μm^2

at pH 7.0, and no aggregates were formed at pH 8.0 in 150 mM NaCl (Fig S3C).

Transmission electron microscopy analysis further confirmed that HAPLN2 incubated at

pH 6.0 in 150 mM NaCl formed amorphous aggregates (Fig S3D).

Many protein aggregates contain cross-β-sheet structures and form protein fibrils.

However, it is impossible to distinguish between oligomeric and fibrillar structures using

diffraction-limited optics [50]. To examine whether HAPLN2 aggregates contain β-sheet-

mediated aggregates that are stained with thioflavin T or Nile Red, we conducted super-

resolution imaging analysis (Fig S3E) [51]. In recombinant HAPLN2 incubated at pH

6.0, some spots, but not the entire aggregate, were stained with thioflavin T and Nile Red,

indicating that the amorphous-appearing HAPLN2 aggregates contain cross-β-sheet

structures (Fig 5C).

To analyze whether the HAPLN2 aggregates formed fibrillar structures, which are

characterized by a long axis and low circularity ratio, we statistically analyzed the

morphology of the Nile Red-positive clustered aggregates, according to a previous report

[52]. The long axis of the Nile Red-positive clusters increased under acidic conditions

(Fig 5D), but the circularity ratio did not change with pH (Fig 5E). These results suggest

that low pH induces the formation of amorphous, rather than fibrillar, aggregates of

HAPLN2 containing partial cross-β-sheet structures.

Figure 5. Aggregates of Recombinant HAPLN2 Formed Under Low pH, High NaCl,

or Hyaluronic Acid Conditions Partially Contain Cross-β-Sheet Structures. (A)

Brightfield microscopic images of recombinant HAPLN2 aggregates incubated in the

indicated buffer conditions without agitation. (B) Phase diagrams showing the

aggregation states of recombinant HAPLN2 under the indicated buffer conditions. (C)

Brightfield and highly inclined and laminated optical sheet (HiLo) images of recombinant

HAPLN2 aggregates incubated in a buffer at the indicated pH with 150 mM NaCl.

Thioflavin T and Nile Red were used at concentrations of 5 µM and 5 nM, respectively to

visualize cross-β-sheet structures. (D, E) Quantitative analyses of the long axis (D) and

circularity ratio (E) of Nile Red-positive aggregates shown in (C). Data were collected

from at least five fields of view. 500 nM Recombinant HAPLN2 monomers were

incubated for 24 hours at 37ºC in all conditions. Error bars represent mean ± S.D. *P*-

values by one-way ANOVA followed by Tukey's *post hoc* tests. Scale bars: 10 µm (A), 1

µm (C).

Treatment with hyaluronidase promoted the clearance of HAPLN2 aggregates in the mouse cerebellar white matter

It has been reported that the volume of the extracellular spaces reduces with age, which is

thought to lead to impairment of protein aggregate clearance [53, 54]. Hyaluronic acid is

the major component of the perinodal extracellular matrix and accumulate in the brain

with aging [55–59]. Furthermore, it has been shown that hyaluronidase treatment

increases the volume of the extracellular space in mouse hippocampal slices [60].

Therefore, we hypothesized that reducing hyaluronic acid would increase the

extracellular space and improve access to HAPLN2 aggregates, thereby enhancing the

clearance of HAPLN2 aggregates.

To test this idea, we injected hyaluronidase into the mouse cerebellum (Fig 6A).

Immunohistochemical analysis revealed that the expression and size of HAPLN2

dramatically decreased 16 hours after hyaluronidase injection (Figs 6B and 6C).

Immunoblot analysis of brain lysates fractionated as in Figure 1A showed that HAPLN2

in the sarkosyl-insoluble P2 fractions in the hyaluronidase-injected brain (ipsilateral)

significantly decreased compared to that in the PBS-injected ipsilateral brain, whereas

HAPLN2 levels in the sarkosyl-soluble S2 fractions were not significantly affected (Figs

6D and 6E). These results suggest that hyaluronidase specifically promoted the clearance

of aggregated HAPLN2.

Figure 6. Hyaluronidase Promotes Clearance of HAPLN2 Aggregates in the Mouse Cerebellum. (A) Schematic illustration of hyaluronidase injection into the mouse cerebellum. (B) Fluorescence immunohistochemistry images of the cerebellum of 12-months-old mice showing HAPLN2, HABP, and DAPI signals. n = 4. (C) Quantitation of the area of HAPLN2-positive puncta in (B). (D) Immunoblot analysis of the cerebellum corresponding to the samples in (B). (E) Densitometric quantification of (D). HAPLN2 Protein levels in the S2 fraction were normalized to α-tubulin in the S2 fractions. HAPLN2 Protein levels in the P2 fraction were normalized to α-tubulin in the S1 fractions. Error bars represent mean ± S.D. *P*-values by one-way ANOVA followed by Tukey's *post hoc* tests.

Microglia regulated the accumulation of HAPLN2 aggregates

Microglia are the resident macrophages in the brain, acting as scavengers that remove debris and protein aggregates from the extracellular space through phagocytosis [61]. To examine whether microglia are involved in the clearance of HAPLN2 aggregates, we treated 4-week-old mice with pexidartinib (PLX3397), a colony-stimulating factor 1 receptor (CSF1-R) inhibitor known to deplete microglia [62]. Immunostaining with ionized calcium-binding adapter molecule 1 (Iba1), a microglia/macrophage marker, confirmed that treatment with PLX3397 for two months successfully depleted microglia (Fig 7A). Immunohistochemical analysis showed that microglial depletion resulted in the accumulation and enlargement of HAPLN2 puncta in the brains of PLX3397-treated mice (Figs 7B and 7C). Furthermore, PLX3397 treatment resulted in HAPLN2 enrichment within the sarkosyl-insoluble fraction of the cerebellum (Figs 7D and 7E). These results suggest that microglia play a critical role in the clearance of HAPLN2 aggregates.

Figure 7. Accumulation of HAPLN2 Aggregates Is Promoted by Microglial Depletion in the Cerebellum. (A) Immunohistochemistry images of the cerebellum from mice fed AIN-76A control chow or AIN-76A containing 290 mg/kg PLX3397 from one

month to three months of age, stained with anti-ionized calcium-binding adaptor molecule 1 (Iba1) antibody as a microglia marker. n = 3. (B) Immunohistochemistry images of the cerebellum from the mice in (A), stained with anti-HAPLN2 antibody. n = 3. (C) Quantification of the area of each HAPLN2-positive punctum in (B). (D) Immunoblot analysis of the cerebellum from the mice in (B), fractionated as described in Figure 1A. n = 3. (E) Densitometric quantitation of (D). Protein expression levels were normalized to α-tubulin in the S1 fractions. Error bars represent mean ± S.D. *P*-values were calculated using Student's *t*-test.

Oligomers and aggregates of HAPLN2 induced microglial activation *in vitro* and *in vivo*

Microglia that phagocytose extracellular protein aggregates exhibit a pro-inflammatory M1 phenotype and secrete cytokines such as interleukin (IL)-6 and tumor necrosis factor (TNF)-α [61]. Recent studies have shown that soluble oligomers of Aβ, tau, and α-synuclein in the extracellular spaces induce inflammatory responses prior to the accumulation of protein aggregates in the brains of patients with neurodegenerative diseases [63].

To investigate whether HAPLN2 triggers microglial inflammation, we purified recombinant proteins of full-length HAPLN2 and each domain from a lipopolysaccharide (LPS)-free *Escherichia coli* strain (Fig. 8A). The N-terminal Ig-like domain is thought to mediate protein interactions, while the Link1 and Link2 domains are believed to bind to hyaluronic acid. First, we performed a filter trap assay to determine whether solubilized full-length HAPLN2 and the N-terminal Ig-like domain exist as monomers or soluble oligomers. Both soluble full-length HAPLN2 and the N-terminal Ig-like domain were trapped on the nitrocellulose membrane after ultracentrifugation, indicating that they form soluble oligomers (Fig. S4A). We also prepared aggregated full-length HAPLN2 and the N-terminal Ig-like domain by incubating soluble each soluble protein at pH 6.0, which was confirmed by filter trap assay (Fig. S4A).

To evaluate the potential to induce microglial inflammatory responses, we added full-

length HAPLN2 oligomers and aggregates as well as the positive control LPS to the MG6

microglial cell line. Both soluble oligomers and aggregates of HAPLN2 increased the

mRNA expression of IL-6 and TNF-α (Fig. 8B). These results suggest that HAPLN2

induces pro-inflammatory microglial activation in both the soluble oligomeric and

aggregated states.

To determine whether specific domains of HAPLN2 induce pro-inflammatory microglial

activation, we added soluble full-length HAPLN2 and its individual domains to MG6

cells. Soluble OxStayGold fluorescent protein, which was not trapped by the

nitrocellulose membrane, was used as a negative control (Fig S4A). Soluble full-length

HAPLN2 significantly induced IL-6 and TNF-α mRNA expression, but none of the

individual domains of HAPLN2 upregulated these cytokines (Fig 8C). These results

suggest that the pro-inflammatory effect is associated with full-length HAPLN2 and not

with specific domains.

Finally, we examined whether HAPLN2 oligomers induce microglial activation in mouse

brains by infusing full-length HAPLN2 and each domain into the hippocampus. Injection

of full-length HAPLN2 increased pro-inflammatory microglia, marked by Iba1, in the

hippocampus (Figs 8E, F, and S4B). In contrast, oxStayGold and each domain of

HAPLN2 did not significantly induce microglial activation, consistent with Figure 8C.

These results suggest that oligomeric full-length HAPLN2 promotes microglial activation

in vivo.

Figure 8. HAPLN2 Induced Microglial Activation *In Vitro* and *In Vivo*. (A) Diagram

of the domains within HAPLN2. (B) Reverse transcription polymerase chain reaction

(RT-qPCR) analysis of pro-inflammatory cytokine mRNA (*Il6* and *Tnf*, normalized to

Actb) in MG6 cells stimulated with 1 μM LPS, and 200 nM recombinant full-length

HAPLN2 in soluble and aggregated states for four hours. PBS served as a control.

Recombinant HAPLN2 was aggregated by incubation in a buffer containing 10 mM

sodium phosphate and 300 mM NaCl (pH 6.0) for 24 hours at 37ºC. (C) RT-qPCR

analysis of pro-inflammatory cytokine mRNA (*Il6* and *Tnf*, normalized to *Actb*) in MG6

cells stimulated with 1 μM LPS, and 200 nM of recombinant proteins (oxStayGold, full-

length HAPLN2, HAPLN2 Ig-like domain, HAPLN2 link 1 domain, and HAPLN2 link 2

domain) for four hours. PBS served as a control. (D) Fluorescence immunohistochemistry

images of the hippocampus of three-months-old mice injected with PBS, 20 μg LPS, and

15 pmol of recombinant proteins (oxStayGold, full-length HAPLN2, HAPLN2 Ig-like

domain, HAPLN2 link 1 domain, and HAPLN2 link 2 domain). The brains were stained

with anti-Iba1 antibody to detect activated microglia. Scale bar, 500 μm. n = 4. (E)

Quantitation of the number of activated microglia labeled by Iba1 from (E). Error bars

represent mean ± S.D. *P*-values were calculated using one-way ANOVA followed by

Tukey's *post hoc* tests.

HAPLN2 accumulated with age in the human cerebellum

Finally, we examined whether HAPLN2 accumulates with age in the human brain. We

stained HAPLN2 in the cerebellum of human brains in their sixties or eighties without

dementia (S3 Table). We observed the increased HAPLN2 puncta in the human brain in

their eighties compared to those in their sixties (Fig 9A). These HAPLN2 puncta did not

colocalize with Caspr, suggesting that these are age-dependent aggregates. In addition,

the area of HAPLN2 puncta increased with age (Figs 9B and 9C). These results suggest

that HAPLN2 accumulation is a common hallmark of aging in both mice and humans.

Figure 9. HAPLN2 Formed Large Puncta with Age in the Human Cerebellum. (A)

Fluorescent immunohistochemistry images for HAPLN2, Caspr, and DAPI in the

cerebellum from human post-mortem brains without dementia. Scale bar, 50 μm. (B)

Quantification of the area of each HAPLN2 puncta in (A). (C) Quantification of the

average area of HAPLN2 puncta across all samples in (A). Error bars represent mean ±

S.D. *P*-values were calculated using Student's *t*-test.

Discussion

Research on age-related protein aggregation in mammals has progressed at a slower pace than that in *C. elegans* due to the difficulty in obtaining a sufficient number of aging individuals. Recently, studies have aimed to identify proteins that become Triton X-100 insoluble in cerebral cortex with age [64]. However, as demonstrated by the evidence that membraneless organelle constituent proteins and functional large protein complexes, such as the cytoskeleton, also become Triton X-100 insoluble, these proteins are not synonymous with aggregated proteins[64]. Therefore, we used sarkosyl to search for proteins that aggregate with aging under conditions that fractionate Aβ and tau protein aggregates.

We identified nine proteins that become sarkosyl-insoluble and accumulate in mouse brains with aging. Of these, seven were extracellular matrix components, immunoglobulin components, and cell adhesion factors, all of which are localized extracellularly or are in contact with the extracellular space. We focused on HAPLN2, a component of the perinodal extracellular matrix, as a protein that aggregates in an age-dependent manner. HAPLN2 aggregates colocalized with hyaluronic acid in the cerebellum white matter (Fig 3D). This observation, coupled with the *in vitro* finding that hyaluronic acid promotes aggregation of HAPLN2 (Fig 5A), suggests a key role for hyaluronic acid in this process. This is consistent with reports that the concentration of hyaluronic acid increases especially in prefrontal cortex and cerebellum with age *in vivo* [55, 59, 65]. Furthermore, it is reported that microglia are fewer in number and more sparsely distributed in the cerebellum compared to other brain regions [66–68]. Therefore, clearance of protein aggregates by microglial phagocytosis may be less efficient in the cerebellum compared to the cerebrum. Moreover, the cerebellum has a significantly higher density of neurons and axons compared to the cerebrum [69]. This implies a greater number of nodes of Ranvier and their associated perinodal extracellular matrix, which could contribute to the high expression of HAPLN2 observed in this region. This may explain the significant HAPLN2 aggregation observed in the cerebellum.

Experiments using recombinant HAPLN2 revealed that the formed amorphous aggregates

partially contained cross-β-sheet structures, characteristic of proteinaceous aggregates

(Fig 5C). This suggests that HAPLN2 forms aggregates through protein-protein

interactions, rather than simply forming complexes with hyaluronic acid. Interestingly,

HAPLN2 extracted from the sarkosyl-insoluble P2 fraction of aged mouse brains

remained insoluble even after enzymatic degradation of hyaluronic acid (Figs 2D and

2E). This suggests that while initial aggregation may be driven by interactions with

hyaluronic acid, HAPLN2 aggregates ultimately acquire protein-protein interactions that

maintain sarkosyl insolubility. Adding another layer to the role of hyaluronic acid, we

also showed that injecting hyaluronidase into the mouse cerebellum led to a decrease in

sarkosyl-insoluble HAPLN2. Hyaluronidase treatment may increase the accessibility of

clearance systems, such as microglia or the glymphatic system, by increasing the volume

of extracellular spaces [60]. This is supported by studies showing that increased

extracellular volume enhances Aβ clearance [70]. Hence, manipulating hyaluronic acid

levels may be a viable approach to modulate HAPLN2 aggregation by influencing

extracellular volume and the accessibility of clearance mechanisms. However, further

research is needed to determine whether microglia or the glymphatic system are

responsible for the enhanced clearance of HAPLN2 aggregates following hyaluronidase

injection. Additionally, recombinant HAPLN2 protein efficiently formed aggregates

under low pH conditions (Fig 5). Given that brain pH decreases with age, from 6.30 to

6.00 in humans and from 6.85 to 6.70 in mice [71], this change in the brain environment

may also contribute to HAPLN2 aggregation. Taken together, our data suggest that

changes in brain pH and hyaluronic acid levels may contribute to the aggregation of

HAPLN2 *in vivo*, highlighting the complex interplay between protein aggregation and the

aging brain environment, particularly for HAPLN2.

Microglia play a critical role in clearing extracellular waste, such as protein aggregates,

through phagocytosis [61]. Recent studies have also shown that microglia phagocytose

the extracellular matrix, including the perineuronal net [72, 73]. However, aging is

associated with a decline in microglial phagocytic capacity due to increased expression of

CD22, a negative regulator of phagocytosis [11, 74]. Our study demonstrated that

microglial depletion promotes the accumulation of HAPLN2 aggregates (Fig 7),

suggesting that age-related impairment of microglial phagocytosis disrupts the clearance

of HAPLN2 and the perinodal extracellular matrix. The accumulation of protein

oligomers and aggregates in the extracellular space activates microglia. This leads to the

secretion of pro-inflammatory cytokines, such as IL-6 and TNF-α, which contribute to

chronic inflammation [75]. Our study showed that excessive HAPLN2 oligomers and

aggregates activate microglia both *in vitro* and *in vivo* (Fig 8). These findings indicate

that while microglia are involved in clearing HAPLN2 aggregates, their phagocytic

capacity can become overwhelmed by abnormal accumulation of HAPLN2 aggregates.

This imbalance may trigger microglial pro-inflammatory activation, further contributing

to chronic inflammation, a key feature of brain aging. Additionally, aging is associated

with a decline in cerebrospinal fluid perfusion, impairing the clearance of extracellular

waste by the glymphatic system [12]. This decline may exacerbate the accumulation of

HAPLN2 aggregates and further promote neuroinflammation in aged brains. However,

further research is needed to elucidate the mechanism linking the glymphatic system with

accumulation of HAPLN2 aggregates and neuroinflammation during aging.

In summary, we have identified HAPLN2, a component of the perinodal extracellular

matrix, as an age-dependent protein aggregate. HAPLN2 aggregates mislocalize and

accumulate in the extracellular space in aged mouse brains, where hyaluronic acid and

microglia regulate their accumulation and clearance. We also observed HAPLN2

aggregates in non-demented human brains. Our study suggests HAPLN2 aggregation as

a potential factor in chronic inflammation in the brain during physiological aging and

sheds new light on our understanding of age-related decline in brain function.

Materials and Methods

Post-mortem brain samples

600 The paraffin-embedded postmortem cerebellum derived from neurologically normal

601 patients in their sixties (n = 4) or eighties (n = 5) were obtained from Japan Brain Bank

Net (JBBN). Information on human donors is provided in S3 Table. All participants, before passing away or their families, provided written informed consent. Experiments using human samples were performed with institutional approval and guidelines from the institutional ethical committee of the Graduate School of Pharmaceutical Sciences, University of Tokyo (No. 4-8).

Animals

C57BL/6N mice were purchased from Clea Japan. All mice were maintained under specific pathogen-free conditions. All animal experiments were performed after obtaining approval from the Institutional Animal Care Committee of the Graduate School of Pharmaceutical Sciences, the University of Tokyo (approval number P26-7). Male mice were used in all experiments. Mice were euthanized by cervical dislocation to preserve the metabolic environment of the brains and to prevent artifacts that could alter the biochemical profiles of the proteome. Mouse brains were bisected down the midline to yield two hemispheres. The brain hemisphere of each animal was quickly frozen on dry ice and stored at -150ºC until use.

Cell culture

The MG6 cell line is a c-Myc-immortalized cell line of mouse microglia. MG6 cells were maintained at 5% CO_2 at 37ºC in Dulbecco's modified Eagle's medium supplemented with 10% (v/v) fetal bovine serum (Thermo Fisher Scientific, 10270-106), 10 μg/mL insulin (Nacalai Tesque, 12878-44), and 0.1 mM 2-mercaptoethanol [76, 77].

Label-free quantitative proteomic analysis

The fractionation of the brain tissues was prepared as previously described [30]. Tissues

were crushed using Multi-Beads-Shocker (Yasui Kikai, MB601U(S)). Then the tissues

were homogenized with a Potter-type homogenizer (AS ONE CORPORATION) at 800

rpm in 1.4 mL of sarkosyl buffer [PBS, 1% sarkosyl (Nacalai Tesque, 20135-14) and 1

mM phenylmethylsulfonyl fluoride (PMSF) (Wako, 020-15372)]. The homogenates were

centrifuged at 15,000 × g for 5 minutes twice, then for 15 minutes once at 4ºC to obtain

supernatant (S1) and pellet (P1) fractions. The supernatants were centrifuged at 100,000

× g for one hour at 4ºC to obtain sarkosyl-soluble (S2) and sarkosyl-insoluble (P2)

fractions. The P2 fraction was resuspended in 140 µL of phase transfer surfactant PTS

[12 mM sodium deoxycholate (SDC), 12 mM sarkosyl, and 200 mM ammonium

bicarbonate] buffer. The resuspended P2 solutions were sonicated with Bioruptor® Plus

(BMBio, B01020001) for 10 cycles, (30"/30" cycles), and incubated for 30 minutes at

60ºC with 5 mM dithiothreitol (DTT) for solubilization and reduction of the proteins. The

samples were then alkylated with 200 mM methyl methanethiosulfonate for 30 minutes at

room temperature in the dark. The samples were purified with methanol-chloroform

extraction and solubilized with 50 µL of Rapigest buffer [0.1% Rapigest SF (Waters,

186002123) and 50 mM tetraethylammonium bicarbonate (Thermo Fisher Scientific,

90114)]. The alkylated samples were sonicated and were digested overnight with 1 µg of

Trypsin Gold (Promega, V5280) at 37ºC. Peptides were purified from the supernatant by

GL-tip SDB (GL Sciences, 7820-11200) and eluted with QE-B solution [70%

Acetonitrile and 0.1% formic acid] (Kanto Chemical Co. Inc., 01922-64). Peptides were

concentrated with Concentrator Plus™ for 45ºC and resuspended with 30 µL of QE-A

solution [0.1% formic acid] (Kanto Chemical Co. Inc., 16245-12). The concentration of

peptides was measured with Pierce Quantitative Colorimetric Peptide Assay (Thermo

Fisher Scientific, 23275). 500 ng of peptides from each fraction were subjected to

proteomic analysis using an EASY-nLC 1200 system connected online to Q-Exactive

(Thermo Fisher Scientific). Samples were loaded onto Acclaim PepMap 100 C18 column

(75 µm × 2 cm, 3 µm particle size and 100 Å pore size; 164946, Thermo Fisher Scientific)

and separated using capillary column, (75 µm × 12.5 cm, 3 µm particle size, NTCC-

360/75-3-125, Nikkyo Technos) in an EASY-nLC 1200 system (Thermo Fisher

Scientific). Data were analyzed using Proteome Discoverer (v.2.4). The mass tolerances

for the precursor and fragment ions were 10 ppm and 0.02 Da, respectively, and peptide

identification was filtered at a false discovery rate <0.01.

Tandem-mass tag (TMT) proteomics

The TMT proteomics was conducted as described previously [34]. Tissues were crushed

and homogenized in 400 µL of PTS buffer. The homogenates were centrifuged at 15,000

× g for 15 minutes at 4ºC to yield the supernatant and pellet fractions. 30 µg of the

supernatant was reduced with 5 mM DTT for 15 minutes at room temperature. The

samples were then alkylated and digested by Trypsin Gold. The samples were centrifuged

at 15,000 × g and the supernatants were labeled with TMT sixplex™ Mass Tag Labeling

Kits and Reagents (Thermo Fisher Scientific, 90061). The labeled samples were added

with 0.1% trifluoroacetate (Wako, 192-09912), incubated for 45 minutes at 37ºC, and

centrifuged at 15,000 × g for 15 minutes at 4ºC. The supernatant was fractionated with

Pierce High pH Reversed-Phase Peptide Fractionation Kit (Thermo Fisher Scientific,

84868). The eluates were concentrated at 45ºC and resuspended with 40 µL of QE-A. 500

ng of peptides were centrifuged at 15,000 × g for 15 minutes at 4ºC and subjected to

proteomic analysis with Q-Exactive.

GO-term analysis

676 In the GO analysis of proteomic data, we used the Database for Annotation, Visualization

677 and Integrated Discovery (DAVID) [78].

Antibodies

Primary antibodies for immunoblotting (IB) and immunohistochemistry (IHC) are listed

in S4 Table. All secondary antibodies used for immunoblotting were purchased from

Jackson ImmunoResearch Laboratories: Peroxidase AffiniPure™ Rabbit Anti-Mouse IgG

+ IgM (H+L) (315-035-048), Peroxidase AffiniPure™ Goat Anti-Rabbit IgG (H+L)

(111-035-144), and Peroxidase AffiniPure™ Donkey Anti-Goat IgG (H+L) (705-035-

147). All the secondary antibodies used for immunohistochemistry were purchased from

ThermoFisher Scientific: Alexa Fluor 488 Goat Anti-Mouse IgG (H+L) (A-28175),

Alexa Fluor 488 Goat Anti-Rabbit IgG (H+L) (A-11008), Alexa Fluor 555 Mouse Anti-

Mouse IgG (H+L) (A-28180), Alexa Fluor 647 Streptavidin (S21374).

690   Immunoblot analysis

Sample preparation, sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-

PAGE), and immunoblot analysis were performed as described previously [79]. Each

fraction of the brain lysates was mixed with SDS lysis buffer [50 mM Tris-HCl (pH 8.0),

150 mM NaCl, 0.1% SDS, 1% Triton X-100, 0.5% SDC, and 1 mM PMSF] and

centrifuged at $20,000 \times g$ for 20 minutes at 4ºC. The supernatants were separated by

SDS-PAGE (12.5% acrylamide, 7×8 cm, 1 mm thick). Proteins were electrotransferred

to Immobilon®-P polyvinylidene difluoride (PVDF) (Merck Millipore, IPVH00010), at

30 V for one hour at room temperature. For immunoblot analysis, the membranes were

incubated with Blocking One (Nacalai Tesque, 03953-95) for one hour at room

temperature. The membranes were incubated with primary antibodies in TBS-T buffer

[20 mM Tris-HCl (pH8.0), 150 mM NaCl, and 0.1% Tween-20] overnight at 4ºC, washed

with TBS-T buffer and incubated with horseradish-peroxidase conjugated secondary

antibody (1:20,000) for 30 minutes at room temperature. The membranes were immersed

with ImmunoLightning Plus (PerkinElmer, NEL105001EA) or ImmunoSTAR LD

(Wako, 290-69904), and the images were acquired with FUSION FX7.EDGE (VILBER).

For band quantification, the intensity of each band was measured using Fiji software

(ImageJ2, version 2.14.0/1.54h) [80].

709   Filter trap assay

The S1 fractions of the young and aged mouse brains were serially diluted two-fold with sarkosyl buffer. The 50 µL of each sample were filtered under vacuum through a pre-wet 0.45-µm pore size nitrocellulose (Bio-Rad, 1620167) and Immobilon®-P, using a Bio-dot® Microfiltration Apparatus (Bio-Rad, 1706545). Each well was washed three times with the sarkosyl buffer. The membranes were then processed following the same immunoblotting procedure from the blocking step onward.

Immunohistochemical analysis

The paraffin sections (10 µm) were boiled in 10 mM sodium citrate buffer (pH 6.0) for one minute in a microwave oven for heat-induced antigen retrieval. The sections were incubated for one hour in Blocking One Histo (Nacalai Tesque, 06349-64), then overnight at 4ºC with primary antibodies diluted in Blocking One Histo. After rinsing, sections were incubated with secondary antibodies and 1 µg/mL 4',6-diamido-2-phenylindole dihydrochloride (DAPI) (Nacalai Tesque, 11034-56) diluted in Blocking One Histo for one hour at room temperature. The sections were washed by TBS-T three times and by PBS three times, then immersed in TrueBlack® Lipofuscin Autofluorescence Quencher (Biotium Inc., 23007) for 20 minutes to reduce autofluorescence from lipofuscin. The slices were mounted with Prolong™ Glass Antifade Mountant (Invitrogen™, P36980).

For the staining of Iba1, TSA plus amplification kit (Akoya Biosciences, NEL741001KT) was used in combination with subsequent conventional immunofluorescence staining for other antigens. To block endogenous peroxidase activity, brain sections were treated with 3% H_2O_2 in methanol for 15 minutes at room temperature. After washing, sections were treated with Blocking one Histo for one hour at room temperature, and then incubated with 0.1% anti-Iba1 antibody (Wako, 013-27691) in 5% Blocking one Histo in TBS-T overnight at 4ºC. The sections were incubated with 0.01% Peroxidase AffiniPure™ Goat Anti-Rabbit IgG (H+L) (Jackson ImmunoResearch Laboratories, 111-035-144) diluted in TNB buffer at 4ºC overnight. After washing with TNT buffer (0.1% Triton X-100, 100 mM Tris-HCl (pH 7.5), 150 mM NaCl) three times, the sections were reacted with

Fluorescein Plus amplification reagent for 10 minutes at room temperature, washed with

TNT.

Images were captured by BZ-X800 (Keyence) using 100X objectives or Leica DMi8

Thunder (Leica Microsystems) using 10X or 63X objectives. The images were processed

and analyzed with Fiji software. Colocalization analyses were performed with the JaCoP

plugin in Fiji software[81].

Protein purification

Recombinant HAPLN2 was purified using the method to purify the versican core protein

G1 domain as previously described [82]. The HAPLN2 coding sequence, nucleotides 82–

1026 was amplified from mouse brain cDNA by PCR and cloned into pGEX6p-1

(Addgene, 27-4597-01). The glutathione-S-transferase tag was removed by PCR. The

construct was transformed into ClearColi® BL21(DE3) Electrocompetent Cells (LGC

Biosearch Technologies, 60810-1). The cells were grown in LB Miller broth containing

100 µg/mL ampicillin at 37ºC with shaking at 150 rpm. The *E. coli* cultures were induced

with 0.8 mM isopropyl β-D-thiogalactopyranoside (Nacalai Tesque, 06289-67), grown

for another 20 hours at 20ºC, and harvested by centrifugation at $10,000 \times g$ for 20

minutes at 4ºC (Beckman Coulter, Avanti J-25). Inclusion bodies were purified and

solubilized in urea and the protein was refolded, essentially as described previously [82].

Briefly, inclusion bodies were isolated with an *E. coli* lysis buffer [1% (w/v) Nonidet P-

40, 1% (w/v) Triton X-100, 1 mM PMSF, and 1 mM DTT in PBS] solubilized in a

denaturation buffer [8 M urea in 50 mM Tris-HCl (pH 7.5)] and clarified by

centrifugation. The supernatant was purified by incubating with Ni-NTA agarose

(QIAGEN, 30210) overnight and eluted with elution buffer [6 M urea, 500 mM

imidazole, and 10 mM HCl] and then diluted 66-fold into refolding buffer [1 mM

ethylenediaminetetraacetic acid, 500 mM L-arginine, 1 mM cysteine, 2 mM cystine, and

20 mM ethanolamine (pH 11.0)] by dropwise addition at 4ºC with gentle stirring,

followed by incubation overnight without stirring. After refolding, the protein was

concentrated to 2 mL with Amicon Ultra-15 Centrifugal Filter 10 kDa MWCO

(Millipore, 36100101), and dialyzed overnight into a storage buffer [10 mM sodium
phosphate (pH 8.0)].

Protein aggregation assay

Recombinant HAPLN2 proteins (500 nM) were subjected to a series of pH (6.0–8.0),
NaCl concentration (0–300 mM), and hyaluronic acid (0–1 mg/mL) in a total volume of
100 µL. The samples were deposited on µ-Slide 18-Well Glass Bottom (Ibidi, 81817) and
incubated for 24 hours at 37ºC before being imaged on Eclipse Ti2 microscope (Nikon
Corporation) with Micromanager software [83] in which bright field images were taken
using a 20X air magnification objective. Aggregate formation was imaged by collecting a
series of images in the bright field channel. Fiji software was used in all image
processing.

In vitro single-aggregate imaging

Super-resolution imaging was conducted based on point accumulation for imaging
nanoscale topography (PAINT). Borosilicate glass coverslips (VWR International, 22 ×
22 mm, 631-0124) were cleaned with an argon plasma cleaner (Harrick Plasma, PDC-
002) for at least one hour to remove fluorescent residues. Before use, each batch of cover
slides was tested for fluorescent artifacts. A drop of 10 µL of the mixture of HAPLN2
used in the protein aggregation assay was placed over a coverslip together with
Thioflavin T and Nile Red dyes diluted in PBS to achieve final concentrations of 5 µM
for Thioflavin T and 5 nM for Nile Red. Imaging was performed using a CFI
Apochromat TIRF 100x Oil immersion objective (Nikon Corporation), mounted on an
Eclipse Ti2 microscope (Nikon Corporation). Thioflavin T was excited using a 405 nm
laser (Cobolt #0405-06-01-0120-100) with a beam power of 20 mW. Emitted
fluorescence was collected by the same objective and separated from the excitation light
by a dichroic (Semrock #DI03-R405/488/532/635-T1-25X36), subsequently passed

through an emission filter (Semrock #FF01-510/84-25) and focused onto a CMOS
camera (ORCA-Flash4.0 V3 Digital CMOS; Hamamatsu). Nile Red was excited by a
532nm laser (Cobolt #0532-06-91-0100-100) with a beam power of 80 mW. Both lasers
were positioned at a highly inclined and laminated optical sheet (HiLo) illumination
angle to image large amorphous aggregates at the surface of the coverslip (exciting ~5
μm deep in the sample) while minimizing background fluorescence. Emitted light was
separated using a dichroic (DC/ZT532rdc-UF2; Chroma) and passed through an emission
filter (Semrock #FF01-650/150-25) before being focused onto the CMOS camera. The
image pixel size was 65 nm. A z-stack of 5.0 μm (10 planes with 0.5 μm spacing) was
acquired in all channels. The microscope was controlled with Micromanager software,
and the frame interval was 50 ms. Each analyzed image corresponds to an average of 10
images in the bright-field channel, 50 images in the thioflavin T, and 1,000 images in the
Nile Red channel. The Nile Red images were reconstructed with the ThunderSTORM
plug-in in Fiji software [84]. The morphological analysis of Nile Red-positive structures
from reconstructed images was conducted by LocAlization Microscopy Analyzer
(LAMA) [52] and MATLAB (Mathworks, ver. R2023b).

Negative-stained transmission electron microscopy

1.5 μL of recombinant samples were deposited onto glow-discharged formvar-coated
300-mesh copper grids (Stork Veco, Eerbeek, The Netherlands) for 20 minutes at room
temperature. The grids were blotted with filter paper (Toyo Roshi Kaisha, Ltd., Tokyo,
Japan), washed twice with PBS, washed twice with distilled water, stained with 2% (w/v)
uranyl acetate for 30 seconds, blotted and air-dried. The prepared specimens were finally
examined with an electron microscope (HT7700; Hitachi, Tokyo, Japan) at 100 kV.

Stereotaxic injection

Mice were anesthetized by intraperitoneal injection with 5.0 mL/kg of a non-narcotic

anesthetic combination (Me/Mi/Bu) [30 µg/mL medetomidine (Zenoaq), 0.4 mg/mL

midazolam (Astellas), and 0.5 mg/mL butorphanol (Meiji Seika Pharma)] and fixed in a

stereotaxic surgery frame.

For intracerebellar injection, 20 mg/mL hyaluronidase from ovine testes (FUJIFILM

Wako Pure Chemical Corporation, 37326-33-3) in a total volume of 1.0 µL was injected

at the following coordinates relative to the bregma: anterior-posterior (A/P) -5.5 mm

medial-lateral (M/L) -1.2 mm, and dorsal-ventral (D/V) -2.5 mm. Mice were sacrificed

one day post-surgery.

For intrahippocampal injection, either 15 pmol of protein or 5 mg/mL LPS solution, each

in a total volume of 2.0 µL, was injected at the following coordinates relative to the

bregma: A/P -1.7 mm, M/L -1.6 mm, and D/V -1.9 mm. Mice were sacrificed one week

post-surgery.

Depletion of microglia by PLX3397 treatment

PLX3397 [85] was synthesized as described previously [86] and formulated at 290 mg/kg

in AIN-76A standard chow by Research Diets Inc. The chows were given to 4-week-old

wild-type mice (C57/BL6J) for two months. Depletion of microglia was confirmed by

immunohistochemistry for a microglial marker Iba1.

RNA isolation and quantitative real-time PCR

Total RNA was extracted from MG6 cells grown in 12-well tissue culture plates (7×10^4

cells/well) by High Pure RNA Isolation Kit (Roche, 11828665001). The isolated total

RNA was then reverse-transcribed with ReverTra Ace (TOYOBO, FSQ-201). cDNAs

were amplified with THUNDERBIRD Probe qPCR mix (Toyobo, QPS-101), and

846 Universal Probe Library probe (Roche) the following specific primer sets. mRNA levels
847 were measured with LightCycler® 480 (Roche).

Gene	Forward (5'–3')	Reverse (5'–3')
Il6	TGTGAAATCAGGATGCTCTGG	CACTTTATTGGGCTCTATACAATATGC
Tnf	TCTTCTCATTCCTGCTTGTGG	GGTCTGGGCCATAGAACTGA
Actb	ACCAGAGGCATACAGGGACA	CTAAGGCCAACCGTGAAAAG

Statistics and reproducibility

850 All data were derived from at least three independent experiments and are presented as
851 mean ± S.D. unless otherwise stated in the figure legends. Statistical analyses were
852 performed with GraphPad Prism 9.0 software. An unpaired two-tailed Student's *t*-test or
853 one-way ANOVA followed by Tukey's *post hoc* test was performed as described in the
854 figure legends. All replication attempts were successful with similar results.

Acknowledgments

857 We thank members of the Laboratory of Morphology and Image Analysis, Biomedical
858 Research Core Facilities, and Juntendo University Graduate School of Medicine for
859 technical assistance with microscopy.

Author Contributions

862 A.W. did most of the experimental work *in vivo* and *in vitro*, designed the experiments,
863 analyzed data, and wrote the paper. S.H. conceived the project, designed the experiments,
864 performed fractionation experiments, MS analysis, stereotaxic injection, and
865 immunohistochemical analysis of HAPLN2, analyzed data, and wrote the paper. K.I.

performed fractionation experiments, stereotaxic injection, and immunohistochemical analysis of HAPLN2, and analyzed data. S. Marchese, P. E., and V.M. provided technical support for HiLo imaging and analyzed data. H.K. performed EM analysis, analyzed data, and wrote the paper. S.T. provided technical support for stereotaxic injection, discussed the results, and analyzed data. M.K. provided technical support for EM analysis, designed the experiments, and analyzed data. T.S. and M.T. provided post-mortem human brains. J.A.V. provided technical support for HiLo imaging, analyzed data, discussed the results, and wrote the paper. T.T. provided technical support for stereotaxic injection and analyzed data. S. Murata conceived the project, designed the experiments, analyzed data, and wrote the paper. All authors reviewed the final version and agreed to its publication.

References

[1]   Chung CG, Lee H, Lee SB. Mechanisms of protein toxicity in neurodegenerative diseases. *Cellular and Molecular Life Sciences* 2018; 75: 3159–3180.

[2]   Chou A-H, Yeh T-H, Ouyang P, et al. Polyglutamine-expanded ataxin-3 causes cerebellar dysfunction of SCA3 transgenic mice by inducing transcriptional dysregulation. *Neurobiol Dis* 2008; 31: 89–101.

[3]   Lustbader JW, Cirilli M, Lin C, et al. ABAD Directly Links Aβ to Mitochondrial Toxicity in Alzheimer's Disease. *Science (1979)* 2004; 304: 448–452.

[4]   Kastelic M, Kalyuzhnyi Y V., Hribar-Lee B, et al. Protein aggregation in salt solutions. *Proceedings of the National Academy of Sciences* 2015; 112: 6766–6770.

[5]   Cui Q, Liu Z, Bai G. Friend or foe: The role of stress granule in neurodegenerative disease. *Neuron* 2024; 112: 2464–2485.

[6]   Bachiller S, Jiménez-Ferrer I, Paulus A, et al. Microglia in Neurological Diseases: A Road Map to Brain-Disease Dependent-Inflammatory Response. *Front Cell Neurosci* 2018; 12: 488.

[7]   Kelmer Sacramento E, Kirkpatrick JM, Mazzetto M, et al. Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol* 2020; 16: e9596.

[8]   Shibata M, Lu T, Furuya T, et al. Regulation of intracellular accumulation of mutant huntingtin by beclin 1. *Journal of Biological Chemistry* 2006; 281: 14474–14485.

[9]   Lipinski MM, Zheng B, Lu T, et al. Genome-wide analysis reveals mechanisms modulating autophagy in normal brain aging and in Alzheimer's disease. *Proceedings of the National Academy of Sciences* 2010; 107: 14164–14169.

[10]   Pan J, Ma N, Yu B, et al. Transcriptomic profiling of microglia and astrocytes throughout aging. *J Neuroinflammation* 2020; 17: 97.

[11] Pluvinage J V., Haney MS, Smith BAH, et al. CD22 blockade restores homeostatic microglial
phagocytosis in ageing brains. *Nature* 2019; 568: 187–192.

[12] Da Mesquita S, Herz J, Wall M, et al. Aging-associated deficit in CCR7 is linked to worsened
glymphatic function, cognition, neuroinflammation, and β-amyloid pathology. *Sci Adv* 2021; 7:
4601–4622.

[13] Vilchez D, Saez I, Dillin A. The role of protein clearance mechanisms in organismal ageing and
age-related diseases. *Nat Commun* 2014; 5: 5659.

[14] Safaiyan S, Kannaiyan N, Snaidero N, et al. Age-related myelin degradation burdens the clearance
function of microglia during aging. *Nat Neurosci* 2016; 19: 995–998.

[15] Crystal H, Dickson D, Fuld P, et al. Clinico-pathologic studies in dementia: nondemented subjects
with pathologically confirmed Alzheimer's disease. *Neurology* 1988; 38: 1682–1687.

[16] Iseki E, Tsunoda S, Suzuki K, et al. Regional quantitative analysis of NFT in brains of
non-demented elderly persons: Comparisons with findings in brains of late-onset Alzheimer's
disease and limbic NFT dementia. *Neuropathology* 2002; 22: 34–39.

[17] Schweighauser M, Arseni D, Bacioglu M, et al. Age-dependent formation of TMEM106B amyloid
filaments in human brains. *Nature* 2022; 605: 310–314.

[18] Kenyon C, Chang J, Gensch E, et al. A C. elegans mutant that lives twice as long as wild type.
Nature 1993; 366: 461–464.

[19] Lin K, Dorman JB, Rodan A, et al. daf-16 : An HNF-3/forkhead Family Member That Can
Function to Double the Life-Span of Caenorhabditis elegans. *Science (1979)* 1997; 278: 1319–
1322.

[20] Hsu A-L, Murphy CT, Kenyon C. Regulation of Aging and Age-Related Disease by DAF-16 and
Heat-Shock Factor. *Science (1979)* 2003; 300: 1142–1145.

[21] Morley JF, Morimoto RI. Regulation of Longevity in Caenorhabditis elegans by Heat Shock Factor
and Molecular Chaperones. *Mol Biol Cell* 2004; 15: 657–664.

[22] David DC, Ollikainen N, Trinidad JC, et al. Widespread Protein Aggregation as an Inherent Part of
Aging in C. elegans. *PLoS Biol* 2010; 8: e1000450.

[23] Walther DM, Kasturi P, Zheng M, et al. Widespread proteome remodeling and aggregation in aging
C. elegans. *Cell* 2015; 161: 919–932.

[24] Fares H, Greenwald I. Genetic Analysis of Endocytosis in Caenorhabditis elegans : Coelomocyte
Uptake Defective Mutants. *Genetics* 2001; 159: 133–145.

[25] Gallotta I, Sandhu A, Peters M, et al. Extracellular proteostasis prevents aggregation during
pathogenic attack. *Nature* 2020; 584: 410–414.

[26] Diner I, Hales CM, Bishof I, et al. Aggregation Properties of the Small Nuclear Ribonucleoprotein
U1-70K in Alzheimer Disease. *Journal of Biological Chemistry* 2014; 289: 35296–35313.

[27] Hales CM, Dammer EB, Deng Q, et al. Changes in the detergent-insoluble brain proteome linked to
amyloid and tau in Alzheimer's Disease progression. *Proteomics* 2016; 16: 3042–3053.

[28] Guo Q, Dammer EB, Zhou M, et al. Targeted Quantification of Detergent-Insoluble RNA-Binding
Proteins in Human Brain Reveals Stage and Disease Specific Co-aggregation in Alzheimer's
Disease. *Front Mol Neurosci* 2021; 14: 623659.

[29]   Woltjer RL, Cimino PJ, Boutté AM, et al. Proteomic determination of widespread detergent
insolubility, including Aβ but not tau, early in the pathogenesis of Alzheimer's disease. *The FASEB*
Journal 2005; 19: 1923–1925.

[30]   Diner I, Nguyen T, Seyfried NT. Enrichment of Detergent-insoluble Protein Aggregates from
Human Postmortem Brain. *J Vis Exp* 2017; 2017: 55835.

[31]   Gozal YM, Duong DM, Gearing M, et al. Proteomics Analysis Reveals Novel Components in the
Detergent-Insoluble Subproteome in Alzheimer's Disease. *J Proteome Res* 2009; 8: 5069–5079.

[32]   Mawuenyega KG, Sigurdson W, Ovod V, et al. Decreased Clearance of CNS β-Amyloid in
Alzheimer's Disease. *Science (1979)* 2010; 330: 1774–1774.

[33]   Hier DB, Azizi S, Thimgan MS, et al. Tau kinetics in Alzheimer's disease. *Front Aging Neurosci*
2022; 14: 1055170.

[34]   Masuda T, Tomita M, Ishihama Y. Phase transfer surfactant-aided trypsin digestion for membrane
proteome analysis. *J Proteome Res* 2008; 7: 731–740.

[35]   Cuanalo-Contreras K, Schulz J, Mukherjee A, et al. Extensive accumulation of misfolded protein
aggregates during natural aging and senescence. *Front Aging Neurosci* 2023; 14: 1090109.

[36]   Chang E, Kuret J. Detection and quantification of tau aggregation using a membrane filter assay.
Anal Biochem 2008; 373: 330–336.

[37]   Xu G, Gonzales V, Borchelt DR. Rapid Detection of Protein Aggregates in the Brains of Alzheimer
Patients and Transgenic Mouse Models of Amyloidosis. *Alzheimer Dis Assoc Disord* 2002; 16:
191–195.

[38]   Sugitani K, Egorova D, Mizumoto S, et al. Hyaluronan degradation and release of a hyaluronan-
aggrecan complex from perineuronal nets in the aged mouse brain. *Biochim Biophys Acta Gen Subj*
2021; 1865: 129804.

[39]   Bekku Y, Vargová L, Goto Y, et al. Bral1: Its Role in Diffusion Barrier Formation and Conduction
Velocity in the CNS. *The Journal of Neuroscience* 2010; 30: 3113–3123.

[40]   Ximerakis M, Lipnick SL, Innes BT, et al. Single-cell transcriptomic profiling of the aging mouse
brain. *Nat Neurosci* 2019; 22: 1696–1708.

[41]   Elbaz B, Darwish A, Vardy M, et al. The bone transcription factor Osterix controls extracellular
matrix- and node of Ranvier-related gene expression in oligodendrocytes. *Neuron* 2024; 112: 247-
263.e6.

[42]   Grundke-Iqbal I, Iqbal K, Quinlan M, et al. Microtubule-associated protein tau. A component of
Alzheimer paired helical filaments. *Journal of Biological Chemistry* 1986; 261: 6084–6089.

[43]   Fawcett JW, Oohashi T, Pizzorusso T. The roles of perineuronal nets and the perinodal
extracellular matrix in neuronal function. *Nat Rev Neurosci* 2019; 20: 451–465.

[44]   Dohm-Hansen S, English JA, Lavelle A, et al. The 'middle-aging' brain. *Trends Neurosci* 2024; 47:
259–272.

[45]   Babinchak WM, Haider R, Dumm BK, et al. The role of liquid–liquid phase separation in
aggregation of the TDP-43 low-complexity domain. *Journal of Biological Chemistry* 2019; 294:
6306–6317.

[46]   Qamar S, Wang G, Randle SJ, et al. FUS Phase Separation Is Modulated by a Molecular Chaperone
980   and Methylation of Arginine Cation-π Interactions. *Cell* 2018; 173: 720-734.e15.

981   [47]   Maharana S, Wang J, Papadopoulos DK, et al. RNA buffers the phase separation behavior of prion-
982   like RNA binding proteins. *Science (1979)* 2018; 360: 918–921.

983   [48]   Patel A, Lee HO, Jawerth L, et al. A Liquid-to-Solid Phase Transition of the ALS Protein FUS
984   Accelerated by Disease Mutation. *Cell* 2015; 162: 1066–1077.

985   [49]   Van Lindt J, Bratek-Skicki A, Nguyen PN, et al. A generic approach to study the kinetics of liquid–
986   liquid phase separation under near-native conditions. *Commun Biol* 2021; 4: 77.

987   [50]   Danial JSH, Klenerman D. Single molecule imaging of protein aggregation in Dementia: Methods,
988   insights and prospects. *Neurobiol Dis* 2021; 153: 105327.

989   [51]   Lim D, Chu KK, Mertz J. Wide-field fluorescence sectioning with hybrid speckle and uniform-
990   illumination microscopy. *Opt Lett* 2008; 33: 1819.

991   [52]   Malkusch S, Heilemann M. Extracting quantitative information from single-molecule super-
992   resolution imaging data with LAMA – LocAlization Microscopy Analyzer. *Sci Rep* 2016; 6: 34486.

993   [53]   Bondareff W, Narotzky R. Age Changes in the Neuronal Microenvironment. *Science (1979)* 1972;
994   176: 1135–1136.

995   [54]   Syková E, Voříšek I, Antonova T, et al. Changes in extracellular space size and geometry in APP23
996   transgenic mice: A model of Alzheimer's disease. *Proceedings of the National Academy of*
997   *Sciences* 2005; 102: 479–484.

998   [55]   Reed MJ, Vernon RB, Damodarasamy M, et al. Microvasculature of the Mouse Cerebral Cortex
999   Exhibits Increased Accumulation and Synthesis of Hyaluronan With Aging. *J Gerontol A Biol Sci*
1000   *Med Sci* 2017; 72: 740–746.

1001   [56]   Cargill R, Kohama SG, Struve J, et al. Astrocytes in aged nonhuman primate brain gray matter
1002   synthesize excess hyaluronan. *Neurobiol Aging* 2012; 33: 830.e13-830.e24.

1003   [57]   Reed MJ, Damodarasamy M, Pathan JL, et al. Increased Hyaluronan and TSG-6 in Association
1004   with Neuropathologic Changes of Alzheimer's Disease. *Journal of Alzheimer's Disease* 2019; 67:
1005   91–102.

1006   [58]   Su W, Foster SC, Xing R, et al. CD44 Transmembrane Receptor and Hyaluronan Regulate Adult
1007   Hippocampal Neural Stem Cell Quiescence and Differentiation. *Journal of Biological Chemistry*
1008   2017; 292: 4434–4445.

1009   [59]   Zakusilo FT, Kerry O'Banion M, Gelbard HA, et al. Matters of size: Roles of hyaluronan in CNS
1010   aging and disease. *Ageing Res Rev* 2021; 72: 101485.

1011   [60]   Godin AG, Varela JA, Gao Z, et al. Single-nanotube tracking reveals the nanoscale organization of
1012   the extracellular space in the live brain. *Nature Nanotechnology 2016 12:3* 2016; 12: 238–243.

1013   [61]   Wang WY, Tan MS, Yu JT, et al. Role of pro-inflammatory cytokines released from microglia in
1014   Alzheimer's disease. *Annals of Translational Medicine* 2015; 3: 136.

1015   [62]   Elmore MRP, Najafi AR, Koike MA, et al. Colony-Stimulating Factor 1 Receptor Signaling Is
1016   Necessary for Microglia Viability, Unmasking a Microglia Progenitor Cell in the Adult Brain.
1017   *Neuron* 2014; 82: 380–397.

[63] Subhramanyam CS, Wang C, Hu Q, et al. Microglia-mediated neuroinflammation in
neurodegenerative diseases. *Semin Cell Dev Biol* 2019; 94: 112–120.

[64] Molzahn C, Kuechler ER, Zemlyankina I, et al. Shift of the insoluble content of the proteome in the
aging mouse brain. *Proc Natl Acad Sci U S A* 2023; 120: e2310057120.

[65] Zolla V, Nizamutdinova IT, Scharf B, et al. Aging-related anatomical and biochemical changes in
lymphatic collectors impair lymph transport, fluid homeostasis, and pathogen clearance. *Aging Cell*
2015; 14: 582–594.

[66] Tan Y-L, Yuan Y, Tian L. Microglial regional heterogeneity and its role in the brain. *Mol*
Psychiatry 2020; 25: 351–367.

[67] Savchenko VL, Nikonenko IR, Skibo GG. Microglial cells of the rat brain in postnatal period
(comparative immunocytochemical analysis). *Neurophysiology* 1997; 29: 145–152.

[68] Verdonk F, Roux P, Flamant P, et al. Phenotypic clustering: a novel method for microglial
morphology analysis. *J Neuroinflammation* 2016; 13: 153.

[69] Keller D, Erö C, Markram H. Cell Densities in the Mouse Brain: A Systematic Review. *Front*
Neuroanat; 12. Epub ahead of print 23 October 2018. DOI: 10.3389/fnana.2018.00083.

[70] Xie L, Kang H, Xu Q, et al. Sleep Drives Metabolite Clearance from the Adult Brain. *Science*
(1979) 2013; 342: 373–377.

[71] Decker Y, Németh E, Schomburg R, et al. Decreased pH in the aging brain and Alzheimer's
disease. *Neurobiol Aging* 2021; 101: 40–49.

[72] Crapser JD, Arreola MA, Tsourmas KI, et al. Microglia as hackers of the matrix: sculpting
synapses and the extracellular space. *Cell Mol Immunol* 2021; 18: 2472–2488.

[73] Tansley S, Gu N, Guzmán AU, et al. Microglia-mediated degradation of perineuronal nets
promotes pain. *Science (1979)*. Epub ahead of print 26 May 2022. DOI:
10.1126/SCIENCE.ABL6773.

[74] Aires V, Coulon-Bainier C, Pavlovic A, et al. CD22 Blockage Restores Age-Related Impairments
of Microglia Surveillance Capacity. *Front Immunol* 2021; 12: 43–57.

[75] López-Otín C, Blasco MA, Partridge L, et al. Hallmarks of aging: An expanding universe. *Cell*
2023; 186: 243–278.

[76] Nakamichi K, Saiki M, Kitani H, et al. Suppressive effect of simvastatin on interferon-β-induced
expression of CC chemokine ligand 5 in microglia. *Neurosci Lett* 2006; 407: 205–210.

[77] Takenouchi T, Ogihara K, Sato M, et al. Inhibitory effects of U73122 and U73343 on Ca2+ influx
and pore formation induced by the activation of P2X7 nucleotide receptors in mouse microglial cell
line. *Biochim Biophys Acta* 2005; 1726: 177–186.

[78] Sherman BT, Hao M, Qiu J, et al. DAVID: a web server for functional enrichment analysis and
functional annotation of gene lists (2021 update). *Nucleic Acids Res* 2022; 50: W216–W221.

[79] Iriki T, Iio H, Yasuda S, et al. Senescent cells form nuclear foci that contain the 26S proteasome.
Cell Rep 2023; 42: 112880.

[80] Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image
analysis. *Nature Methods 2012 9:7* 2012; 9: 676–682.

[81] Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light microscopy.
J Microsc 2006; 224: 213–32.

[82] Clark SJ, Keenan TDL, Fielder HL, et al. Mapping the Differential Distribution of
Glycosaminoglycans in the Adult Human Retina, Choroid, and Sclera. *Investigative Opthalmology*
& Visual Science 2011; 52: 6511.

[83] Edelstein AD, Tsuchida MA, Amodaj N, et al. Advanced methods of microscope control using
µManager software. *J Biol Methods* 2014; 1: e10.

[84] Ovesný M, Křížek P, Borkovec J, et al. ThunderSTORM: A comprehensive ImageJ plug-in for
PALM and STORM data analysis and super-resolution imaging. *Bioinformatics* 2014; 30: 2389–
2390.

[85] Elmore MRP, Najafi AR, Koike MA, et al. Colony-stimulating factor 1 receptor signaling is
necessary for microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron*
2014; 82: 380–397.

[86] Tap WD, Wainberg ZA, Anthony SP, et al. Structure-Guided Blockade of CSF1R Kinase in
Tenosynovial Giant-Cell Tumor. *New England Journal of Medicine* 2015; 373: 428–437.

[87] Braak H, Alafuzoff I, Arzberger T, et al. Staging of Alzheimer disease-associated neurofibrillary
pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* 2006; 112: 389–
404.

[88] Mirra SS, Heyman A, McKeel D, et al. The Consortium to Establish a Registry for Alzheimer's
Disease (CERAD). *Neurology* 1991; 41: 479–486.

[89] Thal DR, Rüb U, Orantes M, et al. Phases of Aβ-deposition in the human brain and its relevance for
the development of AD. *Neurology* 2002; 58: 1791–1800.

[90] McKeith IG, Boeve BF, Dickson DW, et al. Diagnosis and management of dementia with Lewy
bodies. *Neurology* 2017; 89: 88–100.

[91] Braak H, Tredici K Del, Rüb U, et al. Staging of brain pathology related to sporadic Parkinson's
disease. *Neurobiol Aging* 2003; 24: 197–211.

[92] Saito Y, Ruberu NN, Sawabe M, et al. Staging of Argyrophilic Grains: An Age-Associated
Tauopathy. *J Neuropathol Exp Neurol* 2004; 63: 911–918.

[93] Nelson PT, Dickson DW, Trojanowski JQ, et al. Limbic-predominant age-related TDP-43
encephalopathy (LATE): consensus working group report. *Brain* 2019; 142: 1503–1527.

1087

Supporting information

S1 Fig. Protein Extraction by Phase Transfer Surfactant (PTS) Buffer. Coomassie
brilliant blue staining of young (3-month-old) and aged (28-month-old) mouse brains.
Whole brains were lysed with PTS buffer and centrifuged at 15,000 × *g*.

S2 Fig. Detailed Immunohistochemistry Analysis of HAPLN2 in Mouse Brains. (A)

Magnified immunohistochemistry images of the mouse cerebellar white matter stained

with anti-HAPLN2 antibody or biotinylated hyaluronic acid-binding protein (HABP),

which probes hyaluronic acid. Specific staining of HAPLN2 and HABP was observed

without crossover. (B, C) Fluorescence immunohistochemistry images of the mouse

cerebellar white matter showing staining for ubiquitin (B) and p62 (C). Scale bars, 10

μm.

S3 Fig. Aggregation Propensity of Recombinant HAPLN2 Protein. (A, B) Brightfield

microscopic images showing aggregates of recombinant HAPLN2 incubated under

various buffer conditions in the absence (A) or presence (B) of hyaluronic acid. Scale

bars represent 10 μm. (C) Histogram depicting the area of individual aggregates of

recombinant HAPLN2 after 24 hours of incubation in the indicated buffer. Aggregate

areas were measured using ImageJ (version 1.53t). (D) Transmission electron microscopy

images of recombinant HAPLN2 aggregates formed under 150 mM NaCl (pH 6.0). Scale

bars represent 1 μm (black) and 100 nm (white). (E) Schematic representation of cross-β-

positive oligomers and fibrils stained with thioflavin T and Nile Red.

S4 Fig. Microglial Activation Induced by Injecting Recombinant HAPLN2 Protein

into the Mouse Hippocampus. (A) Filter trap assay and immunoblot analysis of

recombinant full-length HAPLN2, HAPLN2 Ig-like domain, and oxStayGold proteins.

These recombinant proteins (2 μM each) were incubated overnight at 37ºC, followed by

centrifugation at $100,000 \times g$ for one hour at 4ºC. (B) Immunohistochemistry images of

the coronal brain section from Figure 8E, stained with anti-Iba1 antibody to probe

activated microglia. Scale bar, 500 μm.

S1 Table. 31 proteins enriched in the sarkosyl-insoluble P2 fraction of aged mouse

brains in Figure 1A.

S2 Table. 45 proteins enriched in the PTS-soluble supernatant of aged mouse brains

in Figure 1B.

S3 Table. Human donors applied for the immunohistochemical analysis in Figure 9.

The stages of neurodegenerative diseases in post-mortem brain samples were examined

using the following criteria. AT8: Tangle pathology (AT8 Stage)[87], CERAD: The

Consortium to Establish a Registry for Alzheimer's Disease, assessing neuritic

plaques[88], Thal: Amyloid-beta deposition (Thal Amyloid Phase)[89], DLB (4th

Consensus): The Dementia with Lewy Bodies criteria (4th Consensus Report)[90], LB

Braak: Lewy body pathology stages (Braak Staging for Lewy Bodies)[91], Grain:

Argyrophilic grain pathology (Saito Scale)[92], Joseph-TDP: TDP-43 proteinopathy

(Josephs' Staging)[93].

S4 Table. Primary Antibodies Used in Immunoblot and Immunohistochemistry

Analyses.

A

B

C

Protein	Fold change (aged)/(young)	
	Fig 1A	Fig 1B
Hyaluronan and proteoglycan link protein 2 (HAPLN2)	48.65	2.121
Immunoglobulin kappa constant (IGKC)	22.89	1.991
Bridging integrator 1 protein (BIN1)	6.384	1.990
Immunoglobulin mu chain C region (IGHM)	19.66	1.936
Aggrecan core protein (ACAN)	9.354	1.724
Claudin-11 (CLDN11)	13.19	1.550
Versican core protein (VCAN)	35.73	1.522
Hyaluronan and proteoglycan link protein 1 (HAPLN1)	7.786	1.284
Alpha-crystallin B chain (CRYAB)	8.017	1.247

D

A

	3 months	6 months	12 months	18 months	
30 kDa					IB: HAPLN2, S1 fraction
30 kDa					IB: HAPLN2, P2 fraction: Sarkosyl insoluble
50 kDa					IB: α-tubulin, S1 fraction

B   **C**

A

| 150 mM NaCl | − | + | − | + |
| 1 mg/mL Hyaluronic acid | − | − | + | + |

pH 6.0

pH 7.0

pH 8.0

B

1 mg/mL Hyaluronic acid	−			+		
	0 mM	150 mM	300 mM	0 mM	150 mM	300 mM
pH 6.0	N	A	A	O	A	A
pH 7.0	N	O	A	N	A	A
pH 8.0	N	N	O	N	O	A

NaCl (mM)

A: amorphous aggregates
O: oligomer
N: not detected

C

	Brightfield	Thioflavin T	Nile Red	Merge
pH 6.0				
pH 7.0				
pH 8.0				

D

The long axis of Nile Red-positive aggregates (nm)

<0.0001

<0.0001

pH: 6.0, 7.0, 8.0

E

Circularity of Nile Red-positive aggregates

0.8681

0.7388

pH: 6.0, 7.0, 8.0

A

Contralateral Ipsilateral

B

DAPI + HAPLN2 + HABP

PBS		Hyaluronidase (HAse)	
Contralateral	Ipsilateral	Contralateral	Ipsilateral

1 mm

C

0.8024 <0.0001

<0.0001

Area of HAPLN2 (µm²)

Contralateral	Ipsilateral	Contralateral	Ipsilateral
PBS		HAse	

D

	PBS				HAse			
Mouse Replicate	1	2	3	4	1	2	3	4

Fraction

S2   IB: HAPLN2
P2

S2   IB: α-tubulin

E

Relative protein levels HAPLN2 in the S2 fraction

0.9990   0.6598

0.9721

Contralateral	Ipsilateral	Contralateral	Ipsilateral
PBS		HAse	

Relative protein levels HAPLN2 in the P2 fraction

0.6996   0.0011

0.1210

Contralateral	Ipsilateral	Contralateral	Ipsilateral
PBS		HAse	

A HAPLN2 + Caspr + DAPI

60–69 years old   80–89 years old

B

<0.0001

Area of HAPLN2 (µm²)

60–69   80–89

Age (years)

C

0.0207

Average area of HAPLN2 (µm²)

60–69   80–89

Age (years)