1

Aggregation of HAPLN2, a component of the perinodal extracellular matrix, is a hallmark of physiological brain aging in mice

Ayaka Watanabe[1][¶], Shoshiro Hirayama[1][¶], Itsuki Kominato[1], Sybille Marchese[2], Pietro Esposito[2], Vanya Metodieva[2], Taeko Kimura[3], Hiroshi Kameda[4], Terunori Sano[5], Masaki Takao[5], Sho Takatori[3], Masato Koike[4], Juan Alberto Varela[6], Taisuke Tomita[3], Shigeo Murata[1][*]

[1] Laboratory of Protein Metabolism, Graduate School of Pharmaceutical Sciences, the University of Tokyo, Tokyo, Japan

[2] School of Biology, University of St Andrews, St Andrews, United Kingdom

[3] Laboratory of Neuropathology and Neuroscience, Graduate School of Pharmaceutical Sciences, the University of Tokyo, Tokyo, Japan

[4] Department of Cell Biology and Neuroscience, Juntendo University Graduate School of Medicine, Tokyo, Japan

[5] Department of Clinical Laboratory, National Centre of Neurology and Psychiatry, Tokyo, Japan

[6] School of Physics and Astronomy, University of St Andrews, St Andrews, United Kingdom

[*] Corresponding author

Email: smurata@g.ecc.u-tokyo.ac.jp (SMurata)

[¶]These authors contributed equally to this work.

Abstract

Protein aggregation is a hallmark of neurodegenerative diseases and is also observed in the brains of elderly individuals without such conditions, suggesting that aging drives the accumulation of protein aggregates. However, the comprehensive understanding of age-dependent protein aggregates involved in brain aging remains unclear. Here, we investigated proteins that become sarkosyl-insoluble with age and identified hyaluronan and proteoglycan link protein 2 (HAPLN2), a hyaluronic acid-binding protein of the extracellular matrix at the nodes of Ranvier, as an age-dependent aggregating protein in mouse brains. Elevated hyaluronic acid levels and impaired microglial function reduced the clearance of HAPLN2, leading to its accumulation. HAPLN2 oligomers induced microglial inflammatory responses both *in vitro* and *in vivo*. Furthermore, age-associated HAPLN2 aggregation was also observed in the human cerebellum. These findings suggest that HAPLN2 aggregation results from age-related decline in brain homeostasis and may exacerbate the brain environment by activating microglia. This study provides new insights into the mechanisms underlying cerebellar aging and highlights the role of HAPLN2 in age-associated changes in the brain.

Author Summary

To identify age-dependent protein aggregates unrelated to disease, we analyzed the sarkosyl-insoluble proteome of young and aged mouse brains. We discovered that hyaluronan and proteoglycan link protein 2 (HAPLN2), a hyaluronic acid-binding protein existing in the nodal extracellular matrix, accumulated with age. Age-dependently accumulated HAPLN2 formed large protein aggregates that cannot be solubilized by the anionic detergent sarkosyl or by hyaluronidase digestion. In addition, HAPLN2 formed irregularly shaped puncta that were mislocalized from the nodes of Ranvier in the cerebellar white matter of not only aged mice but also aged human brains. Oligomers of full-length HAPLN2 specifically induced microglial activation *in vitro* and *in vivo*. Our findings suggest that the accumulation of HAPLN2 aggregates is a new hallmark of brain aging and a possible factor contributing to brain inflammation.

Introduction

Protein aggregation is associated with a wide range of physiological declines, including neurodegenerative diseases, with amyloid β (Aβ) and tau being prominent examples. These proteins aggregate to cause neurodegenerative diseases such as Alzheimer's disease, disrupting neuronal function and ultimately causing cell death. Intracellular aggregates interfere with essential cellular processes such as protein folding, trafficking, and degradation, resulting in the accumulation of misfolded proteins and cellular stress [1]. This disruption impairs crucial neuronal functions, including transcription [2], mitochondrial function [3], and stress granule dynamics [4, 5]. In neurodegenerative diseases, the accumulation of protein aggregates is also observed in the extracellular space. These aggregates activate microglia, triggering the release of pro-inflammatory cytokines and neurotoxic factors, leading to chronic inflammation detrimental to neuronal health [6].

The impairment of clearance systems in both intracellular and extracellular spaces is thought to be responsible for the accumulation of aberrant protein aggregates. The main intracellular pathways for clearing abnormal proteins are the ubiquitin-proteasome system [7] and the autophagy-lysosome system [8, 9]. Additionally, extracellular mechanisms such as phagocytosis, secretion of proteases, and digestive exophagy by microglia [10–12], and the glymphatic system contribute to the clearance of protein aggregates from the brain [13]. However, aging is associated with a decline in the capacity of these mechanisms to maintain proteostasis, making organisms more susceptible to protein aggregation. This decline is attributed to factors such as reduced efficiency of protein degradation pathways, decreased expression of chaperone proteins, impaired cellular stress responses, and microglial senescence [14, 15].

Interestingly, aggregates of wild-type Aβ, tau, and TMEM106B have been observed in the brains of non-demented elderly individuals [16–18]. This suggests that protein aggregation is not merely a consequence of neurodegenerative diseases but also an inherent aspect of the aging process itself, raising the possibility that age-dependent protein aggregates contribute to the broader physiological decline during aging.

Indeed, the detrimental effects of age-related protein aggregation have been demonstrated

in model organisms. In *Caenorhabditis elegans*, the accumulation of both intracellular

and extracellular protein aggregates is associated with reduced lifespan, regardless of

disease state. In contrast, removing these aggregates extends lifespan. For example,

deficiency in *daf-16*, a transcription factor that regulates several antioxidant and

chaperone genes, or in *hsf-1*, a master transcription factor of the heat shock response,

induces protein aggregation and shortens lifespan [19–21]. Conversely, reduced function

of *daf-2*, a homolog of the mammalian insulin/insulin-like growth factor-1 receptor that

suppresses *daf-16* activity, suppresses protein aggregation and extends lifespan [22].

Furthermore, analysis of RIPA buffer-insoluble and sodium deoxycholate-insoluble

proteins by mass spectrometry has revealed that protein aggregation occurs extensively

during physiological aging in the absence of disease [23, 24]. These findings suggest that

the clearance of age-related protein aggregates in *C. elegans* is crucial for maintaining

individual health.

An extracellular protein quality control mechanism exists that disposes of extracellular

waste in *C. elegans*. Coelomocytes remove extracellular debris through non-specific

phagocytosis [25]. Moreover, overexpression of extracellular chaperones in the

pseudocoelom suppresses the accumulation of aggregates and extends lifespan [26].

These findings highlight the importance of extracellular protein quality control and

emphasize the essential role of maintaining proteostasis in the extracellular space to

ensure healthy aging.

In contrast to these studies about *C. elegans*, less is known about how age-dependent

protein aggregates contribute to the aging process in mammals. To address this gap, we

employed proteomic analyses to identify proteins that aggregate and accumulate in the

mouse brain with age. Our investigation identified hyaluronan and proteoglycan link

protein 2 (HAPLN2), a perinodal extracellular matrix constituent protein, as an age-

dependent protein aggregate. We found that HAPLN2 accumulates in the cerebellar white

matter and is mislocalized from its typical location at the nodes of Ranvier in aged mouse

brains. Further investigation using recombinant protein revealed that HAPLN2 forms

aggregates under acidic conditions and in the presence of hyaluronic acid, mimicking

age-related changes in the brain microenvironment. Importantly, we also observed age-dependent HAPLN2 aggregation in human brain sections. These findings suggest that HAPLN2 aggregation may be a conserved hallmark of brain aging across mammalian species. Our study highlights the importance of investigating age-related protein aggregation beyond the context of specific neurodegenerative diseases and suggests a potential role for HAPLN2 aggregates in brain aging, possibly by inducing microglial activation and neuroinflammation.

Results

Proteomic analyses of aged mouse brains identified proteins that accumulate in the sarkosyl-insoluble fraction with age

To identify proteins that aggregate in an age-dependent manner, we enriched detergent-insoluble proteins from whole brains of young (2-month-old) and aged (24-month-old) mice using 1% N-lauryl sarcosine (sarkosyl) fractionation (Fig 1A). Sarkosyl is a well-established detergent that solubilizes natively folded proteins while enriching protein aggregates associated with neurodegenerative diseases in its insoluble P2 fraction (Fig 1A) [27–32]. Proteins in the sarkosyl-insoluble P2 fraction were subjected to label-free quantitative proteomics. We identified 57 proteins that showed more than a 5-fold increase in the brains of aged mice compared to young mice and considered them candidates for age-dependent protein aggregates (S1–S3 Tables). To elucidate the characteristics, we conducted gene ontology (GO) term enrichment analysis on these proteins (Fig 1B). The term "hyaluronic acid binding" was significantly enriched, encompassing aggrecan core protein (ACAN), hyaluronan and proteoglycan link protein (HAPLN) 1, HAPLN2, and versican core protein (VCAN), which were previously reported to increase with age in rodent brains [33–35]. In addition, tumor necrosis factor-inducible gene 6 protein (TNFAIP6), which also possesses hyaluronic acid binding ability and is known to be involved in extracellular matrix remodeling and inflammatory responses, was detected among these proteins [36].

Age-related accumulation of proteins associated with neurodegenerative diseases is

attributed to their inherent resistance to degradation due to aggregation and age-related

declines in clearance mechanisms [37, 38]. Based on this, we hypothesized that proteins

prone to aggregation in the aging mouse brain may also exhibit increased total abundance

compared to those in the brains of younger mice. To determine whether these proteins

also increase in total abundance, we performed further mass spectrometry analyses.

Whole brains from young and aged mice were lysed with phase transfer surfactant (PTS)

buffer (0.5% deoxycholic acid and 0.33% sarkosyl), to increase the solubility of

hydrophobic proteins, including proteins that remain insoluble in sarkosyl alone, and to

enhance their trypsin digestion [39]. The PTS-soluble supernatant, which contains most

of the brain proteome (Fig S1) and mainly corresponds to the proteins shown in S1 and

partly in S2 fractions of Figure 1A, was subjected to label-free proteomic analysis (Fig

1C). We identified 28 proteins that significantly increased by more than 2-fold in the

aged mouse brains compared to young mouse brains (S4–6 Tables). Finally, we narrowed

down the candidates to seven proteins showing an age-dependent increase in both

proteomic analyses (Fig 1D). Although HTRA1 showed the largest fold change in Figure

1A, we could not obtain antibodies to validate this result. Additionally, a reliable

antibody against VCAN could not be obtained for use in western blotting or

immunohistochemistry. Therefore, we focused on ACAN, HAPLN1, and HAPLN2 as

candidates for proteins that aggregate and accumulate with age.

Figure 1. Mass Spectrometry Analyses Revealed Sarkosyl-Insoluble Proteins

Accumulated in Aged Mouse Brains. (A) Workflow (left) illustrating the fractionation

of young (2-month-old) and aged (24-month-old) mouse brains to obtain sarkosyl-

insoluble protein aggregates. A volcano plot (right) represents the results of label-free

quantitative proteomic analysis. The cut-off was set at a false discovery rate (FDR) of

0.01. The original results table and results with imputation of the missing values are listed

in S1 and S2 Tables, respectively. The colored region in the volcano plot highlights

proteins with a fold change > 5.0 and *P*-value < 0.01, as listed in S3 Table. n = 5. Specific

proteins are highlighted: ACAN (green), HAPLN1 (orange), HAPLN2 (red), and VCAN

(blue). (B) Bar chart showing gene ontology (GO) terms in the molecular function

category enriched among the proteins identified in (A). GO enrichment analysis was

performed using the Database for Annotation, Visualization, and Integrated Discovery

(DAVID). Proteins included in the enriched categories are indicated on the right side of

the chart. (C) Label-free proteomic analysis of phase transfer surfactant (PTS)-soluble

supernatant fractions from young (2-month-old) and aged (24-month-old) mouse brains.

The cut-off was set at FDR = 0.01. The original results table and results with imputation

of missing values are listed in S4 and S5 Tables, respectively. Proteins with a fold change

> 2.0 and *P*-value < 0.05 are highlighted in the colored regions of the volcano plots and

listed in S6 Table. n = 4. (D) A list of proteins significantly increased in both proteomic

analyses described in (A) and (C). *P*-values by multiple unpaired *t*-test and two-stage

step-up (Benjamini, Krieger, and Yekutieli) was used to correct for multiple comparisons.

Data underlying this figure are available from the ProteomeXchange Consortium via the

jPOST partner repository (https://repository.jpostdb.org/) with the dataset identifier

JPST003565 and PXD060043 for (A), and JPST003863 and PXD064908 for (C).

191   HAPLN1 and HAPLN2 formed salkosyl-insoluble protein assemblies in aged mouse

192   brains

To validate the proteomic findings, we performed immunoblotting on each fraction

prepared as shown in Figure 1A. Consistent with the proteomic analyses, marked age-

dependent increases in ACAN, HAPLN1, and HAPLN2 were observed in the sarkosyl-

insoluble P2 fraction (Figs 2A and 2B). As expected from previous studies, ubiquitinated

proteins also increased in the P2 fraction with age (Figs 2A and 2B), confirming the

successful enrichment of protein aggregates [33]. These results also demonstrated the

sarkosyl-insolubility of ACAN, HAPLN1, and HAPLN2 in aged mouse brains.

Characteristics of protein aggregates include not only detergent-insolubility but also the

formation of high-molecular-weight assemblies trapped on nitrocellulose membranes and

aberrant localization as large deposits *in vivo* [34, 35]. Therefore, we performed a series

of experiments to determine whether ACAN, HAPLN1, and HAPLN2 are aggregated in
aged mouse brains.

To investigate whether ACAN, HAPLN1, and HAPLN2 form sarkosyl-insoluble large
protein assemblies, we performed a filter trap assay using the S1 fraction. To
simultaneously detect large protein assemblies and total protein, we used a dual
membrane filter trap assay by sequentially layering a nitrocellulose membrane (0.45-μm
pore size) on top of a PVDF membrane (0.2-μm pore size) [40–42]. The nitrocellulose
membrane captures large protein assemblies, while proteins passing through are trapped
by the PVDF membrane, which has a higher protein-binding capacity. Our results
showed that HAPLN1 and HAPLN2 trapped on the nitrocellulose membrane increased
markedly with age, whereas ACAN did not differ between young and aged mice (Fig
2C). In contrast, all three proteins increased on the PVDF membrane in aged mice. These
findings indicate that HAPLN1 and HAPLN2 form sarkosyl-insoluble large complexes
with age, while ACAN does not. ACAN, HAPLN1, and HAPLN2 are known to form the
extracellular matrix via non-covalent association with hyaluronic acid. Previous studies
showed that ACAN from aged mouse brain is enriched in the Triton X-100-insoluble
fraction but becomes soluble after degrading high-molecular-weight hyaluronic acid [43].
Therefore, we next investigated whether treatment with hyaluronidase solubilizes ACAN,
HAPLN1, and HAPLN2 from the sarkosyl-insoluble P2 fraction. We treated the sarkosyl-
insoluble P2 fraction with hyaluronidase, followed by ultracentrifugation to yield the
supernatant hyaluronidase-extractable S3 and the pellet hyaluronidase-resistant P3
fractions (Fig 2D). ACAN was recovered in the S3 fraction after hyaluronidase treatment
(Fig 2E). In contrast, the majority of HAPLN1 and HAPLN2 remained in the P3 fraction,
with only a portion shifting to the S3 fraction (Fig 2E). We also performed a filter trap
assay on the S3 and P3 fractions. HAPLN1 and HAPLN2 in the P3 fraction were
captured on a nitrocellulose membrane, whereas ACAN in the P3 fraction was not
captured on a nitrocellulose membrane (Fig 2F). These results suggest that HAPLN1 and
HAPLN2 form high molecular weight protein assemblies that remain insoluble in
sarkosyl despite hyaluronic acid degradation. We focused on HAPLN2 for further
investigation because HAPLN2 showed the highest fold change in western blot analysis

among ACAN, HAPLN1, and HAPLN2, and its aggregates were not solubilized by

hyaluronidase treatment (Fig 1D).

Figure 2. Extracellular Matrix Component Proteins Accumulate in the Sarkosyl-

Insoluble Fraction of Mouse Brains with Age. (A) Immunoblot analyses of ACAN,

HAPLN1, HAPLN2, and ubiquitin in the S1, P1, S2, and P2 fractions, as outlined in

Figure 1A, from mouse brains. n = 2. (B) Densitometric quantitation of (A). The

expression levels of the indicated proteins were normalized to α-tubulin in the S1

fractions. Error bars represent mean ± S.E.M. *P*-values by two-tailed Student's *t*-test.

***p < 0.001, ****p < 0.0001. n = 4. (C) Filter trap assay and immunoblot analysis

of the S1 fraction from young (3-month-old) and aged (21-month-old) mouse brains. n =

2. (D) Diagram illustrating hyaluronidase treatment of the sarkosyl-insoluble P2 fraction.

The P2 fraction was treated with hyaluronidase and centrifuged at $100,000 \times g$. (E)

Immunoblot analyses of ACAN, HAPLN1, and HAPLN2 in the hyaluronidase-

extractable S3 and hyaluronidase-resistant P3 fractions from 27-month-old mouse brains.

n = 3. (F) Filter trap assay of the fractions described in (E). n = 3.

HAPLN2 was deposited in the extracellular space in the cerebellar white matter of

aged mice

To investigate whether HAPLN2 forms aberrant deposits in aged mouse brains, we

examined the distribution and morphological structure of HAPLN2 in mouse brains.

Previous studies have shown that HAPLN2 is expressed in the cerebellar white matter of

young adult mice [44, 45]. Immunoblot analysis confirmed that HAPLN2 protein was

predominantly detected in the cerebellum of 24-month-old mice, whereas it was only

faintly detected in the cerebellum of 4-month-old mice (Fig 3A). Immunohistochemical

analysis revealed that the expression of HAPLN2 increased with age in highly myelinated

regions, such as brain stem, corpus callosum, and the cerebellar white matter, which is

mainly composed of myelinated axons expressing myelin basic protein (MBP) and is

surrounded by granule layers stained with 4',6-diamidino-2-phenylindole (DAPI) (Figs

3B and S2A). These results demonstrate that HAPLN2 accumulates in highly myelinated

regions of aged mice.

We next examined its detailed structure in the cerebellar white matter of aged mice,

where the age-dependent accumulation of HAPLN2 was most prominent.

Immunohistochemistry analysis showed significantly enlarged puncta of HAPLN2 in the

aged mouse brains compared to young mouse brains (Figs 3C and 3D). We then

investigated whether the HAPLN2 deposits are localized to the intracellular or

extracellular space. Hyaluronic acid is a major component of the extracellular matrix,

which is visualized by staining with hyaluronic acid binding protein (HABP). In young

mouse samples, HABP staining was weak, and colocalization of HAPLN2 and HABP

was unclear. In contrast, in aged mouse brains, both HABP and HAPLN2 staining

showed significantly enlarged puncta, and colocalization of HAPLN2 and HABP was

clearly observed (Figs 3E–G, and S2B). These results indicate that HAPLN2 accumulates

as large deposits with hyaluronic acid in the extracellular space of the cerebellar white

matter of aged mice.

To determine which cells in the cerebellum express HAPLN2 and whether the

accumulation of HAPLN2 is due to increased mRNA expression or not, we referred to

the results of a previous single-cell transcriptome analysis. This data revealed that *Hapln2*

mRNA expression was specifically detected in oligodendrocytes and that mRNA levels

remained constant between 2–3 months and 21–22 months of age (Fig 3H) [46]. These

results suggest that the age-dependent accumulation of HAPLN2 protein is not due to the

induction of mRNA expression.

Since the single-cell RNA-seq result indicates that HAPLN2 is expressed in

oligodendrocytes, we performed co-staining of HAPLN2 and the oligodendrocyte marker

MBP [47]. However, the HAPLN2 deposits did not colocalize with oligodendrocytes (Fig

3I). We also examined co-staining of HAPLN2 with ubiquitin and p62, as protein

aggregates that initially form intracellularly and are later deposited extracellularly, as is

typically observed with tau, often colocalize with these proteins [48]. However, HAPLN2

did not colocalize with ubiquitin and p62 (Figs S2C and S2D). These results suggest that

HAPLN2 is secreted from oligodendrocytes and subsequently accumulated in the

extracellular space.

We next investigated whether enlarged HAPLN2 puncta retain their physiological

function. HAPLN2 is normally localized to the perinodal extracellular matrix, which

surrounds the nodes of Ranvier and contains hyaluronic acid [49]. Therefore, we

examined the colocalization between HAPLN2 and contactin-associated protein (Caspr),

which localizes to the paranodal junction. The colocalization of HAPLN2 with Caspr

decreased with age (Figs 3J and 3K). In addition, Mander's overlap coefficient (MOC)

M1, which indexes the proportion of HAPLN2 colocalized with Caspr, decreased with

age (Fig 3L), suggesting that HAPLN2 was abnormally localized outside the nodes of

Ranvier in aged mice. In contrast, MOC M2, an index of the proportion of Caspr

colocalized with HAPLN2, did not change with age (Fig 3M), suggesting that most of

Caspr colocalizes with functional HAPLN2 and that the nodes of Ranvier are not

dysregulated in aged mice. Additionally, the area of HAPLN2 mislocalized from Caspr

was larger than HAPLN2 colocalized with Caspr in 24-month-old mice (Fig 3N). This

further supports that enlarged HAPLN2 puncta have lost their function.

To investigate whether the accumulation of HAPLN2 is associated with age-related

demyelination, brain slices from young (3-month-old) and aged (18-month-old) mice

were stained for HAPLN2, calbindin 1, a marker of Purkinje cells, the most abundant

neuronal type in the cerebellum, and MBP. Immunohistochemical analysis showed that,

with aging, MBP-negative regions increasingly appeared as hole-like structures within

the white matter, and these regions exhibited a corresponding increase in calbindin 1

staining, as indicated by the magenta arrows in the inset figures (Figs S3A and 3B). This

pattern was consistent with the demyelination-associated changes observed by Luxol Fast

Blue (LFB) staining [50, 51]. Additionally, the calbindin 1-positive area was increased in

aged mice compared to young mice, suggesting that demyelination exposes the axons of

Purkinje cells (Fig S3C). Specifically, while the calbindin 1-positive area accounted for

approximately 1.2% in young mice, it increased nearly threefold in aged mice to

approximately 3.3%.

To directly assess the relationship between demyelination severity and HAPLN2

accumulation, we compared cerebellar regions in aged mice exhibiting high versus low

calbindin 1 staining (Figs S3D and S3E). Specifically, calbindin 1-high and calbindin 1-

low regions were defined as square areas measuring $50\,\mu m \times 50\,\mu m$ in which the

calbindin 1-positive area was more than 5% or less than 1%, respectively. These

thresholds were determined based on the quantification shown in Figure S3C, where the

average calbindin 1-positive area in 18-month-old mice was 3.3%, nearly threefold higher

than the 1.2% observed in 3-month-old mice. Quantitative analysis revealed no

significant differences in HAPLN2-positive puncta size or density between these regions.

These results demonstrate that HAPLN2 accumulation occurs independently of age-

related demyelination severity.

Furthermore, we investigated whether other proteins are deposited in the aged brain.

Immunostaining of brain slices from young and aged mice for HAPLN1 and ACAN

revealed that, in aged mice, HAPLN1 formed puncta that were mislocalized relative to

ACAN staining in the cerebellar nuclei, as indicated by the magenta arrows (Figs S4A

and S4B). In addition, the area of HAPLN1 not colocalized with ACAN increased with

age, whereas the ACAN-positive areas showed no significant age-related changes (Fig

S4C). These findings suggest that HAPLN1 undergoes age-related deposition and loses

its function in the perineuronal net in the aged brain.

Figure 3. HAPLN2 Forms Aggregate-Like Structures in the Cerebellar White

Matter of Aged Mice. (A) Immunoblot analysis of the S1 fraction from the cerebrum

and cerebellum of mice. n = 2. (B) Representative images of fluorescence

immunohistochemistry of the mouse cerebellum showing HAPLN2, myelin basic protein

(MBP, a marker for myelinated axons), and 4',6-diamidino-2-phenylindole (DAPI). n = 3.

(C) Magnified fluorescence immunohistochemistry images for HAPLN2 corresponding

to the boxed areas in the left panels. n = 3. (D) Quantitation of the mean area of

HAPLN2-positive puncta in (C). (E) Representative images of fluorescence

immunohistochemistry for HAPLN2 and biotinylated hyaluronic acid-binding protein

(HABP), a probe for hyaluronic acid. n = 3. (F) Quantitation of the mean area of HABP-

positive puncta in (E). (G) Pearson's correlation coefficients for colocalization between

HAPLN2 and HABP. (H) Single-cell transcriptomics data from Ximerakis et al., *Nat*

Neurosci (2019). TPM: transcripts per million. ASC: astrocyte; MG: microglia; NEUR:

mature neuron; OLG: oligodendrocyte. Error bars indicate maximum values, boxes

indicate the first and third quartiles, and solid and dashed lines represent the median and

mean, respectively. (I) Representative images of fluorescence immunohistochemistry for

HAPLN2 and MBP as an oligodendrocyte marker in the cerebellar white matter. n = 3.

(J) Representative images of fluorescence immunohistochemistry for HAPLN2 and

Caspr, a marker for the nodes of Ranvier. n = 3. (K–M) Colocalization analysis between

HAPLN2 and Caspr using Pearson's correlation coefficient (K) and Mander's overlap

coefficients (MOC) M1 (L) and M2 (M). MOC M1 represents the proportion of HAPLN2

overlapping Caspr, while MOC M2 represents the reverse. (N) Quantitation of the mean

area of HAPLN2 punctum, categorized by whether it colocalizes with Caspr in 24-month-

old mice. All coefficient calculations were performed using ImageJ (version 1.53t) with

the JACoP plugin. Scale bars: 1 mm (B and C (left)) and 10 µm (C (right); E; I; J). Error

bars represent mean ± S.D. *P*-values by two-tailed Student's *t*-test. **$p < 0.01$,

***$p < 0.001$, ****$p < 0.0001$, ns = not significant.

Deposition of HAPLN2 aggregates starts in middle age in mouse brains

Research in humans and rodents has highlighted middle age as a period marked by the

onset of changes associated with brain aging [52]. To investigate when HAPLN2 starts to

accumulate, we examined the age-related progression of HAPLN2 accumulation in the

mouse brain. We performed sarkosyl fractionation followed by immunoblot analysis

using young (3- and 6-month-old), middle-aged (12-month-old), and aged (18-month-old)

mice. HAPLN2 began to accumulate in the S1 and P2 fractions from 12-month-old mice

and was further enriched in 18-month-old mice (Fig 4A). Correspondingly, the puncta of

HAPLN2 became larger in 12- and 18-month-old mice, consistent with the immunoblot

analysis (Figs 4B and 4C). These results indicate that HAPLN2 accumulates gradually

from middle age to old age, further confirming the age-dependent accumulation of

HAPLN2 aggregates.

Figure 4. Time-Course Analysis of Age-Dependent HAPLN2 Accumulation in

Mouse Brains. (A) Immunoblot analyses of the S1 and P2 fractions from the mouse

cerebellum at different ages. n = 3. (B) Representative images of fluorescence

immunohistochemistry of the mouse cerebellum showing HAPLN2 and DAPI staining. n

= 3. Scale bars: 1 mm (B) and 10 µm (B, inlet). (C) Quantitation of the mean area of

HAPLN2-positive structures shown in (B). Error bars represent mean ± S.D. *P*-values by

one-way ANOVA followed by Tukey's *post hoc* test. **$p < 0.01$, ***$p < 0.001$.

Low pH, high NaCl concentration, and hyaluronic acid promoted HAPLN2

aggregation

To investigate the factors affecting HAPLN2 aggregation, we explored the aggregation

propensity of recombinant HAPLN2 protein. Various factors have been reported to

influence the formation of protein aggregates, including pH and NaCl concentration. For

instance, the TAR DNA-binding protein of 43 kDa (TDP-43) aggregates at high pH and

NaCl concentration [53], while fused in sarcoma (FUS) aggregates at low concentrations

of RNA [54] and in the presence of dextran [55, 56] and NaCl [57]. Therefore, we

incubated recombinant HAPLN2 protein in buffers with varying pH and NaCl

concentration, and in the presence or absence of hyaluronic acid. The formation of

amorphous aggregates of HAPLN2 was promoted at low pH, high NaCl concentration,

and in the presence of hyaluronic acid (Figs 5A, 5B, S5A, and S5B). Quantitative area

measurements revealed that recombinant HAPLN2 formed amorphous aggregates larger

than 10 µm^2 at pH 6.0, while it formed small oligomer-like aggregates less than 10 µm^2

at pH 7.0, and no aggregates were formed at pH 8.0 in 150 mM NaCl (Fig S5C).

Transmission electron microscopy analysis further confirmed that HAPLN2 incubated at

pH 6.0 in 150 mM NaCl formed amorphous aggregates (Fig S5D).

Many protein aggregates contain cross-β-sheet structures and form protein fibrils.

However, it is impossible to distinguish between oligomeric and fibrillar structures using

diffraction-limited optics [58]. To examine whether HAPLN2 aggregates contain β-sheet-

mediated aggregates that are stained with thioflavin T or Nile Red, we conducted super-

resolution imaging analysis (Fig S5E) [59]. In recombinant HAPLN2 incubated at pH

6.0, some spots, but not the entire aggregate, were stained with thioflavin T and Nile Red,

indicating that the amorphous-appearing HAPLN2 aggregates contain cross-β-sheet

structures (Fig 5C).

To analyze whether the HAPLN2 aggregates formed fibrillar structures, which are

characterized by a long axis and low circularity ratio, we statistically analyzed the

morphology of the Nile Red-positive clustered aggregates, according to a previous report

[60]. The long axis of the Nile Red-positive clusters increased under acidic conditions

(Fig 5D), but the circularity ratio did not change with pH (Fig 5E).

To determine whether aggregation of HAPLN2 induced by low pH results in an increase

in β-sheet structures, we analyzed the circular dichroism (CD) spectra of the protein in

both the soluble state and the aggregated state. Upon protein aggregation and transition to

β-sheet conformation, a red shift of the positive band is observed in the CD spectrum.

[61]. We found that recombinant HAPLN2 proteins that aggregated at low pH exhibited a

shift in positive ellipticity from approximately 190 nm to 200 nm, suggesting an increase

in β-sheets (Fig 5F). Secondary structure analysis of the CD spectra using BeStSel further

confirmed that the proportion of β-sheet structures was significantly increased in the

aggregated state compared to the soluble state, while the α-helix content was reduced

(Figs 5G and 5H) [62].

These results suggest that low pH induces the formation of amorphous, rather than

fibrillar, aggregates of HAPLN2 containing partial cross-β-sheet structures.

Figure 5. Aggregates of Recombinant HAPLN2 Formed under Low pH, High NaCl,

or Hyaluronic Acid Conditions Partially Contain Cross-β-Sheet Structures. (A)

Brightfield microscopic images of recombinant HAPLN2 aggregates incubated in the indicated buffer conditions without agitation. (B) Phase diagrams showing the aggregation states of recombinant HAPLN2 under the indicated buffer conditions. (C) Brightfield and highly inclined and laminated optical sheet (HiLo) images of recombinant HAPLN2 aggregates incubated in a buffer at the indicated pH with 150 mM NaCl. Thioflavin T and Nile Red were used at concentrations of 5 µM and 5 nM, respectively to visualize cross-β-sheet structures. (D, E) Quantitative analyses of the long axis (D) and circularity ratio (E) of Nile Red-positive aggregates shown in (C). Data were collected from at least five fields of view. 500 nM Recombinant HAPLN2 monomers were incubated for 24 hours at 37ºC in all conditions. Error bars represent mean ± S.D. *P*-values by one-way ANOVA followed by Tukey's *post hoc* tests. Scale bars: 10 µm (A) and 1 µm (C). (F) CD spectra of soluble (blue) and aggregated (red) recombinant HAPLN2 protein. (G, H) Secondary structure estimation of α-helix (G) and β-sheet (H) content based on BeStSel analysis of CD spectra of (F). n = 3. Error bars represent the mean ± S.D. *P*-values were calculated using a two-tailed Student's *t*-test. *$p < 0.05$, ****$p < 0.0001$, ns = not significant.

Treatment with hyaluronidase promoted the clearance of HAPLN2 aggregates in the mouse cerebellar white matter

It has been reported that the volume of the extracellular spaces reduces with age, which is thought to lead to impairment of protein aggregate clearance [63, 64]. Hyaluronic acid is the major component of the perinodal extracellular matrix and accumulates in the brain with aging [65–69]. Furthermore, it has been shown that hyaluronidase treatment increases the volume of the extracellular space in mouse hippocampal slices [70]. Therefore, we hypothesized that reducing hyaluronic acid would increase the extracellular space and improve access to HAPLN2 aggregates, thereby enhancing their clearance.

To test this idea, we injected hyaluronidase into the mouse cerebellum (Fig 6A). Immunohistochemical analysis revealed that the expression and size of HAPLN2

dramatically decreased 16 hours after hyaluronidase injection (Figs 6B, 6C). Immunoblot

analysis of brain lysates fractionated as in Figure 1A showed that HAPLN2 and HAPLN1

in the sarkosyl-insoluble P2 fractions in the hyaluronidase-injected brain (ipsilateral)

significantly decreased compared to that in the PBS-injected ipsilateral brain, whereas

HAPLN2 levels in the sarkosyl-soluble S2 fractions were not significantly affected (Figs

6D, 6E, S6A, and S6B).

Since hyaluronidase administration resulted in a marked reduction of HAPLN2

aggregates, we next investigated the underlying mechanism. A recent study has shown

that hyaluronidase injection increased the volume and decreased the viscosity of the

extracellular space, thereby potentially facilitating glymphatic flow [70]. Therefore, we

analyzed changes in the levels of HAPLN2 in the cerebrospinal fluid (CSF) in the

presence and absence of hyaluronidase treatment. Capillary western blot analysis

demonstrated that HAPLN2 protein levels in the CSF were substantially increased three

hours after hyaluronidase administration (Figs 6F, 6G). This observation suggests that

HAPLN2 aggregates may be cleared from the white matter via pathways such as

glymphatic flow, although the involvement of other clearance or degradation mechanisms

cannot be completely excluded. Collectively, these results indicate that hyaluronidase

treatment facilitates the removal of HAPLN2 aggregates from the cerebellar white matter.

Aging is also associated with a decline in motor function, which is attributed to cerebellar

functional decline [71]. Therefore, to investigate the physiological effects of

hyaluronidase treatment, we analyzed the motor function of 12-month-old mice three

months after hyaluronidase injection (Figs 6H–P).An open field test was conducted for

five minutes, during which the following parameters were measured: total locomotor

activity indicators such as walking distance and mean velocity; indicators of burst activity

including maximum velocity; proportions of time spent in different locomotor states

represented by time spent moving and time spent immobile; anxiety-related spatial

preference assessed by time spent incorners; and parameters related to turning behavior

and balance, including mean meander and mean turn angle. The open-field test analysis

revealed that hyaluronidase injection significantly increased walking distance, mean

velocity, and time spent moving, while it decreased time spent immobile, mean meander,

and mean turn angle, suggesting improvements in gait and balance. In contrast, maximum

velocity and time spent in corners were not significantly affected, indicating that

momentary motor abilities, anxiety levels, and exploratory behavior remained unchanged.

These results suggest that age-dependent protein aggregates, including HAPLN2, are

responsible for the decline in the health of the brain.

Figure 6. Hyaluronidase Promotes Clearance of HAPLN2 Aggregates in the Mouse

Cerebellum. (A) Schematic illustration of hyaluronidase injection into the mouse

cerebellum. Created by BioRender. (B) Fluorescence immunohistochemistry images of

the cerebellum of 12-month-old mice showing HAPLN2, HABP, and DAPI signals. n =

4. Scale bar: 1 mm (B), 10 µm (B, inlet). (C) Quantitation of the mean area of HAPLN2-

positive puncta in (B). (D) Immunoblot analysis of the cerebellum corresponding to the

samples in (B). (E) Densitometric quantification of (D). HAPLN2 Protein levels in the S2

fraction were normalized to α-tubulin in the S2 fractions. HAPLN2 protein levels in the

P2 fraction were normalized to α-tubulin in the S1 fractions. (F) Capillary western blot

analysis of cerebrospinal fluid in (B). (G) Densitometric quantification of (F). HAPLN2

protein levels in CSF were normalized to total protein. (H) Schematic illustration of the

time course of the hyaluronidase injection followed by the open field test. (I–P) Open

field test analysis of the hyaluronidase-injected mice. The assessed parameters included:

distance (I), mean velocity (J), maximum velocity (K), time spent moving (L), time spent

immobile (M), time spent in corners (N), mean meander (O), and mean turn angle (P).

The duration of the measurement was five minutes. Four mice were analysed in each

trial. n = 5. Error bars represent mean ± S.D. (C) and S.E.M. (E, G, I–P) *P*-values by one-

way ANOVA followed by Tukey's *post hoc* tests. *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$, ****$p < 0.0001$, ns = not significant.

Microglia regulated the accumulation of HAPLN2 aggregates

Microglia are the resident macrophages in the brain, acting as scavengers that remove debris and protein aggregates from the extracellular space through phagocytosis [72]. To examine whether microglia are involved in the clearance of HAPLN2 aggregates, we treated 4-week-old mice with pexidartinib (PLX3397), a colony-stimulating factor 1 receptor (CSF1-R) inhibitor known to deplete microglia [73]. Immunostaining with ionized calcium-binding adapter molecule 1 (Iba1), a microglia/macrophage marker, confirmed that treatment with PLX3397 for two months successfully depleted microglia (Fig 7A). Immunohistochemical analysis showed that microglial depletion resulted in the accumulation and enlargement of HAPLN2 puncta in the brains of PLX3397-treated mice (Figs 7B and 7C). Furthermore, PLX3397 treatment resulted in HAPLN2 enrichment within the sarkosyl-insoluble fraction of the cerebellum (Figs 7D and 7E). These results suggest that microglia play a critical role in the clearance of HAPLN2 aggregates.

To determine whether microglial depletion induces myelin damage and subsequently contributes to HAPLN2 aggregate accumulation, we compared brain sections from mice treated with PLX3397 to deplete microglia and from untreated controls. Immunostaining was performed using antibodies against Iba1, a microglial marker; calbindin 1, a marker of Purkinje cells; and MBP, a myelin marker. Quantitative analysis revealed no significant changes in calbindin 1-positive areas following microglial depletion, indicating that the depletion protocol used here did not affect the extent of myelin damage or Purkinje cell axon exposure. However, microglial depletion significantly reduced HAPLN2 aggregate clearance. These results suggest that microglia contribute to the clearance of HAPLN2 aggregates through mechanisms distinct from their canonical phagocytic pathway for damaged myelin (Figs S7A–C).

Figure 7. Accumulation of HAPLN2 Aggregates is Promoted by Microglial Depletion in the Cerebellum. (A) Immunohistochemistry images of the cerebellum from mice fed AIN-76A control chow or AIN-76A containing 290 mg/kg PLX3397 from one month to three months of age, stained with anti-ionized calcium-binding adaptor molecule 1 (Iba1) antibody as a microglia marker. n = 3. (B) Immunohistochemistry

images of the cerebellum from the mice in (A), stained with anti-HAPLN2 antibody. n =

3. (C) Quantification of the mean area of HAPLN2-positive puncta in (B). (D)

Immunoblot analysis of the cerebellum from the mice in (B), fractionated as described in

Figure 1A. n = 3. (E) Densitometric quantitation of (D). Protein expression levels were

normalized to α-tubulin in the S1 fractions. Scale bars: 1 mm (A, up; B, up), 50 µm (A,

down), and 100 µm (B, down), 10 µm (B, down, inlet). Error bars represent mean ± S.D.

P-values were calculated using two-tailed Student's *t*-test. *$p < 0.05$,

****$p < 0.0001$,

Oligomers and aggregates of HAPLN2 induced microglial activation *in vitro* and *in*

vivo

Microglia that phagocytose extracellular protein aggregates exhibit a pro-inflammatory

M1 phenotype and secrete cytokines such as interleukin (IL)-6 and tumor necrosis factor

(TNF)-α [72]. Recent studies have shown that soluble oligomers of Aβ, tau, and α-

synuclein in the extracellular spaces induce inflammatory responses prior to the

accumulation of protein aggregates in the brains of patients with neurodegenerative

diseases [74].

To investigate whether HAPLN2 triggers microglial inflammation, we purified

recombinant proteins of full-length HAPLN2 and each domain from a lipopolysaccharide

(LPS)-free *Escherichia coli* strain, ClearColi™ BL21(DE3) (Fig 8A) [75]. First, we

performed a filter trap assay to determine whether solubilized full-length HAPLN2 and

the N-terminal Ig-like domain exist as monomers or soluble oligomers. Both soluble full-

length HAPLN2 and the N-terminal Ig-like domain were trapped on the nitrocellulose

membrane after ultracentrifugation, indicating that they form soluble oligomers (Fig

S6A). We also prepared aggregated full-length HAPLN2 and the N-terminal Ig-like

domain by incubating each soluble protein at pH 6.0, which was confirmed by filter trap

assay (Fig S8A).

To evaluate the potential to induce microglial inflammatory responses, we added full-

length HAPLN2 oligomers and aggregates, as well as the positive control LPS, to the

MG6 microglial cell line. Both soluble oligomers and aggregates of HAPLN2 increased

the mRNA expression of IL-6 and TNF-α (Fig 8B). These results suggest that HAPLN2

induces pro-inflammatory microglial activation in both the soluble oligomeric and

aggregated states.

To determine whether specific domains of HAPLN2 induce pro-inflammatory microglial

activation, we added soluble full-length HAPLN2 and its individual domains to MG6

cells. Soluble oxStayGold fluorescent protein, which was not trapped by the

nitrocellulose membrane, was used as a negative control (Fig S8A). Soluble full-length

HAPLN2 significantly induced IL-6 and TNF-α mRNA expression, but none of the

individual domains of HAPLN2 upregulated these cytokines (Fig 8C). These results

suggest that the pro-inflammatory effect is associated with full-length HAPLN2 and not

with specific domains.

Finally, we examined whether HAPLN2 oligomers induce microglial activation in mouse

brains by infusing full-length HAPLN2 and each domain into the hippocampus. Injection

of full-length HAPLN2 increased pro-inflammatory microglia, marked by Iba1, in the

hippocampus (Figs 8E, 8F, S8B, and S8C). In contrast, oxStayGold and each domain of

HAPLN2 did not significantly induce microglial activation, consistent with Figure 8C.

These results suggest that oligomeric full-length HAPLN2 promotes microglial activation

in vivo.

Figure 8. HAPLN2 Induced Microglial Activation *In Vitro* and *In Vivo*. (A) Diagram

of the domains within HAPLN2. (B) Reverse transcription polymerase chain reaction

(RT-qPCR) analysis of pro-inflammatory cytokine mRNA (*Il6* and *Tnf*, normalized to

Actb) in MG6 cells stimulated with 1 µM LPS, and 200 nM recombinant full-length

HAPLN2 in soluble and aggregated states for four hours. PBS served as a control.

Recombinant HAPLN2 was aggregated by incubation in a buffer containing 10 mM

sodium phosphate and 300 mM NaCl (pH 6.0) for 24 hours at 37°C. (C) RT-qPCR

analysis of pro-inflammatory cytokine mRNA (*Il6* and *Tnf*, normalized to *Actb*) in MG6

cells stimulated with 1 µM LPS, and 200 nM of recombinant proteins (oxStayGold, full-

length HAPLN2, HAPLN2 Ig-like domain, HAPLN2 link 1 domain, and HAPLN2 link 2

domain) for four hours. PBS served as a control. (D) Fluorescence immunohistochemistry

images of the hippocampus of 3-month-old mice injected with PBS, 20 µg LPS, and 15

pmol of recombinant proteins (oxStayGold, full-length HAPLN2, HAPLN2 Ig-like

domain, HAPLN2 link 1 domain, and HAPLN2 link 2 domain). The brains were stained

with anti-Iba1 antibody to detect activated microglia. Scale bar: 500 µm. n = 4. (E)

Quantitation of the number of activated microglia labeled by Iba1 from (E). Error bars

represent mean ± S.D. (C) and S.E.M. (E). *P*-values were calculated using one-way

ANOVA followed by Tukey's *post hoc* tests. *$p < 0.05$, ***$p < 0.001$, ns = not

significant.

HAPLN2 accumulated with age in the human cerebellum

Finally, we examined whether HAPLN2 accumulates with age in the human brain. We

stained HAPLN2 in the cerebellum of human brains in their sixties or eighties without

dementia (S7 Table). We observed the increased HAPLN2 puncta in the human brain in

their eighties compared to those in their sixties (Fig 9A). These HAPLN2 puncta did not

colocalize with Caspr, suggesting that these are age-dependent aggregates. In addition,

the mean count and area of HAPLN2 puncta increased with age (Figs 9B and 9C). These

results suggest that HAPLN2 accumulation is a common hallmark of aging in both mice

and humans.

Figure 9. HAPLN2 Formed Large Puncta with Age in the Human Cerebellum. (A)

Fluorescent immunohistochemistry images for HAPLN2, Caspr, and DAPI in the

cerebellum from human post-mortem brains without dementia. Scale bar: 50 µm. (B)

Quantification of the mean counts of HAPLN2 puncta in (A). (C) Quantification of the

average area of HAPLN2 puncta across all samples in (A). Error bars represent mean ±

S.E.M. *P*-values were calculated using two-tailed Student's *t*-test. *$p < 0.05$.

Discussion

Research on age-related protein aggregation in mammals has progressed at a slower pace

than that in *C. elegans* due to the difficulty in obtaining a sufficient number of aging

individuals. Recently, studies have aimed to identify proteins that become Triton X-100

insoluble in the cerebral cortex with age [76]. However, as demonstrated by the evidence

that membraneless organelle constituent proteins and functional large protein complexes,

such as the cytoskeleton, also become Triton X-100 insoluble, these proteins are not

synonymous with aggregated proteins [76]. Therefore, we used sarkosyl to search for

proteins that aggregate with aging because sarkosyl fractionation is known to isolate

protein aggregates, including Aβ and tau, into the $100,000 \times g$ pellet fraction. We

identified seven proteins that become sarkosyl-insoluble and accumulate in mouse brains

with aging. Of these, five were extracellular matrix components, and a protease, all of

which are localized extracellularly. We focused on HAPLN2, a component of the

perinodal extracellular matrix, as a protein that aggregates in the highly myelinated

regions, such as cerebellum white matter and corpus callosum in an age-dependent

manner. HAPLN2 aggregates colocalized with hyaluronic acid in the cerebellum white

matter in aged mice (Fig 3D). This observation, coupled with the *in vitro* finding that

hyaluronic acid promotes aggregation of HAPLN2 (Fig 5A), suggests a key role for

hyaluronic acid in this process. This is consistent with reports that the concentration of

hyaluronic acid increases, especially in the prefrontal cortex and cerebellum, with age *in*

vivo [69, 77, 78]. Furthermore, it is reported that microglia are fewer in number and more

sparsely distributed in the cerebellum compared to other brain regions [79–81].

Therefore, clearance of protein aggregates by microglial phagocytosis may be less

efficient in the cerebellum compared to the cerebrum. Moreover, the cerebellum has a

significantly higher density of neurons and axons compared to the cerebrum [82]. This

implies a greater number of nodes of Ranvier and their associated perinodal extracellular

matrix, which could contribute to the high expression of HAPLN2 observed in this

region. This may explain the significant HAPLN2 aggregation observed in the

cerebellum.

Experiments using recombinant HAPLN2 revealed that the formed amorphous aggregates

partially contained cross-β-sheet structures, characteristic of proteinaceous aggregates

(Fig 5C). While HAPLN2 binds hyaluronic acid to form a functional perinodal

extracellular matrix, these findings indicate that during aggregation, HAPLN2 forms

aggregates primarily through protein-protein interactions rather than merely forming

hyaluronic acid complexes. Notably, HAPLN2 isolated from the sarkosyl-insoluble P2

fraction of aged mouse brains remained insoluble even after enzymatic digestion of

hyaluronic acid (Figs 2D and 2E). This implies that while hyaluronic acid interactions

may initiate aggregation, HAPLN2 eventually develops protein-protein interactions that

stabilize its sarkosyl-insoluble state.

Adding another layer to the role of hyaluronic acid, we also showed that injecting

hyaluronidase into the mouse cerebellum led to a decrease in sarkosyl-insoluble HAPLN2

by promoting its clearance into the CSF (Fig 6). This reduction may be attributed to

increased accessibility of clearance systems, such as the glymphatic system, resulting

from the expansion of extracellular spaces [70]. This finding is supported by studies

showing that increased extracellular volume enhances Aβ clearance [83]. Furthermore,

aging reduces CSF flow, and this decline impairs the efficiency of the glymphatic system

in clearing extracellular waste [13]. Nonetheless, it remains unclear whether the

glymphatic system can effectively transport large protein aggregates, such as HAPLN2

aggregates. Thus, a critical unanswered question is whether HAPLN2 clearance occurs

via transport of intact aggregates or requires prior disaggregation into a solubilized form.

Future studies will address this mechanistic ambiguity, which has broad implications for

understanding how the glymphatic system handles pathological protein assemblies.

Additionally, recombinant HAPLN2 protein efficiently formed aggregates under low pH

conditions (Fig 5). Although the decrease in brain pH with age is relatively small, ranging

from 6.30 to 6.00 in humans and from 6.85 to 6.70 in mice [84], it cannot be completely

ruled out as a contributing factor to HAPLN2 aggregation. While this modest change in

acidity alone may not be sufficient to induce aggregation, it is possible that it interacts

with other aging-related factors to promote HAPLN2 aggregation. Taken together, our

data suggest that changes in brain pH, hyaluronic acid levels, and the glymphatic flow

may contribute to the aggregation of HAPLN2 *in vivo*, highlighting the complex interplay

between protein aggregation and the aging brain environment, particularly for HAPLN2.

Microglia play a critical role in clearing extracellular waste, such as protein aggregates,

through phagocytosis [72]. Recent studies have also shown that microglia phagocytose

the extracellular matrix, including the perineuronal net [85, 86]. However, aging is

associated with a decline in microglial phagocytic capacity due to increased expression of

CD22, a negative regulator of phagocytosis [11, 45], and decreased expression of

TREM2 and ABCA7, which are well known to mediate phagocytic clearance of Aβ [87,

88]. Our study demonstrated that microglial depletion promotes the accumulation of

HAPLN2 aggregates (Fig 7), suggesting that age-related impairment of microglial

phagocytosis disrupts the clearance of HAPLN2 and the perinodal extracellular matrix.

The accumulation of protein oligomers and aggregates in the extracellular space activates

microglia. This leads to the secretion of pro-inflammatory cytokines, such as IL-6 and

TNF-α, which contribute to chronic inflammation [89]. Our study showed that excessive

HAPLN2 oligomers and aggregates activate microglia both *in vitro* and *in vivo* (Fig 8).

These findings indicate that while microglia are involved in clearing HAPLN2

aggregates, their phagocytic capacity can become overwhelmed by the abnormal

accumulation of HAPLN2 aggregates. This imbalance may trigger microglial pro-

inflammatory activation, further contributing to chronic inflammation, a key feature of

brain aging. While we demonstrated microglial activation both *in vitro* and *in vivo*

following exposure to recombinant HAPLN2 oligomers and aggregates, these findings

remain correlative and do not establish a causal relationship, as we did not manipulate

endogenous HAPLN2 expression *in vivo*. Therefore, the physiological relevance of

HAPLN2 aggregation in the aged brain remains to be fully elucidated. To overcome these

limitations, future studies will employ genetic and pharmacological approaches to clarify

the causal role of HAPLN2 in neuroinflammation. We believe our current findings

provide a solid foundation for these next steps and offer valuable insight into the potential

contribution of HAPLN2 to brain aging.

In summary, we have identified HAPLN2, a component of the perinodal extracellular

matrix, as an age-dependent protein aggregate. HAPLN2 aggregates mislocalize and

accumulate in the extracellular space in aged mouse brains, where hyaluronic acid and

microglia regulate their accumulation and clearance. We also observed HAPLN2

aggregates in non-demented human brains. Our study suggests HAPLN2 aggregation as

a potential factor in chronic inflammation in the brain during physiological aging and

sheds new light on our understanding of age-related decline in brain function.

Materials and Methods

Post-mortem brain samples

The paraffin-embedded postmortem cerebellum derived from neurologically normal

patients in their sixties (n = 4) or eighties (n = 5) were obtained from Japan Brain Bank

Net (JBBN). Information on human donors is provided in S7 Table. All participants,

before passing away or their families, provided written informed consent. Experiments

using human samples were performed with institutional approval and guidelines from the

institutional ethical committee of the Graduate School of Pharmaceutical Sciences,

University of Tokyo (No. 4-8).

Animals

C57BL/6N mice were purchased from Clea Japan. All mice were maintained under

specific pathogen-free conditions. All animal experiments were performed after obtaining

approval from the Institutional Animal Care Committee of the Graduate School of

Pharmaceutical Sciences, the University of Tokyo (approval number P26-7). Male mice

were used in all experiments. Mice were euthanized by cervical dislocation to preserve

the metabolic environment of the brains and to prevent artifacts that could alter the

biochemical profiles of the proteome. Mouse brains were bisected down the midline to

yield two hemispheres. The brain hemisphere of each animal was quickly frozen on dry

ice and stored at -150ºC until use.

Cell culture

The MG6 cell line is a c-Myc-immortalized cell line of mouse microglia. MG6 cells were

maintained at 5% CO_2 at 37ºC in Dulbecco's modified Eagle's medium supplemented

with 10% (v/v) fetal bovine serum (Thermo Fisher Scientific, 10270-106), 10 µg/mL

insulin (Nacalai Tesque, 12878-44), and 0.1 mM 2-mercaptoethanol [90, 91].

Label-free quantitative proteomic analysis of sarkosyl-insoluble fraction

The fractionation of the brain tissues was prepared as previously described [31]. Tissues

were crushed using Multi-Beads-Shocker (Yasui Kikai, MB601U(S)). Then the tissues

were homogenized with a Potter-type homogenizer (AS ONE CORPORATION) at 800

rpm in 1.4 mL of sarkosyl buffer [PBS, 1% sarkosyl (Nacalai Tesque, 20135-14) and 1

mM phenylmethylsulfonyl fluoride (PMSF) (Wako, 020-15372)]. The homogenates were

centrifuged at $15,000 \times g$ for 5 minutes twice, then for 15 minutes once at 4ºC to obtain

supernatant (S1) and pellet (P1) fractions. The supernatants were centrifuged at 100,000

$\times g$ for one hour at 4ºC to obtain sarkosyl-soluble (S2) and sarkosyl-insoluble (P2)

fractions. The P2 fraction was resuspended in 140 µL of phase transfer surfactant PTS

[12 mM sodium deoxycholate (SDC), 12 mM sarkosyl, and 200 mM ammonium

bicarbonate] buffer. The resuspended P2 solutions were sonicated with Bioruptor® Plus

(BMBio, B01020001) for 10 cycles, (30"/30" cycles), and incubated for 30 minutes at

60ºC with 5 mM dithiothreitol (DTT) for solubilization and reduction of the proteins. The

samples were then alkylated with 200 mM methyl methanethiosulfonate for 30 minutes at

room temperature in the dark. The samples were purified with methanol-chloroform

extraction and solubilized with 50 µL of Rapigest buffer [0.1% Rapigest SF (Waters,

186002123) and 50 mM tetraethylammonium bicarbonate (Thermo Fisher Scientific,

90114)]. The alkylated samples were sonicated and were digested overnight with 1 µg of

Trypsin Gold (Promega, V5280) at 37ºC. Peptides were purified from the supernatant by

GL-tip SDB (GL Sciences, 7820-11200) and eluted with QE-B solution [70%

Acetonitrile and 0.1% formic acid] (Kanto Chemical Co. Inc., 01922-64). Peptides were

concentrated with Concentrator Plus™ for 45ºC and resuspended with 30 µL of QE-A

solution [0.1% formic acid] (Kanto Chemical Co. Inc., 16245-12). The concentration of

peptides was measured with Pierce Quantitative Colorimetric Peptide Assay (Thermo

Fisher Scientific, 23275). 500 ng of peptides from each fraction were subjected to

proteomic analysis using an EASY-nLC 1200 system connected online to Q-Exactive

(Thermo Fisher Scientific). Samples were loaded onto Acclaim PepMap 100 C18 column

(75 µm × 2 cm, 3 µm particle size and 100 Å pore size; 164946, Thermo Fisher

Scientific) and separated using capillary column, (75 µm × 12.5 cm, 3 µm particle

size, NTCC-360/75-3-125, Nikkyo Technos) in an EASY-nLC 1200 system (Thermo

Fisher Scientific). Raw data were analyzed using Proteome Discoverer (version 2.4,

Thermo Scientific). Quantification was performed at the precursor ion level, and data

were normalized using the "Total Peptide Amount" normalization method. The mass

tolerances for the precursor and fragment ions were 10 ppm and 0.02 Da, respectively,

and peptide identification was filtered at a false discovery rate < 0.01. Data were

processed using the R package MSnbase [92]. The missing values were computed using

the method of minDet by replacing values with the minimal value observed in the sample.

Label-free proteomics of the PTS-soluble fraction

Tissues were crushed and homogenized in 400 µL of PTS buffer. The homogenates were

centrifuged at 15,000 × g for 15 minutes at 4ºC to yield the supernatant and pellet

fractions. 30 µg of the supernatant was reduced with 5 mM DTT for 15 minutes at room

temperature. The samples were then alkylated and digested by Trypsin Gold. The

samples were centrifuged at $15,000 \times g$, and the supernatants were added with 0.1%

trifluoroacetate (Wako, 192-09912), incubated for 45 minutes at 37ºC, and centrifuged at

$15,000 \times g$ for 15 minutes at 4ºC. Peptides were purified from the supernatant by GL-tip

SDB (GL Sciences, 7820-11200) and eluted with QE-B solution [70% Acetonitrile and

0.1% formic acid] (Kanto Chemical Co. Inc., 01922-64). The eluates were concentrated

at 45ºC and resuspended with 40 µL of QE-A. 500 ng of peptides were centrifuged at

$15,000 \times g$ for 15 minutes at 4ºC and subjected to proteomic analysis with Q-Exactive.

GO-term analysis

811   In the GO analysis of proteomic data, we used the Database for Annotation, Visualization

812   and Integrated Discovery (DAVID) [93].

Antibodies

Primary antibodies for immunoblotting (IB) and immunohistochemistry (IHC) are listed

in S8 Table. All secondary antibodies used for immunoblotting were purchased from

Jackson ImmunoResearch Laboratories: Peroxidase AffiniPure™ Rabbit Anti-Mouse IgG

+ IgM (H+L) (315-035-048), Peroxidase AffiniPure™ Goat Anti-Rabbit IgG (H+L)

(111-035-144), and Peroxidase AffiniPure™ Donkey Anti-Goat IgG (H+L) (705-035-

147). All the secondary antibodies used for immunohistochemistry were purchased from

ThermoFisher Scientific: Alexa Fluor 488 Goat Anti-Mouse IgG (H+L) (A-28175),

Alexa Fluor 488 Goat Anti-Rabbit IgG (H+L) (A-11008), Alexa Fluor 555 Mouse Anti-

Mouse IgG (H+L) (A-28180), Alexa Fluor 647 Streptavidin (S21374).

Immunoblot analysis

Sample preparation, sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-
PAGE), and immunoblot analysis were performed as described previously [94]. Each
fraction of the brain lysates was mixed with SDS lysis buffer [50 mM Tris-HCl (pH 8.0),
150 mM NaCl, 0.1% SDS, 1% Triton X-100, 0.5% SDC, and 1 mM PMSF] and
centrifuged at $20,000 \times g$ for 20 minutes at 4ºC. The supernatants were separated by
SDS-PAGE (12.5% acrylamide, 7×8 cm, 1 mm thick). Proteins were electrotransferred
to Immobilon®-P polyvinylidene difluoride (PVDF) (Merck Millipore, IPVH00010), at
30 V for one hour at room temperature. For immunoblot analysis, the membranes were
incubated with Blocking One (Nacalai Tesque, 03953-95) for one hour at room
temperature. The membranes were incubated with primary antibodies in TBS-T buffer
[20 mM Tris-HCl (pH8.0), 150 mM NaCl, and 0.1% Tween-20] overnight at 4ºC, washed
with TBS-T buffer and incubated with horseradish-peroxidase conjugated secondary
antibody (1:20,000) for 30 minutes at room temperature. The membranes were immersed
with ImmunoLightning Plus (PerkinElmer, NEL105001EA) or ImmunoSTAR LD
(Wako, 290-69904), and the images were acquired with FUSION FX7.EDGE (VILBER).
For band quantification, the intensity of each band was measured using Fiji software
(ImageJ2, version 2.14.0/1.54h) [95].

Dual membrane filter trap assay

The S1 fractions from young and aged mouse brains were subjected to two-fold serial
dilution using sarkosyl buffer. Fifty microliters of each diluted sample were loaded onto a
pre-wet 0.45-µm pore size nitrocellulose membrane (Bio-Rad, 1620167), which was
layered over a pre-wet Immobilon®-P (PVDF) membrane (0.2-µm pore size) in a Bio-
dot® Microfiltration Apparatus (Bio-Rad, 1706545). Vacuum filtration was performed
under controlled conditions to ensure sequential passage of the protein solution through
the nitrocellulose membrane and then through the PVDF membrane. Each well
underwent three washing cycles with sarkosyl buffer. Subsequently, both membranes
were processed using standard immunoblotting protocols starting from the blocking step
onward.

Immunohistochemical analysis

The paraffin sections (10 µm) were boiled in 10 mM sodium citrate buffer (pH 6.0) for one minute in a microwave oven for heat-induced antigen retrieval. The sections were incubated for one hour in Blocking One Histo (Nacalai Tesque, 06349-64), then overnight at 4ºC with primary antibodies diluted in Blocking One Histo. After rinsing, sections were incubated with secondary antibodies and 1 µg/mL 4',6-diamido-2-phenylindole dihydrochloride (DAPI) (Nacalai Tesque, 11034-56) diluted in Blocking One Histo for one hour at room temperature. The sections were washed by TBS-T three times and by PBS three times, then immersed in TrueBlack® Lipofuscin Autofluorescence Quencher (Biotium Inc., 23007) for 20 minutes to reduce autofluorescence from lipofuscin. The slices were mounted with Prolong™ Glass Antifade Mountant (Invitrogen™, P36980).

For the staining of Iba1, TSA plus amplification kit (Akoya Biosciences, NEL741001KT) was used in combination with subsequent conventional immunofluorescence staining for other antigens. To block endogenous peroxidase activity, brain sections were treated with 3% H_2O_2 in methanol for 15 minutes at room temperature. After washing, sections were treated with Blocking one Histo for one hour at room temperature, and then incubated with 0.1% anti-Iba1 antibody (Wako, 013-27691) in 5% Blocking one Histo in TBS-T overnight at 4ºC. The sections were incubated with 0.01% Peroxidase AffiniPure™ Goat Anti-Rabbit IgG (H+L) (Jackson ImmunoResearch Laboratories, 111-035-144) diluted in TNB buffer at 4ºC overnight. After washing with TNT buffer (0.1% Triton X-100, 100 mM Tris-HCl (pH 7.5), 150 mM NaCl) three times, the sections were reacted with Fluorescein Plus amplification reagent for ten minutes at room temperature, washed with TNT.

For ACAN staining, brain sections were treated with chondroitinase ABC (ChABC, 0.1 U/ml; Sigma, C2905) at 37°C for three hours before blocking. Images were captured by BZ-X800 (Keyence) using 100X objectives or Leica DMi8 Thunder (Leica Microsystems) using 10X or 63X objectives. The images were processed and analyzed

884 with Fiji software. Colocalization analyses were performed with the JaCoP plugin in Fiji

885 software [96].

Protein purification

Recombinant HAPLN2 was purified using the method to purify the versican core protein

G1 domain as previously described [97]. The HAPLN2 coding sequence, nucleotides 82–

1026 was amplified from mouse brain cDNA by PCR and cloned into pGEX6p-1 (27-

4597-01; Addgene). The glutathione-S-transferase tag was removed by PCR. The

construct was transformed into ClearColi® BL21(DE3) Electrocompetent Cells (60810-1,

LGC Biosearch Technologies). The cells were grown in LB Miller broth containing 100

µg/mL ampicillin at 37ºC with shaking at 150 rpm. The *E. coli* cultures were induced

with 0.8 mM isopropyl β-D-thiogalactopyranoside (06289-67; Nacalai Tesque), grown

for another 20 hours at 20ºC, and harvested by centrifugation at $10,000 \times g$ for 20

minutes at 4ºC (Avanti J-25; Beckman Coulter). Inclusion bodies were purified and

solubilized in urea and the protein was refolded, essentially as described previously [97].

Briefly, inclusion bodies were isolated with an *E. coli* lysis buffer [1% (w/v) Nonidet P-

40, 1% (w/v) Triton X-100, 1 mM PMSF, and 1 mM DTT in PBS] solubilized in a

denaturation buffer [8 M urea in 50 mM Tris-HCl (pH 7.5)] and clarified by

centrifugation. The supernatant was purified by incubating with Ni-NTA agarose

(QIAGEN, 30210) overnight and eluted with elution buffer [6 M urea, 500 mM

imidazole, and 10 mM HCl] and then diluted 66-fold into refolding buffer [1 mM

ethylenediaminetetraacetic acid, 500 mM L-arginine, 1 mM cysteine, 2 mM cystine, and

20 mM ethanolamine (pH 11.0)] by dropwise addition at 4ºC with gentle stirring,

followed by incubation overnight without stirring. After refolding, the protein was

concentrated to 2 mL with Amicon Ultra-15 Centrifugal Filter 10 kDa MWCO

(Millipore, 36100101), and dialyzed overnight into a storage buffer [10 mM sodium

phosphate (pH 8.0)].

Protein aggregation assay

Recombinant HAPLN2 proteins (500 nM) were subjected to a series of pH (6.0–8.0), NaCl concentration (0–300 mM), and hyaluronic acid (0–1 mg/mL) in a total volume of 100 µL. The samples were deposited on µ-Slide 18-Well Glass Bottom (Ibidi, 81817) and incubated for 24 hours at 37℃ before being imaged on Eclipse Ti2 microscope (Nikon Corporation) with Micromanager software [98] in which bright field images were taken using a 20X air magnification objective. Aggregate formation was imaged by collecting a series of images in the bright field channel. Fiji software was used in all image processing.

In vitro single-aggregate imaging

Super-resolution imaging was conducted based on point accumulation for imaging nanoscale topography (PAINT). Borosilicate glass coverslips (VWR International, 22 × 22 mm, 631-0124) were cleaned with an argon plasma cleaner (Harrick Plasma, PDC-002) for at least one hour to remove fluorescent residues. Before use, each batch of cover slides was tested for fluorescent artifacts. A drop of 10 µL of the mixture of HAPLN2 used in the protein aggregation assay was placed over a coverslip together with Thioflavin T and Nile Red dyes diluted in PBS to achieve final concentrations of 5 µM for Thioflavin T and 5 nM for Nile Red. Imaging was performed using a CFI Apochromat TIRF 100x Oil immersion objective (Nikon Corporation), mounted on an Eclipse Ti2 microscope (Nikon Corporation). Thioflavin T was excited using a 405 nm laser (Cobolt #0405-06-01-0120-100) with a beam power of 20 mW. Emitted fluorescence was collected by the same objective and separated from the excitation light by a dichroic (Semrock #DI03-R405/488/532/635-T1-25X36), subsequently passed through an emission filter (Semrock #FF01-510/84-25) and focused onto a CMOS camera (ORCA-Flash4.0 V3 Digital CMOS; Hamamatsu). Nile Red was excited by a 532nm laser (Cobolt #0532-06-91-0100-100) with a beam power of 80 mW. Both lasers were positioned at a highly inclined and laminated optical sheet (HiLo) illumination

angle to image large amorphous aggregates at the surface of the coverslip (exciting ~5 µm deep in the sample) while minimizing background fluorescence. Emitted light was separated using a dichroic (DC/ZT532rdc-UF2; Chroma) and passed through an emission filter (Semrock #FF01-650/150-25) before being focused onto the CMOS camera. The image pixel size was 65 nm. A z-stack of 5.0 µm (10 planes with 0.5 µm spacing) was acquired in all channels. The microscope was controlled with Micromanager software, and the frame interval was 50 ms. Each analyzed image corresponds to an average of 10 images in the bright-field channel, 50 images in the thioflavin T, and 1,000 images in the Nile Red channel. The Nile Red images were reconstructed with the ThunderSTORM plug-in in Fiji software [99]. The morphological analysis of Nile Red-positive structures from reconstructed images was conducted by LocAlization Microscopy Analyzer (LAMA) [60] and MATLAB (Mathworks, ver. R2023b).

Circular Dichroism (CD) spectroscopy

CD spectra were recorded using a J-820 CD spectrometer (JASCO) at room temperature. Recombinant HAPLN2 protein (400 ng/µl) was used in its soluble state, and aggregation was induced by adjusting the pH to 6.0 with HCl, followed by overnight incubation at 37°C. The aggregated samples were sonicated for ten minutes prior to measurement. To minimize buffer absorption, the 400 ng/µl samples were diluted 1:8 with deionized water. CD spectra were acquired from 190 nm to 250 nm at a scan rate of 50 nm/min, with a data pitch of 0.1 nm. For each sample, three to four spectra were averaged and smoothed using binomial approximation. Secondary structure content was estimated using BeStSel [62].

Negative-stained transmission electron microscopy

A volume of 1.5 µL of recombinant samples was deposited onto glow-discharged formvar-coated 300-mesh copper grids (Stork Veco) for 20 minutes at room temperature.

The grids were blotted with filter paper (Toyo Roshi Kaisha, Ltd., Tokyo, Japan), washed twice with PBS, washed twice with distilled water, stained with 2% (w/v) uranyl acetate for 30 seconds, blotted and air-dried. The prepared specimens were finally examined with an electron microscope (HT7700; Hitachi) at 100 kV.

Stereotaxic injection

Mice were anesthetized by intraperitoneal injection with 5.0 mL/kg of a non-narcotic anesthetic combination (Me/Mi/Bu) [30 µg/mL medetomidine (Zenoaq), 0.4 mg/mL midazolam (Astellas), and 0.5 mg/mL butorphanol (Meiji Seika Pharma)] and fixed in a stereotaxic surgery frame.

For intracerebellar injection, 20 mg/mL hyaluronidase from ovine testes (FUJIFILM Wako Pure Chemical Corporation, 37326-33-3) in a total volume of 1.0 µL was injected at the following coordinates relative to the bregma: anterior-posterior (A/P) -5.5 mm medial-lateral (M/L) -1.2 mm, and dorsal-ventral (D/V) -2.5 mm. Mice were sacrificed one day post-surgery.

For intrahippocampal injection, either 15 pmol of protein or 5 mg/mL LPS solution, each in a total volume of 2.0 µL, was injected at the following coordinates relative to the bregma: A/P -1.7 mm, M/L -1.6 mm, and D/V -1.9 mm. Mice were sacrificed one week post-surgery.

CSF collection

CSF was collected as described in previous studies [100]. Briefly, mice were anesthetized with a non-narcotic anesthetic combination (Me/Mi/Bu), and the head was secured using a stereotaxic frame (Narishige, SGM-4). The mouse's head was then lowered to create a 45° angle between the cisterna magna and the nose. Under a dissection microscope, the scalps and the three overlapping layers of neck muscle were carefully dissected to expose

the cisterna magna. A borosilicate glass capillary with filament (Narishige, GD-1.2) was

prepared by pulling with a micropipette puller (Narishige, PC100) and connected to a 1

mL syringe via polyethylene tubing (Saint-Gobain, LMT-55). The capillary was mounted

onto a three-dimensional micromanipulator (Narishige, MM3) and gently inserted into

the cisterna magna. Using the micromanipulator, the capillary was inserted into the

cisterna magna and then retracted until CSF began to flow into the capillary. Negative

pressure was applied until approximately 10 µL of CSF was collected, after which the

capillary was fully withdrawn from the cisterna magna. Throughout the experiment, the

body temperature of the mouse was maintained at 37°C using a temperature control

system (Muromachi Kikai Co., Ltd., NS-TC10).

1003

Capillary western blot of HAPLN2 in CSF

1005   Capillary western blot analysis was conducted using Jess (Protein Simple). A volume of 1

1006   µL of cerebrospinal fluid was used for the experiment. Band intensities were calculated

1007   using Fiji software.

1008

Label-free proteomics of the CSF

From 5 µL of CSF, proteins were extracted by methanol-chloroform precipitation and the

resulting pellets were resuspended in 50 µL of PTS-buffer. The samples were reduced

with 5 mM DTT for 15 minutes at room temperature. The samples were then alkylated

and digested by Trypsin Gold. The samples were centrifuged at $15,000 \times g,$ and the

supernatants were added with 0.1% trifluoroacetate (Wako, 192-09912), incubated for 45

minutes at 37ºC, and centrifuged at $15,000 \times g$ for 15 minutes at 4ºC. Peptides were

purified from the supernatant by GL-tip SDB (GL Sciences, 7820-11200) and eluted with

QE-B solution [70% Acetonitrile and 0.1% formic acid] (Kanto Chemical Co. Inc.,

01922-64). The eluates were concentrated at 45ºC and resuspended in 40 µL of QE-A.

500 ng of peptides was centrifuged at 15,000 × *g* for 15 minutes at 4ºC and subjected to
proteomic analysis with Q-Exactive.

1021

Open field test

1023 The behavior of mice was recorded using the Noldus system and analyzed with
1024 EthoVision XT software (Noldus Information Technology). The test was conducted in a
1025 high-walled open-field box (30 × 30 cm), and the animals were observed for five
1026 minutes.

1027

Depletion of microglia by PLX3397 treatment

1029 PLX3397 [101] was synthesized as described previously [102] and formulated at 290
1030 mg/kg in AIN-76A standard chow by Research Diets Inc. The chows were given to 4-
1031 week-old wild-type mice (C57/BL6J) for two months. Depletion of microglia was
1032 confirmed by immunohistochemistry for the microglial marker Iba1.

1033

RNA isolation and quantitative real-time PCR

Total RNA was extracted from MG6 cells grown in 12-well tissue culture plates (7×10^4
cells/well) by High Pure RNA Isolation Kit (Roche, 11828665001). The isolated total
RNA was then reverse-transcribed with ReverTra Ace (TOYOBO, FSQ-201). cDNAs
were amplified with THUNDERBIRD Probe qPCR mix (TOYOBO, QPS-101) and
Universal Probe Library probe (Roche) with the following specific primer sets. mRNA
levels were measured with LightCycler® 480 (Roche).

Gene	Forward (5'–3')	Reverse (5'–3')
Il6	TGTGAAATCAGGATGCTCTGG	CACTTTATTGGGCTCTATACAATATGC

| *Tnf* | TCTTCTCATTCCTGCTTGTGG | GGTCTGGGCCATAGAACTGA |
| *Actb* | ACCAGAGGCATACAGGGACA | CTAAGGCCAACCGTGAAAAG |

1041

Statistics and reproducibility

1043 All data were derived from at least three independent experiments and are presented as

1044 mean ± S.D. or ± S.E.M. as stated in the figure legends. Statistical analyses were

1045 performed with GraphPad Prism 9.0 software. An unpaired two-tailed Student's *t*-test or

1046 one-way ANOVA followed by Tukey's *post hoc* test was performed as described in the

1047 figure legends. All replication attempts were successful with similar results.

1048

Acknowledgments

We thank members of the Laboratory of Morphology and Image Analysis, Biomedical

Research Core Facilities, and Juntendo University Graduate School of Medicine for

technical assistance with microscopy. ChABC was generously provided by Dr. Shinji

Miyata (Tokyo University of Agriculture and Technology). We are grateful to Mr.

Hayato Etani for instruction in stereotaxic injection. Figs 6A and S5E were created by

BioRender (https://www.biorender.com/)

1056

Author Contributions

A.W. did most of the experimental work *in vivo* and *in vitro*, designed the experiments,

analyzed data, and wrote the paper. S.H. conceived the project, designed the experiments,

performed fractionation experiments, MS analysis, stereotaxic injection, and

immunohistochemical analysis of HAPLN2, analyzed data, and wrote the paper. K.I.

performed fractionation experiments, stereotaxic injection, and immunohistochemical

analysis of HAPLN2, and analyzed data. S. Marchese, P. E., and V.M. provided technical

support for HiLo imaging and analyzed data. T.K. collected mouse CSF. H.K. performed

EM analysis, analyzed data, and wrote the paper. S.T. provided technical support for stereotaxic injection, discussed the results, and analyzed data. M.K. provided technical support for EM analysis, designed the experiments, and analyzed data. T.S. and M.T. provided post-mortem human brains. J.A.V. provided technical support for HiLo imaging, analyzed data, discussed the results, and wrote the paper. T.T. provided technical support for stereotaxic injection and analyzed data. S. Murata conceived the project, designed the experiments, analyzed data, and wrote the paper. All authors reviewed the final version and agreed to its publication.

References

[1]   Chung CG, Lee H, Lee SB. Mechanisms of protein toxicity in neurodegenerative diseases. *Cellular and Molecular Life Sciences* 2018; 75: 3159–3180.

[2]   Chou A-H, Yeh T-H, Ouyang P, et al. Polyglutamine-expanded ataxin-3 causes cerebellar dysfunction of SCA3 transgenic mice by inducing transcriptional dysregulation. *Neurobiol Dis* 2008; 31: 89–101.

[3]   Lustbader JW, Cirilli M, Lin C, et al. ABAD Directly Links Aβ to Mitochondrial Toxicity in Alzheimer's Disease. *Science (1979)* 2004; 304: 448–452.

[4]   Kastelic M, Kalyuzhnyi Y V., Hribar-Lee B, et al. Protein aggregation in salt solutions. *Proceedings of the National Academy of Sciences* 2015; 112: 6766–6770.

[5]   Cui Q, Liu Z, Bai G. Friend or foe: The role of stress granule in neurodegenerative disease. *Neuron* 2024; 112: 2464–2485.

[6]   Bachiller S, Jiménez-Ferrer I, Paulus A, et al. Microglia in Neurological Diseases: A Road Map to Brain-Disease Dependent-Inflammatory Response. *Front Cell Neurosci* 2018; 12: 488.

[7]   Kelmer Sacramento E, Kirkpatrick JM, Mazzetto M, et al. Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss and aggregation. *Mol Syst Biol* 2020; 16: e9596.

[8]   Shibata M, Lu T, Furuya T, et al. Regulation of intracellular accumulation of mutant huntingtin by beclin 1. *Journal of Biological Chemistry* 2006; 281: 14474–14485.

[9]   Lipinski MM, Zheng B, Lu T, et al. Genome-wide analysis reveals mechanisms modulating autophagy in normal brain aging and in Alzheimer's disease. *Proceedings of the National Academy of Sciences* 2010; 107: 14164–14169.

[10]   Pan J, Ma N, Yu B, et al. Transcriptomic profiling of microglia and astrocytes throughout aging. *J Neuroinflammation* 2020; 17: 97.

[11]   Pluvinage J V., Haney MS, Smith BAH, et al. CD22 blockade restores homeostatic microglial phagocytosis in ageing brains. *Nature* 2019; 568: 187–192.

[12] Jacquet RG, González Ibáñez F, Picard K, et al. Microglia degrade Alzheimer's amyloid-beta
deposits extracellularly via digestive exophagy. *Cell Rep* 2024; 43: 115052.

[13] Da Mesquita S, Herz J, Wall M, et al. Aging-associated deficit in CCR7 is linked to worsened
glymphatic function, cognition, neuroinflammation, and β-amyloid pathology. *Sci Adv* 2021; 7:
4601–4622.

[14] Vilchez D, Saez I, Dillin A. The role of protein clearance mechanisms in organismal ageing and
age-related diseases. *Nat Commun* 2014; 5: 5659.

[15] Safaiyan S, Kannaiyan N, Snaidero N, et al. Age-related myelin degradation burdens the clearance
function of microglia during aging. *Nat Neurosci* 2016; 19: 995–998.

[16] Crystal H, Dickson D, Fuld P, et al. Clinico-pathologic studies in dementia: nondemented subjects
with pathologically confirmed Alzheimer's disease. *Neurology* 1988; 38: 1682–1687.

[17] Iseki E, Tsunoda S, Suzuki K, et al. Regional quantitative analysis of NFT in brains of
non□demented elderly persons: Comparisons with findings in brains of late□onset Alzheimer's
disease and limbic NFT dementia. *Neuropathology* 2002; 22: 34–39.

[18] Schweighauser M, Arseni D, Bacioglu M, et al. Age-dependent formation of TMEM106B amyloid
filaments in human brains. *Nature* 2022; 605: 310–314.

[19] Lin K, Dorman JB, Rodan A, et al. daf-16□: An HNF-3/forkhead Family Member That Can
Function to Double the Life-Span of Caenorhabditis elegans. *Science (1979)* 1997; 278: 1319–
1322.

[20] Hsu A-L, Murphy CT, Kenyon C. Regulation of Aging and Age-Related Disease by DAF-16 and
Heat-Shock Factor. *Science (1979)* 2003; 300: 1142–1145.

[21] Morley JF, Morimoto RI. Regulation of Longevity in Caenorhabditis elegans by Heat Shock Factor
and Molecular Chaperones. *Mol Biol Cell* 2004; 15: 657–664.

[22] Kenyon C, Chang J, Gensch E, et al. A C. elegans mutant that lives twice as long as wild type.
Nature 1993; 366: 461–464.

[23] David DC, Ollikainen N, Trinidad JC, et al. Widespread Protein Aggregation as an Inherent Part of
Aging in C. elegans. *PLoS Biol* 2010; 8: e1000450.

[24] Walther DM, Kasturi P, Zheng M, et al. Widespread proteome remodeling and aggregation in aging
C. elegans. *Cell* 2015; 161: 919–932.

[25] Fares H, Greenwald I. Genetic Analysis of Endocytosis in Caenorhabditis elegans□: Coelomocyte
Uptake Defective Mutants. *Genetics* 2001; 159: 133–145.

[26] Gallotta I, Sandhu A, Peters M, et al. Extracellular proteostasis prevents aggregation during
pathogenic attack. *Nature* 2020; 584: 410–414.

[27] Diner I, Hales CM, Bishof I, et al. Aggregation Properties of the Small Nuclear Ribonucleoprotein
U1-70K in Alzheimer Disease. *Journal of Biological Chemistry* 2014; 289: 35296–35313.

[28] Hales CM, Dammer EB, Deng Q, et al. Changes in the detergent-insoluble brain proteome linked to
amyloid and tau in Alzheimer's Disease progression. *Proteomics* 2016; 16: 3042–3053.

[29] Guo Q, Dammer EB, Zhou M, et al. Targeted Quantification of Detergent-Insoluble RNA-Binding
Proteins in Human Brain Reveals Stage and Disease Specific Co-aggregation in Alzheimer's
Disease. *Front Mol Neurosci* 2021; 14: 623659.

[30] Woltjer RL, Cimino PJ, Boutté AM, et al. Proteomic determination of widespread detergent
insolubility, including Aβ but not tau, early in the pathogenesis of Alzheimer's disease. *The FASEB*
Journal 2005; 19: 1923–1925.

[31] Diner I, Nguyen T, Seyfried NT. Enrichment of Detergent-insoluble Protein Aggregates from
Human Postmortem Brain. *J Vis Exp* 2017; 2017: 55835.

[32] Gozal YM, Duong DM, Gearing M, et al. Proteomics Analysis Reveals Novel Components in the
Detergent-Insoluble Subproteome in Alzheimer's Disease. *J Proteome Res* 2009; 8: 5069–5079.

[33] Cuanalo-Contreras K, Schulz J, Mukherjee A, et al. Extensive accumulation of misfolded protein
aggregates during natural aging and senescence. *Front Aging Neurosci* 2023; 14: 1090109.

[34] Chang E, Kuret J. Detection and quantification of tau aggregation using a membrane filter assay.
Anal Biochem 2008; 373: 330–336.

[35] Xu G, Gonzales V, Borchelt DR. Rapid Detection of Protein Aggregates in the Brains of Alzheimer
Patients and Transgenic Mouse Models of Amyloidosis. *Alzheimer Dis Assoc Disord* 2002; 16:
191–195.

[36] Baranova NS, Foulcer SJ, Briggs DC, et al. Inter-α-inhibitor impairs TSG-6-induced hyaluronan
cross-linking. *Journal of Biological Chemistry* 2013; 288: 29642–29653.

[37] Mawuenyega KG, Sigurdson W, Ovod V, et al. Decreased Clearance of CNS β-Amyloid in
Alzheimer's Disease. *Science (1979)* 2010; 330: 1774–1774.

[38] Hier DB, Azizi S, Thimgan MS, et al. Tau kinetics in Alzheimer's disease. *Front Aging Neurosci*
2022; 14: 1055170.

[39] Masuda T, Tomita M, Ishihama Y. Phase transfer surfactant-aided trypsin digestion for membrane
proteome analysis. *J Proteome Res* 2008; 7: 731–740.

[40] Wong I, Lohman TM. A double-filter method for nitrocellulose-filter binding: Application to
protein-nucleic acid interactions. *Proc Natl Acad Sci U S A* 1993; 90: 5428–5432.

[41] Laferrière F, Claverol S, Bezard E, et al. Similar neuronal imprint and no cross-seeded fibrils in α-
synuclein aggregates from MSA and Parkinson's disease. *NPJ Parkinsons Dis* 2022; 8: 1–12.

[42] Chang E, Kuret J. Detection and quantification of tau aggregation using a membrane filter assay.
Anal Biochem 2008; 373: 330–336.

[43] Sugitani K, Egorova D, Mizumoto S, et al. Hyaluronan degradation and release of a hyaluronan-
aggrecan complex from perineuronal nets in the aged mouse brain. *Biochim Biophys Acta Gen Subj*
2021; 1865: 129804.

[44] Bekku Y, Vargová L, Goto Y, et al. Bral1: Its Role in Diffusion Barrier Formation and Conduction
Velocity in the CNS. *The Journal of Neuroscience* 2010; 30: 3113–3123.

[45] Aires V, Coulon-Bainier C, Pavlovic A, et al. CD22 Blockage Restores Age-Related Impairments
of Microglia Surveillance Capacity. *Front Immunol* 2021; 12: 43–57.

[46] Ximerakis M, Lipnick SL, Innes BT, et al. Single-cell transcriptomic profiling of the aging mouse
brain. *Nat Neurosci* 2019; 22: 1696–1708.

[47] Elbaz B, Darwish A, Vardy M, et al. The bone transcription factor Osterix controls extracellular
matrix- and node of Ranvier-related gene expression in oligodendrocytes. *Neuron* 2024; 112: 247-
263.e6.

| 1179 | [48] | Grundke-Iqbal I, Iqbal K, Quinlan M, et al. Microtubule-associated protein tau. A component of |
| 1180 | | Alzheimer paired helical filaments. *Journal of Biological Chemistry* 1986; 261: 6084–6089. |

[49] Fawcett JW, Oohashi T, Pizzorusso T. The roles of perineuronal nets and the perinodal
extracellular matrix in neuronal function. *Nat Rev Neurosci* 2019; 20: 451–465.

[50] Hassanzadeh S, Jameie SB, Soleimani M, et al. Coenzyme Q10 Influences on the Levels of TNF-α
and IL-10 and the Ratio of Bax/Bcl2 in a Menopausal Rat Model Following Lumbar Spinal Cord
Injury. *Journal of Molecular Neuroscience* 2018; 65: 255–264.

[51] Thau-Zuchman O, Pallier PN, Savelkoul PJM, et al. High phenylalanine concentrations induce
demyelination and microglial activation in mouse cerebellar organotypic slices. *Front Neurosci*
2022; 16: 926023.

[52] Dohm-Hansen S, English JA, Lavelle A, et al. The 'middle-aging' brain. *Trends Neurosci* 2024; 47:
259–272.

[53] Babinchak WM, Haider R, Dumm BK, et al. The role of liquid–liquid phase separation in
aggregation of the TDP-43 low-complexity domain. *Journal of Biological Chemistry* 2019; 294:
6306–6317.

[54] Qamar S, Wang G, Randle SJ, et al. FUS Phase Separation Is Modulated by a Molecular Chaperone
and Methylation of Arginine Cation-π Interactions. *Cell* 2018; 173: 720-734.e15.

[55] Maharana S, Wang J, Papadopoulos DK, et al. RNA buffers the phase separation behavior of prion-
like RNA binding proteins. *Science (1979)* 2018; 360: 918–921.

[56] Patel A, Lee HO, Jawerth L, et al. A Liquid-to-Solid Phase Transition of the ALS Protein FUS
Accelerated by Disease Mutation. *Cell* 2015; 162: 1066–1077.

[57] Van Lindt J, Bratek-Skicki A, Nguyen PN, et al. A generic approach to study the kinetics of liquid–
liquid phase separation under near-native conditions. *Commun Biol* 2021; 4: 77.

[58] Danial JSH, Klenerman D. Single molecule imaging of protein aggregation in Dementia: Methods,
insights and prospects. *Neurobiol Dis* 2021; 153: 105327.

[59] Lim D, Chu KK, Mertz J. Wide-field fluorescence sectioning with hybrid speckle and uniform-
illumination microscopy. *Opt Lett* 2008; 33: 1819.

[60] Malkusch S, Heilemann M. Extracting quantitative information from single-molecule super-
resolution imaging data with LAMA – LocAlization Microscopy Analyzer. *Sci Rep* 2016; 6: 34486.

[61] Gorantla NV, Das R, Chidambaram H, et al. Basic Limonoid modulates Chaperone-mediated
Proteostasis and dissolve Tau fibrils. *Sci Rep* 2020; 10: 4023.

[62] Kardos J, Nyiri MP, Moussong É, et al. Guide to the structural characterization of protein
aggregates and amyloid fibrils by CD spectroscopy. *Protein Sci* 2025; 34: e70066.

[63] Bondareff W, Narotzky R. Age Changes in the Neuronal Microenvironment. *Science (1979)* 1972;
176: 1135–1136.

[64] Syková E, Voříšek I, Antonova T, et al. Changes in extracellular space size and geometry in APP23
transgenic mice: A model of Alzheimer's disease. *Proceedings of the National Academy of*
Sciences 2005; 102: 479–484.

[65] Reed MJ, Vernon RB, Damodarasamy M, et al. Microvasculature of the Mouse Cerebral Cortex
Exhibits Increased Accumulation and Synthesis of Hyaluronan With Aging. *J Gerontol A Biol Sci*
Med Sci 2017; 72: 740–746.

[66] Cargill R, Kohama SG, Struve J, et al. Astrocytes in aged nonhuman primate brain gray matter
synthesize excess hyaluronan. *Neurobiol Aging* 2012; 33: 830.e13–24.

[67] Reed MJ, Damodarasamy M, Pathan JL, et al. Increased Hyaluronan and TSG-6 in Association
with Neuropathologic Changes of Alzheimer's Disease. *Journal of Alzheimer's Disease* 2019; 67:
91–102.

[68] Su W, Foster SC, Xing R, et al. CD44 Transmembrane Receptor and Hyaluronan Regulate Adult
Hippocampal Neural Stem Cell Quiescence and Differentiation. *Journal of Biological Chemistry*
2017; 292: 4434–4445.

[69] Zakusilo FT, Kerry O'Banion M, Gelbard HA, et al. Matters of size: Roles of hyaluronan in CNS
aging and disease. *Ageing Res Rev* 2021; 72: 101485.

[70] Godin AG, Varela JA, Gao Z, et al. Single-nanotube tracking reveals the nanoscale organization of
the extracellular space in the live brain. *Nature Nanotechnology 2016 12:3* 2016; 12: 238–243.

[71] Radulescu CI, Cerar V, Haslehurst P, et al. The aging mouse brain: cognition, connectivity and
calcium. *Cell Calcium* 2021; 94: 102358.

[72] Wang WY, Tan MS, Yu JT, et al. Role of pro-inflammatory cytokines released from microglia in
Alzheimer's disease. *Annals of Translational Medicine* 2015; 3: 136.

[73] Elmore MRP, Najafi AR, Koike MA, et al. Colony-Stimulating Factor 1 Receptor Signaling Is
Necessary for Microglia Viability, Unmasking a Microglia Progenitor Cell in the Adult Brain.
Neuron 2014; 82: 380–397.

[74] Subhramanyam CS, Wang C, Hu Q, et al. Microglia-mediated neuroinflammation in
neurodegenerative diseases. *Semin Cell Dev Biol* 2019; 94: 112–120.

[75] Mamat U, Woodard RW, Wilke K, et al. Endotoxin-free protein production—ClearColi[TM]
technology. *Nat Methods* 2013; 10: 916–916.

[76] Molzahn C, Kuechler ER, Zemlyankina I, et al. Shift of the insoluble content of the proteome in the
aging mouse brain. *Proc Natl Acad Sci U S A* 2023; 120: e2310057120.

[77] Zolla V, Nizamutdinova IT, Scharf B, et al. Aging□related anatomical and biochemical changes in
lymphatic collectors impair lymph transport, fluid homeostasis, and pathogen clearance. *Aging Cell*
2015; 14: 582–594.

[78] Reed MJ, Vernon RB, Damodarasamy M, et al. Microvasculature of the Mouse Cerebral Cortex
Exhibits Increased Accumulation and Synthesis of Hyaluronan With Aging. *J Gerontol A Biol Sci*
Med Sci 2017; 72: 740–746.

[79] Tan Y-L, Yuan Y, Tian L. Microglial regional heterogeneity and its role in the brain. *Mol*
Psychiatry 2020; 25: 351–367.

[80] Savchenko VL, Nikonenko IR, Skibo GG. Microglial cells of the rat brain in postnatal period
(comparative immunocytochemical analysis). *Neurophysiology* 1997; 29: 145–152.

[81] Verdonk F, Roux P, Flamant P, et al. Phenotypic clustering: a novel method for microglial
morphology analysis. *J Neuroinflammation* 2016; 13: 153.

[82]   Keller D, Erö C, Markram H. Cell Densities in the Mouse Brain: A Systematic Review. *Front*
Neuroanat 2018; 12: 83.

[83]   Xie L, Kang H, Xu Q, et al. Sleep Drives Metabolite Clearance from the Adult Brain. *Science*
(1979) 2013; 342: 373–377.

[84]   Decker Y, Németh E, Schomburg R, et al. Decreased pH in the aging brain and Alzheimer's
disease. *Neurobiol Aging* 2021; 101: 40–49.

[85]   Crapser JD, Arreola MA, Tsourmas KI, et al. Microglia as hackers of the matrix: sculpting
synapses and the extracellular space. *Cell Mol Immunol* 2021; 18: 2472–2488.

[86]   Tansley S, Gu N, Guzmán AU, et al. Microglia-mediated degradation of perineuronal nets
promotes pain. *Science* 2022; 377: 80–86.

[87]   Hickman SE, Kingery ND, Ohsumi TK, et al. The microglial sensome revealed by direct RNA
sequencing. *Nat Neurosci* 2013; 16: 1896–1905.

[88]   Lyssenko NN, Praticò D. ABCA7 and the altered lipidostasis hypothesis of Alzheimer's disease.
Alzheimer's and Dementia 2021; 17: 164–174.

[89]   López-Otín C, Blasco MA, Partridge L, et al. Hallmarks of aging: An expanding universe. *Cell*
2023; 186: 243–278.

[90]   Nakamichi K, Saiki M, Kitani H, et al. Suppressive effect of simvastatin on interferon-β-induced
expression of CC chemokine ligand 5 in microglia. *Neurosci Lett* 2006; 407: 205–210.

[91]   Takenouchi T, Ogihara K, Sato M, et al. Inhibitory effects of U73122 and U73343 on Ca2+ influx
and pore formation induced by the activation of P2X7 nucleotide receptors in mouse microglial cell
line. *Biochim Biophys Acta* 2005; 1726: 177–186.

[92]   Gatto L, Lilley KS. MSnbase-an R/Bioconductor package for isobaric tagged mass spectrometry
data visualization, processing and quantitation. *Bioinformatics* 2012; 28: 288–289.

[93]   Sherman BT, Hao M, Qiu J, et al. DAVID: a web server for functional enrichment analysis and
functional annotation of gene lists (2021 update). *Nucleic Acids Res* 2022; 50: W216–W221.

[94]   Iriki T, Iio H, Yasuda S, et al. Senescent cells form nuclear foci that contain the 26S proteasome.
Cell Rep 2023; 42: 112880.

[95]   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image
analysis. *Nature Methods 2012 9:7* 2012; 9: 676–682.

[96]   Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light microscopy.
J Microsc 2006; 224: 213–32.

[97]   Clark SJ, Keenan TDL, Fielder HL, et al. Mapping the Differential Distribution of
Glycosaminoglycans in the Adult Human Retina, Choroid, and Sclera. *Investigative Opthalmology*
& Visual Science 2011; 52: 6511.

[98]   Edelstein AD, Tsuchida MA, Amodaj N, et al. Advanced methods of microscope control using
μManager software. *J Biol Methods* 2014; 1: e10.

[99]   Ovesný M, Křížek P, Borkovec J, et al. ThunderSTORM: A comprehensive ImageJ plug-in for
PALM and STORM data analysis and super-resolution imaging. *Bioinformatics* 2014; 30: 2389–
2390.

[100] DeMattos RB, Bales KR, Parsadanian M, et al. Plaque-associated disruption of CSF and plasma
amyloid-β (Aβ) equilibrium in a mouse model of Alzheimer's disease. *J Neurochem* 2002; 81:
229–236.

[101] Elmore MRP, Najafi AR, Koike MA, et al. Colony-stimulating factor 1 receptor signaling is
necessary for microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron*
2014; 82: 380–397.

[102] Tap WD, Wainberg ZA, Anthony SP, et al. Structure-Guided Blockade of CSF1R Kinase in
Tenosynovial Giant-Cell Tumor. *New England Journal of Medicine* 2015; 373: 428–437.

[103] Braak H, Alafuzoff I, Arzberger T, et al. Staging of Alzheimer disease-associated neurofibrillary
pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* 2006; 112: 389–
404.

[104] Mirra SS, Heyman A, McKeel D, et al. The Consortium to Establish a Registry for Alzheimer's
Disease (CERAD). *Neurology* 1991; 41: 479–486.

[105] Thal DR, Ru☐b U, Orantes M, et al. Phases of Aβ-deposition in the human brain and its relevance
for the development of AD. *Neurology* 2002; 58: 1791–1800.

[106] McKeith IG, Boeve BF, Dickson DW, et al. Diagnosis and management of dementia with Lewy
bodies. *Neurology* 2017; 89: 88–100.

[107] Braak H, Tredici K Del, Rüb U, et al. Staging of brain pathology related to sporadic Parkinson's
disease. *Neurobiol Aging* 2003; 24: 197–211.

[108] Saito Y, Ruberu NN, Sawabe M, et al. Staging of Argyrophilic Grains: An Age-Associated
Tauopathy. *J Neuropathol Exp Neurol* 2004; 63: 911–918.

[109] Nelson PT, Dickson DW, Trojanowski JQ, et al. Limbic-predominant age-related TDP-43
encephalopathy (LATE): consensus working group report. *Brain* 2019; 142: 1503–1527.

1319

Supporting information

S1 Fig. Protein Extraction by Phase Transfer Surfactant (PTS) Buffer. Coomassie

brilliant blue staining of young (3-month-old) and aged (28-month-old) mouse brains.

Whole brains were lysed with PTS buffer and centrifuged at $15,000 \times g$. n = 3.

1324

S2 Fig. Detailed Immunohistochemistry Analysis of HAPLN2 in the Mouse Brain.

(A) Immunohistochemistry images of the mouse whole brain, cerebellum, and corpus

callosum stained with anti-HAPLN2 antibody and anti-myelin basic protein (MBP)

antibody. n = 3. (B) Magnified immunohistochemistry images of the mouse cerebellar

white matter stained with anti-HAPLN2 antibody or biotinylated hyaluronic acid-binding

1330 protein (HABP), which probes hyaluronic acid. Specific staining of HAPLN2 and HABP

1331 was observed without crossover. n = 3. (C, D) Fluorescence immunohistochemistry

1332 images of the mouse cerebellar white matter showing staining for ubiquitin (C) and p62

1333 (D). n = 3. Scale bars: 1 mm (A), 50 µm (B, C, D), and 10 µm (A (inlet)).

1334

S3 Fig. The Association between Age-Dependent Demyelination and HAPLN2

Aggregation in the Mouse Brain. (A) Immunohistochemistry images of the mouse

cerebellum stained with anti-HAPLN2 antibody, anti-calbindin 1 antibody, and anti-MBP

antibody. n = 3. (B) Magnified fluorescence immunohistochemistry images for HAPLN2,

calbindin 1, and MBP corresponding to the boxed areas in (A). Regions indicated by

magenta arrows showed MBP staining voids accompanied by increased calbindin 1

staining. (C) Quantitation of the mean area of calbindin 1 in (B) comparing young (3-

month-old) and aged (18-month-old) mice. (D) Magnified fluorescent

immunohistochemistry images of calbindin 1–high and –low regions in the cerebellar

white matter of 18-month-old mice. High and low regions were defined as those with

more than 5% and less than 1% calbindin 1–positive area, respectively. (E) Quantitation

of the mean area of calbindin 1 (left) and HAPLN2-positive puncta (right) in calbindin 1–

high region and –low region in 18-month-old mice, as shown in (D). Scale bars: 1 mm

(A), 50 µm (B), and 10 µm (B (inlet), D). Error bars represent mean ± S.D. *P*-values were

calculated using two-tailed Student's *t*-test. **$p < 0.01$, ****$p < 0.0001$, ns = not

significant.

1351

S4 Fig. Age-Dependent Aggregation of HAPLN1 in the Mouse Cerebellar Nuclei.

(A) Immunohistochemistry images of the mouse cerebellum stained with anti-HAPLN1

antibody and anti-ACAN antibody. n = 3. (B) Magnified fluorescence

immunohistochemistry images for HAPLN1 and ACAN corresponding to the boxed

areas in (A). The magenta arrows indicate HAPLN1 that are mislocalized relative to

ACAN staining. (C) Quantitation of the mean area of HAPLN1 (left) and ACAN-positive

area (right) in (B). Scale bars: 1 mm (A), 50 µm (B). Error bars represent mean ± S.D. *P*-

values were calculated using two-tailed Student's *t*-test. ***$p < 0.001$, ns = not

significant.

1361

S5 Fig. Aggregation Propensity of Recombinant HAPLN2 Protein. (A, B) Brightfield

microscopic images showing aggregates of recombinant HAPLN2 incubated under

various buffer conditions in the absence (A) or presence (B) of hyaluronic acid. Data

were collected from at least five fields of view. (C) Histogram depicting the area of

individual aggregates of recombinant HAPLN2 after 24 hours of incubation in the

indicated buffer. Aggregate areas were measured using ImageJ (version 1.53t). (D)

Transmission electron microscopy images of recombinant HAPLN2 aggregates formed

under 150 mM NaCl (pH 6.0). Data were collected from at least five fields of view. (E)

Schematic representation of cross-β-sheet-positive oligomers and fibrils stained with

thioflavin T and Nile Red. Created by BioRender. Scale bars: 10 µm (A, B), 1 µm (D,

black), and 100 nm (D, white).

1373

S6 Fig. Clearance of Cerebellar HAPLN1 after Intracerebellar Hyaluronidase

injection. (A) Immunoblot analysis of the cerebellum corresponding to the samples

shown in Figure 6D. n = 4. (B) Densitometric quantification of (A). HAPLN1 protein

levels in the S2 fraction were normalized to α-tubulin in the S2 fractions. HAPLN1

protein levels in the P2 fraction were normalized to α-tubulin in the S1 fractions. n = 4.

Error bars represent mean ± S.E.M. *P*-values by one-way ANOVA followed by Tukey's

post hoc test. ***$p < 0.001$, ****$p < 0.0001$, ns = not significant.

1381

1382 **S7 Fig. Association between HAPLN2 Accumulation Induced by PLX3397**

1383 **Treatment and Demyelination.The mean area of calbindin 1 did not show a**

1384 **significant difference after PLX3397 treatment.** (A) Immunohistochemistry images of

1385 the mouse cerebellum stained with anti-Iba1 antibody, anti-calbindin 1 antibody, and

1386 anti-MBP antibody. n = 3. (B) Magnified fluorescence immunohistochemistry images for

Iba1, calbindin 1, and MBP corresponding to the boxed areas in (A). (C) Quantitation of the mean area of calbindin 1 (B). Scale bars: 1 mm (A), 50 µm (B), and 10 µm (B, inlet). Error bars represent mean ± S.D. *P*-values were calculated using two-tailed Student's *t*-test. ns = not significant.

S8 Fig. Microglial Activation Induced by Injecting Recombinant HAPLN2 Protein into the Mouse Hippocampus. (A) Filter trap assay and immunoblot analysis of recombinant full-length HAPLN2, HAPLN2 Ig-like domain, and oxStayGold proteins. These recombinant proteins (2 µM each) were incubated overnight at 37℃, followed by centrifugation at $100,000 \times g$ for one hour at 4℃. (B) Immunohistochemistry images of the coronal brain section from Figure 8E, stained with anti-Iba1 antibody to probe activated microglia. (C) Magnified fluorescence immunohistochemistry images for Iba1, corresponding to Figure 8D. Scale bars: 500 µm (B) and 50 µm (C).

S1 Table. Label-free proteomics data of the sarkosyl-insoluble fraction from young and aged mouse brains related to Figure 1A.

S2 Table. Imputed data of label-free proteomics of the sarkosyl-insoluble fraction of young and aged mouse brains corresponding to S1 Table.

S3 Table. 57 proteins enriched in the sarkosyl-insoluble P2 fraction of aged mouse brains in Figure 1A.

S4 Table. Label-free proteomics data of the PTS-soluble fraction from young and aged mouse brains related to Figure 1C.

S5 Table. Imputed data of label-free proteomics of the sarkosyl-insoluble fraction of young and aged mouse brains corresponding to S2 Table.

S6 Table. 27 proteins enriched in the PTS-soluble supernatant of aged mouse brains in Figure 1C.

S7 Table. Human donors applied for the immunohistochemical analysis in Figure 9.

The stages of neurodegenerative diseases in post-mortem brain samples were examined using the following criteria. AT8: Tangle pathology (AT8 Stage) [103], CERAD: The Consortium to Establish a Registry for Alzheimer's Disease, assessing neuritic plaques [104], Thal: Amyloid-beta deposition (Thal Amyloid Phase) [105], DLB (4th Consensus): The Dementia with Lewy Bodies criteria (4th Consensus Report) [106], LB Braak: Lewy body pathology stages (Braak Staging for Lewy Bodies) [107], Grain: Argyrophilic grain pathology (Saito Scale) [108], Joseph-TDP: TDP-43 proteinopathy (Josephs' Staging) [109].

S8 Table. Primary antibodies used in immunoblot and immunohistochemistry analyses.

S1 Data. Raw data and corresponding summary statistics supporting each figure.

S1 Raw Images. Raw images for all IB.

A

B

GO:0005540~hyaluronic acid binding

HAPLN2/VCAN/HAPLN1/ACAN/TNFAIP6

GO:0005539~glycosaminoglycan binding

HAPLN2/VCAN/JCHAIN/HAPLN1/ACAN/IGHM/TNFAIP6

GO:0051015~actin filament binding

BIN1/MYO1D/TPM4/PLS1/FERMT2

GO:0030021~extracellular matrix structural constituent conferring compression resistance

VCAN/HAPLN1

C

D

Protein	Fig 1A		Fig 1C	
	Fold change (aged/young)	*P*-value	Fold change (aged/young)	*P*-value
HTRA1	56.0	2.04E-06	5.01	7.52E-04
HAPLN2	53.2	2.45E-06	2.06	9.48E-04
ACAN	48.8	4.88E-03	16.39	4.88E-02
VCAN	43.9	1.98E-05	3.37	2.30E-04
PLS1	10.5	6.52E-03	2.10	9.41E-04
CRYAB	9.8	2.60E-07	2.36	2.36E-02
HAPLN1	6.5	2.11E-03	2.07	2.03E-05

A

30 kDa — IB: HAPLN2, S1 fraction

30 kDa — IB: HAPLN2, P2 fraction: Sarkosyl insoluble

50 kDa — IB: α-tubulin, S1 fraction

3 months 6 months 12 months 18 months

B

3 months 6 months 12 months 18 months

HAPLN2

HAPLN2 + DAPI

C

A Control PLX3397

DAPI

Iba1

Iba1 + DAPI

B Control PLX3397

DAPI

HAPLN2

HAPLN2 + DAPI

C

Mean area of HAPLN2 (µm²)

****

PLX3397 − +

D

Cerebellum

PLX3397 − +

IB: HAPLN2, P2 fraction: sarkosyl insoluble

IB: HAPLN2, S2 fraction: sarkosyl extractable

IB: α-tubulin, S2 fraction

E

Relative protein levels of HAPLN2 in the P2 fraction

*

PLX3397 − +

A

HAPLN2 + Caspr + DAPI

60–69 years old   80–89 years old

B

Density of HAPLN2 (/mm^2)

*

Age (years)

60–69   80–89

C

Mean area of HAPLN2 (µm^2)

*

Age (years)

60–69   80–89