c-Myc inhibits macrophage antimycobacterial response in *Mycobacterium tuberculosis*

infection

Edoardo Sarti[1,¶], Cédric Dollé[1,¶], Rebekka Wolfensberger[1], Katharina Kusejko[1,2], Doris Russenberger[1], Simon

Bredl[1], Roberto F. Speck[1], Melanie Greter[3], Jan H. Rueschoff[4], Lucas Boeck[5], Dat Mai[6], Alan H. Diercks[6],

Peter Sander[7,8], Gregory S. Olson[5,9], Johannes Nemeth[1*]

[1] Department of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of

Zurich, Zurich, Switzerland

[2] Institute of Medical Virology, University of Zurich, Zurich, Switzerland

[3] Institute of Experimental Immunology, University of Zurich, Zurich, Switzerland

[4] Department of Pathology and Molecular Pathology, University Hospital Zurich, Zurich, Switzerland

[5] Department of Biomedicine, University of Basel, Basel, Switzerland

[6] Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, Washington,

United States of America

[7] Institute of Medical Microbiology, University of Zurich, Zurich, Switzerland

[8] National Reference Laboratory for Mycobacteria, University of Zurich, Zurich, Switzerland

[9] Department of Laboratory Medicine and Pathology, University of Washington, Seattle, Washington, United

States of America

[¶] These authors contributed equally

[*] Corresponding author and lead contact

E-mail: johannes.nemeth@usz.ch

Abstract

Mycobacterium tuberculosis (MTB) is a major global cause of mortality worldwide, responsible for over a million deaths annually. Despite this burden, natural immunity prevents disease in more than 90% of exposed individuals. Previous studies have identified interferon-gamma (IFN-γ) as a key regulator of innate immune defense against MTB. Here, we investigate the impact of IFN-γ timing on macrophage-mediated control of MTB infection. We demonstrate that IFN-γ exposure before infection enhances macrophage antibacterial activity, whereas post-infection exposure does not. Further investigation into this phenotype revealed a strong association between c-Myc signaling and macrophage function in MTB control, as identified using unbiased in vitro systems approaches. Given the challenge of perturbing c-Myc in primary cells, we developed a lentiviral system for c-Myc inhibition and overexpression. Using a tetracycline-inducible Omomyc system - a small peptide inhibitor of c-Myc - we show that c-Myc inhibition promotes a pro-inflammatory macrophage phenotype with enhanced antimycobacterial activity. Mechanistically, c-Myc inhibition induces metabolic reprogramming via increased mTORC1 activity, leading to upregulated inducible nitric oxide synthase and improved bacterial control. In vivo analyses, including murine models and human clinical histopathology, reveal a strong correlation between c-Myc expression and MTB persistence, as well as active tuberculosis (TB), suggesting a role for c-Myc in immune evasion. These findings reveal c-Myc as a potential mediator of immune privilege in MTB infection and highlight its role as a promising target for novel TB therapies aimed at enhancing macrophage function.

Introduction

Mycobacterium tuberculosis (MTB) remains the leading cause of death from a single infectious agent worldwide, responsible for approximately 1.25 million deaths annually, according to the World Health Organization (WHO)[1]. The emergence and spread of antibiotic-resistant strains of MTB demand alternative tools beyond antibiotics to achieve the objectives of global tuberculosis (TB) control efforts.

Although MTB infection is either cleared or effectively controlled by the host immune system in most cases, no host-directed therapies exist for the treatment of active TB[2]. Observations from animal models, human studies, natural infections with HIV-1, and new anti-inflammatory therapies (e.g. anti-TNF therapy) have identified pro-inflammatory macrophages and T cells to be critical components of an effective host immune response[3]. The ensuing immune response may result in profound heterogeneity in the outcomes of MTB infection[4], ranging from deadly disease to complete clearance without any sequelae[2,5,6]. This variability is observed not only across populations but also within individual infected organisms across different species, including mice, non-human primates, and humans[6–8].

To investigate competent host immunity, we have previously focused on asymptomatic MTB infection, the most common, yet underexplored, outcome of MTB infection. Asymptomatic infection appears to confer protection against reinfection - a phenomenon observed in mouse models, non-human primates, and humans[9–11]. In our studies, mice initially received an intradermal inoculation, leading to an asymptomatic primary infection localized within the draining lymph node (contained MTB infection, CMTB). Compared to uninfected controls, mice with CMTB demonstrated a significantly faster and stronger immune response to a secondary, pulmonary challenge. This response included heightened activation of alveolar macrophages (AMs) and rapid recruitment of inflammatory monocytes and T cells, driven by low-grade systemic interferon (IFN)-γ. Consequently, these mice exhibited a substantially lower bacterial burden in the lungs than those without prior infection[9].

IFN-γ is a key cytokine in both innate and adaptive immunity, primarily produced by NK cells, CD4$^+$ T cells, and CD8$^+$ T cells[12]. Under stress or infection, IFN-γ enhances macrophage activation by increasing proinflammatory cytokines and suppressing anti-inflammatory responses, which supports both innate and

adaptive immunity[13,14]. It also boosts macrophage phagocytic and cytolytic abilities, facilitating pathogen

elimination - a process termed classical macrophage activation. Additionally, IFN-γ triggers signaling cascades

that upregulate genes for reactive oxygen and nitrogen species, while also influencing metabolic pathways to

meet the energy demands of activated macrophages in response to external stressors, including

mycobacterial infections[13].

Interestingly, in cases of active TB in humans, there are often high levels of IFN-γ present at the infection

site[15,16]. This counterintuitive observation suggests that, despite the abundance of this potent pro-inflammatory

cytokine, certain tissues may not be responding adequately to the IFN-γ stimulus[4]. This impaired response

could indicate a form of immune resistance or dysfunction[17], where macrophages and other immune cells fail

to fully activate or sustain their antimicrobial functions even in the presence of IFN-γ[2,3,18].

Based on this conundrum, the starting point for the following work was the hypothesis that the timing of IFN-γ

exposure on macrophages relative to MTB infection is crucial for their effective response to MTB infection.

Thus, we hypothesized that the sequence of cytokine exposure and MTB infection could significantly influence

macrophage activation states and, consequently, their ability to control bacterial growth. Therefore, we

decided to investigate how the timing of IFN-γ exposure - whether before or after in vitro infection - affects

macrophage activation, gene expression, and antibacterial activity.

Results

The sequence of infection and activation determines the outcome of *Mycobacterium tuberculosis* infection in macrophages

We hypothesized that the timing of IFN-γ exposure is a critical factor in determining infection outcomes. To test this, we conducted an in vitro experiment using C57BL/6 murine bone marrow-derived macrophages (BMDMs) divided into four groups. To assess bacterial growth, we used a colony-forming unit (CFU) assay, in which macrophages were lysed, and cell lysates were plated on Middlebrook 7H10 agar to quantify viable MTB by counting bacterial colonies.

The experimental groups included: (1) macrophages continuously exposed to IFN-γ throughout the 6-day experiment of infection with the virulent MTB strain H37Rv, including the pretreatment period, (2) macrophages pre-treated with IFN-γ starting 24 hours prior to infection, but thereafter removed, (3) macrophages treated with IFN-γ immediately post-infection (approximately 1 hour after phagocytosis) and maintained for 5 days, and (4) a control group with no IFN-γ treatment (Figure 1A).

Results from CFU assays revealed that only macrophages exposed to IFN-γ prior to infection, with or without continued exposure, showed improved control of MTB growth. In contrast, macrophages treated with IFN-γ post-infection did not exhibit enhanced antimycobacterial activity, as indicated by higher CFU counts (Figure 1B + Supplementary Figure 1).

To assess both functionality and survival of phagocyted MTB, as well as the transcriptional state at the single-cell level, we utilized the MTB live-dead strain[19]. This strain constitutively expresses mCherry, enabling identification of cells containing phagocytosed bacteria, and expresses GFP under the control of a tetracycline-inducible promoter. Bacteria capable of activating GFP expression in response to doxycycline (DOX), a synthetic version derived from tetracycline, are defined as transcriptionally active (mCherry$^+$/GFP$^+$), while those that do not are considered inactive (mCherry$^+$/GFP$^-$) (Figure 1C).

While the live-dead strain has primarily been used in fluorescence microscopy[19–21], we developed a new application of this tool in flow cytometry (Supplementary Figure 2). To validate this approach, we compared

the live-dead strain results with CFU assays using isoniazid-treated MTB H37Rv, finding a comparable

reduction in transcriptional activity and CFU counts (Supplementary Figure 3).

Using flow cytometry and the live-dead strain, we confirmed that macrophages exposed to IFN-γ prior to

infection suppress transcriptional activity in approximately 50% of intracellular MTB. In contrast, macrophages

treated with IFN-γ post-phagocytosis were unable to inhibit MTB transcriptional activity (Figure 1D).

Simultaneously, this assay allowed us to evaluate macrophage activation, with enhanced antimycobacterial

activity correlating with upregulation of MHC-II, CD86 and CD80 surface markers (Figures 1E-1F +

Supplementary Figure 4). These results suggest that the increased bacterial control observed in CFU assays

was due to the heightened activation of viable BMDMs, rather than an artifact from macrophage death or

extracellular bacteria.

c-Myc associated transcriptional programs mirror the in vitro phenotype in mature macrophages

To assess the transcriptome underlying the different infection timepoints and activation states, we conducted

bulk RNA sequencing (RNA-seq) at 6 hours and 24 hours post-infection in the previously described groups of

BMDMs. Principal Component Analysis (PCA) revealed that as early as 6 hours post-infection, macrophages

activated prior to infection began clustering distinctly from those activated after infection (Figure 2A).

Specifically, macrophages activated before infection clustered more closely with fully treated cells (MTB-

controlling cells), while those activated post-infection clustered nearer to untreated cells (MTB-permissive

cells). This distinction became more pronounced at 24 hours, with macrophages activated prior to infection

clustering with fully treated cells, whereas those activated post-infection formed a separate cluster, distinct

from both fully activated and untreated cells.

These findings indicate that as early as 6 hours post-infection, macrophages begin to display trajectories

predictive of the infection outcome. The observation that cells receiving IFN-γ post-infection diverge from

unstimulated cells but do not cluster with cells that received IFN-γ prior to infection suggests that while these

cells can respond to IFN-γ, their trajectory differs significantly from those pre-activated with IFN-γ.

At the single gene level, macrophages treated before infection and macrophages treated after infection have

a clearly different expression signature, with 311 significantly upregulated genes and 396 significantly

downregulated genes (Figure 2B).

We performed Gene Set Enrichment Analysis (GSEA) for functional categorization. Comparisons of gene

expression profiles between macrophages activated prior to infection versus macrophages activated after

infection revealed a relative downregulation of multiple pro-inflammatory pathways in cells that received IFN-

γ after infection (Figure 2C). Notably, however, the most significant changes, in terms of fold changes, were

not among the classical pro-inflammatory genes but were instead related to cell cycle genes, including E2F

targets, G2M targets, and MYC-associated genes, which are upregulated in macrophages treated after

infection (Supplementary Figure 5). Additional comparisons of pre- and post-treated conditions with untreated,

uninfected BMDMs confirmed an activation of IFN-γ–responsive pathways in both treated conditions,

indicating that IFN-γ elicited a strong transcriptional response regardless of timing (Supplementary Figure 6).

Focusing on all significantly up- and downregulated pathways, it is striking that only MYC-associated genes

switch regulation between pre- and post-treatment conditions (Figure 2D).

Among all transcriptional regulators and pathways analyzed, c-Myc was the only transcription factor whose

associated gene expression profile directly mirrored the in vitro antimycobacterial capacity of macrophages.

Specifically, MYC-associated genes were the only gene set that fully recapitulated the phenotype

distinguishing MTB-permissive from MTB-controlling cells, and thus represent a potentially targetable axis.

Macrophages that effectively restricted MTB replication exhibited downregulation of leading-edge genes in the

c-Myc-associated pathway (Figure 2E), suggesting that suppression of MYC activity is linked to enhanced

antimycobacterial function. By contrast, cells stimulated with IFN-γ after infection showed transcriptional

evidence of IFN-γ signaling but did not demonstrate improved mycobacterial control (Figure 1), indicating that

this response alone was insufficient to drive effective restriction of MTB.

These findings suggest that at the transcriptional level, the ability to downregulate c-Myc-associated

transcriptional programs is strongly associated with a macrophage's capacity to mount an effective immune

response that controls in vitro MTB infection.

c-Myc inhibition leads to a gain of antimycobacterial function via a distinct transcriptional program

Based on the strong associations reported above, we decided to experimentally perturb c-Myc in macrophages to better understand its role in mycobacterial infections.

To achieve this, we developed a lentiviral system featuring constitutive expression of a truncated nerve growth factor receptor (NGFR) and tetracycline (Tet)-inducible GFP (Supplementary Figure 7). This system was used to transduce bone marrow cells from C57BL/6 mice. After successful transduction, cells were magnetically sorted and differentiated into BMDMs. When exposed to doxycycline these cells expressed GFP, confirming the functionality of our inducible system (Supplementary Figure 8).

Next, we replaced GFP with murine c-Myc or Omomyc, a small peptide inhibitor of c-Myc[22,23]. RNA-seq analysis then confirmed that, as expected, all transduced cells expressed the truncated NGFR transduction marker (Figure 3A).

To ensure that c-Myc perturbation did not disrupt macrophage function, we assessed phagocytic activity and phagosome acidification in transduced and untransduced BMDMs. We used pHrodo Green *E. coli* bioparticles and mCherry-expressing MTB H37Rv. pHrodo Green is a pH-sensitive fluorescent dye that becomes increasingly fluorescent in acidic environments, allowing selective detection and quantification of active phagocytosis and phagosome acidification[24]. No significant differences were observed between Omomyc-transduced and wildtype macrophages in either the extent of phagocytosis or phagosome acidification (Supplementary Figure 9).

With the system validated, we exposed the transduced BMDMs (Supplementary Figure 10) to doxycycline for 24 hours and performed RNA-seq. The transcriptomic analysis revealed that both c-Myc overexpression and Omomyc-mediated inhibition had profound effects on the cells' transcriptome (Figure 3B). While the significant changes induced by c-Myc overexpression were anticipated - given c-Myc's established role in reprogramming somatic cells into induced pluripotent stem cells[25] - it was particularly striking that Omomyc inhibition also caused extensive transcriptomic alterations.

Subsequent functional enrichment analysis using GSEA indicated that Omomyc-expressing cells exhibited a

distinct transcriptional response (Figure 3C). Notably, these cells showed a pronounced upregulation of

inflammatory responses such as IFN-γ, tumor necrosis factor (TNF)-α or IFN-α response, and a general

increase of metabolic associated pathways, such as an increase of the reactive oxygen species pathway.

These findings suggests that c-Myc plays a critical role in the regulation of gene expression in mature

macrophages in the short term (24 hours), and its up- or downregulation is sufficient to induce changes both

in key inflammatory transcriptional responses and the metabolism in macrophages.

To directly assess how c-Myc inhibition impacts MTB growth control, we infected Omomyc- and c-Myc-

expressing macrophages with MTB H37Rv. Based on our previous data, we hypothesized that c-Myc inhibition

would enhance the macrophages' ability to control MTB growth. Indeed, Omomyc-expressing macrophages

demonstrated that c-Myc inhibition led to a significant gain of antimycobacterial activity, comparable to the

effect observed with IFN-γ pretreatment (Figure 3D). As an additional control, we tested a commercially

available chemical c-Myc inhibitor, 10058-F4, which also led to an increase in antimycobacterial activity,

confirming our findings. At the same time, the construct expressing c-Myc had no significant effect on MTB

growth. In order to exclude possible antibiotic side effects of doxycycline on bacterial growth, we also

compared all tested conditions with and without the addition of doxycycline, thus also eliminating any

overexpression of Omomyc and c-Myc and detected no differences in MTB H37Rv growth depending on the

presence of doxycycline, as was also shown in other studies[26] (Supplementary Figure 11). Finally, the addition

of IFN-γ showed that all conditions responded with similar dynamics in bacterial growth when pre-treated,

further demonstrating that overexpression or inhibition of the key transcription factor c-Myc is tolerated in the

presented system (Supplementary Figure 12).

To understand how these transcriptional changes impacted the macrophages at a cellular level over time, we

infected BMDMs with mCherry-expressing MTB H37Rv (Figure 3E) and monitored key functional markers by

flow cytometry. As predicted by the transcriptional profile, the Omomyc-expressing macrophages displayed

an upregulation of TNF-α on the protein level (Figure 3F). Also, the cells mounted a robust inducible nitric

oxide synthase (iNOS) response (Figure 3G). These findings indicated that the macrophages had gained

enhanced antimycobacterial function, consistent with the prediction based on transcriptional changes.

Importantly, the Omomyc mediated inhibition of c-Myc did not significantly affect macrophage survival over

time compared to control cells as measured with cell viability staining and cell counting, suggesting that the

observed changes were specific to the modulation of c-Myc activity rather than a general cytotoxic effect

(Supplementary Figure 13).

In conclusion, our data demonstrate that c-Myc is a crucial regulator of macrophage function after MTB

infection in vitro. Inhibition of c-Myc via Omomyc induces a unique transcriptional profile that enhances the

antimycobacterial activity of macrophages.

217 mTORC1 mediated metabolic programming partially explains enhanced antimycobacterial function in

218 c-Myc-inhibition in macrophages

We observed that inhibition of c-Myc via the Omomyc peptide led to significant changes in macrophage

functionality, notably an enhancement in antimycobacterial activity. This finding prompted us to investigate the

underlying mechanisms, particularly focusing on metabolism. iNOS, which is essential for the production of

nitric oxide (NO), a key antimicrobial effector[27], requires a substantial energy investment, suggesting a need

for metabolic reprogramming to support this heightened immune response. Considering the extensive

evidence on c-Myc's central role in cellular metabolism, we postulated that c-Myc inhibition might facilitate a

metabolic shift that supports enhanced macrophage function. Our transcription analysis supported this

hypothesis and showed a significant increase in several metabolic pathways in c-Myc-inhibited macrophages,

including mTORC1 signaling (Figure 4A-B). Therefore, we proposed that the gain of function observed through

c-Myc inhibition is driven by metabolic changes mediated by mTORC1.

To investigate this, we conducted in vitro infections using mCherry-expressing MTB H37Rv and measured

iNOS production with or without mTORC1 inhibition via rapamycin. Prior to these experiments, we determined

a concentration of rapamycin that effectively blocked the mTORC1 pathway without impairing macrophage

viability (Supplementary Figure 14). In line with our hypothesis, rapamycin treatment significantly reduced

Omomyc-induced iNOS production (Figure 4C-D), supporting the idea that mTORC1 plays a key role in the

increased iNOS expression seen in c-Myc-inhibited BMDMs. We then infected BMDMs with MTB H37Rv. As

expected, inhibiting mTORC1 reversed in part the enhanced antimycobacterial function caused by c-Myc

inhibition (Figure 4E). This suggests that the improved antimycobacterial activity in c-Myc-inhibited

macrophages is partially due to mTORC1 driven changes in metabolism.

Overall, these findings indicate that c-Myc inhibition leads to metabolic reprogramming including mTORC1,

resulting in enhanced macrophage antimicrobial activity.

c-Myc expression in myeloid cells is associated with MTB persistence in the contained MTB infection

model

Given the central role c-Myc played in in vitro infections of BMDMs, we aimed to determine whether similar

signals could be observed in vivo. To this end, we employed the CMTB model to investigate an in vivo system

of asymptomatic MTB infection. As published previously, mice were intradermally infected with MTB H37Rv

and allowed to develop a chronic infection over a six-week period[9]. After this time, lymph nodes were

harvested, and B and T cells were removed via positive selection using magnetic beads. The remaining cells

were subjected to single-cell RNA sequencing (scRNA-seq) using the 10X Genomics platform, with

subsequent deconvolution of cell populations performed using the Seurat package (Figure 5A +

Supplementary Figure 15) as published previously[28]. We observed a substantial increase of monocytes in

infected lymph nodes (Supplementary Figure 16). Functional enrichment analysis focused on monocytes and

macrophages, utilizing the AddModuleScore function to calculate the average expression levels of GSEA

hallmark c-Myc-related gene programs at the single-cell level (Figure 5B).

Our analysis demonstrated a significant increase in c-Myc transcriptional programs in monocytes and

macrophages during CMTB in mice (Figure 5C). In sum, single-cell transcriptomic analysis of myeloid cells

from the CMTB model revealed elevated expression of c-Myc-associated transcriptional programs in

monocytes and macrophages, linking c-Myc activity to mycobacterial persistence in vivo.

c-Myc is associated with increased lesion size in the standard lung infection model in mice

To examine c-Myc expression in lung tissue, we conducted immunohistochemical staining with anti-c-Myc antibodies on archived lung tissue sections from prior mouse infection studies (Figure 5D). These studies compared MTB reinfection in mice with CMTB, and control mice not pre-exposed to MTB from our previously published experiments[9]. Automated image analysis identified c-Myc positive cells and normalized this count to lesion size. At 3 months post-infection, a significant increase in c-Myc-positive cell density was observed within the lesions. Further analysis revealed a positive correlation between lesion size and the number of c-Myc-positive cells. Additionally, a significant interaction was detected between lesion size and experimental group (control vs. CMTB), suggesting that the relationship between lesion size and c-Myc-positive cell count varies by group (Figure 5E). Specifically, as lesion size increases, the control group showed a greater increase in c-Myc-positive cells than the CMTB group (Figure 5F). This indicates that prior MTB exposure in the CMTB model may reduce the accumulation of c-Myc-positive cells or reduce the expression of c-Myc in cells already present as lesions enlarge, potentially contributing to reduced lesion growth. However, while this interpretation aligns with our data, causality cannot be confirmed by these findings alone. In sum, immunohistochemical analysis of lung tissue revealed that c-Myc expression positively correlates with lesion size during standard MTB lung infection model.

c-Myc expression is spatially enriched in the immune-privileged core of human tuberculosis granulomas

A key limitation of the mouse model is the absence of granulomas, a hallmark of mycobacterial infections in humans. Granulomas with central necrosis are a defining feature of mycobacterial infections in humans and are structured to contain the pathogen[3]. These granulomas generally have a necrotic core surrounded by an outer layer of immune cells, with the inner granuloma known to be an immune-privileged site. We therefore

hypothesized that c-Myc expression may be elevated in the inner granuloma compared to surrounding tissues

in humans.

To test this, we analyzed c-Myc expression in human granulomas by comparing its levels relative to the

necrotic center. Tissue samples were collected from seven patients diagnosed with active tuberculosis during

routine clinical procedures at the University Hospital Zurich (Supplementary Table 1). We used a semi-

automated tissue analysis, in which inner and outer granulomas were manually segmented followed by

automatic nuclei detection and classification of mean c-Myc staining intensity (Figure 6A). We found that the

inner region of the granuloma contained a significantly higher proportion - up to 30% more - of c-Myc-

expressing cells compared to the outer region (Figure 6B). Moreover, the frequency of c-Myc-positive cells

declined with increasing distance from the granuloma core (Figure 6C), and among those positive cells, the

intensity of c-Myc expression also progressively decreased with distance. In summary, c-Myc expression is

spatially associated with the immune-privileged environment of the human granuloma.

Discussion

MTB remains a major cause of mortality worldwide. However, over 90% of exposed individuals do not develop active disease, indicating a significant variability in clinical outcomes likely influenced by host immune factors[29]. Based on prior findings and clinical observations, we hypothesized that the activation state of macrophages, particularly the sequence of activation relative to infection, may play a critical role in determining the outcome of MTB exposure. To test this hypothesis, we employed a controlled in vitro model to examine the effect of IFN-γ activation timing on macrophage responses to infection. Using an unbiased systems approach, we identified c-Myc as a crucial regulator of antimycobacterial activity in vitro. Inhibiting c-Myc induced a proinflammatory macrophage phenotype with enhanced antimycobacterial properties. Mechanistically, c-Myc inhibition shifted macrophage metabolism including increased mTORC1 activation, which improved MTB control.

In line with these findings, inflammation has been linked to suppression of c-Myc associated gene programs[30]. Also, very recent data suggest that overexpression of c-Myc leads to a state of chronic oxidative stress for the macrophage[31]. c-Myc is a crucial regulator in cancer and stem cell biology, known for its role in controlling cell proliferation, growth, and metabolism[32–35]. c-Myc has multiple functions in different cells populations[36–38]. In macrophages, c-Myc has been implicated in the regulation of macrophage polarization. While c-Myc has been associated with the outdated M2 macrophage polarization concept as a potential marker of M2 macrophages[39–41], its specific functions in mature macrophages have not been thoroughly explored.

We tackled the experimental challenge of c-Myc inhibition by employing a tetracycline-inducible Omomyc system, which enabled us to differentiate macrophages into their mature form without disruption and subsequently inhibit c-Myc activity post-maturation. Omomyc, a synthetic dominant-negative regulator of c-Myc, competes with c-Myc to form dimers with Max and binds to c-Myc DNA target sequences (E-boxes) preventing the transactivation of the target genes[22]. This approach has several advantages over chemical inhibitors. It removes the need for solvents and avoids delivery challenges, as the inhibitor is produced directly inside the target cells.

Our data indicate that c-Myc signaling in mature macrophages significantly contributes to macrophage

functionality even in the fully matured cell. Surprisingly, the primary differences between MTB-permissive and

MTB-controlling macrophages occur not only in proinflammatory "effector" pathways but also in canonical cell

cycle pathways. This finding challenges traditional views of c-Myc being key primarily for the development of

cells but positions it also a key regulator of mature macrophage responses[30,42–44] Inhibition of c-Myc does not

immediately compromise cell viability but instead induces a TNF-α, iNOS, and mTORC1-driven transcriptional

program known to be interconnected and to enhance macrophage function during mycobacterial infection[45].

Alveolar macrophages, the cells first to be infected with MTB, and BMDMs represent distinct macrophage

populations that differ in ontogeny, tissue localization, and baseline activation states. AMs are tissue-resident

macrophages originating from fetal precursors and adapted to the lung environment, whereas BMDMs are

derived from circulating monocytes and typically generated ex vivo under defined cytokine conditions[46,47].

Despite these differences, comparative analysis of transcriptional responses across these cell types can

reveal conserved mechanisms of host defense. Complementing our findings in BMDMs, a review of previously

published RNA-seq data from AMs during MTB infection[9] indicated that AMs with increased bactericidal

activity downregulate c-Myc-associated gene programs[28]. Notably, the transcriptional signature of these more

effective AMs closely mirrors the profile observed in MTB-controlling BMDMs, including suppression of c-Myc

target genes and engagement of inflammatory and metabolic pathways. This convergence suggests that c-

Myc suppression may represent a generalizable feature of macrophage-mediated control of MTB, operating

across ontogenically and functionally distinct macrophage subsets both in vitro and in vivo.

We found that c-Myc expression and its related transcriptional profiles are strongly associated with MTB

infection and active tuberculosis in two different mouse models of MTB infection as well as in human samples

from patients with confirmed active TB. Importantly, the spatial pattern of c-Myc - with up to 30% of cells in

the inner granuloma expressing c-Myc during active TB - aligns with the well-established structure of

granulomas, where the core is myeloid cell rich and immune-privileged[3]. Since c-Myc is recognized in cancer

research as a key regulator of immune privilege, this observation appears intuitive, as granulomas are locally

immune-suppressed regions[34]. Additionally, previous studies have identified c-Myc as an essential factor in

the formation of multinucleated giant cells, another feature of mycobacterial infections[48]. While we did not

specifically examine multinucleated giant cells in this study, and instead focused on c-Myc expression per

nucleus rather than per cell, our findings suggest that c-Myc could be a critical mediator of mycobacterial

persistence in the host.

The crucial role of c-Myc in survival and inflammatory response makes it a target for specialized intracellular

pathogens that must prevent pro-inflammatory activation and promote survival of the infected cells in order to

proliferate. Indeed, upregulation of c-Myc has been reported in cells infected with *Chlamydia trachomatis*[49–51],

Salmonella typhimurium[52] and the protozoan pathogen *Leishmania donovani*[53], demonstrating that this

mechanism evolved in different intracellular pathogens. Furthermore, repression of c-Myc signaling alone has

been proved to reduce the growth of *Chlamydia*[50,51] and *Leishmania*[53]. These results suggest that c-Myc may

play a significant role in intracellular infections in general.

Our data indicate that c-Myc inhibition leads to upregulation of several metabolic pathways including mTORC1.

Given that mTORC1 directly and indirectly regulates numerous cellular processes[45] - including iNOS and TNF-

α expression, metabolic reprogramming, autophagy, and trained immunity - it is not surprising that reduced

mTORC1 activity diminishes antimycobacterial responses. This partial reversal of the enhanced effects

observed with c-Myc inhibition suggests that mTORC1 activity is a critical mediator of the functional gain

induced by c-Myc suppression. The metabolic profile resembles the trained immunity phenotype, which

likewise leads to a gain of antimicrobial functions in myeloid cells[54]. Interestingly, we did not observe a

canonical switch towards glycolysis, which is typically considered a hallmark of activated, pro-inflammatory

macrophages with enhanced antimicrobial activity[45,55,56]. Together these findings suggest that metabolic

adaptations may influence the course of MTB infections, emphasizing the importance of metabolic

reprogramming for defense. They also highlight the interaction between c-Myc and mTORC1, which could

enhance the function of macrophages. Furthermore, the presented results provide a potential explanation for

the observation that despite the presence of IFN-γ or IFN-γ-producing cells at the site of infection [15,16], active

disease can occur. Being that higher expression of c-Myc – as seen in granulomas - limits the macrophages

ability to respond properly to the pathogen.

The primary limitation of this study is the absence of in vivo data to demonstrate the effects of c-Myc

perturbation on MTB infection, such as, for example, in a mouse model. Although future studies will address

this, we have observed strong in vivo associations, including the immune privileged core of human

granulomas, where MTB is located[3,8,17].

In conclusion, our data show that c-Myc signaling plays a major role in influencing antimycobacterial responses

in macrophages in vitro. The strong in vivo association in both small animal models and in human tissues from

patients suffering from active TB support further investigation into the interactions between c-Myc and

mycobacterial infections.

Figure 1. Antimycobacterial properties of IFN-γ depend on the sequence of infection and IFN-γ exposure

A) Experimental workflow. BMDMs were isolated from C57BL/6 mice and differentiated for 6 days. Cells were either pre-activated with IFN-γ (25 ng/mL) 24 h before infection or treated immediately after infection. Infection was performed with MTB H37Rv (MOI 5). **B)** Kinetics of intracellular bacterial burden. CFU/mL of intracellular MTB H37Rv were measured immediately after phagocytosis (day 0) and at days 3 and 5 post-infection. Data are mean ± SD of five independent experiments, normalized to day 0. **C)** A representative flow cytometry plot showing viable F4/80$^+$CD11b$^+$ BMDMs infected with MTB H37Rv live-dead strain is shown. Infected (mCherry$^+$) cells are subdivided into "permissive" or "restrictive" populations based on GFP induction (1 µg/mL doxycycline) after 24 h. **D)** Percentage of "permissive" macrophages (GFP$^+$) at days 0, 3 and 5 post-infection. **E–F)** Activation marker expression. Median fluorescence intensity (MFI) of MHC-II (E) and CD86 (F) in infected BMDMs at days 0, 3 and 5, with representative plots (full-treatment condition shown). Unless otherwise stated, data are mean ± SD (n = 3). Statistical analysis: *P* values were determined by unpaired two-tailed Student's t-test (before vs. after infection) and corrected for multiple testing using the two-stage linear step-up method of Benjamini–Krieger–Yekutieli.

Figure 2: Changes in c-Myc associated transcriptional programs mirror MTB permissive versus controlling macrophages in vitro

A) Principal component analysis (PCA) of differently activated/infected BMDMs, 6 and 24 h after infection with MTB H37Rv (MOI 5). Each point represents an individual sample, with colors indicating the treatment. PC1 and PC2 are plotted with the percent variance explained indicated. **B)** Volcano plot of differential expression in BMDMs 24 h post-infection, comparing IFN-γ after vs. IFN-γ before conditions. Genes are plotted by $\log_2$ fold-change (x-axis) against $-\log_{10}$(adjusted P) (y-axis). Red dots mark genes meeting $|\log_2FC| > 2$ and adjusted $P < 0.01$; dashed lines denote these thresholds. **C)** Gene set enrichment analysis (GSEA) dot plot of Hallmark pathways from BMDMs 24 h post-infection, comparing IFN-γ administered before infection versus IFN-γ added after infection. Dots represent normalized enrichment scores (adjusted $P < 0.05$), with positive values indicating enrichment in the before condition. NES values were computed by fgsea on MSigDB Hallmarks (v2024.1.Mm). **D)** Connected-dot plot of GSEA comparing IFN-γ–activated BMDMs before and after relative to untreated, uninfected controls. Each point represents one gene set's NES: orange circles mark sets significantly enriched in the before vs. control comparison (adjusted $P < 0.05$), blue circles mark sets significantly enriched in the after vs. control comparison (adjusted $P < 0.05$). The y-axis shows NES values (positive = up-regulation; negative = down-regulation). Gray dashed lines connect the same gene set's NES across the two contrasts; the solid black line highlights those gene sets that reverse their direction of regulation between pre- and post-infection. **E)** Heatmap of leading-edge genes (top 40 genes) of the MYC_TARGETS_V1 set across all four conditions 24 h after infection. The differential expression is calculated as mean $\log_2$ fold-change per condition compared to untreated, uninfected BMDMs (n = 3). All data shown were derived from bulk RNA sequencing.

Figure 3: c-Myc inhibition leads to a gain in anti-mycobacterial function and induces a pro-inflammatory phenotype in macrophages in vitro

A) RNA expression of rtTa-NGFR (transduction marker) was measured by bulk RNA sequencing as transcripts per million (TPM) in transduced BMDMs after 24 h (n = 3). **B)** Principal component analysis (PCA) of BMDMs expressing either c-Myc or Omomyc versus WT 24 h after Tet-on induction with doxycycline (100ng/mL). Each point represents an individual sample, with colors indicating the experimental group. PC1 and PC2 are plotted with the percent variance explained on the axes. **C)** Gene set enrichment analysis (GSEA) dot plot of Hallmark pathways from BMDMs expressing Omomyc compared to BMDMs expressing c-Myc 24 h after Tet-on induction with doxycycline. Dots represent normalized enrichment scores (adjusted $P < 0.01$), with positive values indicating enrichment in the Omomyc group. NES values were computed by fgsea on MSigDB Hallmarks (v2024.1.Mm). **D)** Kinetics of intracellular bacterial burden in WT BMDMs, BMDMs treated with the chemical c-Myc inhibitor 10058-F4, and BMDMs expressing c-Myc or Omomyc. CFU/mL of intracellular MTB H37Rv were measured immediately after phagocytosis (day 0) and at days 3 and 5 post-infection. Data are mean ± SD of three independent experiments, normalized to day 0. **E)** Representative flow cytometry plots of fully mature BMDMs infected with mCherry-expressing MTB, gated on mCherry[+] cells, showing intracellular TNF-α and iNOS expression (Omomyc-transduced condition is shown). **F–G)** Quantification of the proportion of mCherry[+] BMDMs expressing (F) TNF-α and (G) iNOS at days 0, 3 and 5 post-infection. WT BMDMs, WT pre-treated with IFN-γ, and BMDMs expressing c-Myc or Omomyc were infected and analyzed by flow cytometry. Data are mean ± SD of three independent experiments. Statistical analysis: P values were determined by unpaired two-tailed Student's t-test (Omomyc transduced vs. WT untreated) and corrected for multiple testing using the two-stage linear step-up method of Benjamini–Krieger–Yekutieli.

Figure 4: c-Myc inhibition associated gain of antimycobacterial function is partly mediated by a shift

towards mTORC1 signaling

A) Bar plot of Hallmark metabolic pathways significantly enriched by gene set enrichment analysis (GSEA)

(adjusted *P* < 0.01). Bars show normalized enrichment scores (NES), positive NES (red bars) denote

enrichment in the Omomyc group. **B)** Heatmap of leading-edge genes from the MTORC1_SIGNALING

Hallmark pathway. Rows are genes contributing most to the enrichment score, columns are individual

451 Omomyc-expressing or WT BMDM samples. Values are log$_2$ fold-change relative to the average of WT

452 samples. **C)** Representative flow cytometry plots showing the proportion of iNOS$^+$ cells in mCherry$^+$ BMDMs

453 on day 3 after infection with mCherry-expressing MTB (MOI 5). WT cells, Omomyc-expressing cells and

454 Omomyc-expressing cells treated with 10 µM rapamycin are displayed. All samples received 0.1% DMSO. **D)**

455 Proportion of infected (mCherry$^+$) BMDMs expressing iNOS at days 0, 3 and 5 post-infection in WT, Omomyc-

456 expressing, Omomyc-expressing + rapamycin, and WT + IFN-γ conditions. Data are mean ± SD (n = 3). **E)**

457 Kinetics of intracellular bacterial burden. CFU/mL of MTB H37Rv were measured in WT BMDMs, Omomyc-

458 expressing BMDMs, Omomyc-expressing + rapamycin, and WT + IFN-γ BMDMs immediately after

459 phagocytosis (day 0) and at days 3 and 5 post-infection. Data represent mean relative changes (range) ± SD

460 from three independent experiments, normalized to day 0. Statistical analysis: *P* values were determined by

461 unpaired two-tailed Student's t-test (Omomyc transduced vs. Omomyc transduced + rapamycin) and corrected

462 for multiple testing using the two-stage linear step-up method of Benjamini–Krieger–Yekutieli.

Figure 5: c-Myc expression is associated with MTB infection in the contained MTB infection (CMTB) mouse model of tuberculosis and in the pulmonary infection model

A) Lymph nodes from control and CMTB mice were depleted of CD3[+]/CD20[+] cells and subjected to scRNA-seq. Major clusters were annotated by ImmGen matching, with the monocyte/macrophage cluster circled. **B)** UMAP of the circled monocyte/macrophage cluster showing clear segregation of infected versus non-infected cells. **C)** Single-cell c-Myc gene scores in infected versus control cells, calculated using AddModuleScore (aggregated expression of control gene set subtracted). **D)** Representative immunohistochemistry pathology slides (previously published[9]) from mice with and without CMTB and stained with an anti c-Myc antibody. Lesion size and number of c-Myc-expressing cells were analyzed using a semiautomated pipeline. **F)** Boxplot showing the density of c-Myc-expressing cells measured as number c-Myc positive cells by lesion area (million pixels) in CMTB mice compared to control. **G)** Correlation between number of c-Myc-expressing cells in a lesion and area of the lesion, in CMTB mice and in control mice. Data were analyzed using a two-tailed Student's t-test.

A

B

Figure 6: c-Myc expression is associated with the immune privileged niche in the human granuloma

during active TB

A) Immunohistochemical stain for c-Myc (i-ii) in inner or outer granuloma. Inner and outer granulomas were

manually segmented (iii) followed by automatic nuclei detection and classification of mean c-Myc staining

intensity into negative (blue) or positive: weak (yellow), moderate (orange), or strong (red) (iv). **B)** Boxplot

showing percent of c-Myc -positive nuclei in granuloma regions; inner granulomas had a higher mean

percentage of nuclei positive for c-Myc staining (32.3%) compared to outer granulomas (23.1%). The

horizontal grey line shows the percentage of positive nuclei outside granulomas (17.1%). **C)** Individual nuclei

were grouped by distance to the rim of the nearest inner granuloma into 100 µm bins and the percentages of

nuclei with positive staining by intensity were calculated; negative distances indicate the nucleus is within

the inner granuloma. The total number of nuclei within each bin is represented at the top. Data were

analyzed using a two-tailed Student's t-test.

Materials

Key Resource Table

REAGENT OR RESOURCE	SOURCE	IDENTIFIER

496 Cells

Bone marrow-derived macrophages	*Produced by authors from C57BL/6J mice*	n/a
Conditionally immortalized macrophages	*Produced by authors from C57BL/6J mice*	n/a
HEK 293T	ATCC	CRL-1573

498 Mouse Strains

C57BL/6J wildtype	Charles River Laboratories	C57BL/6J

500 Bacterial Strains

Escherichia coli Stbl2™ competent cells	Invitrogen	10268019
Mycobacterium tuberculosis H37Rv	*Provided by Prof. Alan Aderem*	n/a
Mycobacterium tuberculosis H37Rv – live-dead reporter	*Provided by Prof. Sarah Fortune*	n/a
Mycobacterium tuberculosis H37Rv - mCherry	*Provided by Prof. Alan Aderem*	n/a

502   Antibodies and Fluorescent Dyes

Anti-human CD271 (NGFR)-APC	Biolegend	345108
Anti-human CD271 (NGFR)-PE	Biolegend	345106
Anti-mouse CD16/CD32-TruStain FcX	Biolegend	101320
Anti-mouse CD206-AF700	Biolegend	141734
Anti-mouse CD80-BV421	Biolegend	104725
Anti-mouse CD86-APC	Biolegend	105114
Anti-mouse F4/80-BV785	Biolegend	123141
Anti-mouse F4/80-PerCP-Cy5.5	Biolegend	123127
Anti-mouse iNOS-APC	Thermo Fisher Scientific	17-5920-82
Anti-mouse MHC-II (I-A/I-E)-PE-Cy7	Biolegend	107630
Anti-mouse TNFa-FITC	Biolegend	506304
Anti-mouse TNFa-PE-Dazzle 594	Biolegend	506346
Anti-mouse/human CD11b-BV421	Biolegend	101236
Anti-mouse/human CD11b-BV605	Biolegend	101237
Zombie NIR™ Fixable Viability Kit	Biolegend	423105

504   Chemicals for Bacterial Culture

BBL™ Middlebrook OADC Enrichment	BD Life Science	212351
Difco™ Middlebrook 7H10 Agar	BD Life Science	262710

Difco™ Middlebrook 7H9 Broth	BD Life Science	271310
Glycerol 87%	AppliChem	A3739
Hygromycin B Gold	InvivoGen	HGG-40-01
Kanamycin	Calbiochem	420311
Polysorbate 80	Hänseler	267380-02

506 Chemicals, Peptides, and Recombinant Proteins for Cell Culture

Brefeldin A	Biolegend	420601
c-Myc inhibitor (10058-F4)	Sigma-Aldrich	475956
DMEM	Sigma-Aldrich	D5796
DMSO	Sigma-Aldrich	41640
EDTA	ITW Reagents	A4892
FBS	Thermo Fisher Scientific	26140
HEPES	Sigma-Aldrich	H3537
Isoniazide	Cayman	20378
Lenti-X GoStix Plus	Takara Bio	631280
L-glutamine	Thermo Fisher Scientific	25030081
LPS	Sigma-Aldrich	SL8274
Mouse GM-CSF Recombinant Protein	Biolegend	576304
Mouse IFN-γ Recombinant Protein	Biolegend	575306
Mouse M-CSF Recombinant Protein	Biolegend	576406
Mouse SCF Recombinant Protein	Biolegend	579704

Mouse TPO Recombinant Protein	Biolegend	593304
Opti-MEM™	Thermo Fisher Scientific	11058021
PBS	Sigma-Aldrich	D8537
Penicillin-Streptomycin	Thermo Fisher Scientific	15140148
PFA	Santa Cruz Biotechnology	sc-281692
pHrodo™ Green *E. coli* BioParticles™ Conjugate for Phagocytosis	Thermo Fisher Scientific	P35366
Polyethylenimine	Thermo Fisher Scientific	BMS1003-A
Protamine sulfate	Sigma-Aldrich	P4380
PVA	Sigma-Aldrich	P8136
Rapamycin	Sigma-Aldrich	553210
RPMI-1640	Sigma-Aldrich	R8758
Sodium azide	Sigma-Aldrich	S2002
Sodium pyruvate	Sigma-Aldrich	S8636
StemSpan™ SFEM	STEMCELL Technologies	9600
β-Estradiol	Sigma-Aldrich	E8875
β-Mercaptoethanol	Life Technologies	31350010

508 Commercial Kits and Materials

Cyto-Fast™ Fix/Perm Buffer Set	Biolegend	426803
Direct Lineage Cell Depletion Kit (mouse)	Miltenyi Biotec	130-110-470
MACSelect™ LNGFR MicroBeads	Miltenyi Biotec	130-091-330
QIAshredder	Qiagen	79656
RNase-Free DNase Set	Qiagen	79254
RNeasy Mini Kit	Qiagen	4106

510 DNA Sequences and Plasmids

c-Myc	ENSEMBL	ENSMUSG00000022346
ER-HoxB8	Wang et al., 2006	n/a
Omomyc	Soucek et al., 1998	n/a
P8.9Δ	AddGene	132929
pMD2.G	AddGene	187440
pTRE3G-GSX2	AddGene	96964
pVPX	Pertel et al., 2011	n/a

512 Software and Algorithms

FlowJo™ Software (10.0.8)	FlowJo LLC, Ashland, OR: Becton, Dickinson and Company; 2023
GraphPad Prism (10.3.1)	GraphPad Software, Inc.
R Studio (2025.05.0+496)	Posit Team, RStudio: Integrated Development Environment for R. Posit Software, PBC, Boston, MA http://www.posit.co/

Experimental models

Mycobacterium tuberculosis **strains**

MTB strain H37Rv and H37Rv mCherry fluorescent MTB strains were kindly provided by Prof. Alan Aderem. MTB H37Rv live-dead reporter strain was provided by Prof. Sarah Fortune. All strains were cultured in Difco Middlebrook 7H9 supplemented with 10%(v/v) BBL Middlebrook OADC and polysorbate 80 in roller bottles or Difco Middlebrook 7H10 Agar supplemented with 10%(v/v) BBL Middlebrook OADC and 0.5% (v/v) Glycerol 87%. Kanamycin (30 µg/mL) and hygromycin B (50 µg/mL) were added to the MTB H37Rv mCherry strain and the MTB H37Rv live-dead strain respectively. Bacteria were quantified by spectrophotometry (OD 600nm). To assess transcriptional activity of the MTB H37Rv live-dead strain[20] (after macrophage infection), doxycycline (1 µg/mL) was added to the culture medium for 24 hours.

Mouse strains

C57BL/6J female mice were bought from Charles River Laboratories and were used as donor for bone marrow macrophages production. All procedures were approved by the cantonal animal experimentation authorities and followed international guidelines for laboratory animal experimentation (Canton of Zurich, licenses ZH138/19 and ZH041/21). All mice were maintained on a 12-hour light-dark cycle under clean, conventional conditions and with free access to water and food.

Methods

Cell culture and cell isolation from bone marrow

HEK 293T cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2mM L-glutamine, and 1% Penicillin-Streptomycin (10'000U/mL; P/S). Bone marrow cells were harvested as described by Amend et al.[57]. To induce differentiation into BMDMs, bone marrow cells were cultured in BMDM differentiation medium, consisting of Roswell Park Memorial Institute (RPMI) 1640 medium supplemented with 10% FBS, 2mM L-glutamine, 1% P/S, and 50 ng/mL macrophage colony-stimulating factor (M-CSF), at a density of 2×10^6 cells/mL. The medium was refreshed on day 3. On day 6, non-adherent

suspension cells were discarded, and adherent cells were detached (with cold PBS + 5 mM EDTA by pipetting)

for downstream experiments. To isolate hematopoietic stem cells (HSCs) and immune progenitors from the

bone marrow, negative magnetic selection was performed using the Direct Lineage Cell Depletion Kit following

the manufacturer's protocol. Bone marrow hematopoietic progenitors were grown in 6-well plates with HSC

medium, consisting of StemSpan SFEM medium supplemented with 1 mg/mL polyvinyl alcohol (PVA), 50

ng/mL thrombopoietin (TPO), and 100 ng/mL stem cell factor (SCF), adapted from Wilkinson et al[58]. For one

proof of concept experiment, lineage depleted cells were immortalized using an estrogen regulated HoxB8

(ER-HoxB8) lentiviral vector as described by Wang et al.[59]. Those cells can be kept in a monocyte precursor

stage for over 850 generations in immortalization medium consisting of RPMI 1640 medium containing 10%

FBS, 2mM L-glutamine, 1% P/S, 10 mM HEPES, 50 µM β-mercaptoethanol, 1 mM sodium pyruvate, 2 µM β-

estradiol, and 20 ng/mL granulocyte-macrophage colony-stimulating factor (GM-CSF). Differentiation into

mature macrophages occurs within 4 to 6 days in complete differentiation medium.

Cloning of the Tet-on system into lentiviral vectors

rtTA V16, an improved version of the reverse tetracycline transactivator used in Tet-on systems was

developed by Das et al.[60]. TRE3G inducible promoter was obtained from plasmid pTRE3G-GSX2. Omomyc

peptide sequence was obtained from Soucek et al.[61], and inserted in the construct. The coding sequence of

the murine version of c-Myc was obtained from the ENSEMBL database.

Production of lentiviral particles

Lentiviruses were generated by transfecting HEK 293T cells with lentiviral vectors in Opti-MEM using

polyethylenimine (PEI) as the transfection reagent. The transfection mix included p8.9Δ (containing the coding

sequences for gag/pol, rev, and tat under the control of the immediate/early promoter enhancer of

cytomegalovirus), pMD2.G (encoding vesicular stomatitis virus glycoprotein), and a VPX-expressing plasmid

(as described by Pertel et al.[62]) in a 1:3 DNA-to-PEI ratio. The medium was replaced 16 hours post-transfection

with RPMI supplemented with 5% FBS. Supernatants containing lentiviral particles were collected and filtered

48 hours after transfection. Lentiviral production was confirmed using the Lenti-X GoStix Plus assay. Viral

particles were subsequently concentrated by centrifugation at 20'000 g for 2 hours, resuspended in PBS, and

stored at −80°C until further use.

Transduction of mouse HSCs and myeloid progenitors

HSCs and immune progenitors were isolated from mice as described above and cultured overnight in HSC

medium (described above). For transduction, the medium was prepared by supplementing the HSC medium

with 10 µg/mL protamine and the appropriate lentiviruses at a multiplicity of infection (MOI) of 10. Cells were

resuspended in the transduction medium at a density of $1×10^6$ cells/mL. After 12 hours of incubation, the cells

were pelleted by centrifugation, washed twice with PBS, and resuspended in BMDM differentiation medium to

promote differentiation into macrophages. After 5-6 days of differentiation, the transduced cells expressing the

nerve growth factor receptor (NGFR) were magnetically enriched using MACSelect LNGFR MicroBeads

according to the manufacturer's instructions. NGFR$^+$ cells were counted and used for downstream

experiments.

Macrophage infection assays

BMDMs were seeded in differentiation medium in 96-well plates, $5×10^4$ cells per well. After 24 hours, the cells

were treated with IFN-γ 25ng/mL according to conditions. On the day of infection, bacteria in log-phase (OD

0.1-0.5) were pelleted by centrifugation and resuspended in RPMI and used to infect BMDMs. BMDMs were

infected at a MOI of 5 bacilli per macrophage for 1 hour at 37°C and washed twice with PBS to remove

extracellular bacteria. The cells were provided fresh BMDM differentiation medium and with or without IFN-γ

according to conditions. For assays without IFN-γ but involving Tet-on constructs (c-Myc- and Omomyc-

expressing BMDMs), the differentiation medium was supplemented with doxycycline (100 ng/mL) 24 hours

prior to infection and maintained throughout the experiment. For rapamycin experiments, cells were pretreated

for 30 minutes in BMDM differentiation medium supplemented with or without 10 µM rapamycin, depending

on the experimental condition. All conditions received the same concentration of DMSO (0.1%). Doxycycline

(final concentration 100 ng/mL) was then added where indicated. Rapamycin treatment was also maintained

throughout the experiment. For all experiments medium was changed latest every second day according to

the conditions. At day 0 (1 hour after infection), 3 and 5 the macrophages were detached from the wells with

cold PBS + 5 mM EDTA by pipetting gently. Half of the suspension with the cells were either used for flow

cytometry staining or lysed with 0.05% SDS in water, and the lysate was plated on 7H10 agar plates in serial

dilutions. After 3 weeks of incubation at 37°C colonies were counted to measure the anti-bacterial activity of

macrophages.

Flow cytometry staining

Staining was performed in FACS buffer (PBS + 2 mM EDTA + 2% FBS + 0.1% sodium azide) containing

CD16/CD32 blocker (FcX), Zombie NIR dye and the fluorescent antibodies (= master mix). Suspension cells

were pelleted by centrifugation at 300 g for 5 minutes and washed once prior to staining. BMDMs were

detached by removing the media, washing with PBS, and incubating the cells for 5-10 minutes in cold PBS +

5 mM EDTA, followed by gentle pipetting to detach the cells. Cells were incubated in the master mix for 20

minutes in the fridge. Cells were washed and fixed in 2–4% paraformaldehyde (PFA) at room temperature for

20 minutes, with particular emphasis on MTB-infected cells. For intracellular staining, the Cyto-Fast Fix/Perm

Buffer Set was used according to the manufacturer's instructions before fixation. When stimulation was

performed, cells were stimulated for 6 hours with 10 mg/mL lipopolysaccharide (LPS) and 1X Brefeldin A

(BFA) before being stained. Acquisition was conducted either at a CytoFLEX S (Beckman Coulter) or LSR

Fortessa (BD Biosciences).

Assay with heat-killed MTB and pHrodo Green

BMDMs differentiated and transduced as described above were incubated with heat-killed MTB at a ratio of

1:2 (BMDM:bacteria) for 1 hour at 37°C. Heat inactivation was performed by incubating mCherry-expressing

MTB H37Rv cultures at 80°C for 20 minutes using a thermoblock. After incubation, the cells were washed

twice with PBS and fresh BMDM differentiation medium was added. After 24 hours, the medium was replaced

with 100 μL of fresh differentiation medium. pHrodo Green *E. coli* bioparticles (pre-aliquoted at 1 mg/mL) were

added at 5 μL per well. Cells were incubated for 1 hour at 37°C in an incubator without CO_2, as CO_2 can alter

the pH of the medium and interfere with pHrodo fluorescence. After incubation, the cells were washed with

PBS and resuspended in FACS buffer. Samples were immediately acquired by flow cytometry without fixation

and pHrodo fluorescence was detected in the GFP channel.

Mouse infection experiments

All mouse experiments regarding the CMTB model have been previously reported in Nemeth *et al.*[9]. For

completeness, we briefly describe the establishment of CMTB using intradermal ear injection and aerosol

infection. Intradermal infections involved administering 10'000 CFU of logarithmic-phase MTB H37Rv in 10 µL

PBS using a 10 µL Hamilton syringe into mice anesthetized with ketamine, following a modified protocol.

Aerosol infections were performed using a diluted frozen stock of kanamycin-resistant MTB H37Rv in an

aerosol infection chamber (Glas-Col). Bacterial load was assessed by plating serial dilutions of homogenized

tissue on kanamycin-containing and antibiotic-free plates to differentiate between the sources of infection.

RNA sequencing

For bulk RNA-seq experiments, 1×10^6 BMDMs that had differentiated for 6 days were seeded in 12-well plates.

Depending on the experimental conditions, addition of IFN-γ or doxycycline, as well as in vitro infections, were

performed as described. For RNA isolation, samples were collected 6 and/or 24 hours post-infection or post-

induction. RNA was isolated using the RNeasy Mini Kit, in combination with QIAshredder homogenization

columns and the RNase-Free DNase Set, according to the manufacturer's instructions. RNA quality was

assessed using a Fragment Analyzer (Agilent), and concentrations were measured using a Qubit (1.0)

Fluorometer (Life Technologies). Only samples with 260/280 ratios between 1.8 and 2.1 and 28S/18S ratios

of 1.5–2 were used for downstream processing.

Library construction, sequencing, and parts of the data analysis were performed by the Functional Genomics

Center Zurich. mRNA libraries were prepared using the TruSeq Stranded mRNA Library Prep Kit (Illumina),

which includes poly-A selection, reverse transcription, fragmentation, end-repair, and adapter ligation.

Fragments with adapters on both ends were enriched via PCR using unique dual indexing (UDI). Library

quality and concentration were validated using a Qubit (1.0) Fluorometer (Life Technologies) and Fragment

Analyzer (Agilent). Libraries were normalized to 10 nM in 10 mM Tris-Cl (pH 8.5) with 0.1% Tween 20.

Sequencing was performed as single-end 100 bp reads on the Illumina NovaSeq 6000 platform.

Raw reads were quality filtered and trimmed using fastp. Transcript quantification was performed by pseudo-

aligning reads to the mouse genome (GRCm38.p6 with GENCODE release M23) using Kallisto. Differential

expression analysis was performed in R using "DESeq2"[63]. PCA, heatmaps, and other visualizations were

generated in RStudio using packages "ComplexHeatmap"[64], "circlize"[65], "EnhancedVolcano", "patchwork",

and "tidyverse"[66]. GSEA was conducted with "fgsea"[67] using the Mouse Hallmark gene set from MSigDB[68–70]

(v2024.1.Mm) and pathway-gene annotations were supplied by "org.Mm.eg.db".

Single cell RNA sequencing

Single cell RNA sequencing was performed by our collaborators in the United States. Lymph nodes were

obtained from mice with and without CMTB (n = 5 per condition). Following isolation, samples underwent

methanol fixation according to protocol "CG000136 Rev. D" from 10X Genomics. Briefly, samples were filtered

using 70 µm filters, and red blood cells were lysed with ACK lysis buffer. Post-lysis, cells were resuspended

in 1 mL of ice-cold PBS with a wide-bore pipette, then transferred to a 1.5 mL low-bind Eppendorf tube. After

centrifugation at 700 g for 5 minutes at 4°C, the supernatant was carefully discarded, and the cell pellet was

washed twice with PBS, counted, and resuspended in 200 µL of ice-cold PBS per 1×10^6 cells. To achieve an

80% final methanol concentration, 800 µL of ice-cold methanol was gradually added to the suspension. The

samples were incubated at -20°C for 30 minutes and stored at -80°C for up to 6 weeks prior to rehydration.

For rehydration, samples were thawed at 4°C, pelleted at 1'000 g for 10 minutes at 4°C, and resuspended in

50 µL of a wash-resuspension buffer (0.04% BSA, 1mM DTT, 0.2U/µL Protector RNAse Inhibitor in 3X SSC

buffer) to achieve a final concentration of approximately 1'000 cells/µL, accounting for an estimated 75% cell

loss.

Libraries were constructed using the Next GEM Single Cell 3' Reagent Kits v3.1 (Dual Index) from 10X

Genomics, following the manufacturer's protocol. Sequencing data were aligned to the mouse genome

(mm10), and unique molecular identifier (UMI) counts were calculated using the Cell Ranger pipeline. Data

processing, integration, and downstream analysis were conducted in Seurat v3, with quality control filtering to

exclude droplets containing fewer than 200 detected genes, more than 4'000 detected genes (to discriminate

against doublets), or more than 5% mitochondrial gene expression. Genes detected in fewer than three cells

across all samples were excluded. Cell-type annotation leveraged the Immgen database as a reference,

implemented with the "SingleR"[71] package. Additionally, pseudotime trajectory analysis was performed using

"SeuratWrappers" and "Monocle3"[72]. Data visualization was facilitated using the "Seurat"[73], "tidyverse"[66],

"cowplot", and "viridis" R packages.

Mouse histology

Immunohistochemistry for c-Myc was performed at the University of Washington's Histology and Imaging

Core. The scanned images were exported as TIFF files and uploaded to ImageJ. Areas of consolidation were

identified on images converted to 8-bit grayscale and thresholded using the "Triangle" method. After signal

noise reduction with a median filter, sequential minimum and maximum filters were used to identify areas of

true consolidation. c-Myc$^+$ cells were identified by color-thresholding to identify brown (DAB$^+$) cells against the

background of blue nuclei and these were automatically counted within each consolidated lesion by using the

analyze particles command. The c-Myc$^+$ cell density within each consolidated lesion was calculated as the

number of c-Myc$^+$ cells per million pixels.

Human histology

Patients with confirmed tuberculosis (culture- and/or PCR-positive) who had a histological specimen available

and had given written consent for their routinely obtained specimens to be used for scientific research were

included after ethical clearance was given by the Zurich Ethics Committee (BASEC no. 2023-00800). The

specimens were stained using routine clinical protocols with hematoxylin and eosin (H&E) and an anti-c-Myc

antibody. The tissues were fixed, photographed and transformed into NDPI files, which were loaded into

QuPath and granulomas (both caseating and non-caseating) were identified on H&E stained slides[74]. Manual

segmentation with QuPath's polygon tool was used to create annotations for inner and outer; the inner region

was subtracted from the outer region to create non-overlapping annotations. To calculate the percentage of

non-epithelial cells outside granulomas that express c-Myc, an automatic threshold was applied to the c-Myc

stained slides based on the average RGB intensity to identify all tissue. Areas of epithelium and the previously

defined granuloma regions were excluded from the background tissue region. Additional annotations were

created to exclude areas of artifact (e.g., folded tissue). Positive cell detection with the following parameters

('{"detectionImageBrightfield":"Opticaldensitysum","requestedPixelSizeMicrons":0.5,"backgroundRadiusMicr

ons":15.0,"backgroundByReconstruction":true,"medianRadiusMicrons":0.0,"sigmaMicrons":1.5,"minAreaMicr

ons":10.0,"maxAreaMicrons":400.0,"threshold":0.12,"maxBackground":2.0,"watershedPostProcess":true,"ex

cludeDAB":false,"cellExpansionMicrons":2.0,"includeNuclei":true,"smoothBoundaries":true,"makeMeasurem

ents":true,"thresholdCompartment":"Nucleus: DAB   OD

mean","thresholdPositive1":0.18,"thresholdPositive2":0.3,"thresholdPositive3":0.5,"singleThreshold":false}')

was used to automatically identify nuclei and classify them into negative, positive (weak), positive (moderate),

or positive (strong) c-Myc staining. Separate quantifications at the annotation (i.e. granuloma region) level and

the detection (i.e. nuclei) level were exported and analyzed using R.

Lead contact and material availability

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Johannes Nemeth (johannes.nemeth@usz.ch). RNA data are available from NCBI Gene Expression Omnibus under accession number GSEXXXXX (tba).

Author contributions

JN conceived the study and co-designed the experiments with ES and CD. ES, CD, RW, DM, and DR performed the experiments. KK, JHR, LB, and GSO analyzed (single-cell) RNA sequencing and histology data and provided critical insights. SB advised on plasmid design and lentivirus production. MG, PS, AHD, and RFS provided critical advice on project progression and data analysis. The manuscript was prepared by CD, JN, and ES and reviewed by all authors.

Acknowledgements

The work was supported by the Swiss National Science Foundation (SNSF) grant 310030_200407 to JN. Work in the laboratory of PS is supported by SNSF grant 310030_197699/1, the Federal Office of Public Health (3632001500), and University of Zurich. The work of collaborators from the United States was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health, under awards U19AI135976 and 75N93019C00070. The authors used OpenAI's ChatGPT to support language editing and improve the clarity of the manuscript. After using this tool, the authors reviewed and edited the content as necessary and take full responsibility for the content of the published article. Transcriptomics was performed at the Functional Genomics Centre Zurich (FGCZ) of University of Zurich and ETH Zurich.

Declaration of interest

JN received honoraria for presentations and advisory boards from Oxford Immunotec, Gilead and ViiV. All other authors have declared that no competing interests exist.

References

1.   WHO. *Global Tuberculosis Report. Geneva, Switzerland: World Health Organization*. (2024).

2.   Flynn, J. L. & Chan, J. Immune cell interactions in tuberculosis. *Cell* **185**, 4682–4702 (2022).

3.   Cohen, S. B., Gern, B. H. & Urdahl, K. B. The Tuberculous Granuloma and Preexisting Immunity. *Annu*
Rev Immunol **40**, 589–614 (2022).

4.   Lin, P. L. *et al.* Sterilization of granulomas is common in active and latent tuberculosis despite within-host
variability in bacterial killing. *Nat Med* **20**, 75–79 (2014).

5.   Cadena, A. M., Fortune, S. M. & Flynn, J. L. Heterogeneity in tuberculosis. *Nat Rev Immunol* **17**, 691–702
(2017).

6.   Lenaerts, A., Barry, C. E. & Dartois, V. Heterogeneity in tuberculosis pathology, microenvironments and
therapeutic responses. *Immunol Rev* **264**, 288–307 (2015).

7.   Plumlee, C. R. *et al.* Ultra-low Dose Aerosol Infection of Mice with Mycobacterium tuberculosis More
Closely Models Human Tuberculosis. *Cell Host Microbe* **29**, 68-82.e5 (2021).

8.   Gideon, H. P. *et al.* Multimodal profiling of lung granulomas in macaques reveals cellular correlates of
tuberculosis control. *Immunity* **55**, 827-846.e10 (2022).

9.   Nemeth, J. *et al.* Contained Mycobacterium tuberculosis infection induces concomitant and heterologous
protection. *PLoS Pathog* **16**, e1008655 (2020).

10.   Cadena, A. M. *et al.* Concurrent infection with Mycobacterium tuberculosis confers robust protection
against secondary infection in macaques. *PLoS Pathog* **14**, e1007305 (2018).

11.   Andrews, J. R. *et al.* Risk of Progression to Active Tuberculosis Following Reinfection With Mycobacterium
tuberculosis. *Clinical Infectious Diseases* **54**, 784–791 (2012).

12.   Ivashkiv, L. B. IFNγ: signalling, epigenetics and roles in immunity, metabolism, disease and cancer
immunotherapy. *Nat Rev Immunol* **18**, 545–558 (2018).

13.   Flynn, J. L. *et al.* An essential role for interferon gamma in resistance to Mycobacterium tuberculosis

infection. *J Exp Med* **178**, 2249–2254 (1993).

14.   Hu, X. & Ivashkiv, L. B. Cross-regulation of Signaling Pathways by Interferon-γ: Implications for Immune

Responses and Autoimmune Diseases. *Immunity* **31**, 539–550 (2009).

15.   Huaman, M. A., Deepe, Jr. , G. S. & Fichtenbaum, C. J. Elevated Circulating Concentrations of Interferon-

Gamma in Latent Tuberculosis Infection. *Pathog Immun* **1**, 291 (2016).

16.   Aggarwal, A. N. *et al.* Unstimulated Pleural Fluid Interferon Gamma for Diagnosis of Tuberculous Pleural

Effusion: a Systematic Review and Meta-analysis. *J Clin Microbiol* **59**, (2021).

17.   McCaffrey, E. F. *et al.* The immunoregulatory landscape of human tuberculosis granulomas. *Nat Immunol*

23, 318–329 (2022).

18.   Ramakrishnan, L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev Immunol* **12**, 352–366

(2012).

19.   Martin, C. J. *et al.* Efferocytosis Is an Innate Antibacterial Mechanism. *Cell Host Microbe* **12**, 289–300

(2012).

20.   Bryson, B. D. *et al.* Heterogeneous GM-CSF signaling in macrophages is associated with control of

Mycobacterium tuberculosis. *Nat Commun* **10**, 2329 (2019).

21.   Irvine, E. B. *et al.* Fc-engineered antibodies promote neutrophil-dependent control of Mycobacterium

tuberculosis. *Nat Microbiol* **9**, 2369–2382 (2024).

22.   Beaulieu, M.-E. *et al.* Intrinsic cell-penetrating activity propels Omomyc from proof of concept to viable

anti-MYC therapy. *Sci Transl Med* **11**, (2019).

23.   Whitfield, J. R., Beaulieu, M.-E. & Soucek, L. Strategies to Inhibit Myc and Their Clinical Applicability. *Front*

Cell Dev Biol **5**, (2017).

24.   Lenzo, J. C., O'Brien-Simpson, N. M., Cecil, J., Holden, J. A. & Reynolds, E. C. Determination of Active

772   Phagocytosis of Unopsonized Porphyromonas gingivalis by Macrophages and Neutrophils Using the pH-

773   Sensitive Fluorescent Dye pHrodo. *Infect Immun* **84**, 1753–1760 (2016).

774   25.   Okita, K., Ichisaka, T. & Yamanaka, S. Generation of germline-competent induced pluripotent stem cells.

775   *Nature* **448**, 313–317 (2007).

776   26.   Gengenbacher, M. *et al.* Tissue Distribution of Doxycycline in Animal Models of Tuberculosis. *Antimicrob*

777   *Agents Chemother* **64**, (2020).

778   27.   Herbst, S., Schaible, U. E. & Schneider, B. E. Interferon Gamma Activated Macrophages Kill Mycobacteria

779   by Nitric Oxide Induced Apoptosis. *PLoS One* **6**, e19105 (2011).

780   28.   Mai, D. *et al.* Exposure to Mycobacterium remodels alveolar macrophages and the early innate response

781   to Mycobacterium tuberculosis infection. *PLoS Pathog* **20**, e1011871 (2024).

782   29.   Chandra, P., Grigsby, S. J. & Philips, J. A. Immune evasion and provocation by Mycobacterium

783   tuberculosis. *Nat Rev Microbiol* **20**, 750–766 (2022).

784   30.   Liu, L. *et al.* Proinflammatory signal suppresses proliferation and shifts macrophage metabolism from Myc-

785   dependent to HIF1 **α** -dependent. *Proceedings of the National Academy of Sciences* **113**, 1564–1569

786   (2016).

787   31.   Yabaji, S. M. *et al.* Lipid Peroxidation and Type I Interferon Coupling Fuels Pathogenic Macrophage

788   Activation Causing Tuberculosis Susceptibility. *bioRxiv* (2024) doi:10.1101/2024.03.05.583602.

789   32.   Pelengaris, S., Khan, M. & Evan, G. c-MYC: more than just a matter of life and death. *Nat Rev Cancer* **2**,

790   764–776 (2002).

791   33.   Dejure, F. R. & Eilers, M. MYC and tumor metabolism: chicken and egg. *EMBO J* **36**, 3409–3420 (2017).

792   34.   Casey, S. C., Baylot, V. & Felsher, D. W. MYC: Master Regulator of Immune Privilege. *Trends Immunol*

793   **38**, 298–305 (2017).

35.   Das, S. K., Lewis, B. A. & Levens, D. MYC: a complex problem. *Trends Cell Biol* **33**, 235–246 (2023).

36.   Jha, R. K., Kouzine, F. & Levens, D. MYC function and regulation in physiological perspective. *Front Cell*
Dev Biol **11**, (2023).

37.   Pelengaris, S. & Khan, M. The many faces of c-MYC. *Arch Biochem Biophys* **416**, 129–136 (2003).

38.   van Riggelen, J., Yetil, A. & Felsher, D. W. MYC as a regulator of ribosome biogenesis and protein
synthesis. *Nat Rev Cancer* **10**, 301–309 (2010).

39.   Pello, O. M. Macrophages and c-Myc cross paths. *Oncoimmunology* **5**, e1151991 (2016).

40.   Pello, O. M. *et al.* Role of c-MYC in alternative activation of human macrophages and tumor-associated
macrophage biology. *Blood* **119**, 411–421 (2012).

41.   Luiz, J. P. M. *et al.* MEK5/ERK5 signaling mediates IL-4-induced M2 macrophage differentiation through
regulation of c-Myc expression. *J Leukoc Biol* **108**, 1215–1223 (2020).

42.   Bowen, H. *et al.* c-Myc Represses and Miz-1 Activates the Murine Natural Resistance-associated Protein 1
Promoter. *Journal of Biological Chemistry* **277**, 34997–35006 (2002).

43.   Wu, J. *et al.* EST12 regulates Myc expression and enhances anti-mycobacterial inflammatory response via
RACK1-JNK-AP1-Myc immune pathway. *Front Immunol* **13**, (2022).

44.   Bae, S. *et al.* MYC-mediated early glycolysis negatively regulates proinflammatory responses by
controlling IRF4 in inflammatory macrophages. *Cell Rep* **35**, 109264 (2021).

45.   Weichhart, T., Hengstschläger, M. & Linke, M. Regulation of innate immune cell function by mTOR. *Nat*
Rev Immunol **15**, 599–614 (2015).

46.   Guan, F. *et al.* Tissue macrophages: origin, heterogenity, biological functions, diseases and therapeutic
targets. *Signal Transduct Target Ther* **10**, 93 (2025).

47.   Toda, G., Yamauchi, T., Kadowaki, T. & Ueki, K. Preparation and culture of bone marrow-derived
macrophages from mice for functional analysis. *STAR Protoc* **2**, 100246 (2021).

48.   Herrtwich, L. *et al.* DNA Damage Signaling Instructs Polyploid Macrophage Fate in Granulomas. *Cell* **167**,
1264-1280.e18 (2016).

49.   Al-Zeer, M. A. *et al.* Chlamydia trachomatis Prevents Apoptosis Via Activation of PDPK1-MYC and

820   Enhanced Mitochondrial Binding of Hexokinase II. *EBioMedicine* **23**, 100–110 (2017).

821   50.   Rajeeve, K. *et al.* Reprogramming of host glutamine metabolism during Chlamydia trachomatis infection

822   and its key role in peptidoglycan synthesis. *Nat Microbiol* **5**, 1390–1402 (2020).

823   51.   Vollmuth, N. *et al.* c-Myc plays a key role in IFN-γ-induced persistence of Chlamydia trachomatis. *Elife* **11**,

824   (2022).

825   52.   Ye, Z., Petrof, E. O., Boone, D., Claud, E. C. & Sun, J. Salmonella Effector AvrA Regulation of Colonic

826   Epithelial Cell Inflammation by Deubiquitination. *Am J Pathol* **171**, 882–892 (2007).

827   53.   Colineau, L., Lambertz, U., Fornes, O., Wasserman, W. W. & Reiner, N. E. c-Myc is a novel Leishmania

828   virulence factor by proxy that targets the host miRNA system and is essential for survival in human

829   macrophages. *Journal of Biological Chemistry* **293**, 12805–12819 (2018).

830   54.   Cheng, S.-C. *et al.* mTOR- and HIF-1α–mediated aerobic glycolysis as metabolic basis for trained

831   immunity. *Science (1979)* **345**, (2014).

832   55.   Braverman, J., Sogi, K. M., Benjamin, D., Nomura, D. K. & Stanley, S. A. HIF-1α Is an Essential Mediator

833   of IFN-γ–Dependent Immunity to *Mycobacterium tuberculosis. The Journal of Immunology* **197**, 1287–

834   1297 (2016).

835   56.   Shapouri-Moghaddam, A. *et al.* Macrophage plasticity, polarization, and function in health and disease. *J

836   Cell Physiol* **233**, 6425–6440 (2018).

837   57.   Amend, S. R., Valkenburg, K. C. & Pienta, K. J. Murine Hind Limb Long Bone Dissection and Bone

838   Marrow Isolation. *Journal of Visualized Experiments* (2016) doi:10.3791/53936.

839   58.   Wilkinson, A. C. *et al.* Long-term ex vivo haematopoietic-stem-cell expansion allows nonconditioned

840   transplantation. *Nature* **571**, 117–121 (2019).

841   59.   Wang, G. G. *et al.* Quantitative production of macrophages or neutrophils ex vivo using conditional Hoxb8.

842   *Nat Methods* **3**, 287–293 (2006).

60.   Das, A. T., Tenenbaum, L. & Berkhout, B. Tet-On Systems For Doxycycline-inducible Gene Expression.

Curr Gene Ther **16**, 156–167 (2016).

61.   Soucek, L. *et al.* Design and properties of a Myc derivative that efficiently homodimerizes. *Oncogene* **17**,

2463–2472 (1998).

62.   Pertel, T., Reinhard, C. & Luban, J. Vpx rescues HIV-1 transduction of dendritic cells from the antiviral

state established by type 1 interferon. *Retrovirology* **8**, 49 (2011).

63.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data

with DESeq2. *Genome Biol* **15**, 550 (2014).

64.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional

genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

65.   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. *circlize* implements and enhances circular visualization in

R. *Bioinformatics* **30**, 2811–2812 (2014).

66.   Wickham, H. *et al.* Welcome to the Tidyverse. *J Open Source Softw* **4**, 1686 (2019).

67.   Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* (2016) doi:10.1101/060012.

68.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting

genome-wide expression profiles. *Proceedings of the National Academy of Sciences* **102**, 15545–15550

(2005).

69.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Syst* **1**, 417–425

(2015).

70.   Castanza, A. S. *et al.* Extending support for mouse data in the Molecular Signatures Database (MSigDB).

Nat Methods **20**, 1619–1620 (2023).

71.   Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic

macrophage. *Nat Immunol* **20**, 163–172 (2019).

72.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496–502

867   (2019).

868   73.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21 (2019).

869   74.   Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci Rep* **7**, 16878

870   (2017).

871   75.   Kim, J. H. *et al.* High Cleavage Efficiency of a 2A Peptide Derived from Porcine Teschovirus-1 in Human

872   Cell Lines, Zebrafish and Mice. *PLoS One* **6**, e18556 (2011).

Supplementary Materials

Supplementary Figure 1. IFN-γ activation before infection improves control of MTB growth in BMDMs

Quantification in colony forming units per milliliter (CFU/mL) of intracellular MTB after IFN-γ activation at

different timepoints. Measurements were taken immediately after phagocytosis (day 0) and at 3 and 5 days

post-infection. Data of one experiment is shown. *P* values were determined by unpaired two-tailed Student's

t-test (before vs. after infection) and corrected for multiple testing using the two-stage linear step-up method

of Benjamini–Krieger–Yekutieli.

Supplementary Figure 2. Gating strategy

Flow cytometry gating strategy of infected BMDMs infected with **A)** mCherry-expressing MTB H37Rv and **B)**

MTB H37Rv live-dead reporter strain.

Supplementary Figure 3. Validation of MTB survival analysis using flow cytometry and CFU assays

To validate the GFP-based flow cytometry assay against the colony forming unit (CFU) assay, we infected 50'000 BMDMs per sample with the MTB H37Rv live-dead strain (MOI 5). Cells were incubated for 24 h, followed by activation with doxycycline (1 μg/mL) for an additional 24 h. Subsequently, one half of the cell samples were lysed for CFU quantification, while the remaining cells were analyzed for GFP⁺ infected cells by flow cytometry. CFU ratios, comparing counts at baseline to those at the specified time point, were plotted to assess bacterial survival and for comparison with the MTB H37Rv live-dead strain. To establish a baseline, BMDMs infected with doxycycline-preactivated live-dead bacteria were immediately subjected to either CFU or flow cytometry analysis post-infection. MTB survival was assessed by exposing infected cells to varying concentrations of the antimycobacterial drug isoniazid (100 ng/mL, 10 ng/mL, and no antibiotic), providing a range of survival percentages for comparative analysis.

Supplementary Figure 4. IFN-γ activation before infection enhances CD80 expression in BMDMs

Median fluorescence intensity (MFI) of CD80 in infected BMDMs was measured by flow cytometry at days 0, 3 and 5 post-infection. Data are shown as mean ± SD (n = 3). *P* values were determined by unpaired two-tailed Student's t-test (before vs. after infection) and corrected for multiple testing using the two-stage linear step-up method of Benjamini–Krieger–Yekutieli.

Supplementary Figure 5. Distinct transcriptional profiles in BMDMs depending on IFN-γ activation

Heatmaps of leading-edge genes (top 40 genes) of downregulated **A)** and upregulated **B)** Hallmark gene sets

across all four conditions 24 h after infection. The differential expression is calculated as mean $\log_2$ fold-

change per condition compared to untreated, uninfected BMDMs (n = 3)

Supplementary Figure 6. Gene set enrichment analysis of IFN-γ-treated BMDMs

Dot plot of Hallmark pathways enriched in BMDMs treated with **A)** IFN-γ administered after infection versus

untreated, uninfected BMDMs and **B)** IFN-γ before after infection versus untreated, uninfected BMDMs 24 h

post-infection. Dots indicate normalized enrichment scores (adjusted $P < 0.05$), with positive values indicating

enrichment in the IFN-γ-treated condition. NES values were computed by fgsea on MSigDB Hallmarks

(v2024.1.Mm).

Supplementary Figure 7. Structure of lentiviral constructs containing the Tet-on system

Structure of the Tet-on constructs expressing different genes: **A)** enhanced green fluorescent protein (EGFP)

B) Omomyc. **C)** murine c-Myc. The Human phosphoglycerate kinase 1 promoter (hPGK) is constitutively active

in most human and mouse cell types. It regulates the transcription of the rtTAV16-P2A-NGFR construct.

rtTAV16 is an improved version of reverse tetracycline-controlled transactivator effective at low doses of

doxycyline (100ng/mL). Truncated human nerve growth factor receptor (NGFR) is the extracellular domain of

the human NGFR, used in this study as marker of transduction. Porcine teschovirus 2A peptide (P2A) is a

self-cleaving peptide that allows the production of two separate proteins from one transcript[75].

Supplementary Figure 8. Doxycycline-inducible GFP expression in transduced BMDMs

Representative flow cytometry plots of BMDMs transduced with lentivirus expressing Tet-on regulated GFP,

with or without doxycycline (100ng/mL). Bone marrow cells were sorted 24 h after transduction based on the

expression of the transduction reporter NGFR. After 6 days, cells were exposed to doxycycline for 24 h and

assessed using flow cytometry.

Supplementary Figure 9. c-Myc perturbation does not alter BMDM phagocytosis or phagosomal acidification

A) Quantification of the proportion of BMDMs that have internalized heat-killed, mCherry-expressing MTB and that exhibit an acidic phagosomal compartment (pHrodo-labelled bioparticles) across all experimental conditions, as assessed by flow cytometry (n = 3). **B)** Proportion of BMDMs with acidic phagosomes (pHrodo$^+$) in each condition, measured by flow cytometry (n = 3). **C)** Representative flow-cytometry plots showing gating for mCherry$^+$ (phagocytosis) and pHrodo$^+$ (acidification) populations.

Supplementary Figure 10. Transduction efficiency of BMDMs across independent experiments

Scatter plot showing the percentage of NGFR⁺ BMDMs, used as reporter for transduction, measured by flow cytometry at each timepoint. BMDMs were lentivirally transduced with either Omomyc or c-Myc constructs. Each data point represents the mean proportion of NGFR⁺ cells from an independent experiment (3 independent experiments).

Supplementary Figure 11. Presence of doxycycline does not affect bacterial growth in CFU assays

 Bar plot of relative changes in CFU at day 3 (normalized to day 0) post-infection in the presence or absence

959 of doxycycline (100 ng/mL). Filled and open circles indicate conditions with expected inhibitor-induced kinetics

960 versus those exhibiting standard CFU kinetics, respectively.

Supplementary Figure 12. Transduced and c-Myc perturbed BMDMs show comparable CFU growth to IFN-γ activation

Quantification in colony forming units per milliliter (CFU/mL) of intracellular MTB at day 0 (immediately after phagocytosis) and days 3 and 5 post-infection in IFN-γ–activated WT, c-Myc transduced, Omomyc transduced, and inhibitor-exposed BMDMs.

Supplementary Figure 13. BMDM viability remains stable by Omomyc-mediated c-Myc inhibition over time

Proportion of infected (mCherry⁺) live cells among infected BMDMs in WT, c-Myc–expressing, Omomyc–expressing, or IFN-γ–treated WT conditions, as assessed by flow cytometry. Live cell percentages were measured immediately after phagocytosis (day 0) and at days 3 and 5 post-infection.

Supplementary Figure 14. Rapamycin does not substantially influence survival in macrophages

A) Proportion of live CD11b[+] F4/80[+] conditionally immortalized macrophages at days 0, 3 and 5 following treatment with increasing concentrations of rapamycin, as assessed by flow cytometry. B) Median fluorescence intensity (MFI) of TNF-α in LPS (10 ng/mL) stimulated for 6 h with Brefeldin A, and unstimulated macrophages, treated with increasing concentrations of rapamycin, measured by flow cytometry.

Supplementary Figure 15. Cluster-defining genes in scRNA-seq

Cluster-defining genes were used to deconvolute scRNA-seq data and generate clusters using the Seurat

package[28].

Supplementary Figure 16. Identification of cell populations by scRNA-seq

Identification of cell populations by scRNA-seq in lymph nodes from mice with contained MTB infection (CMTB) and uninfected controls. Relative abundance of each cell population is shown; note that T and B cells were depleted before sequencing and therefore appear at very low frequency.

994 **Supplementary Table 1. Patient characteristics**

	Sex	Age	Diagnosis	Mode of diagnosis
1	female	38	Retroperitoneal tuberculosis	PCR positive
2	male	21	Lymph node tuberculosis	Culture, PCR positive
3	male	73	Lymph node tuberculosis	PCR positive
4	female	28	Pleural tuberculosis	Culture positive
5	male	42	Lymph node tuberculosis	Culture positive
6	male	22	Lymph node tuberculosis	Culture positive
7	male	58	Laryngeal tuberculosis	Culture, PCR positive