Biomimetic hydrogel platform reveals active force transduction from retinal pigment epithelium to photoreceptors

Authors

Sanna Korpela[1], Maija Kauppila[1], Viivi Karema-Jokinen[1], Lassi Sukki[1], Pasi Kallio[1], Heli Skottman[1], Soile Nymark[1*], Teemu O. Ihalainen[1*]

1 Tampere University, Department of medicine and health technology
* Equal contribution

Abstract

Retinal pigment epithelium (RPE) is an epithelial monolayer located at the back of the eye underneath the neural retina. RPE and retina interact closely, with RPE performing multiple tasks to ensure proper retinal physiology and forming a tight, interlocked structure with photoreceptor outer segments (POS). Essential for retinal welfare, RPE enables the daily renewal of POS by phagocytosing their aged tips. Current RPE *in vitro* cell culture systems lack the contact between RPE and retina, possibly limiting not only RPE maturation but also hindering the studies addressing the RPE-retina interactions. We constructed a retina-mimicking structure from a soft polyacrylamide hydrogel with Matrigel coating and introduced it to RPE cells' apical side during culturing to model the RPE-retina interface *in vitro*. RPE cells adhered to the hydrogels during culture, enabling further studies of the biophysical properties of the RPE-hydrogel interface with methods utilizing rheology and traction force microscopy. During phagocytosis, RPE cells generated considerable traction forces that were followed using a hydrogel coated with POS particles. Disruption of RPE's actin cytoskeleton with Cytochalasin D significantly reduced the force generation. These results support the view of RPE as a mechanically active partner for retina. In addition, our retina-mimicking structure offers a tool for further biophysical modeling of the RPE-retina interface.

Keywords

Retinal pigment epithelium; Hydrogel; Retina; Photoreceptor outer segment phagocytosis; In vitro cell culture; Biomimetic;

Abbreviations

AA	Acrylamide
APS	Ammonium persulfate
BIS	N,N'-Methylenebisacrylamide
BSA	Bovine serum albumin
ECM	Extracellular matrix
EGFP	Enhanced green fluorescent protein
FTTC	Fourier transform traction cytometry
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
hESC	Human embryonic stem cell
hiPSC	Human induced pluripotent stem cell
IPM	Interphotoreceptor matrix
L-DOPA	3,4-dihydroxy-L-phenylalanine
PA	Polyacrylamide
PBS	Phosphate buffered saline
PFA	Paraformaldehyde
PIV	Particle image velocimetry
POS	Photoreceptor outer segment
RPE	Retinal pigment epithelium
TEER	Transepithelial electrical resistance
TFM	Traction force microscopy
VEGF	Vascular endothelial growth factor

1. Introduction

Retinal pigment epithelium (RPE) is an epithelial monolayer located at the back of the eye beneath the retina. Unlike a typical epithelium, RPE cells do not face an empty lumen on their apical side but are structurally and functionally in physical contact with retinal photoreceptor outer segments (POS) (1,2). The apical side membrane protrusions of RPE, microvilli, locate between POS, forming an interlocked structure. This close proximity of the retina enables the RPE to efficiently perform one of its primary roles, retinal maintenance. RPE cells play several crucial roles in supporting retinal health, including formation of the blood-retinal barrier, nutrient and waste transportation between retina and underlying choroid, maintenance of ion homeostasis, growth factor secretion, and regeneration of the visual pigments (1–3). In addition, RPE cells are responsible for the essential daily renewal of POS in a process called phagocytosis (1,2,4). The close interaction between the RPE and photoreceptors is mediated by interphotoreceptor matrix (IPM), a specialized and compartmentalized extracellular matrix (ECM), that participates in retina-RPE adhesion (5–7). In addition, IPM partakes in retinoid, oxygen and nutrient transport, and contains enzymes and growth factors. Unlike typical ECM, IPM lacks the collagen meshwork, and is mainly composed of glycosaminoglycans, proteoglycans, and hyaluronan, which are large, hydrophilic molecules. (7,8) The IPM physically bridges the RPE and POS, connecting the tissue layers together. Furthermore, RPE-retina adhesion is mediated by $\alpha_v\beta_5$ integrins on the apical surface of RPE, intraocular pressure, metabolism, and net fluid transport from the retina (9,10). Even though the details of the interactions' strength and dynamics are not known, enzymatic degradation of IPM (11) and inhibition of IPM synthesis (5) have been linked to retinal detachment, and IPM's role in photoreceptor degenerative diseases has been discussed (7,8,12).

The RPE-retina adhesion needs to be dynamic to allow RPE cells to internalize and digest parts of the POS in the daily phagocytosis process (1,2,4). The POS continuously grow from their base, and older tips of outer segments, which have participated in the phototransduction process for longer, are engulfed and recycled by the RPE. This recycling prevents the accumulation of photo-oxidative compounds to photoreceptor cells and IPM, and is vital for retinal welfare. (13) The internalization of POS particles by the RPE is a highly dynamic process, involving actin cytoskeleton reorganization and formation of phagocytic cups that scission parts of the POS for ingestion (14,15). This 'nibbling' of the outer segments requires the interplay of actin and myosin motors and thus it highly likely leads to tensile or compressive force generation locally at the RPE cell membrane (14,16,17). However, whether forces are needed in the POS phagocytosis process and if the forces are transduced from the RPE to the retina is not known. While live cell imaging methods have recently enabled studying the dynamics of phagocytosis by RPE in more detail (14,18,19), characterization of the RPE generated forces in both RPE-retina adhesion and phagocytosis remains a challenge. Furthermore, current RPE *in vitro* cell culture systems (1,20,21) are missing the RPE-retina interface, limiting studies

addressing the biophysical characteristics of the interaction. Based on the biomechanical properties of the retina (22), we hypothesized that the force transduction and biophysics of the RPE-retina interface could be modelled utilizing a soft hydrogel as a retina-mimicking structure. Bringing this hydrogel into contact with the RPE apical side would allow the RPE cells to form connections and transmit forces to the hydrogel.

Hydrogels are versatile tools for mimicking extracellular environments in cell culture conditions (23). Here, we show that the biomimetic retina hydrogel platform can be constructed from elastic polyacrylamide (PA) hydrogel. PA gels are clear, synthetic, and non-degradable hydrogels consisting of cross-linked acrylamide polymer chain network, and their surface can be functionalized with extracellular matrix proteins to facilitate cell adhesion (24–26). We show that human embryonic stem cell (hESC) and human induced pluripotent stem cell (hiPSC) –derived RPE cells can be cultured with the biomimetic PA hydrogel on the apical side of RPE without changing the morphology or functionality of the cells. Also, as RPE cells adhered on the hydrogels on their apical side, we conducted further studies of the biophysical properties of the RPE-hydrogel interaction by using rheology and traction force microscopy. Our results indicate that the RPE-hydrogel adhesion is reversible and actin -mediated. Moreover, by embedding POS particles to the hydrogel, we were able to show that RPE cells exerted considerable forces to the retina-mimicking hydrogel in a scale of $pN/\mu m^2$ during phagocytosis, which supports the emerging view of RPE being a mechanically active partner for the retina. In addition, our retina-mimicking structure offers an inexpensive, reproducible, and accessible tool for further biophysical modeling of the RPE-retina interface.

2. Results

2.1. A retina-mimicking hydrogel can be constructed from polyacrylamide

To study the RPE-retina interaction *in vitro*, we constructed a soft hydrogel to mechanically mimic the photoreceptors interacting with RPE (Fig 1A). Individual photoreceptor cells have been reported to have an elastic modulus (Young's modulus, E) of 0.2 – 1.0 kPa (27) and the whole photoreceptor layer approximately 26 kPa (28). Moreover, the IPM between RPE and the photoreceptors has a high prevalence of hydrophilic glycoproteins and proteoglycans (7). Based on these properties we prepared a soft deformable PA hydrogel with E = 2.8 kPa which mechanically resembles the photoreceptor cell layer. PA hydrogels are resistant to protein binding and, therefore, non-adhesive also for cells without adhesive protein coating. Thus, we modeled the IPM layer by coating the PA hydrogel with Matrigel, a commercially available solubilized basal membrane, using L-DOPA treatment of the gel (25,29). By combining the soft PA hydrogel and Matrigel coating, we were able to construct a system which mechanically resembles the photoreceptor cell layer and has the important IPM components for cell adhesion. The retina-mimicking hydrogel was then used together with hESC- and hiPSC - derived RPE cells.

Fig 1. RPE can be cultured with retina-mimicking hydrogel without affecting its morphology or functionality.

(A) Left: A brightfield image of a transverse section from mouse outer retina in paraffin, showing the close interaction between RPE cells' apical side and photoreceptor cells. Scalebar 5 μm. Right: A schematic illustration of the RPE-hydrogel interaction. The PA hydrogel is coated with Matrigel and placed on the apical side of hESC- or hiPSC-derived RPE cells cultured in Transwell insert on a semipermeable membrane. During culture, RPE cells adhere to the hydrogel on their apical side. (B) Timeline of the cell culture experiments. hESC- and hiPSC-RPE cells were plated on 24-well inserts at week 0. After 5 weeks, the hydrogel was placed on the RPE apical side, and the cells were

cultured with the hydrogel for two weeks until week 7. Alternatively, the hydrogels were placed on 7-week-old cells and cultured until RPE cells were 9 weeks old. (C) Confocal images of Na⁺/K⁺-ATPase, CRALBP, and claudin-19 RPE markers in 7-week-old hESC-RPE cells cultured without the hydrogel (control) and with the hydrogel for 2 weeks. hESC-RPE cells were labeled with immunofluorescence staining after removing the hydrogel. XY sections: maximum intensity projections. YZ sections: maximum intensity projections of 5 images with location indicated as a white line in the XY section. Scalebars 10 μm. (D) Vascular endothelial growth factor (VEGF) secretion was analyzed with ELISA assay from hESC-RPE cells cultured with the hydrogel for 2 weeks (controls cultured without the hydrogel). Data from two ELISA assays, with one datapoint in the plot containing multiple inserts (n = 41). Statistical testing with 2-way ANOVA (E) TEER was measured from hESC-RPE cells cultured 2 weeks with the hydrogel (controls cultured without the hydrogel). Statistical testing with 2-way ANOVA. Significance codes: '*' P < 0.05, '**' P < 0.01 and '***' P < 0.001.

First, the hESC- and hiPSC-RPE cells were plated on 24-well Transwell inserts (Fig 1B). After 5 weeks of culture, when RPE cells had formed a monolayer and started to form pigment but were still considered immature (20,30), the hydrogels were placed on RPE apical side, allowing the cells to interact with the retina-mimicking hydrogel. The cells were cultured further with the hydrogel for 2 to 4 weeks. Control RPE cells were cultured without the hydrogel using the conventional culture method. Additionally, to test the hydrogel with more mature RPE, we used 7-week-old RPE cells and cultured them with the hydrogel until the cells were 9 weeks old. During culture, RPE cells were able to bind the Matrigel-coated PA hydrogels and adhered firmly to the hydrogel surface on their apical side.

Next, we wanted to investigate the hydrogels' effect on RPE cell physiology and morphology. The hydrogels were carefully removed from the inserts, and cells were fixed with 4 % paraformaldehyde (PFA) and labelled with immunofluorescence staining. Labelling of the characteristic RPE markers Na⁺/K⁺-ATPase, CRALBP, and claudin-19 (Fig 1C) and Bestrophin, ZO-1, Ezrin, $Ca_V1.3$, Connexin43, and vimentin (Fig S1) showed no major differences between the RPE cells cultured under the hydrogel for 2 weeks and control RPE cells cultured without the hydrogel. To test one of the physiological functions of the RPE, vascular endothelial growth factor (VEGF) secretion was quantified with ELISA assay from both apical and basal side cell media (Fig 1D). VEGF secretion is known to be more active from the basal side of RPE *in vivo* (30) and the same was observed in our RPE cultures with and without the hydrogel. In addition, VEGF secretion increased with time in control samples measured at week 5 (mean ± SD: apical side 1160 ± 430 and basal side 1940 ± 290 pg/ml), week 7 (apical 1330 ± 260 and basal 2450 ± 200 pg/ml), and week 9 (apical 1560 ± 200 and basal 3130 ± 520 pg/ml) after plating the cells (Fig 1D). However, RPE cells cultured with the hydrogel for 2 weeks showed no significant increase in the secreted VEGF between week 7 (apical 1340 ± 400 and basal 2660 ± 220 pg/ml) and week 9 (apical 1690 ± 290 and basal 3470 ± 670 pg/ml). Still, no significant differences at any time points were seen in VEGF secretion between controls and RPE-hydrogel samples. Finally, we followed the epithelial barrier formation under the gel by measuring transepithelial electrical resistance (TEER) values

from control and RPE-hydrogel samples (Fig 1E). In control cells, TEER values increased significantly between weeks 7 and 9 (from 220 ± 90 to 380 ± 70 Ωcm^2, mean $\pm$ SD). In RPE cells cultured with the hydrogel, TEER did not rise significantly (200 ± 80 Ωcm^2 at week 7, and 220 ± 100 Ωcm^2 at week 9). Thus, at week 9 control cells showed higher TEER values compared to RPE-hydrogel samples. However, both control and RPE-hydrogel samples showed cell-cell junction localization of claudin-19, an important RPE tight junction marker (Fig 1C) (31). This together with the measured TEER values indicate that both controls and RPE-hydrogel samples had functional tight junctions. All in all, RPE cells showed characteristic morphology as well as secretion and barrier functionality, demonstrating compatibility of the hydrogel with the RPE cell cultures.

2.2. RPE adheres heterogeneously to the hydrogel

Our results indicated that RPE can adhere to the apically placed hydrogel and that RPE cells' physiology is not disturbed even during long culturing periods with the hydrogel. To further characterize the retina-mimicking system, the RPE-hydrogel samples were fixed with 4 % PFA and labeled with immunofluorescent staining, with the gel still adhered to the cells. Due to the transparency of the PA hydrogels, we were able to image the RPE cells through the hydrogel with a confocal microscope (Fig 2A). This allowed us to observe the RPE monolayer without disturbing the cell-hydrogel interface. The imaging of RPE cells without removing the hydrogel revealed typical tight junction structures labeled with ZO-1, apical membrane labeled with Ezrin, and actin cytoskeleton labeled with phalloidin. This indicates that the RPE can be easily observed through the hydrogel by using confocal microscopy.

Fig 2. RPE-hydrogel adhesion was characterized with imaging and quantified with rheological tack adhesion test.

(A) Schematic illustration of the imaging setup. RPE cells can be imaged through the hydrogel with a confocal microscope. hESC-RPE cells were fixed with 4% PFA and labeled with immunofluorescence staining with the hydrogel still attached to cells. ZO-1, Ezrin, and actin staining by Phalloidin. Scale bar 25 µm. (B) RPE-hydrogel adhesion was measured with a rheometric tack adhesion test. Live hESC-RPE cells with the hydrogel were placed on a rheometer and the probe was adhered to the hydrogel. Detachment was done by moving the probe up at a constant speed. The force-distance curve is a representative example of one measurement. At the end of the measurement, the force

remained constant as the probe was moving upwards with the detached hydrogel still adhered to it. Adhesion force (maximum force measured, Newtons) was defined from each curve by subtracting this baseline force (indicated as a red line in the graph) from the lowest peak of the curve (detachment point). Adhesion energy (area over the curve, Joules per area) was measured graphically from each curve by calculating the area between the curve and the baseline force. (C) RPE-hydrogel adhesion force and energy were obtained from rheometric tack adhesion tests (n = 14 inserts). In control 1 (n = 3 inserts), RPE-hydrogel samples were treated with actin polymerization inhibitor Cytochalasin D for 1 hour, which detached the hydrogel from RPE cells, and the samples could not be measured (adhesion force and adhesion energy were = 0). For control 2 (n = 6 inserts), RPE-hydrogel samples were first treated with Cytochalasin D for 1 hour, and then fixed with 4 % PFA, which restored the adhesion. Statistical testing with Kruskal-Wallis. Significance codes: '*' P < 0.05.

Previous experiments indicated that the RPE cells are able to adhere to the hydrogel and the hydrogel does not interfere the physiology of the RPE. To further characterize and quantify the adhesion between RPE and the hydrogel, a rheometric tack adhesion test was performed. The membrane with living RPE cells with the hydrogel was cut from the insert and glued on the bottom plate of the rheometer. A cylindrical metal probe was then adhered to the hydrogel with glue and maintained in 0.1 N compression for 2 min. Next, hydrogel detachment from the RPE cells was achieved by moving the probe upwards at a constant speed. As a result, a force-distance curve was obtained, where detachment of the hydrogel typically occurred stepwise instead of an instant detachment (Fig 2B). This indicates that the RPE-hydrogel adhesion is heterogeneous across the cell monolayer, with the weakest adhesion points detaching first. For each measurement, both maximum adhesion force and adhesion energy were determined from the force-distance curve (Fig 2C). After 2 weeks of culture, the average RPE-hydrogel adhesion force was 50 ± 10 mN (mean $\pm$ SD) per sample, corresponding to 1.7 ± 0.4 nN/μm^2. Adhesion energy was 11.2 ± 4.9 μJ/cm^2, corresponding to 112 fJ/μm^2. As a first control, RPE-hydrogel samples were treated before the measurement with Cytochalasin D (10 μg/ml, 1 h), a pharmaceutical actin polymerization inhibitor. The treatment leads to depolymerization of actin fibers and a strong reduction of actomyosin contractility. Cytochalasin D caused complete spontaneous detachment of the hydrogels from RPE cells, corresponding to 0 adhesion force and energy. For a second control, RPE-hydrogel samples were first similarly incubated with Cytochalasin D but then treated with 4 % PFA, a cell fixative and protein crosslinker, which restored the adhesion to 50 ± 20 mN per sample, or 1.9 ± 0.7 nN/μm^2. Adhesion energy was 14.5 ± 7.0 μJ/cm^2, or 145 fJ/μm^2. These controls suggest that the RPE-hydrogel adhesion is reversible and depends on intact actin cytoskeleton. However, actin depolymerization does not physically separate the hydrogel from the RPE in molecular scale, since they can be chemically crosslinked together after the depolymerization.

2.3. RPE can transmit traction forces to the apical hydrogel

The previous experiments indicated that the RPE can adhere to the apically placed hydrogel and that the interaction does not influence the maturation of the RPE monolayer. The tight, actin-cytoskeleton-dependent adhesion thus mimics the close interaction between RPE cells and POS layer of the retina. Thus, the RPE-hydrogel adhesion enabled further studies of the RPE-photoreceptor mimicking interface. We hypothesized that the adhesion between these structures enable the direct transmission of RPE-generated forces from the RPE to the retina-mimicking hydrogel. We investigated this force transmission by conducting traction force microscopy (TFM). In TFM, fluorescent beads are embedded into the linearly elastic cell culturing substrate, e.g., hydrogel, and the forces transmitted across the adhesion between the cells and the substrate can be spatially quantified from the movement of the beads.

Fig 3. RPE-generated forces can be characterized with traction force microscopy.

(A) Schematic illustration of the traction force microscopy (TFM) setup. The hydrogel is embedded with fluorescent beads, and live RPE-hydrogel interface is imaged through the hydrogel with spinning disk confocal microscope. Tractions exerted by RPE cells can be analyzed from the displacement of the beads. (B) Composite image of maximum intensity projections of the beads in RPE-hydrogel surface illustrating their displacement during a 20 min interval. Cyan: fluorescent beads at the start of the interval, magenta: beads after 20 min. Scalebars 50 μm (left) and 20 μm (right). (C) Live hiPSC-derived RPE cells with actin-EGFP expression (green and inverted grayscale color) show that actin inhibition with Cytochalasin D affects actin cytoskeleton of the RPE cells with the hydrogel. Left:

Untreated cells. Right: Cells after actin polymerization inhibition with 10 µg/ml of Cytochalasin D for 30 min. Scalebars 25 µm. (D) Bead displacements were calculated using particle image velocimetry (PIV) analysis and traction stress maps were constructed using Fourier transform traction cytometry (FTTC). Stress (Pascal) represents force per area (N/m²). Representative stress maps of hESC- and hiPSC - derived RPE cells' interaction with the hydrogel, with and without Cytochalasin D. Scalebars 50 µm. (E) Average changes in cellular traction stresses over 20 min were calculated from each image pair. Data from 10 inserts, from which 5 inserts were used for actin inhibition with Cytochalasin D. Statistical testing with Mann-Whitney U. Significance codes: '*' P < 0.05.

Here, we embedded fluorescent beads (diameter 200 nm) into the PA hydrogels prior to polymerization, placed the hydrogels on top of the RPE cells, and cultured for 2 or 4 weeks. RPE-hydrogel interface was then imaged twice with a 20 min interval through the hydrogel with a spinning disk confocal microscope to detect the bead displacements caused by the changes in cell-generated forces during the 20 min period (Fig 3A and 3B).

Our previous experiments indicated that the RPE-hydrogel adhesion can be significantly reduced with the actin polymerization inhibitor Cytochalasin D. To confirm that the Cytochalasin treatment leads to actin cytoskeleton disruption in the cells beneath the hydrogel, we used hiPSC-derived RPE cells with actin-EGFP expression (Fig S2). Live imaging of EGFP-actin hiPSC-RPE cells with the hydrogel showed disruptions in actin organization of the epithelial monolayer during the inhibition compared to the untreated cells (Fig 3C).

To compare the RPE-generated forces with and without an intact actin cytoskeleton, the beads inside the hydrogel were first imaged twice with a 20 min interval, and then RPE cells' actin cytoskeleton was pharmaceutically depolymerized with Cytochalasin D for 40-60 min. The sample was then imaged for the second time, again twice with a 20 min interval. Bead displacement fields were computed from the microscopy image pairs taken 20 min apart with iterative particle image velocimetry (PIV) with a final vector spacing of 0.8 µm (32). Since the rigidity and Poisson's ratio of the PA hydrogels are known, the bead displacement fields were then used to construct traction vector plots (Fig S3A) and stress magnitude maps (Fig 3D) via Fourier transform traction cytometry (FTTC) (32). The stress magnitude maps were heterogeneous and showed areas of low or high stress with local stress peaks, indicating variability in the activity of the RPE cells. The observed stresses generated by RPE tractions were diminished following the actin depolymerization with Cytochalasin D, indicating the importance of actomyosin generated forces in the RPE tractions.

For quantification, average values of the traction stresses were calculated from each image pair, corresponding to the average change in traction stress during the 20 min interval. With the hydrogel on their apical side, RPE cells generated lateral stresses on average of 430 ± 180 Pa (pN/µm², mean ± SD) (Fig 3E). As can also be seen in the representative stress maps of both hiPSC- and hESC-RPE cells (Fig 3D), there was a decrease in traction stresses after inhibition of actin cytoskeleton with Cytochalasin D (on

average to 160 ± 80 pN/μm^2). This confirms that the bead displacements and thus the traction stresses analyzed here are caused by the actin cytoskeleton contractility of the RPE cells. These actin dependent forces are then transmitted to the retina-mimicking hydrogel on the apical side of the RPE cells via the cell-hydrogel contacts.

2.4. RPE actively pulls on photoreceptor outer segments

RPE has an important role in the maintenance of the retina, and it actively recycles the tips of POS via phagocytosis. The process is strongly dependent on an intact actin cytoskeleton and integrin receptors on the apical surface of the RPE cells. (1) Since the RPE can transmit forces to the retina-mimicking hydrogel, we hypothesized that active force transduction happens during the phagocytosis process. This allows the RPE cells to pull the POS to the vicinity of the cell membrane, facilitating the internalization of the particles.

Fig 4. RPE actively generates forces during phagocytosis.

(A) Schematic illustration of the imaging setup to analyze the force transduction during phagocytosis. Thin PA hydrogel (height approximately 100 μm) constructed on a glass coverslip was embedded with fluorescent beads and coated with NHS-acrylic acid and fluorescently labeled photoreceptor outer segment (POS) particles. Insert membrane with RPE cells was cut and placed on the hydrogel with RPE cells' apical side facing towards the POS. Imaging was done through the glass coverslip and the thin hydrogel with a laser scanning confocal microscope. During imaging the RPE cells were kept in media supplemented with 10 % FBS. XY-maximum intensity projection shows RPE cells (grey, actin labeled with live cell dye SiR-actin) and POS particle localization above (green).

*Scale bar 10 µm. YZ image shows RPE cells from the side, with internalized POS particle inside the cell. Scale bar 5 µm. (B) Upper image: YZ-maximum intensity projection of 5 images from the RPE-hydrogel interface showing cellular actin labeled with SiR-actin (grey), fluorescently labeled POS particles (green), and beads embedded in the hydrogel (magenta). Scalebar 10 µm. Lower image: YZ-magnification of a single cell. Apical side microvilli (grey) protrude towards the POS particle (green) during 20 min. Scalebar 5 µm. (C) From each insert, 2-3 separate positions were chosen and imaged alternately. Traction stresses were analyzed from two images taken with an interval of 20 min (Δ20 min). 10 µg/ml of Cytochalasin D was added to the imaging media without moving the sample, and image acquisition in the positions was continued with 20 min intervals (Δ20 min with actin inhibition). In addition, analysis was performed by comparing the bead positions before and after the addition of Cytochalasin D (Relaxation). Average intensity Z-projections of FTTC stress maps from each condition (Δ20 min 21 images from 8 inserts, Relaxation 13 images from 5 inserts, and Δ20 min with actin inhibition 10 images from 5 inserts). Scalebars 10 µm. (D) From each image pair, an average value of the change in traction stresses during 20 min was calculated. Statistical testing with Kruskal-Wallis. (E) Left: Composite image of a traction stress map during cell cytoskeleton relaxation with maximum intensity Z-projection of POS particles (green). The white circles illustrate the areas (d = 10 µm) around the POS particles chosen for the analysis. Scale bar 10 µm. Right: Average traction stresses were calculated from each chosen area in vicinity of POS particles (n = 39 positions from 13 images, from 5 inserts). Control areas were randomized areas within the same images. Statistical testing with Mann-Whitney U. Significance codes: '**' P < 0.01 and 'ns' P > 0.05.*

To study the forces that RPE cells generate during phagocytosis, thin PA hydrogels with fluorescent beads were constructed on glass coverslips. In addition, POS particles were extracted from porcine eyes, labeled with a fluorescent dye, and incorporated on the PA hydrogel surface. Covalent linkage and proper attachment of the POS particles to the PA gel was achieved by mixing NHS-acrylic acid to the PA gel precursor solution. This process yielded a soft hydrogel suitable for TFM, with POS particles on its surface. The hydrogel was not coated with any other ECM proteins, so the POS particles provided the only binding sites for the RPE cells. The insert membrane was then cut and placed on top of the hydrogel so that RPE cells faced down to the POS particles. (Fig 4A). The thin hydrogel allowed imaging through both the glass coverslip and the hydrogel with a laser scanning confocal microscope to achieve better spatial resolution. We used SiR-actin to label the actin cytoskeleton of living hESC-RPE cells, allowing simultaneous imaging of the actin dynamics, bead displacement, and POS particles. During imaging, RPE cells were in close contact with the hydrogel, and RPE cells reached towards the POS particles with apical actin-rich -protrusions. Moreover, RPE cells were able to internalize parts of the POS particles incorporated in the hydrogel. (Fig 4A and 4B)

Images were again acquired at 20 min intervals from different positions of the sample (Fig 4C). Next, Cytochalasin D (10 µg/ml) was added to the imaging media without moving the sample and incubated for at least 60 min. The same positions of the sample were imaged again, and traction stresses were analyzed from the acquired data before and after actin inhibition. Finally, we compared two images taken before and after the addition of Cytochalasin D, to study the change in stress magnitude during complete cell

cytoskeleton relaxation. Analysis was performed by first computing bead displacements with PIV analysis from image pairs with 20 min intervals, and then calculating the traction vector plots (Fig S3B) and stress maps with FTTC. Averages of the stress magnitude maps indicate that actin polymerization inhibition decreases traction stress generation in RPE during POS phagocytosis (Fig 4C).

For quantification, average values of the traction stresses were calculated from each image pair, corresponding to the average change in traction stress during the 20 min interval (Fig 4D). During phagocytosis, RPE cells generated stresses on average of 430 ± 200 Pa ($pN/\mu m^2$, mean $\pm$ SD). Interestingly, tractions measured within 20 min intervals did not differ significantly from the ones during relaxation (500 ± 280 $pN/\mu m^2$), suggesting a very dynamic force generation in RPE especially at the early phases of the phagocytosis process. The actin inhibition with Cytochalasin D decreased traction stress generation during the 20 min interval to 120 ± 70 $pN/\mu m^2$. As a control, we used hydrogels without any surface modifications (Fig S4). In the absence of POS particles, hydrogel's NHS-acrylic acid supplementation, and serum the quantified traction forces were only a few tens of $pN/\mu m^2$. Furthermore, the stresses were again negligible with the addition of serum (10 % FBS) and NHS-acrylic acid, demonstrating that POS particles are essential for force transduction and subsequent traction stress, and that RPE cells are unable to attach to the gel without them.

Finally, we wanted to compare the magnitude of the traction stresses around POS particles to areas without POS (Fig 4E). Three circular areas around POS particles (approximately 10 µm in diameter) were chosen in each traction stress map, representing the change in traction stress during cell cytoskeleton relaxation. Control areas were identical areas randomly within the same image. Traction stresses were significantly higher near the POS particles (800 ± 590 $pN/\mu m^2$) compared to the randomly chosen control areas (320 ± 350 $pN/\mu m^2$). This result indicates that the tractions measured here are mainly caused by RPE interacting with the POS particles on the hydrogel surface. Altogether, this data shows that RPE cells are able to focus physical force to the vicinity of POS during early phases of phagocytosis.

3. Discussion

At the back of the eye, the adhesion between the retina and its underlying tissue, the retinal pigment epithelium (RPE), is vital for retinal wellbeing and functionality. This adhesion allows the close interaction between the RPE and photoreceptor outer segments (POS). In addition, microvilli on the RPE apical side protrude between POS and form an interlocked, physical connection. This physical interaction is needed for the RPE cells to internalize and degrade parts of the POS daily in the phagocytosis process (1,2). RPE cells can be cultured *in vitro* on semipermeable membranes in inserts, which mimic the *in vivo* conditions by allowing transportation of nutrients and ions through the apical and basal sides of the cells (1,20,21). However, most current RPE cell culture systems are missing the RPE-retina connections, possibly limiting not only RPE maturation (30,33) but also hindering studies addressing the biophysical characteristics of the RPE-retina interface.

In vivo, the RPE-retina adhesion is mediated by IPM and RPE-retina interdigitations, and further co-regulated by metabolic factors controlling the subretinal pH, ion composition, and retinal oxygenation. In addition, RPE cells actively pump fluids from the subretinal space to the choroid, creating a pressure that pulls RPE and retina together. If these adhesion forces are overwhelmed, retinal detachment can occur. (10,34) Retinal adhesion decreases postmortem and is affected by environmental factors such as pH, temperature, and ion composition (10,35–37), limiting the possibilities to maintain *ex vivo* RPE-retina tissue sections *in vitro* (1). Alternatively, primary RPE cell cultures have been used with *ex vivo* retinas in direct RPE-retina co-cultures, where the cells remained viable for few days (38). Organoids and organ-on-a-chip technologies offer interesting possibilities for *in vitro* culturing, but they are still missing the direct RPE-retina connections. While Gabriel et al. (2021) (39) succeeded to generate RPE cells in optic vesicle-containing brain organoids, contacts between RPE cells and photoreceptors have still been absent in retinal organoids (40,41). Achberger et al. (2019) (42) combined both RPE cells and retinal organoids in a microfluidic system to create a retina-on-a-chip. However, RPE and POS were still separated with hyaluronic acid -based hydrogel, thus restricting direct RPE-retina connections. Moreover, organoids and organ-on-a-chips are limited with the methodologies with which they can be studied. Understanding the biophysical properties of the RPE-retina interface using, for example, high resolution microscopy or rheology, can be challenging in these systems. Hydrogels, on the other hand, are highly tunable and versatile platforms that can be used to mimic tissues and extracellular environments in simplified *in vitro* cell culture conditions.

We showed that a retina-mimicking structure can be constructed from soft polyacrylamide (PA) hydrogel, coated with Matrigel, and cultured on the apical side of hESC- and hiPSC-derived RPE cells in Transwell inserts. We chose PA for the retina-mimicking platform material based on its mechanical properties, well established

polymerization and conjugation chemistry, and compatibility with high-resolution confocal microscopy. PA hydrogels are easy to construct at a low cost, and their elastic modulus can be tuned by simply changing the ratios of the monomer and the crosslinker, making them suitable for high-throughput cell culture experiments (24,26). In addition, cells cannot spontaneously adhere to PA hydrogels, but the surface can be modified by covalently linking selected ECM proteins using, for example, L-DOPA (25). Even though PA hydrogels are a simple and robust platform for cell culture experiments, more *in vivo* - like hydrogel materials could be considered for RPE cells, such as hyaluronic acid, one of the main components of the IPM, the specialized ECM found between RPE and photoreceptor cells in the eye (7). However, mechanical properties, stability and the construction of the hyaluronic acid hydrogels would need to be tailored for the RPE cultures (43).

In addition, we used Matrigel to covalently link ECM proteins to the hydrogel surface. Matrigel, a commercially-available basement membrane ECM secreted by mouse sarcoma, contains some of the main components of the IPM, such as laminin and heparan sulfate proteoglycans together with TGF-ß growth factors (7). The PA hydrogels used in this work have a flat surface, while *in vivo* RPE and POS form tight, interlocked connections with RPE microvilli protruding between the outer segments. Interestingly, we observed different RPE cell responses depending on whether the hydrogels were polymerized on glass coverslips or cell culture plastic. RPE cells cultured with hydrogels polymerized on glass showed abnormal and disrupted apical microvilli, whereas hydrogels polymerized on plastic led to apical actin structures in the RPE that were comparable to control cells without a hydrogel (Fig S5A). Glass coverslips have a smooth surface with a root mean square (RMS) roughness of less than 1 nm. In contrast, cell culture plastics have an RMS roughness of 1–6 nm, a factor known to influence cell adhesion and proliferation (44). We speculate that this nanoscale roughness is replicated on the hydrogel surface and contributes to the RPE response to the apically placed hydrogel. As a proof-of-concept, we also fabricated hydrogels with retina-mimicking topographical features on the PA hydrogel surface using polydimethylsiloxane (PDMS) molds created by photolithography. This approach successfully generated photoreceptor-sized wells on the hydrogel surface (Fig S5B-D, Supplementary Protocol 1), potentially allowing RPE microvilli to protrude into these wells. However, the construction of these soft-topography hydrogels was technically challenging. Therefore, future work should consider alternative materials and topographical fabrication techniques. The remaining experiments were conducted with a hydrogel casted on cell culture plastics.

We first characterized the RPE-retina mimicking platform by studying the effect of the hydrogel on RPE cells' morphology and functionality. RPE cells cultured with the hydrogel on their apical side did not show significant differences in cell morphology, RPE marker expression, epithelial layer integrity, or growth factor secretion, compared to cells cultured without a hydrogel (Fig 1). The measured TEER values of the RPE cells were around 200 Ωcm^2, which in line with values measured from human tissue (45) and porcine primary RPE cell cultures (18). Previously it has been shown that mechanical compressive stress can influence cell phenotype and signaling (46–48). However, our

data indicates that RPE tolerates the hydrogel compression without changing their phenotype. Furthermore, the RPE cells adhered to the hydrogel from their apical side, enabling, to our knowledge for the first time, *in vitro* modeling of the RPE-retina interface.

Our retina-mimicking hydrogel enabled measurements of the adhesive properties on RPE apical side in a simplified *in vitro* setup. We measured the maximal RPE-hydrogel adhesion with a tack test to be approximately 170 mN/cm^2, or 50 mN for an insert with 75000 cells (Fig 2). That corresponds to a gravitational force generated by a weight of 5.1 g or a vertical pulling force of approximately 17 bumblebees (49). As a cell-level comparison, individual cells in endothelial monolayers have been reported to have a cell-substrate adhesion force of 1170 nN measured with fluidic force microscopy, corresponding to approximately 90 mN for 75000 cells (50). Furthermore, the RPE-retina adhesion has been qualitatively measured from tissue sections *in vitro* by peeling the retina from the RPE and measuring the pigment residues in the retina, with more pigment indicating stronger adhesion (51,52). Even though this method allows fast measurements within only minutes after enucleation, it is still subject to changes in the experimental setup, and cannot distinguish differences in adhesion beyond full pigment coverage (37,53). Alternatively, Marmor et al. (1980) (35) quantified the RPE-retina adhesion *in vitro* from dissected eyecups by measuring the peeling force required to separate retina from the RPE with a force transducer. However, more physiologically relevant RPE-retina adhesion measurements can be achieved in an *in vivo* setup, where RPE and retina are separated by subretinal injection while measuring the pressure inside the generated 'bleb' (53,54). Adhesion force can then be calculated according to Laplace's law. Chiang et al. (1995) (53) showed that *in vivo* injection of Cytochalasin D, the same actin polymerization inhibitor we used in our experiments, into the subretinal space weakened the RPE-retina adhesion, and this effect was reversible. Thus, similarly to our *in vitro* model, intact actin cytoskeleton is needed for proper retina-RPE adhesion. However, when we fixed the Cytochalasin D treated samples with a paraformaldehyde crosslinker, we saw the restoration of the adhesion. This suggests that the adhesion is mediated by RPE cells' apical side and actin cytoskeleton depolymerization does not cause separation of the RPE and the hydrogel in molecular scale.

Our setup also enabled studies on force transduction from RPE cells to the retina-mimicking hydrogel. We studied this with traction force microscopy (TFM), an imaging technique where fluorescent beads incorporated in the hydrogel are used to reconstruct the material deformation and subsequently the cell-generated traction forces leading to the deformations (55). Our TFM experiments showed that after the hydrogel was cultured on the RPE apical side for 2-4 weeks, the RPE cells actively transmit forces into the hydrogel (Fig 3). TFM images revealed a speckled pattern of force transmission, with some regions in the field of view showing higher tractions, indicating either heterogeneous adhesion between the RPE and the hydrogel or variability in RPE force production. Moreover, Cytochalasin D considerably decreased the tractions between RPE and the hydrogel, indicating that the forces are indeed rising from the actin cytoskeleton of the RPE cells. Even though the roles of integrin $\alpha_V\beta_5$ (52) and actin (53) in the RPE-

retina adhesion have been addressed previously, our retina-mimicking hydrogel has enabled simple quantifications of mechanical force transmission from the apical side of RPE to the retina-mimicking hydrogel.

As we developed a method to study force transmission from RPE to the apically placed hydrogel, we next aimed to study the possible force transmission during RPE phagocytosis. To quantify the traction forces that RPE cells generate during phagocytosis, we further developed the retina-mimicking hydrogel system. We constructed thin hydrogels on glass coverslips to enable imaging of the hydrogel surface with high-resolution confocal microscopy. The hydrogel was embedded with fluorescent beads to follow the RPE cell generated tractions, and purified POS particles from porcine eyes were incorporated on the hydrogel surface for RPE cells to phagocytose. To study the phagocytosis process from its initiation, instead of culturing the hydrogel on top of RPE cells for several weeks, the insert membrane with RPE cells was cut and placed on the hydrogel with POS particles. During the experiment, RPE cells were able to connect with the POS particles and internalize parts of them. Previous studies have shown that RPE's apical side actin-rich protrusions form phagocytic cups to scission and ingest parts of the POS (14,15,56–58) together with myosin II (17,59), suggesting that RPE cells must generate considerable forces during this process. We measured RPE generated tractions in the scale of hundreds of pN/µm^2 (Fig 4). In highly contractile cells, the traction stresses in individual focal adhesions on the basal side reach a few nN/µm², suggesting that the RPE can generate considerable forces in their apical side during phagocytosis. However, traction stress quantification by PIV relies solely on the in-plane movement of fluorescent beads, meaning that bead displacement directly toward or away from the RPE is not detected. Thus, the forces quantified here represent a lower estimation of the total magnitude of forces generated by the RPE, rather than measurement of total forces. More precise quantification would require three-dimensional analysis with advanced methods (55).

POS phagocytosis by RPE is dependent on αvβ5 integrin (60), and interestingly, αvβ5-rich reticular adhesions have been shown not to associate with actin fibers (61). However, decrease in force generation within 20 min intervals was seen with the addition of Cytochalasin D, showing that actin cytoskeleton participates in the RPE apical side force generation in phagocytosis. This is in line with previous work by Umapathy et al. (2023) (14), where they showed with live cell imaging that during phagocytosis Cytochalasin D did not affect POS binding to the RPE apical surface but prevented the formation of actin-rich phagocytic cups. Furthermore, our results support the emerging view of RPE being a mechanically active participant in POS phagocytosis, rather than passively clearing shed POS particles from the subretinal space (1).

In conclusion, we have developed a retina-mimicking platform from PA hydrogel to be used in *in vitro* cell culture of hESC- and hiPSC-derived RPE cells. We showed that the hydrogel platform enables measurements of the RPE adhesion to the retina-mimicking hydrogel *in vitro* without affecting RPE cell morphology or functionality. We showed, to our knowledge for the first time, that RPE cells exert considerable forces to POS particles

during phagocytosis and that the force generation is dependent on an intact actin cytoskeleton. Together, our results support the view of the RPE as a mechanically active partner for the retina and provide tools for further biophysical *in vitro* modeling of the RPE-retina interface.

5. Materials and methods

5.1. Polyacrylamide hydrogel construction

Polyacrylamide (PA) hydrogel construction was done based on the work of Tse and Engler (2010) (24). For E=2.8kPa PA hydrogel, acrylamide (AA) (stock 40 %, #1610140, Bio-Rad Laboratories, USA) and Bis-acrylamide (Bis) (stock 2 %, #1610142, Bio-Rad Laboratories) were mixed in the ratio of AA 10 % and Bis 0.03 % in PBS in a falcon tube. Solution was degassed in vacuum. To initiate polymerization, 0.2 % v/v of TEMED (#1610800, Bio-Rad Laboratories) and 1 % v/v of ammonium persulfate (APS) stock solution in ddH$_2$0 (10 % w/v, #A3678, Sigma-Aldrich, USA) were added to the PA solution and mixed by tilting the tube couple of times. The PA hydrogel polymerization was done by pipetting either 150 µl (hydrogels used for immunolabelling and traction force microscopy) or 250 µl (hydrogels used for rheology) of PA solution on plastic cell culture dish lid (#130181, Thermo Fisher Scientific, USA). Smaller cell culture dish lid (#353001, Corning, USA) was placed on top of the PA solution drop for polymerization of 20 min in room temperature. Once polymerized, the PA hydrogels were placed in PBS (#14200-083, Gibco) and stored at +4 °C.

For cell culture, PA hydrogels were cut smaller with puncher (d = 6 mm, #504533, World Precision Instruments, USA) to fit inside the 24-well cell culture insert. To facilitate cell adhesion and growth, the PA hydrogels were coated with Matrigel (#356231, Corning). As proteins cannot spontaneously adhere on PA hydrogels, the hydrogels were first coated with chemical linker 3,4-dihydroxy-L-phenylalanine (L-DOPA) (#D9628, Sigma-Aldrich) according to Wouters et al. (2016) (25). First, L-DOPA was dissolved in 10 mM Tris buffer pH 10 (#0497, VWR, USA) for 30 min in dark. The PA hydrogels were incubated with L-DOPA in 24-well plates for 30 min in dark in swing 5 rpm and washed twice with PBS for 10 min. Matrigel was thawed on ice and diluted to 0.1 mg/ml in PBS. The hydrogels were incubated with Matrigel solution for 1 h in dark in swing 5 rpm and washed once with PBS. Kauppila et al. (2023) (29) have shown that this L-DOPA and ECM coating on PA gel surface does not affect local stiffness values significantly. Before cell culture, the hydrogels were sterilized with UV light for 15 min in UV light (UVP CL-1000 Ultraviolet Crosslinker, USA).

5.2. Cell culture

Human embryonic stem cell (hESC) lines Regea08/017, Regea11/013 and Regea08/023 and human induced pluripotent stem cell (hiPSC) line expressing mEGFP tagged beta-actin (actin beta (ACTB) (ID: AICS-0016 cl.184) developed at the Allen Institute for Cell Science (allencell.org/cell-catalog) and available through Coriell (62)) were differentiated and cultured as described previously (20,21). Cells were plated on 24-well plate inserts (#83.3932.101, Sarsted, Germany) at a density of 2.5*10^5 cells/cm^2. Inserts were coated

with 10 µg/cm^2 Collagen IV from human placenta (#C5533, Sigma) and 1,8 µg/cm^2 Laminin 521 (LN521, Biolamina, Sweden). RPE cells were maintained at 37 °C in 5 % CO$_2$ in DM- media, containing knock-out Dulbecco's modified Eagle's medium (#10829018, Gibco, USA) with 15 % knock-out serum replacement (#10828-028 , Gibco), 2mM GlutaMax (#35050038, Gibco), 1 % MEM Non-essential amino acids (#11140050, Gibco) , 0.1 mM 2-mercapthoethanol (#31350010, Gibco) and 50 U/ml penisillin/streptomycin (#15140122, Gibco). The media was changed every 2–3 days.

PA hydrogels were placed on top of the RPE monolayers after 5 to 7 weeks after plating them on the inserts. First, all the media was removed from the insert. The PA hydrogel was gently placed on top of the cells with tweezers. A spacer was added on top of the PA hydrogel to prevent it from moving or floating. Finally, media was added on top of the hydrogel. During culture, the media was changed every 2–3 days.

For vascular endothelial growth factor (VEGF) analysis, media samples were collected from both basal and apical side of the inserts and stored in -80 °C. Media samples were collected before changing the media (2–3 days after the last media change). VEGF concentration was measured with Human VEGF Quantikine ELISA kit (#DVE00, R&D systems, USA) according to manufacturer's instructions. To reduce the effect of variation between cell culture inserts, media samples from 2-3 parallel inserts were pooled together to form one sample for ELISA assay, when possible. Media samples were collected altogether from 41 inserts. In the assay each sample was measured 3 times, and an average was calculated for analysis. In this work altogether 2 ELISA assays were made.

Transepithelial electrical resistance (TEER) was measured with epithelial voltohmmeter (EVOM2, World Precision Instruments, USA) with STX2 electrode (World Precision Instruments). TEER was measured from RPE cells cultured with the hydrogel for 2 weeks and from control cells cultured without a hydrogel. The electrode was warmed up in +37 °C cell culture media before and in between the measurements. Each insert was measured 3 times, and an average was calculated for analysis. Empty insert with and without the hydrogel and with cell culture media were measured and subtracted from the averaged TEER values to remove the background resistance. TEER was measured from hiPSC-RPE cells with actin-EGFP expression by first keeping the cells in RT for 15 min. Each insert was measured 2 times, and the electrode was placed in 1xPBS between each measurement.

5.3. Immunofluorescence staining and confocal imaging

In this work, fixing and immunofluorescent staining of the RPE cells cultured under the hydrogel was done in two different ways. In the first method, the insert membrane was cut with scalpel and the gel was gently removed from the live cells with tweezers. All the following steps were done at RT. Cells were fixed in 4 % PFA in PBS 10 min and washed 3 times in PBS for 5 min. Insert membrane was then cut into 2 or 4 pieces with scalpel. Permeabilization was done in 0.1 % Triton-X-100 (v/v) in PBS for 15 min. Blocking was

done in 3 % bovine serum albumin (BSA) (w/v) in PBS for 1 h. Primary antibody incubations were done in 3 % BSA (w/v) in PBS for 1 h. Samples were washed 4 times in PBS for 5 min. Secondary antibody incubations were done in 3 % BSA (w/v) in PBS for 1 h in dark. Samples were washed 3 times in PBS for 5 min and mounted between two glass coverslips (18x18 mm, # 474030-9000-000, Zeiss, Germany) in Prolong diamond antifade mountant (#P36961, Molecular Probes, USA). Samples were dried overnight at RT and then stored at +4 °C.

RPE cells were also stained with the hydrogel still attached to cells. First, samples were washed twice with PBS and fixed in 4 % PFA (v/v, #157-8, Electron Microscopy Sciences, USA) in PBS for 30 min at room temperature (RT). Samples were washed 3 times with PBS for 15 min. Permeabilization was done overnight at +4 °C in 0.5 % Triton-X-100 (v/v, #T8787, Sigma-Aldrich) and 3 % BSA (w/v, #P06-139210, PAN-Biotech, Germany) in PBS. Primary antibody incubation was done in 3 % BSA (w/v) in PBS for 24 h at 4 °C. Samples were washed in 0.5 % Triton-X-100 (v/v) and 3 % BSA (w/v) in PBS three times for 4 h and then overnight at +4 °C. Secondary antibody incubation was done in 3 % BSA (w/v) in PBS for 24 h at +4 °C. Samples were washed in PBS three times for 4 h and then overnight at +4 °C. Samples were stored in PBS at +4 °C. For imaging, the insert membrane and the hydrogel adhered on the cells were cut with scalpel (nro10, Schwann-Morton, UK) and placed on cell culture dish (#130181, Thermo Fisher Scientific). To avoid the sample from floating, insert membrane was adhered on the dish with 3 drops of rapid glue.

Primary antibodies used in this work were: ZO-1 1:50 (#61-7300, Thermo Fisher Scientific), Claudin-19 1:100 (#MAB6970, R&D Systems), Ezrin 1:100 (and 1:200 in Fig S2) ([3C12], #ab4069, Abcam, UK), $Ca_V1.3$ 1:100 (#ACC-005, Alomone Labs, Israel), Na^+/K^+-ATPase 1:200 ([464.6], #ab7671, Abcam), Bestrophin 1:200 (and 1:100 in Fig S2) (#016-Best1-01, Lagen laboratories, USA), CRALBP 1:200 (and 1:500 in Fig S2) ([B2], #ab15051, Abcam), Connexin43 1:200 (#C6219, Sigma-Aldrich), MERTK 1:50 (#H00010461-M01, Abnova, Taiwan), Rac1 1:100 (#610650, BD Transduction Laboratories, USA), $PanNa_V$ 1:200 (#ASC-003, Alomone Labs), Vimentin (#ab92547, Abcam). Secondary antibodies were goat anti-rabbit Alexa Fluor 488 1:200 (#A-11008, Invitrogen, USA) and goat anti-mouse Alexa Fluor 568 1:200 (#A-11031, Invitrogen). In addition, goat anti-mouse Alexa Fluor 647 (#A-21235, Invitrogen) was used (Fig S2). Cellular actin was labelled with ATTO 643 Phalloidin 1:100 (#AD643-81, ATTO-TEC, Germany).

Samples with the hydrogel still adhered on the cells were imaged through the hydrogel with Nikon Eclipse FN1 upright spinning disk confocal microscope (Nikon Europe BV, Amsterdam, Netherlands) with Nikon CFI Apo NIR 60x/1.0 water dipping objective (Fig 2A). The light source used in imaging was Spectra X (Lumencor, USA) and the camera iXon 888 EMCCD (Andor, UK). Voxel size was set to x = y = 0.22 µm and z = 0.3 µm and 1024 x 1024 pixel sized images were obtained and saved in .nd2 format. Images were

finalized with ImageJ by performing gaussian blur with radius 1.0 and linear brightness and contrast adjustments.

Mounted samples were imaged with two laser scanning confocal microscopes (LSCM). Zeiss LSM 800 LSCM was used with Zeiss Plan-Apochromat 63x/1.40 oil immersion objective with 488nm, 561nm and 640nm lasers (Fig 1C, S1, S2 and S5A). Voxel size was set to x = y = 0.1 µm and z = 0.15 µm. 1024 x 1024 pixel images were obtained with line averaging 2 and saved in .czi format. Images were finalized with ImageJ by performing gaussian blur with radius 1.0 and linear brightness and contrast adjustments.

5.4. Traction force microscopy

For traction force microscopy (TFM) with the hydrogel cushion (Fig 3), 200 nm fluorescent beads were added to the PA gel (3 % vol/vol, #F8807, Invitrogen) prior to polymerization. RPE cells were cultured under the hydrogel for 2 or 4 weeks. Before imaging, the insert membrane was cut and fixed on cell culture dish (#130181, Thermo Fisher Scientific) with rapid glue with the hydrogel still attached to the live cells. During imaging samples were kept in 5 ml of Ames' medium (#A1420, Sigma-Aldrich) with 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid buffer (HEPES) (H3375, Sigma-Aldrich), 5 % knock-out serum replacement (v/v, #10828-028, Gibco) and 0.2 % penicillin/streptomycin (v/v, #15140122, Gibco). Cells were kept at +37 °C during imaging. Imaging was done with Nikon Eclipse FN1 upright spinning disk confocal microscope with Nikon CFI Apo LWD 25x/1.1 water dipping objective. Voxel size was set to x = y = 0.52 µm and z = 0.4 µm. 1024 x 1024 pixel timelapse z-stacks were obtained with interval of 5 min or 10 min and duration of 20 min and saved in .nd2 format. For control, after the first 20 min timelapse actin polymerization inhibitor Cytochalasin D (#C2618, Sigma-Aldrich) was added to the imaging media in the concentration of 5 or 10 µg/ml. Imaging was continued for 2-3 more timelapses for 40-60 min, and the last timelapse was used for the image with actin inhibition.

Images of the fluorescent beads were processed in ImageJ by subtracting background (radius 20.0) and adjusting image brightness and contrast linearly. Maximum intensity projections were made from two timepoints with 20 min interval so that only images from near the gel surface were included in the projections. The images were aligned with ImageJ plugin StackReg (rigid body) (63). Bead displacement fields were calculated with iterative particle image velocimetry (PIV advanced) in ImageJ (32). 1st pass PIV parameters were set so that the interrogation window size was 64 pixels, search window was set to be interrogation window x 2 (128 pixels) and vector spacing was set to be half from the interrogation window size (32 pixels). For 2nd pass the interrogation window was set to be half from the 1st pass, and for the 3rd pass half from the 2nd pass, so that the final vector spacing was 8 pixels (0.8 µm). In the case of ambiguous vectors in the displacement field, normalized median test (noise 0.20, threshold 5.00) or dynamic mean test (C1 0.20, C2 5.00) were performed. Traction forces were calculated with ImageJ

plugin Fourier transform traction cytometry (FTTC) (32). For FTTC, the Poisson ratio of PA gels was set to 0.5, and the gel stiffness was set to 2800 Pa. No regularization was used as our comparison of different λ values did not show a significant effect on the traction magnitudes detected (Fig S6). For quantification, the output force field text file created by FTTC was used. From each file, an average value was calculated from the force magnitudes (in Pascals). For illustrative purposes, the displayed pixel values of the FTTC stress magnitude maps presented in Fig 3 were adjusted. Original vector plots are shown in Fig S3.

5.5. Phagocytosis assay with the PA hydrogel

The POS particles were extracted as previously described (20). In brief, porcine eyes were obtained from the slaughterhouse, opened and the retinas were removed with tweezers under red light. Retinas were homogenized in 0.73 M sucrose phosphate buffer by shaking and then filtered. POS were separated by ultracentrifugation (Optima ultracentrifuge, Beckman Coulter, Inc., Brea, CA) at 112400 RCF at 4 °C in sucrose gradient. The POS layer was removed, centrifuged at 3000 RCF at 4 °C and stored in 73 mM sucrose phosphate buffer at -80 °C.

POS were thawed and centrifuged at 5400 RCF for 4 min. For labelling, POS particles were mixed with 1 µg/ml ATTO 488 NHS ester (#AD488-31, ATTO-TEC) in 9.5 mM $NaHCO_3$ in PBS, incubated for 1 h RT and washed three times with 2800 RCF for 3min centrifugation.

For phagocytosis assay, thin PA gels were constructed on glass coverslips with POS particles incorporated on the hydrogel surface. First, both the top (d = 13 mm, VWR, USA) and bottom coverslips (22 x 22 mm, No. 1.5H, Marienfeld, Germany) for PA gel construction were washed in 2 % HellmaneX III (Hellma, Germany) solution in MQH_2O for 30 min in a sonicator, rinsed with excess water and ethanol and dried with compressed nitrogen. The bottom coverslips were activated for the PA gel to adhere to it. Bind-silane (3-(Trimethoxysilyl)propyl methacrylate, #M6514, Sigma-Aldrich) and glacial acetic acid were diluted with 95 % ethanol in final concentrations of 0.3 % v/v and 5 % v/v, respectively. The solution was let to react for 3 min on the bottom coverslips. Next, the bottom coverslips were washed with ethanol and dried with compressed nitrogen. The top coverslips were incubated with the ATTO labelled POS particles for 1 h at RT and washed by dipping the coverslip into a beaker with PBS. Acrylic acid N-hydroxysuccinimide ester (NHS-AA, #A8060, Sigma-Aldrich) was diluted in water-free dimethyl sulfoxide (DMSO, #D2650, Sigma-Aldrich) in 50 mg/ml and further diluted to 0.1 mg/ml in 9.5 mM NaHCO3 in PBS. NHS-AA solution was let to react on the top coverslips for 30 min at RT, and after the incubation washed by dipping the coverslip into a beaker with PBS. During that time, polyacrylamide solution for E = 2.8 kPa hydrogel was mixed and degassed as described above. 200 nm fluorescent beads were added to the PA gel (2 % v/v, F8810, Invitrogen) prior to polymerization with TEMED and APS (10 % w/v). 13 µl of PA solution was pipetted on bind-silane activated bottom coverslip, and

d = 13 mm top coverslip with POS and NHS-AA coating was added on top. Gels were let to polymerize 15 min in dark in RT. After polymerization gels were stored in PBS at +4 °C overnight. The next day coverslips were gently removed with scalpel. For imaging, PA gels on coverslips were placed on live cell imaging chambers (#SC15022, Aireka Cells, China). Control gels were constructed similarly as described above. For controls with NHS-AA on the surface, NHS-AA was incubated on clean top coverslip at 0.1 mg/ml in 9.5 mM NaHCO3 in PBS for 30 min and rinsed with PBS.

For live cell imaging, cellular actin was labelled with SiR-Actin kit (#CY-SC001, Cytoskeleton Inc., USA). hESC-RPE cells were incubated at +37 °C in 5 % CO_2 in DM-media containing 25 pmol Sir-Actin for 3 h. Cells were washed once with DM- media. Insert membrane was cut with scalpel and placed on the PA gel in live cell imaging chamber with cells facing down to the POS. A slice anchor (Warner Instruments) and a spacer were added on top of the insert to prevent it from floating. Imaging was done in Hibernate-A media (#A1247501, Gibco) with 10 % fetal bovine serum (FBS, # 10500064, Gibco). For controls without serum, just Hibernate media was used. Zeiss LSM 780 LSCM was used with Zeiss LD C-Apochromat 40x/1.10 water immersion objective with 488nm, 561nm and 640nm lasers. Voxel size was set to x = y = 0.1 µm and z = 0.37 µm. 1024 x 1024 pixel timelapse z-stacks with interval of 10 min and duration of 20 min were obtained and saved in .czi format. Cells were kept at +37 °C during imaging. From each insert, 3 separate positions were chosen and imaged alternately. For control, 10 µg/ml Cytochalasin D (#C2618, Sigma-Aldrich) was added to the medium during imaging without touching the sample.

Cellular tractions were analyzed from the bead images as was described in section 5.4. Images of POS and cellular actin were finalized with ImageJ by performing gaussian blur with radius 1.0 and linear brightness and contrast adjustments. To quantify the traction stress magnitudes around POS particles, 3 circular regions with diameter of 100 pixels (10.4 µm) were chosen from each traction stress magnitude image, which illustrate the change in traction stress during cell actin cytoskeleton relaxation with Cytochalasin D. For control, the same areas chosen for the 1st image were used in the 2nd image, thus creating randomized control areas within the same dataset. The intensity of the areas chosen was then measured from the 32-bit FTTC traction stress magnitude images, where the pixel values represent the traction stress magnitude in Pascals. For illustrative purposes, the displayed pixel values of the stress magnitude maps in Fig 4C and Fig 4D have been adjusted. Original vector plot of the stress magnitude map in Fig 4D is shown in Fig S3.

5.6. Rheology

A rheometric tack adhesion test was performed to quantify the RPE-hydrogel adhesion using Discovery Series Hybrid Rheometer II from TA Instruments. First, the insert membrane and the hydrogel adhered to the live cells were cut from the insert with a scalpel. The insert membrane was fixed on a 22 x 22 mm glass coverslip (#0107052,

Marienfeldt, Germany) with rapid glue, and the coverslip was taped on the bottom parallel plate of the rheometer. Then, the hydrogel was glued on the probe (top 8 mm parallel plate). A holding period of 2 min with compressive force of 0.1 N was applied. Then, the hydrogel was separated from the RPE cells by moving the probe upwards at a constant linear rate of 10 µm/s. The measurement was set to last for 360 s. As a result, a force-distance curve was drawn in Origin (2019b). At the end of the measurement, the force remained constant as the probe was moving upwards with the detached hydrogel still adhered to it (baseline, illustrated s a red line in Fig 2B). Adhesion force was defined from each curve by subtracting the baseline force from the lowest peak of the curve (detachment point). Adhesion energy was measured graphically from each curve by calculating the area between the curve and the baseline force. As a control, cells were treated with 10 µg/ml Cytochalasin D (#C2618, Sigma-Aldrich) for 1 h at +37 °C in 5 % CO_2. For a second control, cells were first treated with Cytochalasin D and then fixed with 4 % PFA (v/v, #157-8, Electron Microscopy Sciences, USA) in PBS for 30 min at RT.

5.7. Statistical testing

Statistical tests were performed using the IBM SPSS statistics for Windows, version 28 (IBM Corp.) with a significance level of $P < 0.05$. Graphs were made with Origin (2019b). In the graphs, median is illustrated with a horizontal line and mean with a white box. Boxes consist of 25-75% of the sample range, and the line is range within 1.5IQR. Schematics and final figures were assembled in Adobe Illustrator.

Acknowledgements

This work was funded by the Research Council of Finland and Tampere University doctoral school. We thank Outi Heikkilä, Outi Melin and Hanna Pekkarinen for their contributions with the cell culture work. We thank Jens Möller and Nonappa Nonappa for their expert insight regarding this work. The authors acknowledge the Biocenter Finland (BF) and Tampere Imaging Facility (TIF) for the service.

References

1. Lakkaraju A, Umapathy A, Tan LX, Daniele L, Philp NJ, Boesze-Battaglia K, et al. The cell biology of the retinal pigment epithelium. Prog Retin Eye Res. 2020;78(February).

2. Strauss O. The Retinal Pigment Epithelium in Visual Function. Physiol Rev. 2005;85(3):845–81.

3. Wimmers S, Karl MO, Strauss O. Ion channels in the RPE. Prog Retin Eye Res. 2007;26(3):263–301.

4. Young RW, Bok D. PARTICIPATION OF THE RETINAL PIGMENT EPITHELIUM IN THE ROD OUTER SEGMENT RENEWAL PROCESS. J Cell Biol. 1969;42(2):392–403.

5. Lazarus HS, Hageman GS. Xyloside-induced disruption of interphotoreceptor matrix proteoglycans results in retinal detachment. Invest Ophthalmol Vis Sci. 1992;33(2):364–76.

6. Johnson LV, Hageman GS. Structural and compositional analyses of isolated cone matrix sheaths. Invest Ophthalmol Vis Sci. 1991;32(7):1951–7.

7. Ishikawa M, Sawada Y, Yoshitomi T. Structure and function of the interphotoreceptor matrix surrounding retinal photoreceptor cells. Exp Eye Res. 2015;133:3–18.

8. Mieziewska K. The interphotoreceptor matrix, a space in sight. Microsc Res Tech. 1996;35(6):463–71.

9. Nandrot EF, Chang Y, Finnemann SC. αvβ5 Integrin receptors at the apical surface of the RPE: One receptor, two functions. In: Advances in Experimental Medicine and Biology. 2008. p. 369–75.

10. Ghazi NG, Green WR. Pathology and pathogenesis of retinal detachment. Eye. 2002;16(4):411–21.

11. Yao XY, Hageman GS, Marmor MF. Retinal Adhesiveness Is Weakened by Enzymatic Modification of the Interphotoreceptor Matrix In Vivo. Invest Ophthalmol Vis Sci. 1990;31(10):2051–8.

12. Hagstrom SA, Duyao M, North MA, Li T. Retinal Degeneration in tulp1-/- Mice: Vesicular Accumulation in the Interphotoreceptor Matrix. Investig Opthalmology Vis Sci. 1999;40(12).

13. Moran AL, Fehilly JD, Floss Jones D, Collery R, Kennedy BN. Regulation of the rhythmic diversity of daily photoreceptor outer segment phagocytosis in vivo. FASEB J. 2022;36(10).

14. Umapathy A, Torten G, Paniagua AE, Chung J, Tomlinson M, Lim C, et al. Spatiotemporal Live-Cell Analysis of Photoreceptor Outer Segment Membrane Ingestion by the Retinal Pigment Epithelium Reveals Actin-Regulated Scission. J Neurosci. 2023;43(15):2653–64.

15. Spitznas M, Hogan MJ. Outer Segments of Photoreceptors and the Retinal Pigment Epithelium: Interrelationship in the Human Eye. Arch Ophthalmol. 1970;84(6):810–9.

16. Kwon W, Freeman SA. Phagocytosis by the Retinal Pigment Epithelium: Recognition, Resolution, Recycling. Front Immunol. 2020;11:604205.

17. Strick DJ, Feng W, Vollrath D. Mertk Drives Myosin II Redistribution during Retinal Pigment Epithelial Phagocytosis. Investig Opthalmology Vis Sci. 2009;50(5):2427–35.

18. Toops KA, Tan LX, Lakkaraju A. A detailed three-step protocol for live imaging of intracellular traffic in polarized primary porcine RPE monolayers. Exp Eye Res. 2014;124:74–85.

19. Jiang M, Esteve-Rudd J, Lopes VS, Diemer T, Lillo C, Rump A, et al. Microtubule motors transport phagosomes in the RPE, and lack of KLC1 leads to AMD-like pathogenesis. J Cell Biol. 2015;210(4):595–611.

20. Vaajasaari H, Ilmarinen T, Juuti-Uusitalo K, Rajala K, Onnela N, Narkilahti S, et al. Toward the defined and xeno-free differentiation of functional human pluripotent stem cell-derived retinal pigment epithelial cells. Mol Vis. 2011;17:558–75.

21. Viheriälä T, Sorvari J, Ihalainen TO, Mörö A, Grönroos P, Schlie-Wolter S, et al. Culture surface protein coatings affect the barrier properties and calcium signalling of hESC-RPE. Sci Rep. 2021;11(1):933.

22. Ferrara M, Lugano G, Sandinha MT, Kearns VR, Geraghty B, Steel DHW. Biomechanical properties of retina and choroid: a comprehensive review of techniques and translational relevance. Eye. 2021;35(7):1818–32.

23. Caliari SR, Burdick JA. A practical guide to hydrogels for cell culture. Nat Methods. 2016;13(5):405–14.

24. Tse JR, Engler AJ. Preparation of hydrogel substrates with tunable mechanical properties. Curr Protoc Cell Biol. 2010;10.16.1-10.16.16.

25. Wouters OY, Ploeger DTA, van Putten SM, Bank RA. 3,4-Dihydroxy-L-Phenylalanine as a Novel Covalent Linker of Extracellular Matrix Proteins to Polyacrylamide Hydrogels with a Tunable Stiffness. Tissue Eng Part C Methods. 2016;22(2):91–101.

26. Denisin AK, Pruitt BL. Tuning the Range of Polyacrylamide Gel Stiffness for Mechanobiology Applications. ACS Appl Mater Interfaces. 2016;8(34):21893–902.

27. Zhang R, Li B, Li H. Extracellular-Matrix Mechanics Regulate the Ocular Physiological and Pathological Activities. J Ophthalmol. 2023;2023:1–11.

28. Qu Y, He Y, Zhang Y, Ma T, Zhu J, Miao Y, et al. Quantified elasticity mapping of retinal layers using synchronized acoustic radiation force optical coherence elastography. Biomed Opt Express. 2018;9(9):4054–63.

29. Kauppila M, Mörö A, Valle-Delgado JJ, Ihalainen T, Sukki L, Puistola P, et al. Toward Corneal Limbus In Vitro Model: Regulation of hPSC-LSC Phenotype by Matrix Stiffness and Topography During Cell Differentiation Process. Adv Healthc Mater. 2023;12(29):2301396.

30. Korkka I, Viheriälä T, Juuti-Uusitalo K, Uusitalo-Järvinen H, Skottman H, Hyttinen J, et al. Functional Voltage-Gated Calcium Channels Are Present in Human Embryonic Stem Cell-Derived Retinal Pigment Epithelium. Stem Cells Transl Med. 2019;8(2):179–93.

31. Peng S, Rao VS, Adelman RA, Rizzolo LJ. Claudin-19 and the Barrier Properties of the Human Retinal Pigment Epithelium. Investig Opthalmology Vis Sci. 2011;52(3):1392–403.

32. Tseng Q, Duchemin-Pelletier E, Deshiere A, Balland M, Guillou H, Filhol O, et al. Spatial organization of the extracellular matrix regulates cell–cell junction positioning. Proc Natl Acad Sci. 2012;109(5):1506–11.

33. Korkka I, Skottman H, Nymark S. Heterogeneity of Potassium Channels in Human Embryonic Stem Cell-Derived Retinal Pigment Epithelium. Stem Cells Transl Med. 2022;11(7):753–66.

34. Michael F. M. Mechanisms of Normal Retinal Adhesion. In: Retina. 5th ed. W. B: Saunders; 2013. p. 447–64.

35. Marmor MF, Abdul-Rahim AS, Cohen DS. The effect of metabolic inhibitors on retinal adhesion and subretinal fluid resorption. Invest Ophthalmol Vis Sci. 1980;19(8):893–903.

36. Yao XY, Hageman GS, Marmor F. Retinal Adhesiveness in the Monkey. Investig Opthalmology Vis Sci. 1994;35(2):744–8.

37. Yao XY, Endo EG, Marmor MF. Reversibility of Retinal Adhesion in the Rabbit. Investig Opthalmology Vis Sci. 1989;30(2):220–4.

38. Fietz A, Schnichels S, Hurst J. Co-cultivation of primary porcine RPE cells and neuroretina induces inflammation: a potential inflammatory AMD-model. Sci Rep. 2023 Nov 7;13(1):19345.

39. Gabriel E, Albanna W, Pasquini G, Ramani A, Josipovic N, Mariappan A, et al. Human brain organoids assemble functionally integrated bilateral optic vesicles. Cell Stem Cell. 2021;28(10):1740–57.

40. Chakrabarty K, Nayak D, Debnath J, Das D, Shetty R, Ghosh A. Retinal organoids in disease modeling and drug discovery: Opportunities and challenges. Surv Ophthalmol. 2024;69(2):179–89.

41. Zhao H, Yan F. Retinal Organoids: A Next-Generation Platform for High-Throughput Drug Discovery. Stem Cell Rev Rep. 2024;20(2):495–508.

42. Achberger K, Probst C, Haderspeck J, Bolz S, Rogal J, Chuchuy J, et al. Merging organoid and organ-on-a-chip technology to generate complex multi-layer tissue models in a human retina-on-a-chip platform. eLife. 2019;8:e46188.

43. Luo Z, Wang Y, Li J, Wang J, Yu Y, Zhao Y. Tailoring Hyaluronic Acid Hydrogels for Biomedical Applications. Adv Funct Mater. 2023;33(49).

44. Zeiger AS, Hinton B, Van Vliet KJ. Why the dish makes a difference: Quantitative comparison of polystyrene culture surfaces. Acta Biomater. 2013;9(7):7354–61.

45. Quinn RH, Miller SS. Ion Transport Mechanisms in Native Human Retinal Pigment Epithelium. Investig Opthalmology Vis Sci. 1992;33(13).

46. Tse JM, Cheng G, Tyrrell JA, Wilcox-Adelman SA, Boucher Y, Jain RK, et al. Mechanical compression drives cancer cells toward invasive phenotype. Proc Natl Acad Sci. 2012;109(3):911–6.

47. Tao J, Choudhury MI, Maity D, Kim T, Sun SX, Fan CM. Mechanical compression creates a quiescent muscle stem cell niche. Commun Biol. 2023;6(1):43.

48. Shroff NP, Xu P, Kim S, Shelton ER, Gross BJ, Liu Y, et al. Proliferation-driven mechanical compression induces signalling centre formation during mammalian organ development. Nat Cell Biol. 2024;26(4):519–29.

49. Mountcastle AM, Combes SA. Wing flexibility enhances load-lifting capacity in bumblebees. Proc R Soc B Biol Sci. 2013;280(1759).

50. Sancho A, Vandersmissen I, Craps S, Luttun A, Groll J. A new strategy to measure intercellular adhesion forces in mature cell-cell contacts. Sci Rep. 2017;7(1):46152.

51. Endo EG, Yao XY, Marmor MF. Pigment adherence as a measure of retinal adhesion: dependence on temperature. Invest Ophthalmol Vis Sci. 1988;29(9):1390–6.

52. Nandrot EF, Anand M, Sircar M, Finnemann SC. Novel role for αvβ5-integrin in retinal adhesion and its diurnal peak. Am J Physiol - Cell Physiol. 2006;290(4):1256–62.

53. Chiang RK, Yao XY, Takeuchi A, Dalal R, Marmor MF. Cytochalasin D reversibly weakens retinal adhesiveness. Curr Eye Res. 1995;14(12):1109–13.

54. Kita M, Marmor MF. Retinal Adhesive Force in Living Rabbit, Cat, and Monkey Eyes. Investig Opthalmology Vis Sci. 1992;33(6):1879–82.

55. Denisin AK, Kim H, Riedel-Kruse IH, Pruitt BL. Field Guide to Traction Force Microscopy. Cell Mol Bioeng. 2024;17(2):87–106.

56. Lieffrig SA, Gyimesi G, Mao Y, Finnemann SC. Clearance phagocytosis by the retinal pigment epithelial during photoreceptor outer segment renewal: Molecular mechanisms and relation to retinal inflammation. Immunol Rev. 2023;319(1):81–99.

57. Matsumoto B, Defoe DM, Besharse JC. Membrane turnover in rod photoreceptors: ensheathment and phagocytosis of outer segment distal tips by pseudopodia of the retinal pigment epithelium. Proc R Soc Lond B Biol Sci. 1987;230(1260):339–54.

58. Almedawar S, Vafia K, Schreiter S, Neumann K, Khattak S, Kurth T, et al. MERTK-Dependent Ensheathment of Photoreceptor Outer Segments by Human Pluripotent Stem Cell-Derived Retinal Pigment Epithelium. Stem Cell Rep. 2020;14(3):374–89.

59. Zihni C. Phagocytosis by the retinal pigment epithelium: New insights into polarized cell mechanics. BioEssays. 2025;47(1).

60. Finnemann SC, Bonilha VL, Marmorstein AD, Rodriguez-Boulan E. Phagocytosis of rod outer segments by retinal pigment epithelial cells requires avb5 integrin for binding but not for internalization. Proc Natl Acad Sci U S A. 94:12932–7.

61. Lock JG, Jones MC, Askari JA, Gong X, Oddone A, Olofsson H, et al. Reticular adhesions are a distinct class of cell-matrix adhesions that mediate attachment during mitosis. Nat Cell Biol. 2018 Nov;20(11):1290–302.

62. Roberts B, Haupt A, Tucker A, Grancharova T, Arakaki J, Fuqua MA, et al. Systematic gene tagging using CRISPR/Cas9 in human stem cells to illuminate cell organization. Drubin DG, editor. Mol Biol Cell. 2017;28(21):2854–74.

63. Thevenaz P, Ruttimann UE, Unser M. A Pyramid Approach to Subpixel Registration Based on Intensity. IEEE Trans Image Process. 1998;7(1):27–41.