Involvement of PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1 in COPII

assembly by interacting with SAR1 GTPase

Hui-Fang Lung[a], Jia-Dong Chu[a], Tzu-Yin Liu[a, b,1]

[a]Institute of Bioinformatics and Structural Biology, College of Life Sciences and Medicine,

National Tsing Hua University, Hsinchu 30013, Taiwan.

[b]Department of Life Science, College of Life Sciences and Medicine, National Tsing Hua

University, Hsinchu 30013, Taiwan.

[1]Address correspondence to tzliu@life.nthu.edu.tw

Institute of Bioinformatics and Structural Biology, College of Life Sciences and Medicine,

National Tsing Hua University, Hsinchu 30013, Taiwan

Tel: +886 3 574-2096, Fax: +886 3 571-5934

Corresponding author:

Tzu-Yin Liu

Email: tzliu@life.nthu.edu.tw

Abstract

Inorganic phosphate (Pi) uptake and translocation are crucial for plant growth and development, relying on plasma membrane targeting of PHOSPHATE TRANSPORTER1 (PHT1) transporters. The plant-specific endoplasmic reticulum (ER)-resident PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1 (PHF1) is structurally related to SEC12, which initiates the coat protein complex II (COPII) assembly as a guanine nucleotide exchange factor (GEF) by activating the small GTPase SAR1. In contrast, PHF1 loses the conserved catalytic residues critical for GEF activity and specifically assists the ER exit of the PHT1 transporters. However, the underlying molecular mechanism remains unknown. In this study, we showed that overexpression of *Arabidopsis thaliana* PHT1;1 (*At*PHT1;1) in the tobacco transient expression system caused a portion of *At*PHF1 distribution into *At*SAR1b- and *At*SEC24a-labeled ER exit sites. We demonstrated that *At*PHF1 interacts with *At*SAR1b and *At*SAR1c based on the tripartite split-GFP association in agro-infiltrated tobacco leaves and verified this interaction using miniTurbo-based proximity labeling. We also confirmed its physiological relevance by co-immunoprecipitation of the endogenous *At*PHF1 with *At*SAR1c-GFP in *Arabidopsis* transgenic lines. Importantly, *At*PHF1 preferentially interacts with the GDP-locked *At*SAR1. Therefore, we propose that PHF1 interacts with the SAR1 GTPase to participate in the early step of COPII recruitment for the ER export of PHT1 transporters.

Keywords: PHOSPHATE TRANSPORTER1, PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1, coat protein complex II (COPII), ER exit sites (ERES)

Introduction

As sessile organisms, plants adjust the protein and lipid composition of the plasma membrane (PM) to sense and adapt to the ever-changing environment. Such dynamic cellular processes heavily rely on the modulation of the endomembrane trafficking system (Wang et al., 2020), including the coat protein complex II (COPII)-mediated endoplasmic reticulum (ER)-to-Golgi transport pathway. Although the core COPII machinery, composed of five cytosolic factors, SAR1, SEC23, SEC24, SEC13, and SEC31, is conserved throughout all eukaryotes, the mechanisms underlying COPII recruitment and assembly are most extensively studied in the budding yeast *Saccharomyces cerevisiae* (Barlowe and Miller, 2013). In this model organism, the COPII-related components are encoded by single genes and recruited sequentially to the ER exit sites (ERES) (Kurokawa and Nakano, 2019). The SAR1 GTPase is activated upon the exchange of GDP to GTP by the guanine nucleotide exchange factor (GEF) SEC12, exposing its amphipathic N-terminal helix to be inserted into the ER membrane (d'Enfert, 1991; Paul et al., 2023). SAR1 then interacts directly with SEC23, inducing the formation of the inner coat SEC23–SEC24 complex, which in turn recruits the outer coat SEC13–SEC31 complex (Bi, 2002; Stagg et al., 2006). In addition, SEC16, an initial factor assembled at the ERES, serves as a scaffold protein by concentrating SEC12 and GTP-bound active SAR1 in proximity (Supek et al., 2002; Montegna et al., 2012). By comparison, the plant COPII subunit paralogs are encoded by multi-gene families and outnumber those in other organisms (Chung et al., 2016). Multiple COPII paralogues allow the diversity of COPII transport carriers to adapt to various developmental and stress-related cues (Chung et al., 2016; Wang et al., 2020). For instance, the phytohormone abscisic acid triggered the formation of *Arabidopsis thaliana* SAR1a (*At*SAR1a)-dependent COPII vesicles that carry osmotic stress-related carriers (Li, 2021). The *At*SAR1a-*At*SEC23a pairing was also shown to mediate the ER export of the ER-associated transcription factor bZIP28, which is upregulated in response to ER stress (Zeng et al., 2015).

By contrast, the mechanism by which the COPII machinery regulates protein ER export to cope

with nutrient deficiency remains unknown.

Inorganic phosphate (Pi) is an essential macronutrient for plant growth. Due to its poor

solubility and mobility in soil, Pi bioavailability is limited, thus causing plants to face Pi

deficiency (Shen, 2011). Under Pi-limited conditions, plants enhance the external Pi uptake as

well as the recycling and remobilization of internal Pi, which involves the upregulation of the

PHOSPHATE TRANSPORTER1 (*PHT1*) gene family at the transcript and post-transcriptional

level (Nussaume et al., 2011). While most *PHT1* genes are upregulated by Pi deprivation, the

newly synthesized PHT1 proteins (PHT1s) must exit the ER for the PM targeting. Similar to

the inhibitory effect of SEC12 overexpression on the ER export of the Golgi transmembrane

protein ERD2 in tobacco leaves (Hanton et al., 2007), the overexpression of SEC12 impaired

the PM targeting of *At*PHT1;2 (Bayle et al., 2011). Thus, the ER-to-Golgi transport of PHT1s

is dependent on efficient COPII assembly. Interestingly, the plant-specific SEC12-related

protein PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1 (PHF1) was identified to

facilitate the ER exit of *At*PHT1s (González et al., 2005; Bayle et al., 2011). Despite sharing

the sequence similarity with SEC12, *At*PHF1 lost the conserved catalytic residues critical for

GEF activity and failed to rescue the growth defect of the yeast *sec12* mutant (González et al.,

2005). Instead, the *Arabidopsis* loss-of-function *phf1* mutant showed decreased cellular Pi

content and impaired PM targeting of *At*PHT1;1 (González et al., 2005). As the role of PHF1

in assisting the ER exit of PHT1s resembles that of the yeast Pho86p in the ER exit of Pho84p,

the yeast homolog of PHT1s (Lau, 2000), PHF1 may functionally diverge from SEC12

(González et al., 2005).

The transient protein expression using tobacco leaves is a well-established plant system to study

the dynamic organization of COPII proteins at the ERES (daSilva et al., 2004; Hanton et al.,

2009). In this system, the ERES formation can be monitored by the subcellular distribution of

fluorescent protein-tagged COPII components changing from cytosolic patterns into punctate

structures (daSilva et al., 2004; Hanton et al., 2007; Hanton et al., 2008; Hanton et al., 2009).

Overexpression of COPII-dependent membrane cargoes enhanced the recruitment of YFP-

*At*SEC24a to the ERES as well as induced the *de novo* formation of ERES (Hanton et al., 2007).

Likewise, *At*PHT1;2-CFP overexpression resulted in the recruitment of YFP-*At*SEC24a to

ERES puncta, reinforcing that the ER export of *At*PHT1s is COPII-dependent (Bayle et al.,

2011). Although the ER-resident *At*PHF1 failed to co-localize with *At*SEC24a, thus leading to

the conclusion that *At*PHF1 is not involved in COPII recruitment (Bayle et al., 2011), it remains

unclear whether PHF1 participates in cargo recognition or the selective packaging of PHT1

transporters into COPII vesicles.

Since PHF1 shares sequence similarity with SEC12 yet exhibits a distinct function, we

speculated that PHF1 may engage in the COPII-dependent ER export of PHT1s on a molecular

basis different from that of SEC12. To explore this hypothesis, we aimed to investigate whether

*At*PHT1;1 overexpression can induce the distribution of *At*PHF1 into ERES puncta and whether

*At*PHF1 can interact with COPII-related components. Using agro-infiltrated tobacco leaves, we

showed that overexpression of *At*PHT1;1 triggered a portion of *At*PHF1 distribution into

punctate structures that partially co-localized with *At*SAR1b and *At*SEC24a. Furthermore,

based on the tripartite split-GFP association, *At*PHF1 interacted with *At*SAR1b and *At*SAR1c,

but not with *At*SEC24a or other COPII-related components. Consistently, we verified the

interaction of *At*PHF1 and *At*SAR1b by the proximity labeling in the transient tobacco system

and demonstrated that the endogenous *At*PHF1 and *At*PHT1;1/2/3 were co-immunoprecipitated

with *At*SAR1c-GFP in *Arabidopsis* transgenic lines. More importantly, we showed that, like

*At*SEC12, *At*PHF1 preferentially interacted with the GDP-locked inactive form of *At*SAR1b.

Our results unveiled that *At*PHF1 may act as an early regulator of COPII assembly for the ER

export of *At*PHT1s through interacting with *At*SAR1 proteins (*At*SAR1s).

Results

Overexpression of *At*PHT1;1 triggers partial distribution of *At*PHF1 into ERES-associated punctate structures

Both the mammalian and plant SEC12 proteins are dispersed throughout the ER (Weissman et al., 2001; Bayle et al., 2011). However, the mammalian SEC12 can be concentrated at the ERES by the collagen cargo receptor component cTAGE5 for the ER export of collagen (Saito et al., 2014). We thus wondered whether the SEC12-related *At*PHF1, which was reported to distribute evenly across the ER (González et al., 2005; Bayle et al., 2011), can be induced to concentrate at ERES when the ER export of *At*PHT1;1 is highly demanded. When expressed alone in agro-infiltrated tobacco leaves, *At*PHF1-GFP showed a reticular ER pattern, while a portion of *At*PHF1-GFP displayed punctate-like structures when the split-GFP tagged *At*PHT1;1 (*At*PHT1;1-S11) was co-expressed (**Fig. 1A**). By contrast, the co-expression of *At*PHF1-GFP with the sugar transporter *At*STP1-S11 barely changed the distribution of *At*PHF1-GFP (**Fig. 1A**). Statistical analysis further suggested that the overexpression of *At*PHT1;1-S11 but not of *At*STP1-S11 significantly triggered the partial distribution of *At*PHF1-GFP into punctate structures (**Fig. 1B**).

To ensure that the formation of *At*PHF1 puncta triggered by *At*PHT1;1 overexpression is not due to the ER morphological changes, we examined the ER organization using the ER lumen marker mCherry-HDEL. The overexpression of *At*PHT1;1-S11 did not change the overall ER morphology labeled by the mCherry-HDEL marker (**Supplementary Fig. S1**), but we occasionally observed the formation of mCherry-HDEL- or *At*PHF1-GFP-labeled puncta upon the overexpression of *At*PHT1;1 (**Fig. 1C**). We then classified the *At*PHT1;1-induced puncta into three groups for quantification: the distinct *At*PHF1-labeled puncta that did not overlap with the mCherry-HDEL puncta, the distinct mCherry-HDEL-labeled puncta that did not overlap with the *At*PHF1-labeled puncta, and the overlapping puncta. Statistical analyses indicated that the distinct *At*PHF1-labeled puncta predominated over the overlapping puncta (**Fig. 1D**; detailed analysis of individual experiment in **Supplementary Fig. S2A**), suggesting

that the majority of the *At*PHT1;1 overexpression-induced *At*PHF1 puncta represented a compartment different from the mCherry-HDEL-labeled puncta.

In the tobacco transient expression system, the co-expression of membrane cargoes recruited the COPII components, such as *At*SAR1 and *At*SEC24, to concentrate at the punctate ERES (Hanton et al., 2007; Hanton et al., 2008). Intriguingly, the populations of *At*PHF1 puncta upon *At*PHT1;1 overexpression displayed a size with a median of 0.66–0.77 μm (**Supplementary Fig. S2B**), which is comparable to the previously reported size of ERES puncta with a range of 0.15–0.8 μm across various organisms (Bannykh, 1996; Yang et al., 2005; Zeuschner et al., 2006; Takagi et al., 2020; Weigel et al., 2021). To address whether the punctate structures of *At*PHF1-GFP induced by the overexpression of *At*PHT1;1 might represent ERES, we selected *At*SAR1b as a COPII marker to co-localize with *At*PHF1-mCherry. This is because *AtSAR1b* is the most highly expressed *AtSAR1* gene in roots and is upregulated in response to Pi deprivation (Liu et al., 2016) (**Supplementary Fig. S3**). In addition, the previous proteomic analysis of the Pi overaccumulator *pho2* root has revealed the co-induction of *At*SAR1b and *At*PHT1s (Huang et al., 2013), implying an increased demand for *At*SAR1b to promote the ER-to-Golgi traffic of *At*PHT1s. Considering that the N-terminal tail of SAR1 is inserted into the ER membranes upon activation (d'Enfert, 1991; Paul et al., 2023), we fused the S11 tag to the C terminus of *At*SAR1b to prevent steric hindrance. In tobacco epidermal cells, *At*SAR1b-S11 exhibited both a diffuse cytosolic distribution and an ER membrane-associated punctate pattern, similar to the previously reported results (Yang et al., 2005; Hanton et al., 2008) (**Supplementary Fig. S4A**). *At*SEC24a is involved in cargo selection for COPII vesicles and is also widely used as a reliable ERES marker (Miller, 2002; Hanton et al., 2007); therefore, we generated a construct expressing *At*SEC24a-S11 for co-localization with *At*PHF1-mCherry as well. The subcellular localizations of *At*SEC24a-S11 displayed the cytosolic and punctate patterns (**Supplementary Fig. S4A**). *At*PHT1;1 overexpression induced the formation of *At*PHF1-mCherry punctate structures that colocalized with the *At*SAR1b- and *At*SEC24a-labeled ERES (**Fig. 2A** and **2B**).

Statistical analyses further suggested that the majority of *At*PHF1-mCherry puncta overlapped

with these two ERES markers (**Fig. 2C**; detailed analysis of individual experiment in

Supplementary Fig. S5A and **S5B**). The size of the overlapping puncta was also comparable

to that of the distinct *At*SAR1b- and *At*SEC24a-labeled puncta, with a median of 0.67–0.81 μm

(**Supplementary Fig. S5C** and **S5D**). The co-localization analysis further indicated a better

spatial co-occurrence of *At*PHF1-mCherry puncta with *At*SAR1b than with *At*SEC24a, as

determined by a significantly higher Manders' overlap coefficient for *At*SAR1b (**Fig. 2D**).

Taken together, these results suggested that when *At*PHT1;1 is overexpressed as a COPII-

dependent export membrane cargo, *At*PHF1 can partially distribute into puncta that are

associated with ERES markers.

The interaction of *At*PHF1 with *At*SAR1b/c *in planta* based on tripartite split-GFP

association

As *At*PHT1;1 overexpression triggered partial distribution of *At*PHF1 into punctate structures

associated with *At*SAR1b and *At*SEC24a, we postulated that *At*PHF1 may participate in COPII

recruitment or assembly. To explore this possibility, we examined the interaction of *At*PHF1

with several COPII-related components based on the tripartite split-GFP association in agro-

infiltrated tobacco leaves (Cabantous et al., 2013; Liu et al., 2018). The expression of split-GFP

tagged *At*SEC12 and *At*PHF1 (S10-*At*SEC12 and S10-*At*PHF1) confirmed their localization at

the ER when co-expressed with the cytosolic S11-GFP1–9 (**Supplementary Fig. S4B**). We also

generated constructs expressing the split-GFP-tagged *At*SAR1c, *At*SEC13a, and *At*SEC16a

(*At*SAR1c-, *At*SEC13a-, and *At*SEC16a-S11). As *AtSAR1c* is the second most highly expressed

AtSAR1 gene in roots (Liu et al., 2016) (**Supplementary Fig. S3**) and plays an interchangeable

role with *AtSAR1b* at the protein level in pollen development (Liang et al., 2020), we included

*At*SAR1c-S11 for comparison with *At*SAR1b. As previously reported (Hanton et al., 2008;

Hanton et al., 2009; Takagi et al., 2013), *At*SAR1c-, *At*SEC13a-S11, and *At*SEC16a-S11

predominantly localized to the cytosol and ERES puncta (**Supplementary Fig. S4A**). The expression of 3xHA-S11 or S10-3xHA exhibited cytosolic patterns (**Supplementary Fig. S4**) and was used as a negative control for the interaction (**Fig. 3A–C**). The interaction of S10-*At*SEC12 with *At*SAR1b- or *At*SAR1c-S11 was used as a positive control, yielding the GFP complementation signals (**Fig. 3A**). Notably, like S10-*At*SEC12, S10-*At*PHF1 interacted with *At*SAR1b- and *At*SAR1c-S11 but not with *At*SEC24a-, *At*SEC13a-, and *At*SEC16a-S11 (**Fig. 3B**), suggesting the specificity of *At*PHF1 interaction with *At*SAR1s among the COPII-related proteins.

The interaction of *At*PHF1 with *At*SAR1b *in planta* based on miniTurbo (mTb) proximity labeling

The tripartite split-GFP association detects protein-protein interaction without spurious background signals even when bait and prey proteins are overexpressed (Cabantous et al., 2013; Liu et al., 2018). Still, we employed the proximity labeling method as a complementary approach to validate the *in planta* interaction between *At*PHF1 and *At*SAR1b. Proximity labeling begins after the application of biotin and can be reversibly halted by removing biotin, thereby allowing the capture of transient and dynamic protein-protein interactions at a ten-nanometer scale (Kim et al., 2014; Xu et al., 2023). Due to the superior feasibility of miniTurbo (mTb) in the tobacco system (Mair et al., 2019), the mTb-based proximity labeling was used. For unknown reasons, our initial attempt to express mTb-EYFP-*At*PHF1 in agro-infiltrated tobacco leaves was unsuccessful, so we generated the *At*SAR1b-mTb-S11 construct by fusing mTb to *At*SAR1b (**Fig. 4A**). We used the cytosol-localized mTb-NES-EYFP (**Fig. 4B**), which carries the nuclear export sequence (NES), as a negative control (Mair et al., 2019). Taking advantage that S11 can self-complement with GFP1–10, we could detect the expression and subcellular distribution of *At*SAR1b-mTb-S11 in agro-infiltrated tobacco leaves (**Fig. 4B**). We then co-expressed mTb-NES-EYFP or *At*SAR1b-mTb-S11 with *At*SEC12-mCherry or

*At*PHF1-mCherry and incubated the harvested leaves in a biotin solution, followed by tissue

homogenization and protein extraction. The protein extraction of non-infiltrated tobacco leaves

was used for the background comparison. As the expression of the cytosolic mTb-NES-EYFP

was much greater than the membrane-associated *At*SAR1b-mTb-S11, we adjusted the ratio of

protein amount for *At*SAR1b-mTb-S11: mTb-NES-EYFP to 30: 1 (**Fig. 4C**; left panel). In this

way, the protein amounts of *At*SAR1b-mTb-S11 and mTb-NES-EYFP were comparable,

allowing them to fairly compete with binding to streptavidin beads. When co-expressed with

*At*SAR1b-mTb-S11, both *At*SEC12-mCherry and *At*PHF1-mCherry could be captured by

streptavidin and detected by immunoblot, indicating that they were biotinylated (**Fig. 4C**; left

panel). By contrast, the co-expression of mTb-NES-EYFP with *At*SEC12-mCherry or *At*PHF1-

mCherry did not yield similar effects (**Fig. 4C**; left panel). Consistently, when equal protein

amounts were used for the experiments, even though the expression of *At*SAR1b-mTb-S11 was

about five times lower than that of mTb-NES-EYFP, about two times more *At*SEC12-mCherry

or *At*PHF1-mCherry was captured by streptavidin when co-expressed with *At*SAR1b-mTb-S11

(**Fig. 4C**; right panel). These results again suggested that *At*PHF1 interacts with *At*SAR1s *in*

planta.

To determine which region of *At*PHF1 interacts with *At*SAR1s, we generated four N-terminally

S10-tagged *At*PHF1 truncated variants as follows: the cytoplasmic domain alone (*At*PHF1 N),

the cytoplasmic domain with the transmembrane (TM) domain (*At*PHF1 N–TM), the TM

domain alone (*At*PHF1 TM), and the TM domain with the ER-luminal domain (*At*PHF1 TM–

C) (**Supplementary Fig. S6A**). The expression and subcellular distribution of these *At*PHF1

variants suggested that the S10-*At*PHF1 N variant was present in the cytosol and nucleus, while

the other truncated forms of *At*PHF1 localized to the ER (**Supplementary Fig. S6B**). The

subcellular localization of the C-terminally S11-tagged truncated forms of *At*PHF1 also showed

similar results (**Supplementary Fig. S6C**), suggesting that the TM domain of *At*PHF1 alone is

sufficient for its ER targeting. Of note, the interaction of S10-tagged *At*PHF1 N or *At*PHF1 N–

TM with *At*SAR1b- or *At*SAR1c-S11 displayed both the ER and puncta patterns

(**Supplementary Fig. S7A** and **S7B**). By contrast, the interaction of S10-tagged *At*PHF1 TM

or *At*PHF1 TM–C with *At*SAR1b- or *At*SAR1c-S11 additionally resulted in a few GFP signals

that did not display a typical reticular ER pattern (**Supplementary Fig. S7A** and **S7B**). These

observations suggest that both the cytosolic and the transmembrane domains of *At*PHF1

contribute to the interaction with *At*SAR1s at the ER.

Co-immunoprecipitation of the endogenous *At*PHF1 with *At*SAR1c-GFP in *Arabidopsis*

root

To determine whether the interaction of *At*PHF1 and *At*SAR1s is physiologically relevant, we

used *Arabidopsis* transgenic plants expressing *At*SAR1c-GFP or GFP under the UBQ10

promoter for co-IP analysis (Zeng et al., 2015). The *Arabidopsis* seedlings were subjected to

seven days of Pi starvation to mimic the physiological conditions in which the endogenous

*At*PHF1 is upregulated and the ER export of *At*PHT1 proteins is also highly demanded. A total

root protein extract was used for GFP-Trap co-IP, followed by immunoblot analysis using an

anti-*At*PHF1 antibody. Due to the differences in the expression level between the cytosolic GFP

and the membrane-associated *At*SAR1c-GFP in transgenic plants, we increased the protein

amount for the co-IP of *At*SAR1c-GFP by 50-fold compared to that for the co-IP of GFP.

Immunoblot results suggested that the endogenous *At*PHF1 was co-immunoprecipitated with

*At*SAR1c-GFP but not with GFP (**Fig. 5A**). Conversely, when we used equal amounts of root

total proteins for co-IP analysis, the expression of *At*SAR1c-GFP was ten times lower than that

of GFP but was co-immunoprecipitated about three times more endogenous *At*PHF1 (**Fig. 5B**),

suggesting that *At*SAR1c-GFP exhibits a higher binding affinity toward *At*PHF1. Moreover, the

endogenous *At*PHT1;1/2/3 proteins could also be co-immunoprecipitated with *At*SAR1c-GFP

(**Fig. 5A** and **5B**), indicating the presence of a protein complex containing *At*SAR1s, *At*PHF1,

and *At*PHT1s. These data supported the notion that through interacting with *At*SAR1s, *At*PHF1

participates in the initial phase of COPII assembly for the ER export of *At*PHT1s.

AtPHF1 preferentially interacts with the GDP-locked inactive form of AtSAR1b

Because the assembly and disassembly of COPII is controlled by the SAR1 GTPase cycle (Van

der Verren and Zanetti, 2023), we next asked whether *At*PHF1 interacts with *At*SAR1s in a

specific nucleotide-binding state. Confocal analysis suggested that both the wild-type

*At*SAR1b-S11 and the GTP-locked *At*SAR1b[H74L]-S11 localized to the ER membrane and

cytosol as previously reported (daSilva et al., 2004; Wei and Wang, 2008) (**Supplementary Fig.**

S4A and **S8A**). The GDP-locked *At*SAR1b[T34N]-S11 also displayed a similar subcellular

pattern (**Supplementary Fig. S8A**). Furthermore, to quantify the soluble and membrane-

associated populations of the *At*SAR1b variants, we applied biochemical fractionation.

Although the microsomal fraction was slightly contaminated with the soluble fraction, as

indicated by the detection of a small amount of cytosolic FBPase, the majority of the *At*SAR1b

proteins were membrane-associated (**Supplementary Fig. S8B**). Namely, the membrane-

associated proportions of *At*SAR1b-S11, *At*SAR1b[H74L]-S11, and *At*SAR1b[T34N]-S11

were 76.4%, 96.1%, and 67.7%, respectively, suggesting that the GDP-locked form of *At*SAR1b

is relatively less membrane-associated (**Supplementary Fig. S8B**). These results were in

agreement with the previous finding that upon binding of GTP, SAR1 is activated to insert into

the ER membrane (d'Enfert, 1991; Paul et al., 2023). More importantly, the *in-planta* tripartite

split-GFP assay suggested that, like *At*SEC12, *At*PHF1 preferentially interacts with the GDP-

locked form of *At*SAR1b (**Fig. 6A** and **6B**). The interaction of *At*PHF1 N with the *At*SAR1b or

the GDP-locked *At*SAR1b[T34N] showed cytoplasmic, ER, and puncta patterns (**Fig. 6B**). By

contrast, the interaction of the *At*PHF1 N–TM, *At*PHF1 TM, or *At*PHF1 TM–C variants with

*At*SAR1b or the GDP-locked *At*SAR1b[T34N] showed a reticular ER pattern, and occasionally

punctate structures at the ER (**Fig. 6B**). Interestingly, the interactions of the *At*PHF1 TM or the

*At*PHF1 TM–C with the GTP-locked *At*SAR1b[H74L] were detected at both the ER and the

non-ER compartments (**Fig. 6B**). Overall, the GTP-locked *At*SAR1b[H74L] drastically reduced

the interaction with the full-length *At*PHF1, *At*PHF1 N and *At*PHF1 N–TM variants (**Fig. 6B**).

Compared to the *At*SAR1b, the GTP-locked *At*SAR1b[H74L] slightly increased the interaction

with the *At*PHF1 TM and the *At*PHF1 TM–C, which was likely due to a better membrane

association of *At*SAR1b[H74L] (**Fig. 6B** and **Supplementary Fig. S8B**). We interpret these as

indicating that the removal of the N-terminal cytosolic domain of *At*PHF1 allows a non-specific

interaction of *At*PHF1's transmembrane domain with the GTP-locked *At*SAR1b[H74L]. Taken

together, these results indicate that the *At*PHF1 variants containing the cytosolic domain

preferentially associate with the GDP-bound *At*SAR1 and thus likely participate in recruiting

COPII components to ERES. To further support this finding, we performed co-IP assays in

transgenic plants expressing *dex:At*SAR1b[H74L]-CFP (Lupanga et al., 2020). Upon

dexamethasone (DEX) induction, *At*SAR1b[H74L]-CFP was expressed and detected in the root

total protein extracts (**Fig. 6C**). When equal amounts of expressed *At*SAR1c-GFP and GTP-

locked *At*SAR1b[H74L]-CFP proteins were used for IP analysis, four times more endogenous

*At*PHF1 was co-immunoprecipitated with *At*SAR1c-GFP than with *At*SAR1b[H74L]-CFP (**Fig.**

6C). These results suggest that GTP-locked *At*SAR1b[H74L]-CFP has a lower binding affinity

than *At*SAR1c-GFP toward *At*PHF1 in *Arabidopsis* roots.

Discussion

Although several ER accessory proteins have been identified as participating in the ER exit of

membrane proteins, the underlying molecular mechanisms are largely unclear (Tang et al., 2005;

Bisnett et al., 2020). In *Arabidopsis*, AUXIN RESISTANCE4 (AXR4) was shown to interact

with auxin influx carriers AUX1 and LAX2 to promote their PM targeting (Dharmasiri, 2006;

Tidy et al., 2024). The KAONASHI3 (KNS3) family was recently identified to mediate the ER

exit of the boric acid channel *At*NIP5;1 (Zhang et al., 2024). While *At*PHF1 is required for the

PM targeting of *At*PHT1s, it was proposed to act as an ER-localized chaperone for the protein

stability of *At*PHT1s (González et al., 2005) and not participate in the COPII vesicle formation

(Bayle et al., 2011). On the contrary, our data revealed a role for *At*PHF1 in the early step of

COPII assembly through the interaction with *At*SAR1s. Using three different methods—

tripartite split-GFP complementation, proximity labeling, and co-immunoprecipitation—we

demonstrated that *At*PHF1 interacts with *At*SAR1s *in planta* (**Fig. 3**, **4**, and **5**). Furthermore, the

tripartite split-GFP and the co-IP assays demonstrated that *At*PHF1 preferentially interacts with

the GDP-bound *At*SAR1s (**Fig. 6**). Our findings uncover that *At*PHF1 likely mediates the

assembly of COPII vesicles through interacting with *At*SAR1s, thereby facilitating the ER

export of *At*PHT1s (**Fig. 7**). The recruitment of COPII to ERES is increased in response to the

demand for ER export of membrane cargoes (Hanton et al., 2007). Our results showed that the

transient overexpression of *At*PHT1;1 in *N. benthamiana* leaves also triggered partial

distribution of *At*PHF1 into the punctate structures associated with the ERES markers *At*SAR1b

and *At*SEC24a (**Fig. 1** and **Fig. 2**). In other words, under an increased demand for the COPII-

mediated export of *At*PHT1s, *At*PHF1 may partially relocate to ERES to promote this process.

The yeast ER membrane-localized chaperone Shr3 is specifically required for proper folding of

amino acid permeases and their targeting to the plasma membrane (Kota and Ljungdahl, 2005).

Interestingly, the amino acid permease Gap1 split into N- and C-terminal portions can be

assembled to form a functional permease in an Shr3-dependent manner (Kota et al., 2007). We

also generated the N- and C-terminal portions of *At*PHT1;1 (*At*PHT1;1 TM1–6 half and the

*At*PHT1;1 TM7–12 half) to test whether *At*PHF1 plays a critical role in enabling *At*PHT1;1 to

fold and attain a proper structure competent for ER export. Even upon overexpression of

*At*PHF1, both the *At*PHT1;1 N and C-halves were unable to exit the ER (**Supplementary Fig.**

S9A). In this scenario, neither the N-half nor the C-half of *At*PHT1;1 resulted in a partial

distribution of *At*PHF1 to punctate structures (**Supplementary Fig. S9B** and **S9C**). This

indicates that partial recognition of *At*PHT1;1 by *At*PHF1 is insufficient; instead, *At*PHT1;1

must attain a functional conformation that enables it to interact properly with *At*PHF1, thereby

triggering a signal for ER export.

Although *At*PHF1 lacks the conserved residues required for GEF activity and cannot

complement the yeast *sec12* growth defect (González et al., 2005), we surmised that upon

*At*PHT1;1 overexpression, the distribution of *At*PHF1 to the ERES may facilitate the COPII

recruitment or assembly. We reasoned that without co-expression of *At*PHT1s, observing partial

co-localization of *At*PHF1 with ERES is beyond the detection limit of confocal imaging, which

may explain the discrepancy between our results and those from a previous study (Bayle et al.,

2011). In support of this notion, we demonstrated that endogenous *At*PHF1 and *At*PHT1;1/2/3

proteins were induced and co-immunoprecipitated with *At*SAR1c-GFP in transgenic plants

subjected to Pi starvation (**Fig. 5**), suggesting the physiological relevance of the interaction

between *At*PHF1 and *At*SAR1s. In addition, *At*PHF1 interacted with *At*SAR1b and *At*SAR1c

but not with other COPII-related components (**Fig. 3B**), and preferentially interacted with the

GDP-locked form of *At*SAR1 *in planta* (**Fig. 6**), indicating that *At*PHF1 participates in the early

step of COPII assembly, thus likely indirectly assisting the loading of *At*PHT1s into COPII

vesicles for the ER export.

For unknown reasons, we failed to detect the interaction of *At*PHF1 or *At*SEC12 with *At*SAR1s

in the yeast split-ubiquitin system (**Supplementary Fig. S10A** and **S10B**). Similarly, our

attempt to compare the interaction strength of the cytosolic N-terminal domain of *At*PHF1 with

the GDP- or GTP-bound *At*SAR1b using yeast two-hybrid assays was unsuccessful

(**Supplementary Fig. S11**). The failure to detect the interaction between *At*SEC12 or *At*PHF1

with *At*SAR1s in yeast may reflect the limitations of the heterologous system. Nevertheless,

our findings indicated that *At*PHF1 could link the specific recognition of *At*PHT1s with the

modulation of *At*SAR1s activity for the COPII assembly. Such a mechanism differs from that

reported for SED4, the yeast structural homolog of SEC12, which also lacks GEF activity

(Gimeno, 1995; Kodera et al., 2011). Genetic and biochemical evidence suggested that SED4

and SEC12 are not functionally interchangeable and that SED4 weakly inhibits the GTPase-

activating (GAP) activity of SEC23 toward SAR1 (Saito-Nakano and Nakano, 2000). On the

contrary, an *in vitro* study showed that SED4 had stronger binding to the non-hydrolyzable GTP

analog GMPPNP-bound SAR1 and stimulated SEC23 GAP activity, thereby accelerating the

dissociation of COPII coats from the ER membrane (Gimeno, 1995; Kodera et al., 2011). While

the interaction between SED4 and SEC16 is required for the localization of SEC16 to ERES

(Gimeno, 1995), the interaction between *At*PHF1 and *At*SEC16a was not observed (**Fig. 3B**).

As many COPII-related factors regulate SAR1 activity through direct or indirect interaction

(Van der Verren and Zanetti, 2023), we propose that PHF1 modulates the recruitment efficiency

of SAR1s to ERES, thereby facilitating the formation of COPII vesicles to carry PHT1s (**Fig.**

7). Nevertheless, we do not exclude the possibility that PHF1 plays dual roles, acting as a

membrane chaperone for the correct folding of newly synthesized PHT1s in the ER and also

mediating COPII assembly. Without uncoupling these two potential functions of PHF1, its role

as a chaperone at an earlier step will mask the involvement of PHF1 in the re-localization of

PHT1s to ERES at a later step.

Interestingly, we recently reported that the *Arabidopsis* CORNICHON HOMOLOG 5

(*At*CNIH5) acts as an ER cargo receptor that cycles between the ER and the early Golgi and

selects *At*PHT1s for ER export (Chiu, 2024; Liu et al., 2025). As the *cnih5 phf1* mutant shows

more severe growth defects than the *phf1* single mutant, and *At*PHF1 proteins are increased in

the root of *cnih5* (Chiu, 2024), we suspected that *At*PHF1 and *At*CNIH5 work interdependently.

Namely, while *At*PHF1 couples the recognition of *At*PHT1s and the recruitment of COPII

proteins by interacting with *At*PHT1s and *At*SAR1s, *At*CNIH5 is involved in the selective

packaging of *At*PHT1 into COPII vesicles.

*At*PHF1 is structurally related to SEC12 proteins, which adopt a seven-bladed β propeller

(González et al., 2005; Joiner and Fromme, 2021). Based on the crystal structure, the

catalytically critical K loop within the cytoplasmic domain of yeast SEC12 makes direct contact

with nucleotide-free SAR1 (McMahon et al., 2012; Joiner and Fromme, 2021). By contrast, our

data showed that both the cytosolic and the transmembrane domains of *At*PHF1 interact with

*At*SAR1b/c *in planta* (**Supplementary Fig. S7A** and **S7B**), indicating that the interaction

interface may span relatively large areas. Since the catalytic residues required for GEF activity

are not conserved in *At*PHF1 (González et al., 2005), it remains to be elucidated whether the

interaction interface of PHF1-SAR1s is different from that of SEC12-SAR1s and how PHF1

modulates the recruitment of SAR1s without GEF activity.

Materials and Methods

Plant material and growth conditions

The seeds of *Arabidopsis thaliana* wild-type (WT) (Columbia, Col-0) and *phf1* were obtained

from the Arabidopsis Biological Resource Center (ABRC). Seeds of *A. thaliana* were surface-

sterilized and germinated on half-strength modified Hoagland agar plates (Millner, 1992; Liu

et al., 2012) and grown in the growth chamber at 22°C under a 16-h-light/8-h-dark cycle. The

Pi-sufficient (+P) and Pi-deficient (−P) media were half-strength modified Hoagland nutrient

solutions containing 250 μM and 0 μM KH_2PO_4, respectively. Seeds of *Nicotiana benthamiana*

(*N. benthamiana*) were surface-sterilized and germinated on half-strength Murashige and

Skoog medium and grown in the growth chamber at 24°C under a 16-h-light/8-h-dark cycle.

Plasmid construction

For the self-complementing split-GFP assay (Liu et al., 2018), the constructs encoding

*UBQ10:sXVE:*GFP1–10 (Addgene plasmid #125663), *UBQ10:sXVE:*ER-GFP1–10 (Addgene

plasmid #125664) or *UBQ10:sXVE:*S11-GFP1–9 (Addgene plasmid #125665) was used. For

the tripartite split-GFP association, the constructs encoding *UBQ10:sXVE:*GFP1–9 (Addgene

plasmid #108187) or *35S:*GFP1–9.pMDC201 were used. The CDS of GFP1–9 was amplified

by PCR using *P16ΔS:*GFP1–9 (Addgene plasmid #108186) as the template and subcloned into

pMDC201, yielding *35S:*GFP1–9.pMDC201. For constructing the S10-tagged clones, the CDS

of full-length and truncated *At*PHF1 and *At*PHT1;1 were amplified by PCR and subcloned into

*UBQ10:XVE:*S10-(MCS) (Addgene plasmid #108177), in which the inducible promoter

UBQ10:sXVE was replaced by the *UBQ10* promoter to generate *UBQ10:*S10-(MCS). The CDS

of *At*SEC12 and *At*PHF1 were amplified by PCR and subcloned into *UBQ10:*S10-(MCS). For

constructing the S11-tagged clones, the CDS of *At*PHT1;1 was amplified by PCR and subcloned

into *UBQ10:sXVE:*(MCS)-3xHA-S11 (Addgene plasmid #108180). The CDS of *At*STP1 and

*At*PHT1;1 and truncated *At*PHF1 were amplified by PCR and subcloned into

UBQ10:sXVE:(MCS)-S11 (Addgene plasmid #108179), in which the inducible promoter

UBQ10:sXVE was replaced by the *UBQ10* promoter to generate *UBQ10:*(MCS)-S11. The CDS

of *At*SAR1b, *At*SAR1b[H74L] (Takeuchi, 2000), *At*SAR1b[T34N], *At*SAR1c, *At*SEC24a,

*At*SEC13a, and *At*SEC16a were amplified and subcloned into *UBQ10:*(MCS)-S11. For

constructing the GFP fusion clones, the inducible promoter *UBQ10:sXVE* of

UBQ10:sXVE:(MCS)-GFP (Schlücking et al., 2013) was replaced by the *UBQ10* promoter to

generate *UBQ10:*(MCS)-GFP. The CDS of *At*PHF1 was amplified and subcloned into

UBQ10:(MCS)-GFP. For constructing the mCherry fusion clones, the GFP of

UBQ10:sXVE:(MCS)-GFP (Schlücking et al., 2013) was replaced by mCherry to generate

UBQ10:sXVE:(MCS)-mCherry. The CDS of *At*SEC12 was amplified and subcloned into

UBQ10:sXVE:(MCS)-mCherry, followed by replacing the inducible promoter with the *UBQ10*

promoter to generate *UBQ10:At*SEC12-mCherry. The CDS of mTb was amplified by PCR

using R4pGWB601_UBQ10p-miniTurbo-NES-YFP (Addgene plasmid #127369) as the

template and subcloned into *UBQ10:At*SAR1b-S11, yielding *UBQ10:At*SAR1b-mTb-S11. The

abovementioned constructs were generated either by restriction enzyme cloning or by In-

Fusion® HD cloning (Takara). Primer sequences and the restriction enzyme cutting sites used

are listed in **Supplementary Table S1**.

Agro-infiltration of *Nicotiana benthamiana* leaves

Three-to-four-week-old *Nicotiana benthamiana* plants were used for *A. tumefaciens* (strain EHA105)-mediated infiltration. The bacterial cultures were grown overnight in Luria-Bertani (LB) medium containing the appropriate antibiotics (50 μg/ml spectinomycin, 5 μg/ml rifampicin, or 50 μg/ml kanamycin) at 30°C with shaking at 200 rpm. Cells were pelleted and resuspended in infiltration buffer containing 10 mM $MgCl_2$, 10 mM MES (pH 7.5), and 150 μM acetosyringone to an OD_{600} of 1.0 and incubated at room temperature (RT) for 2–4 h. Leaf infiltration was performed with agrobacterial suspensions at a final OD_{600} of 0.002–1.0. For *UBQ10:sXVE*-inducible expression, 9.18 μM β-estradiol was applied for an additional 24 or 48 h. Leaves were harvested at 48 or 72 h post-infiltration for confocal microscopy or protein extraction.

Confocal image analysis

Confocal images were captured using the Zeiss LSM 800 microscope (Zeiss) equipped with Plan-Apochromat 10×/0.45 M27, 20×/0.8 M27, and 40×/1.3 Oil DIC M27 objectives. The acquisition was performed in multi-track mode with line switching, and the data were averaged over two readings. Excitation/emission wavelengths were 488 nm/410–546 nm and 561 nm/560–617 nm for GFP and mCherry, respectively.

Soluble and microsomal protein extraction

Soluble (S) and microsomal (M) fractions were prepared from agro-infiltrated *N. benthamiana* leaves with the modified low-speed pellet (LSP) method (Yoshimoto et al., 2004). Leaf tissues were ground in liquid nitrogen (N) and resuspended in three volumes (3 ml per gram tissue) of pre-chilled LSP buffer containing 100 mM Tris-HCl (pH 7.5), 400 mM sucrose, 1 mM EDTA, 1% Triton X-100, 0.1 mM phenylmethylsulfonyl fluoride (PMSF), and 1× Protease Inhibitor Cocktail (P9599, Sigma-Aldrich). After centrifugation at 500 × g for 5 min and then at 13,000

× g for 15 min at 4°C, the final supernatant was collected as the soluble fraction. The pellet was

resuspended in pre-chilled LSP buffer lacking 1% Triton X-100 to yield the microsomal fraction.

Proximity labeling assay

The *N. benthamiana* leaves were incubated in 50 μM biotin solution at RT for 30 min, ground

with liquid N, and extracted in two volumes (2 ml per gram tissue) of lysis buffer containing

20 mM HEPES (pH 7.5), 40 mM KCl, 1 mM EDTA, 1% Triton X-100, 0.2 mM PMSF, and 1×

Protease Inhibitor Cocktail (P9599, Sigma-Aldrich). After centrifugation at 4,000 × g for 5 min

and then at 20,000 × g for 15 min at 4°C, the resulting supernatant was collected. A 25–750 μg

of the crude extract was incubated with 5 μl Streptavidin-agarose (S1638, Sigma-Aldrich) at

4°C overnight under 5 rpm end-to-end rotation. The beads collected by centrifugation at 2,500

× g and 4 °C for 5 min were washed with a lysis buffer containing 0.1% Triton X-100 once.

After centrifugation at 2,500 × g for 5 min at 4°C, the beads were eluted with 2× SDS sample

buffer containing 100 mM Tris-HCl (pH 6.8), 20% glycerol, 4% SDS, and 100 mM

dithiothreitol (DTT) with 0.4 M urea.

Co-immunoprecipitation assay

The 11-day-old Col-0, *UBQ10:*GFP, *UBQ10:At*SAR1c-GFP (Zeng et al., 2015), and

*dex:At*SAR1b[H74L]-CFP (Lupanga et al., 2020) seedlings were grown with 7 days of Pi

starvation. The seedlings of *dex:At*SAR1b[H74L]-CFP were treated with 10 μM

dexamethasone (DEX) or DMSO for an additional 24 h before root collection. The roots were

harvested for total protein extraction with lysis buffer containing 25 mM HEPES (pH 7.2), 150

mM NaCl, 0.5% IGEPAL CA-630 (I8896, Sigma-Aldrich), 2 mM EDTA, 0.5 mM MgCl$_2$, and

1× Protease Inhibitor Cocktail (P9599, Sigma-Aldrich). After centrifugation at 16,000 × g for

5 min at 4°C, supernatants were cross-linked with 2.5 mM dithiobis [succinimidyl propionate]

(DSP) at RT for 30 min and quenched with 100 mM Tris (pH 7.4) for 15 min. A 15–750 μg of

protein mixture was incubated with 5 µl of GFP-Trap® Agarose (ChromoTek) at 4°C for 2 h

with end-over-end rotation at 5 rpm, washed with the lysis buffer once, centrifuged at 2,500 ×

g at 4°C for 5 min, and eluted with 2× SDS sample buffer containing 100 mM Tris-HCl (pH

6.8), 20% glycerol, 4% SDS, and 100 mM DTT.

Immunoblot analysis

Protein samples were loaded onto 4–12% Q-PAGE™ Bis-Tris Precast Gel (SMOBIO) and

transferred to polyvinylidene difluoride (PVDF) membranes (IPVH00010 Immobilon-P

Membrane, Merck). The membrane was blocked with 1% BSA in 1× PBS solution containing

0.2% Tween 20 (PBST, pH 7.2) at RT for 1 h and hybridized with primary antibodies: anti-GFP

(1:5,000, AS21 4696; Agrisera) at 4°C for overnight, anti-*At*PHT1;1/2/3 (1:1,000, PHY0753;

PhytoAB) at 4°C for overnight, anti-*At*PHF1 (Huang et al., 2013) at RT for 2 h, anti-PIP2;7

(1:5,000, AS22 4812, Agrisera) at RT for 1 h, anti-cFBPase (1:5000, AS04 043, Agrisera) at

4°C for overnight, anti-mCherry (1:1,000, ab167453, Abcam) at RT for 1 h, and anti-S11

(1:10,000) at RT for 1 h, the anti-S11 polyclonal rabbit antibody was raised against the peptide

of S11 (EKRDHMVLLEYVTAAGITDASC) and produced by LTK BioLaboratories, Taiwan.

The membrane was washed four times with 1× PBST for 5 min, followed by hybridization with

the horseradish peroxidase-conjugated secondary antibody (1:20,000–1:40,000, GTX213110-

01 and 1:10,000, GTX213111-01, GeneTex) in blocking solution at RT for 1 h. After four

washes in 1× PBST for 5 min and a rinse with distilled water, chemiluminescent substrates

(WesternBright™ ECL and WesternBright™ Sirius, Advansta) for signal detection were

applied.

Quantification analyses

For puncta quantification, two non-overlapping regions of interest (ROIs, 1000 µm²) were

randomly selected from a single confocal image (53.24 × 53.24 µm²). For each ROI, the

automatic Triangle threshold was applied. Punctate structures were defined using the ImageJ

(Schneider et al., 2012) 'analyze particles' function with circularity from 0.1 to 1.0 and size

from 0.04 to infinity. Colocalization analysis was performed with a Costes' automatic threshold,

and Manders' overlap coefficients were calculated using the JACoP plugin with object-based

method in Fiji (Bolte and Cordelieres, 2006). The Dunnett's test and Mann-Whitney U test were

performed using JASP 0.95.4.0. Puncta diameters were measured by using the Zeiss ZEN 3.7

software. The box plots were generated by using BoxPlotR (Spitzer et al., 2014).

Chemical treatment

Acetosyringone (150 mM stock in DMSO, D134406, Sigma-Aldrich) was diluted to a final

concentration of 150 μM in ddH_2O. β-estradiol (36.7 mM stock in ethanol, E2758, Sigma-

Aldrich) was diluted to a final concentration of 9.18 μM in ddH_2O. Dithiobis [succinimidyl

propionate] (25 mM stock in DMSO, 22586, ThermoFisher) was diluted to a final concentration

of 2.5 mM in ddH_2O. Dexamethasone (100 mM stock in DMSO, D4902, Sigma-Aldrich) was

diluted to a final concentration of 10 μM in ddH_2O.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative under the

following accession numbers: *At*PHF1 (AT3G52190), *At*SEC12 (AT2G01470), *At*PHT1;1

(AT5G43350), *At*STP1 (AT1G11260), *At*SAR1b (AT1G56330), *At*SAR1c (AT4G02080),

*At*SEC24a (AT3G07100), *At*SEC13a (AT3G01340), *At*SEC16a (AT5G47480).

Supplementary Data

Supplementary Fig. S1. Expression of ER lumen marker in the absence and presence of

*At***PHT1;1 overexpression.** Representative confocal images **(A)** and quantification of

mCherry-HDEL-labeled puncta **(B)** are shown from three independent experiments. Scale bars,

20 μm. Box plots show medians (center lines), interquartile ranges (boxes), and data ranges

(whiskers), along with individual data points (dots). The number of regions of interest (ROI)

used for quantification is shown in parentheses. Not significant, n.s. (Student's t-test against

mCherry-HDEL expressed alone).

Supplementary Fig. S2. Size analysis of *At*PHF1-GFP- and mCherry-HDEL-labeled

puncta in the presence of *At*PHT1;1-S11 overexpression. (A) Quantification of puncta

number upon overexpression with *At*PHT1;1-S11, corresponding to Fig. 1D. Data were

collected from three independent experiments (Exp. 1, Exp. 2, and Exp. 3). **(B)** Size distribution

of overlapping and non-overlapping puncta labeled by *At*PHF1-GFP or the ER luminal marker

mCherry-HDEL, corresponding to (**A**). Box plots show medians (center lines), interquartile

ranges (boxes), and data ranges (whiskers), along with individual data points (dots). The

number of puncta used for quantification is shown in parentheses. Dunnett's test for multiple

comparisons against overlapping puncta. Not significant, n.s..

Supplementary Fig. S3. Expression of *AtSAR1* isoforms in Col-0 under Pi deprivation by

RNA-seq analysis. Expression of *AtSAR1a/b/c/d/e* in the shoot and root of 10-day-old WT

(Col-0) seedlings under Pi-sufficient conditions (+P0) or under one day (–P1) and three days of

Pi starvation (–P3) as previously described (Liu et al., 2016). RPKM stands for reads per

kilobase of transcript per million mapped reads. Numbers represent average RPKM values of

two replicates, with an assigned value of 0.25 for readings below this threshold.

Supplementary Fig. S4. The expression and distribution of COPII-related components in

agro-infiltrated *N. benthamiana* leaves. (A) Expression and distribution of S11-tagged

*At*SAR1b, *At*SAR1c, *At*SEC24a, *At*SEC13a, and *At*SEC16a at the peripheral layer and 3xHA-

S11 at the middle layer of agro-infiltrated tobacco epidermal cells. **(B)** Expression and

distribution of S10-tagged *At*SEC12, *At*PHF1, and 3xHA at the middle layer of agro-infiltrated

tobacco epidermal cells. Scale bars, 5 μm. Representative confocal images are shown from

three independent experiments.

Supplementary Fig. S5. Quantification of puncta sizes upon co-expression of *At*PHF1-

mCherry and *At*PHT1;1 with ERES markers. (A, **B)** Quantification of puncta number upon

overexpression with *At*SAR1b-S11 (**A**) or *At*SEC24a-S11 (**B**), corresponding to Fig. 2C. Data

were collected from three independent experiments (Exp. 1, Exp. 2, and Exp. 3). **(C**, **D)** Size

distribution of overlapping and non-overlapping puncta labeled by *At*PHF1-mCherry or ERES

markers, corresponding to (**A**) and (**B**), respectively. Box plots show medians (center lines),

interquartile ranges (boxes), and data ranges (whiskers), along with individual data points (dots).

The number of puncta used for quantification is shown in parentheses. Dunnett's test for

multiple comparisons against overlapping puncta. Not significant, n.s..

Supplementary Fig. S6. Expression and subcellular distribution of the truncated *At*PHF1

variants in agro-infiltrated *N. benthamiana* leaves. (A) Schematic diagrams of the truncated

*At*PHF1 variants with domain boundaries indicated by amino acid numbers. **(B**, **C)** The

expression and distribution of the S10-tagged (**B**) and the S11-tagged (**C**) truncated *At*PHF1

variants at the middle layer of epidermal cells. Scale bars, 20 μm. Representative confocal

images are shown from three independent experiments.

Supplementary Fig. S7. *At*SAR1b and *At*SAR1c interact with the cytosolic and

transmembrane domains of *At*PHF1 in agro-infiltrated *N. benthamiana* leaves. (A–C)

The interaction analyses of the S10-*At*PHF1 truncated variants with *At*SAR1b-S11 (**A**),

*At*SAR1c-S11 (**B**), and 3xHA-S11 (**C**) by using the tripartite split-GFP assay. The interactions

with 3xHA-S11 were used as negative controls. Scale bars, 10 μm. Representative confocal

images taken at the peripheral layer of the epidermis are shown from at least three independent experiments.

Supplementary Fig. S8. Distribution of *At*SAR1b/[H74L]/[T34N] in the agro-infiltrated *N. benthamiana* leaves. (A) The expression and distribution of the S11-tagged GTP-locked ([H74L]) and GDP-locked *At*SAR1b ([T34N]) at the peripheral layer of epidermal cells. Scale bars, 5 µm. Representative confocal images are shown from three independent experiments. **(B)** Protein expression of *At*SAR1b/[H74L]/[T34N] in the soluble (S) and microsomal (M) fractions isolated using the modified low-speed pellet (LSP) method (Yoshimoto et al., 2004). Anti-S11 antibody was used to detect S11 fusion proteins. The membrane association percentage (%) was calculated as M/(S+M) and is shown as mean ± SE from two independent experiments. Aquaporin PIP2;7 and cytosolic fructose-1,6-bisphosphatase (cFBPase) served as microsomal and soluble controls, respectively. Amido black staining was used as a loading control.

Supplementary Fig. S9. ER export-incompetent *At*PHT1;1 overexpression cannot trigger partial distribution of *At*PHF1 to punctate structures in agro-infiltrated *N. benthamiana* leaves. (A) Expression and distribution of S10-*At*PHT1;1 TM1–6 and *At*PHT1;1 TM7–12-S11 upon co-expression with *At*PHF1-mCherry in agro-infiltrated tobacco epidermal cells. Scale bars, 5 µm. Representative confocal images taken at the peripheral layer of epidermal cells are shown from three independent experiments. **(B)** Distribution of *At*PHF1-GFP in the presence of S10-*At*PHT1;1 TM1–6 or *At*PHT1;1 TM7–12-S11 co-expression. Scale bars, 5 µm. Representative confocal images are shown from three independent experiments. **(C)** Quantification of *At*PHF1-GFP-labeled punctate structures in (**B**). The data for *At*PHF1-GFP expressed alone are the same as those shown in Fig. 1B. Box plots display medians (center lines), interquartile ranges (boxes), and data ranges (whiskers), along with individual data points (dots). The number of regions of interest used for quantification is shown in parentheses.

Dunnett's test for multiple comparisons against *At*PHF1-GFP expressed alone. Not significant,

n.s..

Supplementary Fig. S10. Interaction analysis of *At*PHF1 with *At*SAR1 in the yeast split-

ubiquitin system. (A) Co-expression of *At*PHF1-CubPLV with NubG-*At*PHT1;1, *At*SAR1b-

NubG or *At*SAR1c-NubG. **(B)** Co-expression of *At*SEC12-CubPLV with *At*SAR1b-NubG and

*At*SAR1c-NubG. The co-expression of *At*PHF1-CubPLV or *At*SEC12-CubPLV with NubWT

was used as a positive control, while the co-expression with NubG was used as a negative

control. The CDS of *At*PHF1, *At*SEC12, *At*PHT1;1, *At*SAR1b, and *At*SAR1c were amplified by

PCR, cloned into the pCR8/GW/TOPO, verified by sequencing, and recombined into

destination vectors via LR reactions. *At*PHF1 and *At*SEC12 were cloned into synthesized

MET17-PLVCub_GW (Chiu, 2024) to generate C-terminal Protein A–LexA–VP16–Cub

fusions (*At*PHF1-CubPLV and *At*SEC12-CubPLV); *At*PHT1;1 was cloned into NX32_GW

(ABRC plasmid CD3-1737) to produce NubG-*At*PHT1;1; *At*SAR1b and *At*SAR1c were cloned

into XN22_GW (ABRC plasmid CD3-1735) to produce *At*SAR1b-NubG and *At*SAR1c-NubG.

Yeast transformants were grown on synthetic medium lacking tryptophan and leucine (SC–WL)

for growth detection or on synthetic medium lacking tryptophan, leucine, adenine, and histidine

containing 0.5 µM methionine (SC–WLAH + M) for testing protein-protein interaction.

Representative results are shown from two independent experiments.

Supplementary Fig. S11. Interaction analysis of the N-terminal cytosolic domain of

***At*PHF1 with *At*SAR1b/[H74L]/[T34N] in the yeast two-hybrid assay.** GAL4[BD]-S10-

*At*PHF1 N was co-expressed with GAL4[AD]-*At*SAR1b-S11, GAL4[AD]-*At*SAR1b[H74L]-

S11, or GAL4[AD]-*At*SAR1b[T34N]-S11. The co-expression of Krev1 with RalGDS-wt,

RalGDS-m1, and RalGDS-m2 were strong positive, weak positive, and negative controls,

respectively. The CDS of *At*SAR1b-S11, *At*SAR1b[H74L]-S11, and *At*SAR1b[T34N]-S11 were

subcloned into pCR8/GW/TOPO and recombined into pDEST22 via LR reaction to generate GAL4[AD]-*At*SAR1b/[H74L]/[T34N]-S11; the CDS of S10-*At*PHF1 N was subcloned into pCR8/GW/TOPO and recombined into pDEST32 via LR reaction to generate GAL4[BD]-S10-*At*PHF1 N. Yeast transformants were grown on synthetic medium lacking tryptophan and leucine (SC–WL) for growth detection, and on medium lacking tryptophan, leucine, and uracil (SC–WLU) or lacking tryptophan, leucine, and histidine with 10 mM 3-Amino-1,2,4-triazole (SC–WLH + 3AT) to assess protein-protein interactions. Representative results are shown from two independent experiments.

Supplementary Table S1. Oligonucleotides used for plasmid constructs.

Funding

This work was supported by the Ministry of Science and Technology (MOST 108-2311-B-007-003-MY3) and the National Science and Technology Council (NSTC 112-2313-B-007-001-MY3).

Acknowledgments

We thank Dr. Liwen Jiang at the Chinese University of Hong Kong, People's Republic of China, for sharing the *Arabidopsis* seeds of *UBQ10:At*SAR1c-GFP and *UBQ10:*GFP homozygous lines, Dr. Masaki Takeuchi at the University of Tokyo, Japan, for sharing the clone *35S:At*SAR1b[H74L], Dr. Tzyy-Jen Chiou at Academia Sinica, Taiwan (R.O.C.), for sharing the clone *35S:At*PHF1-mCherry, and Dr. Karin Schumacher at the Cell Biology Research Group of the Centre for Organismal Studies (COS), Heidelberg University, Germany, for sharing the *Arabidopsis* seeds of *dex:At*SAR1b[H74L]-CFP T2 lines. We acknowledge Ms. Wen-Chun Chou for constructing the plasmid encoding *35S:*GFP1–10 and the technical support from the

Confocal Image Core, National Tsing Hua University (sponsored by MOST 108-2731-M-007-

001 and MOST 110-2731-M-007-001).

Author Contributions

T.-Y. L. conceived the original research plan, designed and supervised the experiments, and

wrote the article. H.-F. L. designed and performed the experiments, analyzed the data, wrote

the article, and contributed to the preparation of figures. J.-D. C. designed and performed the

experiments.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

All study data are included in the article and/or Supporting Material.

Figure Legends

Figure 1. *At*PHT1;1 overexpression triggers partial distribution of *At*PHF1 to punctate

structures in agro-infiltrated *N. benthamiana* leaves. **(A)** Distribution of *At*PHF1-GFP in the

absence or presence of *At*PHT1;1-S11 or *At*STP1-S11 co-expression. Arrows, punctate

structures. **(B)** Quantification analysis of **(A)** from three independent experiments. **(C)** Co-

localization analysis of *At*PHF1-GFP puncta and mCherry-HDEL puncta in the presence of

*At*PHT1;1-S11 or *At*STP1-S11 co-expression. Arrows, overlapping puncta; circles, non-

overlapping puncta. **(D)** The quantification of **(C)** from three independent experiments.

Representative confocal images at the peripheral layers of the epidermis are shown in **(A)** and

(C); scale bars, 5 μm. Box plots show medians (center lines), interquartile ranges (boxes), data

ranges (whiskers), and data points (dots) in **(B)** and **(D)**. The number of regions of interest used

for quantification is shown in parentheses. ***, $P < 0.001$ versus *At*PHF1-GFP expression alone in (**B**); *, $P < 0.05$ versus overlapping puncta in (**D**); Dunnett's test for multiple comparisons.

Figure 2. *At*PHT1;1-induced *At*PHF1 puncta partially co-localize with the ERES markers in agro-infiltrated *N. benthamiana* leaves. (A, **B)** Co-localization of *At*SAR1b-S11 (**A**) and *At*SEC24a-S11 (**B**) with *At*PHF1-mCherry in the absence or presence of *At*PHT1;1 overexpression. Representative confocal images were taken at the peripheral layer of the epidermis. Arrows, overlapping puncta; circles, non-overlapping puncta. Scale bars, 5 μm. **(C)** Quantification of co-localization in (**A**) and (**B**). The number of regions of interest used for quantification is shown in parentheses. *, $P < 0.05$; ***, $P < 0.001$; Dunnett's test for multiple comparisons against overlapping with *At*SAR1b or *At*SEC24a. **(D)** Comparison of Manders' overlap coefficient between the co-localization of *At*PHF1-mCherry with *At*SAR1b-S11 and *At*SEC24a-S11. Data were collected from three independent experiments. Box plots show medians (center lines), interquartile ranges (boxes), data ranges (whiskers), and data points (dots). The puncta numbers used for the quantification analysis are shown in parentheses. *, $P < 0.05$; two-tailed Mann-Whitney U test.

Figure 3. *At*PHF1 interacts with *At*SAR1b and *At*SAR1c *in planta* based on the tripartite split-GFP association in agro-infiltrated *N. benthamiana* leaves. The interaction of S10-*At*SEC12 (**A**) or S10-*At*PHF1 (**B**) with the COPII-related components, *At*SAR1b-, *At*SAR1c-, *At*SEC24a-, *At*SEC13a-, and *At*SEC16a-S11. The interactions with 3xHA-S11 in (**A**) and (**B**) and S10-3xHA (**C**) were used as negative controls. Scale bars, 5 μm. Representative confocal images taken at the peripheral layer of the epidermis are shown from three independent experiments.

Figure 4. *At*PHF1 interacts with *At*SAR1b *in planta* based on the proximity labeling in agro-infiltrated *N. benthamiana* leaves. (A) Schematic diagrams of mTb-fused bait protein in the proximity of the prey protein. The reactive intermediate biotinyl-5'-AMP biotinylates proximal prey proteins within the indicated range (grey zone). **(B)** The expression and subcellular distribution of mTb-NES-EYFP and *At*SAR1b-mTb-S11. Scale bars, 20 μm. Representative confocal images are shown from three independent experiments. **(C)** The interaction analyses of *At*PHF1-mCherry and *At*SAR1b by proximity labeling. In the left panel, thirty-fold more protein extract expressing *At*SAR1b-mTb-S11 was used as input for Streptavidin pulldown relative to that expressing mTb-NES-EYFP; in the right panel, equal amounts of protein from the samples were used for pulldown. Anti-S11 antibody was used to detect EYFP-tagged and S11-tagged fusion proteins. Representative results are shown from two independent experiments.

Figure 5. Co-immunoprecipitation of the endogenous *At*PHF1 and *At*PHT1;1/2/3 with *At*SAR1c-GFP in *Arabidopsis* transgenic plants. (A, B) The interaction analyses of *At*SAR1c-GFP with the endogenous *At*PHF1 and *At*PHT1;1/2/3 in 11-day-old wild-type (Col-0), *UBQ10:*GFP, and *UBQ10:At*SAR1c-GFP seedlings subjected to Pi deprivation (0 μM KH_2PO_4, 7 days of starvation). Fifty-fold more protein extract expressing *At*SAR1c-GFP was used as input for GFP-Trap co-immunoprecipitation relative to that expressing free GFP in **(A)**. Equal amounts of protein from the samples were used for co-immunoprecipitation in **(B)**. Anti-S11 antibody was used to detect GFP fusion proteins. Representative results are shown from three independent experiments.

Figure 6. *At*PHF1 preferentially interacts with the GDP-locked form of *At*SAR1b in agro-infiltrated *N. benthamiana* leaves and *Arabidopsis* transgenic plants. (A, B) The interaction analyses of S10-*At*SEC12 **(A)** and S10-*At*PHF1 variants **(B)** with *At*SAR1b/[H74L]/[T34N]-

S11 in agro-infiltrated tobacco leaves. Representative confocal images were taken at the peripheral layer of the epidermis. Scale bars, 2 μm. Quantification indicates the number of cells expressing GFP signals per unit leaf area (mm^2). Data were collected from three independent experiments; the number of confocal images (0.32 x 0.32 mm^2) used for quantification is shown in parentheses. **, $P < 0.01$; ***, $P < 0.001$; Dunnett's test for multiple comparisons against *At*SAR1b[H74L]. Not significant, n.s.. **(C)** Interaction analyses of the WT or GTP-locked form of *At*SAR1 with the endogenous *At*PHF1 were performed in 11-day-old *UBQ10:At*SAR1c-GFP and *dex:At*SAR1b[H74L]-CFP seedlings subjected to Pi deprivation (0 μM KH_2PO_4, 7 days). *dex:At*SAR1b[H74L]-CFP seedlings were treated with 24 hours of 10 μM dexamethasone (DEX) or DMSO. The protein amounts as input for GFP-Trap co-immunoprecipitation are as indicated above the image. An anti-GFP antibody was used to detect GFP and CFP fusion proteins. The relative intensity is shown as mean ± SE from two independent experiments.

Figure 7. Hypothetical model depicting a role for *At*PHF1 in COPII recruitment for the ER export of *At*PHT1 transporters. The GEF SEC12 recruits and activates SAR1 GTPase by exchanging GDP for GTP. The direct interaction of GTP-bound SAR1 with SEC23 recruits SEC23/24 to form the inner coat, followed by SEC13/31 recruitment to form the outer coat. The ER-resident PHF1 concentrates on ERES triggered by increased PHT1 proteins and interacts with GDP-bound SAR1. While the scaffold protein SEC16 at ERES does not interact with PHF1, the interaction of PHF1 and SAR1 may link to the regulation of SAR1 GTPase recruitment, thereby facilitating COPII assembly at ERES for the ER-to-Golgi transport of PHT1 proteins in the budding COPII vesicles.

References

Bannykh SI, Rowe, T., & Balch, W. E. (1996) The organization of endoplasmic reticulum export complexes. The Journal of cell biology **135:** 19-35

Barlowe CK, Miller EA (2013) Secretory protein biogenesis and traffic in the early secretory
pathway. Genetics **193:** 383-410

Bayle V, Arrighi JF, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E, Paz-Ares J,
Nussaume L (2011) Arabidopsis thaliana high-affinity phosphate transporters exhibit
multiple levels of posttranslational regulation. The Plant Cel **23:** 1523-1535

Bi X, Corpina, R. A., & Goldberg, J. (2002) Structure of the Sec23/24-Sar1 pre-budding complex
of the COPII vesicle coat. Nature **419:** 271-277

Bisnett BJ, Condon BM, Lamb CH, Georgiou GR, Boyce M (2020) Export Control: Post-
transcriptional Regulation of the COPII Trafficking Pathway. Front Cell Dev Biol **8:**
618652

Bolte S, Cordelieres FP (2006) A guided tour into subcellular colocalization analysis in light
microscopy. J Microsc **224:** 213-232

Cabantous S, Nguyen HB, Pedelacq JD, Koraichi F, Chaudhary A, Ganguly K, Lockard MA,
Favre G, Terwilliger TC, Waldo GS (2013) A new protein-protein interaction sensor
based on tripartite split-GFP association. Scientific reports **3:** 2854

Chiu CY, Tsai, C. D., Wang, J. Y., Tsai, M. H., McGinness, A. J., Lung, H. F., Kanno, S.,
Kriechbaumer, V. & Liu, T. Y. (2024) Phosphate starvation-induced CORNICHON
HOMOLOG 5 as endoplasmic reticulum cargo receptor for PHT1 transporters in
Arabidopsis. bioRxiv**:** 2024.2006.2020.599911

Chung KP, Zeng Y, Jiang L (2016) COPII paralogs in plants: functional redundancy or diversity?
Trends in Plant Science **21:** 758-769

d'Enfert C, Wuestehube, L. J., Lila, T., & Schekman, R. (1991) Sec12p-dependent membrane
binding of the small GTP-binding protein Sar1p promotes formation of transport
vesicels from the ER. The Journal of cell biology **114:** 663-670

daSilva LL, Snapp EL, Denecke J, Lippincott-Schwartz J, Hawes C, Brandizzi F (2004)
Endoplasmic reticulum export sites and Golgi bodies behave as single mobile secretory
units in plant cells. The Plant Cell **16:** 1753-1771

Dharmasiri S, Swarup, R., Mockaitis, K., Dharmasiri, N., Singh, S. K., Kowalchyk, M., ... &
Estelle, M. (2006) AXR4 is required for localization of the auxin influx facilitator AUX1.
Science **312:** 1218-1220

Gimeno RE, Espenshade, P., & Kaiser, C. A. (1995) SED4 encodes a yeast endoplasmic
reticulum protein that binds Sec16p and participates in vesicle formation. The Journal
of cell biology **131:** 325-338

González E, Solano R, Rubio V, Leyva A, Paz-Ares J (2005) PHOSPHATE TRANSPORTER TRAFFIC
FACILITATOR1 is a plant-specific SEC12-related protein that enables the endoplasmic
reticulum exit of a high-affinity phosphate transporter in Arabidopsis. The Plant Cell **17:**
3500-3512

Hanton SL, Chatre L, Matheson LA, Rossi M, Held MA, Brandizzi F (2008) Plant Sar1 isoforms
with near-identical protein sequences exhibit different localisations and effects on
secretion. Plant Molecular Biology **67:** 283-294

Hanton SL, Chatre L, Renna L, Matheson LA, Brandizzi F (2007) De novo formation of plant
endoplasmic reticulum export sites is membrane cargo induced and signal mediated.
Plant Physiology **143:** 1640-1650

Hanton SL, Matheson LA, Chatre L, Brandizzi F (2009) Dynamic organization of COPII coat
proteins at endoplasmic reticulum export sites in plant cells. The Plant Journal **57:** 963-
974

**Huang TK, Han CL, Lin SI, Chen YJ, Tsai YC, Chen YR, Chen JW, Lin WY, Chen PM, Liu TY, Chen
YS, Sun CM, Chiou TJ** (2013) Identification of downstream components of ubiquitin-
conjugating enzyme PHOSPHATE2 by quantitative membrane proteomics in
Arabidopsis roots. The Plant Cell **25:** 4044-4060

Joiner AMN, Fromme JC (2021) Structural basis for the initiation of COPII vesicle biogenesis.
Structure **29:** 859-872 e856

Kim DI, Birendra KC, Zhu W, Motamedchaboki K, Doye V, Roux KJ (2014) Probing nuclear pore
complex architecture with proximity-dependent biotinylation. Proc Natl Acad Sci U S A
111: E2453-2461

Kodera C, Yorimitsu T, Nakano A, Sato K (2011) Sed4p stimulates Sar1p GTP hydrolysis and
promotes limited coat disassembly. Traffic **12:** 591-599

Kota J, Gilstring CF, Ljungdahl PO (2007) Membrane chaperone Shr3 assists in folding amino
acid permeases preventing precocious ERAD. J Cell Biol **176:** 617-628

Kota J, Ljungdahl PO (2005) Specialized membrane-localized chaperones prevent aggregation
of polytopic proteins in the ER. The Journal of cell biology **168:** 79-88

Kurokawa K, Nakano A (2019) The ER exit sites are specialized ER zones for the transport of
cargo proteins from the ER to the Golgi apparatus. The journal of biochemistry **165:**
109-114

Lau WTW, Howson, R. W., Malkus, P., Schekman, R., & O'Shea, E. K. (2000) Pho86p, an
endoplasmic reticulum (ER) resident protein in Saccharomyces cerevisiae, is required
for ER exit of the high-affinity phosphate transporter Pho84p. Proceedings of the
National Academy of Sciences **97:** 1107-1112

Li B, Zeng, Y., Cao, W., Zhang, W., Cheng, L., Yin, H., ... & Jiang, L. (2021) A distinct giant coat
protein complex II vesicle population in Arabidopsis thaliana. Nature Plants **7:** 1335-
1346

Liang X, Li SW, Gong LM, Li S, Zhang Y (2020) COPII components Sar1b and Sar1c play distinct
yet interchangeable roles in pollen development. Plant Physiology **183:** 974-985

Liu TY, Chou WC, Chen WY, Chu CY, Dai CY, Wu PY (2018) Detection of membrane protein-
protein interaction in planta based on dual-intein-coupled tripartite split-GFP
association. The Plant Journal **94:** 426-438

Liu TY, Huang TK, Tseng CY, Lai YS, Lin SI, Lin WY, Chen JW, Chiou TJ (2012) PHO2-dependent
degradation of PHO1 modulates phosphate homeostasis in Arabidopsis. Plant Cell **24:**
2168-2183

Liu TY, Huang TK, Yang SY, Hong YT, Huang SM, Wang FN, Chiang SF, Tsai SY, Lu WC, Chiou TJ
(2016) Identification of plant vacuolar transporters mediating phosphate storage.
Nature communications **7:** 11095

Liu TY, Tsai, M. H., Wang, J. Y.,, Lung HF, Chiu CY, Chow HX, Lu CA (2025) CORNICHON
HOMOLOG 5-dependent ER export of membrane cargoes in phosphate-starved
Arabidopsis root as revealed by membrane proteomic analysis. bioRxiv**:**
2024.2012.2030.630708

Lupanga U, Rohrich R, Askani J, Hilmer S, Kiefer C, Krebs M, Kanazawa T, Ueda T, Schumacher
K (2020) The Arabidopsis V-ATPase is localized to the TGN/EE via a seed plant-specific
motif. Elife **9**

Mair A, Xu SL, Branon TC, Ting AY, Bergmann DC (2019) Proximity labeling of protein
complexes and cell-type-specific organellar proteomes in Arabidopsis enabled by
TurboID. elife **8**

McMahon C, Studer SM, Clendinen C, Dann GP, Jeffrey PD, Hughson FM (2012) The structure
of Sec12 implicates potassium ion coordination in Sar1 activation. Journal of Biological
Chemistry **287:** 43599-43606

Miller E, Antonny, B., Hamamoto, S., & Schekman, R. (2002) Cargo selection into COPII
vesicles is driven by the Sec24p subunit. The EMBO journal**:** 6105-6113

Millner PD, & Kitt, D. G. (1992) The Beltsville method for soilless production of vesicular-
arbuscular mycorrhizal fungi. Mycorrhiza **2:** 9-15

Montegna EA, Bhave M, Liu Y, Bhattacharyya D, Glick BS (2012) Sec12 binds to Sec16 at
transitional ER sites. PLoS One **7:** e31156

Nussaume L, Kanno S, Javot H, Marin E, Pochon N, Ayadi A, Nakanishi TM, Thibaud MC (2011)
Phosphate import in plants: focus on the PHT1 transporters. Frontiers in plant science
2: 83

Paul S, Audhya A, Cui Q (2023) Molecular mechanism of GTP binding-and dimerization-
induced enhancement of Sar1-mediated membrane remodeling. Proceedings of the
National Academy of Sciences **120:** e2212513120

Saito-Nakano Y, Nakano A (2000) Sed4p functions as a positive regulator of Sar1p probably
through inhibition of the GTPase activation by Sec23p. Genes to Cells **5:** 1039-1048

Saito K, Yamashiro K, Shimazu N, Tanabe T, Kontani K, Katada T (2014) Concentration of Sec12
at ER exit sites via interaction with cTAGE5 is required for collagen export. Journal of
Cell Biology **206:** 751-762

Schlücking K, Edel KH, Koster P, Drerup MM, Eckert C, Steinhorst L, Waadt R, Batistic O, Kudla
J (2013) A new β-estradiol-inducible vector set that facilitates easy construction and
efficient expression of transgenes reveals CBL3-dependent cytoplasm to tonoplast
translocation of CIPK5. Mol Plant **6:** 1814-1829

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis.
Nature methods **9:** 671-675

Shen J, Yuan, L., Zhang, J., Li, H., Bai, Z., Chen, X., ... & Zhang, F. (2011) Phosphorus dynamics:
from soil to plant. Plant physiology **156:** 997-1005

Spitzer M, Wildenhain J, Rappsilber J, Tyers M (2014) BoxPlotR: a web tool for generation of
box plots. Nature methods **11:** 121-122

Stagg SM, Gurkan C, Fowler DM, LaPointe P, Foss TR, Potter CS, Carragher B, Balch WE (2006)
Structure of the Sec13/31 COPII coat cage. Nature **439:** 234-238

Supek F, Madden DT, Hamamoto S, Orci L, Schekman R (2002) Sec16p potentiates the action
of COPII proteins to bud transport vesicles. The Journal of cell biology **158:** 1029-1038

Takagi J, Kimori Y, Shimada T, Hara-Nishimura I (2020) Dynamic Capture and Release of
Endoplasmic Reticulum Exit Sites by Golgi Stacks in Arabidopsis. iScience **23:** 101265

Takagi J, Renna L, Takahashi H, Koumoto Y, Tamura K, Stefano G, Fukao Y, Kondo M,
Nishimura M, Shimada T, Brandizzi F, Hara-Nishimura I (2013) MAIGO5 functions in
protein export from Golgi-associated endoplasmic reticulum exit sites in Arabidopsis.
The Plant Cell **25:** 4658-4675

Takeuchi M, Ueda, T., Sato, K., Abe, H., Nagata, T., & Nakano, A. (2000) A dominant negative
mutant of Sar1 GTPase inhibits protein transport from the ER to the Golgi apparatus in
tobacco and Arabidopsis cultured cells. The Plant Journal **23:** 517-525

Tang BL, Wang Y, Ong YS, Hong W (2005) COPII and exit from the endoplasmic reticulum.
Biochim Biophys Acta **1744:** 293-303

Tidy A, Abu Bakar N, Carrier D, Kerr ID, Hodgman C, Bennett MJ, Swarup R (2024) Mechanistic
insight into the role of AUXIN RESISTANCE4 in trafficking of AUXIN1 and LIKE AUX1-2.
Plant Physiology **194:** 422-433

Van der Verren SE, Zanetti G (2023) The small GTPase Sar1, control centre of COPII trafficking.
FEBS letters **597:** 865-882

Wang X, Xu M, Gao C, Zeng Y, Cui Y, Shen W, Jiang L (2020) The roles of endomembrane
trafficking in plant abiotic stress responses. Journal of integrative plant biology **62:** 55-
69

Wei T, Wang A (2008) Biogenesis of cytoplasmic membranous vesicles for plant potyvirus
replication occurs at endoplasmic reticulum exit sites in a COPI-and COPII-dependent
manner. Journal of virology **82:** 12252-12264

Weigel AV, Chang CL, Shtengel G, Xu CS, Hoffman DP, Freeman M, Iyer N, Aaron J, Khuon S,
Bogovic J, Qiu W, Hess HF, Lippincott-Schwartz J (2021) ER-to-Golgi protein delivery
through an interwoven, tubular network extending from ER. Cell **184:** 2412-2429 e2416

Weissman JT, Plutner H, Balch WE (2001) The mammalian guanine nucleotide exchange factor
mSec12 is essential for activation of the Sar1 GTPase directing endoplasmic reticulum
export. Traffic **2:** 465-475

Xu SL, Shrestha R, Karunadasa SS, Xie PQ (2023) Proximity labeling in plants. Annual review
of plant biology **74:** 285-312

Yang YD, Elamawi R, Bubeck J, Pepperkok R, Ritzenthaler C, Robinson DG (2005) Dynamics of
COPII vesicles and the Golgi apparatus in cultured Nicotiana tabacum BY-2 cells
provides evidence for transient association of Golgi stacks with endoplasmic reticulum
exit sites. Plant Cell **17:** 1513-1531

Yoshimoto K, Hanaoka H, Sato S, Kato T, Tabata S, Noda T, Ohsumi Y (2004) Processing of
ATG8s, ubiquitin-like proteins, and their deconjugation by ATG4s are essential for plant
autophagy. The Plant Cell **16:** 2967-2983

Zeng Y, Chung KP, Li B, Lai CM, Lam SK, Wang X, Cui Y, Gao C, Luo M, Wong KB, Schekman R,
Jiang L (2015) Unique COPII component AtSar1a/AtSec23a pair is required for the
distinct function of protein ER export in Arabidopsis thaliana. Proceedings of the
National Academy of Sciences **112:** 14360-14365

Zeuschner D, Geerts WJ, van Donselaar E, Humbel BM, Slot JW, Koster AJ, Klumperman J
(2006) Immuno-electron tomography of ER exit sites reveals the existence of free
COPII-coated transport carriers. Nat Cell Biol **8:** 377-383

Zhang Z, Nakamura S, Yamasaki A, Uehara M, Takemura S, Tsuchida K, Kamiya T, Shigenobu
S, Yamaguchi K, Fujiwara T, Ishiguro S, Takano J (2024) Arabidopsis KNS3 and its two
homologs mediate endoplasmic reticulum-to-plasma membrane traffic of boric acid
channels. Journal of Experimental Botany **75:** 7046-7065

Figure 1. *At*PHT1;1 overexpression triggers partial distribution of *At*PHF1 to punctate structures in agro-infiltrated *N. benthamiana* leaves. **(A)** Distribution of *At*PHF1-GFP in the absence or presence of *At*PHT1;1-S11 or *At*STP1-S11 co-expression. Arrows, punctate structures. **(B)** Quantification analysis of **(A)** from three independent experiments. **(C)** Co-localization analysis of *At*PHF1-GFP puncta and mCherry-HDEL puncta in the presence of *At*PHT1;1-S11 or *At*STP1-S11 co-expression. Arrows, overlapping puncta; circles, non-overlapping puncta. **(D)** The quantification of **(C)** from three independent experiments. Representative confocal images at the peripheral layers of the epidermis are shown in **(A)** and **(C)**; scale bars, 5 μm. Box plots show medians (center lines), interquartile ranges (boxes), data ranges (whiskers), and data points (dots) in **(B)** and **(D)**. The number of regions of interest used for quantification is shown in parentheses. ***, $P < 0.001$ versus *At*PHF1-GFP expression alone in **(B)**; *, $P < 0.05$ versus overlapping puncta in **(D)**; Dunnett's test for multiple comparisons.

Figure 2. *At*PHT1;1-induced *At*PHF1 puncta partially co-localize with the ERES markers in agro-infiltrated *N. benthamiana* leaves. (A, **B)** Co-localization of *At*SAR1b-S11 (**A**) and *At*SEC24a-S11 (**B**) with *At*PHF1-mCherry in the absence or presence of *At*PHT1;1 overexpression. Representative confocal images were taken at the peripheral layer of the epidermis. Arrows, overlapping puncta; circles, non-overlapping puncta. Scale bars, 5 μm. **(C)** Quantification of co-localization in (**A**) and (**B**). The number of regions of interest used for quantification is shown in parentheses. *, $P < 0.05$; ***, $P < 0.001$; Dunnett's test for multiple comparisons against overlapping with *At*SAR1b or *At*SEC24a. **(D)** Comparison of Manders' overlap coefficient between the co-localization of *At*PHF1-mCherry with *At*SAR1b-S11 and *At*SEC24a-S11. Data were collected from three independent experiments. Box plots show medians (center lines), interquartile ranges (boxes), data ranges (whiskers), and data points (dots). The puncta numbers used for the quantification analysis are shown in parentheses. *, $P < 0.05$; two-tailed Mann-Whitney U test.

Figure 3. *At*PHF1 interacts with *At*SAR1b and *At*SAR1c *in planta* based on the tripartite split-GFP association in agro-infiltrated *N. benthamiana* leaves. The interaction of S10-*At*SEC12 **(A)** or S10-*At*PHF1 **(B)** with the COPII-related components, *At*SAR1b-, *At*SAR1c-, *At*SEC24a-, *At*SEC13a-, and *At*SEC16a-S11. The interactions with 3xHA-S11 in **(A)** and **(B)** and S10-3xHA **(C)** were used as negative controls. Scale bars, 5 μm. Representative confocal images taken at the peripheral layer of the epidermis are shown from three independent experiments.

Figure 4. ***At*PHF1 interacts with *At*SAR1b *in planta* based on the proximity labeling in agro-infiltrated *N. benthamiana* leaves. (A)** Schematic diagrams of mTb-fused bait protein in the proximity of the prey protein. The reactive intermediate biotinyl-5'-AMP biotinylates proximal prey proteins within the indicated range (grey zone). **(B)** The expression and subcellular distribution of mTb-NES-EYFP and *At*SAR1b-mTb-S11. Scale bars, 20 μm. Representative confocal images are shown from three independent experiments. **(C)** The interaction analyses of *At*PHF1-mCherry and *At*SAR1b by proximity labeling. In the left panel, thirty-fold more protein extract expressing *At*SAR1b-mTb-S11 was used as input for Streptavidin pulldown relative to that expressing mTb-NES-EYFP; in the right panel, equal amounts of protein from the samples were used for pulldown. Anti-S11 antibody was used to detect EYFP-tagged and S11-tagged fusion proteins. Representative results are shown from two independent experiments.

Figure 5. Co-immunoprecipitation of the endogenous *At*PHF1 and *At*PHT1;1/2/3 with *At*SAR1c-GFP in *Arabidopsis* transgenic plants. (A, **B)** The interaction analyses of *At*SAR1c-GFP with the endogenous *At*PHF1 and *At*PHT1;1/2/3 in 11-day-old wild-type (Col-0), *UBQ10:*GFP, and *UBQ10:At*SAR1c-GFP seedlings subjected to Pi deprivation (0 µM KH_2PO_4, 7 days of starvation). Fifty-fold more protein extract expressing *At*SAR1c-GFP was used as input for GFP-Trap co-immunoprecipitation relative to that expressing free GFP in (**A**). Equal amounts of protein from the samples were used for co-immunoprecipitation in (**B**). Anti-S11 antibody was used to detect GFP fusion proteins. Representative results are shown from three independent experiments.

A

AtSAR1b-S11 + GFP1–9   AtSAR1b[H74L]-S11 + GFP1–9   AtSAR1b[T34N]-S11 + GFP1–9

S10-AtSEC12

B

S10-AtPHF1

S10-AtPHF1 N

S10-AtPHF1 N–TM

S10-AtPHF1 TM

S10-AtPHF1 TM–C

AtSAR1b   AtSAR1b[H74L]   AtSAR1b[T34N]

No. of cells expressing GFP signals/leaf area (mm²)

*** (n=22)   (n=20)   *** (n=21)

*** (n=22)   (n=23)   *** (n=21)

*** (n=24)   (n=21)   *** (n=23)

n.s. (n=29)   (n=25)   n.s. (n=24)

n.s. (n=22)   (n=22)   ** (n=22)

C

AtSAR1b[H74L]-CFP

DMSO   DEX   AtSAR1c-GFP

Input protein amount: 1x   1x   1x

kDa

57 —

IB: α-GFP

◀ AtSAR1b[H74L]-CFP
◀ AtSAR1c-GFP

42 —

Relative intensity: 1.0   1.2±0.1

IP protein amount: 350x   350x   350x

IB: α-AtPHF1

42 —   ◀ AtPHF1

Relative intensity: 1.0   4.1±0.7

Figure 6. *At***PHF1 preferentially interacts with the GDP-locked form of** *At***SAR1b in agro-infiltrated** *N.* *benthamiana* **leaves and** *Arabidopsis* **transgenic plants. (A**, **B)** The interaction analyses of S10-*At*SEC12 (**A**) and S10-*At*PHF1 variants (**B**) with *At*SAR1b/[H74L]/[T34N]-S11 in agro-infiltrated tobacco leaves. Representative confocal images were taken at the peripheral layer of the epidermis. Scale bars, 2 μm. Quantification indicates the number of cells expressing GFP signals per unit leaf area (mm^2). Data were collected from three independent experiments; the number of confocal images (0.32 x 0.32 mm^2) used for quantification is shown in parentheses. ******, $P < 0.01$; *******, $P < 0.001$; Dunnett's test for multiple comparisons against *At*SAR1b[H74L]. Not significant, n.s.. **(C)** Interaction analyses of the WT or GTP-locked form of *At*SAR1 with the endogenous *At*PHF1 were performed in 11-day-old *UBQ10:At*SAR1c-GFP and *dex:At*SAR1b[H74L]-CFP seedlings subjected to Pi deprivation (0 μM KH$_2$PO$_4$, 7 days). *dex:At*SAR1b[H74L]-CFP seedlings were treated with 24 hours of 10 μM dexamethasone (DEX) or DMSO. The protein amounts as input for GFP-Trap co-immunoprecipitation are as indicated above the image. An anti-GFP antibody was used to detect GFP and CFP fusion proteins. The relative intensity is shown as mean ± SE from two independent experiments.

Figure 7. Hypothetical model depicting a role for *At*PHF1 in COPII recruitment for the ER export of *At*PHT1 transporters. The GEF SEC12 recruits and activates SAR1 GTPase by exchanging GDP for GTP. The direct interaction of GTP-bound SAR1 with SEC23 recruits SEC23/24 to form the inner coat, followed by SEC13/31 recruitment to form the outer coat. The ER-resident PHF1 concentrates on ERES triggered by increased PHT1 proteins and interacts with GDP-bound SAR1. While the scaffold protein SEC16 at ERES does not interact with PHF1, the interaction of PHF1 and SAR1 may link to the regulation of SAR1 GTPase recruitment, thereby facilitating COPII assembly at ERES for the ER-to-Golgi transport of PHT1 proteins in the budding COPII vesicles.