1 **Cold hardiness dynamics predict budbreak and associated low temperature threats in**
2 **grapevine**
3
4
5 **Running Title: Cold hardiness dynamics predict budbreak**
6

7 Francisco Campos-Arguedas[1,†], Erica Kirchhof[1,†], Michael G. North[1], Jason P. Londo[2] , Terry
8 Bates[3], Cornelis van Leeuwen[4], Agnès Destrac-Irvine[4], Benjamin Bois[5] , Al P. Kovaleski[1,*]
9
10
[1]Department of Plant and Agroecosystem Sciences, University of Wisconsin–Madison, Madison,
WI, USA 53706
[2]Cornell University, Cornell AgriTech, Geneva, NY, USA 14456
[3]Cornell Lake Erie Research and Extension Laboratory, Portland, NY, USA 14769
[4]EGFV, Univ. Bordeaux, Bordeaux Sciences Agro, INRAE, ISVV, F-33882 Villenave d'Ornon,
France
[5]Biogéosciences, UMR 6282 CNRS, Université de Bourgogne, 21000 Dijon, France
18
19 † This work represents equal efforts by these authors.
20 * Corresponding author: Al P. Kovaleski, email: al.kovaleski@wisc.edu

21
22
23
24
25 **Author contributions**
26

32
33

Summary

Temperate woody perennial plants form buds that develop into leaves or flowers that emerge in

the following growth season. To survive winter, dormant buds must attain cold hardiness, and

timely lose it in spring to break bud while avoiding damage from low temperatures and late frosts.

Here, we use a process-based model to predict bud cold hardiness of three grapevine varieties (*V.*

vinifera 'Cabernet-Sauvignon' and 'Riesling', and *V. hybrid* 'Concord') from historical

temperature records of eight different locations in North America and Europe (n=329). Based on

those predictions, and thresholds of cold hardiness at budbreak from literature, timing of budbreak

was extracted. Despite the model being untrained to budbreak data, the cold hardiness model

resulted in good predictions (RMSE=7.3d) that were further improved based on expected delays

from cold damage (RMSE=7.2d). Both increasing and decreasing trends in freeze damage risk

were predicted with increasing temperature, depending on the range of mean dormant season

temperature (MDST; 1 Nov–30 Apr) in each location. Predictions in timing of spring phenology

in relation to MDST also showed warming to advance (MDST<10°C) or delay (MDST>10°C)

budbreak. Cold hardiness dynamics represents an advancement in phenological modeling that

provides information for the entirety of the dormant season, as well as budbreak.

51
52
53   **Keywords**

54
55   Budbreak, climate change, cold hardiness, dormancy, phenological modeling,
56   spring phenology, *Vitis*, winter physiology
57

Introduction

Fluctuations and extremes in temperature, photoperiod, precipitation, and numerous other environmental variables determine the distribution and adaptation of species worldwide (Inouye, 2022). In temperate ecosystems, low temperature is considered one of the most significant factors influencing the distribution of woody perennial species, particularly near the cold edge of their distribution range (Chamberlain & Wolkovich, 2021; Körner, 2021; Qian *et al.*, 2022). Seasonal fluctuations in temperature drive phenological events – the timing of which impacts the length of the growing season, overall productivity, and nutrient cycles (Richardson *et al.*, 2013; Gallinat *et al.*, 2021; Marquis & Lajoie, 2024). Understanding the timing of life cycle events such as budbreak has significant practical and ecological implications, as it informs management practices and climate adaptation strategies (Richardson *et al.*, 2013). Predicting timing of spring phenology is also an essential component of Earth system models, which help us understand and forecast ecosystem responses to climate change, including carbon sequestration potential of plants (Cleland *et al.*, 2012). Despite the importance of budbreak, accurately forecasting this critical phase remains a challenge due to our general lack of understanding of temperature responses associated with this phase in plants (Ding *et al.*, 2024). This is particularly relevant in the face of rapidly changing climates and the emergence of non-analogous environmental conditions (Darbyshire et al., 2013; Piao et al., 2019; Wheeler et al., 2024).Therefore, new models that deviate from standard and incorporate physiologically derived temperature responses may improve our predictive ability for budbreak.

Buds are formed in the late summer and early fall and remain dormant [with little developmental activity (Goeckeritz *et al.*, 2023)] throughout winter prior to growth resumption in the spring. Typical phenological models use the experimentally demonstrated negative relationship between accumulation of chill ('chilling') – cumulative exposure to low temperatures – and the accumulation of heat ('forcing') – cumulative exposure to warm temperatures – to predict budbreak (Wang *et al.*, 2020). However, there is likely an overlap in temperature ranges in which chilling and forcing occur (Flynn & Wolkovich, 2018). Therefore, the timing of transition from one phase to another can influence whether mid-dormant season temperatures are contributing to one process or the other (or occasionally both). In most phenology models, the transition from dormant to a growth-responsive state is either classified as sequential or overlapping (Chuine,

2000). In a sequential approach, both chill and heat accumulation occur in separate phases –
chilling and forcing – without accounting for their interaction (Ashcroft *et al.*, 1977). During the
first phase, dormancy release is modulated by exposure to low temperatures (chilling
accumulation) (Egea *et al.*, 2003). The second phase involves exposure to warm temperatures –
accumulation of growing degree days/hours (Fernandez *et al.*, 2021). However, delays in the
transition between the chilling and heat-accumulation phases can significantly influence the
contribution of chill and heat units, and in many cases the timing of this transition is arbitrarily
defined or imposed by the model (Egea et al., 2003; Harrington et al., 2010). These models are
therefore inherently vulnerable to shifting climatic conditions. Warming temperatures may
increase the time exposed to forcing conditions in winter and spring, which would by itself lead to
advances in spring phenology. However, the warming conditions will also decrease chill
accumulation in fall and spring, which can delay spring phenology. Therefore, variable outcomes
in phenological timing can be expected based on future warming (Chamberlain *et al.*, 2019;
Fernandez *et al.*, 2023). As a result, there is considerable interest in developing generalizable
models which can accurately describe phenology under a wide range of conditions, especially
considering the growing body of evidence supporting both phenological advancement and delay
in woody perennials due to climate warming (Hassan *et al.*, 2024) and urbanization (Li *et al.*,
2019).

Previous work has shown that budbreak can be significantly influenced by cold hardiness
dynamics (Kovaleski *et al.*, 2018; Kovaleski, 2022; North *et al.*, 2022; North & Kovaleski, 2024).
Cold hardiness is a plastic trait within a genotype mainly influenced by air temperature, with gains
(acclimation) occurring in autumn and early winter and losses (deacclimation) in late winter and
early spring. Rates of acclimation, deacclimation, and maximum levels of cold hardiness achieved
in a given location vary by genotype and yearly weather conditions (Ferguson *et al.*, 2011, 2014;
Kovaleski *et al.*, 2018, 2023; North *et al.*, 2021). Thus, cold hardiness has been shown to be an
important barrier to plant survival in cold regions, contributing to the suitability of different species
and ecotypes to different climates (Körner *et al.*, 2016; Meier *et al.*, 2018; De Rosa *et al.*, 2021;
Körner, 2021; Baranger *et al.*, 2024). Deacclimation always precedes budbreak in springtime, with
most cold hardiness lost by budbreak (Lenz *et al.*, 2013; Vitra *et al.*, 2017; Chamberlain &
Wolkovich, 2021; Hillmann *et al.*, 2021; Kovaleski, 2022; North *et al.*, 2022). Effects of cold

damage have already been linked to delays in spring phenology within the same year (Qiu *et al.*,
2024) and the subsequent year (Wang *et al.*, 2025), Altogether, this indicates that measuring or
predicting the cold hardiness phenotype and its dynamics may be useful for delivering predictions
of budbreak for species across varying climates.
125
The focus of this study was to test a process-based cold hardiness model for grapevine, NYUS.1
(Kovaleski *et al.*, 2023), for predicting budbreak phenology and exploring other climate-related
growth risks for three different cultivars across eight locations in Europe and North America.
Grapevine (*Vitis* spp.) is both a horticulturally and historically significant crop grown throughout
the world, with areas of cultivation present on all continents except Antarctica. Critically,
viticultural regions also contain the same clonally propagated genotypes (i.e., cultivars or
varieties), facilitating evaluation of plasticity across different climates. The considerable genotypic
variability in climatic adaptation among commonly grown grapevines (Wolkovich *et al.*, 2018)
makes it an interesting model to understand the connections between cold hardiness predictions
and estimation of budbreak timing. Without reparameterization, the NYUS.1 model was applied
to historical daily temperature records in each location and predictions were validated using
longstanding records of phenology observations. Additionally, we used cold hardiness trajectories
to predict instances of freeze damage across cultivars and locations, validating these predictions
based on newspaper and extension records for one location. Using model outputs, we assessed
patterns in freeze risk over both thermal and temporal scales, and estimated phenological
sensitivity as a product of dormant season temperatures.
142

Materials and Methods

Phenological Records
The phenology dataset used in this study includes three grapevine cultivars: *Vitis vinifera* cvs.
Cabernet-Sauvignon and Riesling, and *Vitis labruscana* cv. Concord (hereon referred to by cultivar
name only). These cultivars were used as the only three for which parameters are available for the
NYUS.1 grapevine cold hardiness model (Kovaleski et al., 2023). These cultivars have relatively
different cold hardiness dynamics patterns, and are representative of different groups of cultivars:
Cabernet-Sauvignon is a warm-climate variety, with low cold hardiness and low deacclimation

rates; Riesling is a cool climate variety, with intermediate cold hardiness (or high within *V.*
vinifera), and moderate deacclimation rates; Concord is a Northern hybrid, with high cold
hardiness and high deacclimation rates (Londo & Kovaleski, 2025). Phenology data was retrieved
from eight different locations across Europe and North America (**Fig. 1**): Niagara-on-the-Lake,
ON, CA (Hébert-Haché *et al.*, 2021), Portland, NY, US, Lewisburg, PA, US (Persico *et al.*, 2021),
Bordeaux, FR (Maury *et al.*, 2023), Bergheim, FR, Montreuil-Bellay, FR (Duchêne, 2019),
Marseillan-Plage, FR (Maury *et al.*, 2023), and Veitshöchheim, DE (Bock *et al.*, 2011). For
Bordeaux, FR, we included additional phenology data collected in the same plot for the years
2012-2023. For Veitshöchheim, DE, budbreak data were extracted from Bock et al. (2011) Figure
2a in RStudio using metaDigitiseR version 1.0.1 (Pick *et al.*, 2019).

The full phenology dataset contains a total of 329 observations – recorded as the Julian day of year
(DOY) of budbreak – spanning a cumulative total of 68 seasons (1955-1956 to 2022-2023). An
exhaustive list of locations, cultivars, and observational timespans for each location can be found
in **Supporting Information Table S1**. In most cases, phenology data was explicitly reported as
the day of year to reach 50% of buds at BBCH 07 and BBCH 09 [Biologische Bundesanstalt,
Bundessortenamt und Chemische Industrie; Meier et al. (2009)], or the equivalent stage 4 in the
modified Eichhorn-Lorenz scale for grapevines (Coombe, 1995). Occasionally, observations were
made of 50% of buds at stage BBCH 05 (and in one location, observations by one observer as first
bud at stage BBCH 05).

Figure 1. Locations and cultivars in the phenology dataset and cold hardiness model development.
The phenology dataset used here contains eight locations: four in France (Bergheim, Bordeaux, Marseillan-Plage, and Montreuil-Bellay), one in Germany (Veitshöchheim), two in the United States (Lewisburg, PA, and Portland, NY), and one in Canada (Niagara-on-the-Lake, ON). The NYUS.1 cold hardiness model was developed using grapevine bud cold hardiness data from Geneva, NY (Kovaleski et al., 2023), and used dormancy parameters from 15 unrelated woody perennial species grown in Boston, MA – both in the United States.

Phenology predictions

To predict phenology, we first predicted cold hardiness using the NYUS.1 grapevine cold hardiness model (Kovaleski *et al.*, 2023). This model uses temperature data as inputs to generate cold hardiness predictions as outputs.

Weather Data and Chill Accumulation

Daily maximum and minimum temperature values were retrieved for all eight locations in this study through publicly available databases. For European locations, downscaled (1 km^2 spatial resolution) data were obtained through the easyclimate package in R (Cruz-Alonso *et al.*, 2023) for the period of 1950-2022, which was the most broad timespan available using these resources. In Bordeaux, FR, additional temperature data were accessed from a nearby weather station (station ID: 335500003) for the time period of 1 January 2023 – 30 June 2023.

For two of the three North American locations (Niagara-on-the-Lake, ON, CA; and Lewisburg, PA, US), high resolution (1 km) temperature data were obtained from Daymet version 4 (Thornton *et al.*, 2021) and accessed via *daymetr* version 1.7.1 (Hufkens *et al.*, 2018) for the period of 1980-

2022, the broadest timespan available. For Portland, NY, US, temperature data from the weather

station located at the Cornell Lake Erie Research and Extension Laboratory were used (available

through the Lake Erie Regional Grape Program) for the time period of 1926-2023. We primarily

used gridded climate datasets to ensure consistent, quality-controlled temperature records across

sites and years, incorporating station-based observations only where high-resolution, long-term

data were available.

For evaluations using general yearly weather, such as sensitivity (see below), we characterized

each location:year combination using the mean temperature for the dormant season calculated by

obtaining the average temperature from 1 November to 30 April for each season within each

location, as this is the period of time when most shifts in cold hardiness occur. This is referred to

simply as mean dormant season temperature.

Modeling Cold Hardiness

The model utilized in this study is the NYUS.1 cold hardiness prediction model (Kovaleski *et al.*,

2023) (see **Notes S1** for a detailed description of the model). NYUS.1 integrates (1) *Acclimation*

and (2) *Deacclimation* processes over daily time steps from bud set to any given point during the

dormant season, using daily temperatures as an input:

(1) *Acclimation*: The rate of acclimation for buds at any given day is a function of chilling

accumulation, the difference between the daily minimum temperature and the maximum

temperature that elicits chilling accumulation on any given day, and the cold hardiness of

the previous day. Generally, (i) as chilling accumulates, temperatures must be lower to

result in acclimation, (ii) the lower the temperature, the more cold hardiness it can elicit,

and (iii) the more cold hardy buds are on the day before, the less cold hardiness they can

gain when exposed repeatedly to the same temperature (e.g., a very cold hardy bud exposed

to mild temperatures in late winter will not gain any further cold hardiness).

(2) *Deacclimation*: The effective rate of deacclimation at any given day is determined by a

temperature response, calculated for daily minimum and maximum temperatures, which is

modulated by a chilling response curve – the deacclimation potential. Generally, (i) the

effective rate of deacclimation is the product of the rate of deacclimation and the

deacclimation potential, where (ii) higher temperatures result in higher rates of

deacclimation, and (iii) high chilling accumulation leads to higher deacclimation potential

(e.g., a bud exposed to high temperatures at low levels of chilling accumulation will not

lose any cold hardiness).

Chilling accumulation is an input within NYUS.1. There are many possible models used to

calculate chill as a response to temperature [see Wang *et al.* (2020) for 12 examples of models].

The responses to chilling accumulation within NYUS.1 are based on chill portions calculated

through the Dynamic Model [Fishman *et al.* (1987a,b); model C_9 within Wang *et al.* (2020)], which

therefore serves as a sub-model. The Dynamic Model was used as there was no reparameterization

of the NYUS.1 model, and thus the parameters used to describe dormancy progression are based

on chill portions. The Dynamic Model is a complex chilling accumulation model, often left

undescribed [e.g., within Wang *et al.* (2020)]. Generally, temperatures between ∼−2 and 13 °C

promote chilling accumulation of chill portions. Once a chill portion is accumulated, it is not lost,

but warm temperatures can cause loss of a precursor of a chill portion. However, fluctuations

between low and warm temperatures can slightly enhance chilling accumulation. Chilling

accumulation was calculated in RStudio using *chillR* version 0.75 (Luedeling *et al.*, 2023). Since

the Dynamic Model calculates chill portions in hourly timesteps, daily minimum and maximum

temperatures were transformed into hourly data using the "*stack_hourly_temps*" function in *chillR*.

Chilling accumulation for each location was calculated continuously from 1 July to 30 June for

every year where temperature data was available. To summarize, the cold hardiness of buds at any

given time is calculated as an integration of functions of temperature over time, which are

modulated by chilling accumulation, which is itself a response to temperature. In this process,

acclimation is the strongest driving force for cold hardiness during the fall, and deacclimation is

the strongest driving force during spring, resulting in a general U-shaped curve for cold hardiness.

The 50[th] percentile cold hardiness predictions used the same parameters originally published in

Kovaleski et al. (2023) and are based on field data collected in Geneva, NY, US. To provide

additional detail regarding potential damage, we used the same calibration and validation datasets

to obtain parameters for the 10[th] and 90[th] percentiles for cold hardiness predictions. These

parameters are described in **Supporting Information Table S2**.

Spring Cold Hardiness

Minimum cold hardiness thresholds during spring are not set within the NYUS.1 model, allowing for the projection of cold hardiness freely towards positive temperature values, which here we use for predictions of budbreak (see below). For analyses concerning the estimation of frost damage risk and safety margins, we limited the cold hardiness to thresholds which describe the minimum cold hardiness of growing tissue, obtained from the literature (Gardea, 1987). Such thresholds were obtained by Gardea (1987) through controlled freeze tests, evaluating damage on buds at "fourth flat leaf" stage (BBCH 14) that were exposed to 1ºC temperature steps between 0ºC and -4ºC, and a +4ºC control. Here, we assumed that newly emerged tissues in springtime reach a minimal level of cold hardiness, that this level is then maintained constant over time, and that these values for grapevine do not differ on a genotype-specific basis [as suggested by the little effect of continent of origin reported for species within a genus; see Kirchhof et al. (2025)].

Cold Hardiness at Budbreak

Projected cold hardiness trajectories for each cultivar were used to predict the day of year by which 50% of buds reached the BBCH 07 ("beginning of bud burst: green shoot tips just visible") stage. Two different thresholds for the cold hardiness at budbreak were applied, based on cultivar: a predicted cold hardiness of +10ºC was used for Cabernet-Sauvignon and Riesling, and +5ºC was utilized for Concord. Thresholds were determined from data within Kovaleski et al. (2018) in which rates of cold hardiness loss in forced deacclimation assays were related to different bud developmental stages.

Additional threshold adjustments were made to account for differences in observational habits within phenological records. The first adjustment was used to account for years and locations where phenology observations were determined at the 50% BBCH 05 ("wool stage") stage rather than at BBCH 07. Because buds at BBCH 05 are more cold hardy, we used a +5ºC threshold so that the predicted cold hardiness values aligned with the developmental stage represented in the observation for Riesling and Cabernet-Sauvignon (North & Kovaleski, 2024). An additional adjustment was made specifically for predictions in Bordeaux, FR over the 1981-2008 period. During this time, the phenological record for budbreak was the day of year when the first bud for

each cultivar reached BBCH 05 (as opposed to 50% of buds at a certain stage as in other locations

and years). Because the first bud to reach BBCH 05 corresponds to an earlier developmental stage

with substantially lower cold hardiness, we used a threshold of 0°C for Cabernet-Sauvignon and

Riesling predictions during the 1981-2008 period to align the model output with the stage

represented in the observational habits (see **Supporting Information Fig. S1** for a schematic

representation of threshold setting and use).

Adjustment of Cold Hardiness Predictions to Account for Damage

In years where low temperature damage to buds was predicted, a correction was applied to the cold

hardiness of the remaining buds based on an estimate of the degree of damage (as a percentile

value). This was achieved through a simple linear interpolation. Damage estimates were performed

for days within each season where the minimum air temperature was lower than the predicted cold

hardiness. In seasons where multiple instances of damage were predicted, percentile values for

damaged buds were calculated for each event, based on the existing population immediately prior.

The cold hardiness of the remaining buds was calculated after each damage event, dependent on

both the existing cold hardiness percentile values preceding the event and the percentile estimation

of buds damaged from the freeze event.

When the minimum air temperature fell between the 10th and 50th percentile cold hardiness values,

the following equation was used to estimate the percent of buds damaged from the existing

population:

$\%damage = \left(\frac{50-10}{CH_{50}-CH_{10}}\right) \times (T_{min} - CH_{10}) + 10$; (Eq. 1)

where T_{min} is the minimum temperature of the day, and CH_{10} and CH_{50} are the 10th and 50th

percentile cold hardiness values. When the minimum air temperature fell between the 50th and 90th

percentile cold hardiness values, the following equation was used to estimate the percent of buds

damaged from the existing population:

$\%damage = min\left(\left[\left(\frac{90-50}{CH_{90}-CH_{50}}\right) \times (T_{min} - CH_{50}) + 50\right], 90\right)$; (Eq. 2)

where CH_{50} and CH_{90} are the 50[th] and 90[th] percentile cold hardiness values, and the constants (e.g.

10, 50 and 90), represent the corresponding quantiles associated with each cold hardiness value.

After calculating the estimated percent of buds that were damaged, cold hardiness percentile values

were adjusted to account for the buds remaining in the undamaged population using the following

equations:

$CH_{10}^* = \begin{cases} CH_{10} + \left(\frac{CH_{50}-CH_{10}}{40}\right) \times \left(\left((100 - \% \; damage) \times 0.1 + \% \; damage\right) - 10\right), when \; \%damage \leq 50\% \\ CH_{10} + \left(\frac{CH_{90}-CH_{10}}{80}\right) \times \left(\left((100 - \% \; damage) \times 0.1 + \% \; damage\right) - 10\right), when \; \%damage \geq 50\% \end{cases};$

(Eq. 3)

$CH_{50}^* = CH_{50} + \left(\frac{CH_{90}-CH_{50}}{40}\right) \times \left(\left((100 - \% \; damage) \times 0.5 + \% \; damage\right) - 50\right);$ (Eq. 4)

$CH_{90}^* = CH_{90} + \left(\frac{CH_{90}-CH_{50}}{40}\right) \times \left(\left((100 - \% \; damage) \times 0.9 + \% \; damage\right) - 90\right);$ (Eq. 5)

Where CH_{10}^*, CH_{50}^*, and CH_{90}^* are the damage-adjusted predicted cold hardiness for the 10[th], 50[th],

and 90[th] percentiles, respectively. Damage-adjusted cold hardiness predictions were used in all

analyses presented unless otherwise mentioned (see **Notes S2** for an example illustrating the

application of the equations 1-5 described above).

Validation of Damage Predictions

Given the absence of long, publicly available records of empirically determined cold hardiness

measurements, we validated our predictions of cold damage by comparing them with historical

reports. To do so, we performed this analysis based on Concord cold hardiness modeled in one

location (Portland, NY) over the entire time period of cold hardiness predictions (1926-2023). Our

analysis focused on Concord in the Lake Erie region given its local economic importance,

guaranteeing mentions of cold damage in a local news source, and the availability of searchable

news reports in English. The historical records used were obtained from two sources: *The Erie*

Daily Times (1927-2006) – a local newspaper – and *Veraison to Harvest* (2007-2023) – an

extension publication by Cornell University which reports on New York State's grapevine

production. For newspaper articles, the archival search was done using the keywords 'grape+frost'.

In order to be included in the analysis, the articles and extension reports had to meet the following

criteria: (i) there was a mention of any degree of frost damage to the grapevine crop or (ii) articles

explicitly mentioned that the crop escaped damage. If no records of damage were found for a given

year, it was assumed that no damage occurred to the buds.

A positive detection of damage for a season was considered when at any day of the season the

daily minimum temperature was lower than the predicted 10^{th} percentile cold hardiness for that

day. A binary outcome was then applied to all years in this analysis, regardless of level of damage,

or number of damage events, where 1 indicated damage occurrence during a season, and 0

indicated no damage. A binary outcome was also applied to the mention of damage in records,

wherein 1 indicated presence (records indicated damage occurred within a season) and 0 indicated

absence (records indicated no damage had occurred, or no records of damage were discovered for

a given season). These binary outcomes were then used to construct a confusion matrix to evaluate

prediction performance and calculate sensitivity, specificity, precision, and accuracy of the model.

Effect of Chilling Temperature Range

To examine potential errors associated with chilling range on the budbreak predictions, both as a

whole and by location, we applied a series of phase shifts to change the relative contribution of

temperatures to chilling accumulation in the Dynamic Model. The chilling range was thus shifted

by 1ºC intervals, from -6ºC to +6ºC during chill accumulation estimation (e.g., chill is maximized

at 7.5 ºC within the Dynamic Model, and we shifted this maximum to occur between 1.5 ºC and

13.5 ºC). Budbreak predictions were then produced using the model without any other adjustments.

RMSE for predictions at each phase shift step were then calculated.

Indicators for Climate Suitability and Sensitivity

To evaluate the temporal nature of freeze risks within a season, we used temperature records from

Portland, NY, over the time period of 1926-2023, to predict cold hardiness and subsequently

estimate the frequency of freeze events and potential freeze damage. Safety margins defined as the

difference between the minimum registered air temperature (T_{min}) and the 50^{th} percentile cold

hardiness prediction were calculated daily (CH_{50}):

$Safety\ margin = T_{min} - CH_{50}$; (Eq. 6)

Where a safety margin below zero indicates at least 50% freezing damage is predicted, and positive

values indicate different levels of safety. We defined freeze risk as the number of days when the

safety margin dropped below zero for each cultivar in all locations for all years in which weather

data was available (n=1626 location:cultivar:year combinations). Therefore, safety margin

calculations were also used to determine the number of freeze risk injury days over the range of

mean dormant season temperatures experienced (November-April), providing an overall measure

of how many days during the season there was a risk of damage:

$N_{days} = \Sigma_{i=1}^{n} I(Safety\ margin_i \leq 0)$; (Eq. 7)

The potential effects of temperature changes on phenological timing (sensitivity) were evaluated

by studying the relationship between the predicted day of budbreak and the mean dormant season

temperatures over all locations, years and cultivar combinations (n=1626). Differences in

responses to different mean dormant season temperatures for the entire dataset, and also in three

temperature segments (a) -1ºC to 3ºC, b) 3.01ºC to 10ºC, and c) 10.01ºC to 12ºC) were calculated

in RStudio using the *rms* package, version 6.8-1 (Harrell & Frank, 2024).

Evaluation of budbreak predictions
Accuracy and performance of the budbreak prediction model, and the effect of different variables

(i.e., observer, cold damage, and chill accumulation) were evaluated through determination of

Bias, Root Mean Square Error (RMSE), and Coefficient of determination (r^2), comparing predicted

budbreak day of year to the effectively independent observations of field budbreak day of year. To

evaluate bias, we calculated residuals as the difference between predicted and observed budbreak

day of year ($Residuals = BB_{predicted} - BB_{observed}$). Bias was then defined as the mean of

residuals for each group. Here we used predicted minus observed so that negative values indicate

early predictions, while positive values indicate late predictions. RMSE was calculated to quantify

the overall prediction error, reflecting both the variance and bias of the model.

To evaluate the effect of observational habit changes in Bordeaux, FR, we tested residuals by

observer (three observers) using a one-way ANOVA, with Tukey-adjusted post hoc comparisons

of means. Residuals were compared by observer before and after applying a new threshold for cold

hardiness at budbreak. To evaluate the effect of predicted damage on predictions of budbreak, we

compared three groups to each other: "no damage", "damage uncorrected", and "damage

corrected". Residuals for each pair were compared using a one-way ANOVA. All statistical

analyses were performed using RStudio. A workflow of the analyses performed here is presented

in **Supporting Information Fig. S2**.

Results

Cold hardiness can be used to predict budbreak phenology across cultivars & locations

We calculated cold hardiness for all locations, and for each year in which we gathered weather

data, we used the projected cold hardiness to predict budbreak (**Figs. 1, 2**). The initial budbreak

predictions across locations and cultivars yielded results (n=329, RMSE=7.3 days, bias=-0.8 days,

r^2=0.88; **Fig 3a**) that are comparable to those reported by other phenology modeling studies (García

de Cortázar-Atauri *et al.*, 2009; Sgubin *et al.*, 2018) **(Notes S3).** Given that cold hardiness trajectories

were used to determine budbreak timing (**Figs. 2, 3b**), it was also possible to group predictions

based on the presence or absence of freeze damage to buds. Although our initial predictions did

not account for effects of cold damage, given the known delay that occurs in budbreak when cold

damage occurs in buds (Chamberlain & Wolkovich, 2021; Charrier *et al.*, 2021; Muffler *et al.*,

2024a) we produced corrected budbreak predictions whenever damage occurred prior to initially

predicted budbreak. A freeze damage event was predicted to occur when the minimum air

temperature dropped below (i.e., was more negative than) the cold hardiness prediction associated

the 10[th] percentile of cold hardiness distribution on any given day. For each individual freeze event,

a simple linear interpolation was used to determine the proportion of buds damaged, and a new

80% confidence interval was calculated to represent the cold hardiness of the remaining population

of living buds. Based on the greater cold hardiness in the remaining buds, budbreak predictions

shifted to a later date following freeze damage, with the estimated delay dependent on the

magnitude of cumulative freeze damage throughout the dormant season (**Fig. 3b**).

439
Figure 2. Conceptual overview of the cold hardiness model to predict budbreak. The NYUS.1 cold
hardiness model (Kovaleski *et al.*, 2023) is used to predict the time to budbreak for *V. labrusca* 'Concord'
grapes grown in Portland, NY, US in the 1983-1984 dormant season. Cold hardiness (blue ribbon) is gained
and lost in response to seasonal fluctuations in air temperature (grey ribbon). Two different end trajectories
are shown for the cold hardiness prediction: (1) observed cold hardiness values, which remain negative and
stop decreasing once they reached the minimum cold hardiness reported by Gardea (1987) (blue ribbon);
and (2) projected cold hardiness values, where the model continues to predict loss of cold hardiness into
positive temperature values. Although these do not reflect real values of cold hardiness, they are used for
budbreak predictions (red ribbon and dashed line). When the projected cold hardiness reaches a threshold
determined for cold hardiness at budbreak [established from data in Kovaleski et al. (2018)], budbreak is
predicted to occur.

Grouping of model residuals based on predictions of freeze damage revealed that errors were

significantly influenced by the presence or absence of such damage (No damage vs. Damaged

uncorrected; **Fig. 3c**). In years where damage was predicted to occur, uncorrected budbreak

predictions were on average 2.7 days earlier than observed budbreak (i.e., $Bias_{damaged, uncorrected} = -2.7d$),

significantly contrasting from years where no damage was predicted ($Bias_{no\ damage} = +0.3$ days).

After accounting for damage in budbreak predictions, error in years where damage was corrected

did not significantly differ from years without damage (No damage vs. Damaged corrected; $p =$

0.527). As a result of incorporating a method for predicting and accounting for damage in the

budbreak predictions, overall model outputs improved (RMSE=7.2 days, bias=0.6 days, r^2=0.88;

Fig. 3d).

Because long-term cold hardiness measurements were unavailable, we used historical reports from

extension documents and newspapers for one location as a proxy for evaluating the model (**Fig.**

3e). While these records likely underreport damage, they provide the only multi-decadal criterion

against which to compare predictions, allowing us to assess whether NYUS.1 captures years with

widespread, well-documented events. This approach classified reported damage in the records as

a binary variable (damage vs. no damage). Years were classified as either "damage" or "no

damage", where any mention of damage to grapevine buds was determined to be a "damage" year,

and years where damage was not mentioned were recorded as "no damage" years. The analysis

was performed using Concord cold hardiness predictions in Portland, NY, US over the time period

of 1926-2023 (n=97). The binary validation of damage predictions over the time period where cold

hardiness predictions were made (1926-2023) resulted in model sensitivity of 0.75, specificity of

0.72, precision of 0.57, and accuracy of 0.73 (**Supporting Information Fig. S3**). Sensitivity and

specificity values above 0.70 suggest that the model is reasonably effective at correctly identifying

both positive and negative cases, while the lower precision highlights that some false positives

remain (Bujang & Adnan, 2016) – although some false positives here may be due to underreporting

of damage. Based on the evidence that (i) damage-corrected budbreak predictions better reflect

phenological observations and (ii) predictions of damage are also reflected in historical records,

corrected predictions were incorporated into all further analyses presented. Hereafter, these are

then referred to as freeze damage-corrected budbreak (FDC budbreak).

483
Figure 3. Comparison of budbreak predictions from cold hardiness model with observations. (a)
Relationship between initial predictions and observations of budbreak. (b) Based on expected cold damage
prior to budbreak, predictions of budbreak can be corrected, shifting to a later date as the cold hardiness of
the surviving bud population is greater. (c) Distribution of residuals based on five different groupings, from
left to right: initial predictions (years uncorrected for damage + years where damage was not predicted n =
329); years where no damage was predicted (n = 200), uncorrected predictions in years with predicted

damage (n = 129), predictions corrected to account for damage (n = 129), and full, corrected predictions

(predictions corrected to account for damage + predictions in years where damage is not predicted, n =

329). Residuals for each pair were compared using a one-way ANOVA. (d) Comparison of

predictions to observations after accounting for the consequences of freeze damage. (e) Sample article from

historical records used to qualitatively validate damage predictions (Erie Daily Times, 5 May 1977). In

(a,b), thin solid black line represents 1:1 ratio between observations and predictions, and dotted lines

represent an interval of +/- 7 days from 1:1 line. Heavy black line shows linear relationship between

predicted and observed points.

Observational habits and chilling as significant sources of error

Using FDC budbreak, we explored the effect of different factors contributing to uncertainty in the

model outputs using partial residual plots. We found observer – based on observational habits – to

be a significant source of uncertainty (**Supporting Information Fig. S4**). Year did not show a

significant effect on residuals, and other factors such as minimum yearly temperature in any given

location, maximum predicted cold hardiness, and chill accumulation showed effects in only one

cultivar (**Supporting Information Fig. S5**). Importantly, the effect of chilling on residuals showed

a positive (albeit not significant for two cultivars) slope for the three cultivars when combined,

indicating systematic issues with chill accumulation – be that through modeling of accumulation,

or how chilling is incorporated in the cold hardiness model. However, the predictions using

NYUS.1 remained robust even under climatic conditions that differ substantially from those where

the model was developed – particularly warmer locations with lower chill accumulation (**Fig. 4,**

Supporting Information Fig. S6).

Observed date of budbreak in Bordeaux, FR between 1981-2008 were recorded based on the date

of first bud to reach BBCH05, or first bud to break, differing from the rest of the record in which

dates were recorded based on 50% budbreak to either BBCH05 or BBCH07. We accounted for

these differences by decreasing the cold hardiness threshold at budbreak (see **Supporting**

Information Fig. S1), allowing predictions to better reflect the earlier stage of development that

better corresponds the observations made, reducing this factor to a non-significant source of error

(**Supporting Information Fig. S4**).

Chilling accumulation – calculated via the Dynamic Model (portions) and used to modulate both

cold acclimation and deacclimation as responses to dormancy – led to a tendency for late

predictions at the lower range found in our datasets (i.e., between 85 and 105 portions, $p<0.001$;

Supporting Information Fig. S5d). Chill portions are accumulated when temperatures fall within

a given range, with the amount of chilling accumulated dependent on the frequency and relative

contribution of those temperatures to chilling (**Fig. 4a**). Therefore, chilling accumulation

throughout the dormant season varies based on the climate of a region and yearly weather patterns.

We explored shifting the range of temperatures contributing to chilling to test whether a simple

correction would reduce the model residuals. Both increasing and decreasing the temperature range

for chill accumulation increased the RMSE of predictions across all regions (**Fig. 4b**). These

changes, however, are uneven when observed within climates. In colder locations, where high

levels of chill accumulate within a season due to wide temperature distributions, small shifts in

chill accumulation range cause virtually no change in predictions (Portland, NY, US in **Fig. 4c,d**).

However, in warmer locations where chill accumulation is lower, small shifts towards colder range

of chill accumulation results in significant delays in predicted budbreak (Marseillan-Plage, FR in

Fig. 4c,d). Large shifts towards colder temperature greatly reduce the amount of chill accumulated

in warmer locations, significantly affecting budbreak predictions in warmer locations, whereas

these effects remain minor in cold areas (**Fig. 4e**).

539
540
Figure 4. Phenology predictions in different sites are differentially affected by shifts in how chilling

accumulation is calculated. (a) Density plot of recorded temperatures in Portland, NY, US and Marseillan-

Plage, FR, and the relative contribution of temperatures to chilling accumulation in the Dynamic Model

under unaltered and two sample phase-shift scenarios (-2ºC and -6ºC). (b) RMSE of budbreak phenology

predictions under a range from -6ºC to +6ºC of phase shift scenarios. (c,d,e) Phenology predictions

compared to observations and average chilling accumulation in a cold location (Portland, NY) and warm

location (Marseillan-Plage, FR) under (c) unshifted conditions, (d) a -2ºC phase shift, and (e) a -6ºC phase

shift scenario. In (c,d,e), thin solid black line represents 1:1 ratio between observations and predictions, and

dotted lines represent an interval of +/- 7 days from 1:1 line. Heavy black line shows linear relationship

between predicted and observed points. Horizontal dashed line at DOY 150 in (c,d,e) is used to highlight

changes in axis in (c-e). Points shown as arrowheads in (e) indicate failed predictions where budbreak

would occur over a year after the start of the model.

Cold hardiness as a driver of climate suitability

Because cold hardiness is a plastic trait that responds to temperature, shifting temperature regimes

will influence cold hardiness dynamics to varying degrees, leaving some genotypes more

vulnerable to freeze risk during the winter and early spring months. Therefore, we sought to

examine the ways in which a cold hardiness model (here NYUS.1) can be utilized to explore low

temperature-related growth risks among the three cultivars. To do so, we expanded cold hardiness

and budbreak predictions beyond the year:location:cultivar combinations in which phenology

observations were available. Instead, for all locations, we estimated cold hardiness and budbreak

for the three cultivars, over the range of time for which weather data was available for each location

(1950-2023 for European locations; 1926-2023 for Portland, NY, US; and 1980-2022 for all other

North American locations).

We first examined predicted freeze risk amongst the three cultivars over the season for a cold

location where the longest range of predictions were available (97 seasons; Portland, NY), and

where cold damage is often reported (**Supporting Information Fig. S3**). This was done based on

safety margin calculations as the difference between the minimum temperature and the CH_{50} in

any given day (Safety margin = $T_{min} - CH_{50}$). Instances of both predicted damage [Safety margin

≤0)] and near damage (0<Safety margin ≤2) were calculated for each cultivar. All seasons were

aligned based on each cultivar's specific FDC budbreak for a given year, and showed different

patterns in freeze risk for each cultivar (**Fig 5a**). Cabernet-Sauvignon faced a higher risk of winter

damage (i.e., ~120 to 80 days before budbreak) and a reduced risk of spring damage (i.e., 10 days

prior to and after budbreak) when compared to other cultivars. Conversely, Concord displayed low

risk of winter damage, but a greater susceptibility to spring damage compared to the other two

cultivars. Riesling showed an intermediate risk for both winter damage and spring damage.

To more easily generalize responses to current and future climates, we explored the influence of

dormant season temperature (November-April) on freeze damage risk as the number of days within a season where minimum temperature was lower than the predicted CH_{50} (negative safety margin). For this, predicted cold hardiness for the three cultivars across the eight locations and for all years was used. Cultivar-dependent patterns in risk became evident: colder dormant seasons (mean temperature below ~2.5ºC) resulted in a greater freeze risk for Cabernet-Sauvignon and (to a slightly lesser degree) Riesling, while Concord exhibited negligible risk (**Fig. 5b**). With moderate temperatures (e.g., 5ºC), all cultivars exhibited some level of freeze risk, with risk being highest for Concord, and lower for Riesling, and Cabernet-Sauvignon. When evaluating all cultivars together, we can separate freeze risk broadly into four different temperature bands (**Fig. 5c**). Mean winter and spring temperatures below 1ºC were considered high freeze damage risk temperatures. Moderate freeze damage risk was assigned to mean temperatures ranging from 1ºC to 3ºC. Increased freeze damage risk was identified for temperatures ranging from 3ºC to 8ºC, and temperatures above 8ºC were classified as low freeze damage risk.

Figure 5. Freeze risk shows temporal, temperature, and cultivar effects. (a) Damage risk density plot over time within a season, based on cold hardiness predictions for three grapevine cultivars in Portland, NY, US over the time period of 1927-2023 shows two instances of higher probability of damage: midwinter (120-80 days before budbreak) and around budbreak (15 days before and after budbreak). Red curve represents predicted instances of freeze damage (safety margin less than or equal to 0), and the yellow curve

represents instances where there is a narrow safety margin (between 0ºC and 2ºC). Horizontal axis shows "0" as the mean day of budbreak for each cultivar over the time period. (b,c) Number of predicted freeze damage risk days based on mean dormant season temperature (1 November–30 April), for (b) each of three grapevine cultivars grown in eight locations and (c) for all cultivars combined. Point sizes (b,c) are proportional within each decimal mean dormant season temperature. Risk of damage was separated in four temperature bands based on mean dormant season temperature: [1; blue] high freeze risk injury, particularly for *V. vinifera* cultivars, due to low mid-winter temperatures; [2; green] cool spring days delay budbreak until last frost; [3; orange] risk of freeze damage increases as spring phenology advances faster than date of last frost; [4; red] temperatures are too warm for low temperature injury to be significant. Horizontal colored lines (**c**) show mean dormant season temperature ranges in datasets from each of the eight locations studied.

We also analyzed the sensitivity of simulated budbreak based on mean temperature of the dormant season using the predicted FDC budbreak for all cultivar:year:location points (**Fig. 6a**). At any mean dormant season temperature, Concord has earlier budbreak, followed by Riesling, and by Cabernet-Sauvignon. Date of budbreak advances with increasing mean dormant season temperatures, with sensitivity of –5.8 days/ºC (i.e., 5.8 earlier budbreak DOY with every degree of warming) for all cultivars combined. At the cultivar level, Concord has a slightly faster advancement compared to Riesling and Cabernet-Sauvignon (–6.0 days/ºC, –5.5 days/ºC, and –5.7 days/ºC, respectively). However, the overall response appears to be non-linear, with varying patterns across different temperature ranges and locations. To highlight these differences, mean winter and spring temperatures in Portland, NY, US (one of the coldest locations in our study) and Marseillan-Plage, FR (the warmest location in our study) were explored (**Fig. 6b**). Portland shows an overall phenology advancement of only –3.3 days/ºC, with mean dormant season temperatures ranging from -1ºC to 5ºC (Fig 6B). This contrasts with Marseillan-Plage, FR, where mean dormant season temperatures range from 8ºC to 12ºC, showing a shifting pattern within its temperature range. There is a phenology advancement of –6.8 days/ºC when mean dormant season temperatures ranges from 8ºC to 10ºC. However, as mean temperatures increase further in that location, phenology starts to delay at a rate of +3.8 days/ºC when mean dormant season temperatures are above 10ºC (**Fig. 6b**). When data for all locations is split into three separate groups, sensitivity is highest in moderate temperatures (–6.3 days/ºC for 3-10ºC), and lowest and positive for warmest temperatures (+2.0 days/ºC for 10-12ºC; **Fig. 6a**).

632
Figure 6. Sensitivity of phenology to temperature is uneven across the range of climates studied. (a,b)
Freeze damage corrected predicted day of budbreak for all cultivars as a response to mean dormant season
temperatures (1 November–30April) for (a) all eight locations combined and (b) separately for Portland,
NY, US and Marseillan-Plage, FR. (a) Budbreak advances at an overall rate of 5.8 days/°C for all cultivars,
while showing different magnitudes and directions at different temperature intervals, (b) which is also
affected by local temperature dynamics.
639

Discussion

For temperate woody perennials, spring phenology (here budbreak) occurs after plants have been

released from an annual period of winter dormancy (Lang *et al.*, 1987). Time spent in cold

temperatures, quantified as chilling accumulation, is required for dormancy release. Then, to break

bud and resume growth, time spent in warm temperatures (forcing) is required (Lang *et al.*, 1987;

Chuine & Régnière, 2017). Many attempts to model budbreak rely at least partially on this

relationship, using quantifiable thermal time metrics to elucidate the time required to reach the

stage (Chuine, 2000). Here, the negative chilling and forcing relationship is substituted by winter

dynamics of cold acclimation and deacclimation, thus incorporating a third temperature response

– cold hardiness – to accurately predict spring phenology while allowing for better understanding

of cold temperature-related suitability. Our findings demonstrate that a process-based cold

hardiness model for grapevine [NYUS.1; Kovaleski et al. (2023)], trained on cold hardiness data

for a single location, is generalizable to predict spring phenology (as budbreak) across a wide range

of global climates in which grapevines are typically cultivated (**Figs. 1,3**). Our analysis also shows

that the NYUS.1 model effectively predicts freeze events (**Supporting Information Fig. S3**),

while highlighting genotypic differences in damage patterns across locations and climatic

conditions (**Fig. 5**). These results emphasize the importance of incorporating cold hardiness as a

predictable trait within a phenology model, offering valuable insights beyond just predicting

budbreak that will be consequential as global climate conditions continue to rapidly change.

Cold hardiness has a u-shaped pattern (**Fig. 2**), where its level at any point depends on prior

temperatures experienced and specific species or cultivar. Here we use a cold hardiness threshold

to predict budbreak. Given that colder climates and colder seasons can elicit greater cold hardiness

(**Supporting Information Fig. S7**), different locations can require different degrees of

deacclimation (or forcing) to reach budbreak. This variation helps explain why the same genotypes

show different chilling and heat requirements when evaluated in different locations. Additionally,

reacclimation is possible if colder temperatures occur after a period of deacclimation [i.e., **Fig. 2**,

see deacclimation period in late February, followed by reacclimation in early March; Kalberer et

al. (2006), Arora & Rowland (2011)]. In this context, reacclimation could be considered analogous

to what would be negative forcing, or negative growing degree-days in ordinary spring phenology

models – a concept contrary to growth, and thus never used, yet applicable and realistic within a

cold hardiness dynamics framework.

The suitability of woody perennial species to local climates is shaped by numerous interacting

environmental factors and phenotypic plasticity (Baranger *et al.*, 2024). Among these, exposure to

low temperatures requires particular attention, especially in light of the unprecedented rise in

global temperatures (Masson-Delmotte *et al.*, 2021). Increasing temperatures have been shown to

cause increased risk of late frosts in some areas (Augspurger, 2013; Sgubin *et al.*, 2018; Vitasse

et al., 2018). Our study found that all cultivars exhibited advancement in phenology with warmer

dormant season temperatures (**Fig. 6**), consistent with broader trends in Northern Hemisphere

populations (Li et al., 2019; Piao et al., 2019; Fernandez et al., 2023; Li et al., 2024). Here we used

mean dormant season temperature (Nov-Apr) as NYUS.1 has a complete overlap of chilling and

forcing phases (i.e., acclimation and deacclimation are possible at any time), thus making it not
possible to separate temperatures for each phase. However, it is important to note that warmer
winter temperatures can also lead to increases in freeze damage potential, and the trends depend
on cultivar [**Fig. 5b**; or similarly provenance in Mura et al. (2022)] and where a location lies in
terms of mean dormant season temperature (**Fig 5c**). Our study found that a general pattern occurs,
where risk generally decreases with increasing mean temperatures, but a "bump" appears between
~3ºC and 8ºC. This "bump", represents a mismatch between phenology advances due to warmer
springs and the potential for significant late frosts, increasing the freeze damage risk. When
looking at each cultivar separately, the phenology-temperature mismatch places Concord at a
greater risk for spring frost damage, including the timing right before budbreak, whereas Riesling
and Cabernet-Sauvignon have lower risk (**Fig. 5a,b**). This further demonstrates the advantage of
using a cold hardiness model to predict spring phenology, allowing for simultaneous assessments
of both phenology advances and freeze damage risk that includes pre-budbreak damage.

In addition to risks related to freeze damage, excessive warming reduces chilling accumulation
and delays budbreak based on our predictions (**Figs. 4,6**). Recent work supports these findings,
suggesting that delays in phenology are also possible due to reduced exposure to temperatures
which contribute to chilling accumulation (Kramer *et al.*, 2000; Benmoussa *et al.*, 2017; Gaeta *et*
al., 2018; Li *et al.*, 2019; Fernandez *et al.*, 2023; Mo *et al.*, 2024), something often observed in
areas with marginal chilling conditions where temperate woody perennial crops are introduced
(Malagi *et al.*, 2015; Du *et al.*, 2019; Song *et al.*, 2020; Zeng *et al.*, 2023) and empirically
determined in general chilling-forcing experiments [e.g., Flynn & Wolkovich (2018); Kovaleski
(2022)]. As changing global temperatures shift spring phenology, woody perennials will face novel
risks in their current distributions (Chmura et al., 2011; Ford et al., 2016; Baranger et al., 2024;
Stuke et al., 2024). The degree to which phenology shifts among the different genotypes in a given
location is ultimately an effect of their plasticity and adaptive capacity on separate aspects of
winter physiology: cold hardiness, chilling, and forcing. This highlights the importance of
expanding empirical work with cold hardiness and chilling to include diverse genotypes and
populations, particularly in warm range edges of natural ecosystems and marginal agricultural
conditions, where plants will be more strongly affected (**Fig. 4**).

Grapevine cultivars possess a great diversity in traits that affect responses to climatic conditions
(Flynn & Wolkovich, 2018; Morales-Castilla *et al.*, 2020). This diversity is evident when exploring
cold hardiness dynamics across the cultivars presented in this study. Across all locations, predicted
patterns in acclimation and deacclimation reflected known genotypic differences in hardiness and
development, and are therefore reflective of current distributions of the genotypes studied (**Fig. 1**).
Cabernet-Sauvignon and Riesling – both cultivars of *V. vinifera* – exhibit low and intermediate
mid-winter freeze tolerance (**Figs. 5a, S7**), respectively. These cultivars are thus best suited to
climates with moderate to mild winters and longer growing seasons that are presently characteristic
of many temperate climates with winters moderated by large bodies of water and western European
locations. Concord, a cultivar of the more cold hardy *V. labruscana,* showed the greatest freeze
tolerance among the three genotypes here, making it best adapted to regions with the cold winters
and shorter growing seasons, presently characteristic of many viticultural regions in Northeastern
North America. When freeze risk is evaluated based on mean dormant season temperature,
Concord shows higher risk at moderate temperatures due to its faster deacclimation and advanced
budbreak compared to the other cultivars (**Figs. 5c, 6a**). Genotypic adaptation to local climates has
allowed for successful grapevine cultivation in a wide range of climates, and our assessment of
spring damage risk supports these known differences. Understanding the degree to which cold
hardiness dynamics respond to fluctuations in dormant season temperatures – and, ultimately, how
this influences budbreak phenology – will be imperative to making informed management and
planting decisions as local climates continue to change.

Although the framework presented here removes subjectivity in determining the transition between
chilling and forcing temperatures, exploring alternative approaches to chill calculation could be
beneficial to further improve model outputs. The Dynamic Model has been shown to be one of the
best models for comparing chilling across regions (Fernandez *et al.*, 2020), and particularly
without acknowledging cold hardiness (North & Kovaleski, 2024). NYUS.1 uses chill portions to
calculate deacclimation potential as an increasing logistic response that translates into the negative
exponential relationship of budbreak time (or forcing) to chilling accumulation (Kovaleski, 2022).
Here, small shifts in temperature for chilling accumulation cause large increases in error (**Fig. 4b**).
Chilling accumulation is a crucial component of most phenology models, and in NYUS.1, chilling
accumulation modulates the magnitude of acclimation and deacclimation responses throughout the

season. Given the significance of chilling to the model variability and the expected future declines in chill accumulation in regions like southern France (**Fig. 4c-e**) (Fernandez *et al.*, 2023), improving existing chill models or developing new ones is necessary to better reflect physiological responses to cold temperatures (Gaeta *et al.*, 2018; North & Kovaleski, 2024). Alternatively, reparametrizing chill models such that they reflect species or genotype-specific behaviors in chill accumulation could also potentially improve the performance of this and other phenology models (North *et al.*, 2024). Our work thus adds to a growing body of literature showing that cold hardiness is likely an important factor to be acknowledged in studies of chilling and dormancy, which is further substantiated as even at the molecular level cold hardiness genes have been implicated in mechanisms associated with dormancy [i.e., *CBF* genes directly promoting expression of *DAM/SVL/SVP2*, Ding et al. (2024)].

Although there was no parameter optimization for the cold hardiness model, predictions of budbreak resulted in very good statistics – particularly after correcting for cold damage (RMSE=7.2 days, Bias=0.6 days, r^2=0.88; **Fig. 3**). The NYUS.1 model was developed using only grapevine bud cold hardiness data obtained in Geneva, NY, US (Kovaleski *et al.*, 2023), while parameters which relate cold hardiness dynamics to chilling accumulation were obtained from a study of 15 unrelated woody perennial species grown in Boston, MA, US [**Fig. 2** in Kovaleski (2022)]. This demonstrates that process-based models that use physiologically-derived parameters can be used across climates – even if chill accumulation has a small effect on residuals (**Supporting Information Fig. S5d**). Measurements of cold hardiness, particularly in warmer climates, or using cold hardiness-informed chill models could be helpful in removing chill accumulation as a source of error. Our metrics are comparable to or better than studies using different models that included calibration based on budbreak data. Ferguson *et al.* (2014), found a general RMSE of 7.3 days (n=152) for budbreak predictions of 23 grapevine cultivars in Washington state, USA using the WAUS.2 cold hardiness model. However, their predictions had a low coefficient of determination (r^2=0.45), likely due to the little variability in dates of budbreak for a single region. Considering the poor performance for the WAUS.2 model to predict cold hardiness in different locations (Kovaleski *et al.*, 2023; Wang *et al.*, 2024), we can speculate that budbreak predictions would also be poor. García de Cortázar-Atauri *et al*. (2009) found RMSE ranging from 9.2 to 18.3 days for three models (one with two parameter sets) for Cabernet-

Sauvignon (n=63) and Riesling (n=64) in four regions in France. Sgubin *et al*. (2018) found
RMSEs between 9.7 and 11.1 days for Cabernet-Sauvignon (n=71) and Riesling (n=65) using two
different phenological models in four regions in France. Moreover, we performed a comparison of
model performance using PhenoFlex, considered to be one of the strongest established phenology
modeling approaches within the chilling-forcing framework [Luedeling *et al.* (2021); see
Supporting Information Figs. S8, S9, and **Notes S3** for a full explanation]. PhenoFlex was
parameterized and tested in two different approaches, all of which returned RMSEs that were
comparable to those produced by NYUS.1 (**Notes S3**). The RMSE is assessed by comparing model
outputs with observed budbreak dates. Standardized methods for recording phenology
observations, such as EL and BBCH scales, are widely used. However, some degree of subjectivity
is unavoidable when the phenotyping is based on observations of characteristics that cannot be
easily discretely quantified. In multi-decadal phenology records spanning multiple locations with
different observers, inconsistencies in observation methods are inevitable (Liu *et al.*, 2021). A
further limitation is the lack of information on observation frequency, which may constrain the
achievable accuracy of model-observation comparisons. Here we demonstrated that, depending on
previously known discrepancies in observational habits within phenology records, adjusting the
threshold of the NYUS.1 model can improve predictions for budbreak (**Supporting Information**
Fig. S4). As has been previously described in other organisms (van Strien *et al.*, 2008), our results
reinforce that observations using 50% occurrence of different phenological stages have better
results than using first observations – even with corrections. With expansion of phenology
modeling, the development of standardized, quantifiable methods for observing budbreak,
including the use of imaging [e.g., PhenoCam (Katal *et al.*, 2022)], would enhance model accuracy
across locations, observers, and species – advances that will likely contribute to a deeper
understanding of dormant season physiology.

Previous modeling work on climate suitability for grapevines only included a mid-winter threshold
(Morales-Castilla *et al.*, 2020). However, dynamics of weather are extremely important in
midwinter cold hardiness, and therefore some of the potential viticultural expansion into colder
climates predicted may be overestimated. Our predictions included a significant amount of years
and locations with some level of cold damage (129 out of the 329 total observations). The
equations used to estimate damage and correct subsequent cold hardiness trajectories are based on

known delays in time to budbreak that occurs as a result of low temperature damage (Evans *et al.*,
2019; Chamberlain & Wolkovich, 2021; Muffler *et al.*, 2024b). We demonstrated that correcting for
predicted cold damage (when temperature drops below cold hardiness levels) can improve
predictions of budbreak timing. Importantly, we validated our predictions of damage using
historical records in the location with the most extensive temperature record (Portland, NY, US;
Figs. 3, S3). Besides extensive temperature records for Portland, NY, qualitative records of
damage (newspaper records and extension reports) were also longstanding, publicly available, and
easily accessible. We also assumed that newspaper records would consistently report damage
events, given the economic importance of grapevines in the region. Nonetheless, smaller instances
of damage, those occurring pre-budbreak may have gone unreported. This may have resulted in an
underrepresentation of damage events, in turn, resulting in a higher false positive rate. Because
these historical reports are incomplete and heterogeneous in detail, this component of the
validation should be regarded as preliminary, but our analysis indicates these reports may be a
possible source of data for cold damage.
820
Though this analysis does suggest that NYUS.1 can accurately predict damage, further subjectivity
exists in the historical records, with variable language used to describe the damage and the degree
to which damage was observable. For example, NYUS.1 predicted some instances of only slight
damage when records of damage were not found. A likely explanation is that the damage was light
and localized, and thus not widely reported, resulting in the more conservative nature of the model
based on higher false positive predictions (**Supporting Information Fig. S3**). To further improve
these predictions, experiments may be required, including defining minimum cold hardiness
thresholds for different tissues (Proebsting *et al.*, 1978; Gardea, 1987), and measuring mid-winter
cold hardiness in multiple locations – particularly warmer climates where this trait is often easily
ignored. It is important to note, however, that cold damage is known to delay spring phenology
(Chamberlain & Wolkovich, 2021; Qiu *et al.*, 2024; Muffler *et al.*, 2024a), and ordinary budbreak
models have trained on phenology datasets that contain varying levels of cold damage. Thus, shifts
in damage patterns that are expected depending on increasing temperatures [e.g., **Fig. 5** in our
study, and Lamichhane (2021)] may be a cause of failure for such models as climate changes,
while that is unlikely to affect a model using cold hardiness as its basis.
836

Here, we present a novel method for predicting spring phenology, producing accurate budbreak
predictions using a previously published cold hardiness model (NYUS.1 Kovaleski *et al.*, 2023),
without calibrating its parameters to the budbreak dataset used in this study. Such phenological
predictions can be used for planning of agricultural practices related to different phenological
stages, such as sprays for pest control and pruning. Furthermore, we also fit new NYUS.1
parameters for different damage percentiles, allowing predicted cold damage (as a result of
temperatures dropping below modeled cold hardiness) to shift budbreak predictions and thereby
improve their accuracy.

The good predictions without calibration indicate that the physiologically derived parameters used
within the NYUS.1 framework work well in non-analogous climates, despite having originated
from data for a single location. Through instances where reacclimation occurs, we use a
physiological basis to introduce the equivalent of negative forcing into phenological modeling.
Beyond budbreak, our framework for predictions includes cold hardiness, allowing for corrections
of predictions based on damage incurred, as well as assessments of cultivar suitability to different
locations and climates, both presently and in future conditions based on winter physiology.
Improvements of winter physiological responses are still possible, such as better defining chilling
accumulation effects. Therefore, empirical determination of dormancy and cold hardiness in
different climatic gradients are warranted. Continued work in measuring cold hardiness of many
species and in varied climates [e.g., Charrier et al. (2013); Deslauriers et al. (2021); Kovaleski
(2022)] for empirical validation of cold hardiness predictions will allow for expansion of cold
hardiness and budbreak modeling using a framework such as the one presented here, particularly
in warmer, marginal climates for temperate woody perennials. Future work may then integrate
cold hardiness dynamics into large-scale predictive models to improve forecasts of plant responses
to climate change and niche modeling. These efforts can ultimately enable more reliable
predictions in earth system models, and better understanding of climatic feedbacks in shifting
spring phenology of woody perennials.

Acknowledgements

We thank M. Chen for comments on the work. This work is partially supported by the National

Institute of Food and Agriculture, United States Department of Agriculture, McIntire Stennis

projects 7002619 and 1027327, and by the Office of the Vice Chancellor for Research at the

University of Wisconsin-Madison with funding from the Wisconsin Alumni Research

Foundation.

Competing interests

The authors have no competing interests to declare.

Data availability

The data and code for analyses and figures are available through GitHub

(https://github.com/apkovaleski/VitisPhenology)

Figure Legends

Figure 1. Locations and cultivars in the phenology dataset and cold hardiness model

development. The phenology dataset used here contains eight locations: four in France (Bergheim,

Bordeaux, Marseillan-Plage, and Montreuil-Bellay), one in Germany (Veitshöchheim), two in the

United States (Lewisburg, PA, and Portland, NY), and one in Canada (Niagara-on-the-Lake, ON).

The NYUS.1 cold hardiness model was developed using grapevine bud cold hardiness data from

Geneva, NY (Kovaleski et al., 2023), and used dormancy parameters from 15 unrelated woody

perennial species grown in Boston, MA – both in the United States.

Figure 2. Conceptual overview of the cold hardiness model to predict budbreak. The NYUS.1

cold hardiness model (Kovaleski *et al.*, 2023) is used to predict the time to budbreak for *V. labrusca*

'Concord' grapes grown in Portland, NY, US in the 1983-1984 dormant season. Cold hardiness

(blue ribbon) is gained and lost in response to seasonal fluctuations in air temperature (grey

ribbon). Two different end trajectories are shown for the cold hardiness prediction: real cold

hardiness values that remain negative in value, where CH stops decreasing once it reaches minima

extracted from [70] (blue ribbon); and a second, projected cold hardiness where values are calculated

into non-freezing temperatures (red ribbon and dashed line). When the projected cold hardiness

reaches a threshold determined for cold hardiness at budbreak [established from data in Kovaleski

et al. (2018)], budbreak is predicted to occur.

Figure 3. Comparison of budbreak predictions from cold hardiness model with observations.

(a) Relationship between initial predictions and observations of budbreak. **(b)** Based on expected

cold damage prior to budbreak, predictions of budbreak can be corrected, shifting to a later date as

the cold hardiness of the surviving bud population is greater. **(c)** Distribution of residuals based on

five different groupings, from left to right: initial predictions (years uncorrected for damage +

years where damage was not predicted n = 329); years where no damage was predicted (n = 200),

uncorrected predictions in years with predicted damage (n = 129), predictions corrected to account

for damage (n = 129), and full, corrected predictions (predictions corrected to account for damage

+ predictions in years where damage is not predicted, n = 329). Residuals for each pair were

compared using a one-way ANOVA. **(d)** Comparison of predictions to observations after

accounting for the consequences of freeze damage. (**e**) Sample article from historical records used
to qualitatively validate damage predictions (Erie Daily Times, 5 May 1977). In (**a,b**), thin solid
black line represents 1:1 ratio between observations and predictions, and dotted lines represent an
interval of +/- 7 days from 1:1 line. Heavy black line shows linear relationship between predicted
and observed points.

Figure 4. Phenology predictions in different sites are differentially affected by shifts in how
chilling accumulation is calculated. (**a**) Density plot of recorded temperatures in Portland, NY,
US and Marseillan-Plage, FR, and the relative contribution of temperatures to chilling
accumulation in the Dynamic Model under unaltered and two sample phase-shift scenarios (-2ºC
and -6ºC). (**b**) RMSE of budbreak phenology predictions under a range from -6ºC to +6ºC of phase
shift scenarios. **c,d,e,** Phenology predictions compared to observations and average chilling
accumulation in a cold location (Portland, NY) and warm location (Marseillan-Plage, FR) under
(**c**) unshifted conditions, (**d**) a -2ºC phase shift, and (**e**) a -6ºC phase shift scenario. In (**c,d,e**), thin
solid black line represents 1:1 ratio between observations and predictions, and dotted lines
represent an interval of +/- 7 days from 1:1 line. Heavy black line shows linear relationship
between predicted and observed points. Horizontal dashed line at DOY 150 in (c,d,e) is used to
highlight change in axis in (e), and points shown as an arrowhead at 212 days are failed predictions.

Figure 5. Freeze risk shows temporal, temperature, and cultivar effects. (**a**) Damage risk
density plot over time within a season, based on cold hardiness predictions for three grapevine
cultivars in Portland, NY, US over the time period of 1927-2023 shows two instances of higher
probability of damage: midwinter (120-80 days before budbreak) and around budbreak (15 days
before and after budbreak). Red curve represents predicted instances of freeze damage (safety
margin less than or equal to 0), and the yellow curve represents instances where there is a narrow
safety margin (between 0ºC and 2ºC). Horizontal axis shows "0" as the mean day of budbreak for
each cultivar over the time period. (**b, c**) Number of predicted freeze damage risk days based on
mean dormant season temperature (1 November–30 April), for (**b**) each of three grapevine
cultivars grown in eight locations and (**c**) for all cultivars combined. Point sizes (**b, c**) are
proportional within each decimal mean dormant season temperature. Risk of damage was
separated in four temperature bands based on mean dormant season temperature: [1; blue] high

freeze risk injury, particularly for *V. vinifera* cultivars, due to low mid-winter temperatures; [2;
green] cool spring days delay budbreak until last frost; [3; orange] risk of freeze damage increases
as spring phenology advances faster than date of last frost; [4; red] temperatures are too warm for
low temperature injury to be significant. Horizontal colored lines (**c**) show mean dormant season
temperature ranges in datasets from each of the eight locations studied.
946
Figure 6. Sensitivity of phenology to temperature is uneven across the range of climates
studied. (a, b) Freeze damage corrected predicted day of budbreak for all cultivars as a response
to mean dormant season temperatures (1 November–30April) for (**a**) all eight locations combined
and (**b**) separately for Portland, NY, US and Marseillan-Plage, FR. (**a**) Budbreak advances at an
overall rate of 5.8 days/ºC for all cultivars, while showing different magnitudes and directions at
different temperature intervals, (**b**) which is also affected by local temperature dynamics.
953
954
955

References

Arora R, Rowland LJ. **2011**. Physiological Research on Winter-hardiness: Deacclimation Resistance, Reacclimation Ability, Photoprotection Strategies, and a Cold Acclimation Protocol Design. *HortScience horts* **46**: 1070–1078.

Ashcroft GL, Richardson EA, Seeley SD. **1977**. A Statistical Method of Determining Chill Unit and Growing Degree Hour Requirements for Deciduous Fruit Trees. *HortScience* **12**: 347–348.

Augspurger CK. **2013**. Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing. *Ecology* **94**: 41–50.

Baranger A, Cordonnier T, Charrier G, Delzon S, Larter M, Martin-StPaul NK, Kunstler G. **2024**. Living on the edge – physiological tolerance to frost and drought explains range limits of 35 European tree species. *Ecography* **n/a**: e07528.

Benmoussa H, Ghrab M, Ben Mimoun M, Luedeling E. **2017**. Chilling and heat requirements for local and foreign almond (Prunus dulcis Mill.) cultivars in a warm Mediterranean location based on 30 years of phenology records. *Agricultural and Forest Meteorology* **239**: 34–46.

Bock A, Sparks T, Estrella N, Menzel A. **2011**. Changes in the phenology and composition of wine from Franconia, Germany. *Climate Research* **50**: 69–81.

Bujang MA, Adnan TH. **2016**. Requirements for minimum sample size for sensitivity and specificity analysis. *Journal of clinical and diagnostic research: JCDR* **10**: YE01.

Chamberlain CJ, Cook BI, García de Cortázar-Atauri I, Wolkovich EM. **2019**. Rethinking false spring risk. *Global Change Biology* **25**: 2209–2220.

Chamberlain CJ, Wolkovich EM. **2021**. Late spring freezes coupled with warming winters alter temperate tree phenology and growth. *New Phytologist* **231**: 987–995.

Charrier G, Martin-StPaul N, Damesin C, Delpierre N, Hänninen H, Torres-Ruiz JM, Davi H. **2021**. Interaction of drought and frost in tree ecophysiology: rethinking the timing of risks. *Annals of Forest Science* **78**: 40.

Charrier G, Poirier M, Bonhomme M, Lacointe A, Améglio T. **2013**. Frost hardiness in walnut trees (Juglans regia L.): How to link physiology and modelling? *Tree Physiology* **33**: 1229–1241.

Chmura DJ, Anderson PD, Howe GT, Harrington CA, Halofsky JE, Peterson DL, Shaw DC, Brad St.Clair J. **2011**. Forest responses to climate change in the northwestern United States: Ecophysiological foundations for adaptive management. *Forest Ecology and Management* **261**: 1121–1142.

Chuine I. **2000**. A Unified Model for Budburst of Trees. *Journal of Theoretical Biology* **207**: 337–347.

Chuine I, Régnière J. **2017**. Process-Based Models of Phenology for Plants and Animals. *Annual Review of Ecology, Evolution, and Systematics* **48**: 159–182.

Cleland EE, Allen JM, Crimmins TM, Dunne JA, Pau S, Travers SE, Zavaleta ES, Wolkovich EM. **2012**. Phenological tracking enables positive species responses to climate change. *Ecology* **93**: 1765–1771.

Coombe BG. **1995**. Growth Stages of the Grapevine: Adoption of a system for identifying grapevine growth stages. *Australian Journal of Grape and Wine Research* **1**: 104–110.

Cruz-Alonso V, Pucher C, Ratcliffe S, Ruiz-Benito P, Astigarraga J, Neumann M, Hasenauer H, Rodríguez-Sánchez F. **2023**. The easyclimate R package: Easy access to high-resolution daily climate data for Europe. *Environmental Modelling & Software* **161**: 105627.

Darbyshire R, Webb L, Goodwin I, Barlow EWR. **2013**. Evaluation of recent trends in Australian pome fruit spring phenology. *International Journal of Biometeorology* **57**: 409–421.

Deslauriers A, Garcia L, Charrier G, Buttò V, Pichette A, Paré M. **2021**. Cold acclimation and deacclimation in wild blueberry: Direct and indirect influence of environmental factors and non-structural carbohydrates. *Agricultural and Forest Meteorology* **301–302**: 108349.

Ding J, Wang K, Pandey S, Perales M, Allona I, Khan MRI, Busov VB, Bhalerao RP. **2024**. Molecular advances in bud dormancy in trees. *Journal of Experimental Botany* **75**: 6063–6075.

Du Y, Pan Y, Ma K. **2019**. Moderate chilling requirement controls budburst for subtropical species in China. *Agricultural and Forest Meteorology* **278**: 107693.

Duchêne E. **2019**. Vitis vinifera cv. Riesling developmental stages.

Egea J, Ortega E, Martínez-Gómez P, Dicenta F. **2003**. Chilling and heat requirements of almond cultivars for flowering. *Environmental and Experimental Botany* **50**: 79–85.

Evans KJ, Bricher PK, Foster SD. **2019**. Impact of frost injury incidence at nodes of Pinot Noir on fruitfulness and growth-stage lag. *Australian Journal of Grape and Wine Research* **25**: 201–211.

Ferguson JC, Moyer MM, Mills LJ, Hoogenboom G, Keller M. **2014**. Modeling Dormant Bud Cold Hardiness and Budbreak in Twenty-Three Vitis Genotypes Reveals Variation by Region of Origin. *American Journal of Enology and Viticulture* **65**: 59–71.

Ferguson JC, Tarara JM, Mills LJ, Grove GG, Keller M. **2011**. Dynamic thermal time model of cold hardiness for dormant grapevine buds. *Annals of botany* **107**: 389–396.

Fernandez E, Krefting P, Kunz A, Do H, Fadón E, Luedeling E. **2021**. Boosting statistical delineation of chill and heat periods in temperate fruit trees through multi-environment observations. *Agricultural and Forest Meteorology* **310**: 108652.

Fernandez E, Mojahid H, Fadón E, Rodrigo J, Ruiz D, Egea JA, Ben Mimoun M, Kodad O, El Yaacoubi A, Ghrab M, *et al.* 2023. Climate change impacts on winter chill in Mediterranean temperate fruit orchards. *Regional Environmental Change* **23**: 1–18.

Fernandez E, Whitney C, Luedeling E. **2020**. The importance of chill model selection — a multi-site analysis. *European Journal of Agronomy* **119**: 126103.

Fick SE, Hijmans RJ. **2017**. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**: 4302–4315.

Fishman S, Erez A, Couvillon GA. **1987a**. The temperature dependence of dormancy breaking in plants: Computer simulation of processes studied under controlled temperatures. *Journal of Theoretical Biology* **126**: 309–321.

Fishman S, Erez A, Couvillon GA. **1987b**. The temperature dependence of dormancy breaking in plants: mathematical analysis of a two-step model involving a cooperative transition. *Journal of Theoretical Biology* **124**: 473–483.

Flynn DFB, Wolkovich EM. **2018**. Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytol* **219**: 1353–1362.

Ford KR, Harrington CA, Bansal S, Gould PJ, St. Clair JB. **2016**. Will changes in phenology track climate change? A study of growth initiation timing in coast Douglas-fir. *Global Change Biology* **22**: 3712–3723.

Gaeta L, Stellacci AM, Losciale P. **2018**. Evaluation of three modelling approaches for almond blooming in Mediterranean climate conditions. *European Journal of Agronomy* **97**: 1–10.

Gallinat AS, Ellwood ER, Heberling JM, Miller-Rushing AJ, Pearse WD, Primack RB. **2021**. Macrophenology: insights into the broad-scale patterns, drivers, and consequences of phenology. *American Journal of Botany* **108**: 2112–2126.

García de Cortázar-Atauri I, Brisson N, Gaudillere JP. **2009**. Performance of several models
for predicting budburst date of grapevine (Vitis vinifera L.). *International Journal of*
Biometeorology **53**: 317–326.
Gardea AA. **1987**. Freeze damage of Pinot noir (Vitis vinifera L.) as affected by bud
development, INA-bacteria, and a bacterial inhibitor.
Goeckeritz CZ, Gottschalk C, van Nocker S, Hollender CA. **2023**. Malus Species with
Diverse Bloom Times Exhibit Variable Rates of Floral Development. *Journal of the American*
Society for Horticultural Science **148**: 64–73.
Harrell E, Frank M. **2024**. Package 'rms'. Regression Modeling Strategies.
Hassan T, Gulzar R, Hamid M, Ahmad R, Waza SA, Khuroo AA. **2024**. Plant phenology
shifts under climate warming: a systematic review of recent scientific literature. *Environmental*
Monitoring and Assessment **196**: 36.
Hébert-Haché A, Inglis D, Kemp B, Willwerth JJ. **2021**. Clone and Rootstock Interactions
Influence the Cold Hardiness of Vitis vinifera cvs. Riesling and Sauvignon blanc. *American*
Journal of Enology and Viticulture **72**: 126.
Hillmann L, Elsysy M, Goeckeritz C, Hollender C, Rothwell N, Blanke M, Einhorn T.
2021. Preanthesis changes in freeze resistance, relative water content, and ovary growth preempt
bud phenology and signify dormancy release of sour cherry floral buds. *Planta* **254**: 74.
Hufkens K, Basler D, Milliman T, Melaas EK, Richardson AD. **2018**. An integrated
phenology modelling framework in r. *Methods in Ecology and Evolution* **9**: 1276–1285.
Inouye DW. **2022**. Climate change and phenology. *WIREs Climate Change* **13**: e764.
Kalberer SR, Wisniewski M, Arora R. **2006**. Deacclimation and reacclimation of cold-hardy
plants: Current understanding and emerging concepts. *Plant Science* **171**: 3–16.
Katal N, Rzanny M, Mäder P, Wäldchen J. **2022**. Deep Learning in Plant Phenological
Research: A Systematic Literature Review. *Frontiers in Plant Science* **13**.
Kirchhof E, Campos-Arguedas F, Arias NS, Kovaleski AP. **2025**. Thresholds for spring
freeze: measuring risk to improve predictions in a warming world. *New Phytologist* **n/a**.
Körner C. **2021**. The cold range limit of trees. *Trends in Ecology & Evolution* **36**: 979–989.
Körner C, Basler D, Hoch G, Kollas C, Lenz A, Randin CF, Vitasse Y, Zimmermann NE.
2016. Where, why and how? Explaining the low-temperature range limits of temperate tree
species. *Journal of Ecology* **104**: 1076–1088.
Kovaleski AP. **2022**. Woody species do not differ in dormancy progression: Differences in time
to budbreak due to forcing and cold hardiness. *Proceedings of the National Academy of Sciences*
119: e2112250119.
Kovaleski AP, North MG, Martinson TE, Londo JP. **2023**. Development of a new cold
hardiness prediction model for grapevine using phased integration of acclimation and
deacclimation responses. *Agricultural and Forest Meteorology* **331**: 109324.
Kovaleski AP, Reisch BI, Londo JP. **2018**. Deacclimation kinetics as a quantitative phenotype
for delineating the dormancy transition and thermal efficiency for budbreak in Vitis species. *AoB*
Plants **10**: ply066.
Kramer K, Leinonen I, Loustau D. **2000**. The importance of phenology for the evaluation of
impact of climate change on growth of boreal, temperate and Mediterranean forests ecosystems:
an overview. *International Journal of Biometeorology* **44**: 67–75.
Lamichhane JR. **2021**. Rising risks of late-spring frosts in a changing climate. *Nature Climate*
Change **11**: 554–555.

Lang GA, Early JD, Martin GC, Darnell RL. **1987**. Endo-, para-, and ecodormancy:
physiological terminology and classification for dormancy research. *HortScience* **22**: 371–377.
Lenz A, Hoch G, Vitasse Y, Körner C. **2013**. European deciduous trees exhibit similar safety
margins against damage by spring freeze events along elevational gradients. *New Phytologist*
200: 1166–1175.
Li H-L, Ali A, Luo X, Liao K, Sun N, Xu M-S, Sha L-B, He D, Du Y-J, Sun W-W, *et al.*
2024. China's subtropical deciduous plants are more sensitive to climate change than evergreen
plants by flowering phenology. *Global Change Biology* **30**: e17168.
Li D, Stucky BJ, Deck J, Baiser B, Guralnick RP. **2019**. The effect of urbanization on plant
phenology depends on regional temperature. *Nature Ecology & Evolution* **3**: 1661–1667.
Liu G, Chuine I, Denéchère R, Jean F, Dufrêne E, Vincent G, Berveiller D, Delpierre N.
2021. Higher sample sizes and observer inter-calibration are needed for reliable scoring of leaf
phenology in trees. *Journal of Ecology* **109**: 2461–2474.
Londo JP, Kovaleski AP. **2025**. Integrating cold hardiness and deacclimation resistance
demonstrates a conserved response to chilling accumulation in grapevines. *Journal of*
Experimental Botany: eraf045.
Luedeling E, Caspersen L, Fernandez E. **2023**. chillR: Statistical methods for phenology
analysis in temperate fruit trees, R package version 0.75.
Luedeling E, Schiffers K, Fohrmann T, Urbach C. **2021**. PhenoFlex - an integrated model to
predict spring phenology in temperate fruit trees. *Agricultural and Forest Meteorology* **307**:
108491.
Malagi G, Sachet MR, Citadin I, Herter FG, Bonhomme M, Regnard J-L, Legave JM.
2015. The comparison of dormancy dynamics in apple trees grown under temperate and mild
winter climates imposes a renewal of classical approaches. *Trees* **29**: 1365–1380.
Marquis B, Lajoie G. **2024**. Experimental exposure to winter thaws reveals tipping point in
yellow birch bud mortality and phenology in the northern temperate forest of Québec, Canada.
Climate Change Ecology **7**: 100087.
Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C, Berger S, Caud N, Chen Y,
Goldfarb L, Gomis MI. **2021**. Climate Change 2021: The Physical Science Basis. Contribution
of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change. *IPCC: Geneva, Switzerland*: 65.
Maury O, Quidoz M-C, Garcia de Cortazar Atauri I, Chuine I, El Hasnaoui M, Tromel L.
2023. Portail de données phénologiques du réseau TEMPO / TEMPO Data Portal (INRAE, Ed.).
Meier U, Bleiholder H, Buhr L, Feller C, Hack H, Heß M, Lancashire PD, Schnock U,
Stauß R, Van Den Boom T. **2009**. The BBCH system to coding the phenological growth stages
of plants–history and publications. *Journal für Kulturpflanzen* **61**: 41–52.
Meier M, Fuhrer J, Holzkämper A. **2018**. Changing risk of spring frost damage in grapevines
due to climate change? A case study in the Swiss Rhone Valley. *International Journal of*
Biometeorology **62**: 991–1002.
Mo Y, Chen S, Wu Z, Tang J, Fu Y. **2024**. The Advancement in Spring Vegetation Phenology
in the Northern Hemisphere Will Reverse After 2060 Under Future Moderate Warming
Scenarios. *Earth's Future* **12**: e2023EF003788.
Morales-Castilla I, García de Cortázar-Atauri I, Cook BI, Lacombe T, Parker A, van
Leeuwen C, Nicholas KA, Wolkovich EM. **2020**. Diversity buffers winegrowing regions from
climate change losses. *Proceedings of the National Academy of Sciences* **117**: 2864–2869.

Muffler L, Weigel R, Beil I, Leuschner C, Schmeddes J, Kreyling J. **2024a**. Winter and
spring frost events delay leaf-out, hamper growth and increase mortality in European beech
seedlings, with weaker effects of subsequent frosts. *Ecology and Evolution* **14**: e70028.
Muffler L, Weigel R, Beil I, Leuschner C, Schmeddes J, Kreyling J. **2024b**. Winter and
spring frost events delay leaf-out, hamper growth and increase mortality in European beech
seedlings, with weaker effects of subsequent frosts. *Ecology and Evolution* **14**: e70028.
Mura C, Buttò V, Silvestro R, Deslauriers A, Charrier G, Raymond P, Rossi S. **2022**. The
early bud gets the cold: Diverging spring phenology drives exposure to late frost in a Picea
mariana [(Mill.) BSP] common garden. *Physiologia Plantarum* **174**: e13798.
North MG, Kovaleski AP. **2024**. Time to budbreak is not enough: cold hardiness evaluation is
necessary in dormancy and spring phenology studies. *Annals of botany*.
North M, Workmaster BA, Atucha A. **2021**. Cold hardiness of cold climate interspecific
hybrid grapevines grown in a cold climate region. *American Journal of Enology and Viticulture*
72: 318–327.
North M, Workmaster BA, Atucha A. **2022**. Effects of chill unit accumulation and
temperature on woody plant deacclimation kinetics. *Physiologia Plantarum*: e13717.
North MG, Workmaster BA, Atucha A, Kovaleski AP. **2024**. Cold hardiness-informed
budbreak reveals role of freezing temperatures and daily fluctuation in a chill accumulation
model. *Journal of Experimental Botany* **75**: 6182–6193.
Persico MJ, Smith DE, Centinari M. **2021**. Delaying Budbreak to Reduce Freeze Damage:
Seasonal Vine Performance and Wine Composition in Two Vitis vinifera; Cultivars. *American*
Journal of Enology and Viticulture **72**: 346.
Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X. **2019**. Plant
phenology and global climate change: Current progresses and challenges. *Global Change*
Biology **25**: 1922–1940.
Pick JL, Nakagawa S, Noble DWA. **2019**. Reproducible, flexible and high-throughput data
extraction from primary literature: The metaDigitise r package. *Methods in Ecology and*
Evolution **10**: 426–431.
Proebsting EL, Brummund VP, Clore WJ. **1978**. Critical temperatures for Concord grapes.
Extension Manual 4330. Washington State University, Pullman.
Qian H, Zhang Y, Ricklefs RE, Wang X. **2022**. Relationship of minimum winter temperature
and temperature seasonality to the northern range limit and species richness of trees in North
America. *Journal of Geographical Sciences* **32**: 280–290.
Qiu H, Yan Q, Yang Y, Huang X, Wang J, Luo J, Peng L, Bai G, Zhang L, Zhang R, *et al.*
2024. Flowering in the Northern Hemisphere is delayed by frost after leaf-out. *Nature*
Communications **15**: 9123.
Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M. **2013**.
Climate change, phenology, and phenological control of vegetation feedbacks to the climate
system. *Agricultural and Forest Meteorology* **169**: 156–173.
De Rosa V, Vizzotto G, Falchi R. **2021**. Cold Hardiness Dynamics and Spring Phenology:
Climate-Driven Changes and New Molecular Insights Into Grapevine Adaptive Potential.
Frontiers in Plant Science **12**.
Sgubin G, Swingedouw D, Dayon G, García de Cortázar-Atauri I, Ollat N, Pagé C, van
Leeuwen C. **2018**. The risk of tardive frost damage in French vineyards in a changing climate.
Agricultural and Forest Meteorology **250–251**: 226–242.

Song Z, Song X, Pan Y, Dai K, Shou J, Chen Q, Huang J, Tang X, Huang Z, Du Y. **2020**.
Effects of winter chilling and photoperiod on leaf-out and flowering in a subtropical evergreen
broadleaved forest in China. *Forest Ecology and Management* **458**: 117766.
van Strien AJ, Plantenga WF, Soldaat LL, van Swaay CAM, WallisDeVries MF. **2008**. Bias
in phenology assessments based on first appearance data of butterflies. *Oecologia* **156**: 227–235.
Stuke M, Yun K, Kim S-H. **2024**. Adapting a process-oriented cold hardiness model to
conifers. *Forest Ecology and Management* **553**: 121611.
Thornton PE, Shrestha R, Thornton M, Kao S-C, Wei Y, Wilson BE. **2021**. Gridded daily
weather data for North America with comprehensive uncertainty quantification. *Scientific Data*
8: 190.
Vitasse Y, Schneider L, Rixen C, Christen D, Rebetez M. **2018**. Increase in the risk of
exposure of forest and fruit trees to spring frosts at higher elevations in Switzerland over the last
four decades. *Agricultural and Forest Meteorology* **248**: 60–69.
Vitra A, Lenz A, Vitasse Y. **2017**. Frost hardening and dehardening potential in temperate trees
from winter to budburst. *New Phytologist* **216**: 113–123.
Wang J, Hua H, Guo J, Huang X, Zhang X, Yang Y, Wang D, Guo X, Zhang R, Smith NG,
et al. **2025**. Late spring frost delays tree spring phenology by reducing photosynthetic
productivity. *Nature Climate Change* **15**: 201–209.
Wang H, Moghe GD, Kovaleski AP, Keller M, Martinson TE, Wright AH, Franklin JL,
Hébert-Haché A, Provost C, Reinke M, *et al.* **2024**. NYUS.2: an automated machine learning
prediction model for the large-scale real-time simulation of grapevine freezing tolerance in North
America. *Horticulture Research* **11**: uhad286.
Wang H, Wu C, Ciais P, Peñuelas J, Dai J, Fu Y, Ge Q. **2020**. Overestimation of the effect of
climatic warming on spring phenology due to misrepresentation of chilling. *Nature*
Communications **11**: 4945.
Wheeler KI, Dietze MC, LeBauer D, Peters JA, Richardson AD, Ross AA, Thomas RQ,
Zhu K, Bhat U, Munch S, *et al.* **2024**. Predicting spring phenology in deciduous broadleaf
forests: NEON phenology forecasting community challenge. *Agricultural and Forest*
Meteorology **345**: 109810.
Wolkovich EM, García de Cortázar-Atauri I, Morales-Castilla I, Nicholas KA, Lacombe T.
2018. From Pinot to Xinomavro in the world's future wine-growing regions. *Nature Climate*
Change **8**: 29–37.
Zeng X, Du Y, Vitasse Y. **2023**. Untangling winter chilling and spring forcing effects on spring
phenology of subtropical tree seedlings. *Agricultural and Forest Meteorology* **335**: 109456.

1216

1217

1218   **Supporting Information**
1219
1220
Table S1. Overview of phenological records used in this study.
Table S2. Parameters used for the NYUS.1 cold hardiness model.
Figure S1. Conceptual framework of grapevine bud cold hardiness dynamics, deacclimation, and
phenological thresholds.
Figure S2. Workflow illustration of analysis of cold hardiness and phenology predictions.
Figure S3. Confusion matrix for phenology predictions.
Figure S4. Effects of observational habits on phenology predictions.
Figure S5. Partial residual plots for possible sources of variability in model outputs.
Figure S6. Residuals of budbreak predictions by location.
Figure S7. Maximum cold hardiness achieved in a season decreases with warmer temperatures.
Figure S8. Predicted budbreak using PhenoFlex, trained on a single location.
Figure S9. Predicted budbreak using PhenoFlex, trained on the entire dataset.
Notes S1. The NYUS.1 model used to estimate cold hardiness for each cultivar.
Notes S2. Adjustment of cold hardiness predictions to account for damage.
Notes S3. Comparison of NYUS.1 model outputs with PhenoFlex.
1236

1237

1238
Figure 1. Locations and cultivars in the phenology dataset and cold hardiness model development.
The phenology dataset used here contains eight locations: four in France (Bergheim, Bordeaux, Marseillan-
Plage, and Montreuil-Bellay), one in Germany (Veitshöchheim), two in the United States (Lewisburg, PA,
and Portland, NY), and one in Canada (Niagara-on-the-Lake, ON). The NYUS.1 cold hardiness model was
developed using grapevine bud cold hardiness data from Geneva, NY (Kovaleski et al., 2023), and used
dormancy parameters from 15 unrelated woody perennial species grown in Boston, MA – both in the United
States.
1246

1247

1248
Figure 2. Conceptual overview of the cold hardiness model to predict budbreak. The NYUS.1 cold
hardiness model (Kovaleski *et al.*, 2023) is used to predict the time to budbreak for *V. labrusca* 'Concord'
grapes grown in Portland, NY, US in the 1983-1984 dormant season. Cold hardiness (blue ribbon) is gained
and lost in response to seasonal fluctuations in air temperature (grey ribbon). Two different end trajectories
are shown for the cold hardiness prediction: (1) observed cold hardiness values, which remain negative and
stop decreasing once they reached the minimum cold hardiness reported by Gardea (1987) (blue ribbon);
and (2) projected cold hardiness values, where the model continues to predict loss of cold hardiness into
positive temperature values. Although these do not reflect real values of cold hardiness, they are used for
budbreak predictions (red ribbon and dashed line). When the projected cold hardiness reaches a threshold
determined for cold hardiness at budbreak [established from data in Kovaleski et al. (2018)], budbreak is
predicted to occur.
1260

1261
1262
Figure 3. Comparison of budbreak predictions from cold hardiness model with observations. (a)
Relationship between initial predictions and observations of budbreak. (b) Based on expected cold damage
prior to budbreak, predictions of budbreak can be corrected, shifting to a later date as the cold hardiness of
the surviving bud population is greater. (c) Distribution of residuals based on five different groupings, from
left to right: initial predictions (years uncorrected for damage + years where damage was not predicted n =
329); years where no damage was predicted (n = 200), uncorrected predictions in years with predicted
damage (n = 129), predictions corrected to account for damage (n = 129), and full, corrected predictions
(predictions corrected to account for damage + predictions in years where damage is not predicted, n =

329). (d) Comparison of predictions to observations after accounting for the consequences of freeze
damage. (e) Sample article from historical records used to qualitatively validate damage predictions (Erie
Daily Times, 5 May 1977). In (a,b), thin solid black line represents 1:1 ratio between observations and
predictions, and dotted lines represent an interval of +/- 7 days from 1:1 line. Heavy black line shows linear
relationship between predicted and observed points.

1276
Figure 4. Phenology predictions in different sites are differentially affected by shifts in how chilling
accumulation is calculated. (a) Density plot of recorded temperatures in Portland, NY, US and Marseillan-
Plage, FR, and the relative contribution of temperatures to chilling accumulation in the Dynamic Model
under unaltered and two sample phase-shift scenarios (-2ºC and -6ºC). (b) RMSE of budbreak phenology
predictions under a range from -6ºC to +6ºC of phase shift scenarios. (c,d,e) Phenology predictions
compared to observations and average chilling accumulation in a cold location (Portland, NY) and warm
location (Marseillan-Plage, FR) under (c) unshifted conditions, (d) a -2ºC phase shift, and (e) a -6ºC phase
shift scenario. In (c,d,e), thin solid black line represents 1:1 ratio between observations and predictions, and
dotted lines represent an interval of +/- 7 days from 1:1 line. Heavy black line shows linear relationship
between predicted and observed points. Horizontal dashed line at DOY 150 in (c,d,e) is used to highlight
change in axis in (e), and points shown as an arrowhead at 212 days are failed predictions.

Figure 5. Freeze risk shows temporal, temperature, and cultivar effects. (a) Damage risk density plot over time within a season, based on cold hardiness predictions for three grapevine cultivars in Portland, NY, US over the time period of 1927-2023 shows two instances of higher probability of damage: midwinter (120-80 days before budbreak) and around budbreak (15 days before and after budbreak). Red curve represents predicted instances of freeze damage (safety margin less than or equal to 0), and the yellow curve represents instances where there is a narrow safety margin (between 0°C and 2°C). Horizontal axis shows "0" as the mean day of budbreak for each cultivar over the time period. (b,c) Number of predicted freeze damage risk days based on mean dormant season temperature (1 November–30 April), for (b) each of three grapevine cultivars grown in eight locations and (c) for all cultivars combined. Point sizes (b,c) are proportional within each decimal mean dormant season temperature. Risk of damage was separated in four temperature bands based on mean dormant season temperature: [1; blue] high freeze risk injury, particularly for *V. vinifera* cultivars, due to low mid-winter temperatures; [2; green] cool spring days delay budbreak until last frost; [3; orange] risk of freeze damage increases as spring phenology advances faster than date of last frost; [4; red] temperatures are too warm for low temperature injury to be significant. Horizontal colored lines (**c**) show mean dormant season temperature ranges in datasets from each of the eight locations studied.

1308
Figure 6. Sensitivity of phenology to temperature is uneven across the range of climates studied. (a,b)
Freeze damage corrected predicted day of budbreak for all cultivars as a response to mean dormant season
temperatures (1 November–30April) for (a) all eight locations combined and (b) separately for Portland,
NY, US and Marseillan-Plage, FR. (a) Budbreak advances at an overall rate of 5.8 days/ºC for all cultivars,
while showing different magnitudes and directions at different temperature intervals, (b) which is also
affected by local temperature dynamics.