Title
A major update of the genome assembly of Indian Eri silkmoth, *Samia ricini*

Authors
Jung Lee[1], Mana Okamoto[1], Rin Kawagoe[1], Toru Shimada[1]

Affiliations
1   Gakushuin University, Faculty of Science, Department of Life Science, Mejiro 1-5-1, Toshima-ku, Tokyo, 171-8588, Japan

*Jung Lee is the corresponding author.

Abstract

Indian eri silkmoth, *Samia ricini*, is a wild silkmoth whose silk occupies a significant economic position. In addition to its importance as an economic animal, *S. ricini* is also useful as a model species of Saturniidae. National BioResource of Japan (NBRP) maintains a *S. ricini* strain brought to Japan during WWII via Taiwan. Since we have previously published a draft genome assembly of *S. ricini*, we have attempted to construct a chromosome-level genome assembly to facilitate genetic studies of *S. ricini*. We successfully constructed a chromosome-scale genome assembly by exploiting two long-read-based technologies, HiFi reads and optical genome mapping.
Furthermore, we performed functional annotations of the genome assembly, i.e., repeat annotation, transcriptome-based gene prediction, ATAC-seq, and PIWI-interacting RNA (piRNA)-targeted small RNA-seq. The assembly harbours 16,226 protein-coding genes and 636 piRNA clusters across three tissues: ovaries, testis, and embryos. ATAC-seq data comprehensively detected open chromosome regions, which will benefit when CRISPR/Cas9-mediated genome editing is conducted.

Background & Summary

In India, many silkworm species other than *Bombyx mori* (mulberry silkworm) are reared for silk production. As far as we know, such species belong to the family Saturniidae. *Samia ricini*, eri silkworm, represents one example. Similar to *B. mori*, *S. ricini* is completely domesticated species and have no wild populations. Of the saturniid silkworms reared in India, *S. ricini* is the biggest cocoon producer[1]. In addition to that, there is a culture of eating the pupae of *S. ricini*, in the northern regions of India[2]. Since FAO (Food and Agriculture Organization of the United Nations) reported that insect diets are one of the practical solutions to the food problems[3], there is growing interest in entomophagy. Therefore, *S. ricini* is expected to be in increasing demand not only as a sericultural insect but also as a food insect.

S. ricini also has another side to it as a model organism for the Saturniidae family. *S. ricini* is exceptionally multivoltine species in the family Saturniidae and can be reared throughout the year. Moreover, *S. ricini* is the only insect that genome editing has already been announced as being possible so far[4]. In view of this situation, the development of highly accurate whole-genome annotation information on *S. ricini* would be of great benefit to entomology.

We have published a draft genome sequence of *S. ricini* in 2020[5], where 14 chromosomes of *S. ricini* were split into 155 contigs. We first attempted to improve the assembly accuracy without adding extra sequence data. We re-assembled the *S. ricini* genome using the continuous long read data and short read data from Lee et al., 2020[5] (DRS137258[6] and DRS137260[7]). The following scaffolding was conducted with optical genome mapping data (Table 1)[8]. In the resulting assembly (Sr_NBRP_v2.0; GCA_014132275.2)[9], the number of sequences and chromosomes were equal, and we were able to construct a

'chromosome level' assembly for the time being. However, the BUSCO completeness was
low (Table 2), suggesting that there were many undetermined regions. In addition, the total
gap size was approximately 4.5 Mbp, which seemed to be the performance limit of Noisy
Long reads. Therefore, we obtained approximately 30 Gbase (coverage x60) of data using
HiFi reads and performed assembly and scaffolding. As a result, we have succeeded in
constructing a chromosome-scale assembly containing only 400 bp gaps (Table 1;
Sr_NBRP_v3.0; GCA_XXXXXX).

Transcriptome-based gene prediction and functional annotation were also
performed, identifying 16,226 protein-coding genes in this assembly. For the future
application of the CRISPR/Cas9, we performed embryonic ATAC-seq to associate the
positions of protein-coding genes and open chromatin regions since the enzymatic activity of
Cas9 is less efficient in heterochromatin regions[10]. In addition to ATAC-seq, we performed
piRNA-targeted small RNA-seq to identify piRNA clusters (piCs) since piRNA is involved in the
early development of Lepidoptera[11].

Methods

Insects
S. ricini (White strain) was provided from NBRP silkworm (https://shigen.nig.ac.jp/silkwormbase/). The
S. ricini larvae were fed on fresh castor leaves under a long-day condition (16 h light / 8 h dark) at
25°C.

Genome assembly of Sr_NBRP_v2.0 (GCA_014132275.2)[9]
PacBio continuous long read (CLR) data (DRS137260)[7] were subjected to Flye (v 2.9)[12] with default
setting. After the draft assembly step, raw CLR data was reused for polishing with Racon[13]. In this case,
consensus calling by Racon was converged by one attempt. The resulting polished draft sequences
were subjected to Pilon (v 1.22)[14] for further polishing. For polishing with Pilon, 2 paired-end libraries
(DRR213145 and DRR244931)[15] and 9 mate-pair libraries (DRR213147, DRR213148, DRR213149,
DRR213150, DRR213151, DRR213152, DRR213153, DRR213154, and DRR213155)[15] were used. The
polished assembly was scaffolded with optical genome mapping as previously described[16]: Genomic
DNA was isolated from the pupa immediately after the pupation for the optical genome mapping.
DNA isolation was conducted using the Bionano Prep Animal Tissue DNA Isolation Fibrous Tissue
Protocol (Bionano Genomics). We used two enzymes, namely, DLE-1 and Nt.BsqQI for labelling. The
labelling procedure was conducted according to the Bionano Prep Direct Label and Stain Protocol. The
labelled samples were scanned on the Bionano Saphyr system using Saphyr Chip G2.3. The obtained
data were analyzed using Bionano Access (v 1.8.2)[17] and Bionano Solve (v 3.8.2)[18]. In the two-enzyme
pipeline, the process of creating the .cmap files was the same as in the one-enzyme pipeline. The
resulting two .cmap files are submitted to runTGH.R (also bundled with Bionano Solve) with default
settings. The .cmap files were deposited in DDBJ. The basic statistics of this assembly are summarised
in Table 1.

Genome assembly of Sr_NBRP_v3.0 (AP038896–AP038909)[19–32]
Genomic DNA was extracted from whole body of a male final instar larvae of *S. ricini* using
NucleoBond HMW DNA (MACHEREY-NAGEL). A SMRTbell library was prepared with SMRTbell library
preparation 3.0. HiFi reads were obtained on the Revio platform. The HiFi reads were assembled with
Hifiasm (v 0.16.1-r375)[33]. The resulting draft assembly was scaffolded with the above-mentioned
optical genome mapping data. To close the gaps, which were bridged by optical genome mapping
reads, we attempted to use error-corrected CLR (DRS137260)[7] :the raw CLRs were subjected into the
"correct" module in PECAT (v 0.0.3)[33] with default settings. The gaps in the scaffolded assembly were
patched to the error-corrected CLRs using "GapFiller" module in QuarTeT (v 1.2.0)[34]. To close
remaining gaps, we patched the previous assembly (Sr_NBRP_v2.0; GCA_014132275.2)[9] using
"AssemblyMapper" module in QuarTeT (v 1.2.0)[34]. Finally, we polished the resulting assembly with
HiFi reads with NextPolish2 (v 0.2.1)[35] with default settings. Winnowmap (v 2.03)[36] was used with
default parameters to align HiFi reads to the temporary genome assembly after "AssemblyMapper."
Yak (v 0.1-r69)[33] was used with default parameters to create k-mer hash tables from short read data
(DRR213150)[15] which were mandatory inputs for NextPolish2. Schematic diagram of assembly process

was summarised in Fig. 1. The assembly gaps were visualised in Fig. 2a. All sequence reads data used
in this study were summarised in Table 2.
110
Repetitive elements annotation in the genomes
Repetitive annotation of the *S. ricini* genome was briefly summarised here: repetitive elements in the
genome assembly were identified using RepeatModeler (v 2.0.4)[37] with the "-LTRstruct" option for
performing an LTR structural search. The annotated elements were masked using RepeatMasker (v
4.1.2)[38] with '-lib' option to specify "consensi.fa.classified" file generated by RepeatModeler. The
amount of annotated repetitive elements in the assembly was summarised in Fig. 2b. We realised that
absence of SINE in the genome can characterise the repetitive elements of *S. ricini* genome (Fig. 2b).
Repeat Landscape graph using the divergence data was also generated (Fig. 2c). Among the annotated
repetitive elements, LTR, non-LTR (LINE or SINE), DNA transposons, and rolling circles were extracted
and the density information of those repetitive groups is visualized by circlize (v 0.4.16)[39] (Fig. 2d).
121
RNA sample preparation for RNA-seq, small RNA-seq, and Iso-seq
All RNA samples were prepared precisely as previously described[16]. Total RNA was extracted using
TRIzol reagent (Invitrogen) according to the manufacturer's protocol. Embryos were sampled 48 hours
after oviposition. Testis and ovary-derived RNA samples were subjected to RNA-seq and small RNA-
seq (sRNA-seq), respectively. Embryonic RNA samples were subjected to sRNA-seq and Iso-seq
respectively.
128
Library preparation for RNA-seq, sRNA-seq and Iso-seq
The sRNA-seq library was prepared using TruSeq small RNA kit (illumina) according to the
manufacturer's protocol with a slight modification. To target piRNA, a region of 147-158 nucleotides
was extracted in the purification step of the cDNA construct using BluePippin (Sage Science). The
constructed library was sequenced on the illumina HiSeq 2500 platform (illumina). Except for an RNA
sample from embryos, the RNA-seq libraries were prepared using TruSeq stranded mRNA kit (illumina)
according to the manufacturer's protocol. The embryonic RNA-seq library was prepared using
NEBNext Poly(A) mRNA Magnetic Isolation Module (New England BioLabs) and the NEBNext® Ultra™ II
Directional RNA Library Prep Kit (New England BioLabs) according to the manufacturer's protocol. The
constructed RNA-seq libraries were sequenced on the illumina NovaSeq6000 platform (illumina). For
Iso-seq, the library was constructed using Sequel Iso-seq Express Template Prep (Pacific Bioscience)
according to the manufacturer's protocol. The constructed library was sequenced on the PacBio
Sequel platform (PacBio).
142
Transcriptome-based gene prediction
BRAKER3 (v 3.0.8) was used for gene prediction[40,41]. The RNA-seq and Iso-seq data were submitted to
BRAKER3 separately[42], and Tsebra finally merged their respective prediction[43]. The detailed
information on transcriptome data is summarised in Supplementary Table 2. Quality trimming for
short read data was conducted using fastp (v 0.20.1)[44] with following options: '-q 28 -l 80. ' Trimmed
short read data were submitted to BRAKER3 using the '--rnaseq_sets_ids' option. The short reads
were aligned to the genome assembly by hisat2 (v 2.2.1)[45]. Iso-seq data were generated consensus for
each read cluster according to the following procedure[46]: Iso-seq subreads were converted to circular
consensus sequences (ccs) using ccs v 6.4.0 with options '--minLength 10 --maxLength 100000 --
minPasses 0 --minSnr 2.5 --minPredictedAccuracy 0.0.' lima (v 2.7.1) was used to remove primer
sequences from the CCSs with options '--isoseq --peek-guess --ignore-biosamples.' After the trimming
of adaptors, PolyA tail trimming and concatemer removal were performed by isoseq3 (v 3.8.2) in
'refine' mode with option '--require-polya.' Finally, isoform-level clustering was conducted by isoseq3
in 'cluster' mode with option '--use-qvs.' The resulting clustered.bam file was submitted to BRAKER3.
The basic metrics of gene models were summarised in Table 3. The resulting gene models were also
submitted to BUSCO analysis[30,] scoring 98.4% completeness (Table 4). BUSCO analysis on the genome
assembly was also conducted to compare completeness scores of both gene models and genome
assembly[30] (Table 4).
161
Functional annotation of gene models
The deduced amino acid sequences of gene models were submitted to EnTAP[6] for functional
annotation. A protein similarity search was conducted against the latest complete UniProtKB/TrEMBL

protein data set and complete UniProtKB/Swiss-Prot data set using diamond (v 0.9.14)[47]. A protein
orthology search was also conducted against the EggNOG databases30 to assign Gene Ontology (GO),
KEGG terms, and protein domains from pfam[48] and smart[49]. Additional family and domain search was
performed against TIGRFAM[50], SFLD[51], HAMAP[52], CDD[53], SUPERFAMILY[54], PRINTS[55], PANTHER[56], and
Gene3d[57] using InterProScan (v 5.70-102)[58]. The results of functional annotation are summarized in
Table 5. The top 10 GOs assigned to the gene models are shown in Fig. 3a without distinguishing
between molecular function, biological process, and cellular component. The top 10 GOs for each
category are shown in Fig. 3b–d.
173
ATAC library preparation and data processing
Another batch of early embryo samples was subjected to Iso-seq, and sRNA-seq was subjected to
ATAC-seq. Fragmentation and amplification of the ATAC-seq libraries were conducted according to
Buenrostro et al. (2015)[59]. The constructed libraries were sequenced on the Illumina HiSeq. ATAC-seq
reads were pretreated with fastp and mapped to the genome with bwa-mem2 (v 2.2.1)[60]. Alignments
containing mismatches were then removed using bamutils (v 0.5.9)[61]. Next, we removed duplicated
reads using GATK MarkDuplicates (v 4.1.7)[62]. The resulting bam files were converted to bigwig files
using deepTools bamCoverage (v 3.5.1)[63]. Heatmap was created using deepTools computeMatrix, and
the starting point of the gene model was set to the reference point (Fig. 4).
183
184 **Small RNA mapping**
185 The small RNA reads were trimmed using Trim Galore (v 0.6.6)[64] in small RNA mode. The trimmed
186 small RNA reads were mapped to the assembled transcriptome, allowing up to 3 nucleotide
187 mismatches using Hisat2 (v 2.1.0)[45] and ngsutils (v 0.5.9)[61]. The information for each library is
188 summarised in Table 2.
189
piRNA cluster detection
The piC detection was performed as previously described[16,65]. proTRAC (v 2.4.4)[65] was used with
options '-clsize 5000 -pimin 23 -pimax 29 -1Tor10A 0.3 -1Tand10A 0.3 -clstrand 0.0 -clsplit 1.0 -distr
1.0-99.0 -spike 90-1000 -nomotif -pdens 0.05.' As a result, we successfully identified 560 piRNA
clusters in the three tissues (Fig. 5). The identity of piC is defined by the two nearest gene models. If
multiple piCs were predicted between such two genes, such piCs were treated as a single piC. The
genomic positions of piCs identified in testes, ovaries, and early embryos were visualized by
RIdeogram (v 0.2.2)[66] (Fig. 5a). The aggregation relationship of those piCs was visualized by
ComplexUpset (v 1.3.3)[67] (Fig. 5b).
199

Data Records

The raw sequence data reported in this paper have been deposited in DDBJ. Genomic data for the
draft assembly were deposited under the accession code DRP012461[68]. Genome assemblies referred
in this study were available at accession codes GCA_014132275.2[9] and AP038896-AP038909[19–32].
Except for embryonic Iso-seq data, all transcriptome data were also registered under the accession
code DRP012461[68]. Embryonic Iso-seq data, small RNA-seq data and ATAC-seq data are available
under the accession code DRP009926 [DRR396182, DRR396184, DRR396185, and
DRR396186][69]. Annotated gene models and piRNA distribution maps have been deposited to the
figshare repository[70].
209

Technical Validation

BUSCO (v 5.4.6)[71] with lepidoptera_odb10 lineage dataset was used to assess the quality of gene
models. For comparison, the results are summarised in Table 4, together with the results of BUSCO
analysis for the genome assembly. 98.4% of the complete and single-copy BUSCO sequences were
present in the gene models, while 98.6% of the complete and single-copy BUSCO sequences were in
the genome assembly. BUSCO completeness scores were almost the same between the genome
assembly and the gene model, suggesting that the gene prediction process is highly accurate across all
genome regions.
218

Code Availability

Programs exploited in this study were executed with the default parameters except where otherwise specified in the Methods section. No custom code was used during this study.

Acknowledgments

Insects were donated from Kyushu University and Shinshu University according to a Grant-in Aid "National BioResource Project (NBRP, RR2002), Silkworm Genetic Resources" for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan. This study was funded by Ellie Inc .This study was also supported by JSPS KAKENHI Grant Numbers 20K15535 and 24K17900 to J.L and JSPS KAKENHI Grant Number J18H03949 to T.S.

Author contributions

J.L. designed the research plan, performed RNA extraction, analyzed the obtained data, and wrote the manuscript. T.S. also designed this research plan and performed the data analysis. M.O. and R.K. reared the larvae of *S. ricini* and prepared RNA samples. T.A.

Competing interests

The authors declare no competing interests.

Figure Legends

Fig. 1 Genome assembly pipeline used in this study

The numbers 1 to 5 in the figure identify one of the five files used as input (1, HiFi reads; 2, Optical genome mapping data; 3, continuous long reads; 4, previous version of genome assembly; 5, short reads).

Fig. 2 A chromosome-scale genome assembly of *S. ricini*

(a) Gap distribution on the genome assembly of *S. ricini*.

There is one gap each on both chromosome 2 and 7. There are two gaps in close proximity in the terminal region of chromosome 11.

(b) Repeat contents of the genome assembly. Amount (Mbp; left) and fractions (right) by repeat category were shown.

(c) Repeat landscape graph based on divergence data of repetitive elements.

(d) Summary of *S. ricini* genome characteristics. The outermost to the innermost circle are I. chromosome ideograms; II. GC content; III. GC skew; IV. LTR element density; V. non-LTR retrotransposon density; VI. DNA transposon density; VII. rolling circle density; and VIII. gene model density.

Fig. 3 The top 10 GO assignments of gene models

(a) Overall top 10 GO assignments to gene models

(b) Top 10 "biological process" GO assignments to gene models

(c) Top 10 "cellular component" GO assignments to gene models

(d) Top 10 "molecular function" GO assignments to gene models

Fig. 4 Heatmap around gene bodies of ATAC-seq on early embryos.

Fig. 5 piRNA clusters on *S. ricini* genome

(a) piCs distribution detected in early embryos (box), pupal ovary (circle), and pupal testis (triangle)

(b) UpSet plot visualising piCs that are each assigned to each tissue. The vertical bars correspond to the intersections. When the circles corresponding to tissues are connected by a line, the bar above circles represents the number of piCs commonly identified in concerning tissues. For example, the yellow bar indicates the number of piCs identified in all tissues. The identity of piCs was defined by the nearest two gene models: When comparing piCs identified in different tissues, if the nearest upstream and downstream gene models are the same, those piCs were treated as the same piC.

1.   Central   Silk   Board.   *SERICULTURAL   STATISTICS   IN   INDIA.*
https://texmin.nic.in/sites/default/files/Sericulture%20Statistics.pdf (2023).
2.   Shrivastava, S. K. & Prakash, A. *INSECTS FOR FOOD: AN UNEXPLORED BIORESOURCE*
FOR MAKE IN INDIA. J. Appl. Zool. Res vol. 27 https://cabidigitallibrary.org (2016).
3.   Huis, A. *et al. EDIBLE INSECTS Future Prospects Fo Food and Feed Security.* vol. 171
(2013).
4.   Lee, J., Kiuchi, T., Kawamoto, M., Shimada, T. & Katsuma, S. Accumulation of uric acid
in the epidermis forms the white integument of Samia ricini larvae. *PLoS One* **13**,
e0205758 (2018).
5.   Lee, J. *et al.* The genome sequence of Samia ricini , a new model species of
lepidopteran insect. *Mol Ecol Resour* **21**, 327–339 (2021).
6.   Lee, J. NCBI Sequence Read Archive. *http://identifiers.org/ncbi/insdc.sra:DRS137258*
http://identifiers.org/ncbi/insdc.sra:DRS137258 (2020).
7.   Lee, J. NCBI Sequence Read Archive. *https://identifiers.org/ncbi/insdc.sra:DRS137260*
https://identifiers.org/ncbi/insdc.sra:DRS137260 (2020).
8.   Lee, J. *et al.* The genome sequence of Samia ricini, a new model species of
lepidopteran insect. *Mol Ecol Resour* **21**, 327–339 (2020).
9.   Lee,   J.   &   Shimada,   T.   NCBI   GenBank.
http://identifiers.org/ncbi/insdc.gca:GCA_014132275.2 (2023).
10.   Jain, S. *et al.* TALEN outperforms Cas9 in editing heterochromatin target sites. *Nat*
Commun **12**, 4–13 (2021).
11.   Kiuchi, T. *et al.* A single female-specific piRNA is the primary determiner of sex in the
silkworm. *Nature* **509**, 633–636 (2014).
12.   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads
using repeat graphs. *Nat Biotechnol* **37**, 540–546 (2019).
13.   Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res* **27**, 737–746 (2017).
14.   Walker, B. J. *et al.* Pilon: An integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, (2014).
15.   Lee, J. NCBI Sequence Read Archive. *https://identifiers.org/ncbi/insdc.sra:DRS137258*
(2020).
16.   Lee, J. *et al.* W chromosome sequences of two bombycid moths provide an insight
into the origin of Fem. *Mol Ecol* **33**, 1–12 (2024).
17.   Bionano   Genomics,   I.   Bionano   Access   Software.   Preprint   at
https://bionano.com/access-software/.
18.   Bionano   Genomics,   I.   Bionano   Solve   software.   Preprint   at
https://bionano.com/software-downloads/.
19.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038896*
(2025).
20.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038897*
(2025).
21.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038898*
(2025).
22.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038899*
(2025).
23.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038900*
(2025).
24.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038901*
(2025).
25.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038902*
(2025).

26.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038903*
(2025).

27.   Lee, J. & Shimada, T. NCBI nucleotide. *https://identifiers.org/ncbi/insdc:AP038904*
(2025).

28.   Lee, J. & Shimada, T. NCBI Nucleotide. *https://identifiers.org/ncbi/insdc:AP038905*
(2025).

29.   Lee, J. & Shimada, T. NCBI Nucleotide. *https://identifiers.org/ncbi/insdc:AP038906*
(2025).

30.   Lee, J. & Shimada, T. NCBI Nucleotide. *https://identifiers.org/ncbi/insdc:AP038907*
(2025).

31.   Lee, J. & Shimada, T. NCBI Nucleotide. *https://identifiers.org/ncbi/insdc:AP038908*
(2025).

32.   Lee, J. & Shimada, T. NCBI Nucleotide. *https://identifiers.org/ncbi/insdc:AP038909*
(2025).

33.   Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170–175
(2021).

34.   Lin, Y. *et al.* QuarTeT: A telomere-To-Telomere toolkit for gap-free genome assembly
and centromeric repeat identification. *Hortic Res* **10**, (2023).

35.   Hu, J. *et al.* NextPolish2: A Repeat-aware Polishing Tool for Genomes Assembled
Using HiFi Long Reads. *Genomics Proteomics Bioinformatics* **22**, (2024).

36.   Jain, C., Rhie, A., Hansen, N. F., Koren, S. & Phillippy, A. M. Long-read mapping to
repetitive reference sequences using Winnowmap2. *Nat Methods* **19**, 705–710
(2022).

37.   Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable
element families. *Proc Natl Acad Sci U S A* **117**, 9451–9457 (2020).

38.   Smit, A., Hubley, R. & Green, P. RepeatMasker Open-4.0. 2013–2015.

39.   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. Circlize implements and enhances
circular visualization in R. *Bioinformatics* **30**, 2811–2812 (2014).

40.   Stanke, M., Schöffmann, O., Morgenstern, B. & Waack, S. Gene prediction in
eukaryotes with a generalized hidden Markov model that uses hints from external
sources. *BMC Bioinformatics* **7**, 62 (2006).

41.   Stanke, M., Diekhans, M., Baertsch, R. & Haussler, D. Using native and syntenically
mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**, 637–
644 (2008).

42.   Lomsadze, A., Burns, P. D. & Borodovsky, M. Integration of mapped RNA-Seq reads
into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res* **42**, 1–8
(2014).

43.   Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M. & Stanke, M. TSEBRA: transcript
selector for BRAKER. *BMC Bioinformatics* **22**, 1–12 (2021).

44.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

45.   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods* **12**, 357–360 (2015).

46.   Brůna, T., Gabriel, L. & Hoff, K. J. Navigating Eukaryotic Genome Annotation Pipelines:
A Route Map to BRAKER, Galba, and TSEBRA. (2024).

47.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nat Methods* **12**, 59–60 (2014).

48.   Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**,
D412–D419 (2021).

49.   Letunic, I. & Bork, P. 20 years of the SMART protein domain annotation resource.
Nucleic Acids Res **46**, D493–D496 (2018).

50.   Haft, D. H. *et al.* TIGRFAMs: A protein family resource for the functional identification
of proteins. *Nucleic Acids Res* **29**, 41–43 (2001).

51.   Akiva, E. *et al.* The Structure-Function Linkage Database. *Nucleic Acids Res* **42**, 521–
530 (2014).

52.   Pedruzzi, I. *et al.* HAMAP in 2015: Updates to the protein family classification and
annotation system. *Nucleic Acids Res* **43**, D1064–D1070 (2015).

53.   Wang, J. *et al.* The conserved domain database in 2023. *Nucleic Acids Res* **51**, D384–
D388 (2023).

54.   Pandurangan, A. P., Stahlhacke, J., Oates, M. E., Smithers, B. & Gough, J. The
SUPERFAMILY 2.0 database: A significant proteome update and a new webserver.
Nucleic Acids Res **47**, D490–D494 (2019).

55.   Attwood, T. K. *et al.* The PRINTS database: A fine-grained protein sequence
annotation and analysis resource-its status in 2012. *Database* **2012**, 1–9 (2012).

56.   Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P. D. PANTHER version 14:
More genomes, a new PANTHER GO-slim and improvements in enrichment analysis
tools. *Nucleic Acids Res* **47**, D419–D426 (2019).

57.   Lewis, T. E. *et al.* Gene3D: Extensive prediction of globular domains in proteins.
Nucleic Acids Res **46**, D435–D439 (2018).

58.   Jones, P. *et al.* InterProScan 5: Genome-scale protein function classification.
Bioinformatics **30**, 1236–1240 (2014).

59.   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for
Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, (2015).

60.   Vasimuddin, Md., Misra, S., Li, H. & Aluru, S. Efficient Architecture-Aware
Acceleration of BWA-MEM for Multicore Systems. in *2019 IEEE International Parallel*
and Distributed Processing Symposium (IPDPS) 314–324 (IEEE, 2019).
doi:10.1109/IPDPS.2019.00041.

61.   Breese, M. R. & Liu, Y. NGSUtils: A software suite for analyzing and manipulating next-
generation sequencing datasets. *Bioinformatics* **29**, 494–496 (2013).

62.   van der Auwera, G. & O'Connor, B. D. *Genomics in the Cloud: Using Docker, GATK, and*
WDL in Terra. (O'Reilly Media, Incorporated, 2020).

63.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160–W165 (2016).

64.   Krueger, F. Trim Galore. https://github.com/FelixKrueger/TrimGalore (2020).

65.   Rosenkranz, D. & Zischler, H. proTRAC - a software for probabilistic piRNA cluster
detection, visualization and analysis. *BMC Bioinformatics* **13**, 5 (2012).

66.   Hao, Z. *et al.* RIdeogram: Drawing SVG graphics to visualize and map genome-wide
data on the idiograms. *PeerJ Comput Sci* **6**, 1–11 (2020).

67.   Krassowski Michał. ComplexUpset.

68.   Lee, J. NCBI Sequence Read Archive. *https://identifiers.org/ncbi/insdc.sra:DRP012461*
(2025).

69.   Lee, J. NCBI Sequence Read Archive. *https://identifiers.org/ncbi/insdc.sra:DRP009926*
https://identifiers.org/ncbi/insdc.sra:DRP009926 (2023).

70.   Lee, J. The genome annotation files of the latest genome assembly of Indian eri
silkmoth (Samia ricini) including gene models, its functional annotation result, and
piRNA cluster maps. *FigShare* https://doi.org/10.6084/m9.figshare.27268212 (2025)
doi:10.6084/m9.figshare.27268212.

71.   Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol* **38**,
4647–4654 (2021).

I chromosome ideogram
II GC content
III GC skew
IV LTR density
V non-LTR density
VI DNA transposon
VII rolling circle
VIII gene density

a

GO Term	
GO:0003674	
GO:0008150	
GO:0005575	
GO:0009987	
GO:0008152	
GO:0005488	
GO:0003824	
GO:0071704	
GO:0044464	
GO:0005623	

count

b

biological process

GO Term	
GO:0008150	
GO:0009987	
GO:0008152	
GO:0071704	
GO:0044238	
GO:0044237	
GO:0043170	
GO:0065007	
GO:0050789	
GO:0032501	

Count

c

cellular component

GO Term	
GO:0005575	
GO:0044464	
GO:0005623	
GO:0005622	
GO:0044424	
GO:0043226	
GO:0043229	
GO:0043227	
GO:0043231	
GO:0016020	

Count

d

molecular function

GO Term	
GO:0003674	
GO:0005488	
GO:0003824	
GO:0097159	
GO:1901363	
GO:0043167	
GO:0003676	
GO:0043169	
GO:0016787	
GO:0046872	

Count

a

Embryo piCs
Ovary piCs
Testis piCs

01 02 03 04 05 06 07 08 09 10 11 12 13 14

b

Intersection size

68
49
34
28

Ovary
Testis
Embryo

Set size

500 400 300 200 100 0

piRNA clusters

Assembly	Accession	Strain	Level	No. of sequences	Total bases	N50 [bp]	Gap [bp]	BUSCO completeness [%]	Source
Sr_NBRP_v1.0	GCA_014132275.1	White	Contig	155	450,479,495	21,366,385	0	97.6	Lee et al., 2020
Sr_NBRP_v2.0	GCA_014132275.2	White	Chromosome	**14**	452,483,983	33,135,288	4,566,028	96.4	This Study
Sr_NBRP_v3.0	AP038896–AP038909	White	Chromosome	**14**	453,650,863	**33,371,992**	**400**	**98.8**	This Study

Table 1 Basic statistics of the genome assemblies of *S. ricini*

ID	Run accession	biosample accession	Purpose	No. of spots	Length	Read pairs after trimming	tissue	age	Source
1	DRR596219	SAMD00819509	RNA-seq	49,533,284	101PE	23,747,007	anterior and middle silk gland	5th instar 3rd day larvae	This study
2	DRR596220	SAMD00819510	RNA-seq	62,957,650	101PE	30,706,997	epidermis	3rd instar 0th day larvae	This study
3	DRR596221	SAMD00819511	RNA-seq	57,939,494	101PE	28,209,875	epidermis	4th instar 1st day larvae	This study
4	DRR596222	SAMD00819512	RNA-seq	45,644,604	101PE	22,195,994	epidermis	5th instar 3rd day larvae	This study
5	DRR596223	SAMD00819513	RNA-seq	47,654,200	101PE	23,060,565	midgut	5th instar 3rd day larvae	This study
6	DRR596224	SAMD00819514	RNA-seq	57,351,070	101PE	27,645,513	posterior silk gland	5th instar 3rd day larvae	This study
7	DRR596225	SAMD00819515	RNA-seq	15,522,314	150PE	15,413,240	hemocytes	5th instar larvae	This study
8	DRR596226	SAMD00819516	RNA-seq	15,351,834	150PE	15,260,640	fat body	5th instar larvae	This study
9	DRR596227	SAMD00819517	RNA-seq	16,354,434	150PE	16,260,987	gonads	5th instar larvae	This study
10	DRR596228	SAMD00819518	RNA-seq	14,152,184	150PE	14,062,249	malpighian tubules	5th instar larvae	This study
11	DRR596229	SAMD00819519	RNA-seq	13,758,095	150PE	13,673,816	midgut	5th instar larvae	This study
12	DRR596230	SAMD00819520	RNA-seq	14,007,357	150PE	13,904,310	adult's 9th ganglion	immediately after eclosion	This study
13	DRR596231	SAMD00819521	RNA-seq	14,743,955	150PE	14,639,299	adult's 9th ganglion	after copulation	This study
14	DRR596232	SAMD00819522	RNA-seq	14,351,152	150PE	14,267,229	nerve chord	5th instar larvae	This study
15	DRR596233	SAMD00819523	RNA-seq	15,453,946	150PE	15,184,662	adult's brain	immediately after eclosion	This study
16	DRR596234	SAMD00819524	RNA-seq	15,236,362	150PE	14,781,588	adult's brain	after copulation	This study
17	DRR596235	SAMD00819525	RNA-seq	14,459,033	150PE	14,243,379	midgut	5th instar larvae	This study
18	DRR596236	SAMD00819526	RNA-seq	14,520,390	150PE	14,379,533	midgut	5th instar larvae	This study
19	DRR396182	SAMD00515715	Iso-seq	9,094,853	-	-	early embryo	48 hours post oviposition	This study
20	DRR396184	SAMD00515715	small RNA-seq	48,219,633	50PE	48,219,633	eary embryo	48 hours post oviposition	This study

21	DRR396185	SAMD00515716	small RNA-seq	25,347,659	50PE	25,347,659	Pupal ovary	7 days post pupation	This study
22	DRR396186	SAMD00515717	small RNA-seq	29,625,406	50PE	29,625,406	Pupal testis	7 days post pupation	This study
23	DRR596218	SAMD00819399	Genomic DNA-seq	1,776,772	-	-	whole body	5th instar larvae	This study
24	DRR213158	SAMD00207788	Genomic DNA-seq	777,113	-	-	silk gland	5th instar larvae	Lee et al., 2020
25	DRR213159	SAMD00207788	Genomic DNA-seq	706,515	-	-	silk gland	5th instar larvae	Lee et al., 2020
26	DRR213156	SAMD00207788	Genomic DNA-seq	925,105	-	-	silk gland	5th instar larvae	Lee et al., 2020
27	DRR213157	SAMD00207788	Genomic DNA-seq	853,458	-	-	silk gland	5th instar larvae	Lee et al., 2020
28	DRR213150	SAMD00207786	Genomic DNA-seq	36,348,660	126PE	35,191,810	silk gland	5th instar larvae	Lee et al., 2020

Table 2 Sequence read information used in this study.

Since sequence runs of IDs 19, 23–27 were performed on long read sequencer, 'Length' and 'Read pairs after trimming' cells are empty.

No. of protein coding gene	16,226
Average CDS length [bp]	1512.4
Average exon length [bp]	228.003
Average intron length [bp]	1822.58

Table 3 Basic statistics of the predicted protein-coding genes

	Complete (Single)	Complete (Duplicated)	Fragmented	Missing
Gene model	5137 (97.2%)	65 (1.2%)	13 (0.2%)	71 (1.4%)
Genome assembly	5204 (98.4%)	13 (0.2%)	14 (0.3%)	55 (1.1%)

Table 4 BUSCO completeness scores of the predicted protein-coding genes and the genome assembly

		Similarity search			Ontology search		total	
		EggNOG	TrEMBL	Swiss-prot	EggNOG**	InterPro		
aligned	informative	12843	12003	7872	10886	12,906	annotated***	14430
	uninformative*	0	2101	182	3046	-		
unaligned		3383	2122	8172	2294	3,320	unannotated***	1796

Table 5 Brief summary of functional annotation

*When the query sequences were aligned to sequences whose description contains any of conserved/predicted/unnamed/hypothetical/putative/unidentified/uncharacterized/unknown/uncultured/uninformative, such alignment was categorized as "uninformative," and the query sequence was treated as an unannotated sequence.

**In this column, queries with at least one GO term were treated as "Informative," while queries without GO terms were treated as "Uninformative." "Unaligned" in this column means queries without protein family assignment.

***"Annotated" means at least one match yielded from any of the databases. "Unannotated" means no match yielded from all databases.