Coupling of cell proliferation to the segmentation clock ensures robust

somite scaling

Marek J. van Oostrom[1], Yuting I. Li[2,3], Wilke H. M. Meijer[1], Tomas E. J. C.

Noordzij[1], Charis Fountas[1,$], Erika Timmers[1,¶], Jeroen Korving[1], Wouter M.

Thomas[1], Benjamin D. Simons[2,4,5] and Katharina F. Sonnen[1,*]

[1]Hubrecht Institute-KNAW (Royal Netherlands Academy of Arts and Sciences),

University Medical Center Utrecht, Utrecht, The Netherlands

[2]Department of Applied Mathematics and Theoretical Physics, Centre for

Mathematical Sciences, University of Cambridge, Wilberforce Road,

Cambridge CB3 0WA, UK

[3]Institute of Science and Technology Austria (ISTA), 3400 Klosterneuburg,

Austria

[4]Gurdon Institute, University of Cambridge, Tennis Court Road, Cambridge,

CB2 1QN, UK

[5]Cambridge Stem Cell Institute, Jeffrey Cheah Biomedical Centre, University of

Cambridge, Cambridge, CB2 0AW, UK

[$]present address: Division of Gene Regulation, the Netherlands Cancer

Institute, Oncode Institute, Plesmanlaan 121, 1066CX Amsterdam, the

Netherlands

[¶]present address: Cell Biology, Neurobiology, and Biophysics, Department of

Biology, Faculty of Science, Utrecht University, Padualaan, 3584 CS Utrecht,

the Netherlands

*Correspondence: k.sonnen@hubrecht.eu

While scaling developmental processes is fundamental to maintaining

robust tissue patterning, the mechanisms underlying this process are

enigmatic. Somitogenesis, the periodic segmentation of growing mesodermal

tissue in vertebrate embryos[1], involves precise scaling with the unsegmented

presomitic mesoderm (PSM) over developmental time and under perturbation[2-

4]. Somitogenesis is spatiotemporally regulated by FGF and Wnt morphogen

gradients and the segmentation clock — oscillations in Notch, Wnt, and FGF signalling[5-7]. Here, we find that cell proliferation is distributed throughout the oscillating PSM. Long-term single-cell tracking in mouse embryo tails uncovered a correlation between cell cycle progression and the segmentation clock, with microfluidics-based entrainment indicating coupling between the cell cycle and signalling oscillations, likely through S-phase inducing Cyclins. A theoretical model suggests this coupling ensures uniform PSM growth, uniform morphogen dilution and precise somite formation, which we validated experimentally by blocking cell proliferation. Our findings reveal that coupling cell proliferation to signalling oscillations is crucial for robust somitogenesis and precise somite scaling.

Main

Somitogenesis is highly reproducible across individual animals of a given species[2]. Cells starting in the posterior tip of the embryo tail traverse the PSM bilaterally along the neural tube until they reach the anterior PSM, where differentiation and somite formation occur[1]. Imbalance between growth and segmentation can lead to premature or delayed arrest in somite formation, improperly sized somites or asymmetries between the left and right[3,4,8], resulting in embryonic lethality or congenital disorders.

Scaling of somitogenesis, the relative adjustment of somite size to PSM size, occurs dynamically during mouse development: PSM size increases until embryonic day (E)9.5 and then decreases until somitogenesis terminates at (E)14.5[2]. At a given developmental time, variations in somite size are very low: at the 36-somite stage (E10.5) forming somites have a size of 162 ± 8 μm^2, which corresponds to a maximum variation of only 1-2 cell diameters. Moreover, experimentally decreasing PSM size results in the formation of smaller somites[3,4]. How such accuracy and precision in segmentation of a variably sized tissue is achieved remains unclear.

The potential involvement of cell proliferation in somitogenesis has been a subject of discussion and controversy. While the cell cycle has been suggested to primarily contribute to PSM growth[9-11], cell cycle perturbations do not interfere with segmentation clock oscillations in 2D *in vitro* culture models

of human somitogenesis[12]. On the other hand, in zebrafish embryos, the

absence of proliferation impairs proper somite formation and has been

described as a source of noise for the segmentation clock[2,8,12-14]. A link between

the segmentation clock and cell division has been suggested[13-15], but not

formally shown.

Here, we use methods including long-term cell tracking, microfluidics-

based entrainment, and modelling to analyse how cell cycle dynamics impact

somite scaling. Our data indicate that proliferation and signalling oscillations

are coupled, and important for robust somite scaling in the context of somitic

development.

Results

Cell cycle dynamics along the tailbud-to-somite axis

To understand the role of cell proliferation in somitogenesis, we first

characterized cell cycle dynamics in uncultured mouse embryo tails.

Immunostaining for the cell cycle marker Ki67 completely overlapped with DAPI

staining, showing that all tailbud, PSM and somite cells were actively cycling

(Fig. 1A). Next, we used flow cytometry to determine the distribution of single

cells from uncultured mouse embryo tails over the four cell cycle phases: G1,

S (when DNA replication occurs), G2 and M phase (when cell division occurs).

We specifically analysed unsegmented mesoderm by gating for Tbx6-positive

cells and found that a majority of cells were in S phase (60.4%), while fewer

were either in G1 (22.7%) or G2/M phase (16.9%, Fig. 1B, Fig. S1). Previously,

we found that mitosis occurs in the entire PSM[10]. To quantify mitosis distribution

more precisely along the anteroposterior (AP) axis from tailbud to somite, we

performed immunostaining using the mitotic cell marker pHH3 (phosphorylated

histone H3) in mouse embryonic tails transversally sectioned from posterior to

anterior (Fig. 1C-E, Fig. S2). We confirmed that cells divided within the tailbud,

PSM and somite region of the tail (Fig. 1E). Furthermore, there was a slight

increase in the percentage of pHH3-positive cells within the first two forming

somite pairs compared to the PSM and tailbud.

Next, we quantified the proliferation rate along the AP axis in embryo

tails using pulse chase labelling by intraperitoneal injection of thymidine

analogues 5-Ethynyl-2′-deoxyuridine (EdU) followed by 5-Bromo-2′-

deoxyuridine (BrdU) 1 h later. This treatment was followed by imaging transversally sectioned tails[16,17] (Fig. 1C, F-H, Fig. S3). The differentially labelled cell fractions allowed us to quantify cell cycle and S-phase duration per section (Fig. S3B, E). Cell cycle duration within the tailbud and PSM was approximately 13.5 hr (Fig. 1H, Fig. S3C) and S phase duration 10 hours (Fig. S3F-H). With a segmentation clock period of approx. 2.7 hours this indicates that it takes about 5 segmentation cycles for tissue size to double. Interestingly, variability in cell cycle duration was much lower in the PSM compared to tailbud and somites (Fig. 1H), hinting at tighter cell cycle control in the PSM.

Thus, cells divide along the whole AP axis at similar rates, suggesting constant expansion of the entire tailbud and PSM region. This contrasts with a model in which cells are added in the posterior PSM, while segmentation occurs in the anterior.

Quantification of cell cycle dynamics using live-cell imaging

To study the dynamics of cell cycle progression at a single-cell rather than population level, we established a long-term single-cell tracking pipeline in the developing mouse embryo tail. To make this possible, we combined light-sheet microscopy with tissue stabilization in agarose grooves and mosaic labelling of cells (Fig. 1I, Fig. S4A-C). We crossed mice carrying the inducible Fucci2 cell cycle G1 reporter R26R-mCherry-hCdt1(30/120)[18] (Fucci2-Cdt1) with mice carrying the inducible nuclear marker R26R-iRFPnucmem[19] (iRFPnucmem) and a CreERT2 expression cassette driven by a T/Brachyury promotor (T-CreERT2)[20]. We then induced mosaic labelling by injecting low concentrations of Tamoxifen into pregnant female mice. Recombination resulted in approximately 10% of cells labelled with mCherry, which tracks the G1 phase of the cell cycle, and nuclear iRFP to allow tracking over time. Single cells were tracked within tissue explants and Fucci2-Cdt1 intensity was quantified over time (Fig. 1J, Movie S1). Cell-cycle duration was 13.6 ±1.3 hr, in agreement with the cell cycle duration *in vivo* (Fig. 1K), and cells spent on average 4.4 hr in G1 phase (Fig. 1L). Notably, we did not detect changes in cell density despite cell divisions over 24 hr of real-time imaging (Fig. S4D, E), indicating steady state tissue growth with cells continuously growing and dividing.

To determine whether cell cycle progression in the PSM is governed by
spatiotemporal regulation, we analysed the spatial organization of cells double-
positive for both the Fucci cell cycle reporters mKO2- hCdt1(30/120) (Fucci1-
Cdt1) and mAG-hGem(1/110) (Fucci1-Gem) in mice expressing both
transgenes in all cells. The presence of both Fucci1-Cdt1 and Fucci1-Gem
signals in cells specifically marks the G1-S transition[21]. Notably, we identified 3
distinct phases for double positive cells in fixed tail explants: an absence of
double positive cells in the PSM (Phase 1), a stripe of double positive cells in
the PSM (Phase 2) and a wider spread of double positive cells throughout the
PSM (Phase 3) (Fig. 1M, Fig. S4F). Such patterns are resonant with fixed
images of cyclic segmentation clock genes showing different expression
patterns in different embryos[22,23]. Our data therefore indicate that there might
be a spatiotemporal organization of cell cycle entry in the PSM.
150
The segmentation clock and cell cycle are correlated
To investigate whether there is a correlation between cell proliferation
and segmentation clock oscillations, we simultaneously quantified cell cycle
progression and segmentation clock oscillations in single cells using a mosaic
labelling approach similar to the above. Specifically, we crossed mice carrying
the inducible nuclear marker R26R-H2B-mCherry[24], the T-CreERT2 transgene
and the bright segmentation clock reporter Achilles-Hes7[25], which is based on
the Notch target gene and transcriptional repressor Hes7 (Fig. 2A-D, Fig. S5).
We tracked single cells using the mosaically expressed mCherry signal
(representing H2B), aligned intensity tracks to the timepoint of cell division and
analysed Hes7 expression (Fig. 2C). The two daughter cells of a given mother
remained closely related in terms of oscillation phase for up to 8 hours after
division (Fig. S5C, Fig. 2A). Interestingly, we found that the oscillation phase of
Hes7 one hour before division was not randomly distributed (timepoint 1, Fig.
2C, D). Rather, at the point of cell division, there was a sudden drop in nuclear
Hes7 signal due to nuclear envelope breakdown. After division, individual cells
immediately showed Hes7 oscillations again (Fig. S5A), but aligned tracks were
not synchronized for approximately 4 hr (timepoint 2, Fig. 2C, D), after which
cells started to synchronize again (Fig. 2C, D, Fig. S5E). The timeframe of non-

synchronization after division roughly corresponds to the time cells spend in the
G1 phase of the cell cycle (4.4 hr, Fig. 1L).

The G1 restriction point is a point of no return, after which the cell cycle
can run autonomously, even though later restriction points can prevent cell
cycle progression to ensure the fidelity of cell division[26]. We therefore
addressed whether segmentation clock oscillations were correlated with the
G1-S transition. To this end, we combined Achilles-Hes7 with Fucci2-Cdt1 and
iRFPnucmem and quantified expression in single cells of *ex vivo* cultured
embryo tails after induction of mosaic labelling (Fig. 2E-G, Fig. S6A-D). We first
analysed the period of segmentation clock oscillations along the posterior-
anterior PSM axis and detected an increase in period towards the anterior (Fig.
S7A), consistent with an oscillation period gradient[27]. We then separated cell
tracks into their respective cell cycle phase based on Fucci2-Cdt1 reporter
expression. When directly comparing all cells in G1 to all cells in the S/G2/M
phase, independent of their location in the PSM, we detected a slightly higher
period in S/G2/M phase cells (Fig. S7B). However, if we considered the position
of the cells along the AP axis, we found that the period gradient was less
pronounced in G1 than in S/G2/M phase cells (Fig. S7C,D). This indicates that
segmentation clock oscillations change with both anteroposterior position in the
PSM and cell cycle phase, suggesting an effect of cell proliferation on the
segmentation clock[13,28].

Next, we tested for a potential correlation between segmentation clock
oscillations and the G1-S transition. When we aligned all tracks to the G1-S
transition, defined as the peak in Fucci2-Cdt1 signal (Fig. 1J), we did not
observe any relationship with the phase of Hes7 oscillations (Fig. 2G, Fig. S6E,
F). In mice, the segmentation clock consists of oscillations in Notch, Wnt and
FGF signalling. Previously, we demonstrated that the phase-relationship
between Wnt and Notch signalling oscillations changes along the posterior-
anterior axis of the PSM from out-of-phase in the posterior to in-phase in the
anterior[29] (Fig. 2I), corresponding to a 1π shift in phase-relationship. If an
equivalent shift in phase-relationship exists between the cell cycle and Hes7
oscillations, any correlation would be masked when analysing the signal
averaged along the anteroposterior axis. We therefore separated tracks from
cells residing in the posterior and anterior PSM and analysed whether Fucci2-

Cdt1 correlated with Hes7 dynamics. Notably, we found that the Fucci2-Cdt1

peak and the segmentation clock were correlated in both regions (Fig. 2G, Fig.

S6E, F). In posterior PSM, the phase of Hes7 oscillations was ~1.3π at G1-S

transition, while it was ~0.4π in anterior PSM. This corresponded to a change

in phase-relationship between Notch oscillations and cell cycle of ~0.9π

between posterior and anterior PSM, roughly matching the 1π shift in Wnt-

Notch phase-relationship. This provides indirect evidence that cell cycle

progression correlates with Wnt-signalling oscillations in the PSM.

While the cell cycle in a single cell takes approximately 14 hr and a single

cell spends approximately 4 hr in G1 phase, at the population level we observed

an oscillatory dynamic in the Fucci2-Cdt1 reporter similar to the segmentation

clock. Specifically, when quantifying small regions containing approximately 10-

15 cells, we observed oscillations in Hes7 and Fucci2-Cdt1 signal with a period

of roughly 2.5 hr (Fig. 2J,K, Movie S2). Such behaviour is consistent with a

model in which groups of cells in this region correlate their G1-S transition in a

manner that varies periodically between other neighbouring groups. This in turn

implies that neighbouring cells within these regions were shifted in their G1-S

transition by a multiple of complete segmentation clock periods relative to their

neighbours (Fig. 2L).

Altogether these data reveal that cell cycle progression is correlated with

signalling of the segmentation clock. More specifically, there is a constant

phase-relationship with Wnt-signalling oscillations along the AP axis of the

PSM.

The segmentation clock and cell proliferation are coupled

We next sought to determine whether cell proliferation is merely

correlated to the segmentation clock or whether there is a functional link

between the two. Continuous inhibition of Wnt signalling by genetic or chemical

perturbation leads to impaired PSM growth and a rapid regression of the PSM

due to accelerated differentiation and loss of the morphogen gradient[30,31] (Fig.

S8). Such continuous perturbations are therefore not well suited to reveal a

functional relationship between the segmentation clock and the cell cycle.

Therefore, we aimed for a system to subtly modulate oscillations. To achieve

this, we used our previously established microfluidic system to entrain

oscillations to external pulses of pathway modulator[29,32]. We cultured

embryonic tail tips in fibronectin-coated microfluidic chips, resulting in the

spreading of the embryonic tissue to form a quasi 2D monolayer (Fig. 3A). In

this system, the posterior-anterior embryonic axis is reorganized radially from

centre to periphery[3,29]. Combining this set-up with mosaic labelling allows

single-cell tracking and quantification of dim reporters, even by confocal

microscopy[19]. Using embryos expressing Achilles-Hes7, inducible Fucci2-Cdt1

and iRFPnucmem[18,25,33], we addressed whether a functional link exists

between proliferation and the segmentation clock. Importantly, the correlation

between segmentation clock (Hes7) oscillations and cell cycle (Cdt1) at the

population level was maintained in the 2D cultures (Fig. S9, Movie S3).

We then modulated the segmentation clock by periodically perturbing

Wnt signalling using the GSK3 inhibitor Chiron (CHIR99021) (Fig. 3B, Movies

S4). Our previous experiments had revealed that Chiron efficiently entrains

Wnt-signalling oscillations, which also affects Notch-signalling oscillations of

the segmentation clock[29]. Indeed, we confirmed that Chiron pulses entrained

the segmentation clock using Achilles-Hes7 as a reporter (Fig. 3C-F).

Subsequently, we analysed when the G1-S transition occurred (based on the

peak Fucci2-Cdt1 reporter signal) within one signalling oscillation. We found

that the G1-S transition preferentially coincided with drug pulses (Fig. 3G-I, Fig.

S10). Interestingly, our previous experiments examining the dynamics of the

Wnt-signalling reporter Axin2T2A revealed that this signal also peaked with

Chiron pulses[29], suggesting that the G1-S transition coincides with peaks in

Wnt-signalling oscillations. In contrast, DMSO pulses led to a random

distribution of Fucci2-Cdt1 peaks during the oscillation cycle (Fig 3I). Thus,

modulating segmentation clock oscillations via targeted Wnt signalling

modulation leads to an entrainment of cell cycle oscillations.

All signalling pathways of the segmentation clock are known mitogens

promoting cell cycle entry. Recently, myc and Cyclin D, well-studied for their

role in cell cycle progression and G1-S transition[34], were identified as potential

cycling genes at the mRNA level[6]. Indeed, using hybridization chain reaction

(HCR), we found Cyclin D1 to be expressed in the PSM with a dynamic pattern

overlapping with the Wnt-signalling target gene Axin2 (Fig. 3J, Fig. S11A).

Since cyclins are post-translationally regulated by modulating protein stability,

we stained for Cyclin D1 protein. The staining pattern for the protein also
corresponded to the Axin2 reporter (Fig. 3K, Fig. S11B). To determine a
functional link, we perturbed Wnt signalling and analysed the effect on Cyclin
D1 expression. Since Wnt inhibition leads to an immediate loss of PSM due to
differentiation, we instead activated Wnt signalling using Chiron. This led to an
increase in both Axin2 and Cyclin D1 expression in the entire PSM within 5
hours of activation (Fig. 3L, M).

We conclude that the segmentation clock and cell proliferation are
mechanistically coupled, presumably via periodic induction of cell cycle
regulators, such as Cyclin D1, by Wnt-signalling oscillations.

An extended somitogenesis model suggests that cell cycle-
segmentation clock coupling ensures the fidelity of somite formation
To determine whether the coupling of cell proliferation to signalling
oscillations might have a functional role in maintaining periodic somite
formation, we turned to a conceptual modelling-based approach, extending on
previous models of somitogenesis[4,35-37]. To develop the model, we placed
emphasis on the following assumptions based on our findings and available
literature: (1) Somite formation occurs at a certain phase between Wnt- and
Notch-signalling oscillations and a low threshold of the FGF morphogen
gradient[29,38,39]. (2) All PSM cells have a certain probability of dividing, and the
cell cycle period takes values in the range 13.6 ± 1.3 hr with a refractory period
of around 9.2 ± 0.5 hr after cell division, translating to the estimated length of
the S/G2/M phase of the cell cycle (Fig. 1K, L). (3) Since FGF mRNA is known
to be produced only in the posterior tailbud[39,40], it follows that the FGF
morphogen gradient is diluted by both mRNA degradation and cell division,
during which mRNA is partitioned equally between the two daughter cells. (4)
Only minimal rearrangement of cells in the one-dimensional model of the PSM
was allowed[41]. This last assumption was consistent with quantifications of cell
migration and rearrangement in the PSM, as tracked using light-sheet imaging
(Fig. 4A-C, Fig. S12A-C): Cells in the tailbud moved constantly, with a slight
preference in directionality towards the posterior end of the tissue (Movie S1,
Fig. 4A). The velocity of cells moving away from the tail tip increased with
increasing distance from the tail tip. In contrast, the movement of cells relative

to their neighbours in the PSM was restricted (0.2 µm/ 15 min, Fig. 4B, C); i.e., cells changed their relative position by less than one cell diameter during a single segmentation clock cycle. Moreover, we did not detect a preferential division axis between the anteroposterior or mediolateral axis (Fig. S5D).

Based on these results and assumptions, we modelled the dynamics of cells in the PSM as a one-dimensional line of Kuramoto-type phase oscillators coupled to local FGF density (for details of the model and its dynamics, see Supplementary Theory). Numerical simulations of the model recapitulated features of the continuously growing PSM field, showing the characteristic phase wave dynamic and periodic somite formation. Based on our observation that cell proliferation occurs along the entire PSM, we first asked what effect coupling between cell cycle entry and the segmentation clock has on somitogenesis and tissue patterning. To this end, we considered a line of 60 cells based on the number of cells in a line from posterior to anterior PSM and minimal cell exchange with neighbours (Fig. S12A-C and Fig. 4A-B). We then simulated tissue growth and somite formation (Fig. 4D top panels, Fig. S13). When cell divisions were allowed to occur randomly, decoupled from the segmentation clock, with a period of 13.6 ± 1.3 hr, numerical simulations revealed large variations in the somite scaling ratio, the ratio of the somite length over the PSM length (Fig. 4E, mean = 1.6, CV = 0.22). Moreover, uncoupled growth led to a non-uniform dilution of the FGF morphogen gradient (Fig. 4D, Fig. S13A). Conversely, when we imposed the condition that cell division occurs 1.25 hr after Wnt oscillations peak (5) (Fig 2D), tissue growth and somite formation were more robust, as evidenced by a sharper distribution of the somite scaling ratio (Fig. 4D, E, Fig. S13, B, mean = 1.7, CV = 0.15). This increase in robustness was observed consistently in simulations over a wide range of parameter values (Supplementary Theory). Together, these results suggest that the coupling of cell proliferation to the segmentation clock may be necessary to ensure robust tissue elongation and uniform dilution of the morphogen gradient, regulating the precise spacing of somite formation.

Perturbing cell proliferation interferes with morphogen gradient dilution and scaling of somite formation

To understand how cell proliferation affects the segmentation process,

we arrested cell proliferation in 2D *ex vivo* cultures[3] of embryos expressing

Fucci2-Cdt1, Achilles-Hes7 and iRFPnucmem using different cell cycle

inhibitors. Inhibition resulted in an accumulation of non-dividing cells and an

accumulation or absence of Fucci2-Cdt1, depending on when the arrest

occurred during the cell cycle (Movie S5). Upon inhibition, segmentation clock

oscillations were maintained (Fig. S14A, B), as shown previously[12]. This

indicates that cell cycle progression is not a necessary driver of segmentation

clock oscillations[12].

Based on the minimal model, simulations suggest that in the absence of

proliferation there would be a change in the scaling ratio from 0.171 in

conditions of coupled growth to 0.138 in the absence of cell divisions (Fig. 5A).

To challenge this prediction, we inhibited proliferation in growing embryo tails

using a strong continuous S-phase inhibitor, Aphidicolin (Fig. 5, Movies S6).

This led to a decrease in cell number, as expected, yet tissue size remained

largely unaltered in the timeframe of the experiment due to continued cell

growth, leading to a decrease in overall cell density (Fig. S14C-F). We then

analysed the effect of Aphidicolin on somite formation. This treatment led to the

formation of disorganized somites with undiscernible boundaries (Fig. 5B)[8]. In

control cultures, the scaling ratio was 0.172 with a CV of 0.13, corresponding

to the figure of CV = 0.15 for the model simulation. (Fig. 5C-E, 4E). Notably,

consistent with the predictions of the model under perturbed conditions (Fig.

5A), the ratio changed from 0.172 to 0.109, indicating altered scaling of somite

formation. We note that since (a) cell density differences between the PSM and

somites (Fig. S14E-G) and (b) cell velocity (Fig. S14H) were the same under

control conditions and following cell cycle inhibition, these cannot account for

the change in scaling.

Finally, the model also predicted an expansion of morphogen gradients

in the absence of cell divisions. We quantified morphogen gradients using

T/Brachyury expression as a readout, given that this is directly affected by FGF

and Wnt3a gradients[40,42]. Consistent with the model, the T/Brachyury gradient

was expanded in the unsegmented PSM upon cell cycle inhibition (Fig. 5F-H).

Similarly, the expression patterns of FGF8 and Axin2 mRNA were expanded

and less graded than in controls (Fig. S14I-L). Consistent with the expansion of

morphogen gradients[40], the size of the oscillating field, a measure of the extent

of differentiation, decreased less than in control tails (Fig. 5I-K, Fig. S14M).

Furthermore, we found that the wave velocity did not decrease to the same

extent as in control tissues but remained higher, despite the decrease in cell

number relative to control tissue (Fig. S14M, N). This contrasted with

unperturbed tissues, in which the wave velocity slows with decreasing tissue

size and cell number[3]. While the density of cells decreased, the wave velocity

increased in perturbed conditions. This implies that the relative phase-shift

between oscillations in neighbouring cells decreased in the absence of cell

division. Thus, absence of cell division impairs somite formation and scaling by

affecting tissue gradients.

Discussion

Here, we examined the role of cell proliferation in somitogenesis,

uncovering a tight correlation between cell cycle entry and the phase of the

segmentation clock oscillation. Further, we hypothesized and validated a

mechanism by which the coupling of cell division to the segmentation clock may

ensure uniform PSM growth and dilution of regulatory molecules, thereby

modulating the scaling and accuracy of somite formation.

In contrast to studies that support a posterior growth zone[43-45], we found

that cell proliferation was uniformly spread over the posterior and anterior PSM,

consistent with previous analyses[2,10]. Furthermore, cell proliferation was

coupled to the segmentation clock, likely via Wnt-induced expression of pro-

proliferative factors such as Cyclin D[6], which ensures uniform distribution of

proliferating cells over the PSM. With a proliferation rate of approximately 14 hr

compared to a somitogenesis period of 2.7 hr, we found that cell proliferation is

a key component of mouse somitogenesis. Upon cell cycle inhibition, cell

division was abolished while tissue and cell growth continued, indicating that

cell division in the PSM has a negligible effect on tissue growth within the

timescale of the experiment.

Morphogen gradients are generated by mRNA and protein production in

the posterior tip and levels slowly decrease as cells move down the posterior-

anterior axis of the PSM[39]. Thus, morphogen gradients are modified by mRNA

(and protein) degradation as well as cell division, which leads to a continuous

decrease in morphogen concentration and nuclear effector proteins in cells.

Indeed, inhibiting cell division in the tissue expanded the morphogen gradients.

In agreement with the clock-and-wavefront model[37], regression of the

oscillating field was also reduced upon cell cycle inhibition. Moreover, signalling

wave velocity was increased and somite formation impaired. This suggests that

the principal role of cell division in the PSM may not lie in its contribution to

tissue growth but rather in shaping morphogen gradients for proper somite

formation.

This is consistent with a model (Fig. 5L) in which cell divisions are

uniformly distributed over the tissue by coupling to the segmentation clock. This

leads to an increase in cell number uniformly distributed over the PSM, which

leads to (1) a uniform expansion of the oscillating field and (2) a uniform dilution

of morphogen mRNA and protein as well as downstream gradients. These

gradients are known to affect wave dynamics[46] and both in turn regulate somite

formation[38,40]. Supported by mathematical modelling, this suggests that the

coupling of cell divisions to the segmentation clock ensures the formation of

consistently scaled somites. Scaling of segment size to PSM size has

previously been correlated with scaling of wave dynamics[3] or the FGF gradient[4]

in mouse and zebrafish, respectively, both of which we found to be affected by

cell cycle inhibition. Thus, this constitutes a mechanism by which local cell-

intrinsic regulations, i.e., whether to divide or not, impact the tissue-wide

regulation of morphogenesis to ensure robust pattern scaling and somite

formation.

How our findings translate to other species remains to be addressed. In

zebrafish, continued tissue growth, impaired somite formation and decreased

cell density are observed in the PSM of *emi1* mutants, which stop proliferation

during gastrulation[8]. Moreover, segmentation clock oscillations correlate with

cell divisions in zebrafish[14], even though cell proliferation is thought to play a

minor role for tissue growth[47]. The coupling we observed could explain why

zebrafish segmentation proceeds more quickly than growth and oscillation

regression, leading to a rapid reduction of tissue size and a decrease in

oscillation wave amplitude from 4π to 2π over developmental time[36]. In human

somitogenesis, a correlation between the timepoint of cell division and Hes7

oscillations has so far not been observed[12]. To clarify the role of cell proliferation

in human cells, cell cycle reporters specific for the G1-S transition should be

quantified relative to Hes7 oscillations in posterior vs. anterior PSM cells.

Signalling oscillations have been observed in other tissue types and

developmental contexts. Correlations between such oscillations and cell

division have also been described[48]. For instance, in *in vitro* cultures of the

small intestine, we recently identified Hes1 oscillations that correlated with cell

divisions[49]. In neural stem cells, interactions between Hes1 oscillations and the

cell cycle inhibitor p21 have been described[50]. Whether these other systems

depend on active coupling between the cell cycle and signalling oscillations, as

shown here for somitogenesis, and whether this coupling impacts growth,

differentiation and morphogenesis at the tissue level are exciting questions for

future research.

        Figure Legends

Figure 1 Cell cycle dynamics along the entire E10.5 presomitic mesoderm.
A-H Quantification of cell cycle dynamics in uncultured tails. A 5 µm section of
E10.5 tail with immunostaining for Ki67 and counterstaining with DAPI. **B** Flow
cytometry analysis of embryonic tail cells stained for Tbx6 and DNA using
Hoechst. N=3 experiments. Raw data shown in Fig. S4. **C-E** Quantification of
late mitotic cells in embryonic tails. N=5. **C** Embryonic tails were sectioned into
5 µm sections. Immunostaining against phosphorylated-histone H3 (PHH3) and
counterstaining with DAPI were performed. For the analysis, sections were
divided into tailbud (posterior to neural tube), PSM and somite regions. **D**
Representative image of a section stained against PHH3 and with DAPI (same
section as in Fig. S2A). **E** Quantification of cells in late mitosis (pro-metaphase
to anaphase) in the regions indicated in **B**. **F-H** Quantification of cell cycle
dynamics by EdU-BrdU double-labelling. N=3. **F** Workflow outlining
experimental steps. Embryos were dissected and tails sectioned akin to **B**
before EdU/BrdU staining. **G** Representative image of a section stained for
EdU, BrdU and nuclei (same image as in Fig. S3A). **H** Quantification of cell

cycle length in the regions indicated in **B**. Coefficient of variation CV: CV(Tailbud) = 0.19; CV(PSM) = 0.02; CV(Somite) = 0.16. N = 3. **I-N** Quantification of cell cycle dynamics in *ex vivo* cultured tails. **I** Schematic illustrating single cell tracking pipeline. Pregnant mice were injected with low doses of Tamoxifen on E9.5 for sparse nuclear labelling of cells. Embryos were dissected after 24h of incubation. Embryonic tails were dissected and imaged with an inverted light-sheet microscope. Embryonic tails were placed in agarose grooves for microscopy. Mosaically labelled cells with nuclear marker iRFPnucmem were followed over time in 3D. **J** Representative timeseries data of the dynamics of the G1 marker Fucci2-Cdt1 in a single cell tracked from anaphase to anaphase. **K, L** Quantification of cell cycle length and duration that cells were positive for the G1 marker Fucci2-Cdt1. **M** Patterns of G1-S transition in uncultured E10.5 embryonic tails (same image as in Fig. S4F). Cells double-positive for Fucci1-Cdt1 and Fucci1-Gem were determined in PSM of embryonic tails. Tails were ordered into three phases according to their staining pattern: Phase 1: no double-positive cells in PSM; Phase 2: stripe of double-positive cells in PSM; Phase 3: double-positive cells distributed over PSM. Number in brackets indicates number of tails in which specific staining pattern was observed. Brightfield (*left panel*), merge of Fucci1-Cdt1 and Fucci1-Gem (*middle panel*) and cells double-positive for Fucci1-Cdt1 and Fucci1-Gem (*right panel*) depicted. N=13. Scale bar 100 µm. P indicates the posterior end of the embryo.

Figure 2 Cell cycle dynamics correlate with segmentation clock. A-D Correlation of Hes7 segmentation clock with cell division. **A** Experimental approach for sparse labelling of nuclei with H2B-mCherry combined with Achilles-Hes7. **B** Representative snapshots of timeseries data. Arrowhead points at one cell tracked over time. **C** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal of cells aligned to timepoint of cell division. **D** Schematic representation of a single phase wave. Phase of Achilles-Hes7 oscillations at timepoint 1, as indicated in **C**, (*left panel*) and at timepoint 2, as indicated in **C**, (*right panel*). **E-G** Correlation of Hes7 segmentation clock to G1-S transition. **E** Experimental approach for sparse labelling of nuclei with iRFPnucmem and Fucci2-Cdt1 combined with Achilles-Hes7. **F** Representative snapshots of timeseries data. Arrowhead points at one cell tracked over time. **G** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal of cells aligned to timepoint of G1-S transition. Left panel: all cells; middle panel: cells in posterior PSM; right panel: cells in anterior PSM. **H** Phase of Achilles-Hes7 oscillations at G1-S transition. Left panel: all cells; middle panel: cells in posterior PSM; right panel: cells in anterior PSM. **I** Schematic of the phase change in Wnt and Notch signalling between anterior and posterior PSM. **J** Maximum Intensity projection of E10.5 embryonic tail (purple: Achilles-Hes7, cyan: Fucci2-Cdt1). **K** Quantification of Achilles-Hes7 and Fucci2-Cdt1 in

518 regions of interest (ROIs) as indicated in **J. L** Scheme illustrating correlation of

519 cell cycle (duration 13.6 hr) to segmentation clock oscillations (2.7 hr).

Figure 3 Coupling of cell proliferation to segmentation clock oscillations. A-H Microfluidics to test functional link. **A** Schematic illustrating a 2D tail tip culture spreading over time in the side view and bottom view of the embryo. **B** Experimental approach: Microfluidic system to modulate segmentation clock oscillations in *ex vivo* tailbud cultures. Analysis of segmentation clock and cell proliferation. **C-F** Analysis of segmentation clock: **C,E** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal in anterior PSM upon treatment with DMSO (**B**) and 5 μM Chiron (**E**). **D,F** Phase of Hes7 oscillations plotted against phase of external drug pulses for treatment with DMSO (**D**) and 5 μM Chiron (**F**). **G** Snapshots of

representative fluorescence real-time imaging data highlighting tracking of a
single cell (arrowhead). **H** Representative timeseries data showing Fucci2-Cdt1
reporter intensity of a single cell during microfluidic experiment. Grey shading
indicates external drug pulse. **I** Phase of external drug pulses, at which G1-S
transition occurs in DMSO- (*left panel*) or Chiron-treated samples (*right panel*).
J Staining for mRNA levels of Axin2, Cyclin D1 and Hes7 in E10.5 embryonic
tails using hybridization chain reaction (HCR) (same image as in Fig. S11A). **K**
Immunostaining against Cyclin D1 in Axin2T2AVenus-expressing E10.5
embryonic tails (same as in Fig. S11B). **L-M** Treatment of E10.5 embryonic tails
with DMSO or Chiron (10 µM). HCR for Axin2 (upper panels) or Cyclin D1
(lower panels). **L** representative images. **M** Quantification of Axin2 and Cyclin
D1 in regions of interest as indicated in representative images. $* < 0.05$; $** <$
0.01.
545

Figure 4 Coupling of cell proliferation to the segmentation clock promotes uniform PSM growth and somite formation in simulations. A-C E10.5 embryonic tails were cultured *ex vivo* and fluorescence real-time imaging was performed (corresponds to data from Figure 1,2). **A** Velocity of cells moving relative to the posterior tip (*top panel*). This indicates stretching of the PSM, as cells further away from the posterior have a higher velocity (*bottom panel*). Arrows indicate direction of movement. N = 6 tissues n = 106 tracks. * < 0.05; ** < 0.01; *** < 0.001; **** <0.0001. **B** Plots of single-cell tracks where whole-tissue movement was computationally removed to reveal movement of single cells relative to their surroundings. Colours correspond to different cells. Each dot represents location of a cell at specific timepoint (interval between timepoints 15 min). **C** Histogram of the amount of cell movement given in % of cell diameter (10 µm). N = 9 tissues n = 169 tracks. **D** Simulation of somitogenesis with non-coupled cell proliferation (top panels) or proliferation coupled to segmentation clock (bottom panels H). Example plots for simulations of signalling waves (*left panels*) as well as FGF gradient (gray shading) and somite formation (blue line) (*right panels*). Arrowheads indicate somite

formation. Annotations are: Somites (S), Anterior (A) and Posterior (P). **E** Histogram showing scaling ratios, defined as the ratio of somite length to the PSM length based on 100 simulations (for details of the model and its parameters, see main text and Supplementary Theory). Coefficient of variation CV(non-coupled proliferation) = 3.5 cell diameters. CV(coupled proliferation) = 2.7 cell diameters.

Figure 5 Coupling of cell cycle to segmentation clock affects scaling of somite formation. A Simulations of the somitogenesis model with cell cycle progression coupled to signalling (left) and following perturbation in which cell cycle is inhibited (right). Note the change in the average somite scaling ratio between the two conditions. **B-K** E10.5 embryonic tails were cultured *ex vivo* and treated with DMSO or the cell cycle inhibitor Aphidicolin. **B-E** After 23 hr of incubation, tails were fixed and stained with DAPI. **B** Representative images are shown. **C-D** Quantification of PSM length from tail tip to last formed somite

boundary (**C**), somite length (**D**) and the ratio of somite length to PSM length as

a measure of scaling (**E**) **F-H** After 23 hr of incubation, tails were fixed and

stained for Brachyury and counterstained with DAPI. Scale bar 100 µm. **F**

Representative images. **G** Quantification of the Brachyury signal along the AP

axis of the PSM. Signal was normalized between 0 and 1. n(DMSO) = 6,

n(Aphidicolin) = 11. **H** Quantification of the length of Brachyury gradient (Line:

mean, shading: standard deviation). Intensity normalized between 0 and 1. $^{*} <$

0.05; $^{***} < 0.001$; $^{****} < 0.0001$. **I,J** Fluorescence real-time imaging of embryonic

tails expressing Achilles-Hes7 was performed. **I** Representative kymographs of

control and Aphidicolin-treated samples. **J** Quantification of PSM length by

determining the size of the oscillating field (Line: mean, shading: standard

deviation). Length normalized to the PSM length at experiment start. n(DMSO)

= 9, n(Aphidicolin) = 11. **K** After 23 hr of incubation, tails were fixed and HCR

against Hes7 and Uncx4.1 as well as counterstaining with DAPI were

performed. Representative images are shown (same as in Fig. S14J). **L**

Schematic model of how coupling of cell proliferation to segmentation clock

affects somitogenesis. Morphogen density is indicated in magenta.

Methods

Animals and Housing

All experimental procedures were conducted in accordance with institutional guidelines and approved by the Ethical Review Board of the Hubrecht Institute. Animal protocols were reviewed and authorized by the Animal Experimentation Committee (DEC) of the Royal Netherlands Academy of Arts and Sciences (KNAW). Mice were bred and housed under standard conditions at the Hubrecht animal facility, with a controlled environment maintained at 50–60% humidity, 22–23°C temperature, and a 12-hour light/dark cycle. Mice had ad libitum access to water and standard chow throughout the study.

Mouse Strains

All transgenic mouse lines used in this study have been previously described. The Achilles-Hes7 line was generously provided by the Kageyama laboratory[25]. Fucci [21], The R26R-H2B-mCherry, and R26R-mCherry-hCdt1(30/120)[24] lines were obtained from the RIKEN Center for Biosystems Dynamics Research and the RIKEN BioResource Research Center (accession numbers CDB0204K, RBRC02706, and CDB0229K, respectively) via the RIKEN Mouse Repository. The Tg(T-cre/ERT2)1Lwd/J ("T-CreERT2") line was sourced from The Jackson Laboratory (accession number #025520). The R26R-iRFPnucmem[19] and Axin-T2A-Venus[29] lines were provided by the Aulehla laboratory. All wild-type mice used in this study were F1 hybrids (BALB/cJ × C57BL/6J).

Animal and Embryo Work

EdU/BrdU Incorporation

Female mice were sacrificed at 10.5 days post-coitum (dpc) and embryos were subsequently dissected. For EdU and BrdU incorporation experiments, pregnant females at 10.5 dpc were intraperitoneally injected with 3.3 mg EdU in PBS, followed one hour later by an equimolar injection of 4.0 mg BrdU in PBS[51]. Mice were sacrificed 30 minutes after BrdU injection. Embryos

were dissected in PBS supplemented with 1% Bovine Serum Albumin (Biowest,

P6154) for tail extraction.

<u>Mosaic Labelling</u>

For mosaic labelling experiments, pregnant females were

intraperitoneally injected with tamoxifen (Sigma-Aldrich, T5648, dissolved in

corn oil) on embryonic day E9.5. To label with H2B-mCherry, 1 mg of tamoxifen

was administered, whereas for labelling with iRFPnucmem and Fucci reporters,

0.1 mg of tamoxifen was used.

<u>Flow Cytometry</u>

For flow cytometry analysis, E10.5 mouse tails were extracted and cut

into smaller pieces using a surgical scalpel to facilitate single-cell dissociation.

Tail fragments were washed three times in PBS0 in Eppendorf tubes, followed

by a 15-minute incubation with 400 µl of TrypLE (Thermo Fisher, 12604013).

After incubation, the tissue was passed through a 30G needle to generate a

single-cell suspension. The suspension was then washed twice in PBS0

supplemented with 1% Bovine Serum Albumin (Biowest, P6154) by

centrifugation at 500 g for 5 minutes.

To fix the cells, the pellet was resuspended in 4% paraformaldehyde

(PFA) and incubated at room temperature for 15 minutes. After two additional

washes in PBS with 1% BSA, cells were stored at 4°C until further staining. For

immunofluorescent staining, the single-cell pellet was resuspended in 3% BSA

PBS with 0.25% Triton X-100 and incubated overnight at 4°C with anti-Tbx6

rabbit primary antibody (Abcam, ab38883) at a dilution of 1:1000. After three

washes in 3% BSA PBS, the cells were incubated with secondary donkey anti-

rabbit Alexa Fluor 488 antibody (Thermo Fisher, A-21206) at a dilution of

1:1000 for 2 hours at room temperature.

Following a further two washes, the single-cell suspension was

resuspended in 500 µl of PBS supplemented with Hoechst 34580 (1 µg/ml) and

passed through a 20 µm mesh. Flow cytometry was performed on a CytoFLEX

S Flow Cytometer, and the Tbx6-positive cell population was selected for cell

cycle analysis using FlowJo software.

<u>Culture of Live Embryonic Tissue</u>

Culture of whole mount and spread-out tail tip tissues was performed as
previously described[29]. For perturbation experiments on whole mount and
spread-out tail tips, the following compounds were used: CHIR99021 (Chiron,
Sigma-Aldrich, SML1046) at 10 µM, IWP-2 (Sigma-Aldrich, I0536) at 1 µM,
Aphidicolin (ThermoFisher, J60236.MCR) at 1.6 µg/mL, Thymidine (Sigma-
Aldrich, T9250) at 2 mM, Palbociclib (Selleckchem, PD-0332991) at 1 µM, and
RO-3306 (Sigma-Aldrich, 217699) at 10 µM.
672
Microfluidics Experiment
The full protocol for performing the microfluidics experiment has been
described elsewhere[29,32]. For entrainment of Achilles-Hes7 and Fucci2-Cdt1
reporter lines, a pulsing scheme was employed consisting of 100 minutes of
medium followed by 30 minutes of medium containing CHIR99021 (Chiron,
Sigma-Aldrich, SML1046) at 5 µM or DMSO.
679
Confocal Live Imaging
Confocal live imaging was performed using a Leica TCS SP8 confocal
microscope equipped with a HC PL APO CS2 20×/0.75 DRY objective. For cell
tracking experiments, images were acquired every 10 minutes with a z-step of
5 µm, a stack depth of 50 µm, and a z-resolution of 4 µm. For non-tracking
imaging, images were taken every 10 minutes with a z-step of 10 µm, a stack
depth of 100–150 µm, and a z-resolution of 9 µm. Excitation of Achilles,
mCherry, or iRFP signals was performed using OPSL lasers at 514 nm, 568
nm, or 638 nm, respectively. Movies were recorded at a resolution of 512 × 512
pixels (pixel size 1.517 µm). For wholemount tissues, imaging was conducted
in a humidified atmosphere maintained at 37°C, with 5% CO_2 and 60% O_2. For
2D cultures the O_2 was reduced to 20%.
692
Light-sheet Live Imaging
Live imaging of whole mount embryo tails for single-cell analysis was
performed using a Viventis light sheet microscope (Viventis Microscopy, #LS1
Live). Imaging was conducted in a humidified atmosphere maintained at 37°C,
with 5% CO_2 and 60% O_2. For fluorescence excitation, lasers with wavelengths

of 514 nm, 568 nm, and 687 nm were used for Achilles, mCherry, and iRFP,

respectively. Imaging was carried out with a 25× objective, using a laser beam

of 3.3 μm and acquiring a z-stack of 60 steps, each with a 3 μm increment.

Images were captured at a resolution of 2048 × 2048 pixels every 15 minutes.

To prevent whole mount tissues from moving out of the field of view while still

allowing unperturbed growth, a 2% agarose gel in PBS was cast around a

custom-made insert in the sample holder. This insert created five spaces (1.5

mm × 0.5 mm) for the individual tissues to be loaded.

<u>Paraffin Sectioning of Tissues and Immunofluorescence</u>

Tissues fixed overnight in 4% paraformaldehyde (PFA) at 4°C were

stored in 70% ethanol at 4°C for up to one week prior to embedding. Embryo

tails were then subjected to Xylene treatment, embedded in paraffin in groups

of three, and sectioned transversally into 5 μm thick slices.

For staining, the slides were sequentially immersed in Xylene, followed

by 100%, 95%, 70%, and 50% ethanol, and rinsed with running cold tap water

for 3 minutes each. Antigen retrieval was performed by heating the slides in a

citrate buffer (8.2 mM sodium citrate and 1.8 mM citric acid, pH 6). Slides were

blocked and permeabilized using 1% Bovine Serum Albumin (BSA, Biowest,

P6154) in 0.5% Triton X-100 PBS. Mitotic cells were stained with anti-phospho-

histone H3 (Sigma-Aldrich, 05-806) at a 1:1000 dilution in 1% BSA PBS, and

incubation was carried out overnight at 4°C. Secondary staining was performed

using Donkey anti-Mouse Alexa Fluor 488 (Thermo Fisher, A-21202) at a

1:1000 dilution, supplemented with DAPI (5 μg/mL, Sigma) in 1% BSA PBS,

and incubation was carried out overnight at 4°C. After staining, slides were

mounted with Vectashield (H1200NB, Vector Laboratories). Images were

acquired using a Leica TCS SP8 confocal microscope with an HC PL APO CS2

20×/0.75 DRY objective and OPSL lasers at 365 nm and 488 nm for excitation.

For BrdU and EdU staining, two additional steps were incorporated. Prior

to blocking and after antigen retrieval, slides were incubated at room

temperature in 2 M HCl for 10 minutes. EdU was then visualized using the Alexa

Fluor™ 647 dye from the Click-iT™ EdU Cell Proliferation Kit for Imaging

(Thermo Fisher, C10340). BrdU was detected using an anti-BrdU antibody

(Thermo Fisher, B35128) at a 1:500 dilution, followed by secondary staining

with Donkey anti-Mouse Alexa Fluor 568 (Thermo Fisher, A10037).

Whole Mount Tissue Antibody Staining

Prior to immunostaining, samples were fixed overnight in 4%

formaldehyde at 4°C and stored in PBS at 4°C. Primary antibodies used for

staining included anti-Ki67 (Abcam, ab16667) at 1:1000, anti-T (human/mouse)

(R&D Systems, AF2085-SP) at 1:1000, and anti-Cyclin D1 (Cell Signaling,

2978) at 1:500. Secondary antibodies included Donkey anti-Goat Alexa Fluor

488 (Thermo Fisher, A-11055) at 1:1000 and Donkey anti-Rabbit Alexa Fluor

568 (Thermo Fisher, A10037) at 1:1000.

Immunostaining was carried out by incubating the samples with primary

and secondary antibodies overnight in PBST (0.1% Triton X-100) containing

1% Bovine Serum Albumin (BSA, Biowest, P6154). The secondary antibody

mixture was supplemented with DAPI (Sigma, 5 µg/mL) for nuclear staining.

For imaging, a z-stack consisting of 10-12 planes with a 10 µm distance

between each plane was acquired at a resolution of 1024 × 1024 pixels (pixel

size 0.758 µm) using a Leica TCS SP8 MP confocal microscope with a HC PL

APO CS2 20×/0.75 DRY objective. Excitation was performed with OPSL lasers

at 365 nm, 488 nm, and 568 nm.

Whole Mount Tissue Hybridization Chain Reaction (HCR)

Hybridization chain reaction (HCR) was performed as described

previously[52]. Cyclin D1, Hes7, Uncx4.1 and Fgf8 probe sets were generated

using a probe generator (https://github.com/rwnull/insitu_probe_generator).

Probe sets for Axin2, Hes7, and Uncx4.1 were obtained from Molecular

Instruments. HCR imaging was performed using a Leica TCS SP8 confocal

microscope with a HC PL APO CS2 20×/0.75 DRY objective. The system was

equipped with OPSL lasers for excitation at 365 nm, 488 nm, 568 nm, and 638

nm. Probe sets for Axin2 were obtained from Molecular Instruments. HCR

imaging was performed using a Leica TCS SP8 confocal microscope with a HC

PL APO CS2 20×/0.75 DRY objective. The system was equipped with OPSL

lasers for excitation at 365 nm, 488 nm, 568 nm, and 638 nm.

<u>Image and Data Processing</u>

To quantify the fraction of phospho-histone H3 (pHH3)-positive cells,
skin, neural tube, notochord, and gut were manually removed from the DAPI
channel in section images using Fiji. Subsequently, StarDist, a Fiji plugin, was
used to segment the remaining nuclei, providing an accurate measurement of
the number of nuclei per slice. pHH3-positive cells were manually counted per
slice.

For the quantification of EdU- and BrdU-positive cells, skin, neural tube,
notochord, and gut were similarly excluded from the DAPI channel in Fiji. This
channel was further denoised using Remove Outliers in Fiji (pixel size= 4,
threshold=50) followed by a gaussian blur with a sigma of 2. Using Cellpose
cryo3 model we segmented the remaining nuclei, allowing for an accurate count
of the total number of nuclei per slice. Mean intensities of EdU and BrdU were
calculated within the DAPI masks, and thresholds were applied to categorize
cells as positive or negative for EdU and/or BrdU. These annotations,
expressed as fractions of the total cell count, were used for the quantification
of cell cycle duration, population doubling time, and S-phase duration (see Fig.
1, Fig. S2,3).

To visualize tissue-wide dynamics of Hes7 and Fucci2-Cdt1, regions of
interest (ROIs) were drawn in maximum intensity projections of time-lapse
movies in Fiji. Intensity measurements of both channels were quantified over
time, smoothed, and linearly normalized between 0 and 1. Kymographs of
tissue-wide dynamics were generated using the KymoResliceWide plugin in
Fiji. A line was drawn from the posterior to the anterior end of the tissue, with a
width of 100 µm.

For the quantification of the T/Brachyury gradient, a similar approach
was used. A line plot was generated by measuring the intensity
of T/Brachyury along a line drawn in Fiji. Gradient length was defined as the
distance from the tip of the tail to the point where T/Brachyury intensity reached
10% above background level. To quantify staining intensities, an ROI was
manually drawn within the PSM, and the mean intensity was calculated using
Fiji. For the measurement of PSM and somite lengths, DAPI images were used.
Lines were drawn parallel to the neural tube from the tail tip to the first somite

boundary for PSM size, or from the first to the second somite boundary for
somite size.

For the quantification of tissue density, an ROI was drawn within confocal
or light sheet images, and the number of cells within the selected area was
counted using either the Achilles-Hes7 or DAPI channel.

To track single cells over short time intervals in confocal microscopy
movies, an ROI of 3 × 3 pixels was manually tracked within the Fucci2-
Cdt1 or Achilles-Hes7 channel in Fiji. The peak intensity of Fucci2-Cdt1 was
defined as the moment when a cell transitions from G1 to the S-G2-M phase.
These tracks were used to quantify cell velocity or the timing of cell cycle
transitions.

For long-term single-cell tracking in light sheet images, cells were
manually tracked by following mosaically expressed H2B or Nucmem reporters
using the Fiji plugin Mastodon (https://github.com/mastodon-sc/mastodon).
Spot centre intensities were measured within the tracked spots to
quantify Achilles-Hes7 or Fucci2-Cdt1. These intensities were smoothed using
a moving average of 3 time points and linearly normalized, such that each track
contained intensity values between 0 and 1. For the analysis of Hes7 dynamics,
PyBOAT was used with the following settings: min_period = 60 min,
max_period = 300 min, and TC = 150. All data plotting was performed using
Python. For binning the cell positions along the tailbud-to-somite axis we
tracked the posterior tail tip and the forming somite boundaries over time and
quantified a relative AP position of the cells were at any given time. We
considered 0.0-0.6 as posterior and 0.6-1.0 as anterior.

Statistical Analysis

All experiments were conducted with a minimum of three independent
replicates. Data visualization was performed using Tukey-style boxplots, where
the bar represents the median, the box delineates the 25th and 75th
percentiles, and the whiskers extend to 1.5 times the interquartile range (IQR).
Violin plots were used to display the distribution of data from the maximum to
the minimum, with the bar indicating the median value. For statistical
comparisons between groups, normality was assessed using the Shapiro-Wilk
test. Statistical significance was determined using either an independent t-test

or the Mann-Whitney U test, depending on whether the data followed a normal

distribution. The coefficient of variation CV is calculated from the standard

deviation std and mean as follows: CV = std/mean.

Data and Code availability

All data and custom code used in this study are available upon request

to k.sonnen@hubrecht.eu.

Acknowledgements

We thank members of the Sonnen group for discussions and feedback. We are grateful to Takashi Hiiragi and Juan Garaycoechea for discussions and feedback on the manuscript. We thank Cantas Alev for discussions. We thank Jan-Daniël de Leede for help with data analysis. We thank the animal, FACS and imaging facilities of the Hubrecht Institute for their support. We thank Alexander Aulehla for support and providing mouse lines, in particular the inducible iRFPnucmem line. We thank Andrea Boni for help with setting up the embryo-tail imaging in the light-sheet microscope. The H2B-mCherry mouse line was kindly provided by the Riken Center for Biosystems Dynamics Research and the Fucci line by the Riken BioResource Research Center. The T-Cre/ERT2 line was kindly provided by Jackson Laboratories. The Achilles-Hes7 was kindly provided by Ryoichiro Kageyama. This work was supported by the Hubrecht Institute and received funding from the European Research Council under an ERC starting grant agreement no. 850554 to K.F.S. In addition, this research was supported in part by grant NSF PHY-1748958 to the Kavli Institute for Theoretical Physics (KITP). B.D.S. is supported by the Wellcome (219478/Z/19/Z) and the Royal Society (RP\R\231004).

Author Contributions

M.J.O. and K.F.S. conceived the project together with Y.I.L. and B.D.S.; K.F.S. supervised the project; M.J.O. performed experiments with help from W.H.M.M., T.N., C.F., E.T., J.K. and W.M.T.; M.J.O., Y.I.L. and K.F.S. analysed the data; Y.I.L. and B.D.S. developed the theoretical framework; M.J.O., Y.I.L. and K.F.S. wrote the manuscript with input from all authors.

References

Meijer, W. H. M. & Sonnen, K. F. From signalling oscillations to somite formation. *Current Opinion in Systems Biology* **39**, 100520 (2024). https://doi.org/https://doi.org/10.1016/j.coisb.2024.100520

Tam, P. P. The control of somitogenesis in mouse embryos. *J Embryol Exp Morphol* **65 Suppl**, 103-128 (1981).

Lauschke, V. M., Tsiairis, C. D., Francois, P. & Aulehla, A. Scaling of embryonic patterning based on phase-gradient encoding. *Nature* **493**, 101-105 (2013). https://doi.org/10.1038/nature11804

Ishimatsu, K. *et al.* Size-reduced embryos reveal a gradient scaling-based mechanism for zebrafish somite formation. *Development* **145** (2018). https://doi.org/10.1242/dev.161257

Hubaud, A. & Pourquie, O. Signalling dynamics in vertebrate segmentation. *Nat Rev Mol Cell Biol* **15**, 709-721 (2014). https://doi.org/10.1038/nrm3891

Matsuda, M. *et al.* Recapitulating the human segmentation clock with pluripotent stem cells. *Nature* **580**, 124-129 (2020). https://doi.org/10.1038/s41586-020-2144-9

Dequeant, M. L. *et al.* A complex oscillating network of signaling genes underlies the mouse segmentation clock. *Science* **314**, 1595-1598 (2006). https://doi.org/10.1126/science.1133141

Zhang, L., Kendrick, C., Julich, D. & Holley, S. A. Cell cycle progression is required for zebrafish somite morphogenesis but not segmentation clock function. *Development* **135**, 2065-2070 (2008). https://doi.org/10.1242/dev.022673

Benazeraf, B. *et al.* Multi-scale quantification of tissue behavior during amniote embryo axis elongation. *Development* **144**, 4462-4472 (2017). https://doi.org/10.1242/dev.150557

Bulusu, V. *et al.* Spatiotemporal Analysis of a Glycolytic Activity Gradient Linked to Mouse Embryo Mesoderm Development. *Dev Cell* **40**, 331-341 e334 (2017). https://doi.org/10.1016/j.devcel.2017.01.015

Mongera, A., Michaut, A., Guillot, C., Xiong, F. & Pourquie, O. Mechanics of Anteroposterior Axis Formation in Vertebrates. *Annu Rev Cell Dev Biol* **35**, 259-283 (2019). https://doi.org/10.1146/annurev-cellbio-100818-125436

Diaz-Cuadros, M. *et al.* Metabolic regulation of species-specific developmental rates. *Nature* **613**, 550-557 (2023). https://doi.org/10.1038/s41586-022-05574-4

13 Carrieri, F. A. *et al.* CDK1 and CDK2 regulate NICD1 turnover and the
periodicity of the segmentation clock. *EMBO Rep* **20**, e46436 (2019).
https://doi.org/10.15252/embr.201846436

14 Delaune, E. A., Francois, P., Shih, N. P. & Amacher, S. L. Single-cell-
resolution imaging of the impact of Notch signaling and mitosis on
segmentation clock dynamics. *Dev Cell* **23**, 995-1005 (2012).
https://doi.org/10.1016/j.devcel.2012.09.009

15 Wiedermann, G. *et al.* A balance of positive and negative regulators
determines the pace of the segmentation clock. *Elife* **4**, e05842 (2015).
https://doi.org/10.7554/eLife.05842

16 Kicheva, A. *et al.* Coordination of progenitor specification and growth in
mouse and chick spinal cord. *Science* **345**, 1254927 (2014).
https://doi.org/10.1126/science.1254927

17 Boehm, B. *et al.* The role of spatially controlled cell proliferation in limb
bud morphogenesis. *PLoS Biol* **8**, e1000420 (2010).
https://doi.org/10.1371/journal.pbio.1000420

18 Abe, T. *et al.* Visualization of cell cycle in mouse embryos with Fucci2
reporter directed by Rosa26 promoter. *Development* **140**, 237-246
(2013). https://doi.org/10.1242/dev.084111

19 El Azhar, Y. *et al.* Unravelling differential Hes1 dynamics during axis
elongation of mouse embryos through single-cell tracking. *Development*
151 (2024). https://doi.org/10.1242/dev.202936

20 Anderson, M. J. *et al.* TCreERT2, a transgenic mouse line for temporal
control of Cre-mediated recombination in lineages emerging from the
primitive streak or tail bud. *PLoS ONE* **8**, e62479 (2013).
https://doi.org/10.1371/journal.pone.0062479

21 Sakaue-Sawano, A. *et al.* Visualizing spatiotemporal dynamics of
multicellular cell-cycle progression. *Cell* **132**, 487-498 (2008).
https://doi.org/10.1016/j.cell.2007.12.033

22 Palmeirim, I., Henrique, D., Ish-Horowicz, D. & Pourquie, O. Avian hairy
gene expression identifies a molecular clock linked to vertebrate
segmentation and somitogenesis. *Cell* **91**, 639-648 (1997).
https://doi.org/10.1016/s0092-8674(00)80451-1

23 Evrard, Y. A., Lun, Y., Aulehla, A., Gan, L. & Johnson, R. L. lunatic fringe
is an essential mediator of somite segmentation and patterning. *Nature*
394, 377-381 (1998). https://doi.org/10.1038/28632

24 Abe, T. *et al.* Establishment of conditional reporter mouse lines at
ROSA26 locus for live cell imaging. *Genesis* **49**, 579-590 (2011).
https://doi.org/10.1002/dvg.20753

25 Yoshioka-Kobayashi, K. *et al.* Coupling delay controls synchronized
oscillation in the segmentation clock. *Nature* **580**, 119-123 (2020).
https://doi.org/10.1038/s41586-019-1882-z

26 Pardee, A. B. A restriction point for control of normal animal cell
proliferation. *Proc Natl Acad Sci U S A* **71**, 1286-1290 (1974).
https://doi.org/10.1073/pnas.71.4.1286

27 Tsiairis, C. D. & Aulehla, A. Self-Organization of Embryonic Genetic
Oscillators into Spatiotemporal Wave Patterns. *Cell* **164**, 656-667
(2016). https://doi.org/10.1016/j.cell.2016.01.028

28 Meijer, H. A. *et al.* NOTCH1 S2513 is critical for the regulation of NICD
levels impacting the segmentation clock in hiPSC-derived PSM cells and
somitoids.   *bioRxiv*,   2024.2012.2010.627712   (2024).
https://doi.org/10.1101/2024.12.10.627712

29 Sonnen, K. F. *et al.* Modulation of Phase Shift between Wnt and Notch
Signaling Oscillations Controls Mesoderm Segmentation. *Cell* **172**,
1079-1090 e1012 (2018). https://doi.org/10.1016/j.cell.2018.01.026

30 Aulehla, A. *et al.* Wnt3a plays a major role in the segmentation clock
controlling   somitogenesis.   *Dev Cell*   **4**,   395-406   (2003).
https://doi.org/10.1016/s1534-5807(03)00055-8

31 Greco, T. L. *et al.* Analysis of the vestigial tail mutation demonstrates
that Wnt-3a gene dosage regulates mouse axial development. *Genes &*
„Ä¶ **10**, 313-324 (1996).

32 van Oostrom, M. J., Meijer, W. H. M. & Sonnen, K. F. A Microfluidics
Approach for the Functional Investigation of Signaling Oscillations
Governing   Somitogenesis.   *J Vis Exp* **169**,   1-17   (2021).
https://doi.org/10.3791/62318

33 Masamizu, Y. *et al.* Real-time imaging of the somite segmentation clock:
revelation of unstable oscillators in the individual presomitic mesoderm
cells. *Proc Natl Acad Sci U S A* **103**, 1313-1318 (2006).
https://doi.org/10.1073/pnas.0508658103

34 Morgan, D. O. *The cell cycle : principles of control.* (2007).

35 Pourquie, O. The segmentation clock: converting embryonic time into
spatial   pattern.   *Science*   **301**,   328-330   (2003).
https://doi.org/10.1126/science.1085887

36 Soroldoni, D. *et al.* Genetic oscillations. A Doppler effect in embryonic
pattern   formation.   *Science*   **345**,   222-225   (2014).
https://doi.org/10.1126/science.1253089

37 Cooke, J. & Zeeman, E. C. A clock and wavefront model for control of
the number of repeated structures during animal morphogenesis. *J*

Theor   Biol **58**,   455-476   (1976).   https://doi.org/10.1016/s0022-
5193(76)80131-2

38 Niwa, Y. *et al.* Different types of oscillations in Notch and Fgf signaling
regulate the spatiotemporal periodicity of somitogenesis. *Genes Dev* **25**,
1115-1120 (2011). https://doi.org/10.1101/gad.2035311

39 Dubrulle, J. & Pourquie, O. fgf8 mRNA decay establishes a gradient that
couples axial elongation to patterning in the vertebrate embryo. *Nature*
427, 419-422 (2004). https://doi.org/10.1038/nature02216

40 Dubrulle, J., McGrew, M. J. & Pourquie, O. FGF signaling controls
somite boundary position and regulates segmentation clock control of
spatiotemporal Hox gene activation. *Cell* **106**, 219-232 (2001).
https://doi.org/10.1016/s0092-8674(01)00437-8

41 Mongera, A. *et al.* A fluid-to-solid jamming transition underlies vertebrate
body   axis   elongation.   *Nature*   **561**,   401-405   (2018).
https://doi.org/10.1038/s41586-018-0479-2

42 Naiche, L. A., Holder, N. & Lewandoski, M. FGF4 and FGF8 comprise
the wavefront activity that controls somitogenesis. *Proc Natl Acad Sci U*
S A **108**, 4018-4023 (2011). https://doi.org/10.1073/pnas.1007417108

43 Cambray, N. & Wilson, V. Axial progenitors with extensive potency are
localised to the mouse chordoneural hinge. *Development* **129**, 4855-
4866 (2002). https://doi.org/10.1242/dev.129.20.4855

44 Mathis, L. & Nicolas, J. F. Different clonal dispersion in the rostral and
caudal mouse central nervous system. *Development* **127**, 1277-1290
(2000). https://doi.org/10.1242/dev.127.6.1277

45 Neijts, R., Simmini, S., Giuliani, F., van Rooijen, C. & Deschamps, J.
Region-specific regulation of posterior axial elongation during vertebrate
embryogenesis.   *Dev   Dyn*   **243**,   88-98   (2014).
https://doi.org/10.1002/dvdy.24027

46 Yaman, Y. I. & Ramanathan, S. Controlling human organoid symmetry
breaking reveals signaling gradients drive segmentation clock waves.
Cell   **186**,   513-527   e519   (2023).
https://doi.org/10.1016/j.cell.2022.12.042

47 Thomson, L., Muresan, L. & Steventon, B. The zebrafish presomitic
mesoderm elongates through compaction-extension. *Cells Dev* **168**,
203748 (2021). https://doi.org/10.1016/j.cdev.2021.203748

48 Sabherwal, N. *et al.* Differential phase register of Hes1 oscillations with
mitoses underlies cell-cycle heterogeneity in ER(+) breast cancer cells.
Proc   Natl   Acad   Sci   U   S   A **118**   (2021).
https://doi.org/10.1073/pnas.2113527118

49   Weterings, S. D. C. *et al.* NOTCH-driven oscillations control cell fate
decisions during intestinal homeostasis. *bioRxiv*,
2024.2008.2026.609553 (2024).
https://doi.org/10.1101/2024.08.26.609553

50   Maeda, Y., Isomura, A., Masaki, T. & Kageyama, R. Differential cell-
cycle control by oscillatory versus sustained Hes1 expression via p21.
Cell Rep **42**, 112520 (2023).
https://doi.org/10.1016/j.celrep.2023.112520

51   Zeng, C. *et al.* Evaluation of 5-ethynyl-2'-deoxyuridine staining as a
sensitive and reliable method for studying cell proliferation in the adult
nervous system. *Brain Res* **1319**, 21-32 (2010).
https://doi.org/10.1016/j.brainres.2009.12.092

52   Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**
(2018). https://doi.org/10.1242/dev.165753
1039
1040