Scaling of mouse somitogenesis by coupling of cell cycle to segmentation clock oscillations

Marek J. van Oostrom[1], Yuting I. Li[2,3], Wilke H. M. Meijer[1], Tomas E. J. C. Noordzij[1], Charis Fountas[1,$], Erika Timmers[1], Jeroen Korving[1], Wouter M. Thomas[1], Benjamin D. Simons[2,4,5] and Katharina F. Sonnen[1,*]

[1]Hubrecht Institute-KNAW (Royal Netherlands Academy of Arts and Sciences), University Medical Center Utrecht, Utrecht, The Netherlands

[2]Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK

[3]Institute of Science and Technology Austria (ISTA), 3400 Klosterneuburg, Austria

[4]Gurdon Institute, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QN, UK

[5]Cambridge Stem Cell Institute, Jeffrey Cheah Biomedical Centre, University of Cambridge, Cambridge, CB2 0AW, UK

[$]present address: Division of Gene Regulation, the Netherlands Cancer Institute, Oncode Institute, Plesmanlaan 121, 1066CX Amsterdam, the Netherlands

*Correspondence: k.sonnen@hubrecht.eu

Somitogenesis is the periodic segmentation of a continuously growing tissue in vertebrate embryos[1]. Timing and spacing of somitogenesis is regulated by the segmentation clock, oscillations in Notch, Wnt and FGF signalling in mice, and FGF and Wnt morphogen gradients[2-4]. The size of nascent somites scales with the size of the unsegmented region, the presomitic mesoderm (PSM), over developmental time and upon experimental perturbation of PSM size[5-7]. Here, we addressed how scaling is ensured in the developing mouse embryo. We found that the entire oscillating presomitic mesoderm proliferates. Quantifying the dynamics of signalling and cell cycle progression in single cells, we detected a correlation between the two. Using a

microfluidics-based entrainment approach, we showed that signalling oscillations and cell cycle are coupled, likely via periodic induction of S phase-inducing Cyclins. An updated theoretical somitogenesis model incorporating this coupling suggests that this correlation ensures uniform PSM growth and uniform dilution of morphogen gradients by cell divisions, thereby directly affecting somite formation. Indeed, when blocking cell proliferation, gradient spreading and somite formation as well as somite scaling are impaired. Our data suggests a mechanism by which coupling signalling oscillations to cell proliferation ensures formation of properly sized somites and scaling of somite size.

Main

Somitogenesis is highly reproducible between individual animals. Cells starting in the posterior tip of the embryo tail traverse the PSM bilaterally to the neural tube, until differentiation and somite formation is induced in anterior PSM[1]. Imbalance between growth and segmentation can lead to premature or delayed arrest in somite formation, improperly sized somites or asymmetries between left and right[6-8]. Scaling of somitogenesis, the relative adjustment of somite size to PSM size, occurs during mouse development when PSM size first increases until embryonic day (E)9.5 and then decreases until the somitogenesis terminates[5]. Moreover, experimentally decreasing PSM size results in the formation of smaller somites[6,7]. The potential involvement of cell proliferation in somitogenesis has been a subject of discussion and controversy. Described as the main contributor to PSM growth[9-11], cell cycle inhibition does not interfere with segmentation clock oscillations in 2D *in vitro* culture models of human somitogenesis[12]. In zebrafish embryos, absence of proliferation impairs proper somite formation and has been described as a source of noise for the segmentation clock[5,8,13-15]. A link between segmentation clock and cell division has been suggested[14-16], but not formally shown.

Results

Cell cycle dynamics along the PSM

To understand the role of cell proliferation in somitogenesis, we first
characterized cell cycle dynamics in uncultured mouse embryo tails.
Immunostaining for the cell cycle marker Ki67 implied that all PSM cells were
actively cycling (Fig. 1A). We then used flow cytometry to determine distribution
into the four cell cycle phases: G1, S (when DNA replication occurs), G2 and M
phase (when cell division occurs). We specifically analysed Tbx6-positive PSM
cells and found that most cells were in S phase (60.4%), while fewer were either
in G1 (22.7%) or G2/M phase (16.9%, Fig. 1B, Fig. S1). Previously, we have
found that cell proliferation occurs in the entire PSM[10]. To quantify this precisely
along the anteroposterior (AP) axis, we performed immunostaining using the
mitotic cell marker pHH3 in mouse embryonic tails transversally sectioned from
posterior to anterior (Fig. 1C-E, Fig. S2). We confirmed that cells divided within
the tailbud, PSM and somite region of the tail (Fig. 1E). Furthermore, there was
a slight increase in the percentage of late mitotic cells within the first two forming
somites compared to the PSM.

Next, we quantified the proliferation rate in embryo tails using sequential
pulse labelling with thymidine analogues 5-Ethynyl-2′-deoxyuridine (EdU)
followed by 5-Bromo-2′-deoxyuridine (BrdU) 1 h later and quantified cell cycle
timing in similarly transversally sectioned tails[17,18] (Fig. 1C,F-H, Fig. S3). Cell
cycle   length   was   quantified   using   the   formula $Cell\ cycle\ length =$
$\frac{Total\ cell\ number\ \times\ 1\ h}{EdU^+BrdU^-cell\ number}$. The differentially labelled cell fractions allowed us to
quantify cell cycle and S-phase duration. Cell cycle duration within the tailbud
and PSM was approximately 14 hours (Fig. 1H, Fig. S3D) and S phase duration
10 hours (Fig. S3E-G). With a segmentation clock period of 2.7 hours this
indicates proliferation results in tissue size doubling in around 5 segmentation
cycles.

Within the first two somites, the overall cell cycle length was decreased
relative to the PSM, which is in line with the increase in mitotic cells (Fig. 1E,H).
Interestingly, variability in cell cycle duration was higher in somites, hinting at
tighter cell cycle control in the PSM than in differentiating somites. Thus, cells
divide along the whole AP axis of the PSM, suggesting constant stretching of
the entire PSM. This contrasts with a model in which cells are added in the
posterior PSM, while segmentation occurs in the anterior.

Quantification of cell cycle dynamics in real-time

We subsequently determined real-time cell cycle dynamics in the
growing mouse embryo. To this end, we cultured E10.5 embryo tails expressing
the inducible Fucci2 cell cycle G1 reporter R26R-mCherry-hCdt1(30/120)[19]
(Fucci2-Cdt1) *ex vivo* and quantified reporter expression in single cells over
time. To make this possible, we established single-cell tracking in the growing
embryonic tail by combining mosaic labelling with light-sheet microscopy (Fig.
1I-N, Fig. S4A-C). We combined the inducible Fucci2 reporter[19] with the
inducible nuclear marker R26R-iRFPnucmem[20] (iRFPnucmem) and induced
sparse labelling with injections of low concentrations of Tamoxifen in the
mother. Cell cycle duration was on average 13.6 hr, corresponding to the cell
cycle timing *in vivo* (Fig. 1H), and cells spent on average 4.4 hr in G1 phase
(Fig. 1N). Notably, we did not detect changes in cell density despite cell
divisions over 24 hr of real-time imaging (Fig. S4D,E), indicating that cells grow
and divide during tissue growth.

To determine if there was any spatiotemporal regulation of cell cycle
progression in the PSM, we analysed the spatial organization of cells double-
positive for both the Fucci reporters mKuzabiraOrange2-Cdt1 (Fucci1-Cdt1)
and mAzabiGreen-hGem(1/110) (Fucci1-Gem), which specifically marks cells
at the G1-S transition[21]. Notably, the spatial pattern of double-positive cells
differed between independent embryo tails with some having no double-
positive cells in the PSM and others having stripes at different locations
throughout the PSM (Fig. 1O, Fig. S4F). This behaviour is analogous to fixed
images of cyclic segmentation clock genes that show different expression
patterns in different embryos[22,23]. This implies that there might be a
spatiotemporal organization of cell cycle entry in the PSM.

Segmentation clock and cell cycle are correlated

To investigate if there was a correlation between cell proliferation and
segmentation clock oscillations, we simultaneously quantified cell cycle
progression using the inducible nuclear marker R26R-H2B-mCherry[24] and the
segmentation clock using the reporter Achilles-Hes7[25] (Fig. 2A-D, Fig. S5).
Based on the nuclear marker, we tracked single cells, aligned tracks to the time

of cell division and analysed Hes7 expression (Fig. 2C). Immediately after cell

division, daughter cells of a given mother tended to oscillate more in-phase than

several hours after division (Fig. S5C). Moreover, we did not detect a

preferential division axis along the anteroposterior axis (Fig. S5D). Interestingly,

we found that the Hes7 oscillation phase before division was not randomly

distributed. At the point of cell division, there was a sudden drop in Hes7 signal

presumably due to nuclear envelope breakdown. After division, individual cells

immediately showed Hes7 oscillations again (Fig. S5A), but aligned tracks

showed no synchronized behaviour for approximately 4 hours, when cells

started to resynchronize again (Fig. 2C,D, Fig. S5E). This timeframe of non-

synchronization roughly corresponds to the time cells spent in G1 phase of the

cell cycle (Fig. 1N).

The G1 restriction point is a point of no return, after which the cell cycle

can run autonomously, even though later restriction points can prevent cell

cycle progression to ensure fidelity of cell division[26]. We therefore addressed

whether segmentation clock oscillations were correlated with G1-S transition.

To this end, we combined Achilles-Hes7 with Fucci2-Cdt1 and iRFPnucmem

(Fig. 2E-G, Fig. S6A-D) and quantified expression in single cells of *ex vivo*

cultured embryo tails after induction of mosaic labelling. When analysing the

instantaneous period of Hes7 oscillations along the posterior-anterior PSM

axis, we detected an increase in period towards the anterior (Fig. S7A),

consistent with an oscillation period gradient[27]. We then separated cell tracks

into their respective cell cycle phase cells, based on Fucci2-Cdt1 reporter

expression. In posterior PSM, cells in G1 had a higher oscillation period than

cells in other cell cycle phases (Fig. S7B,C). Moreover, the period gradient was

less clear in G1 than in S/G2/M phase cells. Also, when directly comparing all

cells in G1 to all cells in S/G2/M phase, independent of the location in the PSM,

we detected a slightly higher period in S/G2/M phase cells (Fig. S7D). This

indicates that oscillation parameters change with anteroposterior position in the

PSM and cell cycle phase suggesting an effect of cell proliferation on the

segmentation clock[14,28].

We then addressed a potential correlation between Hes7 oscillations

and G1-S transition. When aligning all tracks to the point of G1-S transition,

defined as the peak in Fucci2-Cdt1 signal (Fig. 1L), we did not observe any

preferred phase of Hes7 oscillations (Fig. 2G, Fig. S6E,F). In mice, the segmentation clock consists of oscillations in Notch, Wnt and FGF signalling. We have previously shown that the phase-relationship between Wnt and Notch signalling changed along the posterior-anterior axis of the PSM from out-of-phase in the posterior to in-phase in the anterior[29], corresponding to a 1π shift in phase-relationship. If a corresponding shift in phase-relationship was present for cell cycle and Hes7 oscillations, any correlation would be masked when analysing the average signal. We therefore separated tracks into cells residing in the posterior and anterior PSM and analysed the correlation of Fucci2-Cdt1 and Hes7 dynamics. Interestingly, we found that in both regions proliferation and the segmentation clock were correlated (Fig. 2G, Fig. S6E,F). In posterior PSM, the phase of Hes7 oscillations was at $\sim 1.3\pi$ at G1-S transition, while it was at $\sim 0.4\pi$ in anterior PSM. This corresponded to a change in phase-relationship between Notch oscillations and cell cycle of $\sim 0.9\pi$ between posterior and anterior PSM. This indirectly indicated that cell cycle progression correlated with Wnt-signalling oscillations in the PSM.

While the cell cycle in a single cell takes approx. 14 hr, at the population level we observed faster dynamics in the Fucci2-Cdt1 reporter: When quantifying small regions containing approximately 10-15 cells, we observed oscillations in Hes7 and G1-S transition with a period of roughly 2.5 hr (Fig. 2H,I, Movie S3). This suggested that in this region one cell after the other showed high expression of the Fucci2-Cdt1 reporter. This in turn implied that neighbouring cells were shifted in their G1-S transition by n x segmentation clock period relative to their neighbours (Fig. 2J).

Thus, cell cycle progression is correlated with the segmentation clock and has a constant phase-relationship with Wnt-signalling oscillations along the AP axis of the PSM.

Segmentation clock and cell proliferation are coupled

We next sought to determine whether cell proliferation is merely correlated to the segmentation clock or whether there is a functional link between the two. Continuous inhibition of Wnt signalling by genetic or chemical perturbation leads to impaired PSM growth and a rapid regression of the PSM due to accelerated differentiation and loss of the morphogen gradient[30,31] (Fig.

S8). Such continuous perturbations are therefore not ideally suited to reveal a

functional relationship between two oscillators. Therefore, we needed a system

to subtly modulate oscillations. To achieve this, we used our previously

established microfluidic system to entrain oscillations to external pulses of

pathway modulation[29,32]. We cultured embryonic tail tips in Fibronectin-coated

microfluidic chips, so that the embryonic tissue spreads in a "quasi-monolayer"

and the embryonic axis is re-organized from posterior-anterior to centre-

periphery[6,29]. This arrangement in combination with mosaic labelling allows for

single-cell tracking even in standard confocal microscopes[20]. Using embryos

expressing Achilles-Hes7, inducible Fucci2-Cdt1 and iRFPnucmem[19,25,33], we

then addressed whether there was a functional link between proliferation and

the segmentation clock. Importantly, the correlation between Hes7 oscillations

and G1-S transition at population level was maintained in these cultures (Fig.

S9, Movie S4).

We modulated the segmentation clock by periodically perturbing Wnt

signalling using the GSK3 inhibitor Chiron (CHIR99021) (Fig. 3A, Movies S5-

S9). Our previous experiments had revealed that Chiron efficiently entrains

Wnt-signalling oscillations, which also affects Notch-signalling oscillations of

the segmentation clock[29]. Indeed, we confirmed that Chiron pulses entrained

the segmentation clock, using Achilles-Hes7 as a reporter (Fig. 3B-E).

Subsequently, we analysed when during one signalling oscillation cycle the

Fucci2-Cdt1 reporter peaked, corresponding to G1-S transition, and found that

G1-S transition preferentially occurred with the drug pulses (Fig. 3F-H, Fig.

S10). This is consistent with dynamics of the Wnt-signalling reporter Axin2T2A

that also peaked with the drug pulses[29]. In contrast, DMSO pulses led to a

random distribution of Fucci2-Cdt1 peaks during the oscillation cycle. This

indicates that modulating segmentation clock oscillations via targeted Wnt

signalling modulation entrains cell cycle oscillations.

All signalling pathways of the segmentation clock are known mitogens

promoting cells to enter the cell cycle. Recently, myc and Cyclin D, genes well

studied for their role in cell cycle progression and G1-S transition[26], have been

identified as potential cyclic genes at the mRNA level[3]. Indeed, using

hybridization chain reaction (HCR), we found Cyclin D1 to be expressed in the

PSM with a dynamic pattern overlapping with the Wnt-signalling target gene

Axin2 (Fig. 3I, Fig. S11A). Since cyclins are post-translationally regulated by
modulating protein stability, we stained for Cyclin D1 protein. This also showed
a corresponding staining pattern to Axin2 (Fig. 3J, Fig. S11B). To determine a
functional link, we perturbed Wnt signalling and analysed the effect on Cyclin
D expression. Since Wnt inhibition leads to an immediate loss of PSM due to
differentiation, we instead activated Wnt signalling using Chiron. This led to an
increase in Cyclin D expression in the entire PSM (Fig. 3K,L).

Thus, the segmentation clock and cell proliferation are coupled,
presumably via periodic induction of cell cycle regulators.

An extended somitogenesis model suggests that cell cycle-
segmentation clock coupling ensures proper somite formation

To determine whether the coupling of cell proliferation to signalling
oscillations had a functional role in maintaining periodic somite formation, we
turned to a conceptual modelling-based approach, extending on previous
models of somitogenesis[7,34-36]. Specifically, we modelled the dynamics of cells
in the PSM as a line of coupled Kuramoto-type phase oscillators and made the
following assumptions based on our findings and available literature: (1) At a
certain phase between Wnt- and Notch-signalling oscillations and a lower
threshold of the FGF morphogen gradient, somite formation occurs[29,37,38]. (2)
All PSM cells have a certain probability to divide, and the cell cycle period takes
values in the range 13.6 ± 1.3 hr (Fig. 1N). (3) Since FGF mRNA is known to
only be produced in the posterior tailbud[38,39], the FGF morphogen gradient is
diluted by mRNA degradation as well as cell division, when the mRNA present
in a cell is partitioned equally into the two daughter cells. (4) Minimal
rearrangement of cells in the one-dimensional model of the PSM was allowed.
This assumption was consistent with quantifications of cell migration and
rearrangement in the PSM cells tracked using light-sheet imaging (Fig. 4A-D,
Fig. S12A,B). While cells in the tailbud moved constantly (Movie S1), the
movement of cells in the PSM were much more restricted (0.1 µm/ 15 min) with
cells moving relative position by less than one cell diameter over the period of
the segmentation clock cycle (Fig. 4C). (5) We imposed the condition that cell
division occurs 1.25 ± hr after Wnt oscillations peak (Fig 2D). Based on these
assumptions, numerical simulations of this minimal one-dimensional model

recapitulated features of the continuously growing PSM field, with periodic
formation of somites (Supplemental discussion).

Based on our finding that cell proliferation occurs along the entire PSM,
we first questioned what effect the coupling of cell cycle entry with the
segmentation clock has. To this end, we considered a line of 50 cell grid points,
each containing 10 cells, based on the number of cells in a line from posterior
to anterior PSM and minimal cell exchange with neighbours (Fig. S12C and Fig.
4A-C), and simulated tissue growth and somite formation (Fig. 4E, Fig. S13).
Assuming that cell divisions occur randomly, decoupled from the segmentation
clock, with a period of 13.6 ± 1.3 hr, numerical simulations revealed large
variations in somite size (Fig. 4E,F, CV = 0.31). Moreover, uncoupled growth
led to a non-uniform dilution of the FGF morphogen gradient (Fig. 4E, Fig.
S13A). In contrast, when the entry into cell cycle was coupled to the phase of
the segmentation clock, tissue growth and somite formation were more uniform
(Fig. 4G,H, Fig. S13,B CV = 0.20). This indicates that the coupling of cell
proliferation to the segmentation clock leads to uniform growth and uniform
dilution of the morphogen gradient regulating the spacing of somite formation.
288

Perturbing cell proliferation interferes with morphogen gradient
dilution and scaling of somite formation

To understand how cell proliferation affected the segmentation process,
we inhibited cell proliferation in 2D *ex vivo* cultures[6] of embryos expressing
Fucci2-Cdt1, Achilles-Hes7 and iRFPnucmem using different cell cycle
inhibitors. Inhibition resulted in the expected effects on Fucci2-Cdt1 levels and
an accumulation of non-dividing cells (Movie S10). Upon inhibition,
segmentation clock oscillations were maintained (Fig. S14A,B), as shown
previously[12]. This indicates that cell cycle progression is not a necessary driver
of segmentation clock oscillations[12].

To challenge the predictions of the model on the nature of somite
formation in the presence of cell proliferation, we inhibited proliferation in
growing embryo tails using Aphidicolin as strong continuous inhibitor (Fig. 5,
Movies S11, S12). This led to a decrease in cell number, as expected, but tissue
size remained largely unaltered due to continued cell growth, leading to a
decrease in overall cell density (Fig. S14C-G). We then quantified morphogen

gradients using as a readout Brachyury expression, which is directly affected by FGF and Wnt3a gradients[39,40]. Consistent with the model, the Brachyury gradient was expanded in the unsegmented PSM upon cell cycle inhibition (Fig. 5A-C). Similarly, the expression pattern of FGF8 and Axin2 mRNA were expanded and less graded than in controls (Fig. S14H-K).

Consistent with the expansion of morphogen gradients[39], the size of the oscillating field, a measure of the extent of differentiation, decreased less than in control tails (Fig. 5D,E, Fig. S14L). Furthermore, we found that the phase wave velocity did not decrease to the same extent as in control tissues but remained higher, despite the decrease in cell number relative to control tissue (Fig. S14L-M). This contrasted with unperturbed tissues, in which the wave velocity slows with decreasing size and cell number[6]. This implies that the relative phase-shift between oscillations in neighbouring cells decreases in the absence of cell division.

Finally, we analysed the effect on somite formation. Aphidicolin treatment led to the formation of disorganized somites with undiscernible boundaries (Fig. 5F)[8]. The size of forming somites relative to PSM size was affected with the ratio changing from 0.172 to 0.109, thereby indicating altered scaling of somite formation (Fig. 5G-J). We note that since (a) cell density between the PSM and somites (Fig. S14D-G) and (b) cell motility (Fig. 5K) were comparable between control and cell cycle inhibition, this cannot account for the effect on scaling.

Thus, cell cycle inhibition impairs somite formation and scaling of somitogenesis.

Discussion

Here, we addressed the role of cell proliferation in somitogenesis, showing a tight correlation between cell cycle entry and the π of the segmentation clock. Further, we presented a mechanism by which the coupling of cell division to the segmentation clock may ensure uniform PSM growth and dilution of regulatory molecules, thereby modulating the scaling and uniformity of somite formation.

In contrast to studies that support a posterior growth zone[41-43], we found

that cell proliferation was uniformly spread over the posterior and anterior

PSM[5,10]. Furthermore, cell proliferation was coupled to the segmentation clock,

likely via Wnt-induced expression of pro-proliferative factors such as Cyclin D[3],

which ensures uniform distribution of proliferating cells over the PSM. With a

proliferation rate of approximately 14 hr compared to a somitogenesis period of

2.7 hr, cell proliferation is a key component of mouse somitogenesis. Upon cell

cycle inhibition, cell division was abolished while tissue and cell growth

continued, indicating that cell division in the PSM has a negligible effect on

tissue growth within this timescale.

Morphogen gradients are generated by mRNA and protein production in

the posterior tip and levels slowly decrease as cells move down the posterior-

anterior axis of the PSM[38]. Thus, morphogen gradients are modified by mRNA

(and protein) degradation as well as cell division, which lead to a continuous

decrease in morphogen concentration and nuclear effector proteins in cells.

Indeed, inhibiting cell divisions expanded the morphogen gradients. In

agreement with the clock-and-wavefront model[36], regression of the oscillating

field was also reduced upon cell cycle inhibition. Moreover, signalling wave

velocity was increased and somite formation impaired. This suggests that the

principal role of cell division in the PSM may lie not in its contribution to tissue

growth but rather in shaping morphogen gradients for proper somite formation.

This is consistent with a model (Fig. 5L) in which cell divisions are

uniformly distributed over the tissue by coupling to the segmentation clock. This

leads to an increase in cell number uniformly distributed over the PSM, which

leads to (1) a uniform expansion of the oscillating field and (2) a uniform dilution

of morphogen mRNA and protein as well as downstream gradients. The

gradient affects wave dynamics[44] and both in turn regulate somite

formation[37,39]. Supported by mathematical modelling, this suggests that the

coupling of cell divisions to the segmentation clock ensures formation of

correctly sized somites. Scaling of segment size to PSM size has previously

been correlated with scaling of wave dynamics[6] or the FGF gradient[7] in mouse

and zebrafish, respectively, both of which we found to be affected by cell cycle

inhibition. Thus, this constitutes a mechanism by which local cell-intrinsic

regulations, i.e., whether to divide or not, is translated into tissue-wide
regulation of morphogenesis to ensure proper somite formation.

How our findings translate to other species remains to be addressed. In
zebrafish, continued tissue growth, impaired somite formation and decreased
cell density are observed in the PSM of *emi1* mutants, which stop proliferation
during gastrulation[8]. Moreover, segmentation clock oscillations correlate with
cell divisions in zebrafish[15], even though cell proliferation is thought to play a
minor role for tissue growth[45]. This could explain why in zebrafish segmentation
proceeds more quickly than growth and oscillation regression leading to a rapid
reduction of tissue size and a decrease in wave amplitude from 4π to 2π over
developmental time[35]. In human somitogenesis, a correlation between the
timepoint of cell division and Hes7 oscillations has not so far been observed[12].
To clarify the role of cell proliferation in human cells, cell cycle reporters specific
for the G1-S transition should be quantified in posterior or anterior PSM cells
relative to Hes7 oscillations.

Signalling oscillations have been observed in other tissue types and
developmental contexts. Correlations between such oscillations and cell
division have been described in several tissues (e.g., [46]). In *in vitro* cultures of
the small intestine, we recently identified Hes1 oscillations that also correlate
with cell divisions[47]. In neural stem cells interactions of Hes1 oscillations with
the cell cycle inhibitor p21 have been described[48]. Whether there is active
coupling in all of these examples and whether this has similar effects on growth,
differentiation and morphogenesis at the tissue level, as we identified here for
somitogenesis, are exciting questions for future research.

Figure Legends

Figure 1 Cell cycle dynamics along the entire E10.5 presomitic mesoderm.

A-G Quantification of cell cycle dynamics in uncultured tails. A 5 µm section of

E10.5 tail with immunostaining for Ki67 and counterstaining with DAPI. **B** Flow

cytometry analysis of embryonic tail cells stained for Tbx6 and DNA using

Hoechst. DNA content in Tbx6-postive PSM cells. N=3 experiments. Raw data

shown in Fig. S4. **C-E** Quantification of late mitotic cells in embryonic tails. N=5.

C Embryonic tails were sectioned into 5 µm sections. Immunostaining against

phosphorylated-histone H3 (PHH3) and counterstaining with DAPI were

performed. For the analysis, tissue was divided into tailbud (posterior to neural

tube), PSM and somite regions. **D** Representative image of a section stained

against PHH3 and with DAPI (same as in Fig. S2A). **E** Quantification of cells in

late mitosis (pro-metaphase to anaphase) in the regions indicated in **B**. **F-H**

Quantification of cell cycle dynamics by EdU-BrdU double-labelling. N=3. **F**

Scheme of experimental setup. Embryos were dissected and tails sectioned akin to **B** before EdU/BrdU staining. **G** Representative image of a section stained for EdU, BrdU and nuclei (same as in Fig. S3A). **H** Quantification of cell cycle length in the regions indicated in **B**. N = 3. **I-N** Quantification of cell cycle dynamics in *ex vivo* cultured tails. **I** Pregnant mice were injected with low doses of Tamoxifen on E9.5 for sparse nuclear labelling of cells. Embryos were dissected after 24h of incubation. **J** Embryonic tails were dissected and imaged with an inverted light-sheet microscope. **K** Brightfield image of tail within agarose groove. **L** Representative image of embryonic tails with sparse-labelling of nuclear marker iRFPnucmem. **M** Representative timeseries data of the dynamics of the G1 marker Fucci2-Cdt1 in a single cell tracked from anaphase to anaphase. **N** Quantification of cell cycle length and duration that cells are positive for the G1 marker Fucci2-Cdt1. **O** Patterns of G1-S transition in uncultured E10.5 embryonic tails (same as in Fig. S4F). Cells double-positive for Fucci1-Cdt1 and Fucci1-Gem were determined in PSM of embryonic tails. Tails were ordered into three phases according to their staining pattern: Phase 1: no double-positive cells in PSM; Phase 2: stripe of double-positive cells in PSM; Phase 3: double-positive cells distributed over PSM. Number in brackets indicates number of tails in which specific staining pattern was observed. Brightfield (*left panel*), merge of Fucci1-Cdt1 and Fucci1-Gem (*middle panel*) and cells double-positive for Fucci1-Cdt1 and Fucci1-Gem (*right panel*) depicted. N=13. Scale bar 100 μm.

Figure 2 Cell cycle dynamics correlate with segmentation clock. A-D Correlation of Hes7 segmentation clock to cell division. **A** Experimental approach for sparse labelling of nuclei with H2B-mCherry combined with Achilles-Hes7. **B** Representative snapshots of timeseries data. Arrowhead points at one cell tracked over time. **C** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal of cells aligned to timepoint of cell division. **D** Phase of Achilles-Hes7 oscillations at timepoint 1, as indicated in **C**, (*left panel*) and at timepoint 2, as indicated in **C**, (*right panel*). **E-G** Correlation of Hes7 segmentation clock to G1-S transition. **E** Experimental approach for sparse labelling of nuclei with iRFPnucmem and Fucci2-Cdt1 combined with Achilles-Hes7. **F** Representative snapshots of timeseries data. Arrowhead points at one cell tracked over time. **G** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal of cells aligned to timepoint of G1-S transition. Left panel: all cells; middle panel: cells in posterior PSM; right panel: cells in anterior PSM. **H** Phase of Achilles-Hes7 oscillations at G1-S transition. Left panel: all cells; middle panel: cells in posterior PSM; right panel: cells in anterior PSM. **I** Maximum Intensity projection of E10.5 embryonic tail (purple: Achilles-Hes7, cyan: Fucci2-Cdt1). **J** Quantification of Achilles-Hes7 and Fucci2-Cdt1 in regions of interest (ROIs) as indicated in **I**. **K** Scheme

455 illustrating correlation of cell cycle (duration 13.6 hr) to segmentation clock

456 oscillations (2.7 hr).

Figure 3 Coupling of cell proliferation to segmentation clock oscillations. A-H Microfluidics to test functional link. **A** Experimental approach: Microfluidic system to modulate segmentation clock oscillations in *ex vivo* tailbud cultures. Analysis of segmentation clock and cell proliferation. **B-E** Analysis of segmentation clock: **B,D** Mean (blue line) and standard deviation (light blue shading) of Achilles-Hes7 signal in anterior PSM upon treatment with DMSO (**B**) and 5 µM Chiron (**D**). **C,E** Phase of Hes7 oscillations plotted against phase of external drug pulses for treatment with DMSO (**C**) and 5 µM Chiron (**E**). **F** Snapshots of representative fluorescence real-time imaging data

highlighting tracking of a single cell (arrowhead). **G** Representative timeseries data showing Fucci2-Cdt1 reporter intensity of a single cell during microfluidic experiment. Grey shading indicates external drug pulse. **H** Phase of external drug pulses, at which G1-S transition occurs in DMSO- (*left panel*) or Chiron-treated samples (*right panel*). **I** Staining for mRNA levels of Axin2, Cyclin D1 and Hes7 in E10.5 embryonic tails using hybridization chain reaction (HCR) (same as in Fig. S11A). **J** Immunostaining against Cyclin D1 in Axin2T2AVenus-expressing E10.5 embryonic tails (same as in Fig. S11B). **K** Treatment of E10.5 embryonic tails with DMSO or Chiron (10 µM). HCR for Axin2 (upper panels) or Cyclin D1 (lower panels). Left panels: representative images. Right panels: Quantification of Axin2 and Cyclin D1 in regions of interest as indicated in representative images. * < 0.05; ** < 0.01.

Figure 4 Uniform PSM growth and somite formation by coupling of cell proliferation to segmentation clock. A-D E10.5 embryonic tails were cultured ex vivo and fluorescence real-time imaging was performed (corresponds to data from Figure 1,2). **A** Snapshot of an embryonic tail imaged in a light-sheet microscope overlaid with tracks of single cells. **B** Plots of single cell tracks shown in A (*left panel*). Colours correspond to different cells. Each dot represents location of a cell at specific timepoint (interval between timepoints 15 min). Whole-tissue movement was computationally removed to reveal movement of single cells relative to their surrounding (*right panel*). **C**

Histogram of the amount of cell movement given in % of cell diameter (10 µm). N = 9 tissues n = 169 tracks. **D** Velocity of cells moving relative to the posterior tip (*top panel*). This indicates stretching of the PSM, as cells further away from the posterior have a higher velocity (*bottom panel*). Arrows indicate direction of movement. N = 6 tissues n = 106 tracks. * < 0.05; ** < 0.01; *** < 0.001; **** <0.0001. **E-H** Simulation of somitogenesis with random growth (E,F) or growth coupled to segmentation clock (G,H). **E,G** Example plots for simulations of signalling waves (*left panels*) as well as FGF gradient (gray shading) and somite formation (blue line) (*right panels*). **F,H** Histogram of somite lengths in 100 simulations. Coefficient of variation CV(random growth) = 0.31. CV(coupled growth) = 0.20.

**Figure 5 Coupling of cell cycle to segmentation clock affects
scaling of somite formation.** E10.5 embryonic tails were cultured *ex vivo* and
treated with DMSO or the cell cycle inhibitor Aphidicolin. **A-C** After 23 hr of
incubation, tails were fixed and stained for Brachyury and counterstained with
DAPI. Scale bar 100 µm. **A** Representative images. **B** Quantification of the
Brachyury signal along the AP axis of the PSM. Signal was normalized between
0 and 1. n(DMSO) = 6, n(Aphidicolin) = 11. **C** Quantification of the length of
Brachyury gradient (Line: mean, shading: standard deviation). Intensity
normalized between 0 and 1. * < 0.05; *** < 0.001; **** <0.0001. **D,E**
Fluorescence real-time imaging of embryonic tails expressing Achilles-Hes7
was performed. **D** Representative kymographs of control and Aphidicolin-

treated samples. **E** Quantification of PSM length by determining the size of the oscillating field (Line: mean, shading: standard deviation). Length normalized to the PSM length at experiment start. n(DMSO) = 9, n(Aphidicolin) = 11? **F** After 23 hr of incubation, tails were fixed and HCR against Hes7 and Uncx4.1 as well as counterstaining with DAPI were performed. Representative images are shown (same as in Fig. S14J). **G-J** After 23 hr of incubation, tails were fixed and stained with DAPI. **G** Representative images are shown. **H-J** Quantification of PSM length from tail tip to last formed somite boundary (**H**), somite length (**I**) and the ratio of somite length to PSM length as a measure of scaling (**J**). **K** Quantification of cell motility in fluorescence real-time imaging data of embryonic tails expressing Achilles-Hes7 (data corresponds to D,E). **L** Schematic model of how coupling of cell proliferation to segmentation clock affects somitogenesis.

Methods

<u>Animals and Housing</u>

All experimental procedures were conducted in accordance with institutional guidelines and approved by the Ethical Review Board of the Hubrecht Institute. Animal protocols were reviewed and authorized by the Animal Experimentation Committee (DEC) of the Royal Netherlands Academy of Arts and Sciences. Mice were bred and housed under standard conditions at the Hubrecht animal facility, with a controlled environment maintained at 50–60% humidity, 22–23°C temperature, and a 12-hour light/dark cycle (inverted). Mice had ad libitum access to water and standard chow throughout the study.

<u>Mouse Strains</u>

All transgenic mouse lines used in this study have been previously described. The Achilles-Hes7 line was generously provided by the Kageyama laboratory[25]. Fucci [21], The R26R-H2B-mCherry, and R26R-mCherry-hCdt1(30/120)[24] lines were obtained from the RIKEN Center for Biosystems Dynamics Research and the RIKEN BioResource Research Center (accession numbers CDB0204K, RBRC02706, and CDB0229K, respectively) via the RIKEN Mouse Repository. The Tg(T-cre/ERT2)1Lwd/J ("T-CreERT2") line was sourced from The Jackson Laboratory (accession number #025520). The R26R-iRFPnucmem[20] and Axin-T2A-Venus[29] lines were provided by the Aulehla laboratory. All wild-type mice used in this study were F1 hybrids (BALB/cJ × C57BL/6J).

<u>Animal and Embryo Work</u>
<u>EdU/BrdU Incorporation</u>

Unless otherwise stated, female mice were sacrificed at 10.5 days post-coitum (dpc) and embryos were subsequently dissected. For EdU and BrdU incorporation experiments, pregnant females at 10.5 dpc were injected with 3.3 mg EdU in PBS, followed one hour later by an equimolar injection of 4.0 mg BrdU in PBS[49]. Mice were sacrificed 30 minutes after BrdU injection. Embryos were dissected in PBS supplemented with 1% Bovine Serum Albumin (Biowest, P6154) for tail extraction.

<u>Sparse Labelling</u>

For sparse labelling experiments, pregnant females were

intraperitoneally injected with tamoxifen (Sigma-Aldrich, T5648, dissolved in

corn oil) on embryonic day E9.5. To label with H2B-mCherry, 1 mg of tamoxifen

was administered, whereas for labelling with iRFPnucmem and Fucci reporters,

0.1 mg of tamoxifen was used.

<u>Flow Cytometry</u>

For flow cytometry analysis, E10.5 mouse tails were extracted and cut

into smaller pieces using a surgical scalpel to facilitate single-cell dissociation.

Tail fragments were washed three times in PBS0 in Eppendorf tubes, followed

by a 15-minute incubation with 400 μl of TrypLE (Thermo Fisher, 12604013).

After incubation, the tissue was passed through a 30G needle to generate a

single-cell suspension. The suspension was then washed twice in PBS0

supplemented with 1% Bovine Serum Albumin (Biowest, P6154) by

centrifugation at 500 g for 5 minutes.

To fix the cells, the pellet was resuspended in 4% paraformaldehyde

(PFA) and incubated at room temperature for 15 minutes. After two additional

washes in PBS with 1% BSA, cells were stored at 4°C until further staining. For

immunofluorescent staining, the single-cell pellet was resuspended in 3% BSA

PBS with 0.25% Triton X-100 and incubated overnight at 4°C with anti-Tbx6

rabbit primary antibody (Abcam, ab38883) at a dilution of 1:1000. After three

washes in 3% BSA PBS, the cells were incubated with secondary donkey anti-

rabbit Alexa Fluor 488 antibody (Thermo Fisher, A-21206) at a dilution of

1:1000 for 2 hours at room temperature.

Following a further two washes, the single-cell suspension was

resuspended in 500 μl of PBS supplemented with Hoechst 34580 (1 μg/ml) and

passed through a 20 μm mesh. Flow cytometry was performed on a CytoFLEX

S Flow Cytometer, and the Tbx6-positive cell population was selected for cell

cycle analysis using FlowJo software.

<u>Culture of Live Embryonic Tissue</u>

Culture of whole mount and spread-out tail tip tissues was performed as

previously described[29]. For perturbation experiments on whole mount and

spread-out tail tips, the following compounds were used: CHIR99021 (Chiron,

Sigma-Aldrich, SML1046) at 10 µM, IWP-2 (Sigma-Aldrich, I0536) at 1 µM,

Aphidicolin (ThermoFisher, J60236.MCR) at 1.6 µg/mL, Thymidine (Sigma-

Aldrich, T9250) at 2 mM, Palbociclib (Selleckchem, PD-0332991) at 1 µM, and

RO-3306 (Sigma-Aldrich, 217699) at 10 µM.

Microfluidics Experiment

The full protocol for performing the microfluidics experiment has been

described elsewhere[29,32]. For entrainment of Achilles-Hes7 and Fucci2-Cdt1

reporter lines, a pulsing scheme was employed consisting of 100 minutes of

medium followed by 30 minutes of medium containing CHIR99021 (Chiron,

Sigma-Aldrich, SML1046) at 5 µM or DMSO.

Confocal Live Imaging

Confocal live imaging was performed using a Leica TCS SP8 confocal

microscope equipped with a HC PL APO CS2 20×/0.75 DRY objective. For cell

tracking experiments, images were acquired every 10 minutes with a z-step of

5 µm, a stack depth of 50 µm, and a z-resolution of 4 µm. For non-tracking

imaging, images were taken every 10 minutes with a z-step of 10 µm, a stack

depth of 100–150 µm, and a z-resolution of 9 µm. Excitation of Achilles,

mCherry, or iRFP signals was performed using OPSL lasers at 514 nm, 568

nm, or 638 nm, respectively. Movies were recorded at a resolution of 512 × 512

pixels (pixel size 1.517 µm).

Light-sheet Live Imaging

Live imaging of whole mount embryo tails for single-cell analysis was

performed using a Viventis light sheet microscope (Viventis Microscopy, #LS1

Live). Imaging was conducted in a humidified atmosphere maintained at 37°C,

with 5% CO_2 and 60% O_2. For fluorescence excitation, lasers with wavelengths

of 514 nm, 568 nm, and 687 nm were used for Achilles, mCherry, and iRFP,

respectively. Imaging was carried out with a 25× objective, using a laser beam

of 3.3 µm and acquiring a z-stack of 60 steps, each with a 3 µm increment.

Images were captured at a resolution of 2048 × 2048 pixels every 15 minutes.

To prevent whole mount tissues from moving out of the field of view while still

allowing unperturbed growth, a 2% agarose gel in PBS was cast around a

custom-made insert in the sample holder. This insert created four spaces (1.5

mm × 0.5 mm) for the individual tissues to be loaded.

Paraffin Sectioning of Tissues and Immunofluorescence

Tissues fixed overnight in 4% paraformaldehyde (PFA) at 4°C were

stored in 70% ethanol at 4°C for up to one week prior to embedding. Embryo

tails were then subjected to Xylene treatment, embedded in paraffin in groups

of three, and sectioned transversally into 5 μm thick slices.

For staining, the slides were sequentially immersed in Xylene, followed

by 100%, 95%, 70%, and 50% ethanol, and rinsed with running cold tap water

for 3 minutes each. Antigen retrieval was performed by heating the slides in a

citrate buffer (8.2 mM sodium citrate and 1.8 mM citric acid, pH 6). Slides were

blocked and permeabilized using 1% Bovine Serum Albumin (BSA, Biowest,

P6154) in 0.5% Triton X-100 PBS. Mitotic cells were stained with anti-phospho-

histone H3 (Sigma-Aldrich, 05-806) at a 1:1000 dilution in 1% BSA PBS, and

incubation was carried out overnight at 4°C. Secondary staining was performed

using Donkey anti-Mouse Alexa Fluor 488 (Thermo Fisher, A-21202) at a

1:1000 dilution, supplemented with DAPI (5 μg/mL, Sigma) in 1% BSA PBS,

and incubation was carried out overnight at 4°C. After staining, slides were

mounted with Vectashield (H1200NB, Vector Laboratories). Images were

acquired using a Leica TCS SP8 confocal microscope with an HC PL APO CS2

20×/0.75 DRY objective and OPSL lasers at 365 nm and 488 nm for excitation.

For BrdU and EdU staining, two additional steps were incorporated. Prior

to blocking and after antigen retrieval, slides were incubated at room

temperature in 2 M HCl for 10 minutes. EdU was then visualized using the Alexa

Fluor™ 647 dye from the Click-iT™ EdU Cell Proliferation Kit for Imaging

(Thermo Fisher, C10340). BrdU was detected using an anti-BrdU antibody

(Thermo Fisher, B35128) at a 1:500 dilution, followed by secondary staining

with Donkey anti-Mouse Alexa Fluor 568 (Thermo Fisher, A10037).

Whole Mount Tissue Antibody Staining

Prior to immunostaining, samples were fixed overnight in 4%

formaldehyde at 4°C and stored in PBS at 4°C. Primary antibodies used for

staining included anti-Ki67 (Abcam, ab16667) at 1:1000, anti-T (human/mouse)

(R&D Systems, AF2085-SP) at 1:1000, and anti-Cyclin D1 (Cell Signaling,
2978) at 1:500. Secondary antibodies included Donkey anti-Goat Alexa Fluor
488 (Thermo Fisher, A-11055) at 1:1000 and Donkey anti-Rabbit Alexa Fluor
568 (Thermo Fisher, A10037) at 1:1000.

Immunostaining was carried out by incubating the samples with primary
and secondary antibodies overnight in PBST (0.1% Triton X-100) containing
1% Bovine Serum Albumin (BSA, Biowest, P6154). The secondary antibody
mixture was supplemented with DAPI (Sigma, 5 µg/mL) for nuclear staining.
For imaging, a z-stack consisting of 10-12 planes with a 10 µm distance
between each plane was acquired at a resolution of 1024 × 1024 pixels (pixel
size 0.758 µm) using a Leica TCS SP8 MP confocal microscope with a HC PL
APO CS2 20×/0.75 DRY objective. Excitation was performed with OPSL lasers
at 365 nm, 488 nm, and 568 nm.

Whole Mount Tissue Hybridization Chain Reaction (HCR)

Hybridization chain reaction (HCR) was performed as described
previously[50]. Cyclin D1 and Fgf8 probe sets were generated using an online
probe generator (https://github.com/rwnull/insitu_probe_generator). Probe sets
for Axin2, Hes7, and Uncx4.1 were obtained from Molecular Instruments. HCR
imaging was performed using a Leica TCS SP8 confocal microscope with a HC
PL APO CS2 20×/0.75 DRY objective. The system was equipped with OPSL
lasers for excitation at 365 nm, 488 nm, 568 nm, and 638 nm.

Image and Data Processing

To quantify the fraction of phospho-histone H3 (pHH3)-positive cells,
skin, neural tube, notochord, and gut were manually removed from the DAPI
channel in section images using Fiji. Subsequently, StarDist, a Fiji plugin, was
used to segment the remaining nuclei, providing an accurate measurement of
the number of nuclei per slice. pHH3-positive cells were manually counted per
slice.

For the quantification of EdU- and BrdU-positive cells, skin, neural tube,
notochord, and gut were similarly excluded from the DAPI channel in Fiji. This
channel was further denoised using Remove Outliers in Fiji (pixel size= 4,
threshold=50) followed by a gaussian blur with a sigma of 2. Using Cellpose

cryo3 model we segmented the remaining nuclei, allowing for an accurate count of the total number of nuclei per slice. Mean intensities of EdU and BrdU were calculated within the DAPI masks, and thresholds were applied to categorize cells as positive or negative for EdU and/or BrdU. These annotations, expressed as fractions of the total cell count, were used for the quantification of cell cycle duration, population doubling time, and S-phase duration (see Fig. 1, Fig. S2,3).

To visualize tissue-wide dynamics of Hes7 and Fucci2-Cdt1, regions of interest (ROIs) were drawn in maximum intensity projections of time-lapse movies in Fiji. Intensity measurements of both channels were quantified over time, smoothed, and linearly normalized between 0 and 1. Kymographs of tissue-wide dynamics were generated using the KymoResliceWide plugin in Fiji. A line was drawn from the posterior to the anterior end of the tissue, with a width of 100 μm.

For the quantification of the T/Brachyury gradient, a similar approach was used. A line plot was generated by measuring the intensity of T/Brachyury along a line drawn in Fiji. Gradient length was defined as the distance from the tip of the tail to the point where T/Brachyury intensity reached 10% above background level. To quantify staining intensities, an ROI was manually drawn within the PSM, and the mean intensity was calculated using Fiji. For the measurement of PSM and somite lengths, DAPI images were used. Lines were drawn parallel to the neural tube from the tail tip to the first somite boundary for PSM size, or from the first to the second somite boundary for somite size.

For the quantification of tissue density, an ROI was drawn within confocal or light sheet images, and the number of cells within the selected area was counted using either the Achilles-Hes7 or DAPI channel.

To track single cells over short time intervals in confocal microscopy movies, an ROI of 3 × 3 pixels was manually tracked within the Fucci2-Cdt1 or Achilles-Hes7 channel in Fiji. The peak intensity of Fucci2-Cdt1 was defined as the moment when a cell transitions from G1 to the S-G2-M phase. These tracks were used to quantify cell velocity or the timing of cell cycle transitions.

For long-term single-cell tracking in light sheet images, cells were manually tracked by following mosaically expressed H2B or Nucmem reporters using the Fiji plugin Mastodon (https://github.com/mastodon-sc/mastodon). Spot centre intensities were measured within the tracked spots to quantify Achilles-Hes7 or Fucci2-Cdt1. These intensities were smoothed using a moving average of 3 time points and linearly normalized, such that each track contained intensity values between 0 and 1. For the analysis of Hes7 dynamics, PyBOAT was used with the following settings: min_period = 60 min, max_period = 300 min, and TC = 150. All data plotting was performed using Python.

<u>Statistical Analysis</u>

All experiments were conducted with a minimum of three independent replicates. Data visualization was performed using Tukey-style boxplots, where the bar represents the median, the box delineates the 25th and 75th percentiles, and the whiskers extend to 1.5 times the interquartile range (IQR). Violin plots were used to display the distribution of data from the maximum to the minimum, with the bar indicating the median value. For statistical comparisons between groups, normality was assessed using the Shapiro-Wilk test. Statistical significance was determined using either an independent t-test or the Mann-Whitney U test, depending on whether the data followed a normal distribution.

Data and Code availability

All data and custom code used in this study are available upon request to k.sonnen@hubrecht.eu.

Acknowledgements

We thank members of the Sonnen group for discussions and feedback. We are

grateful to Takashi Hiiragi and Juan Garaycoechea for discussions. We thank

Jan-Daniël de Leede for help with data analysis. We thank the animal, FACS

and imaging facilities of the Hubrecht Institute for their support. We thank

Alexander Aulehla for support and providing mouse lines, in particular the

inducible iRFPnucmem line. The H2B-mCherry mouse line was kindly provided

by the Riken Center for Biosystems Dynamics Research and the Fucci line by

the Riken BioResource Research Center. The T-Cre/ERT2 line was kindly

provided by Jackson Laboratories. The Achilles-Hes7 was kindly provided by

Ryoichiro Kageyama. This work was supported by the Hubrecht Institute and

received funding from the European Research Council under an ERC starting

grant agreement no. 850554 to K.F.S. In addition, this research was supported

in part by grant NSF PHY-1748958 to the Kavli Institute for Theoretical Physics

(KITP).

Author Contributions

M.J.O. and K.F.S. conceived the project together with Y.I.L. and B.D.S.; K.F.S.

supervised the project; M.J.O. performed experiments with help from

W.H.M.M., T.N., C.F., E.T., J.K. and W.M.T.; M.J.O., Y.I.L. and K.F.S. analysed

the data; Y.I.L. and B.D.S. developed the theoretical framework; M.J.O., Y.I.L.

and K.F.S. wrote the manuscript with input from all authors.

References

Meijer, W. H. M. & Sonnen, K. F. From signalling oscillations to somite formation. *Current Opinion in Systems Biology* **39**, 100520 (2024). https://doi.org/https://doi.org/10.1016/j.coisb.2024.100520

Hubaud, A. & Pourquie, O. Signalling dynamics in vertebrate segmentation. *Nat Rev Mol Cell Biol* **15**, 709-721 (2014). https://doi.org/10.1038/nrm3891

Matsuda, M. *et al.* Recapitulating the human segmentation clock with pluripotent stem cells. *Nature* **580**, 124-129 (2020). https://doi.org/10.1038/s41586-020-2144-9

Dequeant, M. L. *et al.* A complex oscillating network of signaling genes underlies the mouse segmentation clock. *Science* **314**, 1595-1598 (2006). https://doi.org/10.1126/science.1133141

Tam, P. P. The control of somitogenesis in mouse embryos. *J Embryol Exp Morphol* **65 Suppl**, 103-128 (1981).

Lauschke, V. M., Tsiairis, C. D., Francois, P. & Aulehla, A. Scaling of embryonic patterning based on phase-gradient encoding. *Nature* **493**, 101-105 (2013). https://doi.org/10.1038/nature11804

Ishimatsu, K. *et al.* Size-reduced embryos reveal a gradient scaling-based mechanism for zebrafish somite formation. *Development* **145** (2018). https://doi.org/10.1242/dev.161257

Zhang, L., Kendrick, C., Julich, D. & Holley, S. A. Cell cycle progression is required for zebrafish somite morphogenesis but not segmentation clock function. *Development* **135**, 2065-2070 (2008). https://doi.org/10.1242/dev.022673

Benazeraf, B. *et al.* Multi-scale quantification of tissue behavior during amniote embryo axis elongation. *Development* **144**, 4462-4472 (2017). https://doi.org/10.1242/dev.150557

Bulusu, V. *et al.* Spatiotemporal Analysis of a Glycolytic Activity Gradient Linked to Mouse Embryo Mesoderm Development. *Dev Cell* **40**, 331-341 e334 (2017). https://doi.org/10.1016/j.devcel.2017.01.015

Mongera, A., Michaut, A., Guillot, C., Xiong, F. & Pourquie, O. Mechanics of Anteroposterior Axis Formation in Vertebrates. *Annu Rev Cell Dev Biol* **35**, 259-283 (2019). https://doi.org/10.1146/annurev-cellbio-100818-125436

Diaz-Cuadros, M. *et al.* Metabolic regulation of species-specific developmental rates. *Nature* **613**, 550-557 (2023). https://doi.org/10.1038/s41586-022-05574-4

13 Diaz-Cuadros, M. *et al.* Metabolic regulation of species-specific
developmental rates. *bioRxiv*, 2021.2008.2027.457974 (2021).
https://doi.org/10.1101/2021.08.27.457974

14 Carrieri, F. A. *et al.* CDK1 and CDK2 regulate NICD1 turnover and the
periodicity of the segmentation clock. *EMBO Rep* **20**, e46436 (2019).
https://doi.org/10.15252/embr.201846436

15 Delaune, E. A., Francois, P., Shih, N. P. & Amacher, S. L. Single-cell-
resolution imaging of the impact of Notch signaling and mitosis on
segmentation clock dynamics. *Dev Cell* **23**, 995-1005 (2012).
https://doi.org/10.1016/j.devcel.2012.09.009

16 Wiedermann, G. *et al.* A balance of positive and negative regulators
determines the pace of the segmentation clock. *Elife* **4**, e05842 (2015).
https://doi.org/10.7554/eLife.05842

17 Kicheva, A. *et al.* Coordination of progenitor specification and growth in
mouse and chick spinal cord. *Science* **345**, 1254927 (2014).
https://doi.org/10.1126/science.1254927

18 Boehm, B. *et al.* The role of spatially controlled cell proliferation in limb
bud morphogenesis. *PLoS Biol* **8**, e1000420 (2010).
https://doi.org/10.1371/journal.pbio.1000420

19 Abe, T. *et al.* Visualization of cell cycle in mouse embryos with Fucci2
reporter directed by Rosa26 promoter. *Development* **140**, 237-246
(2013). https://doi.org/10.1242/dev.084111

20 El Azhar, Y. *et al.* Unravelling differential Hes1 dynamics during axis
elongation of mouse embryos through single-cell tracking. *Development*
151 (2024). https://doi.org/10.1242/dev.202936

21 Sakaue-Sawano, A. *et al.* Visualizing spatiotemporal dynamics of
multicellular cell-cycle progression. *Cell* **132**, 487-498 (2008).
https://doi.org/10.1016/j.cell.2007.12.033

22 Palmeirim, I., Henrique, D., Ish-Horowicz, D. & Pourquie, O. Avian hairy
gene expression identifies a molecular clock linked to vertebrate
segmentation and somitogenesis. *Cell* **91**, 639-648 (1997).
https://doi.org/10.1016/s0092-8674(00)80451-1

23 Evrard, Y. A., Lun, Y., Aulehla, A., Gan, L. & Johnson, R. L. lunatic fringe
is an essential mediator of somite segmentation and patterning. *Nature*
394, 377-381 (1998). https://doi.org/10.1038/28632

24 Abe, T. *et al.* Establishment of conditional reporter mouse lines at
ROSA26 locus for live cell imaging. *Genesis* **49**, 579-590 (2011).
https://doi.org/10.1002/dvg.20753

25   Yoshioka-Kobayashi, K. *et al.* Coupling delay controls synchronized
oscillation in the segmentation clock. *Nature* **580**, 119-123 (2020).
https://doi.org/10.1038/s41586-019-1882-z

26   Morgan, D. O. *The cell cycle : principles of control*. (2007).

27   Tsiairis, C. D. & Aulehla, A. Self-Organization of Embryonic Genetic
Oscillators into Spatiotemporal Wave Patterns. *Cell* **164**, 656-667
(2016). https://doi.org/10.1016/j.cell.2016.01.028

28   Meijer, H. A. *et al.* NOTCH1 S2513 is critical for the regulation of NICD
levels impacting the segmentation clock in hiPSC-derived PSM cells and
somitoids.   *bioRxiv*,   2024.2012.2010.627712   (2024).
https://doi.org/10.1101/2024.12.10.627712

29   Sonnen, K. F. *et al.* Modulation of Phase Shift between Wnt and Notch
Signaling Oscillations Controls Mesoderm Segmentation. *Cell* **172**,
1079-1090 e1012 (2018). https://doi.org/10.1016/j.cell.2018.01.026

30   Aulehla, A. *et al.* Wnt3a plays a major role in the segmentation clock
controlling   somitogenesis.   *Dev   Cell*   **4**,   395-406   (2003).
https://doi.org/10.1016/s1534-5807(03)00055-8

31   Greco, T. L. *et al.* Analysis of the vestigial tail mutation demonstrates
that Wnt-3a gene dosage regulates mouse axial development. *Genes &*
‚Ä¶ **10**, 313-324 (1996).

32   van Oostrom, M. J., Meijer, W. H. M. & Sonnen, K. F. A Microfluidics
Approach for the Functional Investigation of Signaling Oscillations
Governing   Somitogenesis.   *J   Vis   Exp*   **169**,   1-17   (2021).
https://doi.org/10.3791/62318

33   Masamizu, Y. *et al.* Real-time imaging of the somite segmentation clock:
revelation of unstable oscillators in the individual presomitic mesoderm
cells. *Proc Natl Acad Sci U S A* **103**, 1313-1318 (2006).
https://doi.org/10.1073/pnas.0508658103

34   Pourquie, O. The segmentation clock: converting embryonic time into
spatial   pattern.   *Science*   **301**,   328-330   (2003).
https://doi.org/10.1126/science.1085887

35   Soroldoni, D. *et al.* Genetic oscillations. A Doppler effect in embryonic
pattern   formation.   *Science*   **345**,   222-225   (2014).
https://doi.org/10.1126/science.1253089

36   Cooke, J. & Zeeman, E. C. A clock and wavefront model for control of
the number of repeated structures during animal morphogenesis. *J*
Theor Biol **58**, 455-476 (1976). https://doi.org/10.1016/s0022-
5193(76)80131-2

37   Niwa, Y. *et al.* Different types of oscillations in Notch and Fgf signaling
regulate the spatiotemporal periodicity of somitogenesis. *Genes Dev* **25**,
1115-1120 (2011). https://doi.org/10.1101/gad.2035311

38   Dubrulle, J. & Pourquie, O. fgf8 mRNA decay establishes a gradient that
couples axial elongation to patterning in the vertebrate embryo. *Nature*
427, 419-422 (2004). https://doi.org/10.1038/nature02216

39   Dubrulle, J., McGrew, M. J. & Pourquie, O. FGF signaling controls
somite boundary position and regulates segmentation clock control of
spatiotemporal Hox gene activation. *Cell* **106**, 219-232 (2001).
https://doi.org/10.1016/s0092-8674(01)00437-8

40   Naiche, L. A., Holder, N. & Lewandoski, M. FGF4 and FGF8 comprise
the wavefront activity that controls somitogenesis. *Proc Natl Acad Sci U*
S A **108**, 4018-4023 (2011). https://doi.org/10.1073/pnas.1007417108

41   Cambray, N. & Wilson, V. Axial progenitors with extensive potency are
localised to the mouse chordoneural hinge. *Development* **129**, 4855-
4866 (2002). https://doi.org/10.1242/dev.129.20.4855

42   Mathis, L. & Nicolas, J. F. Different clonal dispersion in the rostral and
caudal mouse central nervous system. *Development* **127**, 1277-1290
(2000). https://doi.org/10.1242/dev.127.6.1277

43   Neijts, R., Simmini, S., Giuliani, F., van Rooijen, C. & Deschamps, J.
Region-specific regulation of posterior axial elongation during vertebrate
embryogenesis. *Dev Dyn* **243**, 88-98 (2014).
https://doi.org/10.1002/dvdy.24027

44   Yaman, Y. I. & Ramanathan, S. Controlling human organoid symmetry
breaking reveals signaling gradients drive segmentation clock waves.
Cell **186**, 513-527 e519 (2023).
https://doi.org/10.1016/j.cell.2022.12.042

45   Thomson, L., Muresan, L. & Steventon, B. The zebrafish presomitic
mesoderm elongates through compaction-extension. *Cells Dev* **168**,
203748 (2021). https://doi.org/10.1016/j.cdev.2021.203748

46   Sabherwal, N. *et al.* Differential phase register of Hes1 oscillations with
mitoses underlies cell-cycle heterogeneity in ER(+) breast cancer cells.
Proc Natl Acad Sci U S A **118** (2021).
https://doi.org/10.1073/pnas.2113527118

47   Weterings, S. D. C. *et al.* NOTCH-driven oscillations control cell fate
decisions during intestinal homeostasis. *bioRxiv*,
2024.2008.2026.609553 (2024).
https://doi.org/10.1101/2024.08.26.609553

48   Maeda, Y., Isomura, A., Masaki, T. & Kageyama, R. Differential cell-
cycle control by oscillatory versus sustained Hes1 expression via p21.

Cell Rep **42**, 112520 (2023). https://doi.org/10.1016/j.celrep.2023.112520

49   Zeng, C. *et al.* Evaluation of 5-ethynyl-2'-deoxyuridine staining as a sensitive and reliable method for studying cell proliferation in the adult nervous system. *Brain Res* **1319**, 21-32 (2010). https://doi.org/10.1016/j.brainres.2009.12.092

50   Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* **145** (2018). https://doi.org/10.1242/dev.165753