Manual versus automatic annotation of transposable elements: case studies in *Drosophila melanogaster* and *Aedes albopictus*, balancing accuracy and biological relevance

Tomas Carrasco-Valenzuela[1,2,3*], Alba Marino[1,4*], Jessica M. Storer[5], Iago Bonnici[4], Camila J. Mazzoni[2,3], Michael C. Fontaine[6,7], Annabelle Haudry[1], Matthieu Boulesteix[1$@], Anna-Sophie Fiston-Lavier[4,8$@]

1 Université Claude Bernard Lyon 1, LBBE, UMR 5558, CNRS, VAS, Villeurbanne, 69622, France.
2 Evolutionary Genetics Department, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany.
3 Berlin Center for Genomics in Biodiversity Research, Berlin, Germany.
4 ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France
5. Institute for Systems Genomics, University of Connecticut, Storrs, CT 06269, USA
6. MIVEGEC, Univ. Montpellier, CNRS, IRD, Montpellier, France
7. Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, The Netherlands
8 Institut Universitaire de France (IUF). Paris.

* First coauthors
$ Colast authors

@Corresponding authors: matthieu.boulesteix@univ-lyon1.fr & anna-sophie.fiston-lavier@umontpellier.fr

Tomas Carrasco-Valenzuela: tomas.carrasco@univ-lyon1.fr/ORCID: 0000-0003-1850-5376
Alba Marino: alba.marino@umontpellier.fr/ORCID: 0009-0005-0984-2524
Jessica M. Storer: jessica.storer@uconn.edu/ORCID: 0000-0002-9619-5265
Camila J. Mazzoni: mazzoni@izw-berlin.de /ORCID: 0000-0001-6758-5427
Iago Bonnici: iago.bonnici@umontpellier.fr /ORCID:
Michael C. Fontaine: michael.fontaine@cnrs.fr/ORCID: 0000-0003-1156-4154
Annabelle Haudry: annabelle.haudry@univ-lyon1.fr/ORCID: 0000-0001-6088-0909
Matthieu Boulesteix: matthieu.boulesteix@univ-lyon1.fr/ORCID: 0000-0003-3909-9522
Anna-Sophie Fiston-Lavier: anna-sophie.fiston-lavier@umontpellier.fr/ORCID: 0000-0002-7306-6532

Abstract:

Transposable elements (TEs) play a pivotal role in genome evolution, yet their detection and annotation remain challenging due to the limitations of current methods. Manual curation is considered the gold standard for generating TE libraries, particularly for TE-focused studies, although it requires extensive training and time. With the rapid increase in the number of genome assembly publications and the growing need for large-scale comparative analyses, automated software for TE annotation has become indispensable. This study compares manual and automated approaches to TE detection and annotation, focusing on two species: *Drosophila melanogaster* and *Aedes albopictus*.

In *D. melanogaster*, a species with a well-annotated TE repertoire and a smaller genome, the differences between manual curation (MCTE) and automated annotation (ATTE) are relatively minor. However, significant differences arise when analysing *Ae. albopictus*, a species with a relatively large genome and high TE diversity. While automated methods identified a greater number of TEs, including many smaller and fragmented elements, manual curation provided more detailed classifications and, on average, larger consensi. Automated pipelines offer a viable alternative for genome-wide analyses such as TE content estimation, particularly when time and resources are limited. However, caution is advised when interpreting results, as finer details of TE dynamics may be overlooked.

This study highlights that the choice of annotation method depends on the intended analysis. Manual curation is more suitable for TE population genomics and studies focusing on recent transposable element activity, whereas automated methods are appropriate for larger comparative analyses or genome assembly projects. Both methods have strengths and limitations, and understanding the specific features of the genome and repeatome under study is essential for selecting the appropriate approach.

69   Introduction

As high-quality genome assemblies become more widely available,
the analysis and annotation of these genomes present a significant
challenge. This is particularly true for identifying transposable elements
(TEs), which are highly mutagenic and prone to losing their sequence
identity over time and across different genomes (McDonald, Weinstein,
and Lambert 1988; Arkhipova 2018). TEs are major contributors to
genome evolution, gene organisation and expression; thus, their proper
identification is crucial for understanding their roles in adaptation, as well
as in conflicts and cooperation with their host genomes (Hayward and
Gilbert 2022; Casacuberta and González 2013).

Several pipelines, such as RepeatModeler2 (Flynn et al. 2020),
REPET (Rafii and Pardo 2013) or EDTA (Ou et al. 2019) to name a few,
have been developed and are widely used by the research community to
identify TEs. Despite their differences, these methods all follow a similar
core principle: they search for repeated loci within the genome. Some
approaches compare these loci to curated TE sequence databases (S.
Orozco-Arias et al. 2023), whereas others generate *de novo* consensus
sequences. For a deeper comparison of their differences and similarities,
see Hoen et al. (2015). Ultimately, the confidence in the resulting TE
library depends on the quality of the reference TE databases and the
accuracy of detecting biologically meaningful repetitions in the genome.

However, automatic detection algorithms usually lack the precision
and exactitude required for certain downstream applications (Goubert et
al. 2022; Lerat 2010; Flynn et al. 2020). The best way to reach such
quality is to manually curate each generated TE consensus: this implies
checking for specific TE indicators and hallmarks, such as structural
repetitive motifs and coding domains. This process is very tedious, time-
consuming, difficult to reproduce, and challenging for non-TE specialists
(Goubert et al. 2022).

However, manual curation serves as a means of validation for
computational predictions, allowing researchers to confirm the accuracy

of TE annotations and refine results on the basis of specific biological
contexts. Platt et al. (2016) started to define a set of recommendations for
manual curation adopted by (Peona et al. 2021; Jebb et al. 2020; Louha et
al. 2020) and reviewed by (J. M. Storer et al. 2021; Goubert et al. 2022).

Very recently, several tools have been developed to automate the
curation process (Baril, Imrie, and Hayward 2022; S. Orozco-Arias et al.
2023; Simon Orozco-Arias et al. 2022): they are promising, as they can
improve the quality of automatically generated TE libraries, as well as aid
manual curators in their task and prevent reproducibility issues.

Despite the evident need for manual curation when focusing on TE
biology, researchers tend to choose an automated approach over a
manual approach or the other way around, depending on time availability,
expertise, and the scope of their project. However, this choice might
ultimately affect the results of downstream analyses.

Studying the contribution of TEs to recent adaptations can lead to a
focus on recently active and abundant TE families. Furthermore, because
genotyping methods involve read mapping at the extremities of TE
insertions, the description of full-length consensi is a priority for this kind
of study.

Such an approach is impractical when a global assessment of the
entire genome is required and/or when numerous taxa are to be
annotated. On the other hand, a "simple" *de novo* annotation can lack the
level of definition needed because of redundancy, fragmentation and
unclassified elements (Baril, Imrie, and Hayward 2022; Rodriguez and
Makałowski 2022). A detection workflow that is time-effective and that will
produce accurate TE annotations is therefore preferable to allow in-depth
comparative analysis of TE accumulation across numerous nonmodel
species. To meet such needs, some authors of this paper have chosen two
different approaches and independently developed two different protocols
—a manual-based protocol and a fully automated protocol—to annotate
the genome of the same species, the Asian tiger mosquito (*Aedes*
albopictus). The former approach involves *de novo* (RepeatModeler2,
EDTA, MITE-Tracker) and homology-based (RepeatMasker used with

RepBase) TE discovery, followed by redundancy removal, consensi extension and extensive manual curation: its product is hereafter referred to as Manually Curated TE (MCTE) library. The latter combines the outputs of two identification methods (RepeatModeler2 and EDTA) to reduce redundancy and perform automatic curation of TE sequences via MCHelper in fully automated mode: its product is hereafter referred to as Automatic Tools TE (ATTE) library.

The *Aedes albopictus* genome is large by Dipteran standards (1.190–1.275 Gb; (Palatini et al. 2020); earlier studies based on sequencing data estimated that repetitive elements make up approximately 50%, with at least 33.58% being TE-derived (Goubert et al. 2015). More recent research based on transcriptomic data suggests that TEs constitute more than 40% of the genome (de Melo and Wallau 2020). This makes it difficult to determine in advance the completeness and accuracy of both the ATTE and MCTE approaches. Therefore, as a control, we included *Drosophila melanogaster* in the analysis, given its compact genome (180 Mb), excellent genome assembly and comprehensive TE annotation (Rech et al., 2022).

Initial comparisons between the MCTE and ATTE approaches for both genomes revealed minimal overlap between the obtained libraries. We explored these discrepancies in *Ae. albopictus* and examined whether they arose from the genome's complexity or non exhaustiveness of the MCTE approach. ATTE tends to identify more sequences, particularly smaller and fragmented elements, whereas MCTE provides longer consensi with more detailed classifications. This difference is especially evident in *Ae. albopictus,* which harbours greater TE diversity.

When comparing TE libraries across two genomes—*D. melanogaster*, a model organism with only 20% TE content, and *Ae. albopictus*, with its large genome (1.3 Gb) and ~40% TE content—we found that both approaches produce quantitatively distinct genome annotations. While automated curation cannot fully replace manual efforts, the two methods are complementary. Automated approaches can

166 guide manual curation or be integrated to generate more exhaustive
167 libraries.

168   Finally, we provide a composite TE library for *Ae. albopictus* by
169 combining both methods, which can be used for future studies.

Materials and methods

TE discovery for manual curation in *Ae. albopictus*.

For the construction of the MCTE library, TE consensi were obtained
from the AalbF2 genome assembly (Aalbo_primary.1; (Palatini et al.
2020)) via three distinct annotation algorithms: EDTA-2.0.0 (Ou et al.
2019), RepeatModeler-2.0.2a (Flynn et al. 2020), and MITE-Tracker
(Crescente et al. 2018). The resulting TE consensi were combined into a
single multifasta file, along with a subset of sequences from RepBase
25.08 (Jurka et al. 2005; Bao, Kojima, and Kohany 2015) (including
sequences from *Anopheles gambiae*, *Drosophila melanogaster*, Invrep -
Invertebrata-, and Invsub -Invertebrata subfamilies-). This file was then
used to mask the genome with the RepeatMasker command:
`RepeatMasker -pa 64 -s -a -inv -nolow -gff -dir . -lib`
`$WORK_DIR/TE_Aealb.raw.fa -cutoff 250 $ASSEMBLY`. RepeatMasker
hits shorter than 80 bps were removed because of the likelihood of
spurious results. After masking the genome, the OneCodeToFindThemAll
pipeline (Bailly-Bechet, Haudry, and Lerat 2014) was applied to
reconstruct TE copies by merging close-by RepeatMasker hits via the
following command:
`build_dictionary.pl --rm Aalb.out --unknown --fuzzy >`
`dico_fuzzy.txt \ one_code_to_find_them_all.pl --rm Aalb.out --`
`ltr dico_fuzzy.txt --fasta --flanking 100 --strict --unknown`
`--insert 80`
A custom script was then used to parse the output of one code to
find them all (Bailly-Bechet, Haudry, and Lerat 2014) and extract a fasta

file containing all the TE copies sequences
(https://github.com/Tcvalenzuela/Manual-versus-automatic-annotation-of-
transposable-elements/octfta_to_fasta.sh).

To group TE copies by families, RepeatMasker-identified insertions
were clustered via cd-hit (Huang et al. 2010; Li and Godzik 2006) with
parameters set to match the 80-80-80 rule as much as possible: identity
of 80% or greater along more than 80% of the sequence in sequences
longer than 80 bp as described in Flutre (2011), as follows: `cd-hit est -`
`i copies.fasta -o consensi.fasta -c 0.8 -G 0 -aS 0.8 -M 90000`
`-d 0`.

After several test runs, to increase the high copy number families,
we selected only clusters with more than seven insertions. Consensi were
called from the remaining clusters with the Refiner tool from
RepeatModeler2 (Flynn et al. 2020) after downsampling 500 sequences
from the largest clusters. To reduce redundancy among family consensi,
these were in turn clustered with cd-hit-est and consensi called again with
Refiner (Flynn et al. 2020). The process was repeated until no redundancy
was detected in the library (total number of iterations: 8). In the end,
23,009 family consensi were obtained at this step.

To increase the likelihood of obtaining full-length TE models, all
23,009 consensi were extended via the following method. First, the
putative TEs were screened for low complexity and simple repeat
sequences via TRF (Benson 1999). To gather sequences for extension, the
filtered sequences were used as a library for RepeatMasker (v4.1.5) on the
AalbF2 genome assembly. An alignment per family is generated in
stockholm (.stk) format, followed by extension, using somewhat relaxed
parameters for extension into the flanking sequence based on the
chromosomal locations present in the .stk files. Following extension,
additional identifying information might have been obtained, such as long
terminal repeats (to allow for endogenous retroviruses identification),
terminal inverted repeats (for DNA family identification), or polyA tails (for
LINE/SINE identification). Therefore, RepeatClassifier (a utility part of
RepeatModeler) is run to take this additional information into account.

The extended consensi was used as a reference in a second RepeatMasker run, this time performed on AalbF3, a deduplicated version of F2 plus optical mapping reference genome assembly GCA_018104305.1 (Boyle et al. 2021). The output of RepeatMasker was used to sort the list of consensi by the total number of full-length insertions, where a full-length insertion was defined as at least 90% of the length of the consensus, with a maximum nucleotide divergence of 20% (Flutre et al. 2011). Using the custom script https://github.com/Tcvalenzuela/Manual-versus-automatic-annotation-of-transposable-elements/detectFullSize.py, the list of consensi was finally sorted by full-length frequency, and manual curation was performed starting from the most frequent consensi. Elements with fewer than three insertions were not considered for manual curation and were not part of the final library either.

Manual curation of TEs from *Ae. albopictus*

The relevant information necessary for manual curations, such as the absolute frequency of insertions, pre- and post-extension length, insertion frequency in each genome assembly (AalbF3 and AalbF5 versions), ratio of extended consensus length to original consensus length, extended consensus coverage, extended consensus median coverage, insertion frequency, and full-length insertion frequency (Supplementary Table 1), was computed for all the consensi. The frequency of full-length insertion was used as a priority list for manual curation.

Manual curation was conducted on the TE consensus of the 800 most frequent full-length insertions, resulting in a final TE library containing 497 TEs. First, from the insertion coordinates from RepeatMasker and using the custom script https://github.com/Tcvalenzuela/Manual-versus-automatic-annotation-of-transposable-elements/GetMultipleAln.sh, each insertion was extended to 2,000 bp on both flanks and extracted, and the 100 longest insertions were clustered together via ClustalO (Sievers and Higgins 2014) for examination of the characteristic component of the respective category of

TEs. Additionally, each TE consensus was examined via a set of manual
curation identification tools, such as TE-Aid (Goubert et al. 2022),
RepeatClassifier (Flynn et al. 2020), and alignment visualisation via
Aliview (Larsson 2014), together with databases as Repbase (Jurka et al.
2005; Kohany et al. 2006; Vladimir V. Kapitonov and Jurka 2008) and CDD
protein domains (Lu et al. 2020; Marchler-Bauer et al. 2015), following the
annotation recommendations from Goubert et al. (2022).

In summary, following the guidelines of Wicker et al. (2007), long
terminal repeats (LTRs) are identified based on the presence of open
reading frames (ORFs) encoding proteins such as capsid protein, aspartic
proteinase, integrase, RNase H, reverse transcriptase, and envelope
protein, depending on the superfamily. Additionally, the LTR element must
be flanked by target site duplications (TSDs) of 4–6 base pairs (Neumann
et al. 2019).

Penelope-like elements can be further distinguished from homing
endonucleases (ENs) by the presence of a conserved CCHH Zn-finger
motif, where two cysteines are located between the GIY and YIG motifs
(Arkhipova 2006). The typical structure of LINEs includes a
retrotranscriptase combined with an apurinic endonuclease or ENs
domain, and some families have an additional ORF containing RNase H
(Wicker et al. 2007). For LINEs and Penelope-like elements, 5' truncation
products are expected on clustering of different insertions (Arkhipova
2006).

Terminal inverted repeats (TIRs) and variable-size TSDs are key
identifiers for most class II elements. Owing to the absence of conserved
protein domains across all class II transposons, the identification of class II
elements relies on TIR and TSD. After the identification of TIRs and TSDs,
their open reading frame (ORF) phylogeny is critical for classification
between Class II families. In eukaryotic transposons, the transposase
encoded is a DDD/E transposase (Kojima 2020). For nonautonomous
copies, it is important to check whether the TIR matches known
autonomous elements. If no match is found and the consensus sequence

is shorter than 800 bp, the element is classified as a MITE (Hu, Zheng, and
Shang 2018). If it exceeds 800 bp, it is categorised as Class II_other.

More complex class II elements, such as Helitrons and Mavericks,
can be classified on the basis of their homology to protein domains owing
to their unique protein sets and lack of TSDs (V. V. Kapitonov and Jurka
2001), although most of these elements are nonautonomous.

While LTRs and TIRs should be detectable via TE-Aids, there are
instances where TIRs are not immediately visible, even when coverage
suggests a class II element. In such cases, manual inspection using a
visualiser such as Aliview (Larsson 2014) to locate the TIR is
recommended.

As in previous analyses of TEs in mosquitos (de Melo and Wallau
2020), there are limitations to TE annotation due to a lack of homology to
TE databases or to the loss of ORFs due to accumulated mutations. For
some elements, further Class I or Class II classification was impossible for
both MCTE and ATTE library construction. To simplify the comparison,
these elements are referred to as ClassI_other and ClassII_other.

Manually annotated library for *D. melanogaster*

The evaluation of the ATTE method also depends on the
completeness of the MCTE library used as a control. Therefore, for *D.
melanogaster*, we used the manually curated TE library from Rech et al.
(2022). They used long-read data to assemble high-quality genomes of 32
D. melanogaster natural strains from five different climatic regions. From
this genomic resource, Rech et al. (2022) manually curated a TE dataset
composed of 165 consensus sequences, thoroughly representing the
genomic TE diversity in this species.

Automated discovery and curation of TEs in *Ae. albopictus* and *D. melanogaster*

The study of TE dynamics in larger numbers of species can require
an automated strategy of TE discovery where automatic quality checks
and curation steps are integrated to obtain better-quality annotations. The

genomes AalbF5 for *Ae. albopictus* and dm6 for *D. melanogaster* (Hoskins
et al. 2015) were annotated with the following automated pipeline to
obtain the respective ATTE libraries. New insertions were identified with
RepeatModeler2 (Flynn et al. 2020) and EDTA (Ou et al. 2019). Each
program was run independently, and the mined sequences were
concatenated on a single multifasta file. Afterward, MCHelper (S. Orozco-
Arias et al. 2023) was used in fully automated mode to eliminate
redundancy, remove false positives, elongate and check for TE
classification.

334   Comparison of TE libraries

To compare the MCTE and ATTE libraries, we performed a
clusterization of the consensi between libraries for each species following
an 80-80 rule (Flutre et al. 2011), using cd-hit as follows:
`cd-hit-est-2d -i <Library MC> -i2 <Library AT> -o <Output`
`Name> -c 0.8 -G 0 -aS 0.8 -n 5 -d 0`
To further compare the MCTE and ATTE libraries, we used the script
"get_family_summary_paper.sh" from (Flynn et al. 2020). Briefly, a target
library is masked with a reference library via RepeatMasker with -nolow
option, and its level of completeness is evaluated based on the reference.
The overlap between the MCTE and ATTE libraries was assessed by using
the ATTE and MCTE libraries as references.
Based on Flynn (2020), we use an in-house script
(https://github.com/Tcvalenzuela/Manual-versus-automatic-annotation-of-
transposable-elements/blob/main/reassign_family_categories.sh),   where
each element of the reference library is assigned to one of the following 5
categories: "Perfect", "Good,", "Present", "Fragmented" or "Absent". The
first 3 categories are defined following Flynn et al. (2020). "Perfect"
families are those for which one sequence in the target library matches
>95% in sequence identity and coverage with a family in the reference
library. "Good" families are those in which multiple overlapping target-
library sequences with alignments >95% similar to the reference
consensus make up >95% sequence coverage of the element from the

reference library. A family is considered "Present" if one or multiple target
sequences align with >80% similarity to and cover >80% of the reference
consensus sequence. In addition, we consider a family to be
"Fragmented" if multiple query sequences with >80% similarity cover
>30% of a consensus, thus accounting for potential fragments that might
match full-length families in the reference library. Otherwise, we consider
a family to be "Absent".

Finally, each library was used to annotate the respective genome
and compare the genomic estimates of TE content obtained via manual
and automated curation. RepeatMasker (Nishimura 2000; Smit, Hubley,
and Green 2015) was run with RMblast as the search engine, and
annotations were used to quantify the overall genomic TE composition
and the proportion occupied by each TE consensus (Supplementary Table
2).

Results

TE library composition

The MCTE library for *Ae. albopictus* included 496 consensi with up to
733 full-length insertions, with an average of 94.5 insertions in the AalbF5
genome assembly. It is composed of 135 LINEs (27.16% of the total
consensi), 129 MITEs, 80 unclassified Class II elements, 79 LTRs, 60 DNA
elements, 8 *Penelope-like* elements, 4 SINEs and 1 Helitron. On the other
hand, the ATTE library for *Ae. albopictus* comprises 7,782 consensi and is
composed of 5,750 LTRs (73.89% of the total), 1,024 DNA elements, 169
LINEs, 795 MITEs, 1 SINE and 9 RCs, and 33 and 1 unclassified element of
Class I and Class II, respectively (Figure 1).

The MCTE library for *D. melanogaster* was obtained from the public
database made available by Rech et al. (2022). Statistics, including
consensi length, frequency of the genome, and other descriptive
statistics, can be found in Supplementary Table 2. This library was
composed of 165 consensi: 91 LTR elements (55.15% of the total), 45
LINEs, 23 DNA elements, five RCs, and one MITE. The *D. melanogaster*

ATTE library is composed of 463 consensi and has a high proportion of LTR
elements, with 225 consensi (48.6% of the total consensi), 129 DNA
consensi, 61 LINEs, 38 MITEs, 7 RCs and 3 unclassified Class I elements
(Figure 1).

In both species, the ATTE library contains more consensi than the
MCTE library does, although this is particularly evident in *Ae. albopictus,*
where the ATTE library contains 15 times more consensi. For *D.*
melanogaster, the consensus lengths of the ATTE library are not
significantly shorter than those of the MCTE library for any TE order
(Figure 2; Wilcoxon test: DNA: W = 1701, p = 0.138; LINE: W = 1512, p =
0.271; LTR: W = 11112, p = 0.111; MITE: W = 38, p = 0.100). In *Ae.*
albopictus, LTR and LINE elements are significantly shorter in the ATTE
library than in their counterparts in the MCTE library (LTRs: average
length: 2,913 bp in the ATTE library vs. 6,738 bp in the MCTE library;
LINEs: average length: 3,670 bp in the ATTE library vs. 4,574 bp in the
MCTE library). This difference in length is statistically significant for both
LTR (Wilcoxon test: W = 66,322, p = 4.07e-27) and LINE elements
(Wilcoxon test: W = 7,881, p = 3.66e-6).

Figure 1. Composition of the TE libraries of (A) *D. melanogaster* and (B) *Ae. albopictus*.
The proportions of TE sequence types are shown both in percentages on the y-axes and
as the number of sequences on the stacked bars. The number of total sequences is
indicated at the bottom of the bar plots for each library. For categories with only one
sequence, the numbers are not displayed.

Figure 2. TE size distributions of the MCTE (bottom panels) and ATTE (top panels) libraries. For each TE library, the lengths of the
consensus sequences were measured, and the distribution was plotted by grouping the elements by their respective TE order.

How similar are libraries automatically generated versus those
manually curated?

While identical, complete copies from start to finish in the MCTE and
ATTE libraries are ideal; achieving this for TEs is seldom possible in the
current state of research. However, we can assess how overlapping the
MCTE and ATTE libraries are by comparing sequence similarity and
coverage.

The libraries were clustered via the 80–80 rule, as detailed in the
methods section. In *Ae. albopictus*, 58.63% of the consensi from the MCTE
library had no corresponding match in the ATTE library (Supplementary
Table 3). Of the remaining 41.37% of clustered sequences, two-thirds
matched with consensi of the same TE order. In contrast, 98.10% of the
ATTE library lacked a match in the MCTE library, with only 0.82%
clustering with sequences of the same TE order.

Similar results were observed in *D. melanogaster*, where 98.27% of
the ATTE library consensi (Supplementary Table 3) did not match the
MCTE library. Of the remaining 1.73% (8 consensi), 7 matched at the TE
order level. On the other hand, 74.55% of the MCTE library had no
equivalent consensi in the ATTE library, whereas the remaining 25.45%
were largely composed of consensi matching the same TE order.

We compared more exhaustively the composition of the libraries by
categorising consensi as "Perfect", "Good", "Present" (as in Flynn (2020)),
and "Fragmented" and "Absent" with respect to the alternative library
(see the Methods section).

Overall, 64% of the *D. melanogaster* MCTE sequences had a hit in
the ATTE library and, in most cases, had the same TE type
(Supplementary Table 4). Overall, 28% of the MCTE sequences fall into the
category "Perfect", meaning that they have > 95% identity over 95% of
the length with one sequence in the ATTE library (Figure 3A;
Supplementary Table 4). These elements correspond mostly to LTR
elements, followed by LINEs. "Good" sequences (those elements with

>95% identity over 95% of the length in multiple sequences) make up
12% of the MCTE library and are predominantly LTRs. In the reverse
comparison, we identify that 75% of the sequences in the ATTE library do
not have any match in the MCTE one, arguably because of the larger size
of the former (2.8 AT/MC ratio). Compared with the MCTE library, a
significantly smaller fraction of the ATTE library has corresponding hits in
the MCTE library (Perfect = 9%, Good = 4%, Present = 4%, Fragmented =
7%): the large majority are LTR elements and have the same classification
in the two libraries.

Figure 3. Comparison between the ATTE and MCTE libraries in *D. melanogaster*.
(A) Level of completeness of the ATTE library ("query") versus the MC library
("reference"): the percentages of TEs in the MCTE library detected in the ATTE library are
shown for the categories Absent, Fragment, Present, Good and Perfect (left, scale of
gray). For each category, the proportion of consensi is displayed by TE order (right,
coloured stacked bars). (B) The same analysis was performed using the ATTE library as
the "reference" and the MCTE library as the "query".

The same analysis was performed on the *Ae. albopictus* libraries
(Figure 4). Here, three-quarters of the MCTE elements have a similar ATTE
consensus: 36.82% "Perfect", 13.88% "Good", 16.10% "Present", 7.65%
'Fragmented', and 25.55% absent elements (Figure 4A). In particular,
59% of the MCTE consensi that have no counterpart in the AT library are

LINEs. Additionally, most consensi of this category overall correspond to
Class II elements, including MITEs (~25%), DNA (17%), and other non
further classified elements (23%). Furthermore, while 42.25% of the MCTE
consensi matched with sequences of equivalent classification, 32.19%
matched with different TE types, with MITEs constituting 46.88% of the
misclassifications three times more than the next category (LINEs with
17.50% of the misclassifications) (Supplementary Table 4).

In contrast, (Figure 4B), we again identified a substantial number of
AT consensi (96%) as absent from the MC library. This could be due to the
ratio of 15.5 AT/MC, hence either the overabundance of AT sequences or
undersampling of MC sequences. 75% of the absent elements are LTRs,
which are also the most abundant in the AT library. Out of the 302
matching AT sequences (~ 4% of the total), 170 had hits with the same TE
order; the rest had hits with sequences with different classifications,
notably LTR and DNA elements.

Figure 4. Comparison between the ATTE and MCTE libraries in *Ae. albopictus*. The level of completion for the automatically generated library versus the manually curated library in *Ae. albopictus* in panel A illustrates the percentage of TE consensi from the MC ("reference") detected in the AT ("query") for the categories Absent, Fragment, Present, Good and Perfect on a scale of gray. For each category, the portions represented by different TE orders are displayed as stacked bars. Panel B represents the same analysis with the consensi from AT ("reference") detected in the MC library ("query").

Are we capturing the full genetic diversity of transposable elements within the genome?

The ability to draw biologically significant conclusions about TEs depends on capturing a substantial proportion of the number and diversity of TEs present in the organisms. For this purpose, it is important to assess the genomic coverage of the different transposable detection alternatives together with the TEs variability detected. We compared the two different library strategies for the two species, masking the genomes and identifying the level of coverage per strategy, as described in the methods section (Figure 5).

For the *Ae. albopictus* genome, the TE coverage obtained with the MCTE library was lower (40.8% of the genome) than that obtained with the ATTE library (75.4% of the genome). This might be due to fragmented copies in the ATTE library or a reflection of the incomplete nature of the MC library in *Ae. albopictus*, which focuses on recent TE families with >3 full-length copies in the genome. In addition, and in conjunction with the results shown in Figure 1, we observed a prevalence of LTRs annotated with the ATTE library. Interestingly, a sharp increase in genome coverage (~ 12-21%) is observed with the MCTE library, which is due to a LINE/RTE-BovB with a full-length consensus of 4,000 bp and 168,000 insertions—between fragmented and full copies—occupying 9% of the genome.

In the *Ae. albopictus* MCTE library, we observed clear size differences across TE categories, reflecting the progressively smaller sizes of its elements. However, this pattern is not evident in the ATTE library for the same genome, where LTR elements dominate the entire size distribution, as shown in Figure 5.

For *D. melanogaster*, the TE genome coverage is quite similar, with 18% for MCTE and 19.2% for ATTE. This aligns with our expectations,

given the non-significant differences in consensi length distribution, despite variations in sequence similarity. Moreover, the relative scarcity of TEs in *D. melanogaster* and the extensive manual curation already performed in its genome have contributed to these results. Additionally, compared with those in *Ae. albopictus*, the TE families in both MCTE and ATTE show less diversity and lower copy numbers.

Figure 5. Comparative cumulative genome coverage and length distribution of TE consensi with the MCTE and ATTE libraries. (A) Levels of cumulative genome coverage are shown by decreasing the consensus length for the *D. melanogaster* (right) and *Ae. albopictus* (left) genomes via the MCTE libraries (bottom) versus the ATTE libraries (top). Boxplots on the right and top of each scatterplot highlight the distributions of the different TE orders along the x- and y-axes. The colors correspond to different TE categories, as per the legend on top.

Discussion

Owing to the limitations of current TE detection methods, manual curation remains the gold standard for generating TE libraries, particularly for TE-focused analyses, although it requires significant training and execution time. However, with the rapid pace of genome assembly publications and the growing need for large-scale comparative analyses, automated software for TE annotation and curation is becoming increasingly indispensable. In this study, we compare manual and automated approaches and show that differences in library composition, TE identification accuracy, and genome annotation are determined not only by the chosen method but also by the specific features of the genome and repeatome being analysed.

Transposable element library composition.

We chose to analyse *D. melanogaster* alongside *Ae. albopictus* as a proof of concept, given the reliability of the extensive manual annotation available for *D. melanogaster*. Although both are classified under the term "MCTE," the *D. melanogaster* database differs from that of *Ae. albopictus*, as the latter is not fully complete, unlike the comprehensive database described by Rech et al. (2022). This comparison allows us to assess the strengths and limitations of the ATTE method.

In *D. melanogaster*, the differences in TE identification between MCTE and ATTE are relatively minor, likely due to their smaller genome, reduced TE diversity, and lower TE content than those of *Ae. albopictus*. Nevertheless, the same discrepancies between the two methods persist, highlighting inherent differences in their outcomes.

Approximately 3 times more consensi were present in the ATTE library from *D. melanogaster* than in its MCTE counterpart. Furthermore, many LTR retrotransposons and DNA transposons were represented by short consensi in this ATTE library. LTR and DNA elements have defined 5'

and 3' ends (direct and inverted repeats, respectively) (Goubert et al.
2022; Makałowski et al. 2019; Wells and Feschotte 2020). This structure
makes them relatively straightforward to detect computationally.
However, identifying full-length consensus sequences remains
challenging, as many insertions are fragmented or incomplete, leading
automatic pipelines to often identify them as fragmented elements.
Although substantial efforts by the community have been carried on in
order to fix this issue, as in LTR_retriver (Ou and Jiang 2018), LTR
annotator (You et al., n.d.), and LTRharvest (Ellinghaus, Kurtz, and
Willhoeft 2008), among others, this issue is still present in modern and
complex automatic pipelines such as MCHelper (S. Orozco-Arias et al.
2023). Owing to their length, complex structure and propensity of their
terminal repeats to recombine, the elements of this order are difficult to
identify and distinguish between autonomous and non-autonomous (Flynn
et al. 2020; Ji and DeWoody 2016). With respect to the library composition
of *D. melanogaster*, despite ATTE sequences being more numerous, the
relative proportions of TE orders are not different from those of the MCTE
library.

In contrast, for *Ae. albopictus*, there were substantial differences
between the ATTE and MCTE libraries. For our MCTE library for *Ae.
albopictus*, we identified 135 LINE elements, constituting 27.16% of the
total number of consensi in the library. With these findings, we identified
23.0% of the genome as LINEs. This percentage exceeds previous reports
for this species and closely related species, such as 12.09% from Goubert
(2015), 15.7% from *Ae. albopictus* and 14.5% from *Aedes aegypti,* the
sister species of *Ae. albopictus,* reported by Melo and Wallau (2020).
However, our findings align with Melo and Wallau's observation that LINEs
are the most abundant TE class.

Furthermore, we identified a high proportion of MITE elements in the
MCTE library (25.96% of consensi) in the *Ae. albopictus* genome. These
elements are poorly characterised in current databases, despite recent
efforts to classify them in insect genomes (Han et al. 2016; Chen et al.

2014; Ye, Ji, and Liang 2016). We found MITEs to be an abundant yet
uncharacterised transposable element category.

Additionally, we identified elements with MITE hallmarks, such as
TIRs and TSDs ranging from 800-3000 bp in size, but devoid of ORFs.
Although very similar to MITEs, the literature only confidently classifies
them as such only up to a threshold of 800 bp, following the guidelines of
MITEFinderII (Hu, Zheng, and Shang 2018). Therefore, longer elements
were flagged as belonging to the most likely transposable origin family,
labelled "originalDNAElement_NonAutonomous", or if determining the
origin family was not possible, simply "Uncharacterised_ClassII".

For simplicity in comparison with the MCHelper annotation, these
are referred to as "Class II_others" in the figures. We also identified eight
full-length insertions of *Penelope*-like elements, which have previously
been reported in Culicinae mosquitoes (de Melo and Wallau 2020). These
findings served as a positive control, confirming the thoroughness of our
non-exhaustive manual curation of *Ae. albopictus*.

In contrast, for the ATTE library of *Ae. albopictus*, we identified
predominantly LTRs, with 73.89% of the total consensi. This might be
related to the detection of fragmented members of the same family of
more complete LTRs as independent elements, as discussed before. As
illustrated in Figure 5, LTRs, along with DNA elements (the second most
abundant class for the ATTE library) and MITEs, are particularly abundant
in shorter sequence lengths.

In addition to these differences, owing to the nature of manual
curation, the MCTE database has more detailed annotations than ATTE
does. For simplification of the comparison in the figures, the classification
of MCTE sequences has been simplified to the superfamily. Nonetheless,
classification at the family level is more abundant in the MCTE database
than in the ATTE database (Supplementary Table 2).

For the *Ae. albopictus* comparison, significant differences in the
consensus size distributions were observed only for the LTR and DNA
elements. For the other TE categories, no notable differences were
detected. Similarly, no significant differences were found in the *D.*

melanogaster comparison. Therefore, if the objective is not to analyse TEs
but rather to clean their assembly, as is sometimes necessary for genome
assembly or gene evolution analysis, an automatic method such as
MCHelper seems to be a good alternative, especially for small and well
characterised genomes and/or for genomes where the diversity of TE
insertions is limited, such as in *D. melanogaster* or humans (Mills et al.
2007).

Using our MCTE library, we identified 40.8% of the *Ae. albopictus*
genome as TEs, a proportion that is consistent with findings from other
studies. Goubert et al. (2015) reported that TEs make up approximately
33.58% of the genome. More recent studies have built upon this,
indicating that TEs account for more than 40% of the genome (de Melo
and Wallau 2020). Nonetheless, using the ATTE library, we identified
73.89% of the genome as TEs. One explanation for the difference in
genome coverage for the libraries of *Ae. albopictus* might be due to either
the lack of exhaustiveness of the MCTE library or the use of the fully
automated mode of MCHelper.

Accordingly, consensi classified as "incomplete" were retained in
the final library (the consensus name has the suffix "_inc"): these include
all sequences that do show homology with a known TE family but do not
conserve all the expected structural or coding domains. Moreover, several
"unconfirmed" sequences are also present (the consensus name has the
suffix "_unconfirmed"), meaning that they can be assigned only TIRs or
LTRs but fail any other validation by homology or coding domains (Simon
Orozco-Arias et al. 2023).

Since the *Ae. albopictus* ATTE library contains many of such
sequences (5,485 incomplete and 436 unconfirmed), it is possible for the
gap between our annotation and the previous annotation may be caused
by the detection of many motifs belonging to non-autonomous and long-
extinct elements. Another possibility is that the automatic pipeline
employed here may overestimate the repeat content due to an
overabundance of consensi in the ATTE library. However, this is unlikely to
be the case, as the estimations of TE content are very consistent between

the method used here and another published annotation pipeline (Baril,
Imrie, and Hayward 2022) across a wide range of genome sizes (Marino et
al. 2023).

On the other hand, the significant differences in the annotation
outcomes between the MCTE and ATTE libraries in the two species reflect
both the varying degrees of completion of the MCTE libraries and the
distinct genomic architectures of the two dipterans. The *D. melanogaster*
genome, at 0.18 Gb, is relatively poor in repetitive sequences, and its TE
repertoire has been well-annotated, making the MCTE library a nearly
exhaustive reference. In contrast, while the manual curation of *Ae.*
albopictus involves three times the number of *D. melanogaster* MCTE
consensi, the *Ae. albopictus* genome is 1.190 -- 1.275 Gb (Palatini et al.
2020), and it is not only rich in TEs but most of them are present in low
copy numbers (Supplementary Table S3). Achieving a comprehensive
description of TE diversity in tiger mosquitoes through manual curation
alone is an extremely labor-intensive process. However, relying solely on
manual curation would result in overlooking a significant portion of the
TEs present in the genome.

How similar are the automatically generated and manually curated
libraries?

For two sets of TE libraries to represent the evolutionary history of
TEs in the same organism, the consensi should be highly similar
sequences and represent the same variability in TE categories. To explore
this idea, we decided to cluster the sequences via cd-hit-est-2d and later
compared them as reciprocal species pairs -MC<->AT- via the approach
presented in Flynn (2020).

Despite the varying degrees of excess sequences associated with
different genomic TE densities, automatic libraries consistently contain far
more "Absent" consensi than their manual counterparts do. As previously
mentioned, this information may be redundant due to the limited
accuracy of the automatic "blast-extract-extend" process (Baril, Imrie, and
Hayward 2022; Simon Orozco-Arias et al. 2023). However, even if the

ATTE libraries retain fragmented sequences and the corresponding full-length consensi are present in the manual library, the similarity should still allow for their classification as "Fragmented" sequences. Some of the unique ATTE sequences may represent genuine "fossil" TEs that were missed—or impossible to recover—during manual curation.

Nevertheless, it is important to consider that various methods exist for curating consensi, which can result in discrepancies between libraries owing to differences in multisequence alignment, consensus generation, sequence trimming, and other factors. This issue likely extends beyond just the automatic vs. manual comparisons and applies to any comparison between different curation protocols.

Moreover, different expertises and sensitivities can also alter the outcome and lead to divergent libraries. While manual curation allows us to confidently verify the validity of sequences, one disadvantage is indeed the susceptibility of such a process to human bias. In this sense, a benefit of an automatic tool is the reproducibility of the curation process.

Furthermore, the use of a single consensus sequence may be limiting, as it may fail to capture the internal variation within an entire TE family. In the future, adopting the practice of storing family information as Hidden Markov Model (HMM) profiles rather than in FASTA format could help address this limitation. HMM profiles retain information about sequence variability at each position, preserving the diversity within the TE family (J. Storer et al. 2021). Both of the libraries produced in this work, *Ae. Albopictus* MCTE and ATTE, are available on DFAM (DFAM accession number) as multifasta files and HMM profiles on the basis of AalbF5.

Which method to choose then?

In most cases, TE annotation for newly sequenced genomes is delegated to either a similarity-based search (Platt, Blanco-Berdugo, and Ray 2016) or a *de novo* approach without further refinement. While this is often sufficient for basic genome masking, additional curation of TE libraries is necessary for in-depth analysis of the mobilome in the growing number of newly sequenced species. Although some examples of manual

curation exist (Martelossi et al. 2023; Osmanski et al. 2023; Rech et al.
2022), carrying out this method for a wide range of species or for large
and TE-rich genomes such as *Ae. albopictus* is time-consuming and labor-
intensive. Moreover, manual curation expertise tends to be confined
within certain research groups, may lack reproducibility, and, importantly,
there is no single universally optimal protocol. The choice of curation
method depends on the type of downstream analysis intended (Goubert et
al. 2022).

In our case, the MCTE library for *Ae. albopictus* was developed with
a focus on the most abundant elements showing signs of recent
transposition. Manual curation was initiated to identify TE insertions
potentially involved in local adaptation across different mosquito
populations. Since TE polymorphisms are more likely to be associated with
recent adaptations, prioritising potentially active, full-length elements was
particularly relevant for this population genomics study. For such studies
that rely on a single reference genome, manual curation remains the best
option. In contrast, manual curation would be impractical for large-scale
comparative genomic analyses, such as gene evolution studies or genome
assemblies, owing to the time constraints and the sheer number of
species typically involved.

Nonetheless, we have demonstrated that both approaches are
complementary, with minimal overlap in consensi identity and contrasting
strengths and weaknesses. Therefore, we provide both libraries, allowing
users to decide which one (or both) they should use, depending on their
specific needs.

As demonstrated in *D. melanogaster*, the TE content and
composition identified via manual versus automated approaches are
comparable, provided that the MCTE library is comprehensive. This finding
indicates that an automated pipeline can be appropriate for assessing TE
content in a comparative framework. However, finer analyses of TE
dynamics are limited, and the results should be interpreted with caution.

Authors contribution

T.C.-V. and A.M. performed the analysis. T.C.-V. performed the
manual curation of *Ae. albopictus.* T.C.-V., A.M., A.-S.F.-L. and M.B.
designed research and wrote the paper. J.S. performed the extension of
the original consensi in the first steps of the MCTE library of *Ae. albopictus*
and contributed with TE characterizations discussion. I.B., A.H., M.C.F.
Participated in designing the automatic detection pipeline. C.J.M. helped
with computational resources and insightful discussions.

Acknowledgement

773   We are indebted to Tomaso Barberis for his help in the initial steps
774 of the construction of the MCTE library of *Ae. albopictus*. This work was
775 performed using the computing facilities of the CC LBBE/PRABI and the
776 Montpellier Bioinformatics Biodiversity platform (MBB).

Funding

779   This work was supported by "Agence Nationale de la Recherche"
780 (ANR) MosquiTEs ANR-21-CE02-0013, NeGA ANR-20-CE02-0008-01 and
781 the "Institut Universitaire de France".

Supplementary files

Supplementary Figure 1. Overall (black) and by-order genomic TE coverage identified via automated and manual curation in *Ae. albopictus* and *D. melanogaster*. To be filled in the final figure.

Supplementary tables

Please see:

[Supplementary Tables](#)

References

Arkhipova, Irina R. 2006. "Distribution and Phylogeny of Penelope-like
Elements in Eukaryotes." *Systematic Biology* 55 (6): 875–85.
———. 2018. "Neutral Theory, Transposable Elements, and Eukaryotic
Genome Evolution." *Molecular Biology and Evolution* 35 (6): 1332–37.
Bailly-Bechet, Marc, Annabelle Haudry, and Emmanuelle Lerat. 2014.
"'One Code to Find Them All': A Perl Tool to Conveniently Parse
RepeatMasker Output Files." *Mobile DNA* 5 (1): 1–15.
Bao, Weidong, Kenji K. Kojima, and Oleksiy Kohany. 2015. "Repbase
Update, a Database of Repetitive Elements in Eukaryotic Genomes."
Mobile DNA 6 (1): 11.
Baril, Tobias, Ryan M. Imrie, and Alex Hayward. 2022. "Earl Grey: A Fully
Automated User-Friendly Transposable Element Annotation and
Analysis Pipeline." *bioRxiv*.
https://doi.org/10.1101/2022.06.30.498289.
Benson, G. 1999. "Tandem Repeats Finder: A Program to Analyze DNA
Sequences." *Nucleic Acids Research* 27 (2): 573–80.
Boyle, John H., Pasi M. A. Rastas, Xin Huang, Austin G. Garner, Indra
Vythilingam, and Peter A. Armbruster. 2021. "A Linkage-Based
Genome Assembly for the Mosquito Aedes Albopictus and
Identification of Chromosomal Regions Affecting Diapause." *Insects* 12
(2). https://doi.org/10.3390/insects12020167.
Casacuberta, Elena, and Josefa González. 2013. "The Impact of
Transposable Elements in Environmental Adaptation." *Molecular
Ecology* 22 (6): 1503–17.
Chen, Jiongjiong, Qun Hu, Yu Zhang, Chen Lu, and Hanhui Kuang. 2014.
"P-MITE: A Database for Plant Miniature Inverted-Repeat Transposable
Elements." *Nucleic Acids Research* 42 (Database issue): D1176–81.
Crescente, Juan Manuel, Diego Zavallo, Marcelo Helguera, and Leonardo
Sebastián Vanzetti. 2018. "MITE Tracker: An Accurate Approach to
Identify Miniature Inverted-Repeat Transposable Elements in Large
Genomes." *BMC Bioinformatics* 19 (1): 348.
Ellinghaus, David, Stefan Kurtz, and Ute Willhoeft. 2008. "LTRharvest, an
Efficient and Flexible Software for de Novo Detection of LTR
Retrotransposons." *BMC Bioinformatics* 9 (1): 18.
Flutre, Timothée, Elodie Duprat, Catherine Feuillet, and Hadi Quesneville.
2011. "Considering Transposable Element Diversification in de Novo
Annotation Approaches." *PloS One* 6 (1): e16526.
Flynn, Jullien M., Robert Hubley, Clément Goubert, Jeb Rosen, Andrew G.
Clark, Cédric Feschotte, and Arian F. Smit. 2020. "RepeatModeler2 for

Automated Genomic Discovery of Transposable Element Families."
Proceedings of the National Academy of Sciences of the United States
of America 117 (17): 9451–57.
Goubert, Clément, Laurent Modolo, Cristina Vieira, Claire ValienteMoro,
Patrick Mavingui, and Matthieu Boulesteix. 2015. "De Novo Assembly
and Annotation of the Asian Tiger Mosquito (Aedes Albopictus)
Repeatome with dnaPipeTE from Raw Genomic Reads and
Comparative Analysis with the Yellow Fever Mosquito (Aedes
Aegypti)." *Genome Biology and Evolution* 7 (4): 1192–1205.
Goubert, Rory J. Craig, Agustin F. Bilat, Valentina Peona, Aaron A. Vogan,
and Anna V. Protasio. 2022. "A Beginner's Guide to Manual Curation of
Transposable Elements." *Mobile DNA* 13 (1): 7.
Han, Min-Jin, Qiu-Zhong Zhou, Hua-Hao Zhang, Xiaoling Tong, Cheng Lu,
Ze Zhang, and Fangyin Dai. 2016. "iMITEdb: The Genome-Wide
Landscape of Miniature Inverted-Repeat Transposable Elements in
Insects." *Database: The Journal of Biological Databases and Curation*
2016 (December). https://doi.org/10.1093/database/baw148.
Hayward, Alexander, and Clément Gilbert. 2022. "Transposable
Elements." *Current Biology: CB* 32 (17): R904–9.
Hoen, Douglas R., Glenn Hickey, Guillaume Bourque, Josep Casacuberta,
Richard Cordaux, Cédric Feschotte, Anna-Sophie Fiston-Lavier, et al.
2015. "A Call for Benchmarking Transposable Element Annotation
Methods." *Mobile DNA* 6 (August):13.
Hoskins, Roger A., Joseph W. Carlson, Kenneth H. Wan, Soo Park, Ivonne
Mendez, Samuel E. Galle, Benjamin W. Booth, et al. 2015. "The
Release 6 Reference Sequence of the Drosophila Melanogaster
Genome." *Genome Research* 25 (3): 445–58.
Huang, Ying, Beifang Niu, Ying Gao, Limin Fu, and Weizhong Li. 2010. "CD-
HIT Suite: A Web Server for Clustering and Comparing Biological
Sequences." *Bioinformatics* 26 (5): 680–82.
Hu, Jialu, Yan Zheng, and Xuequn Shang. 2018. "MiteFinderII: A Novel Tool
to Identify Miniature Inverted-Repeat Transposable Elements Hidden
in Eukaryotic Genomes." *BMC Medical Genomics* 11 (Suppl 5): 101.
Jebb, David, Zixia Huang, Martin Pippel, Graham M. Hughes, Ksenia
Lavrichenko, Paolo Devanna, Sylke Winkler, et al. 2020. "Six
Reference-Quality Genomes Reveal Evolution of Bat Adaptations."
Nature 583 (7817): 578–84.
Ji, Yanzhu, and J. Andrew DeWoody. 2016. "Genomic Landscape of Long
Terminal Repeat Retrotransposons (LTR-RTs) and Solo LTRs as Shaped
by Ectopic Recombination in Chicken and Zebra Finch." *Journal of*
Molecular Evolution 82 (6): 251–63.
Jurka, J., V. V. Kapitonov, A. Pavlicek, P. Klonowski, O. Kohany, and J.
Walichiewicz. 2005. "Repbase Update, a Database of Eukaryotic
Repetitive Elements." *Cytogenetic and Genome Research* 110 (1-4):
462–67.
Kapitonov, Vladimir V., and Jerzy Jurka. 2008. "A Universal Classification of
Eukaryotic Transposable Elements Implemented in Repbase." *Nature*
Reviews. Genetics 9 (5): 411–12; author reply 414.
Kapitonov, V. V., and J. Jurka. 2001. "Rolling-Circle Transposons in

Eukaryotes." *Proceedings of the National Academy of Sciences of the United States of America* 98 (15): 8714–19.

Kohany, Oleksiy, Andrew J. Gentles, Lukasz Hankus, and Jerzy Jurka. 2006. "Annotation, Submission and Screening of Repetitive Elements in Repbase: RepbaseSubmitter and Censor." *BMC Bioinformatics* 7 (October):474.

Kojima, Kenji K. 2020. "Structural and Sequence Diversity of Eukaryotic Transposable Elements." *Genes & Genetic Systems* 94 (6): 233–52.

Larsson, Anders. 2014. "AliView: A Fast and Lightweight Alignment Viewer and Editor for Large Datasets." *Bioinformatics* 30 (22): 3276–78.

Lerat, E. 2010. "Identifying Repeats and Transposable Elements in Sequenced Genomes: How to Find Your Way through the Dense Forest of Programs." *Heredity* 104 (6): 520–33.

Li, Weizhong, and Adam Godzik. 2006. "Cd-Hit: A Fast Program for Clustering and Comparing Large Sets of Protein or Nucleotide Sequences." *Bioinformatics* 22 (13): 1658–59.

Louha, Swarnali, David A. Ray, Kevin Winker, and Travis C. Glenn. 2020. "A High-Quality Genome Assembly of the North American Song Sparrow, Melospiza Melodia." *G3* 10 (4): 1159–66.

Lu, Shennan, Jiyao Wang, Farideh Chitsaz, Myra K. Derbyshire, Renata C. Geer, Noreen R. Gonzales, Marc Gwadz, et al. 2020. "CDD/SPARCLE: The Conserved Domain Database in 2020." *Nucleic Acids Research* 48 (D1): D265–68.

Makałowski, Wojciech, Valer Gotea, Amit Pande, and Izabela Makałowska. 2019. "Transposable Elements: Classification, Identification, and Their Use as a Tool for Comparative Genomics." *Methods in Molecular Biology (Clifton, N.J.)* 1910:177–207.

Marchler-Bauer, Aron, Myra K. Derbyshire, Noreen R. Gonzales, Shennan Lu, Farideh Chitsaz, Lewis Y. Geer, Renata C. Geer, et al. 2015. "CDD: NCBI's Conserved Domain Database." *Nucleic Acids Research* 43 (Database issue): D222–26.

Marino, Alba, Eliette L. Reboud, Emmanuelle Chevalier, Marie-Ka Tilak, Jorge Contreras-Garduño, Benoit Nabholz, and Fabien L. Condamine. 2023. "Genomics of the Relict Species Baronia Brevicornis Sheds Light on Its Demographic History and Genome Size Evolution across Swallowtail Butterflies." *G3 (Bethesda, Md.)* 13 (12): jkad239.

Martelossi, Jacopo, Filippo Nicolini, Simone Subacchi, Daniela Pasquale, Fabrizio Ghiselli, and Andrea Luchetti. 2023. "Multiple and Diversified Transposon Lineages Contribute to Early and Recent Bivalve Genome Evolution." *BMC Biology* 21 (1): 145.

McDonald, John F., I. Bernard Weinstein, and Michael E. Lambert, eds. 1988. *Eukaryotic Transposable Elements as Mutagenic Agents*. Banbury Report S., v. 30. New York, NY: Cold Spring Harbor Laboratory Press.

Melo, Elverson Soares de, and Gabriel Luz Wallau. 2020. "Mosquito Genomes Are Frequently Invaded by Transposable Elements through Horizontal Transfer." *PLoS Genetics* 16 (11): e1008946.

Mills, Ryan E., E. Andrew Bennett, Rebecca C. Iskow, and Scott E. Devine. 2007. "Which Transposable Elements Are Active in the Human

Genome?" *Trends in Genetics: TIG* 23 (4): 183–91.

Neumann, Pavel, Petr Novák, Nina Hoštáková, and Jiří Macas. 2019. "Systematic Survey of Plant LTR-Retrotransposons Elucidates Phylogenetic Relationships of Their Polyprotein Domains and Provides a Reference for Element Classification." *Mobile DNA* 10 (1): 1.

Nishimura, Darryl. 2000. "RepeatMasker." *Biotech Software & Internet Report* 1 (1-2): 36–39.

Orozco-Arias, Simon, Mariana S. Candamil-Cortes, Paula A. Jaimes, Estiven Valencia-Castrillon, Reinel Tabares-Soto, Gustavo Isaza, and Romain Guyot. 2022. "Automatic Curation of LTR Retrotransposon Libraries from Plant Genomes through Machine Learning." *Journal of Integrative Bioinformatics* 19 (3). https://doi.org/10.1515/jib-2021-0036.

Orozco-Arias, Simon, Pío Sierra, Richard Durbin, and Josefa González. 2023. "MCHelper Automatically Curates Transposable Element Libraries across Eukaryotic Species." *bioRxiv*. https://doi.org/10.1101/2023.10.17.562682.

Orozco-Arias, S., P. Sierra, R. Durbin, and J. González. 2023. "MCHelper Automatically Curates Transposable Element Libraries across Species." https://europepmc.org/article/ppr/ppr745508.

Osmanski, Austin B., Nicole S. Paulat, Jenny Korstian, Jenna R. Grimshaw, Michaela Halsey, Kevin A. M. Sullivan, Diana D. Moreno-Santillán, et al. 2023. "Insights into Mammalian TE Diversity through the Curation of 248 Genome Assemblies." *Science* 380 (6643): eabn1430.

Ou, Shujun, and Ning Jiang. 2018. "LTR_retriever: A Highly Accurate and Sensitive Program for Identification of Long Terminal Repeat Retrotransposons." *Plant Physiology* 176 (2): 1410–22.

Ou, Shujun, Weija Su, Yi Liao, Kapeel Chougule, Jireh R. A. Agda, Adam J. Hellinga, Carlos Santiago Blanco Lugo, et al. 2019. "Benchmarking Transposable Element Annotation Methods for Creation of a Streamlined, Comprehensive Pipeline." *Genome Biology* 20 (1): 275.

Palatini, Umberto, Reem A. Masri, Luciano V. Cosme, Sergey Koren, Françoise Thibaud-Nissen, James K. Biedler, Flavia Krsticevic, et al. 2020. "Improved Reference Genome of the Arboviral Vector Aedes Albopictus." *Genome Biology* 21 (1): 215.

Peona, Valentina, Mozes P. K. Blom, Luohao Xu, Reto Burri, Shawn Sullivan, Ignas Bunikis, Ivan Liachko, et al. 2021. "Identifying the Causes and Consequences of Assembly Gaps Using a Multiplatform Genome Assembly of a Bird-of-Paradise." *Molecular Ecology Resources* 21 (1): 263–86.

Platt, Roy N., 2nd, Laura Blanco-Berdugo, and David A. Ray. 2016. "Accurate Transposable Element Annotation Is Vital When Analyzing New Genome Assemblies." *Genome Biology and Evolution* 8 (2): 403–10.

Rafii, Zafar, and Bryan Pardo. 2013. "REpeating Pattern Extraction Technique (REPET): A Simple Method for Music/Voice Separation." *IEEE Transactions on Audio, Speech, and Language Processing* 21 (1): 73–84.

Rech, Gabriel E., Santiago Radío, Sara Guirao-Rico, Laura Aguilera, Vivien Horvath, Llewellyn Green, Hannah Lindstadt, Véronique Jamilloux,

Hadi Quesneville, and Josefa González. 2022. "Population-Scale Long-
Read Sequencing Uncovers Transposable Elements Associated with
Gene Expression Variation and Adaptive Signatures in Drosophila."
Nature Communications 13 (1): 1948.
Rodriguez, Matias, and Wojciech Makałowski. 2022. "Software Evaluation
for de Novo Detection of Transposons." *Mobile DNA* 13 (1): 14.
Sievers, Fabian, and Desmond G. Higgins. 2014. "Clustal Omega, Accurate
Alignment of Very Large Numbers of Sequences." *Methods in*
Molecular Biology 1079:105–16.
Smit, A., R. Hubley, and P. Green. 2015. "RepeatMasker Open-4.0. 2013-
2015http." *Repeatmasker. Org*.
Storer, Jessica, Robert Hubley, Jeb Rosen, Travis J. Wheeler, and Arian F.
Smit. 2021. "The Dfam Community Resource of Transposable Element
Families, Sequence Models, and Genome Annotations." *Mobile DNA* 12
(1): 2.
Storer, Jessica M., Robert Hubley, Jeb Rosen, and Arian F. A. Smit. 2021.
"Curation Guidelines for de Novo Generated Transposable Element
Families." *Current Protocols* 1 (6): e154.
Wells, Jonathan N., and Cédric Feschotte. 2020. "A Field Guide to
Eukaryotic Transposable Elements." *Annual Review of Genetics* 54
(November):539–61.
Wicker, Thomas, François Sabot, Aurélie Hua-Van, Jeffrey L. Bennetzen,
Pierre Capy, Boulos Chalhoub, Andrew Flavell, et al. 2007. "A Unified
Classification System for Eukaryotic Transposable Elements." *Nature*
Reviews. Genetics 8 (12): 973–82.
Ye, Congting, Guoli Ji, and Chun Liang. 2016. "detectMITE: A Novel
Approach to Detect Miniature Inverted Repeat Transposable Elements
in Genomes." *Scientific Reports* 6 (January):19688.
You, F., S. Cloutier, Yunfeng Shan, and R. Ragupathy. n.d. "Ltr Annotator:
Automated Identification and Annotation of Ltr Retrotransposons in
Plant Genomes." *International Journal of*.
https://doi.org/10.17706/ijbbb.2015.5.3.165-174.

Hoskins, Roger A., et al. "The Release 6 reference sequence of the

Drosophila melanogaster genome." Genome research 25.3 (2015):

445-458.

Orozco-Arias, Simon, et al. "Automatic curation of LTR retrotransposon

libraries from plant genomes through machine learning." Journal of

Integrative Bioinformatics 19.3 (2022): 20210036.

Wong, Wai Yee, and Oleg Simakov. "RepeatCraft: a meta-pipeline for

repetitive element de-fragmentation and annotation." Bioinformatics

35.6 (2019): 1051-1052.