Larger larval sea lamprey (*Petromyzon marinus)* have longer survival times when exposed

to the lampricide 3-trifluoromethyl-4-nitrophenol

Allison M. Nalesnik[a*], Emily L. Martin[a], Ian S. Kovacs[a], Connor S. Johnson[a], Emma I. Carroll[a],

Aaron Jubar[b], William Hemstrom[a], Michael P. Wilkie[c], Erin S. Dunlop[d], Maria S. Sepulveda[e,f],

Nicholas S. Johnson[g], Mark R. Christie[a,e]

a. Department of Biological Sciences, Purdue University, 915 W. State St., West Lafayette, Indiana

47907-2054 USA

b. U.S. Fish and Wildlife Service, Ludington Biological Station, 5050 Commerce Drive, Ludington, MI

49431, USA

c. Department of Biology and Laurier Institute for Water Science, Wilfrid Laurier University, Waterloo,

ON, N2L 3C5, Canada

d. Aquatic Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry, c/o

Trent University, 2140 East Bank Drive, Peterborough, ON, K9L 0G2, Canada

e. Department of Forestry and Natural Resources, Purdue University, 715 W. State St., West Lafayette,

Indiana 47907-2054 USA

f. Sustainability Research Centre and Department Ecology and Biodiversity, Faculty of Life Sciences,

Universidad Andres Bello, Santiago, Chile

g. U.S. Geological Survey, Great Lakes Science Center, Hammond Bay Biological Station, 11188 Ray

Road, Millersburg, MI 49759, USA

*corresponding author: Allison Nalesnik; Department of Biological Sciences, Purdue University; 715 W.

State St., West Lafayette, Indiana 47907; analesni@purdue.edu, 301-706-0594

running title: Larval sea lamprey size and TFM survival

Abstract

Invasive sea lamprey (*Petromyzon marinus*) in the Laurentian Great Lakes have negatively
impacted ecological and economically important fishes for nearly a century. To mitigate these
negative effects, the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) is applied annually to
Great Lakes tributaries to kill larval lampreys before they become juveniles, out-migrate to the
lakes, and parasitize other fishes. It has been hypothesized that larval size (e.g., mass, length,
condition) may affect survival time in response to TFM. To test this hypothesis, we conducted an
experiment with 8,611 larvae across four temporal replicates, in which TFM concentrations like
those used in present-day stream treatments were applied for up to 18 hours. We found a
significant, positive correlation between the total length and mass of larval sea lamprey and their
survival times to TFM. Specifically, there was a 4.56 minute increase in survival time for each
one centimeter increase in total length after accounting for the effect of mass. We also identified
a phenotypic trade-off where the survival time of heavier larvae decreased with an increase in
total length and vice versa. The changes in total length and mass of larval sea lamprey stored in
ethanol for four months was also quantified. The observation that five larvae survived well past
the 12-hour time window of a typical TFM field treatment highlights the need for continuous
monitoring and the development of new control strategies to ensure the continued effective
management of this invasive species.

keywords: ethanol storage, invasive species, lampricide, larval size, sea lamprey

Introduction

Sea lamprey (*Petromyzon marinus*) are hematophagous, non-host specific parasites that are native to the North Atlantic Ocean with populations along the coasts of Europe and North America. However, sea lamprey are considered an invasive pest in the Laurentian Great Lakes (Hansen et al., 2016) where they have contributed to substantial mortality and population declines in ecologically and economically important fishes (Coble et al., 1990; Kitchell, 1990). Sea lamprey were first documented in Lake Ontario in the late 1800s, after which they rapidly spread to Lake Erie following modifications to the Welland Canal in the early 1900s (Eshenroder 2014; Docker et al., 2021). By the late 1930s, sea lamprey had spread to Lake Huron, Lake Michigan, and Lake Superior (Christie & Goddard, 2003). In the novel environment of the Great Lakes, the typically anadromous sea lamprey acclimated to life-long residence in an entirely freshwater environment and developed smaller body sizes both at the time of metamorphosis and as adults in comparison to individuals from their native range (Hansen et al., 2016). These changes may be, at least in part, the result of a response to selection in genes responsible for growth, reproduction, and bioenergetics (Yin et al., 2021).

Early control methods were initiated in the 1950s and relied mostly on engineered structures such as physical and electrical barriers and physical traps to prevent adult sea lamprey from accessing reproductive habitat (Hrodey et al., 2021; Smith & Tibbles, 1980), but the devices were labor intensive to operate and blocked valued fishes (McLaughlin et al., 2007). Lampricides were developed as an alternative approach to population control; over 4,000 organic candidate chemicals were tested at the Hammond Bay Biological Station (HBBS, Millersburg, MI), of which ten compounds were found to have greater toxicity for larval sea lampreys than other fishes at comparatively lower concentrations (Applegate et al., 1957, 1958, 1966). The chemical compound 3-trifluoromethyl-4-nitrophenol (hereafter TFM) was identified as the most effective agent and was found to be practical for field application, bearing relatively low toxicity to other species at low concentrations and a low manufacturing cost (Applegate et al., 1961; Smith & Tibbles, 1980).

TFM induces mortality by interfering with the oxygen-dependent production of ATP within cells and tissues (Niblett and Ballantyne 1976; Birceanu et al. 2021, Huerta et al. 2020). It is thought to act as a protonophore that disrupts the proton gradient required by mitochondrial

enzyme ATP synthase to couple ADP with a phosphate group (*i.e.,* ATP synthesis) (Yin et al.

2021; Wilkie et al., 2019; Birceanu et al. 2021). While larval lamprey can use glycogen and

phosphocreatine stores as temporary alternate anaerobic energy sources, these energy stores are

finite and their eventual depletion in the presence of TFM results in larval mortality. Larval sea

lamprey are hypothesized to be particularly susceptible to TFM given their basal location on the

vertebrate phylogeny and limited ability to metabolize TFM through its molecular conjugation to

form TFM glucuronide, a less toxic TFM derivative that is readily excreted (Bussy et al., 2018;

Kane et al. 1994; Lawrence et al., 2022; Lech & Statham, 1975). Since control methods were

implemented in the late 1950s, sea lamprey populations have decreased by approximately 90%

from their historically high abundances, aiding the recovery of some native fish populations such

as lake trout (*Salvelinus namaycush;* Marsden & Siefkes, 2019). The continued control of sea

lamprey in the Great Lakes hinges on effective stream treatments, which can be influenced by

local environmental conditions at treatment sites. Because TFM bioavailability is higher in low

pH and low alkalinity waters, control personnel can use this relationship to modify TFM

treatment concentrations accordingly (Wilkie et al., 2021). Larval TFM sensitivities may also be

influenced by water temperature (Hlina et al., 2021; Muhametsafina et al., 2019), time of year

(Scholefield et al., 2008), and possibly liver glycogen content (although direct mechanisms are

not well described), such that treatment efficacy can depend on water conditions and larval body

conditions (Schueller et al., 2024). Despite TFM being instrumental to controlling sea lamprey

populations, specific quantitative relationships between larval lamprey body size and toxicant

susceptibility remain largely uncharacterized, although TFM uptake is known to be inversely

related to body mass (Tessier et al., 2018). In the present study, we examine how total length,

mass, and their interaction contribute to TFM susceptibility. A better understanding of these

effects is critical for determining appropriate TFM dosages and durations for field applications

over the next few decades, which can be adjusted based on larval size surveys to ensure

continued treatment effectiveness and population control.

Here, we empirically test the relationship between body size and survival in TFM while

holding pH, season, temperature, and source of larvae constant. We predicted that because

oxygen consumption and TFM uptake per unit of mass are greater in smaller mass larvae

(Tessier et al., 2018), larger animals would be more likely to survive for prolonged periods of

time in TFM. To test this hypothesis, we exposed 8,611 larval sea lampreys to TFM

concentrations designed to mimic treatments in the field and measured survival after each hour

of exposure. We measured the total length and mass of each larva and quantified the

relationships between these morphometric measures and survival time. We additionally include

findings from a completely independent, smaller scale, experiment in which similar comparisons

were made between larval lamprey body size and TFM survival times. We also used this

opportunity to describe the relationship between changes in body size in larval sea lamprey and

their storage time in ethanol. To do so, we quantified changes in total length, mass, and condition

of larval sea lamprey over a four-month period.

Materials and Methods

Sample collection

All larvae used in this experiment were collected by crews from the United States Fish and

Wildlife Service and United States Geological Survey over a period of eight days during July

2022 (**Table S1**). Larvae were obtained from the Muskegon River, a tributary to Lake Michigan,

MI, as it harbors one of the largest populations of spawning adult and larval sea lamprey within

the Great Lakes basin (Lavis et al., 2003; Slade et al., 2003). Prior to collecting the larval

lamprey for the present study, the Muskegon River had been treated with TFM in 2017 and 2019.

Given this three-year interval and the growth rate of larvae (Dawson et al., 2021), the larval

lamprey used in this study are unlikely to have been previously exposed to TFM. Pulsed-DC

backpack electrofishing units were used to capture larval sea lamprey from four reaches of the

Muskegon River (**Table S1**). To minimize the effects of seasonality, we condensed collection

times as much as possible. Most larvae collected were between 60-100 mm (range: 40-140 mm;

Figure 1).

To minimize stress experienced by the larvae due to handling and transportation, larvae

were transferred to acclimation tanks at U.S. Geological Survey, Hammond Bay Biological

Station (HBBS; Millersburg, MI) within one day of collection in individually labeled zippered

mesh bags (Tutudow, 40 cm x 50 cm, polyester, 250 µm mesh). Up to 50 larvae were transported

in each bag alongside two liters of sand from the Muskegon River. The sand grains were larger

than the mesh size of the bag such that the sand added to each bag remained in place during

transit. Bags containing larvae and sand were placed in 50 L aerated tanks maintained at a

temperature of 15 $^{\circ}$C overnight (relatively the temperature of the Muskegon River). The day after

capture, larvae held in mesh bags were transported to HBBS in 200 L aerated tanks maintained at
15 $^{\circ}$C.

Upon arrival at HBBS, larvae in each bag were separated from the sand by emptying the
contents of each bag onto a 750 µm screen and using a slow stream of water from Lake Huron at
15 $^{\circ}$C to wash the sand through the screen, leaving the larvae on top of the screen. Larvae from
each specific bag were next moved into a mesh container (Ikea Fyllen Laundry Basket, 79 L
volume, 50 cm high and 45 cm diameter, 500 µm mesh size) and placed inside an aerated 1000 L
tank supplied with water from Lake Huron at ambient temp (16-18 $^{\circ}$C; water was pumped from a
depth of 25 m in Lake Huron). Within each holding tank, six mesh containers were used to hold
at most 282 larvae each. Sand with a 250 µm grain size was added to each mesh container to
achieve a layer of sand five cm thick for the larval lamprey to burrow into. Capture dates and
locations, the dates of transportation, and the locations of acclimation at HBBS were tracked for
each batch of larvae (**Table S1**). All larvae were acclimated at HBBS within the mesh container
they were assigned for two weeks prior to exposure to TFM, during which larvae were fed fresh
baker's dry yeast at a rate of 1 g of yeast per larvae two times per week (Hanson et al., 1974;
Hume et al., 2024).

TFM exposure experiments
Our TFM target concentrations were determined by preliminary range-finding trials to determine
the lethal concentration of TFM required to kill 90% of larval sea lamprey over 12 hours (12-h
LC$_{90}$), based on the local water conditions and the tank systems to be used, following the
standard guidelines for acute toxicity tests in fish (OECD, 2019). All protocols involving the
handling of larvae at the Hammond Bay Biological Station were carried out in accordance with
United States federal guidelines for care and use of animals in accordance with American
Fisheries Society, ARRIVE guidelines, and the U.S. Animal Welfare Act of 1970 (Jenkins et al.,
2014). We aimed to use a 12-hour exposure period to approximate the total duration of TFM
treatments in the field (Middaugh et al., 2014). Our study consisted of four replicate TFM
exposure experiments (replicated two days apart) each using an identical design to measure the
survival of larval sea lamprey exposed to TFM. Each replicate experiment was conducted in two
1500 L tanks containing 624 L of water supplied from Lake Huron. Each tank was dosed with
between 104 and 112 mL of a 15,000 mg/L TFM stock solution at least 12 hours prior to

introducing the larvae to reach target concentrations of 2.3 to 2.5 mg/L TFM. No additional

water was added to these tanks after TFM was added. Water from Lake Huron did not vary

substantially in temperature (μ = mean ± standard deviation; μ = 17.6 ± 0.5 ^{o}C) or alkalinity over

the four replicant trials (μ = 85.6 ± 0.6 mg/L CaCO$_3$; **Table S2**). However, in some cases, there

were minor fluctuations in pH during treatments, as discussed below (μ = 8.1 ± 0.3; **Table S2**).

Each tank was fitted with a 15 cm long aeration stone at the center to facilitate uniform

distribution of TFM throughout the tank and to keep dissolved oxygen (DO) concentrations

higher than 7.6 mg/L (μ = 9.0 ± 0.6 mg/L; **Table S2**). The air stone was set to turbulently aerate

the water for at least 30 minutes after initial dosing and reduced to a moderate level for

subsequent mixing and aeration. The TFM concentration was measured thirty minutes after

initial dosing the day before each experiment began and at least 12 hours later on the day of the

experiment to determine if additional adjustments were required to maintain consistent exposure

conditions between our replicates.

Between 55 and 282 larval sea lampreys were kept in each of six mesh bags during

acclimation (**Table S1**) which were transferred to the experimental tank at the start of each

experiment. When the bags in the acclimation tanks were lifted and twisted, the 250µm sand fell

through the 500µm mesh containers leaving only larval sea lamprey and no sand, eliminating the

stress and potential injuries to larvae if each larva was dug from the sand. The larvae in each bag

were then lowered into the experimental tank nearby. Each bag was assessed for mortality in the

order in which the bags went into the experimental tanks.

Throughout the duration of each experiment, several parameters were monitored on an

hourly basis. Fish mortality within each bag was assessed to determine survival time in TFM. To

assess mortality, teams of two people transferred the larvae from one mesh bag into a double-net

apparatus (**SI video**). Fish that were alive with swimming ability intact swam through the

apparatus and into an empty mesh bag held below. Dead or impaired fish were unable to swim

through the apparatus and required visual inspection to determine status (alive or dead). Larvae

were inspected in a shallow plastic tray filled with the same TFM solution from the respective

experimental tank for up to three minutes. Larvae were declared dead after the cessation of

branchiopore (gill pore) movement and unresponsiveness to tail pinching, then stored in

individually labeled bottles of 95% ethanol. Impaired fish were returned to their original

experiment tank and into the same mesh bag as individuals that swam through the apparatus. In

addition to recording larval sea lamprey mortality at hourly intervals, DO (YSI Pro 20), pH

(Fisherbrand Accumet AP110 Portable), salinity (YSI Pro 20), TFM concentration, water and air

temperature were measured hourly. TFM concentration was measured using a Genesys 6

spectrophotometer (Thermo Electron Corporation, MA, USA) set to a wavelength of 395, using

analytical standards prepared in the same water, based on Standard Operating Procedure No.

LAB 431.0, Hammond Bay Biological Station, Millersburg, MI (GLFC.

https://glfc.org/sop.php). Analytical standards were prepared from TFM (CAS Number 88-30-2;

Millipore-Sigma, St. Louis, MO). Water and air temperature were measured via a laboratory

liquid-in-glass thermometer. The experiments varied in duration from five to eighteen hours until

the last larva had died. At the conclusion of all experiments, preserved larvae were transported

from HBBS to Purdue University (West Lafayette, IN) where the total length (mm) and mass (g)

was measured for all lamprey (n = 8,611 larvae). All experiment samples were stored in 95%

ethanol for at least four months prior to measuring their total length and mass.

To determine what physiological variables impact lamprey survival time in TFM, we fit a

linear mixed effect model to predict survival time, using total length (mm), mass (g), and their

interaction as fixed effects and a nested random effect of experiment in tank in bag as a covariate

using R 4.3.2 (R Core Team, 2023) and the R package lmerTest (Kuznetsova et al., 2017). Based

on diagnostic assessments, the assumptions for a linear mixed effects model were largely met. A

Q-Q plot of the residuals indicated that the data follow a normal distribution, with only minor

deviations observed. Additionally, the residuals versus fitted values plot showed no discernible

patterns, with most points randomly scattered around the zero line. While five data points

exhibited slight deviations from the expected trend, given the dataset's size, their impact on the

model was minimal. Therefore, the assumptions of normality, linearity, and homoscedasticity

were reasonably satisfied. We compared this model to five others with different covariates to

determine which model best explained individual variance in survival (see **Table S3** for

additional models). Condition factor (CF) was calculated as previously outlined for Pacific sea

lamprey (Lampman et al., 2016) as $CF = \frac{m*10^5}{l^{2.6}}$ where m = mass (g) and l = length (mm).

Independent test of the effect of body size on survival to TFM exposure

A second, independent experiment that was much smaller in scale (n = 48 larvae) was also

performed (See **SI Methods** for details). All experiments were approved by the Wilfrid Laurier

Animal Care Committee (AUP R10004) and followed Canadian Council of Animal Care

guidelines. We include the results here because although the larvae were collected from different

tributaries and the experiment was performed in a different laboratory, the results between the

two different exposure experiments are congruent.

Long-term effects of ethanol storage

To assess the effect of long-term ethanol storage on larval sea lamprey morphometrics, we

systematically measured the total length and mass of two different sets of larval sea lamprey over

the course of four months (hereafter storage experiments A and B). None of the larvae used to

quantify changes in body size and mass due to ethanol storage were included in the TFM

exposure experiments; instead, these larvae were used to determine the effect of measuring the

8,611 experimental larvae after a storage period of up to eight months since it was logistically

impractical to measure all individuals during the large TFM exposure experiments. Storage

experiment A: In July, prior to the large TFM survival experiment described above, 554 larval

sea lamprey were transported to Purdue University's Aquaculture Research Laboratory (ARL;

West Lafayette, IN). Sea lamprey were exposed to TFM concentrations ranging from 2.0 mg/L

to 4.0 mg/L in groups ranging from 10 to 25 individuals. If sea lamprey remained alive beyond

36 hours of TFM exposure, they were euthanized with 250 mg/L tricaine methanesulfonate (MS-

222) buffered to pH 7 with sodium bicarbonate. Total length (mm) and mass (g) were measured

immediately after death and each sample was individually stored in a 50 mL conical tube

containing 40 mL of 95% non-denatured ethanol. Purdue University Institutional Animal Care

and Use Committee reviewed experimental and handling procedures for this study, but did not

require animal care protocols for the larval sea lamprey. Storage experiment B: At HBBS, 110

larval sea lampreys that remained in the acclimation tank after the completion of the large TFM

survival experiments were euthanized with MS-222 using the same procedure described above.

We measured the total length (mm) and mass (g) of each larva immediately after death, then

stored every individual in 15 mL conical tubes containing 12 mL of 95% ethanol. Thereafter,

each larval sea lamprey (554 from ARL, 110 from HBBS) was measured for body mass and total

length systematically over a period of 120 days (see **SI Methods** for details on measurement

procedures). We calculated condition (CF) using the same equation as above.

Given that larvae become smaller over time when stored in ethanol, we fit an additional

model to predict the initial size of an individual from a measurement taken after extended storage

in ethanol. We first randomly sampled our dataset of Experiment A and Experiment B

individuals (664 larvae) such that only one sample per individual was used (since any models

containing individual ID as a covariate could not be used to make predictions for new

individuals). We then used the *nls* function in R to fit a nonlinear regression model to determine

the decay rate which best fit our observed data using an exponential decay function:

$$l_0 = \frac{l_t}{e^{-bt}}$$

where l_0 is initial length, l_t is the length at measurement time t, and b is the rate of decay. We

evaluated the ability of this model to predict the initial size of a larval sea lamprey prior to

storage using a hold-one-out cross validation approach (see **Tables S4** and **S5**; Wong, 2015).

Using these adjusted values, we also re-analyzed the data from our large TFM exposure and

body size experiments using the exact statistical models described above.

Results

Results from large-scale TFM exposure experiments

There was a highly allometric relationship between larval total length and larval mass (**Figure**

1). Measured TFM concentrations during exposure averaged 2.52 ± 0.08 mg/L (n = 87) across

our replicates (**Table S2**). The 8,611 lamprey used in our four experimental replicates had total

lengths between 43.0 and 137.5 mm (μ = 85.39 mm; **Figure 2A**; **Figure S1A**), masses between

0.074 and 4.529 g (μ = 0.869 g; **Figure 2B**; **Figure S1B**), and condition factors between 0.0880

and 1.96 (μ = 0.761; **Figure 2C**; **Figure S1C**). The linear mixed effect model including total

length, mass, and their interaction was significant (p value < 0.001) and explained a large portion

of the variation in survival time in TFM (conditional R^2 = 0.7142; extended results in **Table S3**).

Total length was significantly and positively associated with survival time in TFM (**Figure 2A**)

as described by the following equation: y = 0.0321x + 1.88 (R^2 = 0.693; p value < $2 \bullet 10^{-16}$). Mass

was also significantly and positively associated with survival time in TFM (**Figure 2B**) as

described by the following equation: y = 0.961x + 3.79 (R^2 = 0.711; p value < $2 \bullet 10^{-16}$). There

was a 4.56 minute [95% CI: 4.38 - 6.54] increase in survival time for each 10 mm increase in

total length after accounting for the effect of mass. There was an 86.4 minute [95% CI: 67.8-

105] increase in survival time for each one gram increase in total mass after accounting for the

effect of total length. The effect of the interaction between total length and mass on survival time

was negative (*p* value < 0.001): for every 10 mm increase in total length, the impact of

increasing by 10 g of mass reduced survival time by 3.40 minutes [95% CI: 4.86 - 1.92] and vice

versa (*i.e.,* as fish get longer, the benefit to being heavier decreases, and vice versa). There was

no significant relationship between condition factor and survival time (**Figure 2C;** *p* value =

0.940). Five individuals survived beyond 12 hours of TFM exposure (**Figure 3; Figure S3**).

Results from smaller-scale TFM exposure experiments

Measured TFM concentrations during exposure averaged 4.9 ± 0.5 mg/L (n =16; **Table S6**).

After 16 h of TFM exposure, 43 of the 48 animals had suffered mortality. When including the

five surviving larvae, there was a significant relationship between body length and survival time

for larval sea lamprey, with an average increase in survival time of 58.92 minutes per one mm

increase in body length (R^2 = 0.2, *p* value = 0.002; see **Figure 4A**).The relationship between

body mass and survival time was also significant, with an increase in survival time of 197.4

minutes per one g increase in mass (R^2 = 0.238, *p* value = 0.0005; see **Figure 4B**). There was no

significant relationship between condition factor and survival time (**Figure 4C**). The effect of the

interaction between total length and mass on survival time was positive although not significant

(*p* value = 0.4008).

Effects of ethanol storage on body size

HBBS and ARL larvae both declined in length and mass over the 120 day measurement period

(**Figure S2**). For example, the total length of HBBS larvae (n = 110) initially ranged from 63.5

to 127.5 mm (μ = 95.1 mm; **Figure 5A**). The last measured total length (day 120) of HBBS

larvae ranged from 56.0 to 119.5 mm (μ = 88.6 mm; **Figure 5A**). The mass of HBBS larvae (n =

110) initially ranged from 0.43 to 2.72 g (μ = 1.27 g; **Figure 5B**). The last measured mass of

HBBS larvae ranged from 0.185 to 1.906 g (μ = 0.83 g; **Figure 5B**).

The nonlinear least squares model generated using the combined sample set (n = 664; site

was included as a fixed variable to account for differences in ethanol storage volume) yielded a

total length decay estimate as $4.483 \cdot 10^{-4}$ mm/day (**Table S4**). The linear regression model

suggested a strong, positive relationship between predicted values and true values, with an

average change in predicted initial length of 1.035mm for each 1mm increase in observed length

post-storage after accounting for the effects of ethanol (t value = 13.91; p value < $2 \cdot 10^{-16}$; **Table**

S5). The model explained a very large proportion of the variability in the initial length (R^2 =

0.69). We used our model estimates to calculate initial total length, mass, and condition for our

large TFM exposure experiment larvae by dividing the measured value by e raised to the

negative estimate, multiplied by the number of days the larvae were stored in ethanol and found

nearly identical results (**Figure S4**).

Discussion

We describe a significant, positive relationship between the size of larval sea lamprey and their

survival time in TFM. Larger larval sea lamprey exhibit increased survival time in TFM,

highlighting that this variable, if accounted for during field treatments, could improve treatment

success. To conduct TFM treatments with the highest larval sea lamprey mortality, TFM

applicators and personnel can consider larval sizes within a treatment stream and may adjust

treatment duration accordingly. We also describe the predicted survival time for larvae at the

10th, 25th, 50th, 75th, and 90th total length and mass percentiles within our data (**Table S7**),

which can be used to justify the duration of treatment in selected streams based on surveyed

larval sizes. For instance, streams with an average larval size larger than our 90th percentile

would require at least an additional hour of treatment compared to streams with an average larval

size closer to our 10th percentile (Table S7). The observed correlation between sea lamprey size

and survival time in TFM suggests the presence of a mechanism by which larger individuals can

better withstand the toxic effects of the lampricide. Given that our linear mixed effect model

including total length, mass, and their interaction explained a very large percentage of the

variation in survival time (R^2 = 0.7142), we suggest that the response of sea lamprey larvae to

TFM is determined by larval size.

This conclusion is additionally supported by the independent, smaller scale study where

animals that survived for longer than the treatment time of a TFM exposure tended to be larger in

both mass and length. The smaller scale study, which involved only 48 larvae, reported a

significantly higher increase in survival time with respect to both total length and mass.

Specifically, it indicated an increase of 58.92 min per 1mm increase in length and 197.4 min per

1g increase in mass. In contrast, the larger scale study with 5,611 larvae, estimated much smaller

effects, predicting just 4.56 min per 1mm increase in total length and 86.4 min per 1 gram

increase in mass. The substantial discrepancy in these findings is likely attributed to the sample

size and the number of replicates. The larger study's much greater sample size provides a more

reliable estimate, reducing the likelihood of variability and statistical noise influencing the

results. Consequently, we place more confidence in the findings of the larger scale study, as the

increased sample size leads to higher precision and a more accurate reflection of the true

relationship between fish survival time and both length and mass.

In the large scale TFM exposure study, the extended duration of survival for five

individuals well beyond the typical treatment length of TFM in the field, that were neither the

longest nor the heaviest larvae, indicate that some larvae may be capable of surviving treatment

(**Figure 3; Figure S3**). This observation is consistent with underlying variation in the population

in terms of susceptibility to TFM and has implications for the potential for sea lamprey to evolve

resistance to TFM, a subject of management interest (Christie et al., 2019; Dunlop et al., 2021).

The development of resistance is a known consequence of repeated pesticide application (Délye

et al., 2013; Georghiou & Taylor, 1977; Hawkins et al., 2019) and given that TFM has been used

annually in freshwater streams in the Great Lakes Basin since 1958, resistance in sea lamprey is

conceivable and an urgent concern (Christie et al., 2019; Dunlop et al., 2021). Prior research has

suggested that a response to TFM-induced selection may already be underway in Great Lakes:

sea lamprey genomic and transcriptomic comparisons between treated and untreated populations

of larval sea lamprey suggest that an adaptive, genetic response to TFM may drive incipient

resistance (Yin et al., 2021). Sea lamprey may develop resistance to TFM through two broad

mechanisms: (1) physiological adaptations or (2) evolution of their behavior or life history

(Christie et al., 2019; Dunlop et al., 2018, 2021). Physiological resistance may involve the

development of molecular mechanisms to more efficiently detoxify or neutralize TFM or to

counteract the effects of the mitochondrial proton gradient; however, if larger individuals have a

selective advantage for surviving treatment, and large size (or a correlated trait like growth rate)

is heritable, then there is the potential for both physiological and life history resistance to TFM

via selection for increases in body size. For example, larger sea lamprey could store additional

glycogen, which would provide them with larger anaerobic energy reserves to generate ATP

using anaerobic glycolysis. Such features could enable individuals to out-survive a typical TFM

treatment. Although previous studies demonstrated no clear relationship between liver or muscle

glycogen reserves and TFM tolerance, (Muhametsafina et al., 2019; Hlina et al. 2021), a recent

study by Schueller et al. (2024) found a positive relationship between liver glycogen

concentrations and the minimum lethal concentration (MLC = 12 h-LC$^{99.9}$; Bills et al. 2003) of

TFM to larval sea lamprey using stream-side toxicity tests. Larval sea lamprey also demonstrate

a remarkable capacity to recover from TFM exposure when transferred to clean, TFM-free water,

restoring energy stores such as glycogen and phosphocreatine within a few hours (Clifford et al.

2012). Collectively, these studies suggest that there may be a physiological advantage to being

larger at the time of TFM exposure/treatments (Dunlop et al., 2021).

Contrary to our findings here, a handful of previous studies describe that energy stores

and body size have little impact on TFM sensitivity (Hlina et al., 2021; Muhametsafina et al.,

2019), and instead report that larvae exhibit differential TFM sensitivity based on season, which

may be a result of the seasonally warmer waters enhancing their capacity to detoxify TFM (Hlina

et al., 2021; Muhametsafina et al., 2019). Additionally, as pH and alkalinity of water increase,

larvae exhibit increased tolerance as TFM changes its ionization state (reviewed in Wilkie et al.,

2019), resulting in lower TFM uptake in more basic environments (Hlina et al., 2017). Since we

controlled for pH, alkalinity, and water temperature during the experiment and controlled for

seasonality by using larvae collected within eight days in July to assess the contribution of total

length and mass on survival time in TFM, additional experiments may be needed to better weigh

the contribution of environmental and physiological components associated with differential

survival in TFM (*e.g.*, temperature and alkalinity versus body mass and length).

In addition to quantifying the relationship between larval size and survival in TFM, we

also characterized the relationship between larval body size and length after storage in ethanol.

We provide an estimate of the decay rates of length and body mass, which should prove useful

when back calculating the actual sizes of sea lamprey larvae prior to preservation (as

demonstrated in Figure S4). By generating estimates for total length, mass, and condition, we

evaluated the usefulness of each parameter to make accurate predictions of initial state values.

The total length estimates generated from each sample set were of the same magnitude (on the

scale of 10^{-4}). Estimates from mass and condition were less accurate predictors of initial values,

as reflected by smaller adjusted R^2 values, when compared to those from total length. The best
performing nonlinear regression model was derived from our combined sample set and provides
a total length exponential decay estimate of $4.483 \cdot 10^{-4}$ mm/day (**Table S4**). Given that the model
derived from our combined sample set had high predictability, our estimate may be useful across
diverse larval sea lamprey collections. Our model gives researchers additional options, apart
from using formalin (Neave et al., 2006), for correcting for changes in body size after prolonged
storage in 95% ethanol.

Conclusion
Our study provides valuable insights into the relationship between larval sea lamprey size and
survival time in TFM-treated environments. These findings suggest that duration and TFM
dosage for routine TFM treatments within the Great Lakes basin could be improved if larval
length and mass can be taken into consideration. Additionally, studies that store larval sea
lamprey in 95% ethanol can now correct for storage time using a simple equation. We also
emphasize the need for further research to better understand the underlying mechanisms between
larger larval lamprey size and TFM survival to better inform sea lamprey management
throughout the Great Lakes.

Acknowledgments
We thank the USFWS team responsible for collecting sea lamprey larvae for this experiment:
Jordan Andrus, Philip Bennett, John Ewalt, Timothy Granger, Allen Keffer, David Keffer, Callie
Kopp, Daniel McGarry, Lily Smalstig, Jenna Tews, and Matt Lipps. We also thank the USGS
collection team and on-site personnel who provided hands-on assistance during the experiment:
David Coburn, Tyler Bruning, Ed Benzer, Zachary Nordstrom, Meg Spens, Brad Buechel. We
thank Emma Vieregge and Karen Slaght for their support doing water quality analysis and range-
finder assays. We also thank Robert Rode for his advice and assistance at the ARL, as well as
undergraduate support from Lilly Gorney, Leah Kemple, and Grace Bowman. Research
conducted at Wilfrid Laurier University was conducted by Lisa Sorensen and supported by a
Great Lakes Fishery Commission contract awarded to MP, R Bergstedt and D.G. McDonald
(2011). This research was funded by support to MRC, MSS, ESD, and NSJ from the Great Lakes
Fishery Commission (grant number 2022_CHR_541010, 2022). Any use of trade, firm, or

product names is for descriptive purposes only and does not imply endorsement by the U.S.

Government.

Conflict of interest

432 The authors declare that they have no conflict of interest.

Data archiving

434 All code to reproduce the analyses reported in this manuscript are available via GitHub at

435 https://github.com/ChristieLab/sea-lamprey-morphometrics. Data will be made available on

436 Dryad.

References

Applegate, V. C., Howell, J. H., Hall Jr, A. E., & Smith, M. A., 1957. Toxicity of 4,346
chemicals to larval lampreys and fishes (Special Scientific Report – Fisheries No. 207).
US Department of the Interior, Fish and Wildlife Service. 1-157.

Applegate, V. C., Howell, J. H., Moffett, J. W., & Smith, M. A., 1961. Use of 3-trifluormethyl-4-
nitrophenol as a selective sea lamprey larvicide. Technical Report 1. Great Lakes Fishery
Commission. 1–35. https://www.glfc.org/pubs/TechReports/Tr01.pdf

Applegate, V. C., Johnson, B. G. H., & Smith, M. A., 1966. The relation between molecular
structure and biological activity among mononitrophenols containing halogens. Technical
Report 11. Great Lakes Fishery Commission. 1-19.

Applegate, V. C., Howell, J. H., & Smith, M. A., 1958. Use of mononitrophenols containing
halogens as selective sea lamprey larvicides. Science, 127(3294), 336–338.

Bills, T. D., Boogaard, M. A., Johnson, D. A., Brege, D. C., Scholefield, R. J., Westman, R. W.,
& Stephens, B. E., 2003. Development of a pH/alkalinity treatment model for
applications of the lampricide TFM to streams tributary to the Great Lakes. Journal of
Great Lakes Research, 29, 510-520.

Birceanu, O., Tessier, L. R., Huerta, B., Li, W., McDonald, A., & Wilkie, M. P., 2021. At the
intersection between toxicology and physiology: What we have learned about sea
lampreys and bony fish physiology from studying the mode of action of
lampricides. Journal of Great Lakes Research, 47, S673-S689.

Birceanu, O., McClelland, G. B., Wang, Y. S., & Wilkie, M. P., 2009. Failure of ATP supply to
match ATP demand: The mechanism of toxicity of the lampricide, 3-trifluoromethyl-4-
nitrophenol (TFM), used to control sea lamprey (*Petromyzon marinus*) populations in the
Great Lakes. Aquatic toxicology, 94(4), 265-274.

Bussy, U., Chung-Davidson, Y.-W., Buchinger, T., Li, K., Smith, S. A., Jones, A. D., & Li, W.,
2018. Metabolism of a sea lamprey pesticide by fish liver enzymes part A: Identification
and synthesis of TFM metabolites. Analytical and Bioanalytical Chemistry, 410(6),
1749–1761.

Christie, G. C., & Goddard, C. I., 2003. Sea Lamprey International Symposium (SLIS II):
Advances in the integrated management of sea lamprey in the Great Lakes. Journal of
Great Lakes Research, 29, 1–14.

Christie, M. R., Sepúlveda, M. S., & Dunlop, E. S., 2019. Rapid resistance to pesticide control is
predicted to evolve in an invasive fish. Scientific Reports, 9(1), Article 1.

Clifford, A. M., Henry, M., Bergstedt, R., McDonald, D. G., Smits, A. S., & Wilkie, M. P., 2012.
Recovery of larval sea lampreys from short-term exposure to the pesticide 3-
trifluoromethyl-4-nitrophenol: implications for sea lamprey control in the Great
Lakes. Transactions of the American Fisheries Society, 141(6), 1697-1710.

Coble, D. W., Bruesewitz, R. E., Fratt, T. W., & Scheirer, J. W., 1990. Lake trout, sea lampreys,
and overfishing in the upper Great Lakes: A review and reanalysis. Transactions of the
American Fisheries Society, 119(6), 985–995.

Dawson, H. A., Higgins-Weier, C. E., Steeves, T. B., & Johnson, N. S., 2021. Estimating age
and growth of invasive sea lamprey: A review of approaches and investigation of a new
method. Journal of Great Lakes Research, 47, S570-S579.

Délye, C., Jasieniuk, M., & Le Corre, V., 2013. Deciphering the evolution of herbicide resistance
in weeds. Trends in Genetics, 29(11), 649–658.

Docker, M. F., Bravener, G. A., Garroway, C. J., Hrodey, P. J., Hume, J. B., Johnson, N. S.,
Lewandoski, S. A., Ogden, J. L., & Zollweg-Horan, E. C., 2021. A review of sea lamprey
dispersal and population structure in the Great Lakes and the implications for control.
Journal of Great Lakes Research, 47, S549–S569.

Dunlop, E. S., Christie, M. R., McLaughlin, R., & Steeves, T. B., 2021. Life history evolution of
sea lamprey is predicted to reduce the effectiveness of pesticide control. Journal of Great
Lakes Research, 47, S259–S271.

Dunlop, E. S., McLaughlin, R., Adams, J. V., Jones, M., Birceanu, O., Christie, M. R., Criger, L.
A., Hinderer, J. L. M., Hollingworth, R. M., Johnson, N. S., Lantz, S. R., Li, W., Miller,
J., Morrison, B. J., Mota-Sanchez, D., Muir, A., Sepúlveda, M. S., Steeves, T., Walter, L.,
… Wilkie, M. P., 2018. Rapid evolution meets invasive species control: The potential for
pesticide resistance in sea lamprey. Canadian Journal of Fisheries and Aquatic Sciences,
75(1), 152–168.

Eshenroder, R. L., 2014. The role of the Champlain Canal and Erie Canal as putative corridors
for colonization of Lake Champlain and Lake Ontario by sea lampreys. Transactions of
the American Fisheries Society, 143(3), 634-649.

Georghiou, G. P., & Taylor, C. E., 1977. Genetic and biological influences in the evolution of insecticide resistance. Journal of Economic Entomology, 70(3), 319–323.

Hansen, M. J., Madenjian, C. P., Slade, J. W., Steeves, T. B., Almeida, P. R., & Quintella, B. R., 2016. Population ecology of the sea lamprey (*Petromyzon marinus*) as an invasive species in the Laurentian Great Lakes and an imperiled species in Europe. Reviews in Fish Biology and Fisheries, 26(3), 509–535.

Hanson, L. H., King, E. L., Howell, J. H., & Smith, A. J., 1974. A culture method for sea lamprey larvae. The Progressive Fish-Culturist, 36(3), 122–128.

Hawkins, N. J., Bass, C., Dixon, A., & Neve, P., 2019. The evolutionary origins of pesticide resistance. Biological Reviews, 94(1), 135–155.

Hlina, B. L., Birceanu, O., Robinson, C. S., Dhiyebi, H., & Wilkie, M. P., 2021. The relationship between thermal physiology and lampricide sensitivity in larval sea lamprey (*Petromyzon marinus*). Journal of Great Lakes Research, 47, S272–S284.

Hlina, B. L., Tessier, L. R., & Wilkie, M. P., 2017. Effects of water pH on the uptake and elimination of the piscicide, 3-trifluoromethyl-4-nitrophenol (TFM), by larval sea lamprey. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology, 200, 9–16.

Hrodey, P. J., Lewandoski, S. A., Sullivan, W. P., Barber, J. M., Mann, K. A., Paudel, B., & Symbal, M. J., 2021. Evolution of the sea lamprey control barrier program: The importance of lowermost barriers. Journal of Great Lakes Research, 47, S285–S296.

Huerta, B., Chung-Davidson, Y. W., Bussy, U., Zhang, Y., Bazil, J. N., & Li, W., 2020. Sea lamprey cardiac mitochondrial bioenergetics after exposure to TFM and its metabolites. Aquatic Toxicology, 219, 105380.

Hume, J. B., Bennis, S., Bruning, T., Docker, M. F., Good, S., Lampman, R., ... & Johnson, N. S., 2024. Evaluation of larval sea lamprey *Petromyzon marinus* growth in the laboratory: Influence of temperature and diet. Aquaculture Research, 2024(1), 5547340.

Jenkins, J. A., Bart Jr, H. L., Bowker, J. D., Bowser, P. R., MacMillan, J. R., Nickum, J. G., ... & Warkentine, B. E., 2014. Guidelines for the use of fishes in research. Bethesda, Maryland, USA: American Fisheries Society.

Kane, A. S., Kahng, M. W., Reimschuessel, R., Nhamburo, P. T., & Lipsky, M. M., 1994.
UDP☐glucuronyltransferase kinetics for 3☐trifluoromethyl☐4☐nitrophenol (TFM) in
fish. Transactions of the American Fisheries Society, 123(2), 217-222.

Kitchell, J. F., 1990. The scope for mortality caused by sea lamprey. Transactions of the
American Fisheries Society, 119(4), 642–648.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B., 2017. lmerTest Package: Tests in
linear mixed effects models. Journal of Statistical Software, 82(13).

Lampman, R. T., Moser, M. L., Jackson, A., Rose, R., Gannam, A., & Barron, J., 2016. Chapter
twenty two: Developing techniques for artificial propagation and early rearing of Pacific
lamprey (*Entosphenus tridentatus*) for species recovery. Jawless fishes of the world:
Volume 2, 160.

Lavis, D. S., Henson, M. P., Johnson, D. A., Koon, E. M., & Ollila, D. J., 2003. A case history of
sea lamprey control in Lake Michigan: 1979 to 1999. Journal of Great Lakes Research,
29, 584–598.

Lawrence, M. J., Grayson, P., Jeffrey, J. D., Docker, M. F., Garroway, C. J., Wilson, J. M., ... &
Jeffries, K. M., 2022. Variation in the transcriptome response and detoxification gene
diversity drives pesticide tolerance in fishes. Environmental Science &
Technology, 56(17), 12137-12147.

Lech, J. J., & Statham, C. N., 1975. Role of glucuronide formation in the selective toxicity of 3-
trifluoromethyl-4-nitrophenol (TFM) for the sea lamprey: Comparative aspects of TFM
uptake and conjugation in sea lamprey and rainbow trout. Toxicology and Applied
Pharmacology, 31(1), 150–158.

Marsden, J. E., & Siefkes, M. J., 2019. Control of invasive sea lamprey in the Great Lakes, Lake
Champlain, and finger lakes of New York. In M. F. Docker (Ed.), Lampreys: Biology,
Conservation and Control: Volume 2 (pp. 411–479). Springer Netherlands.

McLaughlin, R. L., Hallett, A., Pratt, T. C., O'Connor, L. M., & McDonald, D. G., 2007.
Research to guide use of barriers, traps, and fishways to control sea lamprey. Journal of
Great Lakes Research, 33, 7-19.

Middaugh, C. R., Sepúlveda, M. S., & Höök, T. O., 2014. Growth and behavioral effects of the
lampricide TFM on non-target fish species. Journal of Great Lakes Research, 40(4),
1010-1015.

Muhametsafina, A., Birceanu, O., Hlina, B. L., Tessier, L. R., & Wilkie, M. P., 2019. Warmer
waters increase the larval sea lamprey's (*Petromyzon marinus*) tolerance to the
lampricide 3-trifluoromethyl-4-nitrophenol (TFM). Journal of Great Lakes Research,
45(5), 921–933.

Neave, F. B., Mandrak, N. E., Docker, M. F., & Noakes, D. L., 2006. Effects of preservation on
pigmentation and length measurements in larval lampreys. Journal of Fish
Biology, 68(4), 991-1001.

Niblett, P. D., & Ballantyne, J. S., 1976. Uncoupling of oxidative phosphorylation in rat liver
mitochondria by the lamprey larvicide TFM (3-trifluoromethyl-4-nitrophenol). Pesticide
Biochemistry and Physiology, 6(4), 363-366.

OECD., 2019. Test No. 203: Fish, Acute Toxicity Test. Organisation for economic co-operation
and development. https://www.oecd-ilibrary.org/environment/test-no-203-fish-acute-
toxicity-test_9789264069961-en

R Core Team, 2023. *R: A Language and Environment for Statistical Computing* (R version 4.3.2
(2023-10-31)) [Computer software]. R Foundation for Statistical Computing.
https://www.R-project.org/

Scholefield, R. J., Slaght, K. S., & Stephens, B. E., 2008. Seasonal variation in sensitivity of
larval sea lampreys to the lampricide 3-trifluoromethyl-4-nitrophenol. North American
Journal of Fisheries Management, 28(5), 1609–1617.

Schueller, J. R., Boogaard, M. A., Kirkeeng, C. A., Schloesser, N. A., Wolfe, S. L., Lettenberger,
A. J., King-Heiden, T. C., & Luoma, J. A., 2024. Seasonal differences in larval sea
lamprey (*Petromyzon marinus*) sensitivity to the pesticide TFM. Journal of Great Lakes
Research, 50(1), 102248.

Slade, J. W., Adams, J. V., Christie, G. C., Cuddy, D. W., Fodale, M. F., Heinrich, J. W.,
Quinlan, H. R., Weise, J. G., Weisser, J. W., & Young, R. J., 2003. Techniques and
methods for estimating abundance of larval and metamorphosed sea lampreys in Great
Lakes tributaries, 1995 to 2001. Journal of Great Lakes Research, 29, 137–151.

Smith, B. R., & Tibbles, J. J., 1980. Sea lamprey (*Petromyzon marinus*) in Lakes Huron,
Michigan, and Superior: History of invasion and control, 1936–78. Canadian Journal of
Fisheries and Aquatic Sciences, 37(11), 1780–1801.

Tessier, L. R., Long, T. A. F., & Wilkie, M. P., 2018. Influence of body size, metabolic rate and
life history stage on the uptake and excretion of the lampricide 3-trifluoromethyl-4-
nitrophenol (TFM) by invasive sea lampreys (*Petromyzon marinus*). Aquatic Toxicology,
194, 27–36.

Wilkie, M. P., Hubert, T. D., Boogaard, M. A., & Birceanu, O., 2019. Control of invasive sea
lampreys using the piscicides TFM and niclosamide: Toxicology, successes & future
prospects. Aquatic Toxicology, 211, 235–252.

Wilkie, M. P., Tessier, L. R., Boogaard, M., O'Connor, L., Birceanu, O., Steeves, T. B., &
Sullivan, W. P., 2021. Lampricide bioavailability and toxicity to invasive sea lamprey
and non-target fishes: The importance of alkalinity, pH, and the gill microenvironment.
Journal of Great Lakes Research, 47, S407–S420.

Wong, T. T., 2015. Performance evaluation of classification algorithms by *k*-fold and leave-one-
out cross validation. Pattern Recognition, 48(9), 2839–2846.

Yin, X., Martinez, A. S., Sepúlveda, M. S., & Christie, M. R., 2021. Rapid genetic adaptation to
recently colonized environments is driven by genes underlying life history traits. BMC
Genomics, 22(1), 269.

Figure legends

Figure 1. Relationship between total length (mm) and mass (g) of larval sea lamprey exposed to TFM (2.52 ± 0.08 mg/L; n = 8,611). Total length ranged from 43.0 mm to 137.5 mm (μ = 85.4 ± 12.9). Mass ranged from 0.074 g to 4.529g (μ = 0.869 ± 0.433).

Figure 2. Relationship between total length, mass, and survival for 8,604 larval sea lamprey after experimental exposure to TFM. Relationships between (A) total length (mm), (B) body mass (g; log transformed), and (C) condition factor ($CF = m/l^{2.6} \cdot 10^5$) and survival time of larval sea lamprey exposed to TFM (2.52 ± 0.08 mg/L) for up to 18 hours (n = 8,811 larvae). Larvae that survived beyond eight hours are binned within the group 8+ . Total length and mass are both positively associated with survival time. Best-fit lines are from basic linear regression.

Figure 3. Relationship between body length and survival time of larval sea lamprey exposed to TFM (2.52 ± 0.08 mg/L) for up to 18 hours (n = 8,611 larvae). Total length is both positively associated with survival time. Zero mortality occurred at hours one and 12. Five larval sea lamprey survived beyond 12 hours of TFM exposure (circled in red).

Figure 4. Correlation between morphometric variables measured immediately following death and survival in TFM for a smaller scale study. Relationships between (A) total length (mm), (B) body mass (g), and (C) condition factor ($CF = m/l^{2.6} \cdot 10^5$) and survival of larval sea lamprey exposed to TFM (4.9 ± 0.5 mg/L) over 16 hours (n = 48 larvae). The five larvae which survived the 16 hour exposure (blue circles) and are included in the regression calculation (n = 48 larvae). Best-fit lines are from basic linear regression. Overlapping points are slightly jittered for visual clarity.

Figure 5. Changes in larvae length and mass over time during storage in ethanol. (A) Total length (mm) and (B) mass (g) of larval sea lamprey from HBBS (n = 110) measured over successive days in ethanol storage.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5