Larger larval sea lamprey (*Petromyzon marinus)* have longer survival times when exposed

to the lampricide 3-trifluoromethyl-4-nitrophenol

Allison M. Nalesnik[a*], Emily L. Martin[a], Ian S. Kovacs[a], Connor S. Johnson[a], Emma I. Carroll[a], ,

Aaron Jubar[b], William Hemstrom[a], Michael P. Wilkie[c], Erin S. Dunlop[d], Maria S. Sepulveda[e,f],

Nicholas S. Johnson[g], Mark R. Christie[a,e]

a. Department of Biological Sciences, Purdue University, 915 W. State St., West Lafayette, Indiana

47907-2054 USA

b. U.S. Fish and Wildlife Service, Ludington Biological Station, 5050 Commerce Drive, Ludington, MI

49431, USA

c. Department of Biology and Laurier Institute for Water Science, Wilfrid Laurier University, Waterloo,

ON, N2L 3C5, Canada

d. Aquatic Research and Monitoring Section, Ontario Ministry of Natural Resources, c/o Trent

University, 2140 East Bank Drive, Peterborough, ON, K9L 0G2, Canada

e. Department of Forestry and Natural Resources, Purdue University, 715 W. State St., West Lafayette,

Indiana 47907-2054 USA

f. Sustainability Research Centre and Department Ecology and Biodiversity, Faculty of Life Sciences,

Universidad Andres Bello, Santiago, Chile

g. U.S. Geological Survey, Great Lakes Science Center, Hammond Bay Biological Station, 11188 Ray

Road, Millersburg, MI 49759, USA

*corresponding author: Allison Nalesnik; Department of Biological Sciences, Purdue University; 715 W.

State St., West Lafayette, Indiana 47907; analesni@purdue.edu; 301-706-0594

running title: Larval sea lamprey size and TFM survival

Abstract

Invasive sea lamprey (*Petromyzon marinus*) in the Laurentian Great Lakes have negatively impacted ecological and economically important fishes for nearly a century. To mitigate these effects, the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) is applied annually on a rotating basis to selected Great Lakes tributaries to kill larval lamprey before they become juveniles, out-migrate to the lakes, and parasitize other fishes. It has been hypothesized that larval size (e.g., mass, length) may affect survival time in response to TFM. To test this hypothesis, we conducted an experiment with 8,611 larvae across four temporal replicates, in which TFM concentrations equivalent to those used in present-day stream treatments were applied for up to 18 hours. When examining the survival times of larval lamprey exposed to TFM, we found a significant, positive relationship between length, mass, toxicity, and their interactions. For every one mm increase in total length, a corresponding increase by one gram of mass reduced survival time by 0.4315 minutes [95% CI: 0.5283 – 0.2992] and vice versa (i.e., the significant interaction between length and mass revealed that as larvae increase in mass, the survival benefit to being longer decreases, and vice versa). The changes in total length and mass of larval sea lamprey stored in ethanol for four months was also quantified. The observation that five larvae survived well past the 12-hour time window of a typical TFM field treatment highlights the need for continuous monitoring and the development of new control strategies to ensure the continued effective management of this invasive species.

keywords**:** ethanol storage, invasive species, lampricide, larval size, sea lamprey

Introduction

Sea lamprey (*Petromyzon marinus*) are hematophagous, non-host specific parasites that are native to the North Atlantic Ocean with populations along the coasts of Europe and North America. However, sea lamprey are an invasive pest in the Laurentian Great Lakes (Hansen et al., 2016) where they have contributed to substantial mortality and population declines in ecologically and economically important fishes (Coble et al., 1990; Kitchell, 1990). Sea lamprey were first documented in Lake Ontario in the late 1800s, after which they rapidly spread to Lake Erie following modifications to the Welland Canal in the early 1900s (Docker et al., 2021; Eshenroder 2014). By the late 1930s, sea lamprey had spread to Lake Huron, Lake Michigan, and Lake Superior (Christie and Goddard, 2003). In the novel environment of the Great Lakes, the typically anadromous sea lamprey acclimated to life-long residence in an entirely freshwater environment and developed smaller body sizes both at the time of metamorphosis and as adults in comparison to individuals from their native range (Hansen et al., 2016). These changes may be, at least in part, the result of a response to selection in genes responsible for growth, reproduction, and bioenergetics (Yin et al., 2021). This capacity for adaptation to novel environmental conditions suggests that sea lamprey may also be capable of responding to other selective pressures encountered in the Great Lakes.

Early control methods were initiated in the 1950s and relied mostly on engineered structures such as physical and electrical barriers and physical traps to prevent adult sea lamprey from accessing reproductive habitat (Hrodey et al., 2021; Smith and Tibbles, 1980), but the devices were labor intensive to operate and blocked valued fishes (McLaughlin et al., 2007). Lampricides were developed as an alternative approach to population control; over 4,000 organic candidate chemicals were tested at the Hammond Bay Biological Station (HBBS, Millersburg, MI), of which ten compounds were found to have greater toxicity for larval sea lamprey than other fishes at comparatively lower concentrations (Applegate et al., 1957, 1958, 1966). The chemical compound 3-trifluoromethyl-4-nitrophenol (hereafter TFM) was identified as the most effective agent and was found to be practical for field application, bearing relatively low toxicity to other species at low concentrations and a low manufacturing cost (Applegate et al., 1961; Smith and Tibbles, 1980).

TFM induces mortality by interfering with the oxygen-dependent production of ATP
within cells and tissues (Birceanu et al. 2021; Huerta et al. 2020; Niblett and Ballantyne 1976).
Larval sea lamprey are hypothesized to be particularly susceptible to TFM given their basal
location on the vertebrate phylogeny and limited ability to metabolize TFM through its
molecular conjugation to form TFM glucuronide, a less toxic TFM derivative that is readily
excreted (Bussy et al., 2018; Kane et al. 1994; Lawrence et al., 2022; Lech and Statham, 1975).
Since control methods were implemented in the late 1950s, sea lamprey populations have
decreased by approximately 90% from their historically high abundances, aiding the recovery of
some native fish populations such as lake trout (*Salvelinus namaycush;* Marsden and Siefkes,
2019). The continued control of sea lamprey in the Great Lakes hinges on effective stream
treatments, which can be influenced by local environmental conditions at treatment sites.
Because TFM bioavailability is higher in low pH and low alkalinity waters, control personnel
can use this relationship to modify TFM treatment concentrations accordingly (Wilkie et al.,
2021). Larval TFM sensitivities may also be influenced by water temperature (Hlina et al., 2021;
Muhametsafina et al., 2019), time of year (Scholefield et al., 2008), and possibly liver glycogen
content (although direct mechanisms are not well described), such that treatment efficacy can
depend on water conditions and larval body conditions (Schueller et al., 2024). Despite TFM
being instrumental to controlling sea lamprey populations, specific quantitative relationships
between larval lamprey body size and toxicant susceptibility remain largely uncharacterized,
although TFM uptake is known to be inversely related to body mass (Tessier et al., 2018). In the
present study, we examine how total length, mass, and their interaction contribute to TFM
susceptibility in sea lamprey. A better understanding of these effects is critical for determining
appropriate TFM dosages and durations for field applications over the next few decades, which
can be adjusted based on larval size surveys to ensure continued treatment effectiveness and
population control.

Here, we empirically test the relationship between larval sea lamprey body size and
survival under TFM exposure while accounting for pH, season, temperature, and source of
larvae. We predicted that because oxygen consumption and TFM uptake per unit of mass are
greater in smaller mass larvae (Tessier et al., 2018), larger animals would be more likely to
survive for prolonged periods of time in TFM. To test this hypothesis, we exposed   8,611
larval sea lamprey to TFM concentrations designed to mimic treatments in the field and

measured survival after each hour of exposure. We measured the total length and mass of each

larva and quantified the relationships between these morphometric measures and survival time.

We also used this opportunity to describe the relationship between changes in body size in larval

sea lamprey and their storage time in ethanol. To do so, we quantified changes in total length,

mass, and condition of larval sea lamprey over a four-month period.

Materials and Methods

Sample collection

All larvae used in this experiment were collected by crews from the United States Fish and

Wildlife Service and United States Geological Survey over a period of eight days during July

2022 (ESM **Table S1**). Larvae were obtained from the Muskegon River, a tributary to Lake

Michigan, MI, as it harbors one of the largest populations of spawning adult and larval sea

lamprey within the Great Lakes basin (Lavis et al., 2003; Slade et al., 2003). Prior to collecting

the larval lamprey for the present study, the Muskegon River had been treated with TFM in 2017

and 2019. Given this three-year interval and the growth rate of larvae (Dawson et al., 2021), the

larval lamprey used in this study are unlikely to have been previously exposed to TFM. Pulsed-

DC backpack electrofishing units were used to capture larval sea lamprey from four reaches of

the Muskegon River (ESM **Table S1**). To minimize the effects of seasonality, we condensed

collection times as much as possible. Most larvae collected were between 60-100 mm (range: 40-

140 mm).

To minimize stress experienced by the larvae due to handling and transportation, larvae

were transferred to acclimation tanks at U.S. Geological Survey, Hammond Bay Biological

Station (HBBS; Millersburg, MI) within one day of collection in individually labeled zippered

mesh bags (Tutudow, 40 cm x 50 cm, polyester, 250 µm mesh). Up to 50 larvae were transported

in each bag alongside two liters of sand from the Muskegon River. The sand grains were larger

than the mesh size of the bag such that the sand added to each bag remained in place during

transit. Bags containing larvae and sand were placed in 50 L aerated tanks maintained at a

temperature of 15 $^{\circ}$C overnight (approximately the temperature of the Muskegon River). The day

after capture, larvae held in mesh bags were transported to HBBS in 200 L aerated tanks

maintained at 15 $^{\circ}$C.

Upon arrival at HBBS, larvae in each bag were separated from the sand by emptying the

contents of each bag onto a 750 µm screen and using a slow stream of water from Lake Huron at

15 $^{\circ}$C to wash the sand through the screen, leaving the larvae on top of the screen. Larvae from

each specific bag were next moved into a mesh container (Ikea Fyllen Laundry Basket, 79 L

volume, 50 cm high and 45 cm diameter, 500 µm mesh size) and placed inside an aerated 1000 L

tank supplied with water from Lake Huron at ambient temp (16-18 $^{\circ}$C; water was pumped from a

depth of 25 m in Lake Huron). Within each holding tank, six mesh containers were used to hold

at most 282 larvae each. Sand with a 250 µm grain size was added to each mesh container to

achieve a layer of sand five cm thick for the larval lamprey to burrow into. Capture dates and

locations, the dates of transportation, and the locations of acclimation at HBBS were tracked for

each batch of larvae (ESM **Table S1**). All larvae were acclimated at HBBS within the mesh

container they were assigned for two weeks prior to exposure to TFM, during which larvae were

fed fresh baker's dry yeast at a rate of 1 gram (g) of yeast per larvae two times per week (Hanson

et al., 1974; Hume et al., 2024).

TFM exposure experiments

Our TFM target concentrations were determined by preliminary range-finding trials to determine

the lethal concentration of TFM required to kill 90% of larval sea lamprey over 12 hours (12-h

LC$_{90}$), based on the local water conditions and the tank systems to be used, following the

standard guidelines for acute toxicity tests in fish (OECD, 2019). All protocols involving the

handling of larvae at the Hammond Bay Biological Station were carried out in accordance with

United States federal guidelines for care and use of animals in accordance with American

Fisheries Society, ARRIVE guidelines, and the U.S. Animal Welfare Act of 1970 (Jenkins et al.,

2014). We aimed to use a 12-hour exposure period to approximate the total duration of TFM

treatments in the field (Middaugh et al., 2014). Our study consisted of four replicate TFM

exposure experiments (replicated two days apart) each using an identical design to measure the

survival of larval sea lamprey exposed to TFM. Each replicate experiment was conducted in two

1500 L tanks containing 624 L of water supplied from Lake Huron. Each tank was dosed with

between 104 and 112 mL of a 15,000 mg/L TFM stock solution at least 12 hours prior to

introducing the larvae to reach target concentrations of 2.3 to 2.5 mg/L TFM. No additional

water was added to these tanks after TFM was added. Water from Lake Huron did not vary

substantially in temperature (μ = mean ± standard deviation; μ = 17.6 ± 0.5 ^{o}C) or alkalinity over

the four replicant trials (μ = 85.6 ± 0.6 mg/L CaCO$_3$; ESM **Table S2**). However, in some cases,

there were fluctuations in pH during treatments, as discussed below (μ = 8.1 ± 0.3; ESM **Table**

S2). Each tank was fitted with a 15 cm long aeration stone at the center to facilitate uniform

distribution of TFM throughout the tank and to keep dissolved oxygen (DO) concentrations

higher than 7.6 mg/L (μ = 9.0 ± 0.6 mg/L; ESM **Table S2**). The air stone was set to turbulently

aerate the water for at least 30 minutes after initial dosing and reduced to a moderate level for

subsequent mixing and aeration. The TFM concentration was measured thirty minutes after

initial dosing the day before each experiment began and at least 12 hours later on the day of the

experiment to   determine if additional adjustments were required to maintain consistent

exposure conditions between our replicates.

Between 55 and 282 larval sea lamprey were kept in each of six mesh bags during

acclimation (ESM **Table S1**) which were transferred to the experimental tank at the start of each

experiment. When the bags in the acclimation tanks were lifted and twisted, the 250µm sand fell

through the 500µm mesh containers leaving only larval sea lamprey and no sand, eliminating the

stress and potential injuries to larvae if each larva was dug from the sand. The larvae in each bag

were then lowered into the experimental tank nearby. Each bag was assessed for mortality in the

order in which the bags went into the experimental tanks.

Throughout the duration of each experiment, several parameters were monitored on an

hourly basis. Fish mortality within each bag was assessed to determine survival time in TFM. To

assess mortality, teams of two people transferred the larvae from one mesh bag into a double-net

apparatus (**SI video**). Fish that were alive with swimming ability intact swam through the

apparatus and into an empty mesh bag held below. Dead or impaired fish were unable to swim

through the apparatus and required visual inspection to determine status (alive or dead). Larvae

were inspected in a shallow plastic tray filled with the same TFM solution from the respective

experimental tank for up to three minutes. Larvae were declared dead after the cessation of

branchiopore (gill pore) movement and unresponsiveness to tail pinching, then stored in

individually labeled bottles of 95% ethanol. Impaired fish were returned to their original

experiment tank and into the same mesh bag as individuals that swam through the apparatus. In

addition to recording larval sea lamprey mortality at hourly intervals, DO (YSI Pro 20), pH

(Fisherbrand Accumet AP110 Portable), salinity (YSI Pro 20), TFM concentration, water and air

temperature were measured hourly. TFM concentration was measured using a Genesys 6

spectrophotometer (Thermo Electron Corporation, MA, USA) set to a wavelength of 395, using

analytical standards prepared in the same water, based on Standard Operating Procedure No.

LAB 431.0, Hammond Bay Biological Station, Millersburg, MI (GLFC.

https://glfc.org/sop.php). Analytical standards were prepared from TFM (CAS Number 88-30-2;

Millipore-Sigma, St. Louis, MO). Water and air temperature were measured via a laboratory

liquid-in-glass thermometer. The experiments varied in duration from five to eighteen hours until

the last larva had died. At the conclusion of all experiments, preserved larvae were transported

from HBBS to Purdue University (West Lafayette, IN) where the total length (mm) and mass (g)

was measured for all lamprey (n = 8,611 larvae). All experiment samples were stored in 95%

ethanol for at least four months prior to measuring their total length and mass.

To determine what physiological variables impact lamprey survival time in TFM, we

used model selection to identify the best fitting linear mixed effect models for predicting survival

time (hour), using total length (mm), mass (g), toxicity units, and their interactions. Toxicity

units were calculated by the Henderson-Hasselbalch equation (McDonald & Kolar 2007) as

$\frac{1}{1+10^{pH-pKa}}$ where the pKa (the negative log of the acid dissociation constant, Ka, for an acid-

base reaction) is 6.07 for TFM (Hubert 2003). These variables were fitted as fixed effects; we

also included a nested random effect of experiment in tank in bag as a covariate using R 4.3.2 (R

Core Team, 2023) and the R package lmerTest (Kuznetsova et al., 2017). Based on diagnostic

assessments, the assumptions for a linear mixed effects model were largely met. A Q-Q plot of

the residuals indicated that the data follow a normal distribution, with only minor deviations

observed. Additionally, the residuals versus fitted values plot showed no discernible patterns,

with most points randomly scattered around the zero line. While five data points exhibited slight

deviations from the expected trend, given the dataset's size, their impact on the model was

minimal. Therefore, the assumptions of normality, linearity, and homoscedasticity were

reasonably satisfied. We compared models using total length, mass, toxicity units, and their

interactions (seen ESM **Table S3** for all models). Additionally, one tank (S4) within replicate

one had a large fluctuation in toxicity units (illustrated in ESM **Fig. S1**). We evaluated our linear

mixed effect models with and without this replicate and the best model was robust to its removal.

We removed tank S4 within replicate one from our analyses due to the greater variance in
toxicity units that is dissimilar from other replicates.

Long-term effects of ethanol storage
To assess the effect of long-term ethanol storage on larval sea lamprey morphometrics, we
systematically measured the total length and mass of two different sets of larval sea lamprey over
the course of four months (hereafter storage experiments A and B). None of the larvae used to
quantify changes in body size and mass due to ethanol storage were included in the TFM
exposure experiments; instead, these larvae were used to determine the effect of measuring the
8,611 experimental larvae after a storage period of up to eight months since it was logistically
impractical to measure all individuals during the large TFM exposure experiments. Storage
experiment A: In July, prior to the large TFM survival experiment described above, 554 larval
sea lamprey were transported to Purdue University's Aquaculture Research Laboratory (ARL;
West Lafayette, IN). Sea lamprey were exposed to TFM concentrations ranging from 2.0 mg/L
to 4.0 mg/L in groups ranging from 10 to 25 individuals. If sea lamprey remained alive beyond
36 hours of TFM exposure, they were euthanized with 250 mg/L tricaine methanesulfonate (MS-
222) buffered to pH 7 with sodium bicarbonate. Total length (mm) and mass (g) were measured
immediately after death and each sample was individually stored in a 50 mL conical tube
containing 40 mL of 95% non-denatured ethanol. Purdue University Institutional Animal Care
and Use Committee reviewed experimental and handling procedures for this study, but did not
require animal care protocols for the larval sea lamprey. Storage experiment B: At HBBS, 110
larval sea lamprey that remained in the acclimation tank after the completion of the large TFM
survival experiments were euthanized with MS-222 using the same procedure described above.
We measured the total length (mm) and mass (g) of each larva immediately after death, then
stored every individual in 15 mL conical tubes containing 12 mL of 95% ethanol. Thereafter,
each larval sea lamprey (554 from ARL, 110 from HBBS) was measured for body mass and total
length systematically over a period of 120 days (see **SI Methods** for details on measurement
procedures). We calculated condition (CF) using the same equation as above.

Given that larvae become smaller over time when stored in ethanol, we fit an additional
model to predict the initial size of an individual from a measurement taken after extended storage
in ethanol. We first randomly sampled our dataset of Experiment A and Experiment B

individuals (664 larvae) such that only one sample per individual was used (since any models

containing individual ID as a covariate could not be used to make predictions for new

individuals). We then used the *nls* function in R to fit a nonlinear regression model to determine

the decay rate which best fit our observed data using an exponential decay function:

$$l_0 = \frac{l_t}{e^{-bt}}$$

where l_0 is initial length, l_t is the length at measurement time t, and b is the rate of decay. We

evaluated the ability of this model to predict the initial size of a larval sea lamprey prior to

storage using a hold-one-out cross validation approach (see **Tables S4** and **S5**; Wong, 2015).

Using these adjusted values, we also re-analyzed the data from our large TFM exposure and

body size experiments using the exact statistical models described above.

Results

Results from TFM exposure experiments

The 7,651 lamprey used in our four experimental replicates (seven experiment tanks) survived

between two and 18 hours of TFM exposure (**Fig. 1A**). There was an allometric relationship

between larval total length and larval mass (**Fig. 1B**). Measured TFM concentrations during

exposure averaged 2.503 ± 0.07 mg/L (n = 78) across our replicates (**ESM Table S2; ESM Fig.**

S1). TFM toxicity units averaged 0.03216, ranging from 0.009558 to 0.04813 (calculated from

ESM Table S2). The 7,651 lamprey used in our four experimental replicates had total lengths

between 43.0 and 137.5 mm (μ = 85.53 mm; **Fig. 1B**; **ESM Fig. S2A**) and masses between 0.074

and 4.529 g (μ = 0.8702 g; **Fig. 1C**; **ESM Fig. S2B**). The best fitting linear mixed effect model

had the equation y= 6.711 + 0.02686*(total length, mm) + 0.4590*(mass, g) – 81.76*(toxicity

units) – 0.006892(total length*mass) – 0.7365(total length*toxicity units) – 21.15(mass*toxicity

units) and explained a large portion of the variation in survival time in TFM (conditional R^2 =

0.8790; **Table 1**; extended results in ESM **Table S3**). Total length was significantly and

positively associated with survival time in TFM (**Table 1; ESM Fig. S2A)** (*p* value = $2.592 \cdot 10^{-8}$

). There was a 16.11 minute [95% CI: 10.46-21.80] increase in survival time for each one cm

increase in total length after accounting for the effects of other parameters. Mass was

significantly and positively associated with survival time in TFM (Table 1; ESM **Fig. S2B**) (*p*

value = 0.01455). There was a 27.54 minute [95% CI: 5.504-49.68] increase in survival time for

each one g increase in total mass after accounting for the effects of other parameters. Toxicity

units were negatively associated with survival time, such that an increase in toxicity units leads

to a decrease in survival time (p value = $1.180 \cdot 10^{-16}$). There was a 98.11 minute reduction in

survival time for each 0.02 increase in toxicity units. The effect of the interaction between total

length and mass on survival time was negative (p value < 0.001): for every one mm increase in

total length, a corresponding increase by one g of mass reduced survival time by 0.4315 minutes

[95% CI: 0.5283 – 0.2992] and vice versa (*i.e.,* the significant interaction between total length

and mass revealed that as larvae increase in mass, the survival benefit to being longer decreases,

and vice versa) (**Fig. 2; Table 1**). The effect of the interaction between mass and toxicity units

on survival time was positive (p value = $2.143 \cdot 10^{-7}$): for every one g increase in mass, the impact

of increasing by 0.02 toxicity units increases survival time by 25.38 minutes [95% CI: 15.80–

34.96]. The effect of the interaction between total length and toxicity units on survival time was

negative (p value = $6.577 \cdot 10^{-7}$): for every 10 mm increase in total length, the impact of

increasing by 0.02 toxicity units reduces survival time by 8.838 minutes [95% CI: 12.32– 5.365].

Five individuals survived beyond 12 hours of TFM exposure (**Fig. 3; ESM Fig. S4**).

Effects of ethanol storage on body size

HBBS and ARL larvae both declined in length and mass over the 120 day measurement period

(ESM **Fig. S3**). For example, the total length of HBBS larvae (n = 110) initially ranged from

63.5 to 127.5 mm (μ = 95.1 mm; ESM **Fig. S3A**). The last measured total length (day 120) of

HBBS larvae ranged from 56.0 to 119.5 mm (μ = 88.6 mm; ESM **Fig. S3A**). The mass of HBBS

larvae (n = 110) initially ranged from 0.43 to 2.72 g (μ = 1.27 g; ESM **Fig. S3B**). The last

measured mass of HBBS larvae ranged from 0.185 to 1.906 g (μ = 0.83 g; ESM **Fig. S3B**).

The nonlinear least squares model generated using the combined sample set (n = 664; site

was included as a fixed variable to account for differences in ethanol storage volume) yielded a

total length decay estimate as $4.483 \cdot 10^{-4}$ mm/day (ESM **Table S4**). The linear regression model

suggested a strong, positive relationship between predicted values and true values, with an

average change in predicted initial length of 1.035mm for each 1mm increase in observed length

post-storage after accounting for the effects of ethanol (p value < $2 \cdot 10^{-16}$; ESM **Table S5**). The

model explained a very large proportion of the variability in the initial length (R^2 = 0.69). We

used our model estimates to calculate initial total length, mass, and condition for our large TFM

exposure experiment larvae by dividing the measured value by e raised to the negative estimate,

multiplied by the number of days the larvae were stored in ethanol and found nearly identical

results (ESM **Fig. S5**).

Discussion

We describe a significant relationships between the total length, mass, and their interaction of

larval sea lamprey and their survival time in TFM. Larger larval sea lamprey exhibit increased

survival time in TFM, highlighting that this variable, if accounted for during field treatments,

could improve treatment success. To conduct TFM treatments with the highest larval sea

lamprey mortality, TFM treatment crews may want to consider larval size within a treatment

stream and adjust treatment duration accordingly. We also describe the predicted survival time

for larvae at the 10th, 25th, 50th, 75th, and 90th total length and mass percentiles within our data

(**Fig. 2;** ESM **Table S6**), which can be used to justify the duration of treatment in selected

streams based on surveyed larval sizes. For instance, streams where the average larval mass is

greater than our 90th percentile would require at least an additional 30 minutes of treatment

compared to streams where the average larval size closer to our 10th percentile (**Figure 2;** ESM

Table S6). An earlier set of models incorporated condition factor (CF) as a co-variate, which

was calculated as previously outlined for Pacific sea lamprey (Lampman et al., 2016) as

$CF = \frac{m*10^5}{l^{2.6}}$ where m = mass (g) and l = length (mm). While, CF was not a significant contributor

to reduced model AIC, perhaps because the equation is derived from a different species, the

significant negative interaction between mass and total length reveals that condition is an

important contributor to larval survival. Specifically, shorter and heavier larvae have longer

survival times than longer and lighter-weight larvae (**Fig. 2**).

The observed correlation between sea lamprey size and survival time in TFM suggests

the presence of a mechanism by which larger individuals can better withstand the toxic effects of

the lampricide. Given that our linear mixed effect model including total length, mass, toxicity

units, and their interactions explained a very large percentage of the variation in survival time

(conditional R^2 = 0.8790), we suggest that the response of sea lamprey larvae to TFM is partly

determined by larval size. While our primary focus was on describing the novel effects of larval

size on survival to TFM exposure, we recognized that other variables—namely pH and TFM

concentration —are well established as influential factors in TFM toxicity (Marking and Olsen

1975; Wilkie 2021). The inclusion of toxicity units in our model, despite our efforts to

standardize experimental conditions, allowed us to account for known sources of variability that

affect larval survival. By statistically controlling for these variables, we were able to more

precisely isolate and evaluate the independent contributions of total length, mass, and their

interaction on survival. This approach not only acknowledges the critical role of these

environmental parameters in modulating TFM efficacy but also strengthens the inference that

morphometric characteristics play a distinct and biologically relevant role in determining

survival outcomes. We employed the LC90 target concentration of TFM to select individuals that

exhibited prolonged survival, effectively sampling the right tail of the survival curve. This

approach provided a robust sample size for future analysis of potential genetic factors associated

with TFM survival.

In our study, the extended duration of survival for five individuals well beyond the

typical treatment length of TFM in the field, that were neither the longest nor the heaviest larvae,

indicate that some larvae may be capable of surviving treatment (**Fig. 3; ESM Fig. S4**). This

observation is consistent with underlying variation in the population in terms of susceptibility to

TFM and has implications for the potential for sea lamprey to evolve resistance to TFM, a

subject of management interest (Christie et al., 2019; Dunlop et al., 2021). The development of

resistance is a known consequence of repeated pesticide application (Délye et al., 2013;

Georghiou and Taylor, 1977; Hawkins et al., 2019) and given that TFM has been used annually

in freshwater streams in the Great Lakes Basin since 1958, resistance in sea lamprey is

conceivable and an urgent concern (Christie et al., 2019; Dunlop et al., 2021). Prior research has

suggested that a response to TFM-induced selection may already be underway in the Great

Lakes: sea lamprey genomic and transcriptomic comparisons between treated and untreated

populations of larval sea lamprey suggest that an adaptive, genetic response to TFM may drive

incipient resistance (Yin et al., 2021). Sea lamprey could develop resistance to TFM through two

broad mechanisms: (1) physiological adaptations or (2) evolution of their behavior or life history

(Christie et al., 2019; Dunlop et al., 2018, 2021). Physiological resistance could involve the

development of molecular mechanisms to more efficiently detoxify or neutralize TFM or to

counteract the effects of the mitochondrial proton gradient; however, if larger individuals have a

selective advantage for surviving treatment, and large size (or a correlated trait like growth rate)

is heritable, then there is the potential for both physiological and life history resistance to TFM

via selection for increases in body size. For example, larger sea lamprey could store additional

glycogen, which would provide them with larger anaerobic energy reserves to generate ATP

using anaerobic glycolysis. Such features could enable individuals to out-survive a typical TFM

treatment. Although previous studies demonstrated no clear relationship between liver or muscle

glycogen reserves and TFM tolerance, (Hlina et al. 2021; Muhametsafina et al., 2019), a recent

study by Schueller et al. (2024) found a positive relationship between liver glycogen

concentrations and the minimum lethal concentration (MLC = 12 h-LC$^{99.9}$; Bills et al. 2003) of

TFM to larval sea lamprey using stream-side toxicity tests. Larval sea lamprey also demonstrate

a remarkable capacity to recover from TFM exposure when transferred to clean, TFM-free water,

restoring energy stores such as glycogen and phosphocreatine within a few hours (Clifford et al.

2012). Collectively, these studies suggest that there may be a physiological advantage to being

larger at the time of TFM exposure/treatments (Dunlop et al., 2021).

Contrary in part to our findings here, a handful of previous studies describe that energy

stores and body size have little impact on TFM sensitivity (Hlina et al., 2021; Muhametsafina et

al., 2019) and instead report that larvae exhibit differential TFM sensitivity based on season,

which may be a result of the seasonally warmer waters enhancing their capacity to detoxify TFM

(Hlina et al., 2021; Muhametsafina et al., 2019). Additionally, as pH and alkalinity of water

increase, larvae exhibit increased tolerance as TFM changes its ionization state (reviewed in

Wilkie et al., 2019), resulting in lower TFM uptake in more basic environments (Hlina et al.,

2017). We accounted for pH within our analyses by proxy using toxicity units, controlled for

alkalinity and water temperature during the experiment, and controlled for seasonality by using

larvae collected within eight days in July to assess the contribution of total length and mass on

survival time in TFM. Additional experiments may be needed to better weigh the contribution of

other environmental and physiological components associated with differential survival in TFM

(*e.g.*, temperature and alkalinity versus body mass and length).

In addition to quantifying the relationship between larval size and survival in TFM, we

also characterized the relationship between larval body size and length after storage in ethanol.

This information is valuable because future studies on resistance may involve preserving residual

lamprey samples for subsequent analyses, and understanding the effects of ethanol storage on

body length and mass will be essential. Similarly, accurate knowledge of preservation effects

will benefit investigations into developmental milestones, such as size at metamorphosis, by

informing decisions about sample storage methods. We provide an estimate of the decay rates of

length and body mass, which should prove useful when back calculating the actual sizes of sea

lamprey larvae prior to preservation (as demonstrated in ESM **Fig. S5**). By generating estimates

for total length, mass, and condition, we evaluated the usefulness of each parameter to make

accurate predictions of initial state values. The total length estimates generated from each sample

set were of the same magnitude (on the scale of 10^{-4}). Estimates from mass and condition were

less accurate predictors of initial values, as reflected by smaller adjusted R^2 values, when

compared to those from total length. The best performing nonlinear regression model was

derived from our combined sample set and provides a total length exponential decay estimate of

$4.483 \cdot 10^{-4}$ mm/day (ESM **Table S4**). Given that the model derived from our combined sample

set had high predictability, our estimate may be useful across diverse larval sea lamprey

collections. However, we also observed some variability in sample responses to ethanol

preservation—while many larvae shrank over time, some exhibited minimal change or even

increases in size (see ESM **Fig. S3**). These deviations from the expected dehydration pattern (i.e.

exponentially shrinking in size) introduce a degree of uncertainty in back-calculations.

Therefore, we advise researchers to use our correction equation with awareness of its limitations

in cases where samples stored in ethanol may not follow typical shrinkage trends. Additionally,

we observed much higher variability in our ARL data (ESM **Fig. S3B,F**) compared to the HBBS

data (ESM **Fig. S3A,E**), likely due to different sample sizes and different combinations of

researchers measuring each dataset, although many of the same researchers measured samples

from each group of samples. Nevertheless, our most predictive model was that derived from the

combined dataset of both groups of samples (ESM **Table S5**). Our model gives researchers

additional options, apart from using formalin (Neave et al., 2006), for correcting for changes in

body size after prolonged storage in 95% ethanol.

Conclusion

Our study provides valuable insights into the relationship between larval sea lamprey size and survival time in TFM-treated environments. These findings suggest that duration and TFM dosage for routine TFM treatments within the Great Lakes basin could be improved if larval length and mass can be taken into consideration. Additionally, studies that store larval sea lamprey in 95% ethanol can now correct for storage time using a simple equation. We also emphasize the need for further research to better understand the underlying mechanisms between larger larval lamprey size and TFM survival to better inform sea lamprey management throughout the Great Lakes.

Author contributions

Allison Nalesnik: Formal analysis, Investigation, Data curation, Writing- Original Draft, Writing- Review & Editing, Visualization, Supervision, Project administration. Emily Martin: Investigation, Data Curation, Writing- Review & Editing. Ian Kovacs: Investigation, Data Curation, Writing- Review & Editing. Connor Johnson: Investigation, Data curation, Writing- Review & Editing. Emma Carroll: Investigation, Data curation, Writing- Review & Editing. Aaron Jubar: Methodology, Resources, Writing- Review & Editing, Project administration. William Hemstrom: Formal analysis, Writing- Review & Editing. Michael Wilkie: Formal analysis, Writing- Review & Editing. Erin Dunlop: Conceptualization, Methodology, Writing- Review & Editing, Funding acquisition. Maria Sepulveda: Conceptualization, Methodology, Investigation, Resources, Writing- Review & Editing, Funding acquisition. Nicholas Johnson: Conceptualization, Methodology, Investigation, Resources, Data curation, Writing- Review & Editing, Supervision, Project administration, Funding acquisition. Mark Christie: Conceptualization, Methodology, Investigation, Resources, Writing- Review & Editing, Supervision, Project administration, Funding acquisition.

Acknowledgments

We thank the USFWS team responsible for collecting sea lamprey larvae for this experiment: Jordan Andrus, Philip Bennett, John Ewalt, Timothy Granger, Allen Keffer, David Keffer, Callie Kopp, Daniel McGarry, Lily Smalstig, Jenna Tews, and Matt Lipps. We also thank the USGS collection team and on-site personnel who provided hands-on assistance during the experiment: David Coburn, Tyler Bruning, Ed Benzer, Zachary Nordstrom, Meg Spens, and Brad Buechel.

We thank Emma Vieregge and Karen Slaght for their support doing water quality analysis and
range-finder assays. We thank Catherine Searle for her advice on our models. We also thank
Robert Rode for his advice and assistance at the ARL, as well as undergraduate support from
Lilly Gorney, Leah Kemple, and Grace Bowman. This research was funded by support to MRC,
MSS, ESD, and NSJ from the Great Lakes Fishery Commission (grant number
2022_CHR_541010, 2022). Any use of trade, firm, or product names is for descriptive purposes
only and does not imply endorsement by the U.S. Government.

Conflict of interest

The authors declare that they have no conflict of interest.

Data archiving

All code to reproduce the analyses reported in this manuscript are available via GitHub at

https://github.com/ChristieLab/sea-lamprey-morphometrics. Data will be made available on

Dryad.

References

Applegate, V.C., Howell, J.H., Hall Jr, A.E., Smith, M.A., 1957. Toxicity of 4,346 chemicals to
larval lamprey and fishes. US Fish and Wildlife Service Special Scientific Report -
Fisheries 207: 1-157.

Applegate, V.C., Howell, J.H., Moffett, J.W., Smith, M.A., 1961. Use of 3-trifluormethyl-4-
nitrophenol as a selective sea lamprey larvicide. Great Lakes Fishery Commission
Technical Report 1: 3-35.

Applegate, V.C., Johnson, B.G.H., Smith, M.A., 1966. The relation between molecular structure
and biological activity among mononitrophenols containing halogens. Technical Report
11. Great Lakes Fishery Commission. 1-19.

Applegate, V.C., Howell, J.H., Smith, M.A., 1958. Use of mononitrophenols containing halogens
as selective sea lamprey larvicides. Science 127(3294), 336–338.

Bills, T.D., Boogaard, M.A., Johnson, D.A., Brege, D.C., Scholefield, R.J., Westman, R.W.,
Stephens, B. E., 2003. Development of a pH/alkalinity treatment model for applications
of the lampricide TFM to streams tributary to the Great Lakes. J. Great Lakes Res. 29,
510-520.

Birceanu, O., Tessier, L.R., Huerta, B., Li, W., McDonald, A., Wilkie, M.P., 2021. At the intersection between toxicology and physiology: What we have learned about sea lamprey and bony fish physiology from studying the mode of action of lampricides. J. Great Lakes Res. 47, S673-S689.

Bussy, U., Chung-Davidson, Y.W., Buchinger, T., Li, K., Smith, S.A., Jones, A.D., Li, W., 2018. Metabolism of a sea lamprey pesticide by fish liver enzymes part A: Identification and synthesis of TFM metabolites. Anal. Bioanal. Chem. 410(6), 1749–1761.

Christie, G.C., Goddard, C.I., 2003. Sea Lamprey International Symposium (SLIS II): Advances in the integrated management of sea lamprey in the Great Lakes. J. Great Lakes Res. 29, 1–14.

Christie, M.R., Sepúlveda, M.S., Dunlop, E.S., 2019. Rapid resistance to pesticide control is predicted to evolve in an invasive fish. Sci. Rep. 9(1), Article 1.

Clifford, A.M., Henry, M., Bergstedt, R., McDonald, D.G., Smits, A.S., Wilkie, M.P., 2012. Recovery of larval sea lamprey from short-term exposure to the pesticide 3-trifluoromethyl-4-nitrophenol: implications for sea lamprey control in the Great Lakes. Trans. Am. Fish. Soc. 141(6), 1697-1710.

Coble, D.W., Bruesewitz, R.E., Fratt, T.W., Scheirer, J.W., 1990. Lake trout, sea lamprey, and overfishing in the upper Great Lakes: A review and reanalysis. Trans. Am. Fish. Soc. 119(6), 985–995.

Dawson, H.A., Higgins-Weier, C.E., Steeves, T.B., Johnson, N.S., 2021. Estimating age and growth of invasive sea lamprey: A review of approaches and investigation of a new method. J. Great Lakes Res. 47, S570-S579.

Délye, C., Jasieniuk, M., Le Corre, V., 2013. Deciphering the evolution of herbicide resistance in weeds. Trends Genet. 29(11), 649–658.

Docker, M.F., Bravener, G.A., Garroway, C.J., Hrodey, P.J., Hume, J.B., Johnson, N.S., Lewandoski, S.A., Ogden, J.L., Zollweg-Horan, E.C., 2021. A review of sea lamprey dispersal and population structure in the Great Lakes and the implications for control. J. Great Lakes Res. 47, S549–S569.

Dunlop, E.S., Christie, M.R., McLaughlin, R., Steeves, T.B., 2021. Life history evolution of sea lamprey is predicted to reduce the effectiveness of pesticide control. J. Great Lakes Res. 47, S259–S271.

Dunlop, E.S., McLaughlin, R., Adams, J.V., Jones, M., Birceanu, O., Christie, M.R., Criger, L.
A., Hinderer, J.L.M., Hollingworth, R.M., Johnson, N.S., Lantz, S.R., Li, W., Miller, J.,
Morrison, B.J., Mota-Sanchez, D., Muir, A., Sepúlveda, M.S., Steeves, T., Walter, L., …
Wilkie, M.P., 2018. Rapid evolution meets invasive species control: The potential for
pesticide resistance in sea lamprey. Can. J. Fish. Aquat. Sci. 75(1), 152–168.

Eshenroder, R. L., 2014. The role of the Champlain Canal and Erie Canal as putative corridors
for colonization of Lake Champlain and Lake Ontario by sea lamprey. Trans. Am. Fish.
Soc. 143(3), 634-649.

Fox J, Weisberg S (2019). An R Companion to Applied Regression, Third edition. Sage,
Thousand Oaks CA. <https://www.john-fox.ca/Companion/>.

Georghiou, G.P., Taylor, C.E., 1977. Genetic and biological influences in the evolution of
insecticide resistance. J. Econ. Entomol. 70(3), 319–323.

Hansen, M.J., Madenjian, C.P., Slade, J.W., Steeves, T.B., Almeida, P.R., Quintella, B.R., 2016.
Population ecology of the sea lamprey (*Petromyzon marinus*) as an invasive species in
the Laurentian Great Lakes and an imperiled species in Europe. Rev. Fish Biol. Fish.
26(3), 509–535.

Hanson, L.H., King, E.L., Howell, J.H., Smith, A.J., 1974. A culture method for sea lamprey
larvae. The Progressive Fish-Culturist 36(3), 122–128.

Hawkins, N.J., Bass, C., Dixon, A., Neve, P., 2019. The evolutionary origins of pesticide
resistance. Biol. Rev. 94(1), 135–155.

Hlina, B.L., Birceanu, O., Robinson, C.S., Dhiyebi, H., Wilkie, M.P., 2021. The relationship
between thermal physiology and lampricide sensitivity in larval sea lamprey (*Petromyzon
marinus*). J. Great Lakes Res. 47, S272–S284.

Hlina, B.L., Tessier, L.R., Wilkie, M.P., 2017. Effects of water pH on the uptake and elimination
of the piscicide, 3-trifluoromethyl-4-nitrophenol (TFM), by larval sea lamprey. Comp.
Biochem. Physiol. C: Toxicol. Pharmacol. 200, 9–16.

Hrodey, P.J., Lewandoski, S.A., Sullivan, W.P., Barber, J.M., Mann, K.A., Paudel, B., Symbal,
M.J., 2021. Evolution of the sea lamprey control barrier program: The importance of
lowermost barriers. J. Great Lakes Res. 47, S285–S296.

Hubert, T.D. (2003). Environmental fate and effects of the lampricide TFM: a review. J. Great
Lakes Res. 29, 456-474.

Huerta, B., Chung-Davidson, Y.W., Bussy, U., Zhang, Y., Bazil, J.N., Li, W., 2020. Sea lamprey
cardiac mitochondrial bioenergetics after exposure to TFM and its metabolites. Aquat.
Toxicol. 219, 105380.

Hume, J.B., Bennis, S., Bruning, T., Docker, M.F., Good, S., Lampman, R., ... Johnson, N.S.,
2024. Evaluation of larval sea lamprey *Petromyzon marinus* growth in the laboratory:
Influence of temperature and diet. Aquacult. Res. 2024(1), 5547340.

Jenkins, J. A., Bart Jr, H. L., Bowker, J. D., Bowser, P. R., MacMillan, J. R., Nickum, J. G., ...
Warkentine, B. E., 2014. Guidelines for the use of fishes in research. Bethesda,
Maryland, USA: Am. Fish. Soc.

Kane, A.S., Kahng, M.W., Reimschuessel, R., Nhamburo, P.T., Lipsky, M.M., 1994.
UDP‐glucuronyltransferase kinetics for 3‐trifluoromethyl‐4‐nitrophenol (TFM) in
fish. Transactions of the American Fisheries Society, 123(2), 217-222.

Kitchell, J.F., 1990. The scope for mortality caused by sea lamprey. Trans. Am. Fish. Soc.
119(4), 642–648.

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in linear
mixed effects models. J. Stat. Softw. 82(13).

Lampman, R.T., Moser, M.L., Jackson, A., Rose, R., Gannam, A., Barron, J., 2016. Chapter
twenty two: Developing techniques for artificial propagation and early rearing of Pacific
lamprey (*Entosphenus tridentatus*) for species recovery. A. Orlov, R. Beamish (Eds.),
Jawless Fishes of the World, 2, Cambridge Scholars Publishing (2016), pp. 160-195

Lavis, D.S., Henson, M.P., Johnson, D.A., Koon, E.M., Ollila, D.J., 2003. A case history of sea
lamprey control in Lake Michigan: 1979 to 1999. J. Great Lakes Res. 29, 584–598.

Lawrence, M.J., Grayson, P., Jeffrey, J.D., Docker, M.F., Garroway, C.J., Wilson, J.M., ...
Jeffries, K.M., 2022. Variation in the transcriptome response and detoxification gene
diversity drives pesticide tolerance in fishes. Environ. Sci. Tech. 56(17), 12137-12147.

Lech, J.J., Statham, C.N., 1975. Role of glucuronide formation in the selective toxicity of 3-
trifluoromethyl-4-nitrophenol (TFM) for the sea lamprey: Comparative aspects of TFM
uptake and conjugation in sea lamprey and rainbow trout. Toxicol. Appl. Pharmacol.
31(1), 150–158.

Marking, L.L., and Olson, L.E. 1975. Toxicity of the lampricide 3-trifluoromethyl-4-nitrophenol
(TFM) to non-target fish in static tests. U.S. Fish and Wildlife Service, Invest. in Fish
Control 60.

Marsden, J.E., Siefkes, M.J., 2019. Control of invasive sea lamprey in the Great Lakes, Lake
Champlain, and finger lakes of New York. In M. F. Docker (Ed.), Lamprey: Biology,
Conservation and Control: Volume 2 (pp. 411–479). Springer Netherlands.

McDonald, D.G., Kolar, C.S. (2007). Research to guide the use of lampricides for controlling
sea lamprey. J. Great Lakes Res. 33, 20-34.

McLaughlin, R.L., Hallett, A., Pratt, T.C., O'Connor, L.M., McDonald, D.G., 2007. Research to
guide use of barriers, traps, and fishways to control sea lamprey. J. Great Lakes Res. 33,
7-19.

Middaugh, C.R., Sepúlveda, M.S., Höök, T.O., 2014. Growth and behavioral effects of the
lampricide TFM on non-target fish species. J. Great Lakes Res. 40(4), 1010-1015.

Muhametsafina, A., Birceanu, O., Hlina, B.L., Tessier, L.R., Wilkie, M.P., 2019. Warmer waters
increase the larval sea lamprey's (*Petromyzon marinus*) tolerance to the lampricide 3-
trifluoromethyl-4-nitrophenol (TFM). J. Great Lakes Res. 45(5), 921–933.

Neave, F.B., Mandrak, N.E., Docker, M.F., Noakes, D.L., 2006. Effects of preservation on
pigmentation and length measurements in larval lamprey. J. Fish Biol. 68(4), 991-1001.

Niblett, P.D. Ballantyne, J.S., 1976. Uncoupling of oxidative phosphorylation in rat liver
mitochondria by the lamprey larvicide TFM (3-trifluoromethyl-4-nitrophenol). Pestic.
Biochem. Physiol. 6(4), 363-366.

OECD., 2019. Test No. 203: Fish, Acute Toxicity Test. OECD Guidelines for the Testing of
Chemicals, Section 2, OECD Publishing, Paris.

R Core Team, 2023. R: A Language and Environment for Statistical Computing (R version 4.3.2
(2023-10-31)) [Computer software]. R Foundation for Statistical Computing.
https://www.R-project.org/

Scholefield, R.J., Slaght, K.S., Stephens, B.E., 2008. Seasonal variation in sensitivity of larval
sea lamprey to the lampricide 3-trifluoromethyl-4-nitrophenol. North Am. J. Fish.
Manage. 28(5), 1609–1617.

Schueller, J.R., Boogaard, M.A., Kirkeeng, C.A., Schloesser, N.A., Wolfe, S.L., Lettenberger,
A.J., King-Heiden, T.C., Luoma, J.A., 2024. Seasonal differences in larval sea lamprey

(*Petromyzon marinus*) sensitivity to the pesticide TFM. J. Great Lakes Res. 50(1),

102248.

Slade, J.W., Adams, J.V., Christie, G.C., Cuddy, D.W., Fodale, M.F., Heinrich, J.W., Quinlan,

H.R., Weise, J.G., Weisser, J.W., Young, R.J., 2003. Techniques and methods for

estimating abundance of larval and metamorphosed sea lamprey in Great Lakes

tributaries, 1995 to 2001. J. Great Lakes Res. 29, 137–151.

Smith, B.R., Tibbles, J.J., 1980. Sea lamprey (*Petromyzon marinus*) in Lakes Huron, Michigan,

and Superior: History of invasion and control, 1936–78. Can. J. Fish. Aquat. Sci. 37(11),

1780–1801.

Tessier, L.R., Long, T.A.F., Wilkie, M.P., 2018. Influence of body size, metabolic rate and life

history stage on the uptake and excretion of the lampricide 3-trifluoromethyl-4-

nitrophenol (TFM) by invasive sea lamprey (*Petromyzon marinus*). Aquat. Toxicol. 194,

27–36.

Wilkie, M.P., Hubert, T.D., Boogaard, M.A., Birceanu, O., 2019. Control of invasive sea

lamprey using the piscicides TFM and niclosamide: Toxicology, successes & future

prospects. Aquat. Toxicol. 211, 235–252.

Wilkie, M.P., Tessier, L.R., Boogaard, M., O'Connor, L., Birceanu, O., Steeves, T.B., Sullivan,

W.P., 2021. Lampricide bioavailability and toxicity to invasive sea lamprey and non-

target fishes: The importance of alkalinity, pH, and the gill microenvironment. J. Great

Lakes Res. 47, S407–S420.

Wong, T.T., 2015. Performance evaluation of classification algorithms by *k*-fold and leave-one-

out cross validation. Pattern Recognit. 48(9), 2839–2846.

Yin, X., Martinez, A.S., Sepúlveda, M.S., Christie, M.R., 2021. Rapid genetic adaptation to

recently colonized environments is driven by genes underlying life history traits. BMC

Genomics 22(1), 269.

     Figure legends

Figure 1. Distribution of sea lamprey exposed to TFM. The 7,651 lamprey exposed to TFM

$(2.52 \pm 0.08$ mg/L) survived between two and 18 hours of exposure (A). There was a highly

allometric relationship between larval total length (mm) and larval mass (g), as illustrated by the

805 fish surviving less than three hours of exposure, representing 10.5% of our samples (orange;

referred to as lower quantile) and the 345 fish surviving seven hours or beyond of exposure,

representing the upper 4.51% of our samples (blue; referred to as upper quantile) (B). The lower

quantile lamprey had masses between 0.074 and 1.801 g ($\mu = 0.694$ g; orange bars; C top) and

total lengths between 46.0 and 111.0 mm ($\mu = 80.18$ mm; orange bars; C bottom). The upper

quantile lamprey had masses between 0.333 and 4.529 g ($\mu = 1.129$ g; blue bars; C top) and total

lengths between 69.0 and 137.5 mm ($\mu = 94.27$ mm; blue bars; C bottom).

Figure 2. Predicted survival time (hrs) for the 10th, 25th, 50th , 75th, and 90th mass and total

length quantiles. (A) Relationship between mass (g) quantiles and predicted survival time (hrs)

calculated using Model A (reported in text; see Table 1 for full model results). Line colors

indicate the corresponding total length quantile. (B) Relationship between total length (mm)

quantiles and predicted survival time (hrs) based on Model A (reported in text; see Table 1 for

full model results). Line colors indicate the corresponding mass quantile.

Figure 3. Relationship between body length and survival time of larval sea lamprey exposed to

TFM (2.52 ± 0.08 mg/L) for up to 18 hours (n = 7,651 larvae). Zero mortality occurred at hours

one and 12. Five larval sea lamprey survived beyond 12 hours of TFM exposure (circled in red).

Table 1. Linear mixed effect model results for Model A (reported in text). Model A is a linear

mixed effect model evaluating the contribution of total length (mm), mass (g), toxicity units, and

their interactions to survival time (hour). Experiment, tank, and bag are a nested random

variable. Coefficient Estimate, 95% Confidence Interval, t value, p-value, and degrees of

freedom (df) are from the model summary in R.

Coefficients	Estimate	95% CI	t	p	df
(Intercept)	6.712	[6.010, 7.410]	18.49	5.691E-32	87.46
total_length	0.02686	[0.01744, 0.03633]	5.573	2.592E-08	7633
mass	0.4590	[0.09174, 0.8280]	2.444	0.01455	7640
tox_units	-81.76	[-100.9, -62.35]	-8.304	1.180E-16	7638
total_length:mass	-0.006892	[-0.008805, -0.004987]	-7.076	1.621E-12	7642
total_length:tox_units	-0.7365	[-1.027, -0.4471]	-4.978	6.577E-07	7635
mass:tox_units	21.15	[13.16, 29.13]	5.191	2.143E-07	7638

Figure 1

Figure 2

A

B

Figure 3