Ancestral and recent bursts of transposition shaped the massive genomes of plant pathogenic rust fungi

Authors:

Emma Corre[1], Emmanuelle Morin[1], Sebastien Duplessis[1*] and Cecile Lorrain[2*]

*Corresponding authors: sebastien.duplessis@inrae.fr; cecile.lorrain@usys.ehz.ch

[1] Université de Lorraine, INRAE, UMR 1136 IAM, F-54000, Nancy, France

[2] Plant Pathology Group, Institute of Integrative Biology, ETH Zürich, Zürich, Switzerland

Abstract

Background

Transposable elements (TEs) play a crucial role in genome evolution, influencing gene regulation, diversity, and genome architecture. Rust fungi of the order Pucciniales (Subphylum Pucciniomycotina) are the largest group of obligate biotrophic plant pathogens and harbor some of the largest and most TE-rich genomes - up to 1.2Gb - compared to other fungi. This global genomic expansion contrasts with the smaller genomes and minimal mobilome found in other Pucciniomycotina species. Despite the availability of high-quality genome assemblies, our understanding of TE dynamics in Pucciniales remains limited due to inconsistent and incomplete TE annotations.

Results

We analyzed the mobilomes of 12 Pucciniomycotina species, producing a manually curated TE library for each genome. In Pucciniales, TEs occupy 47–92% of the genome, whereas 2-36% of TEs are detected in other Pucciniomycotina genomes. The comparison of gene and TE repertoires indicates that TEs, particularly LTR-retrotransposons and TIR-DNA transposons, are the primary contributors to the genome expansion of the Pucciniales. We reconstructed the proliferation histories of TEs in the Pucciniales, combining sequence similarity, clustering, and molecular clock approaches. We highlight recent and ancient TE invasions with some LTR-Gypsy elements predating the divergence of Pucciniomycotina (~176 Mya), while most TE accumulation in Pucciniales occurred within the last 50 Mya. However, the TE invasions in the Pucciniales genomes do not seem to result from specific deficiencies in known TE-control mechanisms.

Conclusion

Our findings uncover extensive TE proliferation in Pucciniales, predominantly driven by LTR-Gypsy expansions. The retention of ancestral TEs and the consistently TE-rich genomes

40   observed in Pucciniales highlight TE proliferation as an ancestral genomic feature in rust

41   fungi.

42   **Keywords**: transposable elements, fungal genomics, rust fungi, plant-pathogens

BACKGROUND

Rust fungi (order Pucciniales) are ubiquitous plant pathogens that infect a wide range of vascular plants, including some of the world's most important crops and trees, such as wheat, soybean, poplar, and coffee trees [1, 2]. Epidemics caused by rust diseases can lead to significant yield losses, posing a substantial threat to global food security [2–4]. The Pucciniales are classified within the subphylum Pucciniomycotina, phylum Basidiomycota, in the Fungal Kingdom and represent the largest order of obligate biotrophic fungi, comprising over 8,000 species [5]. The Pucciniales are estimated to have emerged between 175-230 million years ago [5, 6]. Phylogenetic studies suggest that the diversification of rust fungi is closely tied to their obligate host association and coevolution, with occasional host jumps to unrelated species [7]. The obligate host association has shaped the genomic architecture of rust fungi. For instance, Pucciniales genomes show a reduced repertoire of plant cell wall-degrading enzymes, secondary metabolites, and transporter genes, typical traits of obligate biotrophic fungi [8–10]. However, Pucciniales genomes are exceptionally large, ranging from approximately 80 Mb to over 1.2 Gb [11, 12], and are expected to present even greater genome sizes [13], contrasting with the average ~46.5 Mb genome size commonly found in Basidiomycota [14]. These expanded genomes are characterized by lineage-specific multigene families, an abundance of secreted proteins encoding genes, and a high proportion of repetitive elements [8, 9, 11]. The genome size expansion observed in Pucciniales parallels other filamentous plant pathogens. For instance, several Ascomycetes and Oomycetes, such as *Blumeria graminis* and *Phytophthora infestans*, also possess large genomes exceeding 100 Mb, similarly characterized by a significant proportion of repetitive elements [15]. The high repeat content of Pucciniales genomes has posed challenges in generating high-quality genome assemblies, which are crucial for studying their organization and evolutionary dynamics.

The increasing affordability of new-generation sequencing technologies has revolutionized

the sequencing of high-quality reference genomes for numerous non-model organisms,

including species within the Pucciniales order [11, 16, 17]. The most recent genome

assemblies consist of gapless and phased chromosomes generated from dikaryotic spores

[11, 18]. Analyses of the most recent Pucciniales genomes have confirmed that transposable

elements (TEs) are major drivers of genome architecture and evolution in rust fungi [11, 17].

TEs are ubiquitous mobile DNA sequences capable of proliferating within their host

genomes. TEs are classified into two main classes based on transposition mechanisms [19].

Class I elements, or retrotransposons, replicate through an RNA intermediate, enabling their

insertion into new genomic positions. Class II elements, or DNA transposons, move either

through direct excision from double-stranded DNA or via single-strand excision followed by a

rolling-circle replication mechanism [19]. These two major classes are further organized into

Orders and Superfamilies (SFs) based on their sequence architecture [19]. Within each

class, elements are grouped into Families that share phylogenetic ancestry. Both classes

include autonomous elements, which encode the enzymatic machinery necessary for their

own mobilization, and non-autonomous elements, which depend on the transposition

machinery provided by autonomous counterparts. In Class I TEs, autonomous elements

include for instance long terminal repeat (LTR) elements and long interspersed nuclear

elements (LINEs). In Class II TEs, autonomous elements primarily consist of terminal

inverted repeat (TIR) elements and Helitrons [19]. Conversely, non-autonomous elements

encompassing Miniature Inverted-repeat Transposable Elements (MITEs), Short

Interspersed Nuclear Elements (SINEs), Terminal-repeat Retrotransposons in Miniature

(TRIMs), and LARDs (Large Retrotransposon Derivatives), which lack the coding capacity

for independent transposition [19].

TE-mediated genomic plasticity may confer significant adaptive advantages in plant-

pathogen interactions. TEs contribute to genomic plasticity by facilitating chromosomal

rearrangements, promoting compartmentalization, duplicating or deleting genes, and

influencing gene expression [20]. In the soybean rust *Phakopsora pachyrhizi*, a subset of

TEs is expressed alongside effectors during the early stages of infection, suggesting that

transposition activity may be influenced by the plant-rust interaction, as also reported in

other pathosystems [11, 21]. There is also evidence that fungal genomes have co-opted

some TEs [22]. For instance, the effector AvrK1 in the TE-rich *B. graminis* f. sp. *hordei*

genome is derived directly from a LINE retrotransposon [23]. The interplay between TE

expansion and transposition regulation is largely unknown in TE-rich fungal species,

including Pucciniales. However, these species appear to have evolved tolerant genomes

that can accommodate TE activity, unlike other fungal species with compact genomes [24].

This tolerance may play a crucial role in the adaptability of rust fungi, contributing to their

success as plant pathogens.

TE propagation can harm host genomes, and species have evolved defense mechanisms to

regulate TE proliferation. In fungi, TE activity is often controlled through epigenetic silencing

mechanisms [25, 26]. TEs are frequently associated with heterochromatin histone marks and

cytosine DNA methylation (5mC) [27]. Repeat-induced point mutation (RIP) is also a defense

mechanism that has been so far specifically found within Ascomycetes [8]. RIP inactivates

TEs through cytosine-to-thymine mutations and involves the DNA methyltransferases DIM2

and RID [25, 27, 28]. To date, RIP has not been functionally characterized in

Basidiomycetes, although RIP-like mutational signatures have been reported in the genomes

of some smut fungi [29]. Notably, the canonical RIP-associated DNA methyltransferases,

DIM2 and RID, are absent from the gene repertoires of multiple Basidiomycete species,

including the Pucciniales [30]. RNA interference (RNAi) adds another layer of TE control.

The RNAi pathway relies on three core components: Dicer, which processes double-

stranded RNA into small interfering RNAs (siRNAs); Argonaute, which incorporates these

siRNAs to guide the RNA-induced silencing complex (RISC); and RNA-dependent RNA

polymerase (RdRP), which amplifies siRNAs to sustain the silencing response [31]. This

pathway targets TE transcripts for degradation or chromatin modifications, thereby

repressing TE expression and preventing transposition [32]. Despite its demonstrated

importance in plants and animals, RNAi-mediated TE regulation in fungi remains

comparatively less studied. Comparative genomic studies suggest that fungi exhibit

significant diversity in their TE control mechanisms, with some species retaining these

pathways while others have lost them through evolution [25, 27, 31].

The present study investigates the role of TEs in shaping Pucciniales genome architecture

over evolutionary timescales. We tested four key hypotheses: (i) TE invasions in Pucciniales

are the result of an ancestral transposition burst that has been preserved, (ii) TE invasions in

Pucciniales arise from species-specific independent bursts, (iii) TE proliferation reflects a

combination of ancestral bursts retained within the genome and recent species-specific

transposition events, and (iv) TE proliferation is an ancestral characteristic of

Pucciniomycotina evolution, which has been largely lost in other species but retained in

Pucciniales. To achieve this, we generated a comprehensive annotation and curation of

mobilomes across 12 selected Pucciniomycotina species with comparative genomic

analyses, focusing on TE diversity, abundance, and evolutionary dynamics. Additionally, we

compared gene content and patterns of multigene family expansion and contraction across

Pucciniomycotina genomes to detect potential evidence of TE control presence/absence.

RESULTS

The massive genome size of Pucciniales is determined by their extensive mobilomes

and large gene catalogs

To explore the contributors to the genome size expansion in Pucciniales, we selected a

panel of 12 Pucciniomycotina species with reference genomes comprising eight Pucciniales

species and four other Pucciniomycotina species (Fig. 1A; Table S1). We also included the

complete genomes of two *Puccinia graminis* f. sp. *tritici* strains to compare intra-specific

variation. The size of the nine selected Pucciniales genomes ranges from 109 Mb

(*Melampsora larici-populina*) to 1.057 Gb (*P. pachyrhizi MT2006*), contrasting with the much

smaller genomes of the four other Pucciniomycotina genomes, with sizes ranging from 20

Mb (*Rhodosporidium toruloides*) to 26 Mb (*Microbotryum lychnidis-dioicae*) (Table S1; Fig.

1B). Assemblies included either primary scaffolds from unphased builds or fully separated

haplotype-phased scaffolds. BUSCO (basidiomycota_odb10) completeness was uniformly

high (≤ 10 % missing for all, except *Leucosporidiella creatinivora* at 10.6 %), and most

showed low duplication (≤ 10 % duplicated BUSCOs). However, the three Pucciniales

assemblies *M. allii-populina* (22 %), *M. americana* (44 %) and *P. pachyrhizi* (36 %), exhibited

elevated BUSCO duplication, consistent with uncollapsed haplotypes in these dikaryotic,

repeat-rich genomes. To mitigate potential bias for comparisons, phylogenetic analyses

were made on single-copy orthogroups and performed TE comparisons on a global genomic

scale.

We first compared gene and TE contents and their contribution to the genome inflation of

Pucciniales. Pucciniales genomes harbor substantially more genes ranging from 15,984

genes in *Melampsora americana* to 19,618 genes in *P. pachyrhizi,* compared to the other

Pucciniomycotina species with 7,819 genes in *M. lychnidis-dioicae* to 9,854 in *L. creatinivora*

(Table S1). Although gene coverage showed a slight but significant positive relationship with

genome size ($r^2 = 0.42$, $p < 0.05$), TE content appeared as the primary driver of genome

inflation in Pucciniales, correlating very strongly with genome size ($r^2 = 0.91$, $p < 0.001$;

FigS1). In addition, the TE coverage in Pucciniales genomes ranges from 47% to 92.3% in

M. larici-populina and *P. pachyrhizi*, respectively (Fig. 1B; Table S2). We observed variation

of the TE coverage within the Pucciniales order and between closely related species. For

instance, in the Melampsoraceae, TE content represents 47.0% of the 109 Mb genome of *M.*

larici-populina to 83.4-84% of the 239.9-242.3 Mb genome of *Melamspora lini C* and *M. lini*

H, respectively. In the Pucciniaceae, TEs covered from 49.3-49.5% of the 75.6-75.9 Mb

genome of *Puccinia striiformis* f. sp. *tritici* A and B*,* respectively, to 77.1-77.4% of the 123.9-

121.6 Mb genome of *Puccinia triticina* A and B*,* respectively (Fig. 1B). In comparison, the TE

coverage in the other Pucciniomycotina ranges between 2.3% in *R. toruloides* and 16.2% in

M. lychnidis-dioicae. Finally, haplotype-phased Pucciniales assemblies display nearly

identical TE proportions between phases, further validating our global-TE comparison

approach. Together, these results highlight that TE accumulation, more than gene number,

underlies the extensive genome expansion in Pucciniales relative to other Pucciniomycotina.

LTR retrotransposons are the major contributors to the extensive mobilome and

genome size expansion in the Pucciniales

To detect if TE proliferation in the Pucciniales is linked to specific elements, we further

examined the TE landscapes in the genomes of the selected Pucciniomycotina species

using REPET v3.0 followed by three rounds of manual curation to build a high-quality

Pucciniomycotina TE library. In total, we annotated 39, 834 TE consensus sequences,

constituting the Pucciniomycotina TE library. TE annotation before curation detected 20, 958

(52.6%) classified TE consensus sequences, 16,811 (42.2%) unclassified TE consensus

sequences, and 2,065 (5.2%) TE consensus sequences with ambiguous annotations (or

"conflicts"), which occur when the classifier cannot assign a TE to a single category due to

nested or overlapping elements (Fig. S2). After the first round of manual curation, we

reduced the number of ambiguous annotations to 651 (1.6%) TE consensus sequences but

could not resolve unclassified annotations. Therefore, we used MCHelper [33] and improved

the annotation of ~16% (2, 700) of previously unclassified TE consensus sequences (Fig.

S2). We used a final step during which we clustered all TE consensus sequences from the

Pucciniomycotina TE library to reduce redundancy and correct the remaining ambiguous

annotations. Our final curated Pucciniomycotina TE library comprised 28,868 (72.5%)

classified TE consensus sequences and 10,966 (27.5%) unclassified TE consensus

sequences (Tables S2-S4).

On average, the Pucciniales genomes are covered by 33.9% of Class I TEs, ranging from

15.4% (16.9 Mb) in *M. larici-populina* to 51.7% (515.8 Mb) in *P. pachyrhizi*, and 23.5% of

Class II TEs ranging from 13.2% in *M. lini* haplotype C (31.6Mb) to 31.8% in *Puccinia triticina*

haplotype A (39.3 Mb) (Fig. 2A; Table S2). By contrast, the Class I and Class II genome

coverages were drastically lower within the other Pucciniomycotina, with an average of

13.7% and 3.5%, respectively (Fig. 2A; Table S2). Class I elements represent the most

abundant elements in the Pucciniales except for the three genomes of *M. larici-populina* with

15.4% Class I (16.9 Mb), *Melampsora americana* with 18.8% Class I (21.1 Mb) and

P. striiformis f. sp. *tritici* 18.1% Class I (13.7 Mb) (Fig. 2A; Table S2). These results indicate

an important contribution of Class I elements in the genome expansions compared to the

Class II elements in six of the eight Pucciniales species.

The TE Orders content of the Pucciniales genomes predominantly comprises LTR

retrotransposons and TIR DNA transposons. LTRs represent 12.5% of the genome in *M.*

larici-populina (13.1 Mb) to 63.0% in *M. lini* haplotype C (151.1 Mb), respectively (Fig. 2B-C;

Table S3). At the LTR super-family level, we mostly found Copia and Gypsy elements,

representing 2.9% of genome (3.2 Mb of LTR-Copia in *M. larici-populina*) to 12.7% (30.5 Mb

in *M. lini* haplotype C) and 5.5% (6.2 Mb in *M. americana*) to 48.2% (115.6 Mb in *M. lini*

haplotype C), respectively (Fig. 2 D-E; Table S4). The non-LTR retrotransposons, such as

LINE elements, represent from 0.4% to 2.6% of the Pucciniales genomes. The non-

autonomous elements such as SINE represent less than 0.2% of the Pucciniales genomes

(Tables S3 and S4). In the other Pucciniomycotina genomes, the amount of LINE covers up

to 3.3% of the genome of *M. intermedium* (0.77 Mb). TIRs represent most DNA transposons

within the Pucciniales genomes, with 10.2% to 28.1%, followed by non-autonomous MITE

representing 0.6% to 5.0% (Table S3). Interestingly, some TIRs super-family like CACTA,

hAT, MuDR, PIF-harbinger, and Tc1-Mariner are found only in the Pucciniales and not in the

other Pucciniomycotina (Fig. 2 D-E; Table S4). The Helitron elements cover a small

proportion of the genome, from 0.2% in *M. lini* haplotype C (6.2 Mb) to 2.6% in *P. triticina*

haplotype A (3.2 Mb). These results indicate that at the TE-Order level, the Pucciniales

display an overall similar TE landscape with a majority of LTRs and TIR elements,

suggesting ancient proliferation of TEs in the order Pucciniales. However, the variable

amount of LTR and TIR elements found between Pucciniales genomes, even at the

haplotype level, support additional bursts of transposition that occurred throughout the

evolution of the order Pucciniales.

Temporal dynamics of TE activity shaped the genome architecture of Pucciniales

We hypothesized that the proliferation of TEs observed in the Pucciniales could have

emerged through different scenarios throughout the evolution of the Pucciniomycotina. First,

we analyzed genome-wide TE divergence profiles using Kimura 2-parameter (K2P)

substitution rates. In this framework, low K2P values signify recent TE insertions with

minimal divergence from the consensus, whereas higher values denote ancient insertions

indicative of long-term genomic retention and mutation accumulation. Our comparative

landscape analysis across 12 Pucciniomycotina species revealed species-specific TE

activity patterns (Fig. 3A–P). Among non-Pucciniales species, TE proliferation appears

minimal. For instance, *R. toruloides* and *L. creatinivorum* exhibit low recent TE activity, with

less than 3% of their genomic content composed of insertions with K2P < 0.05. Conversely,

M. intermedium and *M. lychnidis-dioicae* demonstrate moderate levels of recent activity, with

8.5% and 10.5% of their genomes, respectively, reflecting young TE insertions (Fig. 3A–D).

Within Pucciniales, two distinct TE proliferation patterns emerge. Species from the

Melampsoraceae family show evidence of dual TE expansion waves with recent bursts (K2P

< 0.05) accompanied by older insertions peaking between 0.4 and 0.6 K2P (Fig. 3E–I).

Notably, *M. lini* presents a prominent accumulation of ancient LTR elements, indicating

substantial historical activity. In contrast, members of the Pucciniaceae exhibit a

predominance of recent insertions with a marked peak at K2P < 0.05 and a flatter

distribution at higher divergence values, suggesting reduced retention of older TE

sequences. In addition, we used a within genome sequence similarity-based approach to

compare TEs from the oldest insertions to the most recent insertions as follows: i) divergent

TE consensus sequences containing copies with less than 85% sequence identity, ii)

intermediate TE consensus sequences with copies of 85% to 95% sequence similarity and

iii) conserved TE consensus sequences comprising copies with more than 95% sequence

identity [11, 12]. The average sequence identity of all TE consensus sequences per genome

ranges from 80.6% (*R. toruloides*) to 90.6% (*M. lychnidis dioicae*). We observed 13.0--66.7%

divergent (*M. lychnidis-dioicae* and *R. toruloides*, respectively), 33.3-76.5% intermediate (*R.*

toruloides and *M. intermedium*, respectively), and 0.0-16.5% conserved (*R. toruloides* and

M. lychnidis-dioicae, respectively) TE consensus sequences in the other Pucciniomycotina

(Table S5). Within the Pucciniales, TE consensus sequences similarity profiles revealed

30.0-66.2% divergent (*P. pachyrhizi* and *M. larici-populina*, respectively), 27.4-65.6%

intermediate (*M. americana* and *P. pachyrhizi*, respectively), and 2.7-7% conserved TE

consensus sequences (*M. allii-populina* and *P. triticina*, respectively) (Table S6). Among the

Pucciniales genomes, we observed that LTRs and TIRs compose most of each relative TE

age category (Table S6). LTR elements represent 17.4-33.6% of the divergent TEs (in *M.*

americana and *M. allii-populina*, respectively), while TIR elements represent 26.1-38.0% of

the divergent TEs (in *P. pachyrhizi* and *M. larici-populina*, respectively). In the conserved

TEs, LTR elements represent 7.2-34.4% (in *M. larici-populina* and *M. lini*, respectively), and

TIR elements represent 7.6-33.3% (in *P. graminis* f. sp. *tritici Ug99 haplotype C* and *M. larici-*

populina, respectively) (Table S6). The heterogeneity in the composition of TEs with different

relative conservation profiles (i.e., relative insertion ages) in the Pucciniales suggests that

multiple waves of TE proliferation occurred since the early divergence between the

Pucciniales and the other Pucciniomycotina species. The subset of highly conserved

elements in all genomes also indicates recent TE transposition events.

Based on sequence similarity among all TE consensus sequences, we employed a

clustering approach to identify TE families shared among or specific to species within the

Pucciniales order. We found no TE family shared across all the Pucciniales species.

However, we identified three clusters that are specific to the Melampsoraceae and one that

are specific to the Pucciniaceae (Fig. S3). These findings suggest that the accumulation of

ancient TE copies predated the divergence of the family Melampsoraceae and the family

Pucciniastraceae (58±17 million years ago, according to Aime and McTaggart 2021 [2])

on one hand and the divergence of the family Pucciniaceae and the family

Sphaerophragmiaceae (approximately 68±17 million years ago, according to Aime and

McTaggart 2021 [2]) on the other hand. Interestingly, the three Melampsoraceae-specific

TE clusters were all TIR elements, whereas the Pucciniaceae-specific clusters comprised

one LTR cluster (Fig. S3; Tables S7 and S8). Additionally, we observed a predominance of

species-specific TE clusters, ranging from 650 clusters (3.8% of the mobilome) in *P.*

graminis f. sp. *tritici* 20-1 to 3,174 clusters (15.6% of the mobilome) in *P. triticina* (Fig. S3).

Within these species-specific clusters, TIR elements accounted for 3.5% of the mobilome in

M. americana (608 clusters) but only 0.05% (9 clusters) in *P. graminis* f. sp. *tritici* 21-0.

Interestingly, we identified haplotype-specific TE clusters in *M. lini* exhibiting 744 clusters

(4.3% of its mobilome), whereas *P. graminis* f. sp. *tritici* showed between 42 clusters in Ug99

and 45 clusters in 20-1 haplotypes (Fig. S3). These findings highlight the variation in TE

composition and specificity across species and even between haplotypes within individuals.

Finally, we retraced the evolutionary history of LTR invasion using their terminal repeats with

Kimura molecular clock estimates. Across all Pucciniales genomes, LTRs were found to

have accumulated over a broad temporal range, from 618 Mya to the present (Fig. S4). The

oldest Gypsy elements were identified in *P. pachyrhizi*. 0.1% of LTRs accumulated before

176 Mya, thus predating the divergence between the Pucciniales and the other

Pucciniomycotina (Fig. S4). However, our analysis suggests that LTR accumulation within

the Pucciniales genomes, more particularly, spiked in the last 50 Mya, with a notable

acceleration reflecting LTR expansions occurring within the last 5 Mya (Fig. 3S-T; Fig. S4).

We also identified two distinct proliferation profiles in the Pucciniales genomes. The first

profile, observed in *M. lini* and other Melampsoraceae species, exhibits sharp bursts of

Gypsy LTR activity, with up to 2,000 elements accumulating during specific intervals (e.g., 7-

9 Mya, 6-4 Mya, and 3-2 Mya) (Fig. 3S; Fig. S4). The second profile, characteristic of

Phakopsoraceae and Pucciniaceae species, demonstrates a more gradual accumulation of

LTRs over time (Fig. 3S-T; Fig. S4). These results collectively indicate that while genome

inflation driven by TE invasion is a shared feature of Pucciniales, multiple waves of

proliferation have shaped their evolutionary trajectories, reflecting diverse temporal

dynamics of TE activity across lineages.

The TE invasion of the Pucciniales appears unrelated to deficiency in TE control

pathways

We investigated whether the TE invasion in Pucciniales compared to the other

Pucciniomycotina, might be due to a loss of control mechanisms. The most studied control

mechanism against TE proliferation in fungi is the induction of cytosine-to-thymine mutations

in repetitive regions through RIP mutations [25]. Using the composite RIP index in TE

sequences and in genome-wide 1-kb sliding windows, we quantified RIP-like signatures in

TE family consensus sequences in the Pucciniales and the other Pucciniomycotina

genomes. In all Pucciniomycotina genomes we detected less than 0.5% of the genomes

containing RIP-like signatures (Table S9). We estimated the average RIP composite index

per TE family ranging from -0.66 in *M. americana* to -0.15 in *L. creatinivora*, indicating the

lack of RIP-like signatures in TE families among the Pucciniales and the other

Pucciniomycotina TE family sequences (Table S9). Additionally, the two DNA

methyltransferase genes associated with RIP (DIM2 and RID) are absent from both the

Pucciniales and the other Pucciniomycotina genomes. We identified two other DNA

methyltransferase genes DNMT5 and DNMT1 present across all Pucciniomycotina

genomes, except for *P. hordei* in which we did not find DNMT1 (Fig. S5). Finally, we

investigated the presence or absence of genes involved in the RNAi pathway, namely

Argonaute, Dicer, and RNA-dependent RNA polymerase RdRP. We identified 25-72 (*P.*

hordei and *P. triticina*) genes related to Dicer-like genes in the Pucciniales and 25-29 Dicer-

like in the other Pucciniomycotina genomes (Fig. S6). Similarly, we identified 17-60 (and *M.*

lini) RdRP-like genes in the Pucciniales and 12-15 RdRP-like genes in the other

Pucciniomycotina genomes (Fig. S6). We observed a remarkable expansion of AGO1 solely

in the *P. pachyrhizi* genome that displays 572 copies compared to a mean of 91 copies in

the other species (Fig. S5). However, it remains to determine whether these copies are truly

functional. All Pucciniomycotina genomes, regardless of their TE content, are equipped with

genes potentially involved in TE control through DNA methylation silencing and RNAi

pathways. We conclude that the TE proliferation observed in the Pucciniales genomes

compared to the other Pucciniomycotina is not attributable to the loss of genes involved in

TE control mechanisms and related to RIP, DNA methylation, or the RNAi pathway.

Finally, we analyzed gene family dynamics within the Pucciniomycotina subphylum to

investigate potential evidence of loss of genes that may relate to TE control mechanisms in

Pucciniales outside of the canonical mechanisms investigated above. For this, we

determined multigene families within the Pucciniomycotina using *Ustilago maydis* as an

outgroup and identified 43,847 orthogroups (Table S10). Of these, 3067 core orthogroups

(17,1758 genes) were shared across all genomes, including *U. maydis*, while 16, 585

orthogroups (85,786 genes) were specific to Pucciniomycotina (Table S10). We observed

substantial variation in species-specific orthogroups ranging from 68 in *M. intermedium* to

1310 in *P. triticina.* We identified 23 orthogroups in contraction and 380 orthogroups in

expansion at the separation of the Pucciniales and the other Pucciniomycotina (Fig S7). The

orthogroups in contraction in the Pucciniales comprise a total of 483 genes, of which 19.5%

(94 genes) are of unknown functions (Table S11; Fig. S8). The five most represented

functions within the remaining orthogroups in contraction in the Pucciniales, include 62

genes related to "Transcription", 52 genes related to "Signal transduction mechanisms", 52

genes with function related to "Amino acid transport and metabolism", 25 genes related to

"Posttranslational modification, protein turnover, chaperones", 23 genes related to

""Secondary metabolites biosynthesis, transport and catabolism" and 21 genes related to

"Chromatin structure and dynamics"(Fig S8; Table S11). However, we identified only 22

orthogroups with a significant accelerated gene gain/loss between the Pucciniales and the

other Pucciniomycotina genomes and all the 22 orthogroups are in expansion in the

Pucciniales (p-value < 0.05; Fig S9). These results show that Pucciniales underwent

extensive gene losses compared to other Pucciniomycotina species. Based on the functional

categories identified in this comparative analysis, it remains possible that certain gene

losses may be associated with the loss of functions related to TE control within the

'chromatin structure and dynamics' category or other, as yet unidentified, functions.

DISCUSSION

In this study, we provide a better understanding of how TEs have profoundly shaped the

genomes of Pucciniales, contributing to their distinctive genome expansions within the

Pucciniomycotina subphylum. Our comprehensive TE annotation across 12

Pucciniomycotina genomes demonstrates that the Pucciniales possess a mobilome

landscape enriched in LTR Gypsy and TIR elements, a feature contrasting sharply with the

smaller genomes and the minimal TE contribution in other Pucciniomycotina species. Our

analyses examined four hypotheses on TE proliferation in Pucciniales and evidenced both

ancient and recent TE bursts, especially from Gypsy elements, aligning with the third

hypothesis: TE proliferation is driven by a combination of ancestral transposition bursts

retained in the genome and recent species-specific independent transposition events

throughout Pucciniales evolution. Hence, we uncover ancient TE proliferation likely predating

the early divergent branch of the order Pucciniales, followed by further species-specific

expansions. In addition, despite the extensive TE presence, our findings indicate no notable

deficiency in known TE regulatory mechanisms in the Pucciniales genomes compared to the

other Pucciniomycotina genomes, suggesting that Pucciniales may have rather evolved

tolerance to TE proliferation or deployed other mechanisms to accommodate their abundant

presence in the genome.

The accumulation of TEs in Pucciniales is an ancestral genomic trait. Pucciniales is a

monophyletic group of species [5], all possessing inflated genomes enriched in TEs. We

identified retrotransposon copies retained in Pucciniales genomes that predate this

divergence, suggesting a long-term persistence of ancestral TEs within rust fungal genomes

[5]. This TE retention pattern is consistent with observations in previous works of rust fungal

genomes, which often harbor highly divergent, long-standing TE families [17]. Interestingly,

we did not find any TE families shared among all *Pucciniales* species in our panel,

suggesting that ancestral copies have diverged substantially over time. However, we

detected TE families consistently retained at the genus level, including one (LTR-Copia)

within the family *Pucciniaceae* and three others (TIR-Unknown, TIR-hat, and TIR-

Tc1Mariner) within the family *Melampsoraceae*. The genus-specific conservation of certain

TEs suggests that ancestral elements have been retained in distinct subgroups, which

mirrors patterns observed in other fungal species. For example, a recent analysis of

transposable element repertoires across 24 arbuscular mycorrhizal fungal genomes

revealed the persistence of ancient TEs over long evolutionary periods [34]. Likewise, in

large, complex plant genomes such as wheat and maize, retrotransposons are retained

across extended timescales [35, 36]. The long-term retention of TEs across such diverse

fungal and plant lineages suggests a common pattern of TE evolutionary dynamics,

underlying their potential role in shaping their host-genome architecture.

Gypsy retrotransposons represent the largest proportion of TE content across Pucciniales

genomes, followed by *Copia* retrotransposons and TIR elements. This abundance of

retrotransposons is a common feature in other TE-rich fungal genomes, such as those of

B. graminis and *Entomophthoraceae* spp. [37, 38]. We hypothesize that ancient *Gypsy*

elements found in the order Pucciniales may have served as progenitors for more recent

lineage-specific TE expansions observed across the Pucciniales genomes. Indeed, the

dynamics of *Gypsy* element accumulation display extreme variation among Pucciniales

species. For example, we observed a burst accumulation pattern in *M. lini*, marked by a

rapid increase in *Gypsy* copies over a few million years. In contrast, species such as *P.*

pachyrhizi exhibit a more continuous pattern of *Gypsy* accumulations. We also detected

haplotype-specific TE families in the phased genomes of the two species *M. lini* and *P.*

graminis f. sp. *tritici*, indicating that TE dynamics vary not only between species but also

within the two haplotypes in dikaryotic Pucciniales. These differences suggest that *Gypsy*

elements may undergo distinct cycles of activity in each species, with some elements

remaining active and driving ongoing TE expansion while others persist as inactive remnants

within the genome. Notably, the presence of still active elements aligns with the observed

expression of certain TEs during host infection in *P. pachyrhizi* [11], further supporting the

hypothesis that active TEs may still be involved in ongoing genomic changes during

pathogen-host interactions.

TE richness is a shared genomic feature among different groups of fungi, which appears as

a trait associated with obligate parasitic and symbiotic lifestyles. Similarly to the Pucciniales,

other obligate biotrophic plant pathogens, such as *B. graminis*, the causal agent of powdery

mildew in cereals, are characterized by a high proportion of TEs [39]. Enrichment in TE

could enable rapid evolution through genomic plasticity in response to fast-evolving host

immune systems and changing environmental conditions [24]. Outside of plant pathogens,

obligate insect parasites within the fungal order Entomophthorales exhibit TE-rich genomes

[38]. TEs are reported as the main drivers for genome expansion in this clade. TE-rich

genomes extend beyond obligate parasites and represent a key feature of beneficial plant

symbionts. Arbuscular mycorrhizal fungi and ectomycorrhizal fungi exhibit substantial TE

content and expansions, although the role of these TEs in their symbiotic functions remains

scarcely understood [34, 40]. While the evolutionary pressures differ between parasitism and

symbiosis, large TE-rich genomes in both contexts may suggest that TEs could reflect an

evolutionary strategy to facilitate genome plasticity and adaptation to the host environment.

We observed no evidence of specific gene loss associated with known TE regulation

mechanisms in the Pucciniales genomes compared to other Pucciniomycotina. First, RIP-

like signatures are absent in the Pucciniomycotina subphylum which aligns with other

studies centered on basidiomycetes and with the absence of DIM2 and RID genes in

basidiomycetes [27, 41]. In addition, we found the DNA methyltransferases DNMT1 and

DNMT5 in all 12 genomes, suggesting that the whole Pucciniomycotina phylum may

possess the potential for TE silencing mediated by DNA methylation. DNMT1 and DNMT5

could play a role in the establishment and/or maintenance of cytosine DNA methylation

(5mC), as found in plants and animals [27]. In addition, Bewick et al. (2019) [27]

demonstrated that species in the Phylum Basidiomycota displayed the highest 5mC levels

across fungal genomes, with a weak correlation between 5mC levels and repeat content in

the selected species. Similarly, we identified the essential genes of the RNAi pathway in all

12 Pucciniomycotina genomes. Additionally, we confirmed the expansion of Argonaute 1

(AGO1) previously reported in *P. pachyrhizi* [11]. Most of these expanded AGO1-like genes

were described as pseudo-genes surrounded by TEs, suggesting a TE-driven inactivation of

expanded AGO1 gene members in the family. A comparable phenomenon occurs in the

arbuscular mycorrhizal fungus *Gigaspora margarita*, where multiple AGO-like copies lack

conserved PIWI catalytic residues and may have diverged to non-canonical roles or become

pseudogenes [42]. This pattern could be similar to the recent loss of DIM2 activity found in

the wheat fungal pathogen *Zymoseptoria tritici* where an expansion of the DIM2 genes,

followed by inactivation, is thought to have led to TE-derepression [43]. In

Rhizophagus irregularis, several AGO loci localize adjacent to TE insertions and

transcriptional profiling has shown that AGO genes are highly expressed in spores but

downregulated during plant colonization, coincident with transient TE activation [44, 45]. As

our comparative genomic approach could not identify a Pucciniales-specific loss of TE

control, other approaches, such as epigenomics or small RNA sequencing, will be required

to better understand TE regulation mechanisms in rust fungi.

Despite advancements in TE annotation and the availability of high-quality rust genomes, the

absence of a standardized and comprehensive TE library for rust fungi remains a significant

obstacle in performing comparative genomics to address further evolutionary studies [46].

Manually curating TE libraries for highly repeated genomes is labor-intensive and time-

consuming. Here, we generated a three-step approach to curate TE annotations, which can

be applied to many other genomes. This curation has resolved all the ambiguous
annotations and reduced the unclassified TEs from 42.4% to 16% after three rounds of
manual curation. We created a unified rust fungal TE library, enabling consistent and precise
TE identification across different rust fungal species. Still, several TE annotations remain
unresolved after our curation, mainly consisting of elements probably too degraded to be
correctly classified and nested elements. Additional manual curation efforts, supported by
additional genome assemblies from closely related species, could support the refinement of
the classification of these elements by offering more diagnostic features. As large-scale
genomic projects generate vast sequencing data, creating a unified TE library becomes
increasingly crucial for advancing our understanding of the roles of TEs in rust fungal biology
and their evolution. Expanding and refining fungal TE libraries is essential to fully capture the
diversity and dynamics of transposable elements in these genomes, enabling more precise
evolutionary and functional insights in future studies.

CONCLUSION

This study describes manually curated TE catalogs in thirteen Pucciniomycotina species and
provides a new fungal TE database. We report the genome expansions in Pucciniales are
mainly guided by TE proliferation. Our work highlights recent bursts of Gypsy LTR in
Pucciniales while the content in such elements remains relatively poor in other
Pucciniomycotina. We also describe a few conserved TEs within the Pucciniales order linked
to older proliferation events. TE accumulations do not seem to result from Pucciniales-
specific loss of TE control mechanisms. We provided new insights into Pucciniales genome
architecture and favored the hypothesis that TE proliferation is an ancestral characteristic of
the Pucciniales evolution.

METHODS

Genomic data

We selected the genomes and gene annotations of 12 Pucciniomycotina species publicly

available in the JGI MycoCosm database and NCBI repositories (Table S1). We included ten

genomes from the order Pucciniales, comprising four genomes from the family

Melampsoraceae (*M. allii-populina*, *M. lini*, *M. larici-populina*, and *M. americana*), four

genomes from the family Pucciniaceae (*P. graminis* f. sp. *tritici Ug99 and 21-0*, *P. triticina*,

and *P. striiformis* f. sp. *tritici*), and one genome from the family Phakopsoraceae (*P.

pachyrhizi*). Additionally, four genomes representing species from outside the order

Pucciniales but within the subphylum Pucciniomycotina were included (*L. creatinivora*,

M. lychnidis-dioicae, *M. intermedium*, and *R. toruloides*; Table S1). We selected genomes

with either haplotype-phased assemblies, long-read assemblies, or assemblies based on

genetic maps. We assessed genome assembly completeness using BUSCO v5.4.7 with the

basidiomycota_odb10 marker set, which provided a measure of the quality of the selected

genome assemblies (Table S1).

TE detection and annotation

We detected and annotated repeats in Pucciniomycotina genomes using REPET v3.0

(https://urgi.versailles.inra.fr/Tools/REPET, [33]). Repeated elements were identified and

annotated for each genome following a two-step process. First, TE families were identified

using the TEdenovo pipeline, which builds TE family sequences based on repetitive

elements detected in a 100 Mb subset of the genome. This subset-based approach was

applied to enhance workflow efficiency in highly repeated genomes, as described in [34].

Subsequently, the whole genomes were annotated with TEannot using the TE families

generated by TEdenovo. A second round of TEannot was performed as a curation step,

focusing exclusively on TE families with at least one full-length copy. For haplotype-phased

genome assemblies, TEdenovo was run separately for each haplotype (also using 100 Mb

subsets), and the resulting TE family sequences were merged before proceeding with the

TEannot steps. This approach allowed for the identification of potential haplotype-specific

TEs.

Curation of TE annotations

After the automated detection and annotation of TEs, we performed additional steps of

curations of libraries to improve the classifications of elements detected as "conflicts" and

"unclassified" TEs by REPETv3.0. First, the "conflicts" ambiguous TE annotations (i.e. the

classifier cannot assign a TE to a single category) were manually curated and reassigned to

the appropriate TE Class or Order based on sequence similarity, applying a 50% similarity

threshold. In parallel, we used the Manual Curator Helper tool MCHelper [33] to address

redundancy, fragmentation, false positives, and unclassified sequences in TE libraries for

each genome. Briefly, MCHelper extends consensus sequences through iterative rounds of

Blast, extract, and extend (BEE) and refines them into subfamilies based on distance

clustering. False positives are filtered out by removing sequences with low-quality fragments

or homology to multi-copy genes, and the remaining sequences are classified through

homology-based methods. We then pooled all TE libraries into a Pucciniomycotina TE library

to perform an additional manual curation step by clustering the Pucciniomycotina TE library

with CD-HIT v4.8.1 [47]. We grouped TE sequences with 80% similarity across 80% of the

sequence length and re-assigned previously unclassified TEs based on annotated TEs of

the same cluster. The workflow to generate curated TE annotations from multiple genomes

combining automated detection and curation results is available at

https://github.com/Raistrawby/TE_Rust/tree/main. The curated library of TE consensus

sequences has been deposited in the Dfam database (https://dfam.org/home).

TE relative insertion age

The relative age of TE insertions in the Pucciniomycotina genomes was estimated with the similarity-based approach of REPETv3.0, as described in Lorrain et al. (2021) [48]. Briefly, we analyzed sequence divergence between TE copies to their cognate family using the extent of sequence similarity as a proxy to the divergence time of the TE copies within the host genome. The TE copies were classified into three categories: i) conserved, with copies similar up to 95% against their family, ii) intermediate, with a similarity between 85% and 95% and, iii) divergent, with similarity of copies below 85%. In addition, we estimated the sequence divergence of TE copies relative to their consensus sequences, we computed Kimura 2-parameter distances. For each annotated TE copy, the aligned region was extracted and compared to its corresponding consensus sequence. K2P distances were calculated based on the frequency of transitions and transversions and correcting for multiple substitutions following the approach described by Baril et al. (2024)[49] and adapted to our TE annotations (https://github.com/Raistrawby/TE_Rust/tree/main). Briefly, for each TE copy, the corresponding genomic sequence was retrieved based on genomic coordinates and strand orientation. Each extracted sequence was then pairwise aligned to its cognate consensus sequence using matcher [50] and the Kimura 2-parameter distance [51] was computed from the alignment. TE copies with incomplete or failed alignments were excluded from downstream analyses. The resulting Kimura distances were used as a proxy for the relative age of TE insertions.

Insertion age of LTR-retrotransposons

The insertion ages of LTR-retrotransposons were assessed using a molecular clock approach, as described in Gupta et al. (2023) [11]. Full-length LTR-retrotransposons were detected using LTRHarvest v 1.6.1 [52] with default parameters (except for -similar 90 (default 85) and -mintsd 5 (default 4)). Gypsy and Copia LTR-retrotransposons were selected based on the annotations from the repbase_23.12 database. Each element's 3' and 5' long-terminal repeats were extracted and aligned using MAFFT v7.471 [53] to calculate

Kimura's P distances [51]. Gypsy and Copia insertion dates (T) were estimated using the

formula T□=□K / 2r, where K is the distance between the 2 LTRs and is calculated with the

Kimura 2-Parameter distance formula -0.5*log(1 - 2p - q) * sqrt (1 - 2 q)) with p = transition

frequency and q transversion frequency and r is the fungal substitution rate of 1.05×10^{-9}

nucleotides per site per year [51].

Gene family annotations and evolution

We predicted orthologs between the Pucciniomycotina genomes and the reference

U. maydis genome [23], a representative of the Ustilaginomycotina (the closest subphylum

to Pucciniomycotina within Basidiomycete), which we used as an outgroup to identify genes

restricted to the Pucciniomycotina subphylum. Orthogroup analysis was performed using the

software OrthoFinder v2.4.0 (i=2 -M MSA -S diamond -a mafft)[54] on the proteomes. We

chose one haplotypic proteome for species with haplotype-resolved annotation. To assess

the gene family gains and losses throughout the Pucciniomycotina genomes, we first needed

to build a phylogenetic tree; to this end, we used the single copy orthogroups (i.e.

orthogroups containing only one protein of each genome) and infer the tree with *U. maydis*

as an outgroup with Raxml and 500 bootstraps [55]. We used this rooted phylogenetic tree

and the number of genes found per family (i.e. orthogroup) to estimate rates of gene families

gains (i.e., families in expansion) and losses (i.e. families in contraction) at the species-,

genus-, family- and order-levels with CAFE v5.0 [56].

Functional gene annotations and identification of DNA methyltransferases and RNA-

induced silencing complex

We used EggNOG-mapper v2.1.12 [57] to predict functional annotations for all

Pucciniomycotina genomes. We retrieved the PFAM, KOG, KEGG, and GO annotations

from EggNOG-mapper annotations to explore the potential functions of gene families gained

or lost throughout the evolution of the Pucciniomycotina. To identify the DNMTs and the

genes involved in the RISC pathway in the Pucciniomycotina genomes, we extracted the

genes with corresponding PFAM domains. For DNA methyltransferases, we selected genes

with PF00145 (C-5 cytosine-specific DNA methylase) and used orthogroups and BLASTp

searches to differentiate DNMT classes. For the RNA-induced silencing complex, we

selected genes with PF05183 (RNA dependent RNA polymerase), PF16486 (N-terminal

domain of argonaute), PF22389 (Protein Dicer, dsRNA-binding domain), PF09692

(Argonaute siRNA chaperone (ARC) complex subunit Arb1), PF22749 (Arb2 domain),

PF00636 (Ribonuclease III domain), PF04992 (RNA polymerase Rpb1, domain6), PF03871

(RNA polymerase Rpb5, N-terminal domain), PF01191 (RNA polymerase Rpb5, C-terminal

domain). We searched for characterized DNMT and RISC proteins to refine the annotations

and performed BLASTp searches on the putative Pucciniomycotina DNMT and RISC

proteins.

RIP signatures analysis

RIP-like signatures were searched by estimating the RIP composite index [25] and GC

content in TE family sequences of all Pucciniomycotina genomes, as described in Feurtey et

al. (2023)[58]. RIP Composite index values were calculated using the formula: (TpA/ApT) –

(CpA + TpG/ApC + GpT) applied per TE family sequence. A region is considered affected by

RIP if the Composite index exceeds 0 [41]. In addition, we performed a 1kb genome-wide

sliding window RIP-like signature detection using TheRIPper [59].

Data availability

The accession numbers and links to publicly available genome assemblies and gene

annotations used in this study can be retrieved from Table S1. The TE annotations and gene

functional predictions generated in this study, are openly available at

https://doi.org/10.5281/zenodo.14616359.

Code availability

The TE annotation and manual curation pipeline, as well as additional scripts used to create the results presented in this manuscript, are available in the following GitHub repository: https://github.com/Raistrawby/TE_Rust/tree/main.

Authors contributions

EC and CL wrote the manuscript. SD and EM revised the manuscript. EC, CL, and SD designed, analyzed, and interpreted the data. EM provided comparative genomics expertise and bioinformatics technical support. CL and SD contributed to the conception of the work.

Acknowledgments

Jana Sperschneider and Peter Dodds are acknowledged for providing early access to the phased assembly and genome annotation of the *M. lini* reference genome.

Funding Declaration

EC was supported by a PhD grant from the French National Institute for Agriculture, Food and Environment (INRAE). EC, EM and SD were supported by the French Plan Investissement d'Avenir (PIA) Lab of Excellence ARBRE [ANR-11-LABX-0002- 01]. CL was funded by an SNSF Ambizione grant (PZ00P3_209022).

AI language model assistance

ChatGPT (developed by OpenAI) was used to refine and improve the clarity and grammar of the text. The authors reviewed and revised all outputs to ensure accuracy and alignment with the intended message.

ETHICS DECLARATIONS

Ethics approval and consent to participate

Not applicable.

649 **Consent for publication**

650 Not applicable.

651 **Competing interests**

652 No competing interests.

REFERENCES

1. Duplessis S, Lorrain C, Petre B, Figueroa M, Dodds PN, Aime MC. Host Adaptation and Virulence in Heteroecious Rust Fungi. Annu Rev Phytopathol. 2021;59:403–22.

2. Aime MC, McTaggart AR. A higher-rank classification for rust fungi, with notes on genera. Fungal Syst Evol. 2021;7:21–47.

3. Carnegie AJ, Pegg GS. Lessons from the Incursion of Myrtle Rust in Australia. Annu Rev Phytopathol. 2018;56 Volume 56, 2018:457–78.

4. Carefoot, Sprott GL ER. Famine on the wind; Man's Battle Against Plant Disease. 1967.

5. Aime MC, Bell CD, Wilson AW. Deconstructing the evolutionary complexity between rust fungi (Pucciniales) and their plant hosts. Stud Mycol. 2018;89:143–52.

6. Wood AR, Aime MC. Phylogenetics of the rust fungi (Pucciniales) of South Africa, with notes on their life histories and possible origins. Mycologia. 2024;0:1–27.

7. Aime MC, McTaggart AR, Mondo SJ, Duplessis S. Chapter Seven - Phylogenetics and Phylogenomics of Rust Fungi. In: Townsend JP, Wang Z, editors. Advances in Genetics. Academic Press; 2017. p. 267–307.

8. Duplessis S, Cuomo CA, Lin Y-C, Aerts A, Tisserant E, Veneault-Fourrey C, et al. Obligate biotrophy features unraveled by the genomic analysis of rust fungi. Proc Natl Acad Sci U S A. 2011;108:9166–71.

9. Presti LL, Lanver D, Schweizer G, Tanaka S, Liang L, Tollot M, et al. Fungal Effectors and Plant Susceptibility. Annu Rev Plant Biol. 2015;66 Volume 66, 2015:513–45.

10. Guerillot P, Salamov A, Louet C, Morin E, Frey P, Grigoriev IV, et al. A Remarkable Expansion of Oligopeptide Transporter Genes in Rust Fungi (Pucciniales) Suggests a Specialization in Nutrient Acquisition for Obligate Biotrophy. Phytopathology®. 2023;113:252–64.

11. Gupta YK, Marcelino-Guimarães FC, Lorrain C, Farmer A, Haridas S, Ferreira EGC, et al. The soybean rust pathogen Phakopsora pachyrhizi displays transposable element proliferation that correlates with broad host-range adaptation on legumes. 2022;:2022.06.13.495685.

12. Lorrain C, Gonçalves Dos Santos KC, Germain H, Hecker A, Duplessis S. Advances in understanding obligate biotrophy in rust fungi. New Phytol. 2019;222:1190–206.

13. Ramos AP, Tavares S, Tavares D, Silva MDC, Loureiro J, Talhinhas P. Flow cytometry reveals that the rust fungus, romyces bidentis (Pucciniales), possesses the largest fungal genome reported—2489 Mbp. Mol Plant Pathol. 2015;16:1006–10.

14. Mohanta TK, Bae H. The diversity of fungal genome. Biol Proced Online. 2015;17:8.

15. Raffaele S, Kamoun S. Genome evolution in filamentous plant pathogens: why bigger can be better. Nat Rev Microbiol. 2012;10:417–30.

16. Henningsen EC, Hewitt T, Dugyala S, Nazareno ES, Gilbert E, Li F, et al. A chromosome-level, fully phased genome assembly of the oat crown rust fungus Puccinia

coronata f. sp. avenae: a resource to enable comparative genomics in the cereal rusts. G3
GenesGenomesGenetics. 2022;12:jkac149.

17. Tobias PA, Schwessinger B, Deng CH, Wu C, Dong C, Sperschneider J, et al.
Austropuccinia psidii, causing myrtle rust, has a gigabase-sized genome shaped by
transposable elements. G3 GenesGenomesGenetics. 2021;11:jkaa015.

18. Liang J, Li Y, Dodds PN, Figueroa M, Sperschneider J, Han S, et al. Haplotype-phased
and chromosome-level genome assembly of Puccinia polysora, a giga-scale fungal
pathogen causing southern corn rust. Mol Ecol Resour. n/a n/a.

19. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified
classification system for eukaryotic transposable elements. Nat Rev Genet. 2007;8:973–82.

20. Dubin MJ, Mittelsten Scheid O, Becker C. Transposons: a blessing curse. Curr Opin
Plant Biol. 2018;42:23–9.

21. Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D. Stress-Driven
Transposable Element De-repression Dynamics and Virulence Evolution in a Fungal
Pathogen. Mol Biol Evol. 2020;37:221–39.

22. Oggenfuss U, Croll D. Recent transposable element bursts are associated with the
proximity to genes in a fungal plant pathogen. PLOS Pathog. 2023;19:e1011130.

23. Amselem J, Vigouroux M, Oberhaensli S, Brown JKM, Bindschedler LV, Skamnioti P, et
al. Evolution of the EKA family of powdery mildew avirulence-effector genes from the ORF 1
of a LINE retrotransposon. BMC Genomics. 2015;16:917.

24. Fouché S, Oggenfuss U, Chanclud E, Croll D. A devil's bargain with transposable
elements in plant pathogens. Trends Genet. 2022;38:222–30.

25. Gladyshev E. Repeat-Induced Point Mutation and Other Genome Defense Mechanisms
in Fungi. Microbiol Spectr. 2017;5.

26. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten
things you should know about transposable elements. Genome Biol. 2018;19:199.

27. Bewick AJ, Hofmeister BT, Powers RA, Mondo SJ, Grigoriev IV, James TY, et al.
Diversity of cytosine methylation across the fungal tree of life. Nat Ecol Evol. 2019;3:479–90.

28. van Wyk S, Wingfield BD, De Vos L, van der Merwe NA, Steenkamp ET. Genome-Wide
Analyses of Repeat-Induced Point Mutations in the Ascomycota. Front Microbiol.
2021;11:622368.

29. Duhamel M, Hood ME, Rodríguez de la Vega RC, Giraud T. Dynamics of transposable
element accumulation in the non-recombining regions of mating-type chromosomes in
anther-smut fungi. Nat Commun. 2023;14:5692.

30. Lorrain C, Oggenfuss U, Croll D, Duplessis S, Stukenbrock E. Transposable Elements in
Fungi: Coevolution With the Host Genome Shapes, Genome Architecture, Plasticity and
Adaptation. In: Zaragoza Ó, Casadevall A, editors. Encyclopedia of Mycology. Oxford:
Elsevier; 2021. p. 142–55.

31. Nicolás FE, Garre V. RNA Interference in Fungi: Retention and Loss. Microbiol Spectr.
2016;4:10.1128/microbiolspec.funk-0008–2016.

32. Muszewska A, Steczkiewicz K, Stepniewska-Dziubinska M, Ginalski K. Transposable
elements contribute to fungal genes and impact fungal lifestyle. Sci Rep. 2019;9:4307.

33. Orozco-Arias S, Sierra P, Durbin R, González J. MCHelper automatically curates
transposable element libraries across eukaryotic species. 2024;:2023.10.17.562682.

34. Oliveira JIN, Lane C, Mugambi K, Yildirir G, Nicol AM, Kokkoris V, et al. Analyses of
transposable elements in arbuscular mycorrhizal fungi support evolutionary parallels with
filamentous plant pathogens. 2024;:2024.11.04.621924.

35. Wicker T, Gundlach H, Spannagl M, Uauy C, Borrill P, Ramírez-González RH, et al.
Impact of transposable elements on genome structure and evolution in bread wheat.
Genome Biol. 2018;19:103.

36. Mirouze M, Vitte C. Transposable elements, a treasure trove to decipher epigenetic
variation: insights from Arabidopsis and crop epigenomes. J Exp Bot. 2014;65:2801–12.

37. Frantzeskakis L, Kracher B, Kusch S, Yoshikawa-Maekawa M, Bauer S, Pedersen C, et
al. Signatures of host specialization and a recent transposable element burst in the dynamic
one-speed genome of the fungal barley powdery mildew pathogen. BMC Genomics.
2018;19:381.

38. Stajich JE, Lovett B, Lee E, Macias AM, Hajek AE, de Bivort BL, et al. Signatures of
transposon-mediated genome inflation, host specialization, and photoentrainment in
Entomophthora muscae and allied entomophthoralean fungi. eLife. 12:RP92863.

39. Wicker T, Oberhaensli S, Parlange F, Buchmann JP, Shatalina M, Roffler S, et al. The
wheat powdery mildew genome shows the unique evolution of an obligate biotroph. Nat
Genet. 2013;45:1092–6.

40. Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, et al. Large-scale
genome sequencing of mycorrhizal fungi provides insights into the early evolution of
symbiotic traits. Nat Commun. 2020;11:5125.

41. van Wyk S, Harrison CH, Wingfield BD, De Vos L, van der Merwe NA, Steenkamp ET.
The RIPper, a web-based tool for genome-wide quantification of Repeat-Induced Point (RIP)
mutations. PeerJ. 2019;7:e7447.

42. Silvestri A, Turina M, Fiorilli V, Miozzi L, Venice F, Bonfante P, et al. Different Genetic
Sources Contribute to the Small RNA Population in the Arbuscular Mycorrhizal Fungus
Gigaspora margarita. Front Microbiol. 2020;11.

43. Möller M, Habig M, Lorrain C, Feurtey A, Haueisen J, Fagundes WC, et al. Recent loss
of the Dim2 DNA methyltransferase decreases mutation rate in repeats and changes
evolutionary trajectory in a fungal pathogen. PLOS Genet. 2021;17:e1009448.

44. Oliveira JIN, Corradi N. Strain-specific evolution and host-specific regulation of
transposable elements in the model plant symbiont Rhizophagus irregularis. G3
GenesGenomesGenetics. 2024;14:jkae055.

45. Dallaire A, Manley BF, Wilkens M, Bista I, Quan C, Evangelisti E, et al. Transcriptional
activity and epigenetic regulation of transposable elements in the symbiotic fungus
Rhizophagus irregularis. Genome Res. 2021;31:2290–302.

46. Baril T, Croll D. A pangenome-guided manually curated library of transposable elements
for Zymoseptoria tritici. BMC Res Notes. 2023;16:335.

47. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012;28:3150–2.

48. Lorrain C, Oggenfuss U, Croll D, Duplessis S, Stukenbrock E. Transposable Elements in
Fungi: Coevolution With the Host Genome Shapes, Genome Architecture, Plasticity and
Adaptation. In: Zaragoza Ó, Casadevall A, editors. Encyclopedia of Mycology. Oxford:
Elsevier; 2021. p. 142–55.

49. Baril T, Galbraith J, Hayward A. Earl Grey: A Fully Automated User-Friendly
Transposable Element Annotation and Analysis Pipeline. Mol Biol Evol. 2024;41:msae068.

50. Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open
Software Suite. Trends Genet TIG. 2000;16:276–7.

51. Kimura M. A simple method for estimating evolutionary rates of base substitutions
through comparative studies of nucleotide sequences. J Mol Evol. 1980;16:111–20.

52. Ellinghaus D, Kurtz S, Willhoeft U. LTRharvest, an efficient and flexible software for de
novo detection of LTR retrotransposons. BMC Bioinformatics. 2008;9:18.

53. Rozewicki J, Li S, Amada KM, Standley DM, Katoh K. MAFFT-DASH: integrated protein
sequence and structural alignment. Nucleic Acids Res. 2019;47:W5–10.

54. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019;20:238.

55. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics. 2014;30:1312–3.

56. De Bie T, Cristianini N, Demuth JP, Hahn MW. CAFE: a computational tool for the study
of gene family evolution. Bioinformatics. 2006;22:1269–71.

57. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. eggNOG-
mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the
Metagenomic Scale. Mol Biol Evol. 2021;38:5825–9.

58. Feurtey A, Lorrain C, McDonald MC, Milgate A, Solomon PS, Warren R, et al. A
thousand-genome panel retraces the global spread and adaptation of a major fungal crop
pathogen. Nat Commun. 2023;14:1059.

59. van Wyk S, Harrison CH, Wingfield BD, De Vos L, van der Merwe NA, Steenkamp ET.
The RIPper, a web-based tool for genome-wide quantification of Repeat-Induced Point (RIP)
mutations. PeerJ. 2019;7:e7447.

60. Mondo SJ, Dannebaum RO, Kuo RC, Louie KB, Bewick AJ, LaButti K, et al. Widespread
adenine N6-methylation of active genes in fungi. Nat Genet. 2017;49:964–8.

61. Crowell CR, Wilkerson DG, Beckauri M, Cala AR, McMullen PW, Mondo S, et al. The
Melampsora americana Population on Salix purpurea in the Great Lakes Region Is Highly
Diverse with a Contributory Influence of Clonality. Phytopathology. 2022;112:907–16.

62. Persoons A, Maupetit A, Louet C, Andrieux A, Lipzen A, Barry KW, et al. Genomic
Signatures of a Major Adaptive Event in the Pathogenic Fungus Melampsora larici-populina.
Genome Biol Evol. 2022;14:evab279.

63. Branco S, Badouin H, Rodríguez de la Vega RC, Gouzy J, Carpentier F, Aguileta G, et
al. Evolutionary strata on young mating-type chromosomes despite the lack of sexual
antagonism. Proc Natl Acad Sci U S A. 2017;114:7067–72.

64. Perlin MH, Amselem J, Fontanillas E, Toh SS, Chen Z, Goldberg J, et al. Sex and
parasites: genomic and transcriptomic analysis of Microbotryum lychnidis-dioicae, the
biotrophic and plant-castrating anther smut fungus. BMC Genomics. 2015;16:461.

65. Gupta YK, Marcelino-Guimarães FC, Lorrain C, Farmer A, Haridas S, Ferreira EGC, et
al. Major proliferation of transposable elements shaped the genome of the soybean rust
pathogen Phakopsora pachyrhizi. Nat Commun. 2023;14:1835.

66. Li F, Upadhyaya NM, Sperschneider J, Matny O, Nguyen-Phuc H, Mago R, et al.
Emergence of the Ug99 lineage of the wheat stem rust pathogen through somatic
hybridisation. Nat Commun. 2019;10:5068.

67. Wang J, Xu Y, Peng Y, Wang Y, Kang Z, Zhao J. A fully haplotype-resolved and nearly
gap-free genome assembly of wheat stripe rust fungus. Sci Data. 2024;11:508.

68. Holden S, Bakkeren G, Hubensky J, Bamrah R, Abbasi M, Qutob D, et al. Uncovering
the history of recombination and population structure in western Canadian stripe rust
populations through mating type alleles. BMC Biol. 2023;21:233.

69. Coradetti ST, Pinel D, Geiselman GM, Ito M, Mondo SJ, Reilly MC, et al. Functional
genomics of lipid metabolism in the oleaginous yeast Rhodosporidium toruloides. eLife.
2018;7:e32110.

70. Sperschneider J, Chen J, Anderson C, Morin E, Zhang X, Lewis D, Henningsen E,
Grigoriev IV, Rathjen J, Jones DA, Duplessis S, Dodds PN. A chromosome-scale genome
assembly of the flax rust fungus reveals the two unusually large effector proteins, AvrM3 and
AvrN. submitted.

Fig. 1: **The phylogenetic tree and transposable element content in the**
Pucciniomycotina. A) phylogenetic tree of the 12 selected Pucciniomycotina species based
on a single-copy orthogroup from Orthofinder v2.4.0 and visualized using iTOL
(https://itol.embl.de/). The Pucciniales are represented in yellow. The category *Other*
Pucciniomycotina (blue) refers to the four species of Pucciniomycotina outside the order
Pucciniales. B) The bar plot displays the TE content across the analyzed genomes. Dark
blue bars represent the proportion of the genome covered by TEs, while light blue bars show
the total genome size (in Mb). Percentages on the right indicate the TE coverage relative to
the total genome size. Species are described as follow: Melap (*Melampsora allii-populina*),
Melli (*Melampsora lini*), Mellp (*Melampsora larici-populina*), Melam (*Melampsora*
americana), Phapa (*Phakopsora pachyrhizi*), Pucst (*Puccinia striiformis* f. sp. *tritici*), Puctr
(*Puccinia triticina*), PucgtrUg99 (*Puccinia graminis* f. sp. *tritici* Ug99), Pucgtr201 (*Puccinia*
graminis f. sp. *tritici* 20-1), Micld (*Microbotryum lychnidis-dioicae*), Micin (*Microbotryum*
intermedium), Leucr (*Leucosporidiella creatinivora*), Rhoto (*Rhodosporidium toruloides*).

Fig. 2: **Composition of transposable elements in Pucciniomycotina genomes.** The
stacked bar plots represent the percentage of transposable elements (TEs) annotated at the
A) Class level, B) Class I order level, and C) Class II order level for each Pucciniomycotina
genome. The n values indicate the number of distinct TE families identified in each genome.
The percentage of TE genome coverage at the super-family (SF) level for E) Class I
elements (purple) and F) Class II elements (blue).

Fig. 3: **Transposable element insertion ages.** A-R Kimura distances across the TE
landscapes of Pucciniomycotina genomes. Each distribution shows the proportion of the
genome (y-axis) composed of copies exhibiting varying degrees of divergence (x-axis).
Arrows indicate TE mobilization events. S-T) Number of LTR-Gypsy elements accumulated

through time (million years ago - Mya), based on insertion age for the S) Melampsoraceae

species and T) the Pucciniaceae species.

Fig. S1: Association between genome size and gene/TE content. Genome sizes (Mb) were

plotted against genes or TE coverage for each Pucciniomycotina species. The lines

represent the linear regression.

Fig. S2: **Improvement of TE annotations through manual curation**. The proportion of

annotated TE families after REPET detection and annotation (RawR), after the first round of

curation (Postpct), after the second round of curation with MCHelper (postMC), after a third

round of curation with Pucciniomycotina-guided clustering (PostCDhit).

Fig. S3: **Shared and specific TE families between the Pucciniomycotina species.**

Numbers of TE families shared among the 14 genomes based on CDHIT clustering. The

bars on the left represent the total number of genes per species. The clusters shared among

Pucciniaceae are colored in blue and the clusters shared among Melampsoraceae are

colored in red.

Fig. S4: Number of LTR-Gypsy elements accumulated through time (million years ago -

Mya), based on insertion age for each genome.

Fig. S5: **TE controls involved in methylations are detected in all species**. The number of

genes involved in TE control corresponds to DNMT1 and DNMT5.

Fig. S6: **TE controls involved in RNA-induced silencing complex are detected for all**

genomes. The number of genes involved in TE control corresponds to the RNAi pathway:

Dicer-like (DCL; DCR), RNA-dependent RNA polymerase (RDP; RNT; RPAB; RPB),

Argonaute (AGO; ARB) for each genome.

Fig. S7: **Contraction and Expansion of multigene families in Pucciniomycotina.**

The phylogenetic tree of the selected Pucciniomycotina species based on single-copy

orthogroups from Orthofinder v2.4.0 and infer with Raxml (500 bootstraps) and visualized

using iTOL, rooted with *Ustilago maydis.* Numbers in purple correspond to the number of

orthogroups in contractions at each branch or node and yellow at expansions.

Fig. S8: **Contraction of Multigene Families in Pucciniales.** Functional annotation of

multigene families in contraction in Pucciniales. Genes are re-annotated by EggNOG-

mapper v2.1.12. A threshold is set to kept only species with at least 20 proteins per COG

category. COG categories correspond to: "-": no hit; B: chromatin structure and dynamics; C:

energy production and conversion;　E: amino acid transport and metabolism; G:

carbohydrate transport and metabolism; I: lipid transport and metabolism; K: transcription;

O: posttranslational modification, protein turnover, chaperones, P: inorganic ion transport

and metabolism; Q: secondary metabolites biosynthesis, transport and catabolism; S:

function unknown; T: signal transduction mechanisms.

Fig. S9: **Significant contraction and expansion of multigene families in**

Pucciniomycotina. A) The phylogenetic tree of the selected Pucciniomycotina species

based on a single-copy orthogroup from Orthofinder v2.4.0 and visualized using iTOL,

rooted with *Ustilago maydis.* Numbers in black correspond to the number of significant

orthogroups in expansion/contraction at each branch or node. B) COG category annotation

of genes involved in expansion and contraction between Pucciniales and other

Pucciniomycotina. Contraction and expansion are detected with CAFE v5.0 and orthogroups

with a *p*-value (0.05) are kept. Genes are re-annotated by EggNOG-mapper v2.1.12. The

number of genes per species and categories. COG categories functions correspond to : A:

RNA processing and modification; C: energy production and conversion;　D: cell cycle

control, cell division chromosome partitioning; G: carbohydrate transport and metabolism; I:

lipid transport and metabolism; K: transcription; L: replication recombination and repair; M:

cell wall, membrane, envelope biogenesis; N: cell motility; O: posttranslational modification,

protein turnover, chaperones, Q: secondary metabolites biosynthesis, transport and

catabolism; S: function unknown; T: signal transduction mechanisms.

Table S1: **General information of selected genomes** [60–70].

Table S2: **TE annotations at the Class level for the Pucciniomycotina genomes**.

ClassAnn represents the different TE Classes (I, II, Unclassified); nbTE represents the

number of TE families in each Class; ncovP represents the percentage of genome coverage

of each TE class in the species genome; nbcopie represents the number of copies

associated with the TE family for each Class; Sumcov represents the percentage of

coverage by all the TEs per genome.

Table S3: **TE annotations at the Order level for the Pucciniomycotina genomes**.

OrderAnn represents the different TE Orders; nbTE represents the number of TE families in

each Order; ncovP represents the percentage of genome coverage of each TE Order in the

species genome; nbcopie represents the number of copies associated with the TE family for

each Order; Sumcov represents the percentage of coverage by all the TEs per genome.

Table S4: **TE annotations at the Super-family level for the Pucciniomycotina genomes**.

OrderAnn represents the different TE SF; nbTE represents the number of TE families in

each SF; ncovP represents the percentage of genome coverage of each TE SF in the

species genome; nbcopie represents the number of copies associated with the TE family for

each SF; Sommecov represents the percentage of coverage by all the TEs per species.

Table S5: **Average of the TE sequence identity at the Class level**. The meanID

represents the mean sequence identity (%) of individual copies to their respective TE family

consensus sequence averaged per TE class (ClassAnn).

Table S6: **Average of the TE sequence identity at the Order level**. The meanID

represents the mean sequence identity (%) of individual copies to their respective TE family

consensus sequence averaged per TE Order (OrderAnn).

Table S7: **Shared TEs within the three Melampsoraceae genomes**.

Table S8: **Shared TEs within the five Pucciniaceae genomes**.

Table S9: **RIP signature analysis**. Genome-wide RIP signature detection using TheRIPper

[59]. RIP-like signatures were searched by estimating the RIP composite index in TE families

consensus sequences for each Pucciniomycotina genome described in Feurtey et al.

(2023)[58].

Table S10: **Orthogroups summary metrics of the selected Pucciniomycotina genomes.**

Genomes are described as follow: Melap (*Melampsora allii-populina*), Melli (*Melampsora*

lini), Mellp (*Melampsora larici-populina*), Melam (*Melampsora americana*), Phapa

(*Phakopsora pachyrhizi*), Pucst (*Puccinia striiformis* f. sp. *tritici*), Puctr (*Puccinia triticina*),

PucgtrUg99 (*Puccinia graminis* f. sp. *tritici* Ug99), Micld (*Microbotryum lychnidis-dioicae*),

Micin (*Microbotryum intermedium*), Leucr (*Leucosporidiella creatinivora*), Rhoto

(*Rhodosporidium toruloides*).

Table S11: **Orthogroups in contraction of the selected Pucciniomycotina genomes.**

A

B

Melap	74.8 %
MelliH	83.99 %
MelliC	83.38 %
Mellp	46.98 %
Melam	55.73 %
Phapa	92.35 %
PucstB	49.54 %
PucstA	49.27 %
PuctrB	77.43 %
PuctrA	77.08 %
Pucgr201B	51.69 %
Pucgr201A	51.06 %
PucgrUg99A	54.35 %
PucgrUg99C	51.44 %
Micld	16.18 %
Micin	10.56 %
Leucr	6.13 %
Rhoto	2.3 %

Categories
non-TE
TEs

Genome size (Mb)

A Percent coverage of TE family in Class

B Percent coverage of TE family in Class I

C Percent coverage of TE family in Class II

D LTR/TIR super-family coverage (Mb)

E