Consistent covariation of dispersal, life history and thermal niches across terrestrial arthropods

Garben Logghe[1,2] (ORCID: 0000-0003-2000-7982)

Femke Batsleer[1] (ORCID: 0000-0001-5893-5966)

Guillermo Fandos[3] (ORCID: 0000-0003-1579-9444)

Dirk Maes[2,4] (ORCID: 0000-0002-7947-3788)

Dries Bonte[1] (ORCID: 0000-0002-3320-7505)*

1 Ghent University, Department of Biology, Karel Lodewijk Ledeganckstraat 35, 9000 Ghent, Belgium

2 Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 Bus 73, 1000 Brussels, Belgium

3 Department of Biodiversity, Ecology and Evolution, Facultad de Ciencias Biológicas, Complutense University of Madrid

4 Radboud Institute for Biological and Ecological Sciences (RIBES), Radboud University, PO Box 9010, Nijmegen NL-6500 GL, The Netherlands

Corresponding author: D. Bonte, Ghent University, Department of Biology, K. L. Ledeganckstraat 35, 9000 Ghent, Belgium. E-mail: dries.bonte@ugent.be

Key words: Arthropoda, syndromes, fast-slow continuum, dispersal, life-history, climate change

Type of article: Letter

Number of words
Abstract: 149

 Main text (excluding abstract, acknowledgements, references, table and figure legends): 4043

Number of references: 86

Number of figures: 4

Number of Tables: 0

Statement of Authorship

Garben Logghe, Femke Batsleer, Dirk Maes and Dries Bonte conceived the framework of the study. Garben Logghe collected the trait data. Guillermo Fandos and Garben Logghe developed the statistical analysis. Garben Logghe wrote the first draft of the manuscript. All authors contributed substantially to interpretation of the results and revision of the manuscript.

Data accessibility statement

Abstract

Arthropods, as ectotherms, are experiencing global declines, with many species facing the need to

either acclimate or disperse in response to climate change. Understanding to which degree life history,

dispersal and thermal niche traits covary is key to improve distribution forecasting under climate

change. We quantified life history, dispersal and thermal range covariation among 4000 Western

European arthropod species spanning eight orders, considering phylogenetic relationships to account

for common ancestry. We demonstrate the existence of two independent axes of life history variation:

the fast-slow continuum and the reproductive strategy axis. Species at the fast end of the continuum

have higher dispersal capacities and broader thermal niches than slower species. The resulting

syndromes were surprisingly consistent across orders. These trait combinations, which generally

enhance range-shifting potential, point to the emergence of two distinct groups of arthropods: those

well-suited and those less equipped to mitigate the effects of future climate change.

Introduction

Climate change is an escalating global problem, threatening numerous species across various taxonomic groups (Bellard *et al.* 2012; Habibullah *et al.* 2022; Maxwell *et al.* 2019). Species may cope with changing climatic conditions by acclimatisation and/or adaptation to the experienced novel conditions, or by tracking their climate window through dispersal (MacLean & Beissinger 2017). The pace at which species expand or shift their distribution will critically depend on their life histories—such as survival, reproduction and development—that determine population growth rate and overall demography. Furthermore, this rate of range shifts will be shaped by their dispersal capacity and ability to acclimate to novel conditions (MacLean & Beissinger 2017). Forecasting biodiversity under climate change (Settele *et al.* 2014; Urban 2019), therefore, relies on our ability to incorporate accurate trait data into extinction risk projections (La Sorte & Jetz 2010; Thomas *et al.* 2004), thereby informing vulnerability assessments (Foden *et al.* 2013; Foden & Young 2016) and refining predictions of emerging community assemblages (Stralberg *et al.* 2009; Williams & Jackson 2007). For arthropods, such an approach seems to be most urgent since any detailed species-specific assessment will fail to provide the needed insights in light of their steep global decline (Wagner 2020), which potentially jeopardizes the crucial ecosystem services these organisms provide (Yang & Gratton 2014).

Species are expected to show intrinsic life history trade-offs, positioning them along a "slow-fast" continuum (Beukhof *et al.* 2019; Gaillard *et al.* 1989; Promislow & Harvey 1990). Slow-living species have longer development times and lifespans, prioritizing resource allocation toward offspring survival rather than fecundity, while fast-living species adopt the opposite strategy (Reynolds 2003). In theory, slow-living species are better adapted to stable environments and have, therefore, greater difficulty recovering from stochastic mortality events (Pianka 1970). Due to similar evolutionary pressures stemming from environmental instability, dispersal rates are expected to be lower in species with slow life histories (Beckman *et al.* 2018; Stevens *et al.* 2013). Body size, however, may have contrasting effects. While smaller body sizes are typically linked to a fast-paced lifestyle, larger body sizes are often

positively correlated with greater dispersal ability (Hillaert *et al.* 2018; Jenkins *et al.* 2007; MacLean &

Beissinger 2017; Sekar 2012). Predicted relationships between species' dispersal and life history traits

are therefore highly variable, both among and within species (Bonte 2025).

Both dispersing towards new areas and staying within a changing environment comes with certain

costs, such as the risk of staying or settling in environments where conditions may be unfavourable

(Bonte *et al.* 2012). Such limitations should be especially pronounced in ecological specialists, as they

will suffer most from local environmental changes and/or experience greater challenges in establishing

populations in unfamiliar habitats (Gilman *et al.* 2010; Hill *et al.* 2001; Pöyry *et al.* 2009; Warren *et al.*

2001). In the context of climate change, thermal tolerance is a particularly important niche component

in shaping distribution patterns. A species' thermal niche encompasses several critical factors,

including its thermal minimum (the lowest temperature it can endure), thermal maximum (the highest

temperature it can tolerate) and thermal range (the span between these extremes) (Gvoždík 2018).

Species with narrow thermal niche breadths – those with limited tolerance to temperature variation –

are generally expected to be more vulnerable to climate change (Grinder & Wiens 2023; Saupe *et al.*

2015), facing greater risks during dispersal as they attempt to track their optimal temperature

conditions. Given this pivotal role of a species' thermal niche in determining both distribution ranges

and dispersal costs, it can be expected to have shaped the evolution of both life histories (Carbonell &

Stoks 2020; Hampe & Jump 2011) and dispersal (Hillaert *et al.* 2015; Travis *et al.* 2012).

Traits underlying life history variation can show complex covariation and are integrated into so-called

syndromes (Dammhahn *et al.* 2018; Polverino *et al.* 2018; Wright *et al.* 2019). Similarly, dispersal

syndromes notify covariations between dispersal and other traits, and have been documented across

various taxonomic groups (Beckman *et al.* 2018; Buoro & Carlson 2014; Ronce & Clobert 2012; Stevens

et al. 2013, 2014). Understanding the organisation of trait syndromes not only allows us to estimate

missing trait data (Stevens *et al.* 2013; Whitmee & Orme 2013) in understudied taxa, but also allows

more accurate forecasting of species distributions under environmental change (Weil *et al.* 2023).

Given the urgency to understand and predict arthropod distributions under future climate change, we

aim to identify covariations between dispersal, life history and thermal niche widths across a wide

range of Western European species. If dispersal and thermal range are independent of life history

traits, they should form separate axes of variation alongside the fast-slow continuum. Conversely, if

dispersal and thermal range covary with life history traits, their inclusion will allow further application

of the current fast-slow continuum into the context of global change. Such trait relationships may

have important implications for species' vulnerability to climate warning. For example, a positive

correlation between thermal niche breadth and dispersal ability would suggest interspecific

differences in vulnerability, as species with high dispersal capacity and broad thermal tolerance are

likely to be more resilient. In contrast, a negative correlation between these traits could indicate trade-

offs that limit persistence – enabling species to respond through either range shifts or thermal

tolerance, but not both. Similar advantages or trade-offs may also emerge from covariation with life

history strategies. Given the vast diversity in arthropod morphology, lifestyles and dispersal modes,

we expect different taxa to exhibit significant variation in both the direction and strength of these trait

covariations. Additionally, we explore whether these syndromes are phylogenetically conserved, as

shared evolutionary history may drive similar trait associations in closely related species, eventually

shaping their responses to environmental changes and range shifts.

Material and Methods

Data description

The data used for this study was sourced from the dataset described by Logghe et al. (2025), which contains information on 28 life history and ecological traits for 4874 terrestrial arthropod species. We focused on eight arthropod orders for which data on both dispersal and life history was available: Araneae, Coleoptera, Hemiptera, Hymenoptera, Isopoda, Lepidoptera, Odonata and Orthoptera. Two orders, Diptera and Opiliones, were excluded due to lacking data. This resulted in a dataset containing 4343 species. From this dataset, we selected nine traits related to demography (life history), dispersal (movement) and physiology (thermal niche), relevant for predicting range shifts under climate change (Estrada *et al.* 2016; Urban *et al.* 2016). Appendix S.1 shows the number of data points for every trait within each order.

We selected body size as a central trait, since it is shown to correlate with both life history and thermal limits (e.g. Berger et al., 2008; Hillaert et al., 2018; Klockmann et al., 2017). In this study, we use the adult body length (measured in millimetres) as a proxy for body size. In contrast to several other highly correlated , relevant measures of body size (i.e., body mass, volume; Ganihar 1997; Martin *et al.* 2014), body length is always reported in the primary literature. To avoid introducing additional and unquantifiable uncertainty in our analyses, we retained this metric rather than transforming it to one of the derived metrics that require assumptions about body shape that are not consistently documented, even within taxonomic orders.

As demographic traits, we selected development time, fecundity and voltinism. Development time is defined as the duration between egg laying and adult emergence, expressed in days. Fecundity is defined as the number of offspring a female could produce during her lifetime, while voltinism refers to the maximum number of generations a species could complete in a single year. Note that the maximum value for voltinism is used, as the number of generations can vary annually and across regions. All these trait values were obtained from primary literature (see Logghe *et al.* 2025).

Next, we derived metrics of dispersal, defined here as the movement of an organism to a location of

reproduction (Clobert *et al.* 2009). In this study, we do not consider dispersal capacity as a specific

distance range as the latter is most often biased by the spatiotemporal extent of the study. Instead,

we translated various recorded dispersal proxies – often highly variable across and even within orders

– to an individual trait representing a species' general motivation to disperse away from its current

location (Bonte & Dahirel 2017). We thus rely on the assumption that dispersal capacity – reflected in

traits such as movement speed, mean dispersal distance, wing load or wing development – is positively

associated with dispersal propensity. This relationship has been experimentally tested in only a few

groups (e.g. Araneae, Diptera), and is for other taxa inferred from expert judgement (see the

supplement of Logghe *et al.* 2025 for a full list of sources). We consider this assumption reasonable, as

typical (inverse power-family) dispersal kernels show a statistical correlation between dispersal

propensity (i.e. frequency of dispersal distances beyond the individual's social neighbourhood) and

mean dispersal distance (Bitume *et al.* 2013; Clobert *et al.* 2009; Cousens *et al.* 2008). Therefore, we

standardized values from the literature onto a standardized scale ranging from 0.1 to 1 (Logghe *et al.*,

2025). On this scale, a value of 0.1 represents a largely philopatric species that rarely disperses, while

a value of 1 corresponds to a highly mobile and dispersive species. Importantly, these scores are always

relative within the same taxonomic order, meaning that for instance a butterfly with score 1 does not

necessarily show the same dispersal capacity (in terms of absolute distance covered) as a spider with

the same score.

Finally, we included measures of each species' thermal niche, specifically thermal mean, minimum,

maximum and range. These metrics were derived by overlaying GBIF species distribution data with a

raster of climate records, allowing us to assess the climate conditions under which species currently

occur. Using zonal statistics for raster values within defined geographic zones, we calculated the

average, lowest and highest annual temperatures (in °C) experienced across each species' geographic

range (see Logghe *et al.* 2025 for technical details). As distribution data were obtained directly from

GBIF, our estimates represent the realised thermal niche, defined here as the range of temperature

conditions a species experiences within its known distribution. While using GBIF data may introduce

observer bias, we expect this bias to be reduced in our study, as we focus on well-documented species

that are more likely to have reliable and comprehensive trait data available. This approach does not

necessarily capture the full physiological (or fundamental) thermal limits of a species. Instead, it

reflects the environmental temperatures species currently encounter – conditions that may be shaped

by factors such as dispersal constraints or biotic interactions. However, we intentionally excluded

species that are endemic to certain geographic distinct locations. Ranges of species strictly engaged in

host-specific interactions may be primarily constrained by the distribution of the host rather than by

its ow thermal niche. We did not include such highly specialised species, and the few considered are

documented to shift hosts within their geographic range (e.g., more specialistic herbivores or

parasitoids). We do acknowledge that experimentally measured physiological limits and optima would

provide a more direct estimate of thermal tolerance, but such data are not available for the included

broad range of arthropod species.

It is important to note that this study does not account for intraspecific variation in trait values. The

literature mostly reports mean values, occasionally accompanied by minimum and maximum ranges,

but rarely with proper variability statistics. We thus developed a species-level meta-analysis, using

mean trait values to explore broad patterns of trait covariation across the phylogenetic tree.

Statistical analysis

To account for the influence of shared evolutionary history on trait similarities, we controlled for

phylogenetic relatedness in all analyses. This approach ensured that observed relationships between

traits reflected ecological or evolutionary patterns, rather than being influenced by common ancestry.

Phylogenetic correction was essential to meet the assumption of statistical independence, as trait

values are often not independent due to phylogenetic relatedness. To achieve this, we constructed a

phylogenetic tree including all species in the dataset, which served as the basis for subsequent

analyses. The tree was built by linking species names to the Open Tree Taxonomy database, which

contains data on known phylogenetic relationships ("rotl" package) (Michonneau *et al.* 2016). Branch

lengths were added to the tree using the "ape" package (Paradis & Schliep 2019) and calculated with

Grafen's method (Grafen 1989), which transforms a cladogram into an ultrametric tree by scaling

internal node heights according to the number of descendant taxa. This approach assumes a

speciational model of divergence, where evolutionary change is linked to speciation events rather than

elapsed time. The resulting tree is appropriate for use with comparative methods that assume

Brownian motion trait evolution (i.e. traits evolve via a random walk with constant variance over time).

To examine how life history and thermal traits are distributed across arthropod orders, we performed

a phylogenetic principal component analysis (PCA) using the "phyl.pca" function from the "phytools"

package (Revell 2012). Unlike standard PCA, this method incorporates phylogenetic relationships,

correcting for the lack of independence between species due to common ancestry. The analysis was

conducted on a subset of the dataset that included only the 200 species for which complete data on

all traits were available.

Bayesian Generalized Linear models were employed to assess whether covariations were present

between the selected traits ("brms" package) (Bürkner 2021). Given the strong phylogenetic signal

across the tree of life, the variation in data quantity and quality among orders, and our aim to detect

and understand potential divergence in the covariations across arthropod orders, we analysed every

order separately. Trait data on body size, fecundity and development time underwent log

transformation before analysis to reduce the skewness of the data. Each trait combination was run

twice for every order, with the dependent and independent variables interchanged. Given the strong

phylogenetic signals among orders, we chose not to construct a single multivariate model, also to avoid

potential biases arising from uneven patterns of missing data among taxa. Additionally, separate

models allowed us to account for different error structures, depending on the nature of each trait.

Each model was, therefore, fitted with a distribution appropriate to its independent variable: zero-

one-inflated beta for voltinism, binomial for dispersal and Gaussian for the remaining variables. The

models were run using 4 chains and 4000 iterations, with a burn-in period of 1000. Non-informative

flat priors were used for the model parameters. Phylogenetic distances between species were derived

from a phylogenetic distance matrix derived from the previously mentioned phylogenetic tree

("geiger" package) (Matthew *et al.* 2014). These distances were incorporated into every Bayesian

model as blocking factors, to ensure that phylogenetic relationships were considered when the

covariation between traits is estimated. Convergence of the models was checked by examining the

effective sample size (>1000) for each chain as well as Gelman-Rubin statistic (Rhat < 1.1) among

chains. Effect sizes were deemed clearly determined when 0 was not included in the 95% credibility

intervals of the posteriors.

Finally, we assessed the phylogenetic signal using Pagel's Lambda for each considered trait and

arthropod order (Pagel 1999). This is a metric that estimates how much closely related species

resemble each other compared to expectations under a Brownian motion model of evolution (Pearse

et al. 2025). Pagel's Lambda typically ranges from 0 (indicating no influence of phylogenetic

relationships on trait values) to 1 (indicating trait values are entirely explained by phylogenetic

relationships). We calculated this metric using the "phylosig" function from the "phytools" package

(Revell 2012).

R version 4.0.4 (R Core Team 2021) was used to perform all data analyses.

Results

Data on dispersal capacity, life history traits and thermal niche were obtained from an existing dataset described by Logghe et al. (2025). From this dataset, we selected eight arthropod orders with sufficiently high data quality, resulting in a refined dataset comprising over 4000 species and nine functional traits. As a first exploration of these data, we conducted a phylogenetically corrected PCA on a subset of species with complete data for all traits. This analysis revealed no distinct separation among orders, suggesting that covariations may be consistent across taxonomic groups (Figure 1). PC1 shows a strong positive relationship with voltinism and dispersal, while it also strongly negatively correlates with development time (Appendix S.2). Additionally, it exhibits a positive correlation with thermal mean and range, though this relationship is not as strong as with the other traits. In contrast, PC2 is strongly positively correlated with fecundity and size. These results indicate the existence of covariations between dispersal, life history, body size and thermal niche across arthropod orders.

Figure 1: phylogenetically corrected PCA results for arthropod species for which data on all traits was available (n=200). Dots represent species with symbol colour and shape according to their taxonomic order. Arrows represent the direction and magnitude of each variable's contribution to the principal components. Ellipses represent 95% confidence intervals around species scores for each arthropod order, illustrating group-level patterns in trait space.

Given the strong phylogenetic clustering of traits across the eight orders (Figure 2), we employed

Bayesian Generalized Linear models to identify trait covariations within each of these. This approach

reduces model complexity and allows us to infer potential convergence of trait covariations in an

understandable, qualitative manner. Each model accounted for phylogenetic relatedness between

species, ensuring that observed relationships reflect ecological and evolutionary patterns. Our results

reveal multiple trait syndromes within various arthropod orders (Appendix S.3). In most orders, there

is a clear positive relationship between fast development, voltinism and dispersal (Figure 3).

Additionally, temperature niche components tend to covary positively with dispersal capacity,

fecundity and size in most orders. Covariations between other trait combinations are always present

within at least one order. However, these are highly order specific and, therefore, not generalizable

across arthropod orders.

Figure 2: Overview of the distribution of trait values across the phylogenetic tree for dispersal (λ = 0.983), fecundity (λ =

0.988), thermal range (λ = 0.863) and body size (λ = 0.993). To facilitate comparison between the trees, we selected only

species with data available for all four traits (n = 297). Fecundity and body length were log-transformed to improve clarity of

the figure by creating a smoother colour gradient.

Figure 3: Overview of trait covariations across each order, represented as a correlation matrix. Each grid of the matrix is divided into eight cells, with each cell representing one order. The taxa are, from

left to right and top to bottom: Araneae, Coleoptera, Hemiptera, Hymenoptera, Isopoda, Lepidoptera, Odonata and Orthoptera. Red cells indicate a clearly determined (0 not included in the 95 credibility

interval) negative relationship, blue cells indicate a clearly determined positive relationship, and grey cells show that the relationship was not clearly determined. White cells denote that this trait

combination was not assessed for the given order, due to either an absence of data for one of the traits or convergence issues. Sample sizes for every covariation were determined by the trait with the

least available data (see Appendix S.1).

To estimate how much trait values are determined by common ancestry between species from the

same order, we calculated the phylogenetic signal for every trait and arthropod order (Figure 4). On

average, body size and development time showed very strong phylogenetic signals, suggesting these

traits are generally strongly influenced by phylogeny. In contrast, thermal mean, thermal range and

fecundity appeared largely independent of phylogeny in most arthropod orders. Finally, the

phylogenetic signal for dispersal displayed considerable variation, indicating that the influence of

phylogenetic relationships differs substantially between orders (Appendix S.4).

Figure 4: Overview of the phylogenetic signal (Pagel's Lambda) for every assessed trait for each arthropod order. Arthropod orders

are from top to bottom: Lepidoptera, Coleoptera, Hymenoptera, Hemiptera, Orthoptera, Odonata, Isopoda and Araneae. Colours

represent the strength of phylogenetic signal, with higher values (darker red) indicating that trait values are more explained by

phylogenetic relationships. The grey squares indicate that the phylogenetic signal was not calculated for that trait and order

due to a lack of data. Sample sizes for every order and trait can be found in Appendix S.1.

Discussion

Our quantitative synthesis across a wide range of Western European arthropods indicates that

dispersal, life history and thermal niche are tightly connected, implying similar evolutionary pressures

on these traits. While there are notable differences between orders, these relationships remain largely

consistent across taxa, suggesting that trait covariations can be generalized to a broad range of

arthropod species. This remarkable consistency across arthropod orders was one of the most

surprising findings of this study, given the extraordinary diversity of dispersal mechanisms and life

history strategies within the phylum. Such uniformity could have important implications for modelling

the future effects of climate change, suggesting that detailed trait information for every species might

not be as necessary as previously suggested (Urban *et al.* 2016). Instead, missing trait values could

potentially be inferred from accessible trait data, facilitating broader, more efficient predictions.

However, we would like to emphasize that these results should not be directly applied to mechanistic

modelling, as such applications require a thoughtful translation from traits to population dynamics or

ecosystem properties.

A considerable amount of interspecific life history trait variation in arthropods can be attributed to the

slow-fast continuum, with a clear distinction between species with fast (i.e. short development times)

and slow (i.e. long development times) lifecycles. Interestingly, development time appears closely tied

to dispersal capacity, as we observed clear covariations between development time, voltinism and

dispersal across various orders. Previous research has shown similar patterns, with high dispersal

ability often linked with fast life history strategies (Beckman *et al.* 2018; Stevens *et al.* 2013; Weil *et al.*

2022). Given that most insects disperse only in their winged adult stages, this dispersal-life history

syndrome may have evolved as a bet-hedging strategy in response to unstable environments, allowing

for rapid relocation when conditions become unfavourable (Snyder 2011; Southwood 1977).

Interestingly, we found limited evidence to suggest that fast-living species produce more offspring, a

pattern that would typically be anticipated within the slow-fast continuum framework (Reynolds

2003). While Lepidoptera and Coleoptera did show a clear negative relationship between fecundity

and development time, our analyses suggest that the reproductive axis of variation is as good as

independent from the developmental axis. We thus show that the independence of these two distinct

axes of life history variation is rather a rule than an exception, as was previously reported for plants

(Salguero-Gómez *et al.* 2016) and vertebrates (Gaillard *et al.* 1989; Sæther 1987; Stearns 1983).

Our results suggest that thermal range has played a role in shaping the evolution of dispersal in

arthropods, with most orders showing a positive relationship between realised thermal niche breadth

and dispersal ability. This relationship likely arises from the costs associated with dispersing to new

environments, where the risk of encountering unfavourable climatic conditions is high (Bonte *et al.*

2012). A broader thermal niche breadth reduces this risk by increasing the chances that newly

colonised areas fall within the species' tolerable thermal range. However, the causal relationship could

also work in reverse, as limited dispersal capacity may constrain a species' realised thermal niche by

restricting access to potentially suitable regions, despite physiological tolerance to those climates (Sax

et al. 2013). In other words, the realised thermal niche, as considered in this study, would be narrower

than the fundamental thermal niche, simply because of dispersal limitations. Implementation of other

niche components such as diet or habitat specialisation could have provided a more holistic view on

the relationship between dispersal and ecological generalisation. It is worthwhile to notice that the

retrieved thermal niche-body size relationships were predominantly negative, which aligns with

experimentally derived patterns of lower thermal tolerance in large-bodied animals (Gunderson 2024).

Besides thermal niche breadth, thermal optima also covary with dispersal ability. Across nearly all

assessed arthropod orders, species adapted to warmer climates exhibit higher dispersal capacities.

Although cold-adapted species may face greater risks from climate change and thus might benefit

more from range shifting as a mitigation strategy, their generally lower dispersal capacities make

spreading a less viable option (Tayleur *et al.* 2016). Furthermore, our findings indicate that low

dispersal capacity often covaries with slower life histories, which hampers colonization success due to

slower population growth rates and, therefore, elevated vulnerability to stochastic extinction (Caswell

& Hastings 1980). Consequently, the arthropod species most at risk from temperature changes due to

climate change are also those least equipped to mitigate its effects through range shifts - a trend also

noted in European mammals (Morrison *et al.* 2018).

Phylogenetic signals were highly variable and strongly influenced by both the specific trait and

arthropod order being assessed. Body size and development time showed strong phylogenetic

conservation in most orders, while thermal niche was largely independent of common ancestry.

Dispersal and fecundity present an interesting pattern, being strongly conserved in some orders but

showing minimal phylogenetic influence in others. A strong phylogenetic signal in body size was

expected, as closely related species often share a similar morphology. Development time is typically

linked to size, as larger species require more time to accumulate tissue (Blueweiss *et al.* 1978; Calder

1996). However, our data do not fully support the idea that larger arthropod species develop more

slowly, as only Coleoptera showed a clear positive relationship between body size and development

time. Instead, it appears that closely related arthropod species have evolved similar pace-of-life

strategies due to comparable life cycle characteristics (Taylor & Karban 1986). The weak phylogenetic

signal in thermal traits aligns nicely with earlier findings suggesting directional selection driven by

physiological constraints and experienced thermal conditions within species' range over evolutionary

history (Bennett *et al.* 2021). The variable phylogenetic signal for dispersal capacity likely reflects the

diverse dispersal mechanisms across arthropod orders. For example, dispersal in flying insects like

Lepidoptera and Odonata tends to be body size-dependent, with larger species typically showing

higher mobility (Kuussaari *et al.* 2014; Sekar 2012; Wootton 2020). In Coleoptera and Hemiptera,

dispersal capacity is closely tied to hind wing development, a trait that is also strongly phylogenetically

conserved (Braendle *et al.* 2006; Roff 1986, 1994). For taxa with more passive dispersal mechanisms,

a lower phylogenetic signal in dispersal capacity is expected, as dispersal is less directly linked to

specific morphological traits. In Araneae, for instance, nearly all species show ballooning behaviour

regardless of size, with larger species typically ballooning as juveniles. Dispersal propensity in spiders

is thus more strongly correlated with habitat stability and specialization, factors that are generally

independent of phylogenetic lineage (Bonte *et al.* 2006).

Data accessibility statement

367 The complete dataset on arthropod life histories and ecological traits can be found on GBIF:

368 https://doi.org/10.15468/75g5z9. The filtered dataset specific to this study can be accessed on

369 Zenodo: 10.5281/zenodo.14527806.

References

Beckman, N.G., Bullock, J.M. & Salguero-Gómez, R. (2018). High dispersal ability is related to fast

life-history strategies. *Journal of Ecology*, 106, 1349–1362.

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. (2012). Impacts of climate

change on the future of biodiversity. *Ecology Letters*, 15, 365–377.

Bennett, J.M., Sunday, J., Calosi, P., Villalobos, F., Martínez, B., Molina-Venegas, R., *et al.* (2021).

The evolution of critical thermal limits of life on Earth. *Nature Communications*, 12, 1198.

Berger, D., Walters, R. & Gotthard, K. (2008). What limits insect fecundity? Body size- and

temperature-dependent egg maturation and oviposition in a butterfly. *Functional Ecology*, 22,

523–529.

Beukhof, E., Frelat, R., Pecuchet, L., Maureaud, A., Dencker, T.S., Sólmundsson, J., *et al.* (2019).

Marine fish traits follow fast-slow continuum across oceans. *Scientific Reports*, 9, 1-9.

Bitume, E.V., Bonte, D., Ronce, O., Bach, F., Flaven, E., Olivieri, I., *et al.* (2013). Density and

genetic relatedness increase dispersal distance in a subsocial organism. *Ecology Letters*, 16, 430–

437.

Blueweiss, L., Fox, H., Kudzma, V., Nakashima, D., Peters, R. & Sams, S. (1978). Relationships

between body size and some life history parameters. *Oecologia*, 37, 257–272.

Bonte, D. (2025). Integrating dispersal in life history. In: *Life history evolution: Traits, Interactions,*

and Applications (eds. Segoli, M. & Wajnberg, E.). Wiley, Hoboken, NJ, pp. 175–185.

Bonte, D. & Dahirel, M. (2017). Dispersal: a central and independent trait in life history. *Oikos*,

126, 472–479.

Bonte, D., Van Dyck, H., Bullock, J.M., Coulon, A., Delgado, M., Gibbs, M., *et al.* (2012). Costs of

dispersal. *Biological Reviews*, 87, 290–312.

Bonte, D., Vanden Borre, J., Lens, L. & Maelfait, J.-P. (2006). Geographical variation in wolf spider

dispersal behaviour is related to landscape structure. *Animal Behaviour*, 72, 655–662.

Braendle, C., Davis, G.K., Brisson, J.A. & Stern, D.L. (2006). Wing dimorphism in aphids. *Heredity*,

97, 192–199.

Buoro, M. & Carlson, S.M. (2014). Life-history syndromes: Integrating dispersal through space

and time. *Ecology Letters*, 17, 756–767.

Bürkner, P.-C. (2021). Bayesian Item Response Modeling in R with brms and Stan. *Journal of*

Statistical Software, 100, 1–54.

Calder, W.A. (1996). *Size, Function, and Life History*. Dover Publications, New York.

Carbonell, J.A. & Stoks, R. (2020). Thermal evolution of life history and heat tolerance during

range expansions toward warmer and cooler regions. *Ecology*, 101, e03134.

Caswell, H. & Hastings, A. (1980). Fecundity, developmental time, and population growth rate: An

analytical solution. *Theoretical Population Biology*, 17, 71–79.

Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S. & Massot, M. (2009). Informed dispersal,

heterogeneity in animal dispersal syndromes and the dynamics of spatially structured

populations. *Ecology Letters*, 12, 197–209.

Cousens, R., Dytham, C. & Law, R. (2008). *Dispersal in Plants: A Population Perspective*. Oxford

University Press, Oxford.

Dammhahn, M., Dingemanse, N.J., Niemelä, P.T. & Réale, D. (2018). Pace-of-life syndromes: a

framework for the adaptive integration of behaviour, physiology and life history. *Behavioural*

Ecology and Sociobiology, 72, 62, s00265-018-2473-y.

Estrada, A., Morales-Castilla, I., Caplat, P. & Early, R. (2016). Usefulness of Species Traits in

Predicting Range Shifts. *Trends in Ecology & Evolution*, 31, 190–203.

Foden, W.B., Butchart, S.H.M., Stuart, S.N., Vié, J.-C., Akçakaya, H.R., Angulo, A., *et al.* (2013).

Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based

Assessment of all Birds, Amphibians and Corals. *PLoS ONE*, 8, e65427.

Foden, W.B. & Young, B.E. (2016). *IUCN SSC guidelines for assessing species' vulnerability to*

climate change. Version 1.0. IUCN Species Survival Commission, Cambridge, England.

Gaillard, J.-M., Pontier, D., Allainé, D., Lebreton, J.D., Trouvilliez, J., Clobert, J., *et al.* (1989). An

Analysis of Demographic Tactics in Birds and Mammals. *Oikos*, 56, 59-76.

Ganihar, S.R. (1997). Biomass estimates of terrestrial arthropods based on body length. *Journal*

of Biosciences, 22, 219–224.

Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010). A framework for

community interactions under climate change. *Trends in Ecology & Evolution*, 25, 325–331.

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal society of*

London, Biological Sciences, 326, 119–157.

Grinder, R.M. & Wiens, J.J. (2023). Niche width predicts extinction from climate change and

vulnerability of tropical species. *Global Change Biology*, 29, 618–630.

Gunderson, A.R. (2024). Disentangling physiological and physical explanations for body size-

dependent thermal tolerance. *Journal of Experimental Biology*, 227, jeb245645.

Gvoždík, L. (2018). Just what is the thermal niche? *Oikos*, 127, 1701–1710.

Habibullah, M.S., Din, B.H., Tan, S.-H. & Zahid, H. (2022). Impact of climate change on biodiversity

loss: global evidence. *Environmental Science and Pollution Research*, 29, 1073–1086.

Hampe, A. & Jump, A.A. (2011). Climate relicts: Past, Present, Future. *Annual Review of Ecology,*

Evolution, and Systematics, 42, 313–333.

Hill, J.K., Thomas, C.D. & Huntley, B. (2001). Climate and recent range changes in butterflies. In:

"Fingerprints" of Climate Change (eds. Walther, GR., Burga, C.A. & Edwards, P.J.). Springer,

Boston, pp. 77–88.

Hillaert, J., Boeye, J., Stoks, R. & Bonte, D. (2015). The evolution of thermal performance can

constrain dispersal during range shifting. *Journal of Biological Dynamics*, 9, 317–335.

Hillaert, J., Hovestadt, T., Vandegehuchte, M.L. & Bonte, D. (2018). Size-dependent movement

explains why bigger is better in fragmented landscapes. *Ecology and Evolution*, 8, 10754–10767.

Jenkins, D.G., Brescacin, C.R., Duxbury, C.V., Elliott, J.A., Evans, J.A., Grablow, K.R., *et al.* (2007).

Does size matter for dispersal distance? *Global Ecology and Biogeography*, 16, 415–425.

Klockmann, M., Günter, F. & Fischer, K. (2017). Heat resistance throughout ontogeny: body size

constrains thermal tolerance. *Global Change Biology*, 23, 686–696.

Kuussaari, M., Saarinen, M., Korpela, E., Pöyry, J. & Hyvönen, T. (2014). Higher mobility of

butterflies than moths connected to habitat suitability and body size in a release experiment.

Ecology and Evolution, 4, 3800–3811.

La Sorte, F.A. & Jetz, W. (2010). Projected range contractions of montane biodiversity under

global warming. *Proceedings of the Royal Society B: Biological Sciences*, 277, 3401–3410.

Logghe, G., Batsleer, F., Maes, D., Permentier, T., Berg, M.P., Brosens, D., *et al.* (2025). An in-

depth dataset of northwestern European arthropod life histories and ecological traits.

Biodiversity Data Journal, 13, e146785.

MacLean, S.A. & Beissinger, S.R. (2017). Species' traits as predictors of range shifts under

contemporary climate change: A review and meta-analysis. *Global Change Biology*, 23, 4094–

4105.

Martin, C.A., Proulx, R. & Magnan, P. (2014). The biogeography of insects' length-dry mass

relationships. *Insect Conservation and Diversity*, 7, 413–419.

Matthew, P., Eastman, J.M., Slater, G.J., Brown, J.W., Uyeda, J.C., FitzJohn, R.G., *et al.* (2014).

geiger v2.0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic

trees. *Bioinformatics*, 30, 2216–2218.

Maxwell, S.L., Butt, N., Maron, M., McAlpine, C.A., Chapman, S., Ullmann, A., *et al.* (2019).

Conservation implications of ecological responses to extreme weather and climate events.

Diversity and Distributions, 25, 613–625.

Michonneau, F., Brown, J.W. & Winter, D.J. (2016). rotl: an R package to interact with the Open

Tree of Life data. *Methods in Ecology and Evolution*, 7, 1476–1481.

Morrison, L., Estrada, A. & Early, R. (2018). Species traits suggest European mammals facing the

greatest climate change are also least able to colonize new locations. *Diversity and Distributions*,

24, 1321–1332.

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, 401, 877–884.

Paradis, E. & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*, 35, 526–528.

Pearse, W.D., Davies, T.J. & Wolkovich, E.M. (2025). How to Define, Use, and Interpret Pagel's λ

(Lambda) in Ecology and Evolution. *Global Ecology and Biogeography*, 34, e70012.

Pianka, E.R. (1970). On r- and K-Selection. *The American Naturalist*, 104, 592–597.

Polverino, G., Santostefano, F., Díaz-Gil, C. & Mehner, T. (2018). Ecological conditions drive pace-

of-life syndromes by shaping relationships between life history, physiology and behaviour in two

populations of Eastern mosquitofish. *Scientific Reports*, 8, 1-10.

Pöyry, J., Luoto, M., Heikkinen, R.K., Kuussaari, M. & Saarinen, K. (2009). Species traits explain

recent range shifts of Finnish butterflies. *Global Change Biology*, 15, 732–743.

Promislow, D.E.L. & Harvey, P.H. (1990). Living fast and dying young: A comparative analysis of

life-history variation among mammals. *Journal of Zoology*, 220, 417–437.

R Core Team. (2021). R: A language and environment for statistical computing.

Revell, L.J. (2012). phytools: An R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution*, 3, 217–223.

Reynolds, J.D. (2003). Life histories and extinction risk. In: *Macroecology*. Blackwell Publishing,

Oxford, pp. 195–217.

Roff, D.A. (1986). The Evolution of Wing Dimorphism in Insects. *Evolution*, 40, 1009–1020.

Roff, D.A. (1994). The evolution of flightlessness: Is history important? *Evolutionary Ecology*, 8,

639–657.

Ronce, O. & Clobert, J. (2012). Dispersal syndromes. In: *Dispersal Ecology and Evolution*. Oxford

University Press, Oxford, pp. 119–138.

Sæther, B.-E. (1987). The Influence of Body Weight on the Covariation between Reproductive

Traits in European Birds. *Oikos*, 48, 79-88.

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-Ache, C., *et

al.* (2016). Fast–slow continuum and reproductive strategies structure plant life-history variation

worldwide. *Proceedings of the National Academy of Sciences*, 113, 230–235.

Saupe, E.E., Qiao, H., Hendricks, J.R., Portell, R.W., Hunter, S.J., Soberón, J., *et al.* (2015). Niche

breadth and geographic range size as determinants of species survival on geological time scales.

Global Ecology and Biogeography, 24, 1159–1169.

Sax, D.F., Early, R. & Bellemare, J. (2013). Niche syndromes, species extinction risks, and

management under climate change. *Trends in Ecology & Evolution*, 28, 517–523.

Sekar, S. (2012). A meta-analysis of the traits affecting dispersal ability in butterflies: can

wingspan be used as a proxy?: Factors affecting dispersal ability in butterflies. *Journal of Animal*

Ecology, 81, 174–184.

Settele, J., Scholes, R., Betts, R., Bunn, S.E., Leadley, P., Nepstad, D., *et al.* (2014). Terrestrial and

inland water systems. In: *Climate change 2014: Impacts, adaptation, and vulnerability. Fifth*

assessment report of the Intergovernmental Panel of Climate Change. Cambridge University

Press, New York, NY, pp. 1–153.

Snyder, R.E. (2011). Leaving home ain't easy: non-local seed dispersal is only evolutionarily stable

in highly unpredictable environments. *Proceedings of the Royal Society B*, 278, 739–744.

Southwood, T.R.E. (1977). Habitat, the Templet for Ecological Strategies? *Journal of Animal*

Ecology, 46, 336-365.

Stearns, S.C. (1983). The Influence of Size and Phylogeny on Patterns of Covariation among Life-

History Traits in the Mammals. *Oikos*, 41, 173-187.

Stevens, V.M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J. & Baguette, M. (2013).

Dispersal syndromes and the use of life-histories to predict dispersal. *Evolutionary Applications*,

6, 630–642.

Stevens, V.M., Whitmee, S., Le Galliard, J., Clobert, J., Böhning-Gaese, K., Bonte, D., *et al.* (2014).

A comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial animals. *Ecology*

Letters, 17, 1039–1052.

Stralberg, D., Jongsomjit, D., Howell, C.A., Snyder, M.A., Alexander, J.D., Wiens, J.A., *et al.* (2009).

Re-Shuffling of Species with Climate Disruption: A No-Analog Future for California Birds? *PLoS*

ONE, 4, e6825.

Tayleur, C.M., Devictor, V., Gaüzère, P., Jonzén, N., Smith, H.G. & Lindström, Å. (2016). Regional

variation in climate change winners and losers highlights the rapid loss of cold-dwelling species.

Diversity and Distributions, 22, 468–480.

Taylor, F. & Karban, R. (1986). *The Evolution of Insect Life Cycles*. Proceedings in Life Sciences.

Springer New York, New York.

Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., *et al.*

(2004). Extinction risk from climate change. *Letters to Nature*, 427, 145-148.

Travis, J.M.J., Mustin, K., Bartoń, K.A., Benton, T.G., Clobert, J., Delgado, M.M., *et al.* (2012).

Modelling dispersal: an eco-evolutionary framework incorporating emigration, movement,

settlement behaviour and the multiple costs involved. *Methods in Ecology and Evolution*, 3, 628–

641.

Urban, M.C. (2019). Projecting biological impacts from climate change like a climate scientist.

WIREs Climate Change, 10, e585.

Urban, M.C., Bocedi, G., Hendry, A.P., Mihoub, J.-B., Pe'er, G., Singer, A., *et al.* (2016). Improving

the forecast for biodiversity under climate change. *Science*, 353, aad8466.

Wagner, D.L. (2020). Insect Declines in the Anthropocene. *Annu. Rev. Entomol.*, 65, 457–480.

Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., *et al.* (2001). Rapid responses

of British butterflies to opposing forces of climate and habitat change. *Letters to Nature*, 414, 65–

69.

Weil, S., Gallien, L., Lavergne, S., Börger, L., Hassler, G.W., Nicolaï, M.P.J., *et al.* (2022).

Chameleon biogeographic dispersal is associated with extreme life history strategies. *Ecography*,

2022, e06323.

Weil, S.-S., Gallien, L., Nicolaï, M.P.J., Lavergne, S., Börger, L. & Allen, W.L. (2023). Body size and

life history shape the historical biogeography of tetrapods. *Nature Ecology & Evolution*, 7, 1467–

1479.

Whitmee, S. & Orme, C.D.L. (2013). Predicting dispersal distance in mammals: a trait-based

approach. *Journal of Animal Ecology*, 82, 211–221.

Williams, W.W. & Jackson, S.T. (2007). Novel climates, no-analog communities, and ecological

surprises. *Palaeoecology*, 5, 475–482.

Wootton, R. (2020). Dragonfly flight: morphology, performance and behaviour. *International*

Journal of Odonatology, 23, 31–39.

Wright, J., Bolstad, G.H., Araya-Ajoy, Y.G. & Dingemanse, N.J. (2019). Life-history evolution under

fluctuating density-dependent selection and the adaptive alignment of pace-of-life syndromes.

Biological Reviews, 94, 230–247.

Yang, L.H. & Gratton, C. (2014). Insects as drivers of ecosystem processes. *Current Opinion in*

Insect Science, 2, 26–32.

Acknowledgements

The authors would like to thank all contributors to the original dataset from which this study's data were sampled: Tristan Permentier, Matty P. Berg, Pallieter De Smedt, Jonas Hagge, Jorg Lambrechts, Marc Pollet and Fons Verheyde. The computational resources (Stevin Supercomputer Infrastructure) and services used in this work were provided by the VSC (Flemish Supercomputer Center), funded by Ghent University, FWO and the Flemish Government – department EWI. Finally, the authors extend their gratitude to the Flemish Research Fund for their support of the Scientific Research Network EVENET (W001322N). All arthropod icons used in this paper originate from Noun Project (CC BY-NC-ND 2.0): Spider by Matthew Davis, Beetle by Rachel Siao, Cicada by Alejandro Capellan, wasp by Parkji Sun, isopod by Pham Thanh Lôc, Moth by Parkji Sun, Dragonfly by Hermine Blanquart and Cricket by Ed Harrison.

Funding

Garben Logghe was funded by an FWO-INBO doctoral fellowship (grant nr: 1130223N).

Ethics declarations

The authors declare that they have no competing interests.