Impacts of Habitat Transformation on Amphibian and Reptile Communities in a

Heterogeneous Andean Landscape

Nelson Falcón-Espitia [1,2,*] , Juan Camilo Ríos-Orjuela [1,2,*] , Sebastian Perez-Rojas [3]

, Dennys Plazas-Cardona [4] & Alejandra Arias-Escobar [5]

[1] Laboratorio de Biología Evolutiva de Vertebrados, Departamento de Ciencias Biológicas,

Universidad de los Andes, Bogotá, Colombia. ZIP 111711.

[2] Grupo de Morfología y Ecología Evolutiva, Instituto de Ciencias Naturales, Universidad

Nacional de Colombia, Sede Bogotá, Colombia. ZIP 111321.

[3] Independent researcher.

[4] SELVA: Investigación para la Conservación en el Neotrópico. ZIP 111311.

[5] Independent researcher.

*Corresponding authors: ngfalcone@unal.edu.co, jcriosor@gmail.com.

Orcid Id's: NFE: 0000-0002-0959-7072, JCR: 0000-0001-6976-9131, SPR: 0000-0001-7855-

9101, DPC: 0009-0003-3877-2364, AAE: 0000-0002-1356-2243.

Abstract

The Andean region of Colombia is recognized for its exceptional biodiversity and high levels

of endemism, particularly among amphibians and reptiles. However, this region faces

significant challenges due to habitat loss and fragmentation driven by agricultural activities,

urbanization, and livestock farming. This study evaluates the effects of habitat alterations on

herpetofaunal communities within the Sumapaz province, focusing on three habitat types:

secondary Andean forests, coffee plantations, and open areas associated with low-intensity

livestock farming. We also compiled a comprehensive inventory of 85 species based on

fieldwork conducted (2021-2022) and biodiversity databases, recognizing 35 amphibians and

50 reptiles for the region. We found that forests support the highest richness and diversity of

species, while open areas exhibited greater reptile abundance. Crops showed lower biodiversity

but demonstrated potential as secondary habitats for certain species. These findings highlight

the importance of preserving forested areas and designing mixed-use landscapes to promote

biodiversity and maintain ecological functions. This study provides a baseline for future

conservation strategies in the Sumapaz region.

Keywords: Agroecosystems, habitat use, biodiversity conservation, anthropic disturbance,

herpetofauna inventory.

Introduction

In Colombia, Andean ecosystems have undergone significant transformation, particularly in sub-Andean and Andean forests. Today, only 39.5% of their original area remains as natural ecosystems, while the remaining 60.5% has been altered [1]. The Andean region is home to approximately 53% of the country's total herpetofauna species [2], including around 317 endemic species [3,4]. This high level of biodiversity and endemism makes the region one of the top global priorities for herpetofauna conservation [5–7].

One of the most significant threats to amphibian and reptile community assemblages is land use alteration for human activities, such as urbanization or agricultural purposes. This practices often leads to the loss and degradation of forest habitats, leaving only fragmented remnants within heavily disturbed ecosystems. [8,9]. Given their ecological traits, amphibians and reptiles display diverse responses to environmental disturbances, with potential positive or negative effects on the richness and abundance of herpetofaunal species [9]. Both amphibians and reptiles are significantly affected by urbanization; however, amphibian richness is more heavily impacted by activities such as deforestation and forestry, whereas reptiles are more affected by increased livestock intensity [10]. As a result, habitat alterations resulting from human activity significantly influence the composition, structure, and dynamics of amphibian and reptilian communities, thereby affecting ecological interactions in the ecosystem [11–13].

Different strategies have been proposed to reduce and mitigate the impacts of habitat loss and fragmentation due to changes in land use on species diversity [8,14,15]. Thus, it has been proposed that some agroecosystems can protect wildlife species in human-modified landscapes, maintaining both species diversity and the economic benefits of crops [16]. For example, coffee plantations have been suggested as biodiversity-friendly land uses with relatively low environmental impact [17–19]. Nonetheless, the effects of landscape-scale transformations and human activities on herpetofaunal community assemblages remain largely unknown in Colombia.

Similarly, knowledge about the herpetofauna in Colombian Andes and the Sumapaz province, where agricultural activities, livestock farming, and urbanization are integral parts of the landscape remains limited. This area of great conservation importance lacks comprehensive data on amphibian and reptile species composition and ecology [20]. Such knowledge is fundamental to the evaluation of the impacts of land-use alterations on herpetofaunal communities and the formulation of effective conservation strategies [19].

For all reasons mentioned above, we evaluated the changes in species richness and abundance
of amphibians and reptiles in three different habitats with different degrees of anthropic
intervention in the Eastern Cordillera of Colombia: secondary Andean forests (named hereafter
simply as forest, the most conserved habitats remaining in the region today, representing the
closest approximation to a natural assemblage currently found in the area), crops (specifically
coffee plantations, which dominate the agroforestry activities in the region), and open areas
(primarily associated with low-intensity livestock farming activities in the region). All these
areas are spatially interconnected, with varying degrees of connectivity between natural and
human-altered landscapes.

In addition, we compiled historical and current records of the species present in the Sumapaz
province to consolidate an inventory that will serve as a starting point for the establishment of
conservation strategies for the Sumapaz river basin herpetofauna.

Materials and Methods

Study area

We carried out fieldwork in the municipalities of Tibacuy, Nilo and Viotá located in the
Sumapaz province in Cundinamarca, Colombia, between 645 and 1840 m elevation (Figure 1).
The region has a mostly humid mountain climate in combination with dry enclaves, typical of
the middle lands of the Magdalena basin. The study area is characterized by secondary Andean
forest and agroecosystems in a heterogeneous matrix where it is possible to find forest reserve
areas and private properties used for mixed crops and livestock farming [21]. We conducted
the study in three habitat types: forest (11 localities), with scattered trees over five meters high,
where advanced plant regeneration processes are evident; shaded coffee crops (6 localities) and
open areas (9 localities) used as pastures for permanent livestock.

Figure 1. Map of Study Area in the Eastern Cordillera of the Andes, Colombia. The figure shows the locations of fieldwork within the municipalities of Tibacuy, Nilo, and Viotá, situated in the Eastern Cordillera of the Andes, Cundinamarca, Colombia. Field sites encompass a heterogeneous matrix of diverse habitat types, including secondary forest, shaded coffee plantations, and open pastures.

Data collection

We conducted a total of eight field trips (three effective sampling days each) between June of 2021 and July of 2022 to cover the rainy and dry seasons in the region. During the sampling phase, we used the technique of visual and auditory encounter surveys (VES) for a limited time and fixed band transects (2 X 50 m) in which each encounter was counted individually for abundance estimation during a sampling event [22]. We carried out sampling between 8:00 and

12:00 and between 18:00 and 22:00 h, surveying microhabitats suitable for amphibians and
reptiles such as vegetation, ponds, streams and leaf litter, with a maximum limit of 5 m from
the ground [23]. During all the sampling phase, data as habitat, body measurements and
photographic records for all individuals were collected on the site of observation. The
individuals were released at the same location where they were captured.

In addition to the field trips conducted in 2021 (33 species, 153 individuals), historical records
from surveys between 2015 and 2019 (51 species, 455 individuals, of which 24 species were
not reported in 2021) were also included in the analysis. These records were collected by the
authors of this study and the herpetology group of Universidad Nacional de Colombia
(Herpetos UN) following the VES methodology described above. Finally, we followed Frost
[24] and Uetz *et al* [25] as classification systems for amphibians and reptiles, respectively.

Database searching

As a supplement to the species list obtained in the study area, we consulted the databases of
the Biodiversity Information System of Colombia (SiB). We filtered records that came from
preserved specimens, machine observations and observations. The latter were only taken from
institutions with local environmental authority (Corporación Autónoma Regional de
Cundinamarca - CAR) to avoid errors in species identification. We also reviewed historical
data in biological collections at the Instituto de Ciencias Naturales (ICN) at Universidad
Nacional de Colombia and the Amphibian and Reptiles collections of the Instituto de
Investigación de Recursos Biológicos Alexander von Humboldt (IAvH) for the municipalities
of Tibacuy, Nilo, Silvania and Viotá (Cundinamarca). These supplement species records were
used to compile a complete list of amphibians and reptiles for the Sumapaz Province. However,
they were not included in the assemblage analyses due to a lack of standardization and reliable
information about their association with vegetation cover at the time of capture.

Data analysis

We used a species accumulation curve to determine the representativeness of the sampling
following the guidelines of Villarreal *et al* [22]. For this, we used the Chao 1, ACE and Cole's
Rarefaction estimators with abundance-based data, analyzing the herpetofaunal community as
a whole and the amphibian and reptile communities separately. We calculated curves using
EstimateS 9.1.0 software [26] for the 52 sampling units performed.

We assessed the diversity of reptiles and amphibians in the study area by calculating five diversity indices: species richness, Shannon-Wiener index, Simpson's index, Pielou's equitability, and Margalef's richness index. Such indices were chosen for their ability to reflect both species' richness and evenness in community structure. We used the *vegan* package in R to calculate these diversity indices, as it provides robust and widely accepted methods for ecological data analysis [27].

To evaluate species richness at varying sampling efforts and predict species richness beyond our observed sample size, we estimated the percentage of completeness achieved through sampling coverage curves, as well as the additional sampling necessary to reach the maximum number of species. This analysis estimates the proportion of the total number of individuals in a community that belong to a sampled species [28]. We used the *iNEXT* package [29] to create the 95% confidence intervals (IC) with a resampling of 1000 bootstraps. Furthermore, we plotted rank-abundance curves to compare the species abundance distributions among habitats.

To assess the dominance and community structure, we analyzed the species distribution across the three different habitat types: forest, crops, and open areas (see specific definition of habitat types in Ríos-Orjuela *et al* [19]). Using the *rankabuncomp* function from the *vegan* package [27], we generate rank-abundance curves for reptiles and amphibians which allows for the comparison of species' richness, evenness, and dominance patterns among habitats. We organized species abundance by habitat type, focusing on the three most abundant species per habitat. This subset provided insights into habitat-specific species dominance patterns. Rank-abundance plots were generated using *ggplot2* [30], complemented with *ggrepel* to annotate dominant species and ensure clear visualization of key species. We applied log-transformed abundance values to better illustrate the distribution across ranks. The rank-abundance plots provide detailed insights into how individual species contribute to the overall community structure, helping us evaluate biodiversity patterns beyond simple species counts. We also used stacked bar plots to explore species richness and individual abundance across the three previously mentioned habitats. All analyses were conducted in R version 4.1.0. [31].

Color schemes were standardized to maintain consistency between habitat types, with each habitat type assigned a distinct color palette. We used a color-blind friendly palette to ensure accessibility to all readers.

Results

Based on the information stored in databases, biological collections and literature regarding herpetofauna in the analyzed area of the Sumapaz River province we found records of 58 species (Table S1): 24 amphibians (21 anurans, 1 salamander and 2 caecilians) and 34 reptiles (11 lizards, 21 snakes, 1 tortoise and 1 crocodilian). As a result of our fieldwork, we recorded a total of 57 species (21 amphibians and 36 reptiles), 27 of these species are additional records for the region. Then, we were able to compile a more comprehensive state of knowledge to better understand the diversity of herpetofauna in this area.

A total of 85 species (35 amphibians and 50 reptiles) have been recorded in the municipalities in the study area (Table S1). For amphibians, most species belong to the order Anura, with Hylidae having the highest richness in the region (8 spp.), followed by Dendrobatidae and Strabomantidae (5 spp. each). Caecilians (Gymnophiona) were represented by *Caecilia thompsoni* (Caeciliidae) and *Microcaecilia* sp. (Siphonopidae) and salamanders (Caudata) by *Bolitoglossa muisca* (Plethodontidae). For reptiles, within the order Squamata, the family with the highest richness was Colubridae (23 spp.), followed by Anolidae (6 spp.). The orders Crocodylia and Testudines were represented by a single species each (*Caiman crocodilus* and *Kinosternon* sp.). There are records of two invasive species in the area, the North American Bullfrog *Aquarana catesbeiana* (Ranidae) and the South Asian House Gecko *Hemidactylus frenatus* (Gekkonidae).

Regarding the data from our fieldwork, both Hylidae (5 spp.) and Leptodactylidae (4 spp.) had the highest richness for amphibians, as for reptiles it was Colubridae (23 spp.) and Anolidae (4 spp.). In terms of abundance, we recorded 608 individuals: 391 amphibians and 217 reptiles (Table 1). The most abundant amphibian species were *Boana platanera* (62), *Engystomops pustulosus* (60) and *Pristimantis taeniatus* (55), while for reptiles *Gonatodes albogularis* (53), *Cnemidophorus lemniscatus* (33) and *Hemidactylus frenatus* (17) were the most abundant species in the sampled area (Table S1). Forest sustained the highest numbers of individuals for amphibians and the herpetofauna, but instead reptiles were slightly more abundant in open areas. We found consistently less individuals in crops.

Regarding diversity, the Shannon-Wiener index was highest for the entire herpetofauna community (3.237), followed by reptiles (2.793) and amphibians (2.469), indicating that the herpetofauna community exhibited a more even distribution of species compared to the reptile and amphibian groups individually. Simpson's index showed similarly high diversity across the groups, with values of 0.944 for herpetofauna, 0.894 for reptiles, and 0.896 for amphibians,

reflecting low dominance across species in the assemblages. In terms of species evenness,
Pielou's equitability index was similar across groups, with reptiles at 0.779 and amphibians at
0.811. Margalef's index indicated that the reptile community (6.506) had greater species
richness relative to sample size than the amphibian community (3.351), as shown in Table 1.
In general terms, forests showed higher diversity scores for herpetofauna as a whole and for
amphibians and reptiles separately (Figure 2). However, open areas showed a higher abundance
of reptiles and evenness for amphibians, while evenness in the reptile community in coffee
plantations was higher.

Table 1. Diversity indices for herpetofauna, reptiles, and amphibians. This table presents
the calculated diversity indices, including species richness, Shannon, Simpson, Pielou's
equitability, and Margalef's index for the herpetofauna community, as well as for the reptiles
and amphibians separately. Richness represents the number of species, while the Shannon and
Simpson indices measure diversity and dominance. Equitability reflects how evenly
individuals are distributed among species, and Margalef's index provides a measure of species
richness considering the number of individuals sampled.

	Herpetofauna			Reptiles			Amphibians		
	Forest	Crops	Open areas	Forest	Crops	Open areas	Forest	Crops	Open areas
Richness	45	14	34	24	8	23	21	6	11
Abundance	319	47	242	87	12	118	232	35	124
Simpson_1-D	0.929	0.758	0.922	0.919	0.894	0.841	0.877	0.571	0.848
Shannon_H	3.137	2.034	2.884	2.899	2.199	2.324	2.418	1.2	2.061
Margalef	7.632	3.376	6.012	5.15	2.817	4.611	3.672	1.406	2.075
Equitability_J	0.824	0.771	0.818	0.912	1.057	0.741	0.794	0.67	0.860

	Herpetofauna	Reptiles	Amphibians
Richness	57	36	21
Abundance	608	217	391
Simpson_1-D	0.944	0.894	0.896
Shannon_H	3.237	2.793	2.469
Margalef	8.736	6.506	3.351
Equitability_J	0.801	0.779	0.811

215 The species accumulation and sample coverage curves (Figure 3A and 3B) demonstrate that
216 the diversity of both reptiles and amphibians approached saturation with increased sampling.
217 Amphibians reached the saturation point slightly earlier. Extrapolation analyses suggest that

additional sampling beyond 1000 individuals would not significantly increase species richness, particularly for amphibians. Both amphibian and reptile communities achieved near-complete sample coverage, with amphibians reaching this threshold faster (Figure 3B). The accumulation curves (Figures 3C and 3D) present results from different estimation methods (S(est), ACE, Chao, and Cole's rarefaction). Singleton and doubleton species—those recorded only once or twice—highlight the potential for undiscovered species, especially in reptiles.

Figure 2. Biodiversity Metrics of Herpetofauna, Reptiles, and Amphibians Across Different Habitats. This figure presents radar charts comparing biodiversity metrics (Richness, Abundance, Equitability, Shannon Index, and Simpson Index) across different land use types (Forest, Crops, and Open areas) for the Herpetofauna, Reptiles, and Amphibians communities. Each plot illustrates the relative values of these metrics within each environment type, coded by color — Forest (blue), Crops (red), and Open areas (green). In herpetofauna, forested areas exhibit the highest scores across most biodiversity metrics, indicating greater biodiversity and ecosystem stability relative to other land use types. Open areas maintain intermediate values, whereas cropland shows the lowest biodiversity metrics across all indices. For reptiles, like herpetofauna, natural forested areas show higher values in richness, abundance, and diversity (Simpson and Shannon indices), suggesting forest habitats support more reptilian biodiversity. Cropland presents the lowest scores, with open areas falling in between. For amphibians, forests exhibit the highest biodiversity metric values, with richness and diversity indices significantly outperforming those in crops and open areas. Cropland remains the least supportive habitat for amphibians among the three types.

Figure 3. Interpolation and extrapolation of species richness and sample coverage for
reptiles and amphibians. (A) Species diversity curves show the relationship between species

richness and the number of individuals sampled, with solid lines representing observed data
and dashed lines indicating extrapolated estimates for reptiles (orange) and amphibians (blue).
Shaded regions represent 95% confidence intervals derived from bootstrapped resampling
(1000 iterations). (B) Sample coverage curves demonstrate the proportion of the total species
captured as a function of the number of individuals sampled. The curves suggest the
completeness of sampling for both groups, with the shaded regions showing 95% confidence
intervals. (C & D) Accumulation curves display species richness estimation methods (S(est),
ACE, Chao and Cole rarefaction) across different sample sizes for amphibians (C) and reptiles
(D). Singleton and doubleton species (species recorded only once or twice) are highlighted to
show their impact on richness estimations. (E) Bar plots compare species richness (purple) and
individual abundance (green) across three habitat types: forest, crops, and open areas. The
forest habitat supports higher species richness, while open areas show higher individual
abundance.

Forest areas supported the highest species richness (Figure 4A), with amphibians such as
Engystomops pustulosus and *Pristimantis gaigei* showing high abundance. Crops exhibited the
lowest species richness, but still hosting certain species with few dominants like *P. taeniatus*.
Open areas presented intermediate species richness but higher overall abundance, with key
reptile species such as *Gonatodes albogularis* and *Cnemidophorus lemniscatus* dominating
(Figure 4B). These findings reveal clear habitat preferences among species, with amphibians
more common in forest habitats and reptiles more evenly distributed across the different
environments.

Figure 4. Species abundance and rank distributions across three habitat types. (A) Species abundance curves compare the distribution of species across the three habitat types. Forest habitats (blue circles) have higher abundance and evenness, while crop areas (red triangles)

show steep declines in abundance, indicating lower species diversity. Open areas (green
crosses) display intermediate patterns of species distribution. (B) Rank-abundance plots
display species-specific abundances (log-transformed) within each habitat type. Key
amphibian and reptile species for each habitat are highlighted: *E. pustulosus* and *P. gaigei*
dominate in forests; *P. taeniatus* is prevalent in crop areas; and *G. albogularis* and *C.*
lemniscatus are common in open areas. These plots illustrate differences in community
structure and dominance among the habitats.

Discussion

List of amphibians and reptiles and comments on records

This area of the Sumapaz River province holds 85 species of amphibians and reptiles. The
environmental conditions of the Eastern Andes Cordillera ecoregion considerably influence the
herpetofaunal assemblage [32], as well as the landscape dynamics that have determined habitat
availability for these taxonomic groups. Previous records reviewed in this study from literature
[33–37] and biodiversity databases accounted for around 68% of the diversity present in the
area, therefore our results from this monitoring have increased the knowledge of the local
diversity and allowed us to make different ecological inferences.

It is important to note that most of the "new" records provided by our sampling in the study
area are from species whose distribution in the surrounding localities is well known or at least
are expected to occur (i.e. *Leptodactylus fragilis*, *Rhinella humboldti*, *Cnemidophorus*
lemniscatus or *Bothrops asper*), however, this review provides a more robust base line about
the current knowledge of the biodiversity in the region, which can be helpful for future studies,
local communities and, stakeholders for decision-making and territorial planning.

There are some records for both amphibians and reptiles worth discussing. We were unable to
determine one species of *Anolis* based on the current knowledge of its diversity in the region.
With a preliminary review on lepidosis and morphology we tentatively assigned it as close
relative to *A. granuliceps*, but further work is required to define its identity. This also
underscores the uncharacterized biodiversity of the region that only recently has been studied
deeper. Moreno-Arias *et al* [38] recently described *Anolis tequendamae* whose distribution
spans to this area, *Bolitoglossa muisca* was described with material from Silvania municipality
by López-Perilla *et al* [34] and Acosta-Galvis *et al* [33] described *Hyloxalus arliensis* with

material from several localities in the Magdalena River valley including samples from Viotá.

We now contribute new records of this species in the lowland humid forests of Tibacuy (vereda

El Mango). Also recent papers about the taxonomy and systematics of *Scinax* [39,40] besides

the molecular evidence provided by Guarnizo *et al* [41], opened the discussions to the fact that

specimens usually identified in the local herpetology community as *Scinax ruber* and *S. x-*

signatus in this region are more than likely undescribed lineages highlighting the need to

develop integrative studies to clarify this taxonomical complex in Colombia.

Another significant and concerning record was from the invasive species *Aquarana*

catesbeiana. After its introduction in Colombia around 1986, this frog was initially reported in

Fusagasugá and Silvania (Cundinamarca) in 1993, where was later observed in nearby

municipalities such as Pandi, Nilo and Arbeláez [42]. This means that the species has been in

this area for at least 30 years, but still with limited knowledge of its actual distribution and

population status. Recently CAR [43] have gathered records in Tibacuy and nearby

municipalities in the process of formulating a management plan to control and monitor the

species. Our fieldwork provided one additional record in Tibacuy in 2019, but we did not

observe it in Nilo or Viotá. Besides active sampling, to increase the understanding about *A.*

catesbeiana in Cundinamarca we suggest that future work could benefit from advances in

passive acoustic monitoring and citizen science initiatives.

There are also several relevant historical records in the municipalities studied, especially

toward higher altitudes. Species such as *Atelopus subornatus* and *Hyloxalus ruizi* were

observed in the area surrounding El Alto de San Miguel (within the limit of Sibaté and Silvania

municipalities) by Lynch [35]. Both species are critically endangered [44], and besides

southern records of *A. subornatus* in Tolima [45], there are no recent records in these

municipalities, and we did not register them in this study. Other species currently threatened in

the area worth mentioning are *H. vergeli, P. uisae* and *P. renjiforum* (Table S1).

Overall, our results demonstrate a well-structured herpetofaunal community in this region, with

reptiles exhibiting greater species richness than amphibians. These findings align with previous

studies conducted in similar transitional ecosystems in Cundinamarca, where reptiles are often

more species-rich than amphibians due to their broader adaptability to varied ecological

conditions [46]. The diversity in the study area, as indicated by the Shannon and Simpson

indices, suggests a healthy ecosystem with a balanced distribution of individuals among

species.

Habitat assemblages

Different biodiversity metrics evidenced consistently that forests in this Andean region are key for both amphibians and reptiles. This is expected as natural areas keep higher structural complexity, stable microclimates, several microhabitats and resources availability that suite habitat requirements for species with different life history traits and with less anthropic pressures in general [47,48]. 17 species were only observed in forested areas (7 amphibians and 10 reptiles), although it cannot be assumed that those are forest specialists, some species present traits that make them more specialized in this habitat. For example, riparian forests are essential for species with reproductive modes linked to lotic environments as it is the case for Centrolenidae [49], or they allow for specialized escape behavior as in *B. galeritus* to cross water bodies [50].

The relatively high Margalef's index for reptiles (6.506) compared to amphibians (3.351) reflects the greater species richness within the reptilian community, which may be attributed to the variety of microhabitats available in the transitional forest ecosystem, a trend previously reported by Cervantes-López and Morante-Filho [8]. This region, located between tropical and montane forest zones, provides ideal conditions for reptiles, which are often more tolerant of drier conditions and variable temperatures compared to amphibians, which are more dependent on moisture and stable environmental conditions, making them more susceptible to changes in land use [8,51,52].

Our findings suggest that intermediate habitats, such as croplands, play an important role in the conservation of biodiversity. The presence of species such as *Pristimantis taeniatus* in these environments serves to illustrate the significance of agricultural landscapes in complementing conservation efforts by acting as secondary habitats. These findings lend support to the notion that heterogeneous land-use matrices can serve to enhance biodiversity, as evidenced by recent studies on coffee agroecosystems in the Colombian Andes [19].

The completeness of sample coverage (Figure 3B) highlights the adequacy of our sampling strategy, especially for amphibians, which reached saturation faster than reptiles. This result underscores the need for ongoing monitoring of the group, particularly given the vulnerability of amphibian populations to environmental changes such as habitat loss, climate change and exotic species [23]. The rarefaction and extrapolation curves confirm that our sampling efforts captured most of the species present in the study area. However, the presence of singleton and doubleton species—recorded only once or twice—indicates the potential for undiscovered

species, particularly among reptiles (Figures 3C and 3D). Continued surveys targeting less-
explored microhabitats (e.g., riparian zones or localities at higher altitudes) could reveal
additional species, such as those previously reported from databases or museum specimens,
and which were not observed in our surveys (Table S1).

The differences in richness and abundance between amphibians and reptiles (Fig. 3) show the
expected differential response of both groups to the environmental conditions and perturbation
level present in the study area [9,10]. Amphibian communities presented a lower number of
species, but a higher abundance than reptiles, not only due to the effect of differences in the
ecology or reproductive biology of each group, but also as a consequence of the presence of
species with generalist habits (such as *E. pustulosus, P. taeniatus* and *G. albogularis*) that
colonize semi-natural or disturbed environments with relative ease, from which specialist
species move after disturbance [53], modifying the ecosystem dynamics as a consequence of
changes in dominance at the community level [54].

Although species richness is the most used measure of biodiversity, other measures (such as
functional diversity) are relevant in disturbed environments, where the history of disturbances
in the study ecosystem influences the ecological dynamics within it, reducing functional
diversity and, potentially, the provision of ecosystem services [9,54–57]. Although the
evaluation of functional diversity of amphibian and reptile communities in environmental
gradients for the Colombian Andes is limited, environmental heterogeneity plays an important
role in the degree of association of communities with disturbed environments [58].

Beyond the high richness of amphibians and reptiles observed, the Andean region in Colombia
has undergone a considerable change in the vegetation cover available for the local fauna. As
a result of the intensification of agriculture, cattle ranching and infrastructure development in
the region, the area of forest has decreased considerably [59]. In the Colombian Andes, there
is a differential velocity in land use change along the altitudinal gradient [60] with secondary
forests and grasslands being the most dynamic vegetation units in terms of area gain and loss
[61]. Frequently, such deforested areas go through an intermediate state as crops, which
becomes a constant challenge for the consolidation of amphibian and reptile communities,
favoring ecologically generalist species (such as *E. pustulosus* and *G. albogularis*), which
dominate the transitional environments and relegate species with specific ecological
requirements (such as glass frogs or snakes), resulting in low diversity, homogenization of

ecological functions and a decrease in the ecosystem services provided by amphibians and reptiles within the ecosystem [9,62].

Given the scenario of intensive change in the availability of forested environments in the Andean region and the negative effects of climate change on the populations of amphibians and reptiles that inhabit mountainous areas at intermediate elevations [47,63,64], it is of vital importance to develop long-term monitoring programs to evaluate changes in the distribution of species and ensure the conservation of vulnerable species, mitigating the loss of forest habitats and preserving corridors that promote ecological connectivity [9].

Finally, our study provides evidence about the differential impact of agroforestry systems in transformed landscapes on amphibian and reptile assemblages. In the context of increasing deforestation and expansion of the agricultural frontier, it is of particular interest to identify the characteristics that allow maintaining environments with the ecological quality necessary to sustain stable amphibian and reptile communities, favoring the conservation of forested areas and designing mixed environmental matrices that favor species diversity, and, by extension, the environmental services provided by them.

Supplementary materials

ST1: Species recorded in the municipalities of the study area (Cundinamarca, Colombia).

Document D1: Spanish translation of the present manuscript.

Funding

This work was partially supported by the Asociación Colombiana de Herpetología (ACH) through the Botas al Campo grant (2019).

Data Availability Statement

Data will be available by request to the authors.

Acknowledgements

We thank the Asociación Colombiana de Herpetología for funding though the Botas al Campo grant (2019) and APRENAT for their logistical support. We also acknowledge the cooperation of various individuals and organizations, including the Castillo Urrego family, Bernal Méndez family, Ramos family, Maria Rosa Gaona de Chacón, Mayelly Liévano, Gabriel Garzón, Fray José María Sepúlveda, the Province of Nuestra Señora de la Gracia de Colombia and the

members of Herpetos UN, for their hospitality, collaboration throughout the field phase and for their valuable contributions in the development of this project. To Nataly Casas, Carolina Martínez and Juan Diego Rodríguez for their help during the fieldwork phase and species identification. Finally, we appreciate the ongoing efforts of different communities in Tibacuy, Nilo and Viotá municipalities in conserving their biocultural heritage.

Conflict of interest

The authors declare no conflict of interest.

References

1. Rodríguez, N.; Armenteras, D.; Romero, M. *Ecosistemas de Los Andes Colombianos*; Instituto de Investigación de Recursos Biológicos Alexander von Humboldt: Bogotá, Colombia, 2006; ISBN 9588151309.

2. Romero, M.; Maldonado-Ocampo, J.; K, B.-G.; Usma, J.; Umaña-Villaveces, A.; Murillo, J.; Restrepo-Calle, S.; Álvarez, M.; Palacios-Lozano, M.; Valbuena, M.; et al. *Informe Sobre El Estado de La Biodiversidad En Colombia 2006-2007*; 2009;

3. Armesto, L.; Señaris, J. Anuros Del Norte de Los Andes: Patrones de Riqueza de Especies y Estado de Conservación. *Pap. Avulsos Zool.* **2017**, *57*, 491–526, doi:10.11606/0031-1049.2017.57.39.

4. Bernal, M.; Lynch, J. Review and Analysis of Altitudinal Distribution of the Andean Anurans in Colombia. *Zootaxa* **2008**, 1–25, doi:10.11646/zootaxa.1826.1.1.

5. Bax, V.; Francesconi, W.; Delgado, A. Land-Use Conflicts between Biodiversity Conservation and Extractive Industries in the Peruvian Andes. *J. Environ. Manage.* **2019**, *232*, 1028–1036, doi:10.1016/j.jenvman.2018.12.016.

6. Myers, N.; Mittermeier, R.; Mittermeier, C.; da Fonseca, G.; Kent, J. Biodiversity Hotspots for Conservation Priorities. *Nature* **2000**, *403*, 853–858, doi:10.1038/35002501.

7. Vasconcelos, T.; da Silva, F.; dos Santos, T.; Prado, V.; Provete, D. *Biogeographic Patterns of South American Anurans*; Springer Cham, 2019; ISBN 978-3-030-26295-2.

8. Cervantes-López, M.; Morante-Filho, J. A Global Meta-Analysis on Patterns of

452   Amphibian and Reptile Diversity in Agroforestry Systems. *Glob. Ecol. Conserv.* **2024**,
453   *51*, doi:10.1016/j.gecco.2024.e02914.

454 9. Thompson, M.; Nowakowski, A.; Donnelly, M. The Importance of Defining Focal
455   Assemblages When Evaluating Amphibian and Reptile Responses to Land Use.
456   *Conserv. Biol.* **2015**, *30*, 249–258, doi:10.1111/cobi.12637.

457 10. Cordier, J.; Aguilar, R.; Lescano, J.; Leynaud, G.; Bonino, A.; Miloch, D.; Loyola, R.;
458   Nori, J. A Global Assessment of Amphibian and Reptile Responses to Land-Use
459   Changes. *Biol. Conserv.* **2021**, *253*, doi:https://doi.org/10.1016/j.biocon.2020.108863.

460 11. Ernst, R.; Linsenmair, K.; Rödel, M. Diversity Erosion beyond the Species Level:
461   Dramatic Loss of Functional Diversity after Selective Logging in Two Tropical
462   Amphibian Communities. *Biol. Conserv.* **2006**, *133*, 143–155,
463   doi:10.1016/j.biocon.2006.05.028.

464 12. Gardner, T.; Barlow, J.; Peres, C. Paradox, Presumption and Pitfalls in Conservation
465   Biology: The Importance of Habitat Change for Amphibians and Reptiles. *Biol.*
466   *Conserv.* **2007**, *138*, 166–179, doi:10.1016/j.biocon.2007.04.017.

467 13. Whitfield, S.; Bell, K.; Philippi, T.; Sasa, M.; Bolaños, F.; Chaves, G.; Savage, J.;
468   Donnelly, M. Amphibian and Reptile Declines over 35 Years at La Selva, Costa Rica.
469   *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 8352–8356, doi:10.1073/pnas.0611256104.

470 14. Arroyo-Rodríguez, V.; Fahrig, L.; Tabarelli, M.; Watling, J.; Tischendorf, L.;
471   Benchimol, M.; Cazetta, E.; Faria, D.; Leal, I.; Melo, F.; et al. Designing Optimal
472   Human-Modified Landscapes for Forest Biodiversity Conservation. *Ecol. Lett.* **2020**,
473   *23*, 1404–1420, doi:10.1111/ele.13535.

474 15. Soley, F.; Perfecto, I. A Way Forward for Biodiversity Conservation: High Quality
475   Landscapes. *Trends Ecol. Evol.* **2021**, *36*, 770–773,
476   doi:https://doi.org/10.1016/j.tree.2021.04.012.

477 16. Perfecto, I.; Vandermeer, J. Biodiversity Conservation in Tropical Agroecosystems: A
478   New Conservation Paradigm. *Ann. N. Y. Acad. Sci.* **2008**, *1134*, 173–200,
479   doi:10.1196/annals.1439.011.

480 17. Céspedes, L.; Bayly, N. Over-Winter Ecology and Relative Density of Canada
481   Warbler *Cardellina canadensis* in Colombia: The Basis for Defining Conservation

Priorities for a Sharply Declining Long-Distance Migrant. *Bird Conserv. Int.* **2019**, *29*,

232–248, doi:10.1017/S0959270918000229.

18. González, A.; Wilson, S.; Bayly, N.; Hobson, K. Contrasting the Suitability of Shade

Coffee Agriculture and Native Forest as Overwinter Habitat for Canada Warbler

(*Cardellina Canadensis*) in the Colombian Andes. *Condor* **2020**, *122*, 1–12,

doi:10.1093/condor/duaa011.

19. Ríos-Orjuela, J.C.; Falcón-Espitia, N.; Arias-Escobar, A.; Plazas-Cardona, D.

Conserving Biodiversity in Coffee Agroecosystems: Insights from a Herpetofauna

Study in the Colombian Andes with Sustainable Management Proposal. *Perspect.*

Ecol. Conserv. **2024**, *22*, 196–204, doi:10.1016/j.pecon.2024.04.001.

20. Sguerra, C.; Bejarano, M.; Rodríguez, O.; Blanco, F.; Jaramillo, R.; Sanclemente, Z.

Corredor de Conservación Chingaza-Sumapaz-Guerrero: Resultados Del Diseño y

Lineamientos de Acción; Conservación Internacional Colombia y Empresa de

Acueducto y Alcantarillado de Bogotá ESP, 2011;

21. Peña-Torres, J. Análisis Florístico Del Bosque de Roble (*Quercus Humboldtii*)

Bonpl.(Fagaceae) de La Reserva Protectora Forestal Quininí, Municipio de Tibacuy,

Cundinamarca., 2016.

22. Villarreal, H.; Álvarez, M.; Córdoba, S.; Escobar, F.; Fuga, G.; F., M.; H., O.M.;

Umaña, A. *Manual de Métodos Para El Desarrollo de Inventarios de Biodiversidad*;

2006;

23. Cortés, A.M.; Ramírez-Pinilla, M.P.; Suárez, H.A.; Tovar, E. Edge Effects on

Richness, Abundance and Diversity of Frogs in Andean Cloud Forest Fragments.

South Am. J. Herpetol. **2008**, *3*, 213–222, doi:10.2994/1808-9798-3.3.213.

24. Frost, D. Amphibian Species of the World: An Online Reference. Available online:

https://amphibiansoftheworld.amnh.org/index.php (accessed on 27 October 2024).

25. Uetz, P.; Freed, P.; Aguilar, R.; Reyes, F.; Kudera, J.; Hošek, J. The Reptile Database

Available online: http://www.reptile-database.org/ (accessed on 27 October 2024).

26. Colwell, R. EstimateS: Statistical Estimation of Species Richness and Shared Species

from Samples 2013.

27. Oksanen, F.; Blanchet, G.; Kindt, R.; Legendre, P.; Minchin, P.; O'Hara, R.; Solymos,

P.; Stevens, M.; Szoecs, E.; Wagner, H.; et al. Vegan: Community Ecology Package 2019.

28. Chao, A.; Jost, L. Coverage-Based Rarefaction and Extrapolation: Standardizing Samples by Completeness Rather than Size. *Ecology* **2012**, *93*, 2533–2547, doi:10.1890/11-1952.1.

29. Hsieh, T.; Ma, K.; Chao, A. INEXT: An R Package for Rarefaction and Extrapolation of Species Diversity (Hill Numbers). *Methods Ecol. Evol.* **2016**, *7*, 1451–1456, doi:10.1111/2041-210X.12613.

30. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*; Springer New York, NY, 2009;

31. R Core Team R: A Language and Environment for Statistical Computing 2021.

32. Lynch, J.; Ruiz-Carranza, P.; Ardila-Robayo, M. Biogeographic Patterns of Colombian Frogs and Toads). *Rev. la Acad. Colomb. Ciencias Exactas, Fis. y Nat.* **1997**, *21*, 237–248.

33. Acosta-Galvis, A.; Vargas-Ramírez, M.; Anganoy-Criollo, M.; Ibarra, O.; Gonzáles, S. Description of a New Diminutive *Hyloxalus* (Anura: Dendrobatidae: Hyloxalinae) from the Magdalena Valley of Colombia. *Zootaxa* **2020**, *4758*, 83–102, doi:10.11646/zootaxa.4758.1.3.

34. López-Perilla, Y.; Fernández-Roldán, J.; Meza-Joya, F.; Medina-Rangel, G. A New *Bolitoglossa* (Amphibia, Caudata, Plethodontidae) from the Cordillera Oriental of Colombia. *Zookeys* **2023**, 27–48, doi:10.3897/zookeys.1158.99077.

35. Lynch, J. Two New Species of Poison-Dart Frogs (*Colostethus*) from Colombia. *Herpetologica* **1982**, *38*, 366–374.

36. Cárdenas-Ortega, M.; Gutiérrez-Cárdenas, P.; Cifuentes-Ortegón, M.; Patiño-Gallego, A. *Dendrobates Truncatus* (Cope, 1861). *Catálogo Anfibios y Reptil. Colomb.* **2019**, *5*, 32–41.

37. Ramírez, J. Range Extension of *Pristimantis Uisae* (Lynch, 2003) (Anura: Craugastoridae) in the Cordillera Oriental of Colombia. *Check List* **2017**, *13*, 25–30, doi:10.15560/13.4.25.

38. Moreno-Arias, R.; Méndez-Galeano, M.; Beltrán, I.; Vargas-Ramírez, M. Revealing Anole Diversity in the Highlands of the Northern Andes: New and Resurrected Species of the *Anolis heterodermus* Species Group. *Vertebr. Zool.* **2023**, *73*, 161–188, doi:10.3897/vz.73.e94265.

39. Araujo-Vieira, K.; Lourenço, A.; Lacerda, J.; Lyra, M.; Blotto, B.; Ron, S.; Baldo, D.; Pereyra, M.; Suárez-Mayorga, Á.; Baêta, D.; et al. Treefrog Diversity in the Neotropics: Phylogenetic Relationships of Scinaxini (Anura: Hylidae: Hylinae). *South Am. J. Herpetol.* **2023**, *27*, 1–145, doi:10.2994/SAJH-D-22-00038.1.

40. Araujo-Vieira, K.; Pombal, J.; Caramaschi, U.; Novaes-e-Fagundes, G.; Orrico, V.; Faivovich, J. A Neotype for *Hyla x-signata* Spix, 1824 (Amphibia, Anura, Hylidae). *Pap. Avulsos Zool.* **2020**, *60*, e20206056--e20206056, doi:10.11606/1807-0205/2020.60.56.

41. Guarnizo, C.; Paz, A.; Muñoz-Ortiz, A.; Flechas, S.; Méndez-Narváez, J.; Crawford, A. DNA Barcoding Survey of Anurans across the Eastern Cordillera of Colombia and the Impact of the Andes on Cryptic Diversity. *PLoS One* **2015**, *10*, doi::10.1371/journal.pone.0127312.

42. Rueda-Almonacid, J. Situación Actual y Probleatica Generada Por La Introducción de "Rana Toro" a Colombia. *Rev. la Acad. Colomb. Ciencias Exactas, Físicas y Nat.* **1999**, *23*, 367–393.

43. Corporación Autónoma Regional de Cundinamarca CAR Plan de Prevención, Control y Manejo (PPCM) de Rana Toro (Rana Catesbeiana) En La Jurisdicción CAR 2018, 41.

44. IUCN The IUCN Red List of Threatened Species Available online: https://www.iucnredlist.org/ (accessed on 24 October 2024).

45. Enciso-Calle, M.; Viuche-Lozano, A.; Anganoy-Criollo, M.; Bernal, M. Rediscovery of *Atelopus subornatus* Werner, 1899 (Anura: Bufonidae), with a Redescription of the Tadpole. *Zootaxa* **2017**, *4344*, 160–162, doi:10.11646/zootaxa.4344.1.7.

46. Méndez-Narváez, J. Diversidad de Anfibios y Reptiles En Hábitats Altoandinos y Paramunos de La Cuenca Del Río Fúquene, Cundinamarca, Colombia. *Biota Colomb.* **2014**, *14*, 94–103.

47.   Roach, N.; Urbina-Cardona, N.; Lacher, T. Land Cover Drives Amphibian Diversity
across Steep Elevational Gradients in an Isolated Neotropical Mountain Range:
Implications for Community Conservation. *Glob. Ecol. Conserv.* **2020**, *22*, e00968,
doi:10.1016/j.gecco.2020.e00968.

48.   Cortés-Gómez, A.; Castro-Herrera, F.; Urbina-Cardona, J. Small Changes in
Vegetation Structure Create Great Changes in Amphibian Ensembles in the Colombian
Pacific Rainforest. *Trop. Conserv. Sci.* **2013**, *6*, 749–769,
doi:10.1177/194008291300600604.

49.   Guayasamin, J.; Cisneros-Heredia, D.; McDiarmid, R.; Peña, P.; Hutter, C. Glassfrogs
of Ecuador: Diversity, Evolution, and Conservation. *Diversity* **2020**, *12*,
doi:10.3390/D12060222.

50.   Snyder, R. Bipedal Locomotion of the Lizard *Basiliscus basiliscus*. *Copeia* **1949**,
1949, 129, doi:10.2307/1438487.

51.   Palacios, C.P.; Agüero, B.; Simonetti, J.A. Agroforestry Systems as Habitat for
Herpetofauna: Is There Supporting Evidence? *Agrofor. Syst.* **2013**, *87*, 517–523,
doi:10.1007/s10457-012-9571-z.

52.   Wanger, T.; Iskandar, D.; Motzke, I.; Brook, B.; Sodhi, N.; Clough, Y.; Tscharntke, T.
Effects of Land-Use Change on Community Composition of Tropical Amphibians and
Reptiles in Sulawesi, Indonesia. *Conserv. Biol.* **2010**, *24*, 795–802,
doi:10.1111/j.1523-1739.2009.01434.x.

53.   Herrera-Montes, A.; Brokaw, N. Conservation Value of Tropical Secondary Forest: A
Herpetofaunal Perspective. *Biol. Conserv.* **2010**, *143*, 1414–1422,
doi:10.1016/j.biocon.2010.03.016.

54.   Flynn, D.; Gogol-Prokurat, M.; Nogeire, T.; Molinari, N.; Richers, B.; Lin, B.;
Simpson, N.; Mayfield, M.; DeClerck, F. Loss of Functional Diversity under Land Use
Intensification across Multiple Taxa. *Ecol. Lett.* **2009**, *12*, 22–33, doi:10.1111/j.1461-
0248.2008.01255.x.

55.   Biswas, S.; Mallik, A. Species Diversity and Functional Diversity Relationship Varies
with Disturbance Intensity. *Ecosphere* **2011**, *2*, 1–10, doi:10.1890/ES10-00206.1.

56.   Gagic, V.; Bartomeus, I.; Jonsson, T.; Taylor, A.; Winqvist, C.; Fischer, C.; Slade, E.;

Steffan-Dewenter, I.; Emmerson, M.; Potts, S.; et al. Functional Identity and Diversity
of Animals Predict Ecosystem Functioning Better than Species-Based Indices. *Proc. R.*
Soc. B Biol. Sci. **2015**, *282*, doi:10.1098/rspb.2014.2620.

57. Suazo-Ortuño, I.; Benítez-Malvido, J.; Marroquín-Páramo, J.; Soto, Y.; Siliceo, H.;
Alvarado-Díaz, J. Resilience and Vulnerability of Herpetofaunal Functional Groups to
Natural and Human Disturbances in a Tropical Dry Forest. *For. Ecol. Manage.* **2018**,
426, 145–157, doi:10.1016/j.foreco.2017.09.041.

58. María Galindo-Uribe, D.; Mario Hoyos-Hoyos, J.; Isaacs-Cubides, P.; Corral-Gómez,
N.; Urbina-Cardona, N. Classification and Sensitivity of Taxonomic and Functional
Diversity Indices of Anurans in the Andean Coffee Cultural Landscape. *Ecol. Indic.*
2022, *136*, doi:10.1016/j.ecolind.2022.108650.

59. Vanegas-Cubillos, M.; Sylvester, J.; Villarino, E.; Pérez-Marulanda, L.; Ganzenmüller,
R.; Löhr, K.; Bonatti, M.; Castro-Nunez, A. Forest Cover Changes and Public Policy:
A Literature Review for Post-Conflict Colombia. *Land use policy* **2022**, *114*,
doi:10.1016/j.landusepol.2022.105981.

60. Armenteras, D.; Rodríguez, N.; Retana, J.; Morales, M. Understanding Deforestation
in Montane and Lowland Forests of the Colombian Andes. *Reg. Environ. Chang.*
2011, *11*, 693–705, doi:10.1007/s10113-010-0200-y.

61. Rodríguez Eraso, N.; Armenteras-Pascual, D.; Alumbreros, J.R. Land Use and Land
Cover Change in the Colombian Andes: Dynamics and Future Scenarios. *J. Land Use*
Sci. **2013**, *8*, 154–174, doi:10.1080/1747423X.2011.650228.

62. Valencia-Aguilar, A.; Cortés-Gómez, A.; Ruiz-Agudelo, C. Ecosystem Services
Provided by Amphibians and Reptiles in Neotropical Ecosystems. *Int. J. Biodivers.*
Sci. Ecosyst. Serv. Manag. **2013**, *9*, 257–272, doi:10.1080/21513732.2013.821168.

63. Agudelo-Hz, W.; Urbina-Cardona, N.; Armenteras-Pascual, D. Critical Shifts on
Spatial Traits and the Risk of Extinction of Andean Anurans: An Assessment of the
Combined Effects of Climate and Land-Use Change in Colombia. *Perspect. Ecol.*
Conserv. **2019**, *17*, 206–219, doi:10.1016/j.pecon.2019.11.002.

64. Forero-Medina, G.; Joppa, L.; Pimm, S. Constraints to Species' Elevational Range
Shifts as Climate Changes. *Conserv. Biol.* **2011**, *25*, 163–171, doi:10.1111/j.1523-

1739.2010.01572.x.