Defective transcription of AAGAG satellite DNA causes sex-ratio meiotic drive in

Drosophila

Authors: Tomohiro Kumon[1,2], Mami Nakamizo-Dojo[1], Amelie A Raz[1], Romain Lannes[1,2], Jaclyn

M Fingerhut[1,2], Yukiko M Yamashita[1,2,3,*]

Affiliations: [1]Whitehead Institute for Biomedical Research, Cambridge, MA02142, USA

[2]Howard Hughes Medical Institute, Cambridge, MA02142, USA

[3]Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

* Email: yukikomy@wi.mit.edu

10
11

Abstract:

Male germ cells have complex transcriptomes, with a large fraction of the genome being

transcribed. This includes protein-coding genes (often not translated), non-coding DNA, and

repetitive DNA, such as transposons and satellite DNA, which are normally silenced as

heterochromatin. The significance of such widespread transcription remains unknown. Here, we

show that a heterochromatin protein, HP2, is required for the transcription of AAGAG satellite

DNA in *Drosophila* spermatocytes. HP2 depletion leads to abnormal retention of

heterochromatin histone marks (H3K9me3) and spermatid death during sperm DNA packaging,

leading to a model that transcription of AAGAG satellite DNA facilitates the remodeling of its

heterochromatic nature in preparation for sperm DNA packaging. Strikingly, the severity of the

spermatid death correlates with the amount of AAGAG satellite DNA carried by the spermatids,

leading to preferential death of Y chromosome-containing spermatids over X-containing

spermatids, and hence sex-ratio meiotic drive phenotype. We propose that widespread

spermatocyte transcription may reflect the process of chromatin remodeling to allow sperm DNA

packaging. We further propose that differential composition and amount of satellite DNA on

chromosomes may underlie naturally occurring male meiotic drive.

Introduction:

In a broad range of organisms from *Drosophila melanogaster* to *Homo sapiens*, testis is known to have a high-complexity transcriptome, transcribing a large fraction of the genome. This includes many protein-coding genes (without necessarily being translated), non-coding DNAs, as well as repetitive sequences such as transposons and satellite DNA that are normally silenced as heterochromatin[1-8]. A few hypotheses have been proposed to explain such widespread transcription. For example, one model proposed that transcription-coupled DNA repair in male germ cells serves to lower mutation rates of protein-coding genes[9](but see also [10]), but it does not explain the significance of transcription of non-coding and repetitive sequences. Others proposed that non-coding transcripts function as lncRNA of yet unknown function during spermatogenesis, but many of them showed no phenotype upon deletion[2]. Yet another model proposed that transcription is a by-product of generally open chromatin in testis [5]. Accordingly, there is no consensus model that explains widespread transcription of the testis to date.

Here using *Drosophila* spermatogenesis as a model (Fig. 1a), we investigate the role of satellite DNA transcription. Satellite DNA is non-coding tandem repetitive DNA, and constitutes approximately 30% of the *Drosophila* genome (Fig. 1b)[11-13]. We show that satellite DNA is highly transcribed in spermatocytes along with other transcripts, contributing to widespread transcription. We found that a heterochromatin binding protein, HP2 [14-17], which preferentially binds to AAGAG satellite DNA, is required for its transcription in spermatocytes. HP2 depletion led to subfertility due to defective sperm DNA packaging, associated with aberrant retention of heterochromatin histone mark H3K9me3 and defective incorporation of protamine, the sperm-specific DNA packaging protein[18, 19]. Strikingly, we found that defects in sperm DNA packaging upon HP2 depletion correlate with the abundance of AAGAG satellite DNA contained in the haploid spermatid: spermatids with the Y chromosome, which contains much more AAGAG satellite DNA than the X chromosome, were particularly sensitive to HP2 depletion, leading to skewed sex ratio in progeny. Moreover, spermatids containing a chromosome II variant with a minimal amount of AAGAG satellite DNA were more resistant to HP2 depletion, leading to a partial rescue of the HP2 depletion phenotype. Taken together, we propose that HP2-mediated AAGAG satellite DNA transcription facilitates the opening of AAGAG satellite DNA heterochromatin in preparation for sperm DNA packaging. We speculate that, in more general, widespread transcription broadly observed in male germ cells may reflect the process of chromatin remodeling during sperm DNA packaging. We further propose that the distinct

composition of satellite DNA may have been exploited by meiotic drivers, leading to an

evolutionary arms race between meiotic drivers and satellite DNA.

Results:

Satellite DNA is transcribed in *Drosophila* spermatocytes

We comprehensively analyzed the transcription in the *Drosophila* testis, utilizing multiple

available data sets, and established the nature of widespread transcription. *Drosophila* genome

is composed of ~20% of protein-coding genes, ~30% of satellite DNA, ~20% of transposons,

and ~30% of other non-coding/non-repetitive DNA (Fig. 1b). Previous single-nucleus and single-

cell RNA-seq data from *Drosophila* testis showed the widespread transcription of

polyadenylated transcripts in the spermatocytes[1]. Reanalysis of these data[1] further revealed

that 69% of protein-coding genes are expressed in the testis (Fig. 1c), which is much higher

than in somatic tissues[20]. Comparison of transcriptome[1] and proteome[21] further revealed that

more than half of transcribed protein-coding genes (i.e., 42.1% of total protein-coding genes)

are not translated to proteins (Fig. 1c). lncRNAs and transposons were more abundantly

transcribed in spermatocytes than in the somatic epithelia of the testis (Fig. 1d, e).

Moreover, RNA fluorescence *in situ* hybridization (FISH) revealed marked upregulation

of various satellite DNA transcription in spermatocytes, which cannot be captured by RNA

sequencing that only identified polyadenylated RNAs[1] (Fig. 1f, Supplementary Fig. 1). Most of

satellite DNA found in *Drosophila melanogaster* genome[13, 22-24] were found to be transcribed in

spermatocytes, with many of them exhibiting the signal from both strands (Fig. 1f,

Supplementary Fig. 1). Although a known source of satellite transcripts in the spermatocytes is

satellite DNA embedded in the gigantic introns of several Y-linked genes [24-27], the large amount

of satellite transcripts remained in the spermatocytes from X/O males that lack the Y

chromosome (Fig. 1g, Supplementary Fig. 1), suggesting that satellite DNA from the X

chromosome and autosomes are also transcribed. Taken together, these results establish the

widespread nature of spermatocyte transcription in *Drosophila* spermatocytes, including protein-

coding genes, non-coding RNA, and satellite DNA.

HP2 is required for the transcription of AAGAG satellite DNA in spermatocytes

We found that heterochromatin protein 2 (*Su(var)2-HP2*, hereafter *HP2*), an HP1-

interacting protein[14, 17], is specifically required for the transcription of AAGAG satellite DNA in

the spermatocytes. Depletion of HP2 from late spermatogonia/ spermatocytes using *bam-GAL4*

driver (*bam-GAL4>UAS-HP2^HMS01699*, hereafter *bam>HP2^RNAi*) led to marked reduction of

transcripts from AAGAG satellite DNA in spermatocytes: RNAs from both strands of AAGAG

satellite DNA (i.e., AAGAG RNA and CUCUU RNA) were reduced in *bam>HP2^RNAi*, with a more

severe reduction of AAGAG RNA than CUCUU RNA (Fig. 2a, b). Expression of RNAi-resistant

HP2 construct rescued transcription of AAGAG satellite DNA (both strands) in *bam>HP2^RNAi*

(Supplementary Fig. S2), demonstrating that reduction in AAGAG/CUCUU RNA is indeed

caused by HP2 depletion. *bam>HP2^RNAi* had little effect on the transcription of other satellite

DNA (Supplementary Fig. S3), suggesting that HP2 regulates satellite DNA transcription in a

sequence-specific manner. These results demonstrate that HP2 is required for AAGAG/CUCUU

transcription in spermatocytes.

Analysis of HP2 ChIP-seq data on the modENCODE database[28] showed that HP2 binds

preferentially to AAGAG satellite DNA (Fig. 2c), implying that HP2 may regulate the transcription

of AAGAG satellite DNA via direct binding. Moreover, we found that HP2's localization to

chromatin is dependent on AAGAG satellite DNA (Supplementary Fig. S4). However, given that

HP2 expression decreased at the time of strongest AAGAG/CUCUU transcription in

spermatocytes (Supplementary Fig. S4a), it is possible that HP2's role in AAGAG/CUCUU

transcription can be indirect. If it is direct, HP2's requirement in AAGAG/CUCUU transcription

may represent an example of heterochromatin-dependent transcription[29, 30].

It was previously shown that AAGAG RNA can be depleted by RNAi (*bam-GAL4>UAS-*

AAGAG^shRNA, hereafter *bam>AAGAG^RNAi*)[3, 31]. Although the mechanism of nuclear RNA

degradation by RNAi is not fully established, *bam>AAGAG^RNAi* was previously used to eliminate

AAGAG RNAi from the spermatocytes[3]. We confirmed that *bam>AAGAG^RNAi* specifically

depleted AAGAG RNA without affecting the opposite strand, CUCUU RNA (Fig. 2a, b) as

previously shown[3]. These results suggest that *bam>HP2^RNAi* interferes with the transcription of

AAGAG satellite DNA, whereas *bam>AAGAG^RNAi* cleaves AAGAG RNA after it is transcribed.

Hereafter, we analyzed both *bam>HP2^RNAi* and *bam>AAGAG^RNAi* because their comparison

provided insights into the function of AAGAG transcription vs. AAGAG transcript. In conclusion,

HP2 is required for the transcription of AAGAG satellite DNA.

HP2 depletion leads to subfertility due to sperm DNA packaging defects

We found that *bam>HP2^RNAi* led to a reduction in fertility, whereas *bam>AAGAG^RNAi* led

to complete sterility, as shown previously[3] (Fig. 3a). The majority of *bam>HP2^RNAi* animals still

had mature sperm in their seminal vesicles, corresponding to their residual fertility, whereas

bam>AAGAG^RNAi animals had no sperm in their seminal vesicles (Fig. 3b). Based on the sterility

caused by *bam>AAGAG^RNAi*, a previous study proposed that AAGAG RNA plays an important

role in the process of spermatogenesis. They reported that *bam>AAGAG^RNAi* led to late

spermatogenesis phenotypes, such as nuclear morphology defects and sperm bundling defects

[3]. They further observed that *bam>AAGAG^RNAi* led to defective protamine incorporation into

sperm nuclei, a critical process for sperm DNA packaging. However, we found that protamine

was mostly normally incorporated in *bam>AAGAG^RNAi* testis (Supplementary Fig. S5). We noted

that the disruption of sperm nuclei bundling made it challenging to correctly identify the

differentiation stages of sperm nuclei, potentially leading to the interpretation of defective

protamine incorporation in the previous study[3]. RT-qPCR showed that *bam>AAGAG^RNAi* led to

marked downregulation of a subset of Y-linked fertility genes (e.g., *ORY* and *kl-2*) (Fig. 3c). *kl-2*

encodes a subunit of axonemal dynein, whose depletion is known to lead to the disruption of

sperm nuclei bundling[27], similar to the phenotype of *bam>AAGAG^RNAi*. *ORY* and *kl-2* genes are

known to contain large stretches of AAGAG satellite DNA in their gigantic introns

(Supplementary Fig. S6)[24, 26]. RT-qPCR further revealed a striking reduction of *ORY* transcripts

across AAGAG-containing gigantic introns in *bam>AAGAG^RNAi* animals (Fig. 3d). RNA FISH

further confirmed the reduction of *ORY* mRNA in spermatocytes in *bam>AAGAG^RNAi* animals

(Figure 3e). Moreover, *Pzl*, an autosomal gene that contains gigantic introns with assembly

gaps[32] that likely contain AAGAG satellite DNA (Supplementary Fig. S6), was also

downregulated in *bam>AAGAG^RNAi* (Fig. 3c). These results suggest that *bam>AAGAG^RNAi* leads

to cleavage of AAGAG-containing RNA. Because *ORY* and *kl-2* are required for male fertility[33-

35], these results suggest that the sterility of *bam>AAGAG^RNAi* is caused by RNAi-mediated

knockdown of these genes that contain AAGAG in their introns.

In contrast to *bam>AAGAG^RNAi*, *bam>HP2^RNAi* did not affect the expression of genes that

contain gigantic introns with AAGAG satellite DNA (Fig. 3c, d, e), suggesting that *bam>HP2^RNAi*

impacts spermatogenesis through a mechanism distinct from *bam>AAGAG^RNAi*. We found that

bam>HP2^RNAi led to sperm DNA packaging defects, explaining their subfertility. Although germ

cell development appeared mostly normal, leading to normal meiosis (Supplementary Fig. S7),

bam>HP2^RNAi led to unique cytological phenotypes during spermatid differentiation stage. In wild

type animals, post-meiotic germ cells undergo stereotypical morphological changes, leading to

the formation of mature sperm: a cyst of 64 spermatids, resulting from four mitotic and two

meiotic divisions, develop in synchrony to undergo sperm DNA packaging and axoneme

elongation (Fig. 1a). Sperm DNA is packaged by protamine, sperm-specific DNA packaging

proteins that replace histone-based chromatin, leading to highly compacted DNA (Fig. 4a)[18, 19].

We found that *bam>HP2^RNAi* led to defective sperm DNA compaction during the last stages of

sperm development (Fig. 4b): In some cases (Fig. 4b, ii), a subset of spermatid nuclei within a

cyst lost their typical 'needle' morphology (Fig. 4b, i) and become completely rounded-up, while

other nuclei exhibited normal needle morphology. In other cases, all spermatids within a cyst

(Fig. 4b, iii) exhibited defective sperm nuclear morphology. Expression of RNAi-resistant HP2

rescued sperm DNA compaction defect in *bam>HP2^RNAi* (Fig. 4b, iv), demonstrating that sperm

DNA compaction defect is indeed due to HP2 depletion.

Although the morphology of earlier "canoe" stage spermatids appeared mostly normal in

bam>HP2^RNAi animals, we found that the defects began in these earlier stages. Canoe stage

spermatid is characterized by the removal of histone and incorporation of protamine, such as

Mst77F[13]. In control animals, sperm DNA compaction is accompanied by sequential events of

removing heterochromatin mark H3K9me3, removing histone H3.3 (Supplementary Fig. S8a),

then incorporating protamine such as Mst77F (Fig. 4c, control). However, in *bam>HP2^RNAi*

animals, a subset of spermatids exhibited a failure in the removal of H3K9me3 and deposition of

Mst77F, whereas other nuclei in the same cyst have removed H3K9me3 and deposited Mst77F

(Fig. 4c, *bam>HP2^RNAi*). In wild type, protamine incorporation occurs in synchrony within the

cyst, and all nuclei in a cyst are either Mst77F-positive or -negative (Fig. 4c, control). In contrast,

the spermatids cyst from *bam>HP2^RNAi* animals contained a mixture of Mst77F-positive and -

negative nuclei, demonstrating that a subset of nuclei fails to incorporate Mst77F in a timely

manner. Moreover, when the cyst progressed to the needle spermatid stage, the defective

nuclei in *bam>HP2^RNAi* that failed to compact their DNA still retained H3K9me3 (Fig. 4d). Similar

to the retention of H3K9me3, H3.3 histone was also retained in the defective spermatids in

bam>HP2^RNAi, whereas another histone mark, AcH4, did not exhibit detectable difference

between control and *bam>HP2^RNAi* (Supplementary Fig. S8b). Taken together, these results

suggest that retention of H3K9me3 is associated with the failure in protamine incorporation,

which in turn leads to ultimate failure in sperm DNA compaction, in *bam>HP2^RNAi*.

HP2 depletion preferentially harms Y-chromosome-containing spermatids, leading to sex

ratio meiotic drive

Unexpectedly, we found that *bam>HP2^RNAi* preferentially impacts sperm DNA

compaction of Y-chromosome-bearing spermatids, leading to a sex ratio meiotic drive

phenotype. While counting the number of offspring for the fertility assay (Fig. 3a), we noticed

that *bam>HP2^{RNAi}* males produce significantly more daughters than sons (Fig. 5a), suggesting

that sperm DNA compaction defects may be biased toward Y chromosome-containing

spermatids. Indeed, by DNA FISH using X- and Y-specific probes, we found that defective

nuclei in *bam>HP2^{RNAi}* predominantly contained the Y chromosome (Fig. 5b), demonstrating

that the Y chromosome is more sensitive to HP2 depletion than the X chromosome. Seeking for

the difference(s) between X and Y chromosomes that may explain Y-biased sensitivity to HP2

depletion, we realized that the Y chromosome contains much more AAGAG satellite DNA than

the X chromosome[13, 22] (Fig. 5c). Based on this correlation between the amount of AAGAG on

the chromosome and the tendency to fail in sperm DNA compaction, we hypothesized that HP2-

dependent transcription of AAGAG satellite DNA prepares for protamine incorporation by

remodeling heterochromatin state of AAGAG satellite DNA (Fig. 5d). Accordingly, the

dependence of a chromosome on HP2 to remodel their AAGAG satellite DNA heterochromatin

correlates with the amount of AAGAG satellite DNA on it. This can explain why X chromosome-

containing spermatids are less impacted by HP2 depletion, because the X chromosome has

less AAGAG satellite DNA. Based on these results, we propose that satellite DNA transcription,

instead of its RNA product, is important to prepare for histone-to-protamine transition.

The amount of AAGAG satellite DNA determines sperm DNA compaction defects caused

by HP2 depletion.

We found that the reduction in AAGAG satellite DNA abundance can rescue sperm DNA

compaction defect observed in *bam>HP2^{RNAi}*, further supporting our hypothesis that the

remodeling of satellite DNA via transcription is critical for sperm DNA packaging (Fig. 5d). In

addition to the Y chromosome, chromosome II contains a considerable amount of AAGAG

satellite DNA in its pericentromeric heterochromatin (Fig. 5c)[13, 22], likely contributing to spermatid

DNA compaction defects in *bam>HP2^{RNAi}*. This may also explain why sometimes all spermatids

within a cyst fail to undergo sperm DNA compaction defects (Fig. 4b, iii), because all spermatids

contain a copy of chromosome II.

Consistent with this notion, we found that natural chromosome II variants with minimal

AAGAG satellite DNA (hereafter II^{ΔAAGAG}, 'Ithaca 06' and 'Ithaca16' strains)[36] (Supplementary

Fig. S9) can rescue sperm DNA compaction defects caused by the depletion of *HP2*. When this

chromosome II^{ΔAAGAG} was placed in the background of *bam>HP2^{RNAi}* (*y^+w^+/Y; UAS-HP2^{HMS01699}/*

ΔAAGAG; bam-GAL4/+, hereafter X/Y; II^+/II^{ΔAAGAG} *bam>HP2^{RNAi}*), we observed a considerable

reduction in the frequency of defective spermatids that failed sperm DNA compaction (Fig. 6a,

b), suggesting that the abundance of AAGAG satellite DNA correlates with the severity of sperm

DNA compaction defects due to HP2 depletion.

To further test our hypothesis that AAGAG satellite DNA amount determines the

sensitivity of spermatids to HP2 depletion, we examined whether spermatids containing $II^{\Delta AAGAG}$

survived better than those with II^+ in *bam>HP2^{RNAi}*. Specifically, we analyzed the genotype of

the offspring of X/Y; $II^+/II^{\Delta AAGAG}$ *bam>HP2^{RNAi}* males. From diploid X/Y; $II^+/II^{\Delta AAGAG}$

spermatocytes, there are four possible genotypes of the haploid sperm: X II^+, X $II^{\Delta AAGAG}$, Y II^+,

and Y $II^{\Delta AAGAG}$ (Fig. 6c). Based on the offspring genotype, we deduced the genotype of survived

sperm. In male offspring, where II^+ and $II^{\Delta AAGAG}$ can easily be scored based on the body color

(Method), we observed a small but significant overrepresentation of $II^{\Delta AAGAG}$ over II^+ (Fig. 6c),

suggesting that spermatid containing Y $II^{\Delta AAGAG}$ indeed survived better than Y II^+. This implies

that AAGAG satellite DNA on chromosome II worsens sperm DNA compaction defects in Y-

containing spermatids.

In female offspring, II^+ and $II^{\Delta AAGAG}$ cannot be easily distinguished by visible phenotypic

markers. Thus, a randomly-chosen subset of female offspring was subjected to a PCR-based

assay to distinguish between II^+ and $II^{\Delta AAGAG}$ (Method), which showed no significant difference in

the survival of X II^+ and X $II^{\Delta AAGAG}$ (Fig. 6c). This is not surprising considering that *bam>HP2^{RNAi}*

males with a regular amount of AAGAG (II^+) sired female-biased offspring (Fig. 5a), suggesting

that X-containing spermatids can tolerate the amount of AAGAG on the second chromosome.

These results may reflect the dose-sensitive nature, where HP2-depleted cells can tolerate only

a certain amount of AAGAG satellite DNA (see Discussion): X-containing spermatids may be

able to tolerate AAGAG satellite DNA on the chromosome II, because they contain only a small

amount of AAGAG satellite DNA on the X chromosome. In contrast, Y-containing spermatids,

which carries a large amount of AAGAG satellite DNA, cannot tolerate AAGAG satellite DNA on

the chromosome II under HP2 depletion. Accordingly, sperm DNA compaction defects of Y-

containing spermatids in *bam>HP2^{RNAi}* can be rescued by reducing the dose of AAGAG satellite

DNA on the chromosome II.

These results suggest that the abundance of AAGAG satellite DNA is a determinant that

renders spermatid sensitive to the loss of HP2. The fact that the reduction of AAGAG satellite

DNA markedly rescues the phenotype of *bam>HP2^{RNAi}* implies that AAGAG satellite DNA is the

major target of HP2, although HP2 may also regulate other heterochromatic loci as well.

Additionally, the rescue of the phenotype by reducing the amount of AAGAG satellite DNA

implies that AAGAG RNA is not essential for spermatid development. Taken together, we

propose that HP2 promotes transcription of AAGAG satellite DNA in spermatocytes, facilitating

the remodeling of heterochromatin in preparation of germline DNA for histone-to-protamine

transition.

Discussion:

Through the study of satellite DNA transcription in *Drosophila* spermatocytes, we

propose that the widespread transcription in male germline reflects the process of chromatin

remodeling in preparation for histone-to-protamine transition. According to this model,

transcription, but not transcripts, is important for male germ cell development. Although it is

challenging to distinguish between the function of transcription to remodel the chromatin and

that to generate functional transcripts, several lines of evidence favor the model that

transcription, but not resultant transcripts, is important. First, the expression of AAGAG did not

rescue the *bam>HP2^{RNAi}* phenotype (Supplementary Figure S10), suggesting that the lack of

AAGAG transcript is not the cause of the *bam>HP2^{RNAi}* phenotype. Furthermore, the rescue of

the *bam>HP2^{RNAi}* phenotype by the introduction of chromosome II with little AAGAG satellite

DNA (II$^{\Delta AAGAG}$) (Fig. 6) provides critical support for this notion. If AAGAG RNA has an important

function for spermatid development, it would be expected that deletion of AAGAG satellite DNA

in II$^{\Delta AAGAG}$ worsens the phenotype of *bam>HP2^{RNAi}* because both conditions reduce AAGAG

RNA. Instead, II$^{\Delta AAGAG}$ rescued *bam>HP2^{RNAi}* phenotype, implying that AAGAG RNA is

dispensable for spermatid development. The role of AAGAG RNA indicated by the previous

study by the use of *AAGAG^{RNAi}* [3] likely reflects the function of fertility genes that contain AAGAG

satellite DNA in their introns. Instead of AAGAG RNA serving a function, we propose that

AAGAG RNA is produced as a consequence of transcription-dependent chromatin remodeling

of heterochromatin, which is required for subsequent sperm DNA packaging by protamine. In

this model, satellite DNA's heterochromatic nature is a "burden" for spermatid development,

impeding protamine-mediated sperm DNA compaction process if left unopened.

It is important to note that defective spermatid development in *HP2^{RNAi}* is sensitive to the

dose of total AAGAG carried in the haploid spermatids. We speculate that each spermatid may

have a threshold of the unremodeled chromatin it can tolerate (e.g. the amount of unremoved

histones, and/or the amount of the chromatin that is not packaged by protamine), and the

spermatids that have an above-threshold unremodeled chromatin may be fated for death.

Indeed, the presence of a checkpoint mechanism that monitors the quality of sperm DNA has

been proposed to explain the cellular phenotype of sperm-killing meiotic drivers, such as *SD* [37,]

[38]. Interestingly, the cytological phenotype of *HP2^RNAi^* is reminiscent of the *SD* phenotype. We

speculate that the AAGAG dose-sensitive nature of the *HP2^RNAi^* phenotype may be due to such

a checkpoint mechanism.

HP2-mediated transcription of AAGAG satellite DNA likely represents just one example

of many transcription regulators in the testis that reorganize chromatin for protamine

incorporation. Among many heterochromatic satellite DNA blocks in *Drosophila* genome, HP2

appears to regulate mainly the transcription of AAGAG, whereas other satellite DNA such as

AACATAGAAT, AATAT, AACAC, and AAGAC were not detectably impacted. Therefore, we

speculate that there are other factors that serve the same function as HP2 but for different

satellite DNA. By extending this model, we further propose that the widespread transcription of

the testis, encompassing various genomic elements such as protein-coding (but not translated)

genes and non-coding sequences, is the result of the process that remodels germ cell

chromatin to allow histone-to-protamine transition. We predict that a panel of transcription

factors together contribute to transcription of a large fraction of the genome in male germline.

Discovery of such factors awaits future investigations.

Importantly, our study unexpectedly revealed a condition that caused sex-ratio meiotic

drive. Because the Y chromosome contains more AAGAG satellite DNA, it was more sensitive

to HP2 depletion than the X chromosome, causing HP2-depleted males to sire female-biased

progeny. Our results may provide insights into the molecular mechanisms of evolution of

satellite DNA and their binding proteins, eventually contributing to incompatibility between

species. For example, if a chromosome that happens to have fewer copies of certain satellite

DNA compared to other alleles in the population, such a chromosome can become a selfish

meiotic driver by acquiring a mutation (e.g., dominant negative, or loss of function if haplo-

insufficient) in the gene(s) that mediate transcription/remodeling of the target satellite DNA

(Supplementary Fig. S11). Subsequently, the targeted chromosome may evolve to reduce the

copy number of the satellite DNA, or to change the satellite DNA sequence that can be

transcribed by other transcription factors to avoid being targeted (Supplementary Fig. 9).

Repeated cycles of changing satellite DNA sequence/amount and transcription

336 factors/chromatin remodelers may eventually result in incompatibility between satellite DNA and

337 their regulators, leading to speciation.

339   There are natural variations in satellite DNA abundance at a megabase scale [23], which

340 may lead to biased inheritance of chromosome variants, leading to meiotic drive if left

341 unremodeled in spermatocytes. Accordingly, satellite DNA transcription may serve as a

342 mechanism to prevent male meiotic drive. Interestingly, the well-established meiotic driver in *D.*

343 *melanogaster*, *Segregation Distorter* (*SD)*, involves satellite DNA (*Rsp*) and a RanGAP

344 mutation, where the chromosome II containing a higher amount of *Rsp* satellite DNA is

345 specifically killed with the very same cytological phenotype (sperm DNA compaction defects

346 described in this study)[39-41]. It is tempting to speculate that a RanGAP mutation compromises

347 the nuclear import of proteins responsible for the remodeling of *Rsp* satellite DNA, making the

348 chromosome II with a high amount of *Rsp* satellite DNA fail in sperm DNA packaging. In

349 addition, many of *Drosophila* "hybrid incompatibility loci" that render hybrids lethal or sterile

350 contain satellite DNA (*Zhr*) and encode satellite DNA/heterochromatin-binding proteins (*Hmr,*

351 *Lhr/HP3*, and *OdsH*)[42-46]. Moreover, rapid changes in the binding specificity of heterochromatin

352 proteins have been documented[46]. The present study may provide a hint as to how satellite

353 DNA sequence and its binding proteins may engage in an evolutionary arms race.

355   In summary, the present study provides a model to explain the significance of

356 widespread transcription broadly observed in male germline, where it serves to remodel the

357 chromatin of male germline to make it compatible with histone-to-protamine transition. Such

358 requirement for transcription-mediated chromatin remodeling for sperm DNA packaging may

359 provide a potential link between rapid evolution of satellite DNA and male meiotic drive.

361 **Methods:**

362 *Fly Husbandry*

363 All experimental flies were raised on modified Bloomington *Drosophila* Stock Center (BDSC)

364 cornmeal food (agar 0.65%, cornmeal 6.71%, inactivated yeast 1.59%, soy flour 0.92%, corn

365 syrup 7.0%, tegosept 0.15%; without propionic acid; less dry time) at 25°C, and young flies (1-

366 to 3-day-old adults) were used for all experiments. All fly stocks were maintained at room

367 temperature while not in use for experiments, and stocks that exhibited spontaneous spermatid

368 DNA compaction errors were replaced with new stocks. Control flies were either the sibling from

369 the same genetic cross or the parental *GAL4* stock. The following fly stocks were used: y^1w^1

(BDSC 1495), *nos-GAL4:VP16[47]*, *bam-GAL4:VP16[48]*, *Ubi-GFPS65C:α1-tub84B[49]*, *H3.3A-*

Dendra2 [50], *C(1)RM/C(1;Y)6, y[1]w[1]f[1]/O* (BDSC 9460; males from this stock were crossed with

y[1]w[1] females to generate X/O males), *UAS-TRiP.HMS01699* (BDSC 38255; *UAS-HP2[RNAi]*),

UAS-AAGAG[shRNA] (gift from Gary Karpen[3]), *Su(var)2-HP2-GFP.FPTB* (BDSC 68184; *hp2-gfp*),

UAS-HP2-S (this study), *UAS-HP2-L* (this study), Ithaca I16 and Beijing B52 of Global Diversity

lines (gift from Andrew Clark[36]). Combinations of transgenes were generated using the CyO;

TM6B double balancer strain.

Immunofluorescence (IF) Staining

Testes were dissected in 1X PBS, and fixed for 30 minutes at room temperature (RT) in 1mL

fixative (1X PBS (Invitrogen AM9624; 4mL of 10X PBS), 4% formaldehyde (ThermoFisher

Scientific 28908; 10mL of 16% solution), 0.1% Triton X-100 (Millipore Sigma T9284; 400µL of

10% solution) in H_2O (25.6mL); stored at −20˚C). Testes were rinsed with 1mL PBST (1X PBS

(Invitrogen AM9624; 5mL of 10X PBS), 0.1% Triton X-100 (Millipore Sigma T9284; 500µL of

10% solution) in H_2O (44.5mL); stored at 4˚C), permeabilized for at least 3 hours at RT in 1mL

PBST, blocked for 1 hour at RT in 200µL blocking solution (1X PBS (Invitrogen AM9624; 5mL of

10X PBS), 3% BSA (Fisher Scientific BP1605-100; 1.5g in 25mL H_2O), 0.1% Triton X-100

(Millipore Sigma T9284; 500µL of 10% solution) in H_2O (up to 50mL); stored at −20˚C), and

incubated overnight at 4°C in 200µL blocking solution with primary antibodies. Samples were

rinsed with 1mL PBST, washed for at least 1 hour at RT in 1mL PBST, incubated overnight at

4°C in 200µL blocking solution with secondary antibodies, rinsed and washed as above.

Samples were mounted in VECTASHIELD with DAPI (Vector Laboratories H-1200). Images

were acquired using Leica Stellaris8 confocal microscope with a 63X oil immersion objective

lens (NA = 1.4) and processed/analyzed using ImageJ software. The primary antibodies used

were anti-Mst77F (1:1000 dilution; guinea pig[51]) and anti-H3K9me3 (1:200 dilution; rabbit;

Abcam ab8898). The secondary antibodies used were anti-rabbit IgG conjugated with Alexa

Fluor 647 (1:200 dilution; goat; Abcam ab150079) and anti-guinea pig IgG conjugated with

Alexa Fluor 568 (1:200 dilution; goat; Abcam ab175714).

RNA Fluorescent in situ Hybridization (FISH)

Testes were dissected in 1X PBS, and fixed for 30 minutes at RT in 1mL fixative. Testes were

washed briefly (5 min x 2) at RT in 1mL PBST, rinsed briefly (5 min) at RT with 1mL wash buffer

(2X SSC (Invitrogen AM9770; 5mL of 20X SSC), 10% formamide (Millipore Sigma S4117; 5mL),

0.1% Triton X-100 (Millipore Sigma T9284; 500µL of 10% solution) in H_2O (39.5mL); stored at

4°C), and hybridized overnight at 37°C in 200μL hybridization buffer (10% dextran sulfate

(Millipore Sigma D8906; 1g in 5mL H$_2$O), 1mg/mL yeast tRNA (Millipore Sigma R8759; 1mL of

10mg/mL solution), 0.5% BSA (Invitrogen AM2616; 1mL), 2X SSC (Invitrogen AM9770; 1mL of

20X SSC), 10% formamide (Millipore Sigma S4117; 1mL), 2mM Ribonucleoside Vanadyl

Complex (NEB S1402; 100μL) in H$_2$O (up to 10mL); stored at −20°C). Following hybridization,

samples were washed two times in wash buffer for 30 minutes each at 37°C and mounted in

VECTASHIELD with DAPI (Vector Laboratories H-1200). Images were acquired using Leica

Stellaris8 confocal microscope with a 63X oil immersion objective lens (NA = 1.4) and

processed/analyzed using ImageJ software. RNA FISH signal intensity is semi-quantitative due

to inherent technical (e.g., different tubes) and biological (e.g., testis thickness) variations. RNA

FISH intensities among different genotypes (Figure 2C) were analyzed from images with

roughly the same intensity of internal control (GFP-tubulin). Fluorescently labeled probes were

added to the hybridization buffer to a final concentration of 50nM (1μL of 10μM probe solution in

200μL hybridization buffer). Probes against the repetitive DNA transcripts were from Integrated

DNA Technologies: Cy5-(AAGAG)x6, Cy3-(CTCTT)x6; Cy3-(AATAACATAG)x3, Cy5-

(CTATGTTATT)x3; Cy5-(AATAT)x6, Cy3-(ATATT)x6; Cy3-(AACAC)x6, Cy5-(GTGTT)x6; Cy3-

(AAGAC)x6, Cy5-(GTCTT)x6. *ORY* probes Quasar 670-(TTTTTGGCTTTCTTTCTGTC,

AAAAGTTGAGGCTCCGAGTT, CGTTTAATTCGCGATGCTTC,

CTTCATCTACATACCGACGA, TCCGATGTTGAAGTCAGTTC, CCCCCAACAAATCTTTAAGT,

TGATGCATCTGATTCTTTCC, TCTTGCACTAACTGTTCTCG, CAGTTGTACCACTATTTCGG,

TTAGCTCGAGAAGCCATTTT, ATTCAACGTTTAGGCGTTCG, CCATTGTCTTCATCAAAGCT,

TTTCCATGTGCTTCTTTTTG, CTCTATTCGCAATATCCAGT, AGTTTACGTGTCGTTTCTGA,

GCATTGCCTTATTTAATGCG, GCCTTAACTGCTTTATCATT, AAGCCTGCTCGTTAATTGAG,

TTGTACATTCTTGTTGTCGC, CATCCTTCTCCAGAGAATTA, GACGAATATCATCCGTTCGG,

CACGTTGCAATGTCTCTTTG, ATGTTTAAGTCCGCCATAG, TCTTTCTCTGCTTTGGATTT,

CGTCTTCAGAGAGATTTGGT, TCGTGTAACTTTTCCGTGAG, ACGGACTCCAACTGTTTTTT,

CGCTGAAGCATCATTTTCTC, TTTTACTTCGTTTGCGCTTA, GCCACTAAGTTTTCTTTGTT,

TTTCTCTCATATCCTTACGT, CAATCCGACTAGGTTACGTT, AACAATCTCGTCATTCCGTC,

GGGCATTTTGTGCAATTTGA, AAGGCATTCACTTTCAGTGT, CTCATGTCAGCAGTACTTTT,

TCTTCAGTTAGAGCTCGTAT, CAACGATGTACTCCTGTAGG, AATTTCTTAGGATCTTCGCC,

TATTGACTGCTTTAGGGAGC, AGCCTCACACAGTTTATTTT, GCATATGAGATAGCATCCTT,

TAATGAACGCTCTGCTGCTA, GTCTTGATTTAAGTTCCACC, AATTGACAGCTCTCCGATTT,

TATTTCGTTTTTCCCATCTG, ATCCACATTCCATATACTCT, ATATCCGTTAACTTCGCACA)

Quantification of RNA FISH Signal

To quantify RNA FISH signal (for Fig. 1a, 2b, S2f and S3a), we chose spermatocytes of the

indicated stages and quantified their signal intensity. To minimize artifactual variations in the

signal intensity (such as the depth of the cells within the tissue), images were acquired from

near the surface of the tissue, which showed the brightest signal. Z-stacks (step size: 0.4 μm)

were collected to cover the entire depth of the spermatocyte to be quantified. To quantify the

signals within each spermatocyte, three circular regions per nucleus of 1.2μm in diameter were

chosen, while avoiding the areas that contain the signal from the other cells overlapping with the

cell of interest. Then, the mean intensity of these regions was calculated after subtracting the

cytoplasmic background. These values were then normalized to the mean value of the control.

Analysis of HP2 ChIP seq for satellite DNA binding

HP2 ChIP-seq data were obtained from the SRA (accessions: SRR870205, SRR870204,

SRR870206, SRR870207) (modENCODE ID: 5593)[28]. We analyzed all samples using the k-

seek program, and the resulting data were compiled into a summary table using the k.compile.pl

script [36]. Replicate counts were merged for both ChIP-seq and input samples. Satellites with a

repeat unit length shorter than 4 bp were excluded. We then selected five major satellites of

interest (AAGAG, AACATAGAAT, AATAT, AACAC, AAGAC), while all remaining satellites were

combined into a category labeled "other satellites". For each satellite, enrichment was

calculated as the ratio of the satellite's proportion in the ChIP-seq sample to its proportion in the

input. This ratio served as the enrichment metric for each satellite in HP2. The satellite

proportion in the input was used as a measure of satellite abundance.

Reverse Transcribed Quantitative Polymerase Chain Reaction (RT-qPCR)

Dissected testes from 50 flies were pooled in ice-cold 1mL 1X PBS. Samples were first

homogenized in ice-cold 300μL TRIzol (Invitrogen 15596026), supplemented with 700μL TRIzol,

vortexed vigorously, centrifuged for 5 min at 4˚C. Approximately 1mL of clear supernatant was

transferred to a new tube, and 400μL of phenol:chloroform (Invitrogen AM9722) was added.

Samples were mixed gently by hand for 15 sec, incubated for 3 min at RT, centrifuged for 15

min at 4˚C. Approximately 400μL of the top aqueous phase was transferred to a new tube, and

1μL of GlycoBlue Coprecipitant (Invitrogen AM9515), 1mL of 100% ethanol, and 140μL of 5M

NH$_4$Oac were added. RNA samples were stored at −80˚C until the RT reaction. RNA samples

were centrifuged for 20 min at 4˚C (discard supernatant), washed with 700μL 70% ethanol,

centrifuged for 10 min at 4˚C (discard supernatant), dried for 15 min at RT, and dissolved in

20µL of 95˚C H$_2$O. RNA was denatured for 5 min at 65˚C and put on ice at least for 1 min.

Almost all RNA (20µL, typically 5µg RNA yield) was used for synthesis of cDNA (mixed with

25µL buffer and 5µL SuperScript III (Invitrogen 11752); 25˚C 10min, 50˚C 30min, 85˚C 5min,

4˚C keep; mixed with 2.5µL RNase; 37˚C 20min; stored at −20˚C). Quantitative PCR was done

using PowerUp SYBR Green Master Mix (Applied Biosystems A25742; 10µL reaction volume

with 5µL of 2X PowerUp SYBR Green Master Mix, 3.5µL H$_2$O, 0.5µL cDNA, and 1µL of 5µM

forward and reverse primer mix; Primers from IDT were dissolved with H$_2$O to make 100µM

stock, and 10µL of 100µM forward primer and 10µL of 100µM reverse primer were mixed with

180µL H$_2$O to make 5µM primer mix) and assessed with QuantStudio 6 Flex system (Applied

Biosystems; 96-well, 0.2mL; standard curve; SYBR Green reagents; standard) with the following

PCR condition: 50˚C 2 min, 95˚C 2 min; 40 cycles of 95˚C 15s, 55˚C 15s, 72˚C 60s (PCR

Stage); continuous cycles of 95˚C 15s, 55˚C 15s, 95˚C 15s (Melt Curve Stage). Expression

values calculated from the $2^{-\Delta\Delta Ct}$ comparative Ct method, $2^{\wedge}-[(Ct^{Target, Experimental} - Ct^{Housekeeping,}$

$^{Experimental}) - (Ct^{Target, Control} - Ct^{Housekeeping, Control})]$. Primers used for this study are as follows: HP2-S

Fwd 5'-ATTTGGGTGACTTATACGGGGACT-3', HP2-L Fwd 5'-ATCCCGAACCCAGCACAT-3',

HP2 Rev 5'-CTGGTTTAACTTGTTCCTTTTTGCCAGT-3', Und (as housekeeping) Fwd 5'-

GCAAGAAAAGCGGTCAGACT-3', Und (as housekeeping) Rev 5'-

CGTGTTGATACGGTCCAGAG-3', kl-5 Fwd 5'-TGTCTATGAGTGCCCAGTTTAC-3', kl-5 Rev

5'-GTCCACTTACTTGCTCTCTCTC-3', kl-3 Fwd 5'-ATGGTGCTGGTTGGGATAAG-3', kl-3 Rev

5'-TTGGCAGCCGTTGAGAAT-3', kl-2 Fwd 5'-TTTACCAGTGTCCCGCATATT-3', kl-2 Rev 5'-

AGTGCAGTACCTCGCTTTATC-3', ORY (Last Exon) Fwd 5'-

CAGCGTATAAACCGACAAATGG-3', ORY (Last Exon) Rev 5'-

GACAGCTCTCCGATTTCACTAA-3', ORY (Exons 4-5) Fwd 5'-ACTGTGCACTTCCCTTTGT-3',

ORY (Exons 4-5) Rev 5'-GAGATGAAATGGCGCAAGAAAT-3', CCY Fwd 5'-

CGGAGCCGTAAAGGATGATT-3', CCY Rev 5'-CGCTGACCTGATAACCCTATTC-3', DIP-λ

Fwd 5'-ACACCCTGGATACTCACAAATG-3', DIP-λ Rev 5'-AGTGAAACACACGCCAGAA-3',

CG44666 Fwd 5'-GCCGATTTGTGCAGTCTTTC-3', CG44666 Rev 5'-

GGACGATGTGGATGCTGTTA-3', Myo81F Fwd 5'-CCTCTTCATCGAGCGACTATTT-3',

Myo81F Rev 5'-ACGATGTTCCAGTAGGTGAATAC-3', Pzl Fwd 5'-

GGATGATATCCTACTACCAGCTTTG-3', Pzl Rev 5'-CCGGATAACAAGCCTCCTTTAT-3', Mitf

Fwd 5'-TCAGATGACTGTTGCGGTATAA-3', Mitf Rev 5'-ATACCCGTGTGGTGAGAATG-3'.

DNA FISH

Adult testes (for meiotic chromosome spread, typically three flies per slide) or third instar larval

brains (for mitotic chromosome spread, typically three larvae per slide) were dissected in PBS.

Samples were fixed for 4 min at RT in 50 µL of modified fixative (45% acetic acid (450µL of

glacial acetic acid) and 2.5% formaldehyde (550µL of fixative described earlier)) placed on

Superfrost Plus microscope slides (ThermoFisher Scientific 22-037-246). Samples were

squashed under a coverslip, and then immediately frozen in liquid nitrogen. Coverslips were

quickly removed and slides were immediately dehydrated in a Coplin jar filled with 100%

ethanol. Then, slides were dried at RT at least for 1 hour. 17 µL of hybridization buffer (10%

dextran sulfate (Millipore Sigma D8906; 1g in 2mL H_2O), 50% formamide (Millipore Sigma

S4117; 5mL), 2X SSC (Invitrogen AM9770; 1mL of 20X SSC), 10mM EDTA (Invitrogen

AM9260G; 200µL of 0.5M EDTA) in H_2O (up to 8.5mL); stored at −20˚C) was mixed with 3µL

probe mix (0.5 µM probe (1µL of 10µM probe solution) up to three colors or substitute with

H_2O). This hybridization mix was applied to samples on a slide and covered with a cover slip.

Samples were incubated at 95°C for 5 min, cooled and wrapped in parafilm, then incubated

overnight at RT in a dark humid chamber. In a Coplin jar filled with 0.1X SSC (5mL of 20X SSC

diluted with H_2O up to 1L), coverslips were removed and slides were washed three times for 15

min each. Slides were dried, sample locations were marked, and samples were mounted in

VECTASHIELD with DAPI (Vector Laboratories H-1200). For the whole testis staining, testes

were dissected in 1X PBS, fixed for 30 minutes at RT in 1mL modified fixative (1mM EDTA

(Invitrogen AM9260G; 2µL of 0.5M EDTA) and 1mL of fixative described earlier), rinsed with

1mL PBST-EDTA (1X PBS (Invitrogen AM9624; 5mL of 10X PBS), 0.1% Triton X-100 (Millipore

Sigma T9284; 500µL of 10% solution), 1mM EDTA (Invitrogen AM9260G; 100µL of 0.5M EDTA)

in H_2O (44.4mL); stored at RT), washed for 1hr at RT in 1mL PBST-EDTA, rinsed with 1mL

PBST (no EDTA), incubated for 10 min at 37˚C in 100µL RNase A (2mg/mL in PBST), washed

for 10 min at RT in 1mL PBST-EDTA, rinsed with 1mL SCCT-EDTA (50mL SCCT (5mL of 20X

SSC and 250µL of 20% Tween in 45mL H_2O; stored at RT) and 100µL of 0.5M EDTA; stored at

RT), washed for 15 min at RT in 100µL SCCT-20%F (800µL SCCT and 200µL formamide,

freshly made), washed for 15 min at RT in 100µL SCCT-40%F (600µL SCCT and 400µL

formamide, freshly made), washed for 15 min at RT in 100µL SCCT-50%F (500µL SCCT and

500µL formamide, freshly made), washed for 30 min at RT in 100µL SCCT-50%F, and put in

100µL hybridization mix (85µL hybridization buffer mixed with 5µL of 10µM probe up to three

colors or substitute with H_2O). Samples were denatured for 5 min at 95˚C and hybridized

overnight at 37˚C. Following hybridization, samples were washed two times in 1mL SCCT-

EDTA for 30 minutes each at RT and mounted in VECTASHIELD with DAPI (Vector

Laboratories H-1200). Images were acquired using Leica Stellaris8 confocal microscope with a

63X oil immersion objective lens (NA = 1.4) and processed/analyzed using ImageJ software.

Probes used for this study are as follows: Cy3-(TAGA)x8, Cy3/Cy5-(AATAC)x6,

Cy5/AlexaFluor-488-(AAGAG)x6.

Genotyping

X/Y; II+/II$^{\Delta AAGAG}$ (*bam>HP2^{RNAi}*) were mated with *y^{1}w^1* females for 5 days, and the eclosed flies

was collected a week later for genotyping. The *y$^+$* transgene on II$^+$ was used to genotype male

offspring by the *y$^+$* body color. Because X/Y; II$^+$/II$^{\Delta AAGAG}$ (*bam>HP2^{RNAi}*) carry a wild-type X

chromosome with the endogenous *y$^+$* gene, female offspring was genotyped by PCR using *y$^+$*

transgene-specific sequences. Each fly was homogenized in 50μL extraction buffer (1X TE

(Sigma Aldrich 93283; 49μL), 25mM NaCl (0.5μL of 2.5M NaCl), 200μg/mL proteinase K (0.5μL

of 20mg/μL), incubated for 30 min at 37˚C followed by 1 min at 95˚C. Genomic DNA was stored

in −20˚C. 1μL of genomic DNA was added in PCR mixture (DreamTaq DNA Polymerase

(ThermoFisher Scientific EP0705; 0.2μL), 0.5μM primer (2μL of 5μM forward and reverse primer

mix), 200μM dNTP (0.4μL of 10mM dNTP), DreamTaq Green Buffer (ThermoFisher Scientific

B71; 2μL) in H$_2$O (14.4μL)). Target is amplified by touchdown PCR (95˚C 3 min; 15 cycles of

95˚C 30s, 70˚C (−1˚C/cycle) 45s, 72˚C 30s; 20 cycles of 95˚C 30s, 55˚C 45s, 72˚C 30s; 72˚C 5

min and 10˚C keep). 10μL of PCR reaction was analyzed by gel electrophoresis of 1.5% TAE

gel 120V for 45min with 2μL of 100bp ladder. Primers used for this study are as follows: Fwd 5'-

CTAGAGTAAGTAGTTCGCCAGTTAAT-3' and Rev 5'-GCTGAATGAAGCCATACCAAAC-3'.

Fertility Assay and sex ratio scoring

One male (control or *bam>HP2^{RNAi}*) and three *y^{1}w^1* virgin females (1-3 days old) are mated for 4

days in the 25˚C incubator. After 4 days, parents were discarded and the number of eclosed

flies is counted a week later. After 4 days, parents were discarded and the numbers of eclosed

male and female progeny was counted for each mating pair. The sex ratio was calculated as the

number of female offspring/the total number of offspring.

Transgenic Fly Generation

The codon-modified fragment of HP2 (RNAi target sequence CAGGAGAGGAATGAAGAACAA

was changed to CAAGAACGCAACGAGGAGCAG) was synthesized by the IDT gBlocks Gene

Fragments service (1000ng; suspended in 100μL TE to make 10ng/μL solution and incubated at

50˚C for 20 minutes; stored at −20˚C) and cloned into UFO12514 (pUAST-based vector with

HP2-S cDNA, obtained from *Drosophila* Genomics Resource Center (DGRC) Stock 1648698)

using In-Fusion cloning. Briefly, the UFO12514 vector was amplified by standard PCR (95˚C 2

min; 30 cycles of 95˚C 20s, 55˚C 20s, 72˚C 30s/kb using Herculase II Fusion DNA Polymerase;

72˚C 3 min and 10˚C keep); 0.5µL vector (at least 50ng), 0.5µL insert (at least a few fold excess

of insert), 3µL H_2O, and 1µL In-Fusion Snap Assembly Master Mix (Takara Bio 638947) were

mixed and incubated for 15 min at 50˚C followed by 4˚C keep; 50µL DH5α (NEB C2987) and

5µL In-Fusion reaction mixture was mixed and incubated for 15 min on ice, and the bacteria was

heat shocked for 45 sec at 42˚C, on ice for 1 min, recovered in 200µL SOC (briefly if ampicillin

resistant, 1h at 37˚C if non-ampicillin resistant), and cultured on a LB plate with 100µg/mL

carbenicillin or other antibiotics overnight at 37˚C; Each colony is dissolved in 10µL H_2O, and

1µL was used for colony PCR (mixed with19µL DreamTaq PCR mixture used for genotyping

followed by touchdown PCR that amplify the insert) and 9µL was added in 200µL LB (stored at

4˚C during colony PCR, and cultured in 3mL LB overnight at 37˚C); 500µL of cultured *E. coli*

was mixed with 500µL of 50% glycerol and stored at −80˚C, and the rest of cultured *E. coli* was

used to collect plasmids using Miniprep (QIAGEN 27104; culture in 2mL LB overnight;

centrifuge for 3 min; dissolve in 250µL RNase solution; add 250µL SDS solution; gently invert 5

times; incubate at RT for 3min; add 350µL neutralizing solution; gently invert 5 times; centrifuge

for 10min; transfer 700µL supernatant to the spin column; add 700µL salt wash solution;

centrifuge for 30sec; add 700µL ethanol wash solution; centrifuged for 30sec; centrifuge for

another 1min; elute in 50µL water); The entire plasmid was sequenced by Plasmidsaurus Inc;

Small cloning errors were corrected by QuikChange Lightning Site-Directed Mutagenesis Kit

(Agilent 210518). HP2-L fragments were amplified from testis cDNA library: Briefly, 1µL of cDNA

and 5µL of 5µM forward and reverse primer mix for each fragment were added in PCR mixture

(Herculase II Fusion DNA Polymerase (Agilent 600675; 1µL), 250µM dNTP (1.25µL of 10mM

dNTP), 5x Buffer (10µL) in H_2O (31.75µL)). For each fragment, two tubes with 50µL PCR

reaction were prepared and amplified by touchdown PCR, adjusting the extension time (30s/kb).

The PCR products were run on 0.7% TAE gel and extracted by gel purification (QIAGEN 28506;

mix gel and binding solution in 1:1 ratio; 50°C for 5-10min; spin down; add 700µL binding

solution; centrifuge for 1min; add 700µL ethanol wash solution; centrifuge for 1min; centrifuge

for another 1min; elute in 20µL water). The two HP2-L fragments were fused by overlap

extension PCR: Briefly, 20µL PCR reaction (0.5µL of Herculase II Fusion DNA Polymerase,

0.5µL of 10mM dNTP, 4µL of 5x Buffer, ([length of fragment (bp) ÷ 10] ng of each fragment with

H_2O) was incubated in a thermocycler (95˚C 3 min; 14 cycles of 95˚C 30s, 72˚C (−0.5˚C/cycle)

45s, 72˚C 30s/kb; 72˚C 5 min and 10˚C keep). 30µL spike-in mixture (17.75µL H_2O, 5µL of 5µM

forward and reverse primer mix for the entire fragment, 6µL of 5x Buffer, 0.75µL of 10mM dNTP,

0.5µL of Herculase II Fusion DNA Polymerase) was added in each PCR reaction (total 50µL)

and incubated in a thermocycler (95°C 3 min; 14 cycles of 95°C 30s, 72°C (−0.5°C/cycle) 45s,

72°C 30s/kb; 20 cycles of 95°C 30s, 65°C 45s, 72°C 30s/kb; 72°C 5 min and 10°C keep). The

fused product was gel purified in 20µL H_2O and inserted into pCR-Blunt II-TOPO vector

(ThermoFisher Scientific 450245; 4µL gel purified DNA, 1µL salt solution, and 1µL TOPO vector

were incubated for 5 min at RT; 5µL reaction was transformed in 50µL DH5α and cultured on a

LB plate with 50µg/mL kanamycin). The fused HP2-L fragment was subcloned into the HP2-S

plasmid (amplified by long template (>10kb) PCR: 50µL Herculase PCR mixture was incubated

in a thermocycler (92°C 2 min; 10 cycles of 92°C 20s, 55°C 20s, 68°C 30s/kb; 20 cycles of 92°C

20s, 55°C 20s, 68°C 30s/kb +20s/cycle; 68°C 8 min and 10°C keep)) using In-Fusion cloning.

The PhiC31 integrase-mediated integration was conducted by BestGene Inc. The pUAST

construct with RNAi resistant HP2-S is inserted into the VK27 site on 3R (BDSC 9744). The

pUAST construct with RNAi resistant HP2-L is inserted into the attP2 site on 3L (BDSC 8622).

Both constructs were balanced with TM3.

Analysis of spermatocyte-enriched DNA-binding proteins

The list of proteins annotated as transcription factors or as containing a DNA-binding domain

were obtained from FlyBase, using the associated "Gene Group" lists. These lists were merged

and redundancies removed. The Fly Cell Atlas testis dataset[1] was used to identify

spermatocyte-enriched genes using the FindMarkers tool in Seurat v5.1 with default

paremeters, with spermatocytes defined as per the previous annotation (clusters C-G, as in

Figure 4A, Raz et al[1]). Bona fide markers were considered those with an adjusted p-value < 10^{-9}.

These lists were then intersected to obtain a catalog of putative transcription factors in

spermatocytes.

Comparison of transcribed and translated protein-coding genes

Mass-spec data from Gärtner 2019 (accessed from Project PXD010627 Proteomics

Identifications Database, folder MaxQuant_Output.zip, subfolder proteinGroups.txt) was sorted

by annotated "protein groups". We binned the data with any protein group with a specific

corresponding gene ID and at least one peptide across in all three adult testis replicates

considered "translated", and all others considered absent. Protein presence in non-adult stages

was not considered. Genes were considered transcribed if they had an average expression of

>=0.1 ln(UMIs-per-10,000 + 1) across the entire testis sequencing dataset[1]. These lists were

intersected to find genes transcribed (present in the sequencing dataset but not the mass-spec

dataset), translated (present in the mass-spec dataset but not the sequencing dataset), neither

or both.

Identification of transcribed transposable elements and lncRNAs

We generated a new reference file that contains both the DM6 reference and the consensus

sequences of *Drosophila* transposable elements (https://github.com/bergmanlab/drosophila-

transposons). Specifically, a .gtf file was generated from the consensus TE .fa file

(D_mel_tranasposon_sequence_set.fa) with every TE sequence considered an "exon". New .fa

and .gtf files were then generated by concatenating the respective files from DM6 and the TE

consensus. The Fly Cell Atlas testis dataset was then remapped to this new reference using

default parameters in the cellranger 8.0.1 tool. The resulting mapped file was processed,

dimensionality reduced, and clustered according to previous parameters[1]; and annotations were

transferred from the previous publication. lncRNAs, as identified in DM6, or TEs, as identified in

the consensus file, were considered "expressed" in spermatocytes if they had an average

expression of >=15 ln(UMIs-per-10,000 + 1) across all spermatocytes, with spermatocytes

annotated as above.

Statistics and Reproducibility

Information of statistics is provided in respective figure legends or Source Data file. For some

imaging results where quantification is not provided, experiments were repeated at least 2

times, each with 10 pairs of testes, and each testis contained hundreds of spermatocytes

(Figure 1i, 3e).

Data availability statement: The previously published data sets utilized are, Raz et al[1], NCBI
Gene Expression Omnibus GSE220615; Gartner et al.[21], PXD010627 (ProteomeCentral), and
modEncode (2010)[28] (www.modencode.org/publications/integrative_fly_2010/) as detailed in [28].
Source Data are available with this paper.

References

References

1. Raz, A.A. *et al.* Emergent dynamics of adult stem cell lineages from single nucleus and
single cell RNA-Seq of Drosophila testes. *Elife* **12** (2023).
2. Wen, K. *et al.* Critical roles of long noncoding RNAs in Drosophila spermatogenesis.
Genome Res **26**, 1233-1244 (2016).
3. Mills, W.K., Lee, Y.C.G., Kochendoerfer, A.M., Dunleavy, E.M. & Karpen, G.H. RNA
from a simple-tandem repeat is required for sperm maturation and male fertility in
Drosophila melanogaster. *Elife* **8** (2019).
4. Lawlor, M.A., Cao, W. & Ellison, C.E. A transposon expression burst accompanies the
activation of Y-chromosome fertility genes during Drosophila spermatogenesis. *Nat*
Commun **12**, 6854 (2021).
5. Soumillon, M. *et al.* Cellular source and mechanisms of high transcriptome complexity in
the mammalian testis. *Cell Rep* **3**, 2179-2190 (2013).
6. Chen, Y. *et al.* Single-cell RNA-seq uncovers dynamic processes and critical regulators
in mouse spermatogenesis. *Cell Res* **28**, 879-896 (2018).
7. Guo, J. *et al.* The adult human testis transcriptional cell atlas. *Cell Res* **28**, 1141-1157
(2018).
8. Wang, D. *et al.* A deep proteome and transcriptome abundance atlas of 29 healthy
human tissues. *Mol Syst Biol* **15**, e8503 (2019).
9. Xia, B. *et al.* Widespread Transcriptional Scanning in the Testis Modulates Gene
Evolution Rates. *Cell* **180**, 248-262 e221 (2020).
10. Liu, H. & Zhang, J. Higher Germline Mutagenesis of Genes with Stronger Testis
Expressions Refutes the Transcriptional Scanning Hypothesis. *Molecular biology and*
evolution **37**, 3225-3231 (2020).
11. Walker, P.M. Origin of satellite DNA. *Nature* **229**, 306-308 (1971).
12. Yunis, J.J. & Yasmineh, W.G. Heterochromatin, satellite DNA, and cell function.
Structural DNA of eucaryotes may support and protect genes and aid in speciation.
Science **174**, 1200-1209 (1971).
13. Lohe, A.R., Hilliker, A.J. & Roberts, P.A. Mapping simple repeated DNA sequences in
heterochromatin of Drosophila melanogaster. *Genetics* **134**, 1149-1174 (1993).
14. Shaffer, C.D. *et al.* Heterochromatin protein 2 (HP2), a partner of HP1 in Drosophila
heterochromatin. *Proc Natl Acad Sci U S A* **99**, 14332-14337 (2002).
15. Shaffer, C.D. *et al.* The large isoform of Drosophila melanogaster heterochromatin
protein 2 plays a critical role in gene silencing and chromosome structure. *Genetics* **174**,
1189-1204 (2006).
16. Stephens, G.E., Xiao, H., Lankenau, D.H., Wu, C. & Elgin, S.C. Heterochromatin protein
2 interacts with Nap-1 and NURF: a link between heterochromatin-induced gene
silencing and the chromatin remodeling machinery in Drosophila. *Biochemistry* **45**,
14990-14999 (2006).
17. Stephens, G.E., Slawson, E.E., Craig, C.A. & Elgin, S.C. Interaction of heterochromatin
protein 2 with HP1 defines a novel HP1-binding domain. *Biochemistry* **44**, 13394-13403
(2005).
18. Rathke, C., Baarends, W.M., Awe, S. & Renkawitz-Pohl, R. Chromatin dynamics during
spermiogenesis. *Biochim Biophys Acta* **1839**, 155-168 (2014).
19. Loppin, B. & Berger, F. Histone Variants: The Nexus of Developmental Decisions and
Epigenetic Memory. *Annu Rev Genet* **54**, 121-149 (2020).
20. Li, H. *et al.* Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly.
Science **375**, eabk2432 (2022).

21. Gartner, S.M.K. *et al.* Stage-specific testes proteomics of Drosophila melanogaster
identifies essential proteins for male fertility. *Eur J Cell Biol* **98**, 103-115 (2019).
22. Jagannathan, M., Warsinger-Pepe, N., Watase, G.J. & Yamashita, Y.M. Comparative
Analysis of Satellite DNA in the Drosophila melanogaster Species Complex. *G3*
(Bethesda) **7**, 693-704 (2017).
23. Wei, K.H. *et al.* Variable Rates of Simple Satellite Gains across the Drosophila
Phylogeny. *Molecular biology and evolution* **35**, 925-941 (2018).
24. Chang, C.H. & Larracuente, A.M. Heterochromatin-Enriched Assemblies Reveal the
Sequence and Organization of the Drosophila melanogaster Y Chromosome. *Genetics*
211, 333-348 (2019).
25. Bonaccorsi, S., Pisano, C., Puoti, F. & Gatti, M. Y chromosome loops in Drosophila
melanogaster. *Genetics* **120**, 1015-1034 (1988).
26. Bonaccorsi, S. & Lohe, A. Fine mapping of satellite DNA sequences along the Y
chromosome of Drosophila melanogaster: relationships between satellite sequences and
fertility factors. *Genetics* **129**, 177-189 (1991).
27. Fingerhut, J.M., Moran, J.V. & Yamashita, Y.M. Satellite DNA-containing gigantic introns
in a unique gene expression program during Drosophila spermatogenesis. *PLoS*
genetics **15**, e1008028 (2019).
28. mod, E.C. *et al.* Identification of functional elements and regulatory circuits by Drosophila
modENCODE. *Science* **330**, 1787-1797 (2010).
29. Lu, B.Y., Emtage, P.C., Duyf, B.J., Hilliker, A.J. & Eissenberg, J.C. Heterochromatin
protein 1 is required for the normal expression of two heterochromatin genes in
Drosophila. *Genetics* **155**, 699-708 (2000).
30. Wei, X., Eickbush, D.G., Speece, I. & Larracuente, A.M. Heterochromatin-dependent
transcription of satellite DNAs in the Drosophila melanogaster female germline. *Elife* **10**
(2021).
31. Castel, S.E. & Martienssen, R.A. RNA interference in the nucleus: roles for small RNAs
in transcription, epigenetics and beyond. *Nat Rev Genet* **14**, 100-112 (2013).
32. Hoskins, R.A. *et al.* The Release 6 reference sequence of the Drosophila melanogaster
genome. *Genome Res* **25**, 445-458 (2015).
33. Carvalho, A.B., Lazzaro, B.P. & Clark, A.G. Y chromosomal fertility factors kl-2 and kl-3
of Drosophila melanogaster encode dynein heavy chain polypeptides. *Proceedings of*
the National Academy of Sciences of the United States of America **97**, 13239-13244
(2000).
34. Hafezi, Y., Sruba, S.R., Tarrash, S.R., Wolfner, M.F. & Clark, A.G. Dissecting Fertility
Functions of Drosophila Y Chromosome
Genes with CRISPR. *Genetics* **214**, 977 (2020).
35. Carvalho, A.B., Dobo, B.A., Vibranovski, M.D. & Clark, A.G. Identification of five new
genes on the Y chromosome of Drosophila melanogaster. *Proceedings of the National*
Academy of Sciences of the United States of America **98**, 13225-13230 (2001).
36. Wei, K.H., Grenier, J.K., Barbash, D.A. & Clark, A.G. Correlated variation and population
differentiation in satellite DNA abundance among lines of Drosophila melanogaster. *Proc*
Natl Acad Sci U S A **111**, 18793-18798 (2014).
37. Ridges, J.T. *et al.* Overdrive is essential for targeted sperm elimination by Segregation
Distorter. *bioRxiv* (2024).
38. Courret, C., Wei, X. & Larracuente, A.M. New perspectives on the causes and
consequences of male meiotic drive. *Curr Opin Genet Dev* **83**, 102111 (2023).
39. Merrill, C., Bayraktaroglu, L., Kusano, A. & Ganetzky, B. Truncated RanGAP encoded by
the Segregation Distorter locus of Drosophila. *Science* **283**, 1742-1745 (1999).
40. Larracuente, A.M. & Presgraves, D.C. The selfish Segregation Distorter gene complex of
Drosophila melanogaster. *Genetics* **192**, 33-53 (2012).

41. Herbette, M. *et al.* Distinct spermiogenic phenotypes underlie sperm elimination in the
Segregation Distorter meiotic drive system. *PLoS genetics* **17**, e1009662 (2021).
42. Maheshwari, S. & Barbash, D.A. The genetics of hybrid incompatibilities. *Annu Rev*
Genet **45**, 331-355 (2011).
43. Presgraves, D.C. & Meiklejohn, C.D. Hybrid Sterility, Genetic Conflict and Complex
Speciation: Lessons From the Drosophila simulans Clade Species. *Front Genet* **12**,
669045 (2021).
44. Ferree, P.M. & Barbash, D.A. Species-specific heterochromatin prevents mitotic
chromosome segregation to cause hybrid lethality in Drosophila. *PLoS Biol* **7**, e1000234
(2009).
45. Satyaki, P.R. *et al.* The hmr and lhr hybrid incompatibility genes suppress a broad range
of heterochromatic repeats. *PLoS genetics* **10**, e1004240 (2014).
46. Bayes, J.J. & Malik, H.S. Altered heterochromatin binding by a hybrid sterility protein in
Drosophila sibling species. *Science* **326**, 1538-1541 (2009).
47. Van Doren, M., Williamson, A.L. & Lehmann, R. Regulation of zygotic gene expression
in Drosophila primordial germ cells. *Curr Biol* **8**, 243-246 (1998).
48. Chen, D. & McKearin, D.M. A discrete transcriptional silencer in the bam gene
determines asymmetric division of the Drosophila germline stem cell. *Development* **130**,
1159-1170 (2003).
49. Rebollo, E., Llamazares, S., Reina, J. & Gonzalez, C. Contribution of Noncentrosomal
Microtubules to Spindle Assembly in Drosophila Spermatocytes. *PLoS Biol* **2**, E8 (2004).
50. Shindo, Y. & Amodeo, A.A. Dynamics of Free and Chromatin-Bound Histone H3 during
Early Embryogenesis. *Curr Biol* **29**, 359-366 e354 (2019).
51. Park, J.I., Bell, G.W. & Yamashita, Y.M. Derepression of Y-linked multicopy protamine-
like genes interferes with sperm nuclear compaction in D. melanogaster. *Proc Natl Acad*
Sci U S A **120**, e2220576120 (2023).
52. Misra, S. *et al.* Annotation of the Drosophila melanogaster euchromatic genome: a
systematic review. *Genome Biol* **3**, RESEARCH0083 (2002).
53. Merel, V., Boulesteix, M., Fablet, M. & Vieira, C. Transposable elements in Drosophila.
Mobile DNA **11**, 23 (2020).
802

Acknowledgements: We thank Dr. Gary Karpen, and Dr. Andrew Clark for reagents, and the

members of the Yamashita lab for discussions and comments on the manuscript. We thank the

FlyBase, Bloomington *Drosophila* Stock Center, and *Drosophila* Genomics Resource Center for

reagents and information. **Funding:** This research is funded by the Howard Hughes Medical

Institute, Gordon and Betty Moore Foundation, Whitehead Institute for Biomedical Research (to

Y.M.Y), and National Institute of General Medical Sciences (F32GM143850 and K99GM154143

to A.A.R).

Author Contributions: Conceptualization, T.K. and Y.M.Y.; methodology, T.K.; investigation,

T.K. and M.N-D; software, A.R. and R. L.; resources, J.M.F. and Y.M.Y.; writing – original draft,

T.K.; writing – editing, T.K. and Y.M.Y.; funding acquisition, Y.M.Y.; supervision, Y.M.Y.

Competing Interests: The authors declare no competing interests.

Figure legends

Fig. 1: *Drosophila* spermatocytes exhibit widespread transcription

(a) Schematics of *Drosophila* spermatogenesis. Germline stem cells at the apical tip of the testis

(left in the cartoon) produce differentiating spermatogonia, which divide four times before

becoming spermatocytes. While spermatocytes grow in size, they exhibit widespread

transcription, with many protein-coding and non-coding transcripts. After meiosis, spermatid

nuclei undergo morphological changes through round, leaf, canoe, and needle stages, during

which histone is replaced by protamine. Mature sperm are stored in seminal vesicles and

prepared for ejaculation.

(b) Composition of *Drosophila melanogaster* genome[13, 24, 52, 53]. AAGAG satellite DNA occupies

approximately 10% of the genome.

(c) Transcription of protein-coding genes (12,760 genes) in the *Drosophila* testis. Approximately

70% of total protein-coding genes are transcribed (30% translated (dark blue), and 40% not

translated (blue)), while 30% are not transcribed (black).

(d-e) Transcription of lncRNAs (1764 genes) and transposons (122 transposons) in

spermatocytes vs. somatic cells (somatic epithelia surrounding each cyst). Note that the RNA

sequencing data [1] were obtained by polyA selection, thus do not capture transcripts that are not

polyadenylated such as satellite RNA.

(f, g) RNA FISH of AAGAG/CUCUU satellite transcripts in male germ cells in various stages of

differentiation (germline stem cells/spermatogonia, spermatocytes, meiotic cells). f) low

magnification images capturing the course of germ cell differentiation. g) individual cells at

various stages (signal intensities are not modified relative to each other).

(h) Quantification of AAGAG/CUCUU satellite RNA at various stages of male germ cell

differentiation based on FISH signal (see Methods). n = 50 cells were counted for each cell

type. The black bar in each row represents the mean value. The dotted black line at 1 indicates

the reference level. Signal intensities are shown as relative values normalized to the control

(early spermatocytes), whose average value was set to 1 (as described in the Methods). Two

biological replicates were performed.

(i) RNA FISH of AAGAG/CUCUU satellite transcripts in the spermatocyte of X/O males lacking

the Y chromosome, demonstrating these transcripts are not solely from the satellite DNA-rich Y

chromosome.

Fig. 2: HP2 is required for the transcription of AAGAG satellite DNA in spermatocytes

(a) RNA FISH of AAGAG/CUCUU satellite transcripts in control, *bam>HP2^RNAi* and

bam>AAGAG^RNAi spermatocytes. Spermatocyte nuclei are indicated by dotted circle.

(b) Relative signal intensities of AAGAG/CUCUU transcripts in control, *bam>HP2^RNAi* and

bam>AAGAG^RNAi spermatocytes (n = number of spermatocytes examined). Red line: mean. The

p-value is calculated by an unpaired t-test (two-tailed). #: CUCUU RNA was slightly but

significantly (p<0.05) upregulated in *bam>AAGAG^RNAi*. Each dot represents the RNA FISH

intensity in a single spermatocyte (Method). Exacat *p*-values are provided in the Source Data

file.

(c) HP2 enrichment on satellite DNA (right) and the abundance of satellite DNA in the

Drosophila melanogaster genome calculated from the ChIP input (left). HP2 ChIP-seq data

(modENCODE ID: 5593)[28] were analyzed using the k-Seek method [36] to quantify simple satellite

repeat abundance from short-read sequencing (Method).

Fig. 3: HP2 depletion leads to subfertility

(a) Fertility assay of wild type, *bam>HP2^RNAi* and *bam>AAGAG^RNAi* males. Each dot represents

the number of offspring from a single mating pair (n = 29 mating pairs for control, n = 27 for

bam>HP2^RNAi and *bam>AAGAG^RNAi*); red line, mean. The exact *p*-value is 0.000000071, which

was calculated by an unpaired t-test (two-tailed).

(b) Seminal vesicles of wild type, *bam>HP2^RNAi* and *bam>AAGAG^RNAi* males stained with DAPI.

Only the seminal vesicle was cropped for clarity using Fiji/ImageJ. An example of sperm

nucleus is marked with red.

(c) Cytological map of AAGAG satellite DNA and fertility genes on the Y chromosome[13, 22, 24]. *Pzl*

is an autosomal gene with large intronic AAGAG satellite DNA (Supplementary Figure S4). RT-

qPCR using the primers to amplify the last exon of indicated genes in *bam>HP2^RNAi* and

bam>AAGAG^RNAi compared to control with technical duplicates for each genotype.

(d) RT-qPCR using the primers to amplify exon-exon junction across AAGAG satellite DNA-

containing introns of *ORY* gene with technical duplicates for each genotype.

(e) RNA FISH of *ORY* mRNA in spermatocytes in control, *bam>HP2^RNAi* and *bam>AAGAG^RNAi*.

Cytoplasmic *ORY* mRNA granules (arrowheads) are formed in control and *bam>HP2^RNAi*, but

not in *bam>AAGAG^RNAi*.

Fig. 4: HP2 depletion leads to defective histone-to-protamine transition in spermatids

(a) Schematics of sperm development and histone-to-protamine transition. Histone-to-protamine

transition occurs in the canoe stage spermatids.

(b) Needle stage spermatids in control and *bam>HP2^RNAi* animals, and *bam>HP2^RNAi* animals

with HP2 rescue construct (RNAi-resistant HP2-S transgene) stained with DAPI. Spermatid

nuclei with defective sperm DNA compaction is indicated by arrowheads in ii. In some cysts (iii),

all spermatids within a cyst failed to compact DNA.

(c) Immunofluorescence staining of H3K9me3 and Mst77F in canoe spermatids from control

and *bam>HP2^RNAi* animals. Arrowheads indicate a spermatid nucleus that failed to undergo

histone-to-protamine transition (n=15 testes for each genotype).

(d) Immunofluorescence staining of H3K9me3 in needle spermatids from control and

bam>HP2^RNAi animals. Arrowhead indicates spermatid nuclei in *bam>HP2^RNAi* animals that failed

to undergo sperm DNA compaction and retained the H3K9me3 histone mark. Frequencies of

cysts with H3K9me3 histone mark in needle stage spermatid in control and *bam>HP2^RNAi*

animals is shown. (n = 10 testes for control, n = 13 testes for *bam>HP2^RNAi*). The exact *p*-value

with the Mann-Whitney U test (two-tailed) is 0.00000175.

Fig. 5: HP2 depletion preferentially harms Y-chromosome-containing spermatids,

leading to sex ratio meiotic drive

(a) Sex ratio of the offspring of control vs. *bam>HP2^RNAi* males. Each dot represents the sex

ratio of offspring from a single mating pair (n = 29 for control, n = 26 for *bam>HP2^RNAi*); red line,

geometric mean. The exact *p*-value is 0.0052, which was calculated by an unpaired t-test (two-

tailed).

(b) DNA FISH on needle stage spermatids in *bam>HP2^RNAi* spermatids using X- and Y-specific

satellite DNA X: TAGA satellite DNA. Y: AATAC satellite DNA. The frequency of defective

spermatids with X vs. Y chromosome is shown (n=232 spermatids from 14 testes). The exact *p*-

value is 0.0001, which was calculated by the chi-square test for goodness of fit for deviations

from 50% X and 50% Y.

(c) Schematics of *Drosophila melanogaster* karyotype with AAGAG satellite DNA location

(orange block, size is proportional to the estimate[13]). Large AAGAG satellite blocks are found in

intergenic regions of the Y chromosome and pericentromeric region of the chromosome II.

(d) Model for the functional significance of transcription.

Fig. 6: The amount of AAGAG satellite DNA determines sperm DNA compaction defects

caused by HP2 depletion

(a) Needle stage spermatids stained with DAPI from *bam>HP2^{RNAi}* animals with animals with

indicated chromosome II genotypes (II$^+$/II$^+$ and II$^{\Delta AAGAG}$/II$^+$).

(b) The frequency of cysts with defective spermatids (at needle stage) was scored for the

indicated genotypes. Each data point represents an individual testis. The red bar indicates the

median, and the black bar indicates the interquartile range. The p-value was calculated by the

Mann-Whitney U test (two-tailed). n = number of testes examined. The exact *p*-values were

0.0003 between control and ithaca16, 0.0001 between control and ithaca06.

(c) Genotype of spermatids generated from X/Y; II+/II$^{\Delta AAGAG}$ *bam>HP2^{RNAi}* spermatocytes, and

their frequency represented in the offspring. n=1145 male offspring and n=296 female offspring

were scored. The exact *p*-values were 0.00235 (in males) 0.727 (in females), calculated by the

chi-square test for goodness of fit for deviations from 50% II$^+$ and 50% II$^{\Delta AAGAG}$.

a

Nuclear Morphology

Round/Leaf Spermatids — **Canoe Spermatids** — **Needle Spermatids**

Histone-based → Protamine-based

b

Control

(i)

DAPI

10μm

bam>HP2^{RNAi}

(ii)

(iii)

HP2 Rescue

(iv)

c

Control — *bam>HP2^{RNAi}*

DAPI Merge

10μm

H3K9me3

Mst77F

d

Control — *bam>HP2^{RNAi}*

DAPI H3K9me3

10μm

H3K9me3

% cysts with defective H3K9me3+ spermatids
/total cysts in a testis (needles stage)

0%   50%   100%

Control

bam>HP2^{RNAi}

Mann-Whitney U test
p<0.01

a

Control

bam>HP2^RNAi

Female (%)

p<0.05

b

DAPI **X (TAGA)** **Y (AATAC)**

10µm

Deviation from
50% X-50% Y

p<0.01

Defective spermatids
in *bam>HP2^RNAi*

X
27 (12%)

Y
205 (88%)

c

AAGAG

Cen

X   Y   II   III   IV

d

Spermatogonium   Spermatocyte   Spermatid

AAGAG satellite DNA / with   AAGAG transcription ⌒ by HP2   Protamine
heterochromatin mark H3K9me3 ●   heterochromatin remodeling   incorporation

NOT compatible with   Compatible with
protamine incorporation   protamine incorporation