1    ***Euprymna berryi* as a comparative model host for *Vibrio fischeri* light organ symbiosis**
2

3    Avery M. Imes[1,2], Morgan N. Pavelsky[3], Klodia Badal[3], Derrick L. Kamp[4], John L. Briseño[4],
4    Taylor Sakmar[5], Miranda A. Vogt[5], Spencer V. Nyholm[4], Elizabeth A. C. Heath-Heckman[6], Bret
5    Grasse[5], Alecia N. Septer[3], Mark J. Mandel[1,2],*
6

[1] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
Madison, WI USA
[2] Genetics Training Program, University of Wisconsin-Madison, Madison, WI USA
[3] Department of Earth, Marine & Environmental Sciences, University of North Carolina, Chapel
Hill, NC USA
[4] Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT USA
[5] Eugene Bell Center, Marine Biological Laboratory, Woods Hole, MA USA
[6] Department of Integrative Biology, Michigan State University, East Lansing, MI USA
15

16   Running Title: *Euprymna berryi* light organ colonization
17

Keywords: bobtail squid; *Aliivibrio fischeri*; biofilm; motility; Type 6 Secretion System; apoptosis;
laboratory models

ORCIDs:
AM Imes: https://orcid.org/0000-0003-1978-8750
M Pavelsky: https://orcid.org/0009-0006-8516-665X
DL Kamp: https://orcid.org/0000-0003-1787-1825
JL Briseño: https://orcid.org/0000-0002-2702-5078
T Sakmar: https://orcid.org/0000-0001-6874-2703
MA Vogt: https://orcid.org/0009-0004-8720-052X
B Grasse: https://orcid.org/0000-0001-6799-8715
SV Nyholm: https://orcid.org/0000-0001-5597-1945
EAC Heath-Heckman: https://orcid.org/0000-0001-5265-9032
AN Septer: https://orcid.org/0000-0002-9309-2930
MJ Mandel: https://orcid.org/0000-0001-6506-6711
33

34   * Correspondence to:
35

Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu

**ABSTRACT**

Functional studies of host-microbe interactions benefit from natural model systems that enable exploration of molecular mechanisms at the host-microbe interface. Bioluminescent *Vibrio fischeri* colonize the light organ of the Hawaiian bobtail squid, *Euprymna scolopes*, and this binary model has enabled advances in understanding host-microbe communication, colonization specificity, *in vivo* biofilms, intraspecific competition, and quorum sensing. The hummingbird bobtail squid, *Euprymna berryi,* can be generationally bred and maintained in lab settings and has had multiple genes deleted by CRISPR approaches. The prospect of expanding the utility of the light organ model system by producing multigenerational host lines led us to determine the extent to which the *E. berryi* light organ symbiosis parallels known processes in *E. scolopes*. However, the nature of the *E. berryi* light organ, including its microbial constituency and specificity for microbial partners, have not been examined. In this report, we isolate bacteria from *E. berryi* animals and tank water. Assays of bacterial behaviors required in the host, as well as host responses to bacterial colonization, illustrate largely parallel phenotypes in *E. berryi* and *E. scolopes* hatchlings. This study reveals *E. berryi* to be a valuable comparative model to complement studies in *E. scolopes*.

**IMPORTANCE**

Microbiome studies have been substantially advanced by model systems that enable functional interrogation of the roles of the partners and the molecular communication between those partners. The *Euprymna scolopes-Vibrio fischeri* system has contributed foundational knowledge, revealing key roles for bacterial quorum sensing broadly and in animal hosts, for bacteria in stimulating animal development, for bacterial motility in accessing host sites, and for *in vivo* biofilm formation in development and specificity of an animal's microbiome. *Euprymna berryi* is a second bobtail squid host, and one that has recently been shown to be robust to laboratory husbandry and amenable to gene knockout. This study identifies *E. berryi* as a strong symbiosis model host due to features that are conserved with those is *E. scolopes*, which will enable extension of functional studies in bobtail squid symbioses.

**INTRODUCTION**

Symbiotic microbes play critical roles in animal development, immunity, and health. Invertebrate
animal models have played foundational roles in elucidating the mechanisms by which
microbiomes are established and maintained, and the bobtail squid-*Vibrio fischeri* system has
been extensively studied with regard to host, symbiont, and the interaction between the two (1–
5). Many bobtail squid species harbor a light-emitting organ that is colonized by luminescent
bacteria, the light from which camouflages the host's silhouette as the animal forages under
moonlight (6, 7). During each generation, squid hatch aposymbiotically and acquire their
symbiotic bacteria from the seawater. Over the course of hours, the colonization initiation
process selects for the correct symbionts. Host-microbe communication continues into the
persistence phase as the host provides nutrients to the bacteria, which in turn produce light for
the host. A majority of symbionts are expelled each day and the remaining bacteria continue on
a diel cycle of regrowth and quorum sensing and luminescence (8, 9).

Most bobtail squid studies have focused on the Hawaiian bobtail squid, *Euprymna scolopes*.
Studies in *E. scolopes* have revealed bacterial factors that are required for colonization initiation
(10, 11), including flagellar motility (12, 13) and symbiotic biofilm formation (14–16).
Luminescence, which is produced at high cell density in response to quorum sensing, is
required during the persistence phase; while dark mutants can colonize successfully at 24 h,
beginning at 48 h they exhibit a defect and are not detectable after a few weeks (17–19). Work
in *E. scolopes* has additionally revealed principles regarding host responses to bacterial
colonization. Ciliated epithelial appendages that recruit the symbionts undergo apoptosis and
regression in response to release of the bacterial peptidoglycan fragment tracheal cytotoxin
(TCT), with apoptotic nuclei detectable as early as 24 h post infection (20–22). Additionally, TCT
triggers the migration of macrophage-like hemocytes into the light organ, where the hemocytes
begin a light organ maturation process that results in the stable tolerance of the *V. fischeri*
symbiont (23–25). As a result of the cited studies and many others, it is clear that there is a rich
dialogue between host and microbe during the colonization process (26).

The availability of multiple bacterial strains has facilitated comparative approaches that have
yielded a broad understanding of microbiome dynamics and evolution. Interbacterial competition
takes place inside the host, as revealed from the function of a Type 6 Secretion System that is
present in approximately half of known strains and enables those strains to kill competitors that
lack the system (27). Comparisons between fish and squid symbionts revealed differential
biofilm regulation as a key basis for host colonization specificity in the North Pacific Ocean (28).
However, there have not been extensive functional host studies conducted outside of *E.*
*scolopes*. Pilot experiments in *Euprymna tasmanica* have served as the basis for colonization
studies in the laboratory, and the light organ of *Sepiola robusta* has been characterized (29–31).
At the same time, a variety of cephalopod models have been cultivated both within and outside
the squid-*Vibrio* field; *E. scolopes* have been used to build cephalopod reference genomes (32,
33), multiple *Euprymna* species have revealed novel biology in neuroscience (34, 35), and a
broad range of cephalopods have been used to study transcriptome plasticity (36).

*Euprymna berryi*, the hummingbird bobtail squid, has been suggested as a cultivable model for
symbiosis studies. Recent efforts have expanded the aquaculture viability of valuable
cephalopod models including *E. berryi*. Previous work culturing *E. berryi* noted similarity
between *E. berryi* and *E. scolopes* and suggested that *E. berryi* could make a promising model
system due to higher survivorship and fecundity in aquaculture compared to *E. scolopes (37)*.
The first CRISPR-mediated knockouts in bobtail squids were performed in *E. berryi* to remove
pigmentation from the organism, demonstrating the ability to genetically manipulate *E. berryi*
(38). Therefore, the focus of this study is to evaluate the suitability of lab-raised *E. berryi* for
mechanistic studies in an animal-microbe symbiosis.

**RESULTS**

***Euprymna berryi* bobtail squid harbor luminescent bacteria in a light organ**. Given the
prominence of *E. scolopes* as a model host for light organ symbioses, we directly compared the
adult and hatchling *E. berryi* animals with their *E. scolopes* counterparts. While similar in overall
appearance, the *E. berryi* animals were larger than the *E. scolopes* animals (**Fig. 1, Fig. S1**),
and the *E. berryi* animals were darker brown in color compared to *E. scolopes* adults when the
animals were placed on the same substrate. The eggs laid by *E. berryi* animals were also
darker and had more defined structures outside of the eggs. However, the hatchling animals of
both species were similar in color, with the *E. berryi* hatchlings notably larger (**Fig. 1**).

**Figure 1. Comparison of *Euprymna* species.** (A) *E. berryi* (left) and *E. scolopes* (right) eggs immediately prior to hatching. (B) An *E. berryi* and *E. scolopes* hatchling 24 hours post-hatching. (C) An adult female *E. berryi*. (D) An adult female *E. scolopes*.

133
134

**Supplemental Figure S1.** *E. berryi* **males have larger mantle lengths than** *E. scolopes***.** Mantle lengths of adult *Euprymna* held in laboratory aquaculture. Mantle lengths were estimated in FIJI. Circles represent individual animals and the median is plotted.

We used GFP-expressing *V. fischeri* strains as biological probes to detect a possible light organ in *E. berryi.* The strains used included the *E. scolopes* light organ isolate ES114 and the *E. berryi* isolate BW2 that we describe in detail later. *E. berryi* were reported to contain a luminous organ when first identified (39), so we asked whether this organ was colonized by bioluminescent bacteria as observed in closely related *Euprymna* species. We introduced the bacteria to water containing *E. berryi* hatchlings, washed the hatchlings at 3 and 24 h, and assayed luminescence of the animals at 48 h. All animals were luminous and were fixed for imaging. Confocal imaging of the animals revealed that the bacteria of both strains were localized to the light organ in the mantle cavity in a structure that strongly resembled the light organ of *E. scolopes* (40). Representative images from BW2 are presented in Fig. 2 and quantification of the distribution of both strains are presented in Fig. S2. The bilobed organ of *E. berryi* contains three crypts on each side, with a dedicated exterior pore connecting to each crypt. The posterior pore leads to Crypt 2 (c2), middle pore to c1, and anterior pore to c3 (Fig. 2), which matches the morphology observed in *E. scolopes* (40). The relative crypt size is conserved with *E. scolopes*, with c1 as the dominant crypt, and c2 as the next smallest followed by c3 (Fig. S2). c1 contains approximately 70% of the bacteria by volume (Fig. S2). Furthermore, we observed antechambers and host anatomical bottleneck structures that were first described in *E. scolopes* (Fig. 2C) (40). Overall, imaging suggested strong conservation of light organ structures between *E. scolopes* and *E. berryi*.

**Figure 2. The light organ of juvenile *E. berryi.*** Images are representative of all symbiotic animals (n=10). (A) Representative image of an *E. berryi* light organ. Image is a maximum intensity projection of confocal z-stack approximately 150 µm deep. DAPI staining (blue) of host tissue shows that each side of the bi-lobed organ has a long anterior appendage (aa) and shorter posterior appendage (pa). At the base of each anterior appendage are three pores (p). Three crypts (c1-3) are on each side. Each crypt is visualized by GFP-expressing *V. fischeri* (green). The hindgut (hg) lies in the anterior end. (B) A single lobe of the light organ, counterstained with Alexa640-conjugated phalloidin to visualize actin cytoskeleton. (C) Enlarged image of boxed region in B. Pores (p) lead to crypts. A tight bottleneck (bn) separates the antechamber (ac) from crypt 1 (c1). (D) Maximum intensity projection and orthogonal views of crypts from one lobe of the light organ. Locations of orthogonal slices represented by dashed line. Each of the three crypts (c1, c2, c3) is fully colonized by GFP-expressing *V. fischeri.* (E) Three-dimensional model of the crypts, each a different color. Crypt 1 (c1) is the largest crypt, with branching structures. The smaller crypts 2 (c2) and 3 (c3) lay dorsally to crypt 1. Bars: A&B = 150 µm, C = 10 µm, D&E = 100 µm.

**Supplemental Figure S2. ES114 occupies a greater crypt volume in crypt 1 than BW2.** Z-stack images of single light organ lobes approximately 150 µm deep were taken at 1.2 µm intervals. Animals were colonized with a pVSV102 GFP-expressing derivative of the indicated strain. Image stacks were processed and analyzed using Arivis software, using the GFP signal to determine the space occupied by the *V. fischeri* and calculate the size of the crypt. The median of each group is shown with a horizontal bar. ES8, an animal that vented its crypt bacteria during processing, is marked with an X and omitted from statistical analysis.

**160**
**161**
**162**
**163**  ***V. fischeri* and other γ-proteobacteria were isolated from *E. berryi* adult tank water and**
**164**  **colonized hatchlings**. We sought to identify bacterial strains that colonize the *E. berryi* light
**165**  organ. We noticed that *E. berryi* raised with access to water pumped in from Eel Pond (Woods
**166**  Hole, MA, USA) readily became luminous. Therefore, we used bacteria isolated from these
**167**  luminous *E. berryi* hatchlings as a source for putative *E. berryi* colonizing strains. As a second
**168**  source for bacterial isolates, we hypothesized that *E. berryi* may expel bacteria in a rhythmic
**169**  daily cycle similar to *E. scolopes* (9, 23, 41), so we isolated bacteria from the tank water of wild-
**170**  caught *E. berryi* adult animals. In both cases, we isolated bacteria on LBS agar, streak-purified
**171**  different colony morphologies, and conducted secondary screening for isolates that were
**172**  detected as luminous in liquid LBS medium. We identified nine (9) distinct strains from the adult
**173**  "*berryi* water" (BW) sampling: two of these isolates represented colony types that were highly
**174**  abundant in our sampling yet minimally luminous (BW4, BW8), while the other seven were
**175**  luminous. From the "*berryi* hatchling" (BH) sampling, we identified two (2) strains, both of which

176 were luminous. For each strain above, we grew a culture in LBS media, obtained whole genome
177 Nanopore sequencing, and determined the likely species by FastANI analysis (42). This
178 analysis identified three *V. fischeri* strains (BH2, BW2, BW9), 7 other *Vibrio spp.* isolates, and
179 one *Pseudoalteromonas flavipulchra*. Figure 3 lists the isolates and shows specific
180 luminescence for each strain when grown in liquid culture.

**Figure 3. Isolation and characterization of bacterial strains.** Hatchling isolates were plated from colonized *E. berryi* hatchlings. Water isolates were plated from tank water housing a wild-caught adult *E. berryi*. Isolate species was identified by average nucleotide identity. Specific luminescence for each strain when grown in liquid culture was plotted with circles representing individual biological replicates. The median is indicated by a vertical bar.

182
183
***E. berryi* is colonized specifically by *V. fischeri* bacteria**. The *E. scolopes* light organ is
colonized only by *V. fischeri* bacteria, yet we isolated a number of bacterial species from the
water and hatchlings. We therefore asked whether *E. berryi* was more permissive to
colonization. We conducted standard 3 h inoculation, 48 h colonization assays that are well-
established for *E. scolopes* and that we adapted for *E. berryi* (43). The key adjustment needed
from our published protocol was to accommodate the larger size of the hatchlings by cutting the
transfer pipet farther up the stem and transferring the animals at a slower pace to ensure that
they remain in the liquid stream as they pass through the pipet opening.

We found that our *E. berryi*-isolated *V. fischeri* strains generally colonized *E. berryi* hatchlings,

with BW9 as the only strain showing slightly reduced CFU counts (Fig. 4A). In contrast, the

other species isolated from *E. berryi* water were unable to colonize, as was the *V. campbellii*

isolate BH1 isolated from a hatchling. These colonization patterns match what has been

reported for colonization of non-*V. fischeri* strains in *E. scolopes* (44, 45).

**Figure 4. *V. fischeri* are the exclusive colonizers of *E. berryi*.** *E. berryi* hatchlings were used in single-strain colonization experiments, where circles represent individual animals. The limit of detection for this assay, represented by the dashed line, is 7 CFU/light organ. Horizontal bars represent the median for each group.

199
200

201     *V. fischeri* isolated from squid light organs in the North Pacific Ocean and the Mediterranean
202     Sea that can robustly colonize *E. scolopes* are similarly able to colonize *E. berryi* (Fig. 4A). This
203     includes *V. fischeri* from *E. scolopes, Euprymna morsei*, *Sepiola robusta, and Sepiola affinis.*
204
The only divergence between colonization patterns in *E. scolopes* and those in *E. berryi* were
noted for the fish isolates MJ11 and CG101. These strains do not robustly colonize the
Hawaiian host (28, 46, 47), yet for both strains we observed a bimodal colonization pattern of *E.*
*berryi* (Fig. 4A). In both cases, most animals had either undetectable levels of bacteria or >10$^3$
CFU per squid. While in the case of MJ11, the bimodality partitioned by experiment, in neither
case did it correlate with the concentration of bacteria in the inoculum (Fig. 4A, Fig. S3).

**Supplemental Figure S3. Inoculum concentration did not positively correlate to squid colonization concentration in strains that exhibited bimodality.** (A) A reproduction of monocolonization data for select strains as presented in Fig4. Each symbol represents an individual animal, where color and shape indicate the replicate group that animal belonged to. (B) The inoculum concentration provided to squid, where each symbol represents the concentration of the indicated strain provided to the squid group within an experiment. Horizontal bars represent the median in both graphs. Colors in panel A and B correspond to the same replicate across experiments.

211
The above results suggest that most colonization patterns for *E. scolopes* strains are conserved
during colonization in *E. berryi*. We therefore asked whether the new *V. fischeri* isolates from *E.*
*berryi* were capable of colonizing *E. scolopes* squid. Not all *V. fischeri* can colonize *E. scolopes,*
although squid isolates of *V. fischeri* (including those from other bobtail squid species) are
typically able to colonize *E. scolopes* (28, 46, 48, 49). All three of the new *V. fischeri* strains

isolated from *E. berryi* samples, BW2, BW9, and BH2, consistently colonized *E. scolopes* squid
(Fig. S4).

**Supplemental Figure S4.** *E. scolopes* **are colonized by** *V. fischeri* **isolated from** *E. berryi.* *E. scolopes* hatchlings were used in single-strain colonization experiments, where circles represent individual animals. The limit of detection for this assay, represented by the dashed line, is 7 CFU/ light organ. Horizontal bars represent the median for each group.

219
220
221

**Figure 5. Colonization-deficient mutants are defective in *E. berryi*.** *E. berryi* hatchlings were used in single-strain colonization experiments, where circles represent individual animals. The limit of detection for this assay, represented by the dashed line, is 7 CFU/light organ. Horizontal bars represent the median for each group. All mutants are constructed from the wildtype strain ES114. The inoculum strain is indicated on the X-axis, with the hour squid were fixed post-inoculation labeled below each group. Treatment groups were compared to the ES114 control group using a non-parametric one-way ANOVA (Kruskal-Wallis) test using Dunn's multiple comparisons test.

222
223
**Bacterial factors required for host colonization in *E. scolopes* are similarly required for**
**colonization of *E. berryi*.** Decades of work in *E. scolopes* have defined bacterial colonization
factors (10, 11, 13–15, 50–52). If the process of colonization is conserved in *E. berryi*, then a
specific prediction is that mutants that fail to undergo key steps in the initiation or persistence of
the symbiosis will exhibit colonization defects in *E. berryi*. To test this hypothesis, we examined
ES114 mutants with initiation defects in the Syp biofilm (Δ*sypF*) or in flagellar motility (Δ*flrA*) and
determined their ability to colonize *E. berryi*. Both strains exhibited significant colonization
defects with a majority of animals completely uncolonized (Fig. 5). In *E. scolopes*, non-
luminescent mutants such as Δ*luxCDABEG* (Δ*lux*) are known to colonize the light organ but fail

to persist. We observed a similar defect in *E. berryi*, with no significant defect at 24 h post
inoculation (hpi), but with a highly significant defect and 4-fold fewer bacteria at 48 hpi (Fig. 5).
This pattern matches that observed in *E. scolopes,* wherein the Δ*lux* mutant first exhibits its
persistence defect at 48 hpi and is eventually lost from the *E. scolopes* light organ over 3-4
weeks (17–19). These data together argue that colonization in *E. berryi* bears large parallels to
that which occurs in *E. scolopes*, especially with regard to specificity for bacterial strains and
genetic requirements during symbiont entry and establishment in the host.
240

**Figure 6. *V. fischeri* strains stimulate variable intensities of host development responses in *E. berryi.*** *E. berryi* hatchlings were inoculated with a single bacterial strain and evaluated at 24 hours post-inoculation, where circles represent individual animals. Horizontal bars represent the median for each group. New bacterial isolates from *E. berryi* are shown as blue circles. (A) The TUNEL+ counts in the anterior appendage were taken as a measurement for cell apoptosis, a marker of colonization-driven host development. (B) Hemocyte counts in the anterior appendage were used to measure hemocyte trafficking into the appendage, which is elevated in colonized *E. scolopes.* P-values were calculated using a Kruskal-Wallis test with Dunn's multiple comparisons test.

241
242
243

***The E. berryi* appendages exhibit developmental phenotypes cued by bacterial**
**colonization**. In *E. scolopes*, colonization by the *V. fischeri* symbiont triggers irreversible host
morphological changes, particularly the dramatic apoptosis of the light organ appendages (21).
In *E. scolopes,* colonized squid have detectable TUNEL staining in the anterior appendage at 24
hpi as a result of DNA fragmentation in apoptotic cells. To determine if similar host development
is triggered by symbiotic colonization in *E. berryi*, we used a TUNEL assay. In *E. berryi,* BW9
induced the highest TUNEL count, followed by BH2 (Fig. 6A). Another measure for host
development stimulated by symbiont colonization is hemocyte trafficking into the light organ
appendages, which is elevated at 24 hpi in *E. scolopes* and minimal in aposymbiotic animals.
Among the strains tested, BW9 and BH2 again promoted the strongest responses but with a
wide range of variability across animals in all groups (Fig. 6B). BW2 was the only strain that did
not yield statistically significant hemocyte trafficking (Fig. 6B). Overall, these results point to
conserved host responses in *E. berryi* compared to *E. scolopes*, and with BH2 and BW9 (along
with ES114) able to elicit both TUNEL staining and hemocyte trafficking in *E. berryi*.
258
259

**Figure 7. *E. berryi* can be used to study T6SS-mediated competition in *V. fischeri*
symbionts.** (A) Coincubation assays between the indicated strain and a GFP-tagged ES114
target strain. The scale bar for images in panel A is 2 mm. (B) Representative images of T6SS2
sheaths in wildtype BW2 and the *tssF* derivative carrying a VipA-GFP expression vector. The
scale bar for images in panel B is 2 µm. (C) Single colonization assays for indicated ES401 strains
at 24 hr. (D) Competitive colonization assays with GFP-tagged ES114 and RFP-tagged ES401-
derived strain. Time points are 24hr post-inoculation. Panel D shows the competitive index (CI) in
log 10 scale for each squid, where the ending strain ratio is divided by the beginning strain ratio.
Statistical comparison was made using a Mann-Whitney test.

260
261

**Type 6 secretion-mediated competition operates in the *E. berryi* host**. Among the
competitive mechanisms influencing squid colonization, some *V. fischeri* strains are known to
use a strain-specific type VI secretion system found on chromosome II (T6SS2), which confers
a contact-dependent ability to kill off competitive strains (27). We aimed to determine if this
mechanism for interbacterial competition operates and influences colonization of *E. berryi*. All
three *V. fischeri* isolates from *E. berryi* encode T6SS2. To determine if the system was
functional, we co-incubated the isolates with GFP-expressing ES114. The control strain for this
assay, the *E. scolopes* isolate ES401, kills the target ES114-GFP and results in no detectable
fluorescence from the resulting colony spot. In the absence of the structural protein TssF,
T6SS2 activity is prevented and ES114-GFP is able to grow and produce fluorescence. This
phenotype was mirrored in the isolates from *E. berryi*: all three strains exhibited functional
T6SS2 activity, and this activity was removed in BW2 by mutation of *tssF* (Fig. 7A). A halo effect
seen in BW2 has been previously observed in other T6SS2$^+$ strains and sometimes occurs
when survivors at the edge of the colony escape T6SS-mediated killing. Moreover, when BW2-
derived strains expressed a GFP-tagged sheath component (VipA), sheaths were observed in
the parent strain but not seen for the *tssF* mutant, which lacks the ability to make T6SS2
sheaths (Fig. 7B) and results in a non-killing phenotype (Fig. 7AB).
279
To ask whether T6SS-mediated competition occurs in *E. berryi* juveniles, as it does in E.
scolopes (53), we conducted competition assays within the squid host. Both ES401 and its non-
killing *tssF* derivative are able to consistently colonize *E. berryi* in single-strain colonizations
(Fig. 7C). To determine if T6SS2 competition functions in the *E. berryi* host, we used a
competitive colonization assay between ES401 and ES114, where strains are differentially
fluorescently labeled and co-inoculated into squid hatchlings. ES401 exhibited a 60-fold
competitive colonization advantage over ES114 (Fig. 7D; comparing the *tssF*$^+$/*tssF*$^-$ results).
This result confirmed that T6SS2 in the *V. fischeri* isolates from *E. berryi* is functional and that
the system provides a competitive advantage in colonization in *E. berryi*.

**DISCUSSION**
290
Overall, this work establishes *E. berryi* as a culturable model for bobtail squid-*V. fischeri*
symbiosis studies. We demonstrate that for the light organ symbiosis, the host anatomy, host-
microbe specificity, bacterial gene requirements, interbacterial competition, and host responses
to symbiosis broadly mirror those processes observed in the well-studied *E. scolopes* model.
Recent work in wild-caught *E. berryi* characterized the microbial community in the female
accessory nidamental gland (ANG), a reproductive organ in which bacteria provide key
defensive roles including provisioning antifungal compounds into the jelly coat of the developing
squid eggs (54–57). Therefore, the ability to use cultured animals for bobtail squid symbiosis
studies, in a host species in which it was recently shown that gene knockouts could be
conducted (38), opens additional resources to investigate how animal microbiomes are
established across both the light organ and ANG microbiomes.
302

One area in which we observed a variable response in *E. berryi* was in the host responses to
symbiotic colonization. Although all strains tested exhibited a significant response in the TUNEL
assay, the overall response was weak compared to *E. scolopes*, which averages around 40
TUNEL+ cells per anterior appendage in colonized animals at the 24 hpi time point (22). Only
strain BW9 exhibited median counts of >20 per appendage (Fig. 6A). Similarly, the average
hemocyte count induced by ES114 has been previously observed to be higher in *E. scolopes*
than we observed in Fig. 6B*,* averaging 15 per anterior appendage (58). One possibility is that
the host responses in *E. berryi* are qualitatively similar but quantitatively distinct from *E.*
*scolopes*. Another possibility is that the timeline for the response in *E. berryi* differs from that in
*E. scolopes*. A future study could examine this question by comparing the responses side-by-
side, and carrying out the timeline to 5-7 d. A general limitation of this study is that most work
was examined only to the 48 h timepoint, so differences at later times are possible and were not
examined in this initial investigation.
316
We expect that the non-*V. fischeri* strains isolated from the *E. berryi* water and hatchlings were
not light organ symbionts, but instead were environmental samples. We did not conduct light
organ dissections for the BH samples, so bacteria present in other tissues could have been
isolated if they were osmotically protected during freezing. Given that the non-*V. fischeri* strains
failed to colonize aposymbiotic *E. berryi* hatchlings when presented at >$10^3$ CFU/ml in
monoassociation suggests that they are not able to colonize the light organ. The *V. fischeri*
isolates BH2, BW2, and BW9 are clearly capable of colonizing the bobtail squid light organs,
both for *E. berryi* and *E. scolopes*, and with BW2 we confirm specific light organ colonization via
direct imaging (Fig. 2). These *V. fischeri* isolates may have originated from the adult *E. berryi* in
the aquaculture system or from the Atlantic Ocean water feeding the tank systems. Strain BH2
colonizes *E. berryi* consistently across replicates (Fig. 4) and also stimulates significant host
responses in both assays tested (Fig. 6), suggesting that it may be a natural symbiont and/or
share characteristics with natural *E. berryi* symbionts. When it is possible to isolate symbionts
from wild *E. berryi* animals, it would be valuable to address the phylogenetic distribution and
phenotypes of those symbionts.
332
In summary, we describe *E. berryi* as a model for *V. fischeri* colonization and light organ
symbiosis. This work therefore will enable use of tools developed for *E. berryi* to be applied for
symbiosis studies and will allow for comparative studies between bobtail squid hosts.

**MATERIALS AND METHODS**

**Animal comparison**
**Eggs/embryos**. *Euprymna* spp. eggs were imaged using a Nikon D810 digital camera at
University of Wisconsin-Madison and measured using FIJI (59). *Euprymna* spp. hatchlings were
imaged within 24 hours post-hatching using a Leica M60 stereomicroscope with Leica FireCam
software. **Adults**. Adult *E. berryi* (120-178 d post-hatching) were raised at Woods Hole Marine
Biological Laboratory and photographed using an iPad 9th generation (model number
MK2K3LL/A). Adult *E. scolopes* were collected from Oahu, Hawaii, housed in an aquaculture
facility for 61 d, and were photographed using a Nikon D810 digital camera. Mantle lengths

were measured using FIJI (59). A non-parametric one-way ANOVA (Kruskal-Wallis) using
Dunn's multiple comparisons test was used to compare the mantle lengths between males and
females of each species.
348
**Bacterial strains and media**
Bacterial strains used in this study are listed in Table 1. All bacteria except for *E. coli* were
grown in Lysogeny broth salt (LBS) medium at 25˚C (25 g Difco LB broth [BD], 10 g NaCl, and
50 mL 1 M Tris buffer [pH 7.5] per liter). An *E. coli* strain, used as the conjugation donor for the
GFP plasmid, was grown in Lysogeny broth (LB) medium at 37˚C (25 g Difco LB broth [BD] per
liter). When required, kanamycin was added to a final concentration of 100 µg/ml (*V. fischeri*) or
50 µg/ml (*E. coli*), and erythromycin to 5 µg/ml (*V. fischeri*). *E. coli* strains requiring erythromycin
were grown in Brain-Heart Infusion (BHI) medium at 37˚C (37 g BBL BHI powder [BD] per liter)
with 150 µg/ml. Growth media was solidified to 1.5% agar as needed by adding 15 g Bacto agar
(BD) per liter.
359
**Squid monocolonizations**
Hatchling squid, either *E. scolopes* or *E. berryi*, were colonized according to our previously
described protocol (43) and summarized below. Bacterial strains were grown overnight with
aeration in LBS at 25˚C. Cultures were diluted 1:80 in LBS and subcultured until an approximate
$OD_{600}$ of 0.3. This culture was used to inoculate 40 mL of artificial seawater containing squid
hatchlings, with the inoculum normalized to approximately $5 \times 10^3$ CFU/mL. Squid remained in
the inoculum water for 3 h at room temperature. Squid were then washed in filter-sterilized
Instant Ocean (FSIO) and transferred to individual vials containing 4 mL FSIO each. Squid were
transferred to fresh FSIO at 24 hpi. At 46-48 hpi, individual squid were measured for
luminescence using a Promega GloMax 20/20 luminometer before euthanasia via freezing. CFU
per squid was enumerated as described, wherein euthanized squid were homogenized,
dilutions were plated on LBS agar, and total light organ counts were calculated from the count
and proportion of colonies that were plated (43). When assaying colonization-deficient mutants,
some squid in the Δ*lux* treatment group (n = 9) were euthanized and frozen at 24 hpi. Mutant
strain CFU counts were compared to the ES114 treatment group using a Kruskal-Wallis with
Dunn's multiple comparisons test.
376
**Light organ imaging**
*E. berryi* hatchlings were colonized as described above. At 48 hpi, animals were fixed in 4%
paraformaldehyde in 1X mPBS (10 mL DI water, 10 mL 16% paraformaldehyde, and 20 mL 2x
mPBS [marine phosphate-buffered saline: 50 mM phosphate buffer and 0.45 M NaCl [pH 7.4]).
Animals remained in fixative for 48 h at 4˚C. Squid were then washed 4 times for 15 minutes
each at 4˚C with 1X mPBS. Mantles and funnels of fixed squid were dissected away to expose
the light organs of symbiotic animals. Samples were stained overnight in the dark in a
permeabilization solution containing mPBS + 0.5% Triton-X 100. An overnight staining of DAPI
(5 µg/µL) and Alexa647-conjugated phalloidin (25 µg/µL) was used to stain host nuclei and host
F-actin, respectively. The excess stain was removed with 3 washes of mPBS. The tissues were
immersed in a mPBS + 90% glycerol solution and allowed to equilibrate overnight in the dark at
4°C. Samples were transferred to 3 mm depression well slides (United Scientific), mounted

ventral side facing up in the mPBS + 90% glycerol solution, and overlaid with a No. 1.5
coverslip.
391
Qualitative sample images were acquired on a Nikon AXR confocal microscope, illuminated with
405-, 488- and 647-nm lasers and imaged with either a 10X 0.45 NA or 20X 0.80 NA PLAN
Apochromat objective. To reconstruct crypts in 3D and determine the volume occupied by
bacteria, 150 µm deep Z-stacks of one lobe of each animal (n=10) were acquired. The 20X 0.80
NA PLAN Apochromat objective was used, and the laser power and gain were adjusted at
different focal depths using the Z Intensity Correction function (NIS Elements; Nikon) to
normalize signal from the GFP-expressing bacteria across all Z-positions. Image stacks were
imported to and analyzed in arivis Pro (Zeiss), where the 3D crypts were reconstructed via
threshold segmentation of the GFP channel. The resulting segmented 3D objects were used to
determine crypt volume.
402
**Bacterial strain isolation and identification**
Bacterial strains were isolated from *E. berryi* using two distinct methods. In the first method,
luminous *E. berryi* hatchlings raised at the Marine Biological Laboratory (MBL) were euthanized,
frozen, homogenized, and plated onto LBS, providing "*berryi* hatchling" (BH) isolates. As a
second approach, the tank water from a wild-caught adult *E. berryi* ("*berryi* water"; BW) was
collected and serial diluted onto LBS agar. In both approaches, bacterial isolates were streak
purified on LBS, screened for luminescence, and saved as 50% glycerol stocks.
410
Genomic DNA was extracted from all isolates with the Qiagen DNeasy Blood & Tissue Kit.
Samples were submitted to Plasmidsaurus (Eugene, OR) for bacterial genome sequencing
using Oxford Nanopore Technology. Plasmidsaurus results included taxonomic assignment
using sourmash (60), though this did not always provide species-level precision. Isolate species
identity was then determined using FastANI v1.34 (42) against the species suggested by
sourmash and likely candidates of our choosing (Table S1). Organisms belonging to the same
species typically result in >95% average nucleotide identity (ANI). All bacterial isolates were
found to have an ANI >95% for the listed species.
419
**Specific luminescence**
Bacterial cultures were incubated overnight in LBS at 25°C. Luminescence was measured 24 h
later in a clear-bottom black 96-well optical plate using a BioTek Synergy Neo2 plate reader.
Strains were diluted 1:2 in 150 mM NaCl SWTO (61). The plate was shaken for 10 seconds
before luminescence was recorded at a gain setting of 135. The plate was again shaken for 10
seconds and $OD_{600}$ was recorded. Luminescence and $OD_{600}$ values were normalized to blanks.
Specific luminescence was calculated as luminescence-135 divided by $OD_{600}$. Each assay was
performed in triplicate, with two biological replicates conducted.
428
**TUNEL counts and hemocyte trafficking**
*E. berryi* were inoculated as described above for monocolonizations. At 24 hpi, animals were
fixed in 4% paraformaldehyde in 1X mPBS. Animals remained in fixative for 24 h at 4°C before
being washed in 1X mPBS for 30 minutes each a total of 4 times at 4°C. Treatment groups had

a minimum of 13 squid (n = 13-15). Apoptosis was assayed using the DeadEnd Fluorometric
TUNEL Assay kit (Promega) according to manufacturer instructions, as described (22).
Hemocyte counts were taken from the anterior appendage of individual *E. berryi*, adapted from
the method described (62)(24). Briefly, we stained fixed animals with FITC-DnaseI overnight,
counterstained with TOTO-3 to visualize nuclei, and then visualized samples on a Lecia Stellaris
5 CLSM. Due to the non-normality of the data, a non-parametric one-way ANOVA (Kruskal-
Wallis) using Dunn's multiple comparisons test was used to compare each symbiotic group to
the aposymbiotic control group.
441
442  **T6SS2 Assays**
443
444  The BW2 *tssF* mutant was made by introducing a disruption into *tssF_2* in BW2 through
445  triparental mating and *E. coli* carrying the *tssF_2* disruption plasmid pAS2038 (27).
446  Fluorescently-tagged strains were made by triparental mating, introducing pVSV102 for GFP-
447  tagged variants and pVSV208 for dsRed-tagged variants. VipA-GFP expressing strains were
448  made by introducing pSNS119 (27) using triparental mating.
449
To visualize T6SS sheath formation, strains carrying the IPTG-inducible VipA_2-GFP
expression vector (pSNS119) were grown in LBS liquid supplemented with kanamycin overnight
with shaking at 24°C. Cultures were then diluted 1:80 into fresh LBS supplemented with 5% w/v
polyvinylpyrrolidone (PVP) and 0.5 mM isopropyl-β-d-thiogalactopyranoside (IPTG) and grown
to an $OD_{600}$ of 1.0. Two microliters of each culture was spotted onto a 35 mm petri dish with #1.5
glass bottom, and a 2% agarose pad with 0.5 mM IPTG was placed on top of the culture. Cells
were imaged using a Nikon Ti2 inverted fluorescence microscope outfitted with a Hamamatsu
Orca Fusion C1440 and CFI plan apo lambda 100X oil objective lens. Images were captured
using NIS-element software and brightness and contrast was adjusted using FIJI.
459
460  Co-incubation assays were performed between the indicated strain and GFP-tagged ES114 as
461  described (63).
462
463  Competitive colonizations were performed as described for single-strain monocolonizations,
464  except both designated strains were provided as inoculum. Strains were differentiated for colony
465  counting by fluorescent label, using ES114-GFP and ES401-RFP, or the respective ES401
466  derivative with RFP.
467
468  **Data analysis**
469  Plotting and statistical analysis were conducted in GraphPad Prism. Affinity designer was used
470  for figure design.
471
472  **ACKNOWLEDGMENTS**
473
474  We are grateful to Joshua Rosenthal for assistance throughout this project and to Ruth Isenberg
475  for construction of MJM3972. Work in the lab of MJM was supported by NIGMS R35GM148385,
476  in the lab of ANS by NIGMS R35GM137886, in the lab of EACH by NIGMS R35GM150478, and

478
479

480    **Table 1.** List of strains used in this study. Strains are listed by name, and lab stock number is
481    shown in parentheses.

| Strain | Description | Source |
|---|---|---|
| MJ11 (MJM1059) | Natural isolate, *Monocentris japonica* light organ | (48) |
| ES114 (MJM1100) | Natural isolate, *E. scolopes* light organ | (41) |
| CG101 (MJM1115) | Natural isolate, *Cleidopus gloriamaris* light organ | (46) |
| ES213 (MJM1117) | Natural isolate, *E. scolopes* light organ | (64) |
| MB11B1 (MJM1130) | Natural isolate, *E. scolopes* light organ | (65) |
| EM17 (MJM1136) | Natural isolate, *E. morsei* light organ | (66) |
| ES114 Δ*luxCDABEG* (MJM1296) | Non-luminescent ES114 (MJM1100) | (18) |
| ES401 (MJM1318) | Natural isolate, *E. scolopes* juvenile light organ | (46) |
| ES114 Δ*flrA::bar* (MJM3782) | Non-motile ES114, barcoded mutant (MJM1100) | (67) |
| ES114 Δ*sypF::bar* (MJM3972) | Syp-deficient ES114, barcoded mutant (MJM1100) | (68) |
| SA1 (MJM3859) | Natural isolate, *Sepiola affinis* light organ | (49) |
| SR5 (MJM3878) | Natural isolate, *Sepiola robusta* light organ | (49) |
| MJM1100 / pVSV102 (MJM1107) | ES114-GFP (Kan$^R$) | (11) |
| MJM5772 / pVSV102 (MJM5781) | BW2-GFP (Kan$^R$) | This study |

| BH1 (MJM5769) | Natural isolate (*Vibrio campbellii)*, *E. berryi* hatchling | This study |
|---|---|---|
| BH2 (MJM5770) | Natural isolate (*Vibrio fischeri)*, *E. berryi* hatchling | This study |
| BW1 (MJM5771) | Natural isolate (*Vibrio campbellii)*, *E. berryi* tank water | This study |
| BW2 (MJM5772) | Natural isolate (*Vibrio fischeri)*, *E. berryi* tank water | This study |
| BW3 (MJM5773) | Natural isolate (*Vibrio campbellii)*, *E. berryi* tank water | This study |
| BW4 (MJM5774) | Natural isolate (*Vibrio atypicus)*, *E. berryi* tank water | This study |
| BW5 (MJM5775) | Natural isolate (*Vibrio campbellii)*, *E. berryi* tank water | This study |
| BW6 (MJM5776) | Natural isolate (*Vibrio campbellii)*, *E. berryi* tank water | This study |
| BW7 (MJM5777) | Natural isolate (*Vibrio jasicida)*, *E. berryi* tank water | This study |
| BW8 (MJM5778) | Natural isolate (*Pseudoalteromonas flavipulchra)*, *E. berryi* tank water | This study |
| BW9 (MJM5779) | Natural isolate (*Vibrio fischeri)*, *E. berryi* tank water | This study |
| ES401 *tssF_2* (ANS2100) | ES401 *tssF_2/vasA_2* mutant (Erm$^R$) | (69) |
| BW2 *tssF_2* (KB100) | ErmR *tssF_2/vasA_2* mutant (Erm$^R$) | This study |
| BW2 / pSNS119 | BW2 with *vipA_2-gfp* expression vector (Kan$^R$) | This study |
| KB100 / pSNS119 | BW2 *tssF_2* mutant with *vipA_2-gfp* expression vector (Kan$^R$, Erm$^R$) | This study |

| ES401 / pVSV208 | ES401-RFP (Cam$^R$) | (27) |
|---|---|---|
|  |  |  |
| **E. coli strains** |  |  |
| CC118 λpir / pEVS104 (MJM534) | *E. coli* carrying the conjugation helper plasmid for triparental matings. | (70) |
| DH5α λpir / pVSV102 (MJM542) | *E. coli* carrying the GFP-expressing plasmid | (71) |

483
484
485    Table 2: Plasmids used in this study.
486

| Plasmid | Description | Source |
|---|---|---|
| pEVS104 | Conjugation helper plasmid (Kan$^R$) | (70) |
| pVSV102 | Constitutive GFP (Kan$^R$) | (71) |
| pVSV208 | Constitutive DsRed (Cam$^R$) | (71) |
| pAS2038 | *tssF_2/vasA_2* disruption vector (Erm$^R$) | (27) |
| pSNS119 | IPTG inducible *vipA_2-gfp* expression vector (Kan$^R$) | (27) |

487
488    Cam, chloramphenicol; Erm, erythromycin; Kan, kanamycin; GFP, green fluorescent protein;
489    DsRed, Discosoma sp. red fluorescent protein; IPTG, isopropyl-β-d-thiogalactopyranoside.
490
491    Supplemental Table 1: Reference genomes used in FastANI analysis for species identification.
492

| Species | Accession number |
|---|---|
| *Pseudoalteromonas flavipulchra* | ASM25911v1 |
| *Pseudoalteromonas maricaloris* | ASM2474685v1 |
| *Vibrio atypicus* (DSM 25292) | ASM981131v1 |
| *Vibrio campbellii* | ASM290647v1 |
| *Vibrio fischeri* (ES114) | ASM1180v1 |

| *Vibrio fischeri* (MB14A3) | ASM164037v1 |
|---|---|
| *Vibrio fischeri* (MJ11) | ASM2084v1 |
| *Vibrio harveyi* | ASM3006043v1 |
| *Vibrio parahaemolyticus* | ASM19609v1 |
| *Vibrio vulnificus* | ASM222426v1 |

493
494

## REFERENCES

1.   Ruby EG. 2008. Symbiotic conversations are revealed under genetic interrogation. Nat Rev
Microbiol 6:752–762.

2.   Bosch TCG, Guillemin K, McFall-Ngai M. 2019. Evolutionary "experiments" in symbiosis:
The study of model animals provides insights into the mechanisms underlying the diversity
of host-microbe interactions. Bioessays 41:e1800256.

3.   Mandel MJ, Dunn AK. 2016. Impact and influence of the natural *Vibrio*-squid symbiosis in
understanding bacterial-animal interactions. Front Microbiol 7:1982.

4.   Nyholm SV, McFall-Ngai MJ. 2021. A lasting symbiosis: how the Hawaiian bobtail squid
finds and keeps its bioluminescent bacterial partner. Nat Rev Microbiol 19:666–679.

5.   Visick KL, Stabb EV, Ruby EG. 2021. A lasting symbiosis: how *Vibrio fischeri* finds a squid
partner and persists within its natural host. Nat Rev Microbiol 19:654–665.

6.   Ruby EG, McFall-Ngai MJ. 1992. A squid that glows in the night: development of an animal-
bacterial mutualism. J Bacteriol 174:4865–4870.

7.   Jones BW, Nishiguchi MK. 2004. Counterillumination in the Hawaiian bobtail squid,
*Euprymna scolopes* Berry (Mollusca: Cephalopoda). Mar Biol 144:1151–1155.

8.   Boettcher KJ, Ruby EG, McFall-Ngai MJ. 1996. Bioluminescence in the symbiotic squid
*Euprymna scolopes* is controlled by a daily biological rhythm. J Comp Physiol A Neuroethol
Sens Neural Behav Physiol 179:65–73.

9.   Wier AM, Nyholm SV, Mandel MJ, Massengo-Tiassé RP, Schaefer AL, Koroleva I, Splinter-
Bondurant S, Brown B, Manzella L, Snir E, Almabrazi H, Scheetz TE, Bonaldo M de F,
Casavant TL, Soares MB, Cronan JE, Reed JL, Ruby EG, McFall-Ngai MJ. 2010.
Transcriptional patterns in both host and bacterium underlie a daily rhythm of anatomical
and metabolic change in a beneficial symbiosis. Proc Natl Acad Sci U S A 107:2259–2264.

10.  Visick KL, Skoufos LM. 2001. Two-component sensor required for normal symbiotic
colonization of *Euprymna scolopes* by *Vibrio fischeri*. J Bacteriol 183:835–842.

11.  Brooks JF 2nd, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler C,

Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the squid
symbiont *Vibrio fischeri*. Proc Natl Acad Sci U S A 111:17284–17289.

12.  Ruby EG, Asato LM. 1993. Growth and flagellation of *Vibrio fischeri* during initiation of the
sepiolid squid light organ symbiosis. Arch Microbiol 159:160–167.

13.  Millikan DS, Ruby EG. 2003. FlrA, a $\sigma^{54}$-dependent transcriptional activator in *Vibrio*
*fischeri*, is required for motility and symbiotic light-organ colonization. J Bacteriol 185:3547–
3557.

14.  Yip ES, Grublesky BT, Hussa EA, Visick KL. 2005. A novel, conserved cluster of genes
promotes symbiotic colonization and σ-dependent biofilm formation by *Vibrio fischeri*. Mol
Microbiol 57:1485–1498.

15.  Yip ES, Geszvain K, DeLoney-Marino CR, Visick KL. 2006. The symbiosis regulator RscS
controls the *syp* gene locus, biofilm formation and symbiotic aggregation by *Vibrio fischeri*.
Mol Microbiol 62:1586–1600.

16.  Fung BL, Esin JJ, Visick KL. 2024. *Vibrio fischeri*: a model for host-associated biofilm
formation. J Bacteriol 206:e0037023.

17.  Visick KL, Foster J, Doino J, McFall-Ngai M, Ruby EG. 2000. *Vibrio fischeri lux* genes play
an important role in colonization and development of the host light organ. J Bacteriol
182:4578–4586.

18.  Bose JL, Rosenberg CS, Stabb EV. 2008. Effects of *luxCDABEG* induction in *Vibrio*
*fischeri*: enhancement of symbiotic colonization and conditional attenuation of growth in
culture. Arch Microbiol 190:169–183.

19.  Koch EJ, Miyashiro T, McFall-Ngai MJ, Ruby EG. 2014. Features governing symbiont
persistence in the squid-vibrio association. Mol Ecol 23:1624–1634.

20.  Montgomery MK, McFall-Ngai M. 1994. Bacterial symbionts induce host organ
morphogenesis during early postembryonic development of the squid *Euprymna scolopes*.
Development 120:1719–1729.

21.  Koropatnick TA, Engle JT, Apicella MA, Stabb EV, Goldman WE, McFall-Ngai MJ. 2004.
Microbial factor-mediated development in a host-bacterial mutualism. Science 306:1186–
1188.

22.  Troll JV, Adin DM, Wier AM, Paquette N, Silverman N, Goldman WE, Stadermann FJ,
Stabb EV, McFall-Ngai MJ. 2009. Peptidoglycan induces loss of a nuclear peptidoglycan
recognition protein during host tissue development in a beneficial animal-bacterial
symbiosis. Cell Microbiol 11:1114–1127.

23.  Nyholm SV, McFall-Ngai MJ. 1998. Sampling the light-organ microenvironment of
*Euprymna scolopes*: description of a population of host cells in association with the
bacterial symbiont *Vibrio fischeri*. Biol Bull 195:89–97.

24.  Koropatnick TA, Kimbell JR, McFall-Ngai MJ. 2007. Responses of host hemocytes during
the initiation of the squid-Vibrio symbiosis. Biol Bull 212:29–39.

25.  Nyholm SV, Stewart JJ, Ruby EG, McFall-Ngai MJ. 2009. Recognition between symbiotic

*Vibrio fischeri* and the haemocytes of *Euprymna scolopes*. Environ Microbiol 11:483–493.

26.  Visick KL, Ruby EG. 2006. *Vibrio fischeri* and its host: it takes two to tango. Curr Opin
Microbiol 9:632–638.

27.  Speare L, Cecere AG, Guckes KR, Smith S, Wollenberg MS, Mandel MJ, Miyashiro T,
Septer AN. 2018. Bacterial symbionts use a type VI secretion system to eliminate
competitors in their natural host. Proc Natl Acad Sci U S A 115:E8528–E8537.

28.  Mandel MJ, Wollenberg MS, Stabb EV, Visick KL, Ruby EG. 2009. A single regulatory gene
is sufficient to alter bacterial host range. Nature 458:215–218.

29.  Soto W, Rivera FM, Nishiguchi MK. 2014. Ecological diversification of *Vibrio fischeri* serially
passaged for 500 generations in novel squid host *Euprymna tasmanica*. Microb Ecol
67:700–721.

30.  Chavez-Dozal AA, Gorman C, Lostroh CP, Nishiguchi MK. 2014. Gene-swapping mediates
host specificity among symbiotic bacteria in a beneficial symbiosis. PLoS One 9:e101691.

31.  Foster JS, von Boletzky S, McFall-Ngai MJ. 2002. A comparison of the light organ
development of *Sepiola robusta* Naef and *Euprymna scolopes* Berry (Cephalopoda:
Sepiolidae). Bull Mar Sci 70:141–153.

32.  Heath-Heckman E, Nishiguchi MK. 2021. Leveraging short-read sequencing to explore the
genomics of sepiolid squid. Integr Comp Biol 61:1753–1761.

33.  Schmidbaur H, Kawaguchi A, Clarence T, Fu X, Hoang OP, Zimmermann B, Ritschard EA,
Weissenbacher A, Foster JS, Nyholm SV, Bates PA, Albertin CB, Tanaka E, Simakov O.
2022. Emergence of novel cephalopod gene regulation and expression through large-scale
genome reorganization. Nat Commun 13:2172.

34.  Howard RB, Lopes LN, Lardie CR, Perez PP, Crook RJ. 2019. Early-life injury produces
lifelong neural hyperexcitability, cognitive deficit and altered defensive behaviour in the
squid *Euprymna scolopes*. Philos Trans R Soc Lond B Biol Sci 374:20190281.

35.  Crook RJ. 2022. Somatotopic organization of mechanosensory afferents in the stellate
ganglion of the squid, *Euprymna*. https://doi.org/10.1101/2022.10.21.513268. bioRxiv.

36.  Liscovitch-Brauer N, Alon S, Porath HT, Elstein B, Unger R, Ziv T, Admon A, Levanon EY,
Rosenthal JJC, Eisenberg E. 2017. Trade-off between transcriptome plasticity and genome
evolution in cephalopods. Cell 169:191–202.e11.

37.  Jolly J, Hasegawa Y, Sugimoto C, Zhang L, Kawaura R, Sanchez G, Gavriouchkina D,
Marlétaz F, Rokhsar D. 2022. Lifecycle, culture, and maintenance of the emerging
cephalopod models *Euprymna berryi* and *Euprymna morsei*. Front Mar Sci 9:1039775.

38.  Ahuja N, Hwaun E, Pungor JR, Rafiq R, Nemes S, Sakmar T, Vogt MA, Grasse B, Diaz
Quiroz J, Montague TG, Null RW, Dallis DN, Gavriouchkina D, Marletaz F, Abbo L,
Rokhsar DS, Niell CM, Soltesz I, Albertin CB, Rosenthal JJC. 2023. Creation of an albino
squid line by CRISPR-Cas9 and its application for *in vivo* functional imaging of neural
activity. Curr Biol 33:2774–2783.e5.

39.  Sasaki M. 1929. A monograph of the dibranchiate cephalopods of the Japanese and

adjacent waters. Journal of the Faculty of Agriculture, Hokkaido Imperial University 1–357.

40. Sycuro LK, Ruby EG, McFall-Ngai M. 2006. Confocal microscopy of the light organ crypts
in juvenile *Euprymna scolopes* reveals their morphological complexity and dynamic function
in symbiosis. J Morphol 267:555–568.

41. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the
sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

42. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun
9:5114.

43. Naughton LM, Mandel MJ. 2012. Colonization of *Euprymna scolopes* squid by *Vibrio
fischeri*. J Vis Exp e3758.

44. McFall-Ngai MJ, Ruby EG. 1991. Symbiont recognition and subsequent morphogenesis as
early events in an animal-bacterial mutualism. Science 254:1491–1494.

45. Bongrand C, Koch E, Mende D, Romano A, Lawhorn S, McFall-Ngai M, DeLong EF, Ruby
EG. 2022. Evidence of genomic diversification in a natural symbiotic population within its
host. Front Microbiol 13:854355.

46. Ruby EG, Lee KH. 1998. The *Vibrio fischeri-Euprymna scolopes* light organ association:
current ecological paradigms. Appl Environ Microbiol 64:805–812.

47. Xu F, Gonzalez-Escalona N, Drees KP, Sebra RP, Cooper VS, Jones SH, Whistler CA.
2017. Parallel evolution of two clades of an Atlantic-endemic pathogenic lineage of *Vibrio
parahaemolyticus* by independent acquisition of related pathogenicity islands. Appl Environ
Microbiol 83:e01168–17.

48. Ruby EG, Nealson KH. 1976. Symbiotic association of *Photobacterium fischeri* with the
marine luminous fish *Monocentris japonica*; a model of symbiosis based on bacterial
studies. Biol Bull 151:574–586.

49. Fidopiastis PM, von Boletzky S, Ruby EG. 1998. A new niche for *Vibrio logei*, the
predominant light organ symbiont of squids in the genus *Sepiola*. J Bacteriol 180:59–64.

50. Brennan CA, Mandel MJ, Gyllborg MC, Thomasgard KA, Ruby EG. 2013. Genetic
determinants of swimming motility in the squid light-organ symbiont *Vibrio fischeri*.
Microbiologyopen 2:576–594.

51. Brooks JF 2nd, Mandel MJ. 2016. The histidine kinase BinK is a negative regulator of
biofilm formation and squid colonization. J Bacteriol 198:2596–2607.

52. Norsworthy AN, Visick KL. 2015. Signaling between two interacting sensor kinases
promotes biofilms and colonization by a bacterial symbiont: Two interacting kinases permit
biofilm formation. Mol Microbiol 96:233–248.

53. Speare L, Woo M, Dunn AK, Septer AN. 2022. A putative lipoprotein mediates cell-cell
contact for type VI secretion system-dependent killing of specific competitors. MBio
13:e0308521.

54.  Gromek SM, Suria AM, Fullmer MS, Garcia JL, Gogarten JP, Nyholm SV, Balunas MJ.
2016. *Leisingera* sp. JC1, a bacterial isolate from Hawaiian bobtail squid eggs, produces
indigoidine and differentially inhibits vibrios. Front Microbiol 7:1342.

55.  Kerwin AH, Gromek SM, Suria AM, Samples RM, Deoss DJ, O'Donnell K, Frasca S Jr,
Sutton DA, Wiederhold NP, Balunas MJ, Nyholm SV. 2019. Shielding the next generation:
Symbiotic bacteria from a reproductive organ protect bobtail squid eggs from fungal fouling.
MBio 10.

56.  McAnulty SJ, Kerwin AH, Koch E, Nuttall B, Suria AM, Collins AJ, Schleicher TR, Rader
BA, Nyholm SV. 2023. "Failure to Launch": Development of a reproductive organ linked to
symbiotic bacteria. MBio 14:e0213122.

57.  Vijayan N, McAnulty SJ, Sanchez G, Jolly J, Ikeda Y, Nishiguchi MK, Réveillac E, Gestal C,
Spady BL, Li DH, Burford BP, Kerwin AH, Nyholm SV. 2024. Evolutionary history influences
the microbiomes of a female symbiotic reproductive organ in cephalopods. Appl Environ
Microbiol 90:e0099023.

58.  McFall-Ngai M, Heath-Heckman EAC, Gillette AA, Peyer SM, Harvie EA. 2012. The secret
languages of coevolved symbioses: insights from the *Euprymna scolopes-Vibrio fischeri*
symbiosis. Semin Immunol 24:3–8.

59.  Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.
Nat Methods 9:676–682.

60.  Titus Brown C, Irber L. 2016. sourmash: a library for MinHash sketching of DNA. J Open
Source Softw 1:27.

61.  Bose JL, Kim U, Bartkowski W, Gunsalus RP, Overley AM, Lyell NL, Visick KL, Stabb EV.
2007. Bioluminescence in *Vibrio fischeri* is controlled by the redox-responsive regulator
ArcA. Mol Microbiol 65:538–553.

62.  Altura MA, Heath-Heckman EAC, Gillette A, Kremer N, Krachler A-M, Brennan C, Ruby
EG, Orth K, McFall-Ngai MJ. 2013. The first engagement of partners in the *Euprymna
scolopes-Vibrio fischeri* symbiosis is a two-step process initiated by a few environmental
symbiont cells. Environ Microbiol 15:2937–2950.

63.  Speare L, Septer AN. 2019. Coincubation assay for quantifying competitive interactions
between *Vibrio fischeri* isolates. J Vis Exp https://doi.org/10.3791/59759.

64.  Boettcher KJ, Ruby EG. 1994. Occurrence of plasmid DNA in the sepiolid squid symbiont
*Vibrio fischeri*. Curr Microbiol 29:279–286.

65.  Wollenberg MS, Ruby EG. 2009. Population structure of *Vibrio fischeri* within the light
organs of *Euprymna scolopes* squid from Two Oahu (Hawaii) populations. Appl Environ
Microbiol 75:193–202.

66.  Nishiguchi MK, Ruby EG, McFall-Ngai MJ. 1998. Competitive dominance among strains of
luminous bacteria provides an unusual form of evidence for parallel evolution in Sepiolid
squid-vibrio symbioses. Appl Environ Microbiol 64:3209–3213.

67. Burgos HL, Burgos EF, Steinberger AJ, Suen G, Mandel MJ. 2020. Multiplexed competition
in a synthetic squid light organ microbiome using barcode-tagged gene deletions.
mSystems 5:e00846–20.

68. Vander Griend JA, Isenberg RY, Kotla KR, Mandel MJ. 2024. Transcriptional pathways
across colony biofilm models in the symbiont *Vibrio fischeri*. mSystems 9:e0081523.

69. Speare L, Smith S, Salvato F, Kleiner M, Septer AN. 2020. Environmental Viscosity
Modulates Interbacterial Killing during Habitat Transition. MBio 11:e03060–19.

70. Stabb EV, Ruby EG. 2002. RP4-based plasmids for conjugation between *Escherichia coli*
and members of the *Vibrionaceae*. Methods Enzymol 358:413–426.

71. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp*- and pES213-derived
tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in
situ. Appl Environ Microbiol 72:802–810.