A systematic screen for genetic factors underpinning transposon defense systems across

the fungal kingdom

Thomas Badet [1][*] & Daniel Croll [1][*]

[1] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel,

Switzerland

[*] Correspondence: daniel.croll@unine.ch & thomas.badet@unine.ch

Keywords: repeat-induced point mutation, fungal genomics, transposable elements, genome evolution

Short title: Transposon defense system screen across fungi

Competing interests: The authors have declared no competing interest.

Author contributions: TB conceived the study and performed analyses with input from DC; DC

provided funding; TB and DC wrote the manuscript. All authors have seen and approved this version

of the manuscript.

Data availability: The data analyzed in the frame of this study are available from the NCBI

repositories as indicated in the Supplementary Table S1.

Funding: DC was supported by the Swiss National Science Foundation (grants 173265 and 201149).

Acknowledgements: We thank group members for fruitful discussions.

Abstract

Fungal genome sizes exhibit more than a 100-fold variation, largely driven by the expansion of repetitive sequences such as transposable elements (TEs). Genomic defenses have evolved to counteract TE proliferation through silencing mechanisms operating at the epigenetic or transcript level. In fungi, repeat-induced point mutation (RIP) targets TEs by recognizing repetitive sequences and inducing mutagenesis. However, the prevalence of RIP across the fungal kingdom and the fidelity of the canonical C-to-T mutation signatures remain unclear. In this study, we address these gaps by tracking shifts in genome architecture across the fungal kingdom. We show that the loss of RIP components is associated with an 80% (~30 Mb) increase in genome size in ascomycetes (class of Leotiomycetes). To track the impact of genome defenses, we designed a quantitative screen for RIP-like mutation signatures. The phylum of ascomycetes was unique in showing enrichment in mutation signatures in non-coding and repetitive sequences, consistent with a phylogenetically restricted occurrence of RIP-like genome defense systems. Then, we gene functions associated in a phylogenetic context with RIP-like mutation signatures. We identified a zinc-finger protein as the strongest candidate underpinning a novel mechanism of genome defenses. Our findings reveal the multifaceted drivers of genome defense systems and their close ties to genome size evolution in fungi, highlighting how proximate molecular mechanisms can shape genome evolution on deep phylogenetic scales.

Introduction

Mechanisms underlying variation in genome size across eukaryotes have attracted significant attention ever since the discovery of the C-value paradox and the extent of non-coding DNA in most genomes [1]. With increasing genome size, the non-coding fraction becomes the dominant constituent of the genome [2]. Non-coding DNA is composed in large parts of transposable elements (TEs) capable of transposing or creating additional copies in genomes [3]. TE content variation is the primary source of genome size variation among species [4]. Beyond inflating genome size, active TEs can impair fitness through deleterious insertions and destabilize genome integrity [5,6]. Genomes evolved multiple defense systems that act post-transcriptionally with the primary aim of controlling the expression of the genes required for TE mobilization [7,8]. Such genomic defense systems are widespread among eukaryotes and are based on DNA or histone methylation, as well as RNA interference (RNAi) [9–11]. However, to what extent genomic defenses effectively contain TE proliferation remains poorly understood.

Despite active genome defense systems, many fungal lineages experienced genome expansions and TEs recurrently escaped from host genome control [12–14]. Variation in repeat content and TE activity have also been found within individual species highlighting the potential rapid turnovers in defense mechanisms [15,16]. Fungi present a highly heterogeneous clade for mechanisms and effectiveness of genome defenses. Most species in the Saccharomycota subphylum (including the Baker's yeast *Saccharomyces cerevisiae*) have small and compact genomes nearly devoid of repeats despite lacking DNA methylation and RNAi systems [17–19]. Small genomes have been associated with the colonization of nutrient-rich environments and the capacity to develop as single-celled yeasts [20,21]. However, yeast-forming fungi have emerged multiple times independently across the fungal kingdom and show convergent loss of important metabolic genes and genome streamlining [21–24]. Host-associated fungi also tend to have smaller genomes than their free-living relatives. The genomes of obligate fungal parasites for instance have been shown to harbor small genomes, suggesting that lifestyle might be a major driver of genome evolution [25,26]. Similar observations were made for obligate endosymbiotic bacteria showing reduced effective population size and inefficient selection [27–29]. Therefore, species with small effective population size are predicted to evolve larger genomes as a consequence of ineffective excision of mildly deleterious insertions such as those induced by TE mobilization [27,30]. However, several species including plant pathogenic fungi experienced TE-mediated genome size increases despite the assumption of large effective population sizes [31–33]. Hence, current models fail to predict genome size evolution with TE defenses likely being a poorly modeled factor.

With few exceptions, most fungi have retained the machinery necessary for DNA and histone methylation as well as RNAi-associated pathways [17,18,34]. These widely shared mechanisms of TE

defense are complemented in fungi by specific mechanisms such as targeted mutagenesis of repeated
DNA sequences [35]. This process, known as repeat-induced point mutation (RIP), has been first
described in *Neurospora crassa* and is responsible for the highest known mutation rate outside viruses
[36,37]. In fungal plant pathogens, RIP-mediated mutation rates underpinned rapid evolution of
pathogenicity genes located near repeats [38]. RIP activity likely contributed to the emergence of
genome compartments with contrasted TE and gene contents [39]. A major cost of the RIP genome
defense mechanism is to hinder the evolution through gene duplication. In the *N. crassa* genome,
most paralogs predate the emergence of RIP activity [40]. RIP is thought to be active in the pre-
meiotic cell containing both parental nuclei and promotes C→T mutations in targeted repeats in a
CpA/TpG dinucleotide context [41,42]. RIP efficiency is influenced by the length and identity of the
matched repeat sequences, as well as by the periodicity of interspersed homology between the repeats
[43–46]. RIP is controlled by two cytosine DNA methylases and a histone methyltransferase that
segregate presence absence in the phylum [47–49]. Many genomes of ascomycete fungi show
signatures of RIP in repetitive sequences, however what underpins the patchy distribution remains
unknown. The trade-off between genome defense benefits and costs for paralog creation are likely
creating complex dynamics.
99
100 In this study, we tested the hypothesis that RIP activity prevents genome expansion in fungi. For that,
101 we first determined shifts in genome architecture across 1,239 genomes covering the fungal kingdom.
102 We then tested whether gains or losses of the RIP machinery predicts TE proliferation rates and
103 genome size evolution. Finally, we designed a quantitative screen for RIP-associated mutation
104 signatures to predict previously unknown genetic determinants of RIP activity.
105

Results

Genome evolution in the fungal kingdom

To determine the action and signatures of genome defense systems in the fungal kingdom, we sampled a total of 1,239 genomes representing five major phyla in addition to 16 outgroup species belonging to the Oomycota and Stramenopiles (Supplementary Table S1). The number of assembled scaffolds ranged from 4 to 83,551 for an average of 1,211 while genome size was on average 35 Mb ranging from 7.3 Mb (*Malassezia restricta*) to 733 Mb (*Gigaspora margarita*) (Figure 1A). Assembly completeness ranged from 46% BUSCO completeness (*Botryozyma nematodophila*) to 100 (eight Stramenopiles), for an average of 95% across all phyla (Supplementary Table S1). Larger genomes contain larger fractions classified as repeats (Figure 1B and D) highlighted by the >100 Mb genomes in species from the Mucoromycota. We identified 21 species for which more than 50% of the assembly is made of repetitive DNA for a genome size ranging from 71,5 to 773 Mb (top 5.2% larger genomes, Supplementary Table S1-S2). Genomes from the Saccharomycota are small and mostly devoid of repeats, with a repetitive fraction accounting for 0-20% of the genome (average of 2%). We computed for each species the number of sequences annotated as repeats and that share strong identity (>95% sequence identity over intervals ranging from 100 bp to 10 kb, which we term "repeat identity"; see Figure 1C). As expected for species with few or no repeats, 75% of the Saccharomycotina genomes have less than 10 sequences larger than 1 kb sharing >95% identity (265/353 species). In contrast, 83-98% of the species in the Agaricomycetes, Dothideomycetes, Eurotiomycetes, Sordariomycetes or Leotiomycetes have more than 10 genomic repeats sharing 80-95% identity over 1 kb in sequence length (Supplementary Table S3).

To investigate the evolution of genome defense mechanisms in a phylogenetic context, we sampled 100 BUSCO genes with a species occupancy ≥50% and built a phylogenetic tree for the phylum (Figure 1E, Methods). Using phylogenetic independent contrasts, we find that most of the measures of assembly contiguity are highly correlated (Figure 1F). To prevent spurious associations with genome assembly quality, we assessed phylogenetic signals for twelve genome assembly metrics. All the tested assembly metrics show some level of phylogenetic signal. Applying a local indicator of phylogenetic association, we find that ~79% of the species show phylogenetic signal for at least one of the genome assembly or architecture metrics (977 / 1239, Supplementary Figure S4; Table S4). However, we find that the phylogenetic signal is evenly distributed along the species tree, suggesting no major taxonomy driven bias in genome assembly quality. We find that genome size strongly correlates with protein numbers and GC dinucleotide content (r = 0.53 and 0.57, respectively). We computed genome segmentation metrics based on the number of segments with distinct GC content. We find that genome size and segmentation are highly correlated with larger genomes being more segmented (r =0.76, Figure 1F). Using the 3' and 5' intergenic distance as a measure of gene density,

we show that ~12.5% of the genomes show compartmentalization as defined by stretches of

contrasted GC content (Supplementary Figure S2). In addition, we find that the proportion of the total

gene pool found in gene-sparse regions (>5 kb at both 3' and 5' ends) is strongly positively correlated

with genome size and genome segmentation (Supplementary Figure S3).

Shifts in genome size associated with lifestyle changes

To dissect drivers of genome evolution, we analyzed shifts in genome architecture in a phylogenetic

context. We parametrized a Brownian motion model of trait evolution assuming monotonous and

heterogeneous evolutionary rates without *a priori* information on the number or position of rate shifts

(*i.e.* Ornstein–Uhlenbeck process). Analyzing five metrics of genome architecture, we identified a

total of 229 shifts across 150 edges of the phylogeny (148 / 229, *i.e* ~65%, Figure 2AB). The highest

number of shifts across the phylogeny were associated with genome size and lifestyle changes, while

most intense shifts were found for the number of highly similar repeat sequences (*i.e.* repeat identity

metric) and the genome segmentation metric (Figure 2AB). Overall, 58% of the shifts co-occurred

with at least one other metric of genome architecture (134 / 229; Figure 2BD). Shifts in genome size

often co-occurred with changes in lifestyle (Figure 2BD, Supplementary Table S5). Nearly 57% of the

shifts are in terminal branches of the tree including four co-occurring shifts at the terminal branch of

Colletotrichum spinosum with a decrease in genome size and repeat identity being associated with an

increase in GC content (Figure 2B, Supplementary Table S5). Shifts tend to occur closer to the tree

tips than in preceding non-terminal edges (Wilcoxon rank sum test *p*-value = 9.34e-12,

Supplementary Figure S5). One notable exception was the ~30 Mb increase in genome size internal to

the order of the Leotiomycetes with 29 offspring tips (Figure 2E). Following this large increase, six

non-terminal offspring branches show shifts in genome architecture, including genome size, repeat

identity and GC content. Five shifts in genome size, lifestyle and repeat identity mapped to a clade of

plant pathogens of the *Rhynchosporium* genus suggesting that the gain of the pathogenic lifestyle

might have been facilitated by changes in genome architecture (Figure 3A).

Furthermore, we found that the 30 Mb genome increase in the Leotiomycetes co-occurred with a loss

of the major determinant of RIP (*i.e. Rid1*; Figure 3A). Ancestral state reconstruction revealed indeed

that *Rid1* was likely absent at the onset of the genome size increase (Supplementary Figure S6). We

found that both the maintenance-type DNA methyltransferase Dnmt5 and the *de novo* DNA

methyltransferase Dim2 were lost in the same clade of 29 species (Figure 3A). We confirmed the loss

of the RIP-essential cytosine methyltransferase in the enlarged genome of *Blumeria graminis f. sp.*

tritici by analyzing the *Rid1* locus of a genome from the sister genus *Cadophora* (Figure 3B). We

identified synteny relationships for multiple protein coding genes surrounding the *Rid1* sequence in

the *Cadophora* genome matching a single scaffold in the *Blumeria graminis* f. sp. *tritici* assembly.

Most of the genes surrounding *Rid1* were conserved among the genomes, with the locus *in B.*

graminis f. sp. tritici clearly lacking the RIP-essential gene (Figure 3B). Across the fungal phylogeny,

Rid1 and *Dnmt5* are encoded in mostly conserved loci. In contrast, the *Dim2* locus is poorly

conserved in agreement with the recurrent gene loss across the phylogeny (Figure 3C, Supplementary

Table S6).

Genes likely to impact RIP were lost in fungi

RIP is controlled by both DNA and histone methylation. We analyzed similar functions across the

fungal kingdom to assess potential correlates with genome size shifts. The full set of 12,837,225

annotated proteins were clustered into 933,315 orthogroups (Figure 4A; Supplementary Tables S7-9).

Overall, 79% of the orthogroups were restricted to single species (*i.e.* singletons), and ~20% were

present in less than 50% of the species (*i.e.* variable set). Only 2,434 core orthogroups (0.2%) were

present in at least 90% of the species making up ~12-68% of each species proteome (Figure 4A). We

next investigated the taxonomic distribution of eight orthogroups encoding proteins known to affect

RIP in *N. crassa*. The heterochromatin protein 1 (Hp1) and the damage-specific DNA binding protein

1 (Ddb1) are highly conserved in fungi with each being present in >90% of the species (Figure 4B).

The histone H3 lysine methyltransferase Dim5 and the ubiquitin ligase components Cullin4 (Cul4)

have homologs identified in 72% and 66% of the species, respectively. The two cytosine DNA

methyltransferases Rid1 and Dim2 have homologs in only 48% and 51% of the species. Dim2 is

nearly absent in Saccharomycetes and Eurotiomycetes and Rid1 is absent in Saccharomycetes (Figure

4B). Species carrying Rid1/Masc1 homologs have more segmented and larger genomes, as well as

lower repeated sequence identity (Figure 4C). Repeat-targeted mutations are linked to DNA and

histone modifications, DNA repair machinery and sexual recombination [50,51]. For an exhaustive

view of relevant molecular machineries present across fungi, we used protein family annotations to

identify proteins related to DNA methylation, heterochromatin formation, RNA interference, DNA

repair and meiotic recombination (Supplementary Table S10). A cytosine DNA methylase domain is

present in 871 genomes and mostly absent from *Saccharomycota* and the *Oomycota* outgroup (Figure

4D, PF00145.18). Proteins with domains associated with DNA repair were found consistently across

most genomes, in concordance with their important role (Supplementary Figure S8). Histone-related

proteins are highly conserved across fungi, with the exception of the histone-lysine N-

methyltransferase E(z), which is missing in species from the *Eurotiomycetes* (Figure 4D,

Supplementary Figure S8). The C2H2 type master regulator of conidiophore development brlA

involved in RNA interference was exclusively found in *Eurotiomycetes* and *Sordariomycetes*.

Furthermore, all ascomycetes lack the RNA exonuclease acting as a negative regulator of RNA

interference (Figure 4D, ERI1). The pool of protein functions showing overlaps with known

components of RIP is taxonomically restricted and shows a patchy distribution.

Widespread nucleotide composition enrichment in repetitive sequences

Mutagenic mechanisms targeting repeats such as TEs should leave strong signatures in repetitive, non-coding sequences. We designed a screen for mutation signatures against TEs, which is agnostic of the specific sequence patterns generated by the mutagenic mechanism. Our approach started by segmenting each genome into coding, non-coding and repeated sequences to assess the abundance of 336 specific *k-mers* (including 16 dimers, 64 trimers and 256 tetramers; Figure 5A). We computed *k-mer* frequencies only for full genome assemblies and after filtering for scaffolds larger than 50 kb to reduce artefacts (Supplementary Figure S9). We found that nearly all species have at least one *k-mer* that is >2-fold enriched in non-coding sequences compared to coding sequences (1,027 / 1,239 species, Figure 5C, Supplementary Table S11). A total of 152 tetramers were overrepresented in non-coding sequences with a >2-fold enriched in 1,027 species (Supplementary Figure S10). We identified 26 *k-mers* with >10-fold enrichment and these *k-mers* were found across 53 species grouped into seven taxonomic classes (Supplementary Figure S11-S12). Finally, we identified a set of 10 *k-mers* >2-fold enriched in non-coding sequences in ~40% of all analyzed species. This broadly enriched set of 10 *k-mers* were all AT-rich tetramers consistent with sequence signatures generated by RIP mutations (*i.e.* CpA dinucleotide contexts).

To expand the screen for RIP-like repeat-directed genome defense mechanisms, we analyzed hallmarks of the rapid loss of sequence identity as expected for heavily mutated TEs. For this, we contrasted TE sequences against the entirety of the non-coding sequences (Figure 5B, Supplementary Table S11). We found that in most species the *k-mer* frequency at repeats largely reflects the frequency at non-coding sequences, suggesting that for most genomes the repeated sequences are not targeted by an active RIP-like mechanism (Figure 5C; Supplementary Figure S13). However, we also identified 68 *k-mers* with a >2-fold enrichment in non-coding sequences, as well as a >2-fold enrichment in TEs compared to all non-coding sequences (Figure 5D, Supplementary Table S11). In species of the Dothideomycetes, Eurotiomycetes, Leotiomycetes and Sordariomycetes, AT-rich *k-mers* were found even >5-fold enriched in repeats compared to all non-coding sequences (Figure 5C, Supplementary Figure S14). In addition, ~60% of the species with a strong *k-mer* enrichment in repeats carry only few large sequences with >95% identity. This matches expectations for a RIP-like mechanism driving the degeneration of repeat sequences (279/466, average number of >5kb sequences with >95% identity = 23). Interestingly, species with large and repetitive genomes carry on average more repeats of high sequence identity, consistent with recent TE proliferation unopposed by a RIP-like mechanism (Supplementary Figure S15).

We assigned each species as being putatively RIP-proficient if the genome showed a proportion of *k-mer* enriched >2-fold in non-coding and repeat regions larger than 0.5 (Supplementary Table S12). Using a threshold of least 10 *k-mers* with a 2-fold enrichment in non-coding regions and repeats, we found 130 species showing strong signatures of recent episodes of repeat-directed mutations (Figure

5E, 22 Dothideomycetes, 45 Eurotiomycetes, 50 Sordariomycetes and 13 Leotiomycetes, respectively). We identified 105 genomes with enriched *k-mers* in non-coding regions and repeats at a ratio of 0.3-0.5. Such genomes experienced most likely only trace activity of a RIP-like mechanism, and the mechanism ceased activity long ago. Using these discrete categories for evidence of RIP-like activity, we found that Ascomycota genomes with trace or ancient RIP activity tend to be smaller than genomes with no traces of RIP (Supplementary Figure S16; Tukey's HSD Test for multiple comparisons $p < 9.3e-7$). We also computed pairwise protein sequence identity per genome and found that genomes with trace or ancient RIP activity encoded fewer proteins with recent duplicates (Supplementary Figure S17). In conjunction, our results strongly suggest that RIP-like genomic defenses share a preference for CpA dinucleotides and were active in only few fungal clades. Furthermore, RIP-like defenses tend to be associated with smaller genomes and lower protein sequence identity suggesting effective defenses against genome expansion at the cost of constraining gene duplications.

RIP-like mutations associated with DNA-binding proteins

We conducted phylogenetically informed association analyses to identify strong candidates for yet unknown components of RIP-like mutation machinery. For this, we associated individual orthogroups with variation in genome size, genome segmentation, lifestyle and *k-mer* enrichment across the phylogeny. Using three distinct methods, we identified a total of 1,348 orthogroups with significant associations (Supplementary Table S13). We conservatively focused on orthogroups with association supported by at least two methods. By this, we found a total of four genome characteristics and 112 associated orthogroups (Figure 6A). All associations were exclusive to individual genomic characteristics with fungal lifestyle accounting for most of the associations (63 orthogroups, Figure 6A). Lifestyle-associated orthogroups encode lipase, cupin and hydrolase activity matching expectations for their niche adaptation (Figure 6B). Most associated orthogroups reflect the species phylogeny and trophic lifestyles (Figure 6C, Supplementary Figure S18). Strikingly, many associated orthogroups were missing from the Saccharomycota species and other monomertrophs in other clades (Supplementary Figure S18). Features of genome segmentation associated with 13 orthogroups found in 62-110 ascomycetes, with the majority belonging to the class of Sordariomycetes, Leotiomycetes and Dothideomycetes (136, 50 and 44 species respectively; Figure 6C). The orthogroups include a total of 1,232 proteins with largely unknown functions with the exception of four orthogroups encoding mostly SKG6, PAN, zinc-finger and CFEM domains (Figure 6B). A further 13 orthogroups are associated with variation in genome size (Figure 6AC). About 20% of all protein sequences included in orthogroups associated with genome size encode AAA+ ATPase domains, which are often identified as TE mobilization proteins. This further supports the strong correlation between genome size and TE activity (Supplementary Table S14).

The strongest association of mutation signature enrichment in repeats was found for a single

orthogroup (OG0000460) shared by Ascomycetes (eta-squared = 0.4; Figure 6C-D). The orthogroup

is shared by 598 species (n = 2,853 proteins; ~97% of Ascomycota excluding Saccharomycota). The

most abundant domain encoded by these proteins is a DNA binding bZIP domain (Figure 6B).

Furthermore, OG0000460 is associated with the strongest *k-mer* enrichment at repeats as well a larger

and more segmented genomes (Supplementary Figure S19). In addition, we found that a clade of six

species including the plant pathogen *Blumeria graminis* f. sp. *tritici* with genome size variation of 41-

140 Mb lack between 1-3 genes associated with variation in genome size (Figure 3A and 6AC). In *N.*

crassa, OG0000460 includes two highly conserved proteins of 260 and 379 amino acids, respectively

(Figure 6E). The best structural match for the two *N. crassa* proteins were found with A0A2J8CDV3

(5 members), which includes five members from the fungal kingdom (A0A1V1TG90) in addition to a

protein encoded by the plant *Digitaria exilis* (*Poaceae* family). Our results show that no single gene

likely controls genome size evolution in fungi. Furthermore, we find that Ascomycetes harbor

numerous clades with strong evidence for yet undescribed RIP mechanisms underpinning convergent

evolution in genome defenses.

Discussion

Eukaryotic genome size variation is most likely driven by TE activity [52–54]. Here, we showed that

genome size evolution can be associated both with the presence of specific orthogroups and sequence

signatures of recent RIP activity. Most fungal genomes showed strong *k-mer* compositional

enrichment at non-coding sequences compared to coding sequences. However, only a third of the

genomes showed a concomitant compositional enrichment both in non-coding and repeated

sequences, suggesting that mechanisms of repeat-driven mutations are not conserved across fungi. In

Ascomycetes, RIP targets repeated sequences to promote C-to-T transitions [43,55]. Consistent with

RIP, we found that all repeat-enriched *k-mer* are AT-rich and reminiscent of RIP signatures. Also, we

found that species with repeat-enriched *k-mers* were almost exclusively Ascomycetes, and this

phylum also encodes the clearest molecular toolset for RIP. We used our quantitative measure of

mutation bias to pinpoint a gene with a strong positive association to AT-rich *k-mers*. Finally, we

recapitulated a 30-Mb increase in genome size in Leotiomycetes 120 Mya ago that was likely

facilitated by the concurrent loss of multiple RIP-related genes.

Bursts in TEs are often attributed to the inactivation of genomic defenses and the de-repression of

active elements [56–58]. Here, we show that RIP activity as observed in *N. crassa* is likely restricted

to a small number of Ascomycetes. Furthermore, we show that *N. crassa* carries sequence signatures

consistent with relict RIP activity, because signatures mostly mapped to non-coding sequences

excluding repetitive DNA. Such a mutation bias is most likely explained by weak or dormant TE
activity in the species. Using these sequence signatures, we show that most species previously
reported to harbor RIP-like mutations do not show biased nucleotide compositions in repeats. For
instance, we found no evidence for recent RIP activity in repeats of the TE -rich genome of the barley
pathogen *Pyrenophora teres f. teres* with *ca.* 30% of the genome predicted to be RIP-affected [59].
Furthermore, the *Pyricularia grisea* lineage carries signatures of ancient RIP activity but no such
evidence in the closely related *P. oryzae*. Even though sexual recombination and RIP-like mutations
were documented in *P. oryzae*, most recent divergence occurred through clonal reproduction, which is
expected to disable RIP due to the lack of meiosis. This is supported by a recent study showing that
recombining and non-recombining populations of the blast fungus *P. oryzae* differ in RIP-like
mutation accumulation [60,61]. The *P. oryzae* example illustrates the limitations of using mutational
signatures within single genomes to infer the occurrence of RIP in a species.

The paradigm for the genes required for RIP is confirmed in seven species, including *N. crassa*,
supporting the existence of a universal toolbox underpinning the genome defense mechanism [60,62–
66]. We find however that near 50% of the species lack one of the two important DNA
methyltransferases. The cytosine methyltransferase Dim2 and adenine methylation were lost in most
Eurotiomycetes. Another cytosine methyltransferase, Dnmt5, was lost in most Sordariomycetes,
including *N. crassa* [17,48]. Among the species with experimental evidence for RIP, two most likely
encode Dnmt5 (*i.e., P. anserina* and *L. maculans*) showing both weaker RIP signatures [62,66].
Interestingly, most experimental evidence for RIP strength outside of *N. crassa* indicated less
effective RIP mutations, in particular in unlinked repeats [60,62–66]. In *P. anserina,* premeiotic
recombination and RIP are much more frequent in a deletion mutants with delayed fruiting-body
development, further highlighting the differences in RIP activity among fungi [67]. As an example,
the DNA methyltransferase Rid1 is essential for RIP but also for normal sexual development in *P.*
anserina but not in *N. crassa* [47,68–71]. This is highlighted by the fact that in *P. anserina* the mild
RIP-like mutations observed upon sexual recombination are not independent from cytosine
methylation of the targeted repeats in contrast to *N. crassa* [66,70].

Our kingdom-wide screen reveals complex dynamics between genome defenses and genome size in
fungi. By analyzing over a thousand fungal genomes, we demonstrate that genome size variation is
intricately linked to TE accumulation and genome compartmentalization. While genes essential for
genome defense against TEs are frequently lost across the fungal kingdom, this loss is not consistently
associated with major TE-driven genome expansion. Instead, ecological factors appear to be the
primary drivers of genome evolution, as most changes in genome size coincide with lifestyle
transitions. Within Ascomycota, ancient signatures of RIP activity correlate with smaller genomes,
indicating effective control against genome inflation. This kingdom-wide perspective underscores that

while RIP mutational signatures are strongly associated with Ascomycetes, RIP presence and
effectiveness are highly dynamic within the phylum. This highlights the value of comparative
genomic studies in uncovering the interplay between genome defense mechanisms and genome
architecture. Integrating intraspecific pangenomes into this approach presents further opportunities to
associate TE dynamics with genome evolution. Evolve-and-resequence experiments may reveal host
genome responses to TE activity and short-term effects of selection on genome defenses mechanisms
[72,73].

Methods

Fungal genomes and annotation procedure

The genomes of 1,342 species were retrieved from two different sources (Supplementary Table S1). The yeast genomes and their respective gene annotation were taken from the study https://doi.org/10.1016/j.cell.2018.10.023 (available on figshare at https://figshare.com/articles/dataset/Tempo_and_mode_of_genome_evolution_in_the_budding_yeast _subphylum/5854692). All other published fungal and Oomycota genomes were retrieved from NCBI together with their respective gene annotation. We calculated assembly statistics using the *stat* function from the *pyfastx* tool version 0.7.0 [74]. As an additional genome metric, we computed variation in GC content across the assemblies using the *GC-Profile* method [75]. Using a segmentation threshold of 100 and a minimum length of 200 bp, we estimated genome segmentation by counting the number of segments identified by GC-Profile.

Inference of species trophisms

We categorized species according to their trophic mode as developed by Hane et al [76]. Using gene annotation of the 1,239 individual species genomes, we searched for carbohydrate-degrading enzymes (CAZymes; dbCAN version 10 [77]). We used the *hmmscan* function from the HMMER package version 3.3.2 to identify CAZYmes in each of the proteomes using the dbCAN hidden Markov models as queries [78]. We then applied the *CATAStrophy* algorithm to classify each species trophism given its CAZymes pool. For each proteome we also extract the fitted CATAStrophy principal component values based on its CAZymes frequencies. The first two axes of the principal component analysis were used as a quantitative measure of trophic lifestyles.

Species phylogeny reconstruction

For the species tree reconstruction, we followed a similar method as in [79]. Briefly, we first identified a set of single-copy orthologous genes in each of the 1,342 genomes using *BUSCO* version 4.1.4 [80] searching the Fungi or Oomycota orthology database version 10, respectively. The method identified a maximum set of 756 BUSCO genes in the genome of the fungus *Colletotrichum plurivorum*. Using a minimal threshold of 50% gene occupancy (> 378 BUSCO genes), we proceeded with 1,263 individual species. BUSCO genes were then translated into protein sequences respecting the species' genetic code (code 12 for Saccharomycota species except for *Pachysolen tannophilus* for which code 26 was used and code 1 for all other species). A random sample of 100 of the resulting protein sequences were then concatenated using the geneStitcher.py script (https://github.com/ballesterus/Utensils) and aligned using the *mafft* alignment tool (version v7.475, --maxiterate 1000 –auto parameters) [81]. The resulting alignment was trimmed using trimal v1.4.rev15 with the -gappyout option [82]. We estimated the best molecular evolutionary models for the

concatenated 100 protein sequences using *partitionfinder* version 2 [83] with the quick option (-q)
while using *RAxML* for the analysis version 8.2.12 [84]. The resulting partitioned model was then
applied for phylogenetic inference using iqtree2 version 2.1.2 after 1,000 replicates for ultrafast
bootstrap and 2 independent runs (-B 1000 --runs 2) [85]. Finally, the tree was rooted using *Achlya*
hypogyna as the oomycete outgroup species with the *root* function in the ape R package v 5.7-1 with
resolve.root = TRUE [86].
412

K-mer **frequency estimates**
We computed the frequencies of all possible 336 *k-mer* with size 2, 3 or 4, across all 1,239 fungal
genome assemblies. First, we used the gene annotation available for each genome assembly to define
coding and non-coding compartments. Using the *subtractBed* function from the *bedtools* suite v2.26.0
[87], we extracted coordinates of all the intergenic segments for each individual assembly ("gene"
was used as the feature to subtract, leaving introns as part of the coding fraction of the genome). Next,
we masked the genome assemblies using either the genic or intergenic coordinates (*maskfasta*
function from the *bedtools* suite). We then counted *k-mer*s in either the gene-masked and intergenic-
masked assemblies using *jellyfish* 2.3.0 [88]. To account for variation in assembly contiguity, we also
counted *k-mer*s after filtering out the scaffolds smaller than 50 kb in length using the *seqtk seq -L*
function, version 1.4-r122 [89]. Finally, to account for the size of the coding and non-coding
compartments in each genome, we normalized the frequency of each *k-mer* by the sum of all *k-mer*
counts in their respective compartment. For each *k-mer*, its non-coding enrichment was represented by
the frequency in non-coding sequences over its frequency in coding sequences (*i.e.,* the *k-mer* is
defined as enriched in non-coding sequences with values > 1).
428

Repeat identification and annotation
To identify putative repetitive elements, we used *RepeatModeler* v2.0.1 using rmblast v2.10.0 in
combination with *LTR_Harvest* and *LTR_retriever* (-engine ncbi and -LTRStruct options) [90–92].
For five small genomes we reduced the sampling size (-genomeSampleSizeMax option) to 8100000
or 810000 (Kazsaulg = 8100000, Canathen = 8100000, Malrestr = 8100000, Psehubei = 8100000,
Kazaerob = 810000, respectively). Using the consensus sequences identified in each genome, we next
annotated repeats with *RepeatMasker* version 4.1.5 using a cutoff value of 250 and skipping bacterial
insertion element (-cutoff 250 and -no_is options) [93]. The resulting repeat annotation was filtered
for simple_repeats and low_complexity regions, interrupted repeats were merged using the helper
script *parseRM_merge_interrupted.pl* and annotations were finally converted to gff3 format using the
rmOutToGFF3.pl script (https://github.com/4ureliek/Parsing-RepeatMasker-Outputs). Based on the
RepeatModeler classification, we assigned each putative repeat to either of five major families,
namely DNA, Long Terminal Repeat (LTR), Rolling Circle (RC), Long Interspersed Nuclear Element
(LINE), Short Interspersed Nuclear Element (SINE), leaving the rest assigned to unclassified

(Unknown). To calculate the *k-mer* frequencies at repeats, we used the repeat annotation of each

genome assembly to recover their sequences using the *bedtools getfasta* tool [87]. As for the coding

and non-coding compartments, we counted *k-mer*s at repeat sequences using jellyfish 2.3.0 and

calculated for each *k-mer* its normalized frequency using the sum of all *k-mer* counts at repeats within

each genome. For each *k-mer*, the repeat enrichment is represented by the frequency at repetitive

sequences over its frequency at non-coding sequences, *i.e* the *k-mer* is enriched in repetitive

sequences for values > 1.

Repeat sequence identity

To estimate to which extent each genome contains long stretches of high sequence identity, we

applied a blast-based strategy. Using the repeat annotation gff files of each genome, we first merged

overlapping repeats with the *bedtools merge* function and extracted the resulting repeated sequences

using *bedtools getfasta* [87]. The set of repeats of each genome was next blasted against itself using

blastn from the BLAST 2.10.0+ suite using a word size of 11 and reporting only the top 10 sequences

with hits of ≥80% sequence identity (-max_target_seqs 10 -perc_identity 80 -word_size 11). The

resulting blastn results were parsed by removing self-hits (query hit against itself) and keeping only

the first hit if for a same target sequence multiple overlapping hits with identical alignment score were

identified (*i.e* same pident, length, mismatch, gapopen and qstart). The remaining hits were then

assigned to the following four categories based on their alignment length and sequence identity score:

alignment length > 100 bp, >1,000 bp, >5,000 bp or >10,000 bp, with sequence identities between 80-

95% or >95%.

Genome compartmentalization

To assess the concordance of genome segmentation metric with the "two-speed" genome concept, we

estimated genome compartmentalization using gene density. For this, we calculated for each gene in

each genome the distance to the nearest gene in the 5' and 3' context. We defined regions where

coding sequences are flanked on both sides by intergenic sequences >5 kb as gene sparse. We

considered genomes with >1% of their gene pool with flanking intergenic regions >5 kb as

compartmentalized.

Phylogenetic signal and independent contrasts

We estimated phylogenetic signals for the 12 genome assembly metrics using the *phylosignal* package

in R (version 1.3, [94]). We first concatenated phylogenetic relationship and trait variables into a

phylo4d object using the *phylo4d* function implemented in the phylobase package and assessed the

signal for each trait using the five implemented methods by the *phyloSignal* function, namely,

Blomberg's K and K*, Abouheif's Cmean, Moran's I, and Pagel's Lambda

(https://github.com/fmichonneau/phylobase). In addition, we computed the correlation between the

same 12 genome assembly metrics using the phylogenetically independent contrasts (method by

Felsenstein 1985 [95]) and implemented in the *ape* package in R version 5.7-1 [86]. We computed the

correlation between each pair of metrics by fitting a linear model between the two independent

contrasts (fit = lm(iy~ix-1)) in R. For each linear model, we used the coefficient of determination as a

measure of correlation between the two variables (adjusted R-squared), and the *p*-value estimates

were adjusted using the false discovery rate method in R.

Identification of putative shifts in genome architecture

To identify the edges in the phylogeny with putative major shifts in genome architecture, we fitted a

modified random walk process of trait evolution modeled by an Ornstein-Uhlenbeck (OU) process

and implemented in the *phylolm* package in R version 2.6.2 [96]. For each variable, we fitted the

model with $\log_{10}$ transformed values except for null values (kept as 0), as well as the PC1 and PC2

values of the CATAStrophy output. For each model, we allowed a maximum of 100 shifts across the

phylogeny (nmax = 100 option). Best models were selected based on log-likelihoods using the

modified Bayes information criterion (mBIC, [97]). The *OUshifts* function outputs the edge on the

tree where the shift occurred and the estimated shifts in the expected value of the trait (pshift and shift

values in the output). For each edge with a putative trait shift, we recovered the underlying tips in the

tree using the corresponding node with the *offspring* function from the *tidytree* package v0.4.5 [98].

Gene orthology analysis

We inferred gene orthology across the 1,239 species based on protein sequence identity. For that we

used *orthofinder* version 2.4.1, which implements *diamond* for homology search (version 0.9.24) [99].

From the entire set of 13,863,658 single proteins, *orthofinder* retrieved 1,008,244 orthogroups. We

assessed pangenome categories based on the frequency of each orthogroup in the dataset. We

considered orthogroups present in >90% of the 1,239 species as "core" orthogroups. Orthogroups

assigned to the "softcore" category were present in >80% but <90% of the species. The "shell"

orthogroups were present in >50% but <80% of the species. Orthogroups present in <50% of the

species were considered "variable" and those present in only one species were called "singletons". We

found major differences in the orthogroup composition across the different taxonomic groups in the

dataset. Saccharomycotes have fewer protein coding sequences than the rest of the dataset but ~54%

were assigned core proteins. Species in the Oomycota outgroup have a large fraction of their

proteome assigned as variable (average of 61% of the proteins). Most orthogroups are well defined in

the dataset, as ~84% are single-copy and likely representing true orthologs (790,441 / 933,315). Six

orthogroups were present in all 1,239 species of the dataset, and all six are multi-copy and associated

with functions related to repeat elements (Supplementary Table S7). The number of proteins assigned

to an orthogroup ranges from 1-43,748 (Supplementary Table S8). We identified 11,070 orthogroups

with ≥10 representatives in at least one genome, and 142,874 orthogroups with more than one

representative in at least one species ("multi-copy", Supplementary Figure S7, Supplementary Table
S9). We identified 15 species for which >40% of the proteome was constituted of multi-copy
orthogroups for genome sizes ranging from ~16Mb to >770Mb (*Martiniozyma abiesophila* and
Gigaspora margarita, respectively).

Annotation of functional domains across proteomes

To identify putative functional domains across the species proteomes, we downloaded the annotated
domains hidden Markov models from the Pfam release 31 [100]. We used the *hmmsearch* function
from the *Hmmer* package version 3.3.2 to scan all the species proteomes for functional domains (--
noali option to speed up the process) [78]. We then filtered the resulting outputs for a minimal
bitscore of 50 and a maximal e-value of $1e^{-17}$ using the HmmPy.py script
(https://github.com/EnzoAndree/HmmPy).

Identification of putative determinants of genome architecture

We applied a hypothesis-free approach to identify putative genetic determinants of trait variation in a
phylogenetic context. We used the *treeWAS* v1 package in R, a method that performs genome-wide
associations between a continuous or binary trait and biallelic genomic loci correcting for the
confounding effects of relatedness. We performed association tests for a total of 27 traits, including
genomic metric, lifestyle, and *k-mer* enrichment (Supplementary Table S15). For each variable, we
performed the association of log_{10} transformed values except for null values that were set to 0 and the
PC1 and PC2 values of the CATAStrophy output, which was kept untransformed. For the genotype
matrix, we encoded binary values representing presence/absence of the 10,474 orthogroups identified
by *orthofinder* with a minimum frequency of 5% in the dataset (see "gene orthology analysis").
Similarly, we performed association analyses using a presence / absence matrix of the 18,259 protein
families (Pfam) identified using *hmmsearch* with the HMM models of the Pfam release 31 (see
"annotation of functional domains across proteomes"). For each trait, we reported the associated
orthogroups and Pfams given the terminal, simultaneous and subsequent association tests performed
by *treeWAS* (Supplementary Table S13). For the associated orthogroups of interest, protein sequences
were aligned using Clustal Omega version 1.2.4 allowing for five iterations (--iterations 5) [101].
Gaps in the resulting alignment were trimmed using *trimAl* v1.4.rev15 and gaps in the resulting
alignment were further excluded using the -gappyout option [82]. The phylogenetic relationship
among proteins was inferred from the trimmed and filtered alignment using FastTree under the
Whelan-And-Goldman 2001 model after 1,000 bootstraps (-boot 1000 and -wag options) [102]. Based
on the annotated HMM domains for each protein assigned to the orthogroup, we also counted the
occurrence of Pfams for each protein sequence. For the most co-occurring Pfams, we illustrated the
protein topology using a random representative using the *drawProteins* package in R [103].

Protein family analysis

We focused on a set of protein families related to DNA biology (DNA methylation and DNA repair). For that, we screened the Hidden Markov Models domains identified across the entire set of proteins given a Pfam identifier (see "annotation of functional domains", Supplementary Table S10). All proteins with a hit for a given Pfam domain were filtered for non-canonical sequences (i.e., no methionine start, in-frame stop codons, https://github.com/milesroberts-123/extract-weird-proteins). The remaining hits were used to assess presence/absence of DNA-repair and DNA methylation related Pfams in each genome (Supplementary Table S10). To estimate presence-absence of the eight *N. crassa* RIP-related genes and their orthogroups, we used the *N. crassa* protein sequences to identify RIP-associated orthogroups. For the list of genes associated with meiotic recombination, heterochromatin, histone biology and RNA interference (Supplementary Table S10), we used *diamond* in sensitive mode to identify the best reciprocal blast hit among the full set of 13,863,658 proteins (version 0.9.24). For each of the query proteins, the best reciprocal hit was used to define presence/absence of the corresponding gene for each genome.

Protein identity analysis

For the 1,239 genomes, we performed *diamond blastp* searches using the set of annotated proteins within a genome as both query and subject (default parameters, version 0.9.24). The resulting alignments were filtered for redundant hits and classified according to their length (<=50, 50<100, 100<500 or >500 amino acids sequence alignment).

Orthogroup synteny analyses

We focused on the set of 4,666 orthogroups present in >5% of the dataset, single-copy in >80% of the genomes and represented in at least 10 different taxonomic classes (Supplementary Table S6). For each of the 4,666 orthogroups, we recovered the 10 neighboring genes in each direction, the orthogroup assignment and counted their occurrence near the focal orthogroup in the dataset. For each pair of focal orthogroups and nearby orthogroups, we estimated their synteny ratio by dividing the number of species with the orthogroup in range of the focal orthogroup by the total number of species carrying the focal orthogroup. The synteny of each of the 4,666 orthogroups was summarized by the mean synteny of all the orthogroups found in the 10+10 neighboring gene range.

Figures

587
Figure 1: Genome architecture across 1,239 fungi. A. Assembly metrics of the 1,239 genomes.
Assembly contiguity varies greatly in the dataset, but except for 13 assemblies from the Oomycota
phylum (100 genes in the *Strametopiles* database), all genomes contain more than 100 single-copy
BUSCO genes. **B.** Cumulative length in base-pairs of annotated repeats across the 1,239 genome
assemblies. **C.** Correlation between repetitive and non-repetitive genome size across the 1,239
assemblies. **D.** Heatmap showing the number of sequences sharing more than 95% sequence identity

over 100 bp, 1 kb, 5 kb and 10 kb across the 1,239 assemblies. Only a small fraction of genomes carry multiple >5 kb repeats with >95% sequence identity. Genomes with >1,000 hits of >95% sequence identity were also among the top 25% largest genomes (31 out of 37 genomes). **E.** Phylogenetic reconstruction of the relationship based on the concatenated alignment of 100 protein-coding genes. The resulting phylogeny recapitulates all major taxonomic classes represented in the dataset with the Saccharomycetes, Sordariomycetes, Eurotiomycetes, Dothideomycetes and Agaricomycetes forming the largest clades. **F.** Correlation of the phylogenetic independent contrasts of different assembly metrics across the 1,239 genome assemblies. We used the reconstructed phylogeny to calculate the phylogenetic independent contrast of each genome assembly metric.

Figure 2: Widespread shifts in genome architecture and lifestyle across fungi. A. Number of shifts identified across the phylogeny of 1239 species and for each genomic trait. **B.** The number of shifts identified for each genomic trait and their co-occurrence. **C.** Estimated intensity of the shifts identified for each genomic trait. **D.** Shifts co-occurring at the same edge in the phylogeny. **E.**

609 Phylogenetic positions of the shifts identified for genome size. Negative values denote an identified
610 decrease in the genomic trait value while positive values denote an increase in the genomic trait value.
611

612
613 **Figure 3: Major shift in genome size detected in Leotiomycetes. A.** Phylogenetic branch in which
614 the ~30Mb increase in genome size occurred in the Leotiomycetes. Presence / absence heatmap of
615 RIP-related genes and genes associated with genome size. **B.** Genomic synteny at the Rid1 locus
616 between *Blumeria graminis* and *Cadophora* sp. *DSE10*. Syntenic links are reported for encoded

amino-acid sequences. **C.** Synteny of 4,666 orthogroups across the 1,239 genomes. The dotted line
depicts the average synteny across all orthogroups.

Figure 4: Patchy distribution of RIP-related protein coding genes across fungi. A. Proportion of
the pangenome categories discovered among the 1,239 genomes. **B.** Presence / absence heatmap for
eight protein coding genes related to RIP in *Neurospora crassa*. **C.** Variation in six measures of
genome architecture in relation to the presence / absence of the eight RIP-related protein coding
genes. Presence of the gene is depicted by shaded boxes. **D.** Presence / absence heatmap for five
protein families (Pfams) related to DNA maintenance.

628
Figure 5: Identification of repeat-enriched mutation signatures in fungal genomes. A. Schematic
of the approach used to assess signatures of repeat-targeted mutations. **B.** Correlation between the
frequencies at repetitive sequences and non-coding sequences of *k-mer*s highly enriched in non-
coding sequences. Data is shown for four major taxonomical classes. **C.** Heatmap for the top 23 *k-mer*
enrichments in non-coding sequences compared to coding sequences. **D.** For each genome, bars show
the number of *k-mer*s that have both repeat and non-coding enrichment values > 2 (upper bars) or that
have only the non-coding enrichment value > 2 (lower panel). **E.** Proportion of *k-mers* enriched >2-
fold in both non-coding and repeat regions across all genomes. Species are grouped by taxonomic
class. Values larger than 0.5 is consistent with recent RIP activity. Dot size represents the number of
k-mers enriched >2-fold at repeats.
639

640
Figure 6: Identification of protein coding genes associated with repeat-enriched ***k-mer***
frequencies. A. Number of orthogroups associated with genome architecture. **B.** Word cloud of the
most represented protein family annotations of orthogroups associated with genome architecture. **C.**
Presence / absence heatmap of the orthogroups associated with genome architecture. **D.** Effect size of
the top associated orthogroups (presence/absence variation) with six genomic traits expressed by eta-
squared values. **E.** Phylogenetic relationship of proteins assigned to the orthogroup OG0000460
associated with *k-mer* enrichment at repeats.

References

1.   Thomas CA. The genetic organization of chromosomes. Annual review of genetics. Annu Rev Genet; 1971. pp. 237–256. doi:10.1146/annurev.ge.05.120171.001321

2.   Hatje K, Mühlhausen S, Simm D, Kollmar M. The Protein-Coding Human Genome: Annotating High-Hanging Fruits. BioEssays. 2019;41: 1900066. doi:10.1002/BIES.201900066

3.   Arkhipova IR. Neutral Theory, Transposable Elements, and Eukaryotic Genome Evolution. Mol Biol Evol. 2018;35: 1332–1337. doi:10.1093/MOLBEV/MSY083

4.   Burns KH, Boeke JD. Human Transposon Tectonics. Cell. 2012;149: 740–752. doi:10.1016/J.CELL.2012.04.019

5.   Payer LM, Burns KH. Transposable elements in human genetic disease. Nature Reviews Genetics 2019 20:12. 2019;20: 760–772. doi:10.1038/s41576-019-0165-8

6.   Hedges DJ, Deininger PL. Inviting instability: Transposable elements, double-strand breaks, and the maintenance of genome integrity. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis. 2007;616: 46–59. doi:10.1016/J.MRFMMM.2006.11.021

7.   Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of the genome. Nat Rev Genet. 2007;8: 272–285. doi:10.1038/nrg2072

8.   Gladyshev E. Repeat-Induced Point Mutation and Other Genome Defense Mechanisms in Fungi. Microbiol Spectr. 2017;5. doi:10.1128/microbiolspec.funk-0042-2017

9.   Buchon N, Vaury C. RNAi: a defensive RNA-silencing against viruses and transposable elements. Heredity 2006 96:2. 2005;96: 195–202. doi:10.1038/sj.hdy.6800789

10. Fedoroff N V. Transposable elements, epigenetics, and genome evolution. Science Nov 9, 2012 pp. 758–767.

11. Zhou W, Liang G, Molloy PL, Jones PA. DNA methylation enables transposable element-driven genome expansion. Proc Natl Acad Sci U S A. 2020;117: 19359–19366. doi:10.1073/PNAS.1921719117/SUPPL_FILE/PNAS.1921719117.SAPP.PDF

12. Tobias PA, Schwessinger B, Deng CH, Wu C, Dong C, Sperschneider J, et al. Austropuccinia psidii, causing myrtle rust, has a gigabase-sized genome shaped by transposable elements. G3 Genes|Genomes|Genetics. 2021;11. doi:10.1093/G3JOURNAL/JKAA015

13. Kelkar YD, Ochman H. Causes and Consequences of Genome Expansion in Fungi. Genome Biol Evol. 2012;4: 13–23. doi:10.1093/GBE/EVR124

14. Stajich JE, Lovett B, Lee E, Macias AM, Hajek AE, Bivort BL de, et al. Signatures of transposon-mediated genome inflation, host specialization, and photoentrainment in Entomophthora muscae and allied entomophthoralean fungi. Elife. 2023;12. doi:10.7554/ELIFE.92863.1

15. Nakamoto AA, Joubert PM, Krasileva K V. Intraspecific Variation of Transposable Elements Reveals Differences in the Evolutionary History of Fungal Phytopathogen Pathotypes. Genome Biol Evol. 2023;15. doi:10.1093/GBE/EVAD206

16. Lorrain C, Feurtey A, Ller MM, Haueisen J, Stukenbrock E. Dynamics of transposable elements in recently diverged fungal pathogens: Lineage-specific transposable element content and efficiency of genome defenses. G3: Genes, Genomes, Genetics. 2021;11. doi:10.1093/g3journal/jkab068

694   17.   Bewick AJ, Hofmeister BT, Powers RA, Mondo SJ, Grigoriev I V., James TY, et al.
695   Diversity of cytosine methylation across the fungal tree of life. Nature Ecology &
696   Evolution 2019 3:3. 2019;3: 479–490. doi:10.1038/s41559-019-0810-9
697   18.   Nakayashiki H, Kadotani N, Mayama S. Evolution and diversification of RNA silencing
698   proteins in fungi. J Mol Evol. 2006;63: 127–135. doi:10.1007/S00239-005-0257-
699   2/FIGURES/4
700   19.   Kim JM, Vanguri S, Boeke JD, Gabriel A, Voytas DF. Transposable Elements and
701   Genome Organization: A Comprehensive Survey of Retrotransposons Revealed by the
702   Complete Saccharomyces cerevisiae Genome Sequence. Genome Res. 1998;8: 464–
703   478. doi:10.1101/GR.8.5.464
704   20.   Stajich JE. Fungal Genomes and Insights into the Evolution of the Kingdom. Microbiol
705   Spectr. 2017;5. doi:10.1128/MICROBIOLSPEC.FUNK-0055-2016/ASSET/F179A154-
706   6318-4BDA-9DC0-7FC2C7D9CA5B/ASSETS/GRAPHIC/FUNK-0055-2016-FIG2.GIF
707   21.   Nagy LG, Ohm RA, Kovács GM, Floudas D, Riley R, Gácser A, et al. Latent homology
708   and convergent regulatory evolution underlies the repeated emergence of yeasts.
709   Nature Communications 2014 5:1. 2014;5: 1–8. doi:10.1038/ncomms5471
710   22.   Shen XX, Opulente DA, Kominek J, Zhou X, Steenwyk JL, Buh K V., et al. Tempo and
711   Mode of Genome Evolution in the Budding Yeast Subphylum. Cell. 2018;175: 1533-
712   1545.e20. doi:10.1016/J.CELL.2018.10.023
713   23.   Naranjo-Ortiz MA, Gabaldón T. Fungal evolution: cellular, genomic and metabolic
714   complexity. Biol Rev Camb Philos Soc. 2020;95: 1198. doi:10.1111/BRV.12605
715   24.   Nguyen TA, Cissé OH, Yun Wong J, Zheng P, Hewitt D, Nowrousian M, et al.
716   Innovation and constraint leading to complex multicellularity in the Ascomycota. Nat
717   Commun. 2017;8: 1–13. doi:10.1038/ncomms14444
718   25.   Keeling PJ, Slamovits CH. Simplicity and complexity of microsporidian genomes.
719   Eukaryot Cell. 2004;3: 1363–1369. doi:10.1128/EC.3.6.1363-1369.2004
720   26.   Katinka MD, Duprat S, Cornillott E, Méténler G, Thomarat F, Prensier G, et al. Genome
721   sequence and gene compaction of the eukaryote parasite Encephalitozoon cuniculi.
722   Nature. 2001;414: 450–453. doi:10.1038/35106579
723   27.   Rispe C, Moran NA. Accumulation of Deleterious Mutations in Endosymbionts:
724   Muller's Ratchet with Two Levels of Selection. Am Nat. 2000;156: 425–441.
725   doi:10.1086/303396
726   28.   Wernegreen JJ, Moran NA. Evidence for genetic drift in endosymbionts (Buchnera):
727   analyses of protein-coding genes. Mol Biol Evol. 1999;16: 83–97.
728   doi:10.1093/OXFORDJOURNALS.MOLBEV.A026040
729   29.   Woolfit M, Bromham L. Increased Rates of Sequence Evolution in Endosymbiotic
730   Bacteria and Fungi with Small Effective Population Sizes. Mol Biol Evol. 2003;20:
731   1545–1555. doi:10.1093/MOLBEV/MSG167
732   30.   Wernegreen JJ. Endosymbiont evolution: Predictions from theory and surprises from
733   genomes. Ann N Y Acad Sci. 2015;1360: 16. doi:10.1111/NYAS.12740
734   31.   McDonald BA, Suffert F, Bernasconi A, Mikaberidze A. How large and diverse are field
735   populations of fungal plant pathogens? The case of Zymoseptoria tritici. Evol Appl.
736   2022;15: 1360–1373. doi:10.1111/EVA.13434
737   32.   Badet T, Oggenfuss U, Abraham L, McDonald BA, Croll D. A 19-isolate reference-
738   quality global pangenome for the fungal wheat pathogen Zymoseptoria tritici. BMC
739   Biol. 2020;18: 1–18. doi:10.1186/s12915-020-0744-3

740   33.   Oggenfuss U, Badet T, Wicker T, Hartmann FE, Singh NK, Abraham L, et al. A
741   population-level invasion by transposable elements triggers genome expansion in a
742   fungal pathogen. Elife. 2021;10: 2020.02.11.944652. doi:10.7554/elife.69249

743   34.   Freitag M. Histone Methylation by SET Domain Proteins in Fungi. Annu Rev Microbiol.
744   2017;71: 413–439. doi:10.1146/ANNUREV-MICRO-102215-095757/1

745   35.   Selker EU. Premeiotic instability of repeated sequences in Neurospora crassa. Annu
746   Rev Genet. 1990;24. doi:10.1146/ANNUREV.GE.24.120190.003051

747   36.   Wang L, Sun Y, Sun X, Yu L, Xue L, He Z, et al. Repeat-induced point mutation in
748   Neurospora crassa causes the highest known mutation rate and mutational burden of
749   any cellular life. Genome Biol. 2020;21: 142. doi:10.1186/S13059-020-02060-
750   W/FIGURES/3

751   37.   Selker EU, Cambareri EB, Jensen BC, Haack KR. Rearrangement of duplicated DNA in
752   specialized cells of Neurospora. Cell. 1987;51: 741–752. doi:10.1016/0092-
753   8674(87)90097-3

754   38.   Rouxel T, Grandaubert J, Hane JK, Hoede C, van de Wouw AP, Couloux A, et al.
755   Effector diversification within compartments of the Leptosphaeria maculans genome
756   affected by Repeat-Induced Point mutations. Nat Commun. 2011;2: 202.
757   doi:10.1038/ncomms1189

758   39.   Frantzeskakis L, Kusch S, Panstruga R. The need for speed: compartmentalized
759   genome evolution in filamentous phytopathogens. Mol Plant Pathol. 2019;20: 3–7.
760   doi:10.1111/MPP.12738

761   40.   Galagan JE, Selker EU. RIP: The evolutionary cost of genome defense. Trends in
762   Genetics. Elsevier; 2004. pp. 417–423. doi:10.1016/j.tig.2004.07.007

763   41.   Selker EU, Garrett PW. DNA sequence duplications trigger gene inactivation in
764   Neurospora crassa. Proceedings of the National Academy of Sciences. 1988;85: 6870–
765   6874. doi:10.1073/PNAS.85.18.6870

766   42.   Selker EU, Cambareri EB, Jensen BC, Haack KR. Rearrangement of duplicated DNA in
767   specialized cells of Neurospora. Cell. 1987;51: 741–752. doi:10.1016/0092-
768   8674(87)90097-3

769   43.   Carlier F, Nguyen TS, Mazur AK, Gladyshev E. Modulation of C-to-T mutation by
770   recombination-independent pairing of closely positioned DNA repeats. Biophys J.
771   2021;120: 4325–4336. doi:10.1016/J.BPJ.2021.09.014

772   44.   Gladyshev E, Kleckner N. Recombination-Independent Recognition of DNA Homology
773   for Repeat-Induced Point Mutation (RIP) Is Modulated by the Underlying Nucleotide
774   Sequence. PLoS Genet. 2016;12: e1006015. doi:10.1371/JOURNAL.PGEN.1006015

775   45.   Gladyshev E, Kleckner N. Direct recognition of homology between double helices of
776   DNA in Neurospora crassa. Nat Commun. 2014;5: 1–11. doi:10.1038/ncomms4509

777   46.   Mazur AK, Gladyshev E. Partition of Repeat-Induced Point Mutations Reveals
778   Structural Aspects of Homologous DNA-DNA Pairing. Biophys J. 2018;115: 605–615.
779   doi:10.1016/j.bpj.2018.06.030

780   47.   Freitag M, Williams RL, Kothe GO, Selker EU. A cytosine methyltransferase
781   homologue is essential for repeat-induced point mutation in Neurospora crassa.
782   Proceedings of the National Academy of Sciences. 2002;99: 8802–8807.
783   doi:10.1073/pnas.132212899

784   48.   Kouzminova E, Selker EU. dim-2 encodes a DNA methyltransferase responsible for all
785   known cytosine methylation in Neurospora. EMBO J. 2001;20: 4309–4323.
786   doi:10.1093/EMBOJ/20.15.4309

49. Gladyshev E, Kleckner N. DNA sequence homology induces cytosine-to-thymine
mutation by a heterochromatin-related pathway in Neurospora. Nat Genet. 2017;49:
887–894. doi:10.1038/ng.3857

50. Gladyshev E. Repeat-Induced Point Mutation and Other Genome Defense
Mechanisms in Fungi. Microbiol Spectr. 2017;5. doi:10.1128/microbiolspec.funk-
0042-2017

51. Gladyshev E, Kleckner N. Recombination-Independent Recognition of DNA Homology
for Repeat-Induced Point Mutation (RIP) Is Modulated by the Underlying Nucleotide
Sequence. PLoS Genet. 2016;12: e1006015. doi:10.1371/JOURNAL.PGEN.1006015

52. Kidwell MG. Transposable elements and the evolution of genome size in eukaryotes.
Genetica. 2002;115: 49–63. doi:10.1023/A:1016072014259

53. Castanera R, López-Varas L, Borgognone A, LaButti K, Lapidus A, Schmutz J, et al.
Transposable Elements versus the Fungal Genome: Impact on Whole-Genome
Architecture and Transcriptional Profiles. Feschotte C, editor. PLoS Genet. 2016;12:
e1006108. doi:10.1371/journal.pgen.1006108

54. Feschotte C, Pritham EJ. DNA Transposons and the Evolution of Eukaryotic Genomes.
2007. doi:10.1146/annurev.genet.40.110405.090448

55. Perkins DD, Margolin BS, Selker EU, Haedo SD. Occurrence of repeat induced point
mutation in long segmental duplications of neurospora. Genetics. 1997;147: 125–136.
doi:10.1093/genetics/147.1.125

56. Oggenfuss U, Croll D. Recent transposable element bursts are associated with the
proximity to genes in a fungal plant pathogen. Thomma B, editor. PLoS Pathog.
2023;19: e1011130. doi:10.1371/JOURNAL.PPAT.1011130

57. Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D. Stress-driven
transposable element de-repression dynamics in a fungal pathogen. Mol Biol Evol.
2019; 633693. doi:10.1101/633693

58. Dubin MJ, Mittelsten Scheid O, Becker C. Transposons: a blessing curse. Current
Opinion in Plant Biology. Elsevier Ltd; 2018. pp. 23–29. doi:10.1016/j.pbi.2018.01.003

59. van Wyk S, Wingfield BD, De Vos L, van der Merwe NA, Steenkamp ET. Genome-Wide
Analyses of Repeat-Induced Point Mutations in the Ascomycota. Front Microbiol.
2021;11: 622368. doi:10.3389/FMICB.2020.622368/BIBTEX

60. Ikeda KI, Nakayashiki H, Kataoka T, Tamba H, Hashimoto Y, Tosa Y, et al. Repeat-
induced point mutation (RIP) in Magnaporthe grisea: implications for its sexual cycle
in the natural field context. Mol Microbiol. 2002;45: 1355–1364. doi:10.1046/J.1365-
2958.2002.03101.X

61. Zong X, Lou Y, Xia M, Zhao K, Chen J, Huang J, et al. Recombination and repeat-
induced point mutation landscapes reveal trade-offs between the sexual and asexual
cycles of Magnaporthe oryzae. Journal of Genetics and Genomics. 2024.
doi:10.1016/J.JGG.2024.03.003

62. Idnurm A, Howlett BJ. Analysis of loss of pathogenicity mutants reveals that repeat-
induced point mutations can occur in the Dothideomycete Leptosphaeria maculans.
Fungal Genetics and Biology. 2003;39: 31–37. doi:10.1016/S1087-1845(02)00588-1

63. Li WC, Huang CH, Chen CL, Chuang YC, Tung SY, Wang TF. Trichoderma reesei
complete genome sequence, repeat-induced point mutation, and partitioning of
CAZyme gene clusters. Biotechnol Biofuels. 2017;10: 1–20. doi:10.1186/S13068-017-
0825-X/FIGURES/7

64.   Cuomo CA, Güldener U, Xu JR, Trail F, Turgeon BG, Di Pietro A, et al. The Fusarium
834   graminearum genome reveals a link between localized polymorphism and pathogen
835   specialization. Science (1979). 2007;317: 1400–1402.
836   doi:10.1126/SCIENCE.1143708/SUPPL_FILE/CUOMO_SOM.PDF

837   65.   Coleman JJ, Rounsley SD, Rodriguez-Carres M, Kuo A, Wasmann CC, Grimwood J, et
838   al. The Genome of Nectria haematococca: Contribution of Supernumerary
839   Chromosomes to Gene Expansion. PLoS Genet. 2009;5: e1000618.
840   doi:10.1371/JOURNAL.PGEN.1000618

841   66.   Graïa F, Lespinet O, Rimbault B, Dequard-Chablat M, Coppin E, Picard M. Genome
842   quality control: RIP (repeat-induced point mutation) comes to Podospora. Mol
843   Microbiol. 2001;40: 586–595. doi:10.1046/J.1365-2958.2001.02367.X

844   67.   Bouhouche K, Zickler D, Debuchy R, Arnaise S. Altering a Gene Involved in Nuclear
845   Distribution Increases the Repeat-Induced Point Mutation Process in the Fungus
846   Podospora anserina. Genetics. 2004;167: 151–159. doi:10.1534/GENETICS.167.1.151

847   68.   Li WC, Chen CL, Wang TF. Repeat-induced point (RIP) mutation in the industrial
848   workhorse fungus Trichoderma reesei. Appl Microbiol Biotechnol. 2018;102: 1567–
849   1574. doi:10.1007/S00253-017-8731-5/TABLES/1

850   69.   Lee DW, Freitag M, Selker EU, Aramayo R. A Cytosine Methyltransferase Homologue
851   Is Essential for Sexual Development in Aspergillus nidulans. PLoS One. 2008;3: e2531.
852   doi:10.1371/JOURNAL.PONE.0002531

853   70.   Grognet P, Timpano H, Carlier F, Ait-Benkhali J, Berteaux-Lecellier V, Debuchy R, et al.
854   A RID-like putative cytosine methyltransferase homologue controls sexual
855   development in the fungus Podospora anserina. PLoS Genet. 2019;15: e1008086.
856   doi:10.1371/JOURNAL.PGEN.1008086

857   71.   Arnaise S, Zickler D, Bourdais A, Dequard-Chablat M, Debuchy R. Mutations in mating-
858   type genes greatly decrease repeat-induced point mutation process in the fungus
859   Podospora anserina. Fungal Genetics and Biology. 2008;45: 207–220.
860   doi:10.1016/J.FGB.2007.09.010

861   72.   Phillips MA, Kutch IC, Long AD, Burke MK. Increased time sampling in an evolve-and-
862   resequence experiment with outcrossing Saccharomyces cerevisiae reveals multiple
863   paths of adaptive change. Mol Ecol. 2020;29: 4898–4912. doi:10.1111/MEC.15687

864   73.   Long A, Liti G, Luptak A, Tenaillon O. Elucidating the molecular architecture of
865   adaptation via evolve and resequence experiments. Nature Reviews Genetics 2015
866   16:10. 2015;16: 567–582. doi:10.1038/nrg3937

867   74.   Du L, Liu Q, Fan Z, Tang J, Zhang X, Price M, et al. Pyfastx: a robust Python package for
868   fast random access to sequences from plain and gzipped FASTA/Q files. Brief
869   Bioinform. 2021;22: 1–8. doi:10.1093/BIB/BBAA368

870   75.   Gao F, Zhang CT. GC-Profile: a web-based tool for visualizing and analyzing the
871   variation of GC content in genomic sequences. Nucleic Acids Res. 2006;34: W686–
872   W691. doi:10.1093/NAR/GKL040

873   76.   Hane JK, Paxman J, Jones DAB, Oliver RP, de Wit P. "CATAStrophy," a Genome-
874   Informed Trophic Classification of Filamentous Plant Pathogens – How Many Different
875   Types of Filamentous Plant Pathogens Are There? Front Microbiol. 2020;10: 492799.
876   doi:10.3389/FMICB.2019.03088/BIBTEX

877   77.   Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. dbCAN: a web resource for automated
878   carbohydrate-active enzyme annotation. Nucleic Acids Res. 2012;40.
879   doi:10.1093/NAR/GKS479

78.   Finn RD, Clements J, Eddy SR. HMMER web server: Interactive sequence similarity
searching. Nucleic Acids Res. 2011;39: W29-37. doi:10.1093/nar/gkr367
79.   Li Y, Steenwyk JL, Chang Y, Wang Y, James TY, Stajich JE, et al. A genome-scale
phylogeny of the kingdom Fungi. Current Biology. 2021;31: 1653-1665.e5.
doi:10.1016/J.CUB.2021.01.074
80.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:
Assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31: 3210–3212. doi:10.1093/bioinformatics/btv351
81.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780.
doi:10.1093/molbev/mst010
82.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:
1972–1973. doi:10.1093/BIOINFORMATICS/BTP348
83.   Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. PartitionFinder 2: New
Methods for Selecting Partitioned Models of Evolution for Molecular and
Morphological Phylogenetic Analyses. Mol Biol Evol. 2017;34: 772–773.
doi:10.1093/MOLBEV/MSW260
84.   Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
thousands of taxa and mixed models. Bioinformatics. 2006;22: 2688–2690.
doi:10.1093/BIOINFORMATICS/BTL446
85.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, et
al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Mol Biol Evol. 2020;37: 1530–1534. doi:10.1093/MOLBEV/MSAA015
86.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics. 2019;35: 526–528.
doi:10.1093/BIOINFORMATICS/BTY633
87.   Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26: 841–842. doi:10.1093/bioinformatics/btq033
88.   Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. Bioinformatics. 2011;27: 764–770.
doi:10.1093/BIOINFORMATICS/BTR011
89.   Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. PLoS One. 2016;11: e0163962.
doi:10.1371/JOURNAL.PONE.0163962
90.   Xu Z, Wang H. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. Nucleic Acids Res. 2007;35: W265–W268. doi:10.1093/nar/gkm286
91.   Ellinghaus D, Kurtz S, Willhoeft U. LTRharvest, an efficient and flexible software for de
novo detection of LTR retrotransposons. BMC Bioinformatics. 2008;9: 1–14.
doi:10.1186/1471-2105-9-18/TABLES/5
92.   Ou S, Jiang N. LTR_retriever: A Highly Accurate and Sensitive Program for
Identification of Long Terminal Repeat Retrotransposons. Plant Physiol. 2018;176:
1410. doi:10.1104/PP.17.01310
93.   Smit, AFA, Hubley, R & Green P. RepeatMasker Open-4.0. 2015.
94.   Keck F, Rimet F, Bouchez A, Franc A. phylosignal: an R package to measure, test, and
explore the phylogenetic signal. Ecol Evol. 2016;6: 2774–2780.
doi:10.1002/ECE3.2051

95.   Felsenstein J. Phylogenies and the Comparative Method. Am Nat. 1985;125: 1–15.

96.   Tung Ho LS, Ané C. A Linear-Time Algorithm for Gaussian and Non-Gaussian Trait
929   Evolution Models. Syst Biol. 2014;63: 397–408. doi:10.1093/SYSBIO/SYU005

930   97.   Bogdan M, Frommlet F, Biecek P, Cheng R, Ghosh JK, Doerge RW. Extending the
931   modified bayesian information criterion (mBIC) to dense markers and multiple
932   interval mapping. Biometrics. 2008;64: 1162–1169. doi:10.1111/J.1541-
933   0420.2008.00989.X

934   98.   Curry E, Zhang H. Data Integration, Manipulation and Visualization of Phylogenetic
935   Trees. Data Integration, Manipulation and Visualization of Phylogenetic Trees. 2022.
936   doi:10.1201/9781003279242

937   99.   Emms DM, Kelly S. OrthoFinder: Phylogenetic orthology inference for comparative
938   genomics. Genome Biol. 2019;20: 466201. doi:10.1186/s13059-019-1832-y

939   100.   Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the
940   protein families database. Nucleic Acids Res. 2014;42: D222.
941   doi:10.1093/NAR/GKT1223

942   101.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation
943   of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst
944   Biol. 2011;7: 539. doi:10.1038/MSB.2011.75

945   102.   Price MN, Dehal PS, Arkin AP. Fasttree: Computing large minimum evolution trees
946   with profiles instead of a distance matrix. Mol Biol Evol. 2009;26: 1641–1650.
947   doi:10.1093/molbev/msp077

948   103.   Brennan P. drawProteins: a Bioconductor/R package for reproducible and
949   programmatic generation of protein schematics. F1000Res. 2018;7.
950   doi:10.12688/F1000RESEARCH.14541.1

951
952