ATTENTION MODULATES STIMULUS REPRESENTATIONS IN NEURAL FEATURE DIMENSION MAPS

Daniel D. Thayer (danielthayer@ucsb.edu) and **Thomas C. Sprague** (tsprague@ucsb.edu)

Department of Psychological and Brain Sciences, University of California, Santa Barbara

Running head: Attention sculpts feature dimension maps

Key words: feature-selective attention, visual system, fMRI, priority map

Word count: 8558

Acknowledgments: This work was supported by NIH R01-EY035300, an Alfred P Sloan Research Fellowship, an Nvidia Hardware Grant, and US Army Research Office Cooperative Agreement W911NF-19-2-0026 for the Institute for Collaborative Biotechnologies

ABSTRACT

Computational theories posit that attention is guided by a combination of spatial maps for individual features that can be dynamically weighted according to task goals. Consistent with this framework, when a stimulus contains several features, attending to one or another feature results in stronger fMRI responses in regions preferring the attended feature. We hypothesized that multivariate activation patterns across feature-responsive cortical regions form spatial 'feature dimension maps', which combine to guide attentional priority. We tested this prediction by reconstructing spatial maps from fMRI activation patterns across retinotopic regions of visual cortex while participants performed a feature-selective attention task. Participants viewed a peripheral visual stimulus at a random location which always contained moving colored dots. On each trial, participants were precued to report the predominant direction of motion or color of the stimulus, or to attend fixation. Stimulus representations in reconstructed maps based on a spatial inverted encoding model were selectively enhanced in color-selective regions when color was attended, and in motion-selective regions when motion was attended. While enhancement was localized to the stimulus position in color-selective regions, modulations in motion-selective regions was consistent with a more global enhancement when motion was task relevant. These results suggest feature-selective cortical regions support 'neural feature dimension maps': spatial maps of different visual features that are dynamically reweighted based on task demands to guide visual behavior to the most relevant locations based on important features

Introduction

The environment typically contains many objects competing for our attention. While gardening, your goal may be to find green weeds in a strawberry bush, so attention will be directed to green objects. However, task-irrelevant objects that are salient may capture your attention, such as a hummingbird darting in the sky or a red strawberry. These two factors contributing to how attention is allocated in the visual field—task-relevant goals and task-irrelevant salience—are core components of priority map theory (Awh et al., 2012; Itti & Koch, 2001; Luck et al., 2020; Serences & Yantis, 2006; Treisman & Gelade, 1980; Wolfe, 1994). Per this theory, these priority signals are integrated to index the most important locations in a scene, and locations with the highest level of priority are then selected for further processing (Carrasco, 2011; Eckstein, 2011).

In this framework, overall priority is computed based on the combination of priority indexed independently for individual feature dimensions, like color, motion, orientation, and shape. Priority within these 'feature dimension maps' is a combination of salience (defined based on image-computable properties, like local feature contrast) and relevance (defined based on task goals, like which location(s) or feature(s) are important; Wolfe & Horowitz, 2004, 2017). For example, a color dimension map would index locations made salient based on their unique color relative to other stimuli, while a motion dimension map would do the same for stimuli moving in a unique direction. Additionally, it would be expected that making one feature task-relevant would sculpt stimulus representations in the corresponding map, leading to a greater impact on downstream priority maps (Bahle et al., 2019; Corbetta & Shulman, 2002; Folk et al., 1992; Leber & Egeth, 2006; Moore & Armstrong, 2003; Poltoratski et al., 2017; Witkowski & Geng, 2023).

Recently, we demonstrated that color- (hV4/VO1/VO2; Bartels & Zeki, 2000; Brewer et al., 2005) and motion- (TO1/TO2; Amano et al., 2009; Huk et al., 2002) selective visual cortical regions index salient locations on the screen based on their preferred feature values, even when those locations are task-irrelevant (Thayer & Sprague, 2023). Importantly, studies investigating the impact of task relevance on feature-selective regions cannot establish whether feature-selective regions satisfy predictions of neural feature dimension maps because they all either measured from a single feature-selective visual region (Bichot et al., 2005; Martínez-Trujillo & Treue, 2002; Mazer & Gallant, 2003; Ogawa & Komatsu, 2004, 2006; Reynolds & Desimone, 2003), include only a single

goal-relevant feature dimension (Beck & Kastner, 2005; Cook & Maunsell, 2002; Moran &

Desimone, 1985), or investigate goal-driven prioritization of a stimulus location without selective

attention to specific features (Bisley & Goldberg, 2003, 2006; Ikkai & Curtis, 2008; Sprague,

Itthipuripat, et al., 2018; Sprague & Serences, 2013). If the same regions selectively encode

representations of salient locations based on their preferred feature value and reweight their

representations based on an observer's task demands, this would provide further support for a

role of compartmentalized 'neural feature dimension maps' in computing attentional priority.

Task demands can sculpt neural responses in visual cortex through spatial attention (Desimone &

Duncan, 1995; Kastner et al., 1999; Sprague, Itthipuripat, et al., 2018) and feature-selective

attention (Chawla et al., 1999; Saenz et al., 2002; Serences & Boynton, 2007; Treue & Trujillo,

1999). Spatial attention improves neural representation of an attended location relative to other

irrelevant locations in the visual field as shown by changes in fMRI BOLD responses (Ikkai & Curtis,

2008; Kastner et al., 1999, 2001; Sprague, Itthipuripat, et al., 2018; Sprague & Serences, 2013; Vo

et al., 2017). Attention to specific feature values has instead been shown to have a spatially global

effect, where all instances of a goal-relevant feature value, such as the color green, are prioritized

irrespective of stimulus location or relevance. Previous work has demonstrated that feature-

selective attention modulates fMRI BOLD activity in color- and motion-selective regions

(Beauchamp et al., 1997; Chawla et al., 1999; McMains et al., 2007; O'Craven et al., 1999; Runeson

et al., 2013; Saenz et al., 2002), by enhancing responses corresponding to stimuli matching the

attended feature value across the entire screen, even at locations containing no stimulus

(Serences & Boynton, 2007). Since both relevance signals can together sculpt neural response

patterns within feature-selective regions of cortex, it is important to characterize their joint

influence on activation profiles supporting neural feature dimension maps.

While both relevance of a stimulus location (spatial attention) and relevance of a specific feature

dimension (feature-selective attention) have each been shown to impact activation levels at

stimulus locations in feature-selective regions, it is unknown whether feature-selective

modulations occur equivalently across the entire visual field, reflecting a *global enhancement* of

processing of the attended feature dimension across the entire map, or instead, are primarily

restricted to the location of the presented stimuli containing the attended feature dimension,

reflecting a *local enhancement* of relevant locations within the attended map (Fig. 1; McMains et
al., 2007).

Here, we used a multivariate inverted encoding model (IEM; Thayer & Sprague, 2023) approach
to determine how attention to a cued feature dimension modulated activity across neural feature
dimension maps encoding an identical visual stimulus. We found that map locations corresponding
to the stimulus location in motion- and color-selective regions had the greatest activation, and
that these stimulus representations were strongest when attending to each region's preferred
feature. Interestingly, feature-specific modulations in color regions showed evidence for local
enhancement while modulations in motion regions more closely resembled global enhancement,
suggesting that each feature dimension map was uniquely modulated by feature-selective
attention. These findings were only observable after implementing IEM and leveraging region-
level spatial activation patterns, demonstrating the necessity to consider the information jointly
encoded across entire neural populations when characterizing the computational properties of
neural feature dimension maps.

Figure 1: How does attention to a specific feature dimension sculpt neural feature dimension maps? **A.** Priority map theory posits that various "feature dimension maps" are used to index the most important locations in the visual field based on computations within their preferred feature dimension, and that activation in these maps should scale with an observer's goals. If detecting or discriminating motion is needed for an ongoing task (e.g., identifying the motion direction of a darting hummingbird), then activation within the corresponding "motion map" would increase the importance associated with the hummingbird's location. There are two ways in which the motion map could prioritize information beyond the local effects expected of spatial attention (e.g., Sprague et al, 2018). Local enhancement could occur, such that only the location of a stimulus with the attended feature is prioritized. Alternatively, global enhancement could occur, such that activation across the entire map is additively scaled, leading to increased sensitivity to the attended feature dimension at any location. This type of modulation would still drive a stronger target representation but would additionally lead to stronger responses in locations that contain no stimulus when motion is the goal-relevant feature dimension. A motion dimension map is depicted here, but modulations would similarly apply for other feature dimensions, such as color. **B.** Evaluating the activation

at the stimulus location and the opposite location within the feature (motion) map can discriminate between the local and global enhancement accounts. Both models predict that the stimulus will have the greatest activation at the stimulus location when the preferred feature dimension is relevant (e.g., motion; left). If enhancement is localized, then activation at the opposite location should not change across conditions (center). However, if there is global enhancement, then the opposite location should have increased activation when motion is task relevant. By computing the difference in activation between the stimulus and opposite locations, the spatial specificity of feature-selective modulations can be assessed (right). If this activation difference (stimulus-opposite) is greater for the attend motion condition in the motion map, enhancement is localized. However, if the activation difference is similar between the attend color and motion conditions, then enhancement is global across the feature dimension map.

Materials & Methods

Participants

We recruited 10 subjects from the University of California, Santa Barbara (UCSB) community to participate in this study (9 female, 18-29 years old). We identified an appropriate sample size for our main effect of interest by conducting a power analysis using pilot data (n = 3). In the pilot study, we had a strong effect (d_z = 1.24) which required a sample of 6 subjects to obtain 80% power with an alpha criterion of α = .05 as indicated by a power analysis conducted in G*Power (Faul et al., 2007). We collected a large number of measurements from each subject to minimize within-subject variance, which can benefit statistical power more than increased sample sizes (Baker et al., 2021). Subjects reported normal or corrected-to-normal vision and did not report neurological conditions. All procedures were approved by the UCSB Institutional Review Board (IRB# 5-24-0030), and the study was registered on ClinicalTrials.gov (NCT06281457). Subjects gave written informed consent before participating and were compensated for their time ($20/h for scanning sessions, $10-20/h for behavioral familiarization/training).

Stimuli and Procedure

Participants performed a 30-minute training session before scanning so that they were familiarized with the instructions. We used this session to establish the initial behavioral performance thresholds used in the first run of the scanning session. In the main task session, participants were scanned for a single two-hour session consisting of at least 4 runs of a spatial mapping task, which were used to independently estimate encoding models for each voxel, and 8 runs of the experimental feature-selective attention task. All participants also underwent

additional anatomical and retinotopic mapping scanning sessions (1-2x 1.5-2 hr sessions) to

identify regions of interest (ROI; see Region of interest definition).

Stimuli were presented using the Psychophysics toolbox (Brainard, 1997; Pelli, 1997) for MATLAB

(The MathWorks, Natick, MA). Visual stimuli were rear-projected onto a screen placed ~110 cm

from the participant's eyes at the head of the scanner bore using a contrast-linearized LCD

projector (1,920×1,080, 60 Hz) during the scanning session. For two of the participants, a CRS

BOLDScreen LCD monitor was used to present stimuli (1,920x1,080, 60 Hz) that was placed ~149

cm from the participant's eyes. Participants who performed the task on the BOLDScreen had

slightly different stimulus dimensions (scaled by a factor of .9 compared to the projector), but all

results are qualitatively identical for the subjects who used either the BOLDScreen monitor or the

projector, and all findings hold considering only the n = 8 participants tested with the projector.

Throughout, we describe stimulus dimensions/units based on the projector stimulus size. In the

behavioral familiarization session, we presented stimuli on a contrast-linearized LCD monitor

(2,560×1,440, 60 Hz) 62 cm from participants, who were seated in a dimmed room and positioned

using a chin rest. For all sessions and tasks (main tasks, localizers, and mapping task), we presented

stimuli on a neutral gray circular aperture (9.15° radius), surrounded by black (only aperture shown

in Fig. 2).

Feature-selective attention task

For each trial of the main task (Fig. 2), participants viewed a colorful moving dot stimulus in the

periphery (similar to that used in Mante et al., 2013) along with a flickering central fixation cross

(0.2° size). The colorful moving dot stimulus was a 1.5°-radius circular disc comprised of several

individual dots centered 5° from fixation at a random location along an invisible ring. Individual

dots were physically isoluminant, colored either cyan (RGB: [0; 200; 200]) or orange (RGB: [255;

165; 0]), and moved clockwise or counterclockwise around the stimulus center, or in a random

direction. Individual dots occupied 0.04° of visual angle, with a dot density was 40 dots/deg^2, and

moved at a speed of 4.5°/s. Dots were randomly replotted every 50 ms or when they exceeded

the stimulus bounds. Both cyan and orange dots were always present. Dots were always moving,

but only some dots on each trial would move clockwise or counterclockwise, while the rest of the

dots moved in a random direction (uniform distribution of planar motion directions).

At the start of each trial, participants were shown a letter cue at fixation (0.4° height) that would

indicate what aspect of the display to attend. The letter cue was presented for 1000 ms in white

(RGB: [180; 180; 180]) Arial font. The cue could be an 'F' (attend fixation), 'C' (attend color), or 'M'

(attend motion). After a 500 ms blank period, with only the aperture and fixation dot present, the

colorful moving dot stimulus and fixation cross was shown for 3000 ms. When cued to attend the

color of the stimulus ('C'), participants reported which individual feature value (cyan or orange)

was most prevalent in the stimulus. When cued to attend the motion of the stimulus ('M'), they

reported the coherent motion direction of the stimulus (clockwise or counterclockwise). Since

motion and color was present on each trial, visual input was constant, and we were able to isolate

changes in the neural stimulus representations across brain regions due to instructed task goals.

We adjusted the difficulty of the attend color/motion tasks between runs by changing the

coherence of the dots to try and equate task performance between conditions (e.g., 80% dot

coherence indicates that 80% of the dots were presented in cyan or were moving clockwise). The

first run of the task in the scanner used coherence values acquired from the behavioral training

session.

As a neutral baseline condition, participants attended the central fixation cross on some trials. The

vertical and horizontal lines of the fixation cross were 0.2° of visual angle long (0.1° from center in

either direction) and flickered at 3 Hz (10 frames on, 10 frames off at 60 Hz). On each trial, one

line of the fixation cross would slightly change in size for a single flicker cycle. When cued to attend

fixation ('F'), participants reported which line increased in length (horizontal or vertical) with a

button press. Since the dot stimulus was ignored on these trials due to fixation task difficulty

(Thayer & Sprague, 2023), the neural response associated with the stimulus on these trials was

purely driven by visual input. We adjusted the difficulty of the fixation task between runs by

altering the degree of size change for vertical/horizontal lines based on behavioral accuracy

(range: 0.21° to 0.32°). The fixation target appeared at a random time during the stimulus

presentation period other than the first or last 500 ms.

Participants completed 8 runs of the task, where each run contained 24 trials, with 8 trials of each

condition (attend fixation, attend stimulus color, attend stimulus motion). All trials were separated

by a randomly selected ITI ranging from 6-9 s with an average ITI of 7.5 s. Trial order was shuffled

within run. Each run started with a 3 s blank period and ended with a 10.5 s blank period, for a

total run duration of 301.5 s. Eye position (right eye) was monitored throughout the experiment

using an Eyelink 1000 eyetracker (SR Research) at 500 Hz. Participants performed a 5-point

calibration at the start of the experiment.

Figure 2: Feature-selective attention task. A. On each trial, participants viewed a dot array comprised of colorful moving dots. All dots were randomly either cyan or orange, and all dots were always moving, but only some would move clockwise or counterclockwise, while the rest moved in random direction. A flickering fixation cross (3 Hz) was also always present, and the vertical or horizontal arm of the cross would increase in size once during the stimulus presentation period. A letter at fixation cued participants to perform one of three attention tasks related to the upcoming stimulus display (F: attend fixation, C/M attend peripheral stimulus color/motion, respectively). On attend fixation trials, participants reported whether the horizontal or vertical arm of the fixation cross lengthened; on attend color/motion trials, they reported the prominent feature value of the cued feature dimension (more cyan/orange dots, or more dots moving clockwise/counterclockwise). B. Accuracy for each attention condition. Performance was slightly worse for the attend color than the attend motion or fixation conditions. Error bars indicate within-subject SEM (Franz & Loftus, 2012). Faint dots indicate individual subject means. * Significant difference based on permuted paired-samples t test ($p < 0.05$).

Spatial mapping task

We also acquired several runs of a spatial mapping task used to independently estimate a spatial

encoding model for each voxel, following previous studies (Sprague et al., 2016; Sprague,

Itthipuripat, et al., 2018; Thayer & Sprague, 2023). On each trial of the mapping task, we presented

a flickering checkerboard at different positions selected from a hexagonal grid spanning the

screen. Participants viewed these stimuli and responded whenever a rare contrast change

occurred (6 out of 43 trials, 13.9%), evenly split between contrast increments and decrements.

The checkerboard stimulus was the same size as the stimulus in the feature-selective attention

task (1.5° radius) and was presented at 70% contrast and 6-Hz full-field flicker. All stimuli appeared

within a gray circular aperture with a 9.15° radius, as in the feature-selective attention task. For

each trial, the location of the stimulus was selected from a hexagonal grid of 37 possible locations

with an added random uniform circular jitter (0.5° radius). The base position of the grid was

rotated by 30° on every other scanner run to increase spatial sampling density. As a result, every

mapping trial was unique, which enabled robust spatial encoding model estimation.

Each trial started with a 3000 ms stimulus presentation period. If a target was present, then the

stimulus would be dimmed/brightened for 500 ms with the stipulation that the contrast change

would not occur in either the first or last 500 ms of the stimulus presentation period. Finally, ITIs

were randomly selected to be either 6 or 8.25 s. All target-present trials were discarded when

estimating the spatial encoding model. Each run consisted of 43 trials (6 of which included targets).

We also included a 3 s blank period at the beginning of the run and a 10.5-s blank period at the

end of the run. Each run totaled 432 s.

Retinotopic mapping task

We used a previously reported task (Mackey et al., 2017; Thayer & Sprague, 2023) to identify

retinotopic regions of interest (ROIs) via the voxel receptive field (vRF) method (Dumoulin &

Wandell, 2008). Each run of the retinotopy task required participants attend several random dot

kinematograms (RDK) within bars that would sweep across the visual field in 2.25 s steps. Three

equally sized bars were presented on each step and the participants reported in which of the two

peripheral bars the motion in the central bar matched with a button press. Participants received

feedback via a red (incorrect) or green (correct) color change at fixation. We used a three-

down/one-up staircase to maintain ~80% accuracy throughout each run so that participants would

continue to attend the RDK bars. RDK bars swept 17.5° (15.8° on BOLDScreen) of the visual field.

Bar width and sweep direction was pseudo-randomly selected from several different widths

(ranging from 2.0° to 7.5°) and four directions (left-to-right, right-to-left, bottom-to-top, and top-

to-bottom).

fMRI acquisition and MRI preprocessing

All scanning was conducted at the UCSB Brain Imaging Center with a 3T Siemens Prisma scanner. fMRI data acquisition and preprocessing pipelines in the current study exactly follow our previous report (Thayer & Sprague, 2023). All fMRI scans for retinotopic mapping, model estimation, and main experimental task were acquired using the CMRR MultiBand accelerated EPI pulse sequences. We acquired all images with the Siemens 64 channel head/neck coil with all elements enabled. We collected 3 T1-weighted and 1 T2-weighted anatomical images using the Siemens product MPRAGE and Turbo Spin-Echo sequences (0.8 mm isotropic voxels, 256 X 240 mm slice FOV). The TE/TR of each anatomical image were: 2.24/2400 ms (T1w) and 564/3200 ms (T2w). These scans were used to coregister all functional images to each participant's native anatomic space. We acquired functional images using a Multiband (MB) 2D GE-EPI scanning sequence with MB factor of 4, acquiring 44 2.5 mm interleaved slides with no gap, with isotropic voxel size being 2.5 mm. The TE/TR of these scans were: 30/750 ms (4x simultaneous-multi-slice acceleration). We used freesurfer's recon-all script (version 6.0) to process all anatomical scans and AFNI's afni_proc.py command to preprocess all functional scans. Preprocessing included unwarping, alignment, rigid-body motion correction (6-parameter affine transform), and smoothing along the cortical surface normal. Retinotopic mapping data was spatially smoothed along the cortical surface (5 mm FWHM) for identifying ROIs, but feature-selective attention and spatial mapping task data was not smoothed beyond projecting to surface space and returning to volume space.

Region of interest definition

We identified 15 ROIs using independent retinotopic mapping data. We fit a vRF model for each voxel in the cortical surface (in volume space) using averaged and spatially smoothed (on the cortical surface; 5 mm FWHM) time series data across all retinotopy runs (8-12 per participant). We used a compressive spatial summation isotropic Gaussian model (Kay et al., 2013; Mackey et al., 2017) as implemented in a customized, GPU-optimized version of mrVista (see Mackey et al., 2017) for detailed description of the model). High-resolution stimulus masks were created (270 x 270 pixels) to ensure similar predicted responses within each bar size across all visual field positions. Model fitting began with an initial high-density grid search, followed by subsequent nonlinear optimization. Retinotopic ROIs (V1, V2, V3, V3AB, hV4, LO1, LO2, VO1, VO2, TO1, TO2,

IPS0-3) were then delineated by projecting vRF best-fit polar angle and eccentricity parameters

with variance explained ≥5% onto each participant's inflated cortical surfaces via AFNI and SUMA.

We drew ROIs on each hemisphere's cortical surface based on previously-established polar angle

reversal and foveal representation criteria (Amano et al., 2009; Mackey et al., 2017; Swisher et al.,

2007; Wandell et al., 2007; Winawer & Witthoft, 2015). Finally, ROIs were projected back into

volume space to select voxels for analysis. For primary analyses (Figs. 4, 6), we focused on

previously identified 'feature dimension maps' (Thayer & Sprague, 2023) which were aggregate

ROIs made up of a combination of all voxels across retinotopic maps with similar feature selectivity

(color: hV4, VO1, and VO2; motion: TO1 and TO2). For completeness, we additionally report

results from each individual ROI (Figs. 5, 7).

Inverted encoding model.

We used a spatial inverted encoding model (IEM) to reconstruct images of stimulus-related

activation patterns measured across entire ROIs (Sprague & Serences, 2013; Fig. 3). To do this, we

first estimated an encoding model, which describes the sensitivity profile over the relevant feature

dimension for each voxel in a region. This requires using data set aside for this purpose, referred

to as the "training set". Here, we used data from the spatial mapping task as the independent

training set. The encoding model across all voxels within a given region is then inverted to estimate

a mapping used to transform novel activation patterns from a "test set" (runs from the feature-

selective attention task) and reconstruct the spatial representation of the stimulus at each

timepoint. For this analysis, we used voxels for which the best-fit vRF model explained ≥5% of the

variance during the retinotopic mapping task.

We built an encoding model for spatial position based on a linear combination of 37 spatial filters

(Sprague, Itthipuripat, et al., 2018; Thayer & Sprague, 2023). Each voxel's response was modeled

as a weighted sum of each identically shaped spatial filter arrayed in a triangular grid (Fig. 3A). The

centers of each filter were spaced by 2.83° and were Cosine functions raised to the 7th power:

$$f(r) = \left(0.5 + 0.5\cos\frac{\pi r}{s}\right)^7 \tag{1}$$

for $r < s$; 0 otherwise

where r is the distance from the filter center and s is a size constant. The size constant reflects the

distance from the center of each spatial filter at which the filter returns to 0. This triangular grid

of filters forms the basis set, or information channels for our analysis. For each stimulus used in

our mapping task, we converted from a contrast mask to a set of filter activation levels by taking

the dot product of the vectorized stimulus mask (n trials × p pixels) and the sensitivity profile of

each filter (p pixels × k channels). We then normalized the estimated filter activation levels such

that the maximum activation was 1 and used this output as C_1 in the following equation, which

acts as the forward model of our measured fMRI signals:

$$B_1 = C_1 W \tag{2}$$

296   B_1 (n trials × m voxels) in this equation is the measured fMRI activity of each voxel during the

297   visuospatial mapping task and W is a weight matrix (k channels × m voxels) which quantifies the

298   contribution of each information channel to each voxel. $\widehat{W}$ can be estimated using ordinary least-

299   squares linear regression to find the weights that minimize the differences between predicted

300   values of B and the observed B_1:

$$\widehat{W} = (C_1^T C_1)^{-1} C_1^T B_1 \tag{3}$$

301   This is computed for each voxel within a region independently, making this step univariate. The

302   resulting $\widehat{W}$ represents all estimated voxel sensitivity profiles. We then used $\widehat{W}$ and the measured

303   fMRI activity of each voxel (i.e., BOLD response) during each trial (using each TR from each trial, in

304   turn) of the feature-selective attention task using the following equation:

$$\hat{C}_2 = B_2 \widehat{W}^T (\widehat{W}\widehat{W}^T)^{-1} \tag{4}$$

Here, $\hat{C}_2$ represents the estimated activation of each information channel (n trials × k channels)

which gave rise to that observed activation pattern across all voxels within a given ROI (B_2; n trials

× m voxels). To aid with visualization, quantification, and coregistration of trials across stimulus

positions, we computed spatial reconstructions using the output of Equation 4. To do this, we

weighted each filter's spatial profile by the corresponding channel's reconstructed activation level

and then sum all weighted filters together (Fig. 3B).

Since stimuli in the feature-selective attention task were randomly positioned on every trial, we

rotated the center position of spatial filters such that the resulting 2D reconstructions of the

stimuli were aligned across trials and participants (Fig. 3C). We then sorted trials based on

condition (attend fixation, attend color, attend motion). Finally, we averaged the 2D

reconstructions across trials within the same condition for individual participants, then across all

participants for our grand-average spatial reconstructions (Fig. 5; Appendix Fig. A2). Individual

values within the 2D reconstructed spatial maps correspond to map activation at the

corresponding relative visual field coordinates based on activation profiles across entire ROIs.

Critically, because we reconstructed all trials from all conditions of the feature-selective attention

task using an identical spatial encoding model estimated with an independent spatial mapping

task, we can compare reconstructions across conditions on the same footing (Sprague, Adam, et

al., 2018; Sprague et al., 2019; Sprague, Itthipuripat, et al., 2018).

Figure 3: Inverted encoding model used to reconstruct maps of the visual field. A. Data from an independent spatial mapping task was used to estimate a spatial IEM for each ROI (for details, see Materials and Methods). **B.** Using this approach, activation patterns across all voxels in a region for each trial can be mapped onto the activation of spatial channels that are then summed to produce reconstructed spatial maps of the visual field. Identical encoding models, estimated with the task in (A), are used to compute reconstructions for each condition, ensuring results can be compared on equal footing. We reconstructed maps for each trial using activation patterns averaged 4.5-7.5 s after cue onset and directly compared the map activation across conditions. **C.** Since maps are in visual field coordinates, we were able to rotate and align reconstructed spatial maps to the known position of the moving dot array, then average across trials within each condition. Computed channel responses are example data from a representative subject's (sub015) aggregate TO1/TO2 ROI.

Quantifying neural responses

First, to establish the impact of feature-selective attention to specific stimulus dimensions on overall activation in each ROI at the broadest spatial scale, we computed the average activation across all visually responsive voxels (defined as those with at least 5% variance explained based on vRF model fits) in each ROI (McMains et al., 2007; Runeson et al., 2013). Second, because we can measure activation across each entire region simultaneously (that is, voxels with the full spread of spatial encoding properties), we performed a multivariate spatial IEM analysis (described above) to transform activation patterns to reconstructed spatial maps for each trial. This analysis leverages all spatial information from all voxels simultaneously to recover the stimulus representation supported by the entire neural population distributed across each retinotopic region or set of regions (Fig. 3C). To quantify stimulus representations in reconstructed spatial maps, we computed the mean map activation of pixels located within a 1.5° radius disk centered at the known position of each stimulus (matching the stimulus radius of 1.5°; see Sprague, Itthipuripat, et al., 2018; Thayer & Sprague, 2023). This provides a single value corresponding to the activation of the stimulus location for a given condition, within each retinotopic ROI. To assess the spatial selectivity of reconstructed spatial maps, we compared the mean map activation at the location of stimulus to map activation at the location opposite fixation using a 1.5° radius disk (Fig. 6 and Appendix Fig. A2).

For univariate and multivariate analyses, we computed an attention modulation index (AMI) to directly compare the impact of attending to color vs motion on stimulus representations, while accounting for the overall impact of spatial attention, similar to previous reports (McMains et al., 2007). AMI was computed as the difference in activation between the attend motion and attend color conditions, where positive values indicate a stronger response to the stimulus when attending motion (we did not normalize by the sum because, in some regions/analyses, the color- or motion-related activation was negative, making a normalized difference challenging to interpret).

In addition to the primary focused analyses on *a priori* feature-selective retinotopic regions (Figs. 4, 5), we also present all results computed for each retinotopic ROI individually (Appendix Figs. A1,

352 A2). All figures contain data that were averaged across participants as well as individual subject
353 means.

Statistical analysis

Parametric statistical tests were conducted for all comparisons (repeated-measures ANOVAs and
t tests). All statistical tests were conducted across participants after averaging all trials of each
condition within individual participants. To account for possible non-normalities in our data, we
generated null distributions for each test using a permutation procedure (see below) to derive p-
values. Throughout the manuscript, we report statistics (F/T-scores, effect sizes) based on
parametric assumptions, while p-values are derived from the shuffling procedure described above.

First, we used a one-way repeated-measures ANOVA (factor: attention condition, with 3 levels:
attend fixation, attend color, attend motion) to determine whether behavioral accuracy on the
fixation task depended on what was attended on each trial (Fig. 2B). We then conducted follow-
up paired-samples *t* tests between all pairs of attention conditions to see which conditions
differed. A paired-samples *t* test was conducted to compare attend color and attend motion RTs.
The attend fixation condition was excluded for this analysis due to differences in stimulus onset
(fixation target appeared briefly at least 500 ms after the dot array stimulus onset, while coherent
color/motion was present throughout the dot stimulus presentation), making comparisons
difficult to interpret.

When analyzing fMRI data, we focused on retinotopic regions that have previously been shown to
be color- or motion-selective (Brewer et al., 2005; Huk et al., 2002; Thayer & Sprague, 2023). For
univariate analyses, we computed two-way ANOVAs with attention condition (3 levels: attend
fixation, color, and motion) and ROI (2 levels: color-selective hV4/VO1/VO2 and motion-selective
TO1/TO2) as factors. Follow-up paired-samples *t* tests were then conducted to determine if
responses differed within each ROI as a function of attention condition (Fig. 4, Appendix Fig. A1).

For multivariate neural responses, we first computed a three-way ANOVA with location (2 levels:
stimulus location and opposite location), attention condition (3 levels: attend fixation, color, and
motion), and ROI (2 levels: color-selective hV4/VO1/VO2 and motion-selective TO1/TO2) as
factors. We then computed separate two-way ANOVAs for each relevant location (stimulus and

opposite location), with attention condition (3 levels: attend fixation, color, motion) and ROI (2

levels: color-selective hV4/VO1/VO2 and motion-selective TO1/TO2) as factors. This analysis

determined how task demands and feature selectivity uniquely modulated responses between

regions at each location. Next, we computed the difference between stimulus and opposite

locations and conducted a two-way ANOVA to determine the *spatial selectivity* of neural

modulations with attention condition (3 levels: attend fixation, color, motion) and ROI (2 levels:

color-selective hV4/VO1/VO2 and motion-selective TO1/TO2) as factors. To directly test whether

feature-selective ROIs represent stimulus locations more strongly when participants direct

attention to their preferred feature dimension, we computed a paired-samples t test on the AMI

computed for color-selective and motion-selective ROIs (Fig. 5C, Appendix Fig. A2C). Lastly, one-

sample t tests for univariate and multivariate AMIs were conducted for each region to determine

if neural responses were feature selective (AMI different from zero). For completeness, we include

data from all individual retinotopic ROIs (Appendix Fig. A5, A7) as well as the corresponding

statistical analyses (Table. 1, 2). However, those results are not discussed further as there were no

a priori hypotheses for individual regions. We note that results from individual feature-selective

ROIs were weaker, but qualitatively similar to the aggregate ROI analyses.

For our shuffling procedure, we used a random number generator that was seeded with a single

value for all analyses. The seed number was randomly selected using an online random number

generator   (https://numbergenerator.org/random-8-digit-number-generator).   Within   each

participant, averaged data within each condition were shuffled across conditions for each

participant individually, and once shuffled, the statistical test of interest was recomputed over

1000 iterations. P values were derived by computing the percentage of shuffled test statistics that

were greater than or equal to the measured test statistic. We controlled for multiple comparisons

using the false discovery rate (Benjamini & Yekutieli, 2001) across all comparisons within an

analysis when necessary. Error bars are standard error, unless noted otherwise.

Data & Code Availability

All data supporting the conclusions of this report and all associated analysis scripts are available

on   Open   Science   Framework

(https://osf.io/3qrpc/?view_only=93818b413458400f9fac93dd878f09a2). To protect participant

privacy, and in accordance with IRB-approved procedures, freely available data is limited to

extracted timeseries for each voxel of each ROI for all scans of the study. Whole-brain 'raw' data

will be made available from the authors upon reasonable request.

Results

Behavior

Participants performed a feature-selective attention task while we used fMRI to measure BOLD

activation patterns across several independently defined retinotopic regions. We cued

participants to covertly attend to either the color or motion of a colored-moving dot stimulus on

every trial to determine which individual feature value was most prevalent. Accuracy differed

across attention conditions as indicated by a permuted one-way repeated measures ANOVA,

$F(2,18) = 9.53$, $p = 0.002$, $\eta_p^2 = 0.51$, where accuracy was worse on attend color than attend motion

trials (paired sample t-test, $t(9) = 4.99$, p < 0.001, $d_z = 1.58$). This indicates that, despite our

attempts to match difficulty, the color task was more difficult than the motion task. Notably, there

was no significant difference in response times (RT; not shown) between the attend color and

motion conditions (mean color and motion RT: 1.65, 1.57; SEM: 0.05, 0.06; $t(9) = 1.42$, $p = 0.270$,

$d_z = 0.42$), suggesting participants were engaged with the peripheral stimuli for a similar duration

across conditions.

Univariate neural responses

First, to determine whether the attended feature dimension impacts aggregate neural responses

across independently mapped retinotopic ROIs selective for color and motion, we compared

univariate neural responses averaged across all visually responsive voxels within each region,

without considering their spatial selectivity with respect to the stimulus location (for individual ROI

results, see Appendix Fig. A1). Hemodynamic response functions (HRFs) for each aggregate

feature-selective ROI show a stimulus-related visually evoked response which differs as a function

of task condition (Fig. 4A). To quantify these attentional modulations, we averaged responses 4.5-

7.5 seconds after the attention cue was presented (3-6 s after stimulus onset; Fig. 4 B-C). A two-

way permuted ANOVA with attention condition (attend fixation, color, and motion), and ROI

(hV4/VO1/VO2 and TO1/TO2) as factors revealed a main effect of ROI ($F(1,9) = 24.19$, $p < 0.001$,

$\eta_p^2 = 0.97$) as well as a two-way interaction (F(2,18) = 54.97, $p < 0.001$, $\eta_p^2 = 0.86$). To determine if

each region's overall activation was modulated by attention to the peripheral stimulus as

compared to attention to the fixation point, we performed follow-up permuted two-sample t tests

within each ROI between each attend-stimulus condition and the attend fixation condition. This

analysis provides insights into how directing attention to the location of the peripheral stimulus

may modulate overall responses across all voxels within each region, possibly by enhancing the

response to goal-relevant feature dimensions. In hV4/VO1/VO2, there was a stronger response

when attending to the stimulus than when attending to fixation (fixation vs color: t(9) = 7.12, $p <$

0.001, d_z = 2.25; fixation vs motion: t(9) = 4.52, p = 0.001, d_z = 1.43), indicating that attention

directed to the stimulus enhanced responses regardless of attended feature. However, TO1/TO2

showed no general modulation when attention was directed to the stimulus for either condition

when compared to attending fixation (fixation vs color: t(9) = 2.12, p = 0.067, d_z = 0.67; fixation vs

motion: t(9) = 0.51, p = 0.598, d_z = 0.16).

Figure 4: BOLD response across all voxels in aggregate feature-selective visual cortex ROIs show motion selectivity for TO1/TO2 and task demand modulation in hV4/VO1/VO2. **A.** HRFs from aggregate feature-selective retinotopic regions TO1/TO2 and hV4/VO1/VO2. Gray lines indicate cue (first) and stimulus (second) onsets. Shaded area indicates time points used for trial averaging across participants. Dashed lines indicate SEM within participants. Dot array stimulus was presented 1.5 to 4.5 seconds (second gray line to start of shaded area). **B.** Mean BOLD response across participants for each attention condition. Motion-selective regions showed no modulation of response when attending to either feature of the stimulus (color or motion) relative to when attending to fixation (color vs fixation: p = 0.07; motion vs fixation: p = 0.59). Color-selective regions were modulated by the task-relevance of the peripheral stimulus, with a greater region-level response when attending to the color ($p < 0.01$) or the motion ($p < 0.01$) relative to attending to fixation. **C.** Attention modulation index for both aggregate feature-selective regions averaged across participants. Only motion-selective regions had a response that differed from zero (p = 0.02). * indicate significant differences based on permuted paired or single sample t tests ($p < 0.05$). Error bars indicate within-subject SEM. Faint dots indicate individual subject means. Col: color, Mot: motion, Fix: fixation.

Next, to assess whether the specific attended feature modulates activation in each region, we

computed an attention modulation index (AMI) by subtracting the neural responses between the

attend motion and attend color conditions, with a positive value indicating a stronger response

when attending to motion (note that this is equivalent to a paired comparison between attend-

motion and attend-color conditions). Permuted one-sample t tests revealed that TO1/TO2 had a

significant positive value ($t(9) = 2.35$, $p = 0.024$, $d_z = 0.74$). Even though motion-selective regions

showed no general modulation when attention was directed to the stimulus as compared to

fixation (Fig. 4B), there was a difference in overall response as a function of attended feature

dimension. In contrast, hV4/VO1/VO2 AMI based on overall activation did not significantly differ

from zero ($t(9) = 1.79$, $p = 0.061$, $d_z = 0.56$). Thus, while color regions were more active when the

peripheral stimulus was relevant, their overall activation did not significantly vary based on what

feature dimension was attended. Finally, a direct comparison of AMI between color- and motion-

selective regions revealed a significant difference (Fig. 4C; $t(9) = 6.07$, $p < 0.001$, $d_z = 1.92$),

consistent with a region-selective modulation of overall activation levels based on attended

feature.

In sum, attending to specific feature dimensions rescales the overall activation levels of feature-

selective retinotopic ROIs, with regions showing greater univariate activation when their preferred

feature dimension is task-relevant.

Multivariate spatial representations

While informative, the univariate results alone cannot speak to the mechanism of attentional

modulation of feature dimension maps, because univariate analyses collapsing across entire

retinotopic maps are unable to discern which parts of the region—corresponding to different

locations in the visual field—are modulated by task demands. Is the re-scaling observed in

univariate analyses (Fig. 4) driven primarily by neural populations encoding the attended location,

or instead, does this reflect a global scaling of activation across the entire ROI (Fig. 1)? To further

evaluate how feature dimension maps encoded within TO1/TO2 and hV4/VO1/VO2 are modulated

by task demands, we computed an inverted encoding model (IEM) for each region (Fig. 3; see

Materials and Methods; Sprague et al., 2018). We used an independent "mapping" task to

estimate a spatial encoding model for each voxel. The model was parameterized as a series of

weights across several smooth, overlapping spatial channels (Fig. 3A). The weights from the spatial

encoding model across all voxels that comprised a feature-selective ROI were then inverted which

allowed us to reconstruct spatial maps based on activation profiles from the feature-selective

attention task (Fig. 3B-C) averaged over 4.5-7.5 s after cue onset.

The reconstructed spatial maps (Fig. 5A) appear to have a stronger response at the aligned

stimulus location than the rest of the modelled visual field (for individual ROI results, see Appendix

Fig. A2). We quantified whether there was a stronger response to the stimulus by averaging map

activation profiles within a 1.5° radius disk (stimulus size) at the aligned stimulus location. We

could similarly assess map activation at the opposite screen location across fixation, which had no

stimulus present (and was never attended; Fig. 5B). A three-way permuted ANOVA with location

(stimulus and opposite), attention condition (attend fixation, color, and motion), and ROI

(hV4/VO1/VO2 and TO1/TO2) revealed a three-way interaction ($F(2,18) = 12.21$, $p < 0.001$, $\eta_p^2 =$

0.58). We conducted two-way permuted ANOVAs for each location with ROI (hV4/VO1/VO2 and

TO1/TO2) and attention condition (attend fixation, color, and motion) as factors to better

determine what was driving the three-way interaction. Like the univariate approach, the aim of

this analysis was to determine whether motion- and color-selective regions were modulated by

task demands and to determine if these regions differed in their representation depending on

which feature dimension was prioritized. Responses at the stimulus location showed a main effect

of attention condition ($F(2,18) = 52$, $p < 0.001$, $\eta_p^2 = 0.89$) and ROI ($F(1,9) = 7.31$, $p < 0.025$, $\eta_p^2 =$

0.78) as well as an interaction ($F(2,18) = 19.43$, $p < 0.001$, $\eta_p^2 = 0.68$). Follow-up permuted paired

sample t tests indicate that color-selective regions hV4/VO1/VO2 have a stronger neural response

when attending to stimulus color than when attending to fixation $t(9) = 7.69$, $p < 0.001$, $d_z = 2.43$.

There was also a stronger response when attending to stimulus motion than fixation ($t(9) = 5.45$,

$p < 0.001$, $d_z = 1.72$), consistent with spatial attention allocated to the stimulus location. Similarly,

motion-selective regions TO1/TO2 responded more strongly to the stimulus when attending to

either feature than when attending to fixation (fixation vs color: $t(9) = 5.38$, $p < 0.001$, $d_z = 1.70$;

fixation vs motion: $t(9) = 5.40$, $p < 0.001$, $d_z = 1.71$), again due to spatial attention.

A two-way ANOVA at the opposite location revealed no effect of ROI ($F(1,9) = 3.8$, $p = 0.131$, $\eta_p^2 =$

0.23) but did show a main effect of attention condition ($F(2,18) = 14.56$, $p < 0.001$, $\eta_p^2 = 0.81$) and

an interaction between ROI and attention condition ($F(2,18) = 3.93$, $p = 0.022$, $\eta_p^2 = 0.30$). Follow-

up permuted two-sample t tests reveal that there was a greater response at the opposite location

in motion-selective regions when attending to fixation than either feature dimension of the

stimulus (fixation vs color: $t(9) = 4.75$, $p = 0.001$, $d_z = 1.50$; fixation vs motion: $t(9) = 4.49$, $p = 0.001$,

$d_z = 1.42$). Color-selective regions had a greater response at the opposite location when attending

to fixation than attending to motion ($t(9) = 3.19$, $p = 0.008$, $d_z = 1.01$), but there was no difference

between the attend fixation and attend color conditions ($t(9) = 2.37$, $p = 0.04$, $d_z = 0.75$,

uncorrected). Thus, particularly for motion-selective regions, there are some task-dependent

modulations occurring at the location opposite of the stimulus in reconstructed maps. This

appears to be driven by an increased response at the opposite location when attending to fixation

than when attending to either feature of the stimulus in motion regions, which is likely due to

shifts of spatial attention (see Discussion).

Figure 5: Reconstructed spatial maps from color- and motion-selective regions are differentially modulated by feature-selective attention. **A.** Reconstructions from motion- and color-selective regions from the attend fixation, color, and motion conditions. Qualitatively, there is a strong response at the aligned location, particularly when attending to either the motion or color of the stimulus. **B.** We determined whether each region was modulated by task demands by assessing the strength of the stimulus representation for each stimulus condition as well as the opposite location. For both regions, the neural response at the stimulus location was stronger when attending to either feature of the stimulus as compared to attend fixation trials, indicating that stimulus representations in these regions track task-relevance of the stimulus location. At the opposite location, responses were stronger when attending to fixation. **C.** Difference in map activation between the stimulus and opposite locations for each stimulus condition, which indexes spatial specificity of map activation profiles. Responses were always greater when attending to the stimulus as compared to fixation for both regions. While color regions hV4/VO1/VO2 show a difference between the attend motion and color conditions ($p = 0.02$), indicating local enhancement, motion regions TO1/TO2 showed no difference ($p = 0.61$), indicating global enhancement. **D.** Difference in map activation at the stimulus location between the attend motion and attend color conditions. Responses reliably differed from zero for both ROIs, where motion-selective areas had stronger response when attending to motion and color

regions had a stronger response when attending to color. Responses differed between regions. * Significant difference based on permuted paired-samples t test ($p < 0.05$). Faint dots indicate individual subject means. Error bars indicate within-subject SEM.

Next, to evaluate the spatial selectivity of these modulations, we computed the difference in map

activation between the stimulus and opposite locations for each attention condition (Fig. 5C). A 2-

way repeated measures ANOVA with attention condition (attend fixation, color, and motion) and

ROI (TO1/TO2 and hV4/VO1/VO2) as factors reveal a main effect of ROI ($F(1,9) = 9.28$, $p = 0.018$,

$\eta_p^2 = 0.87$), attention condition ($F(2,18) = 56.92$, $p < 0.001$, $\eta_p^2 = 0.96$), and an interaction between

ROI and attention condition ($F(2,18) = 12.21$, $p = 0.006$, $\eta_p^2 = 0.58$). Follow up comparisons indicate

that in color-selective regions, the stimulus location was more strongly represented when

attending to a stimulus feature (fixation vs color: $t(9) = 7.35$, $p < 0.001$, $d_z = 2.33$; fixation vs motion:

$t(9) = 7.00$, $p < 0.001$, $d_z = 2.21$). Furthermore, the stimulus location was more strongly represented

when attending to color than when attending to motion ($t(9) = 2.78$, $p = 0.019$, $d_z = 0.88$), indicating

that feature-selective attention resulted in localized enhancement of the location of the stimulus

representation. Motion regions also exhibit a stronger stimulus representation when attention

was directed to the stimulus (attend color vs attend fixation: $t(9) = 8.5$, $p < 0.001$, $d_z = 2.69$

;attention motion vs attend fixation:$t(9) = 9.46$, $p < 0.001$, $d_z = 2.99$). However, there was no

difference between the attend color and motion conditions in the spatially selective stimulus

representation ($t(9) = 0.52$, $p = 0.610$, $d_z = 0.16$). Therefore, while TO1/TO2 is spatially selective,

in the sense that spatial attention to the stimulus location locally enhances its representation,

feature-selective attention results in a global shift in activation across the entire map (including

the opposite location with no stimulus present). This striking and unexpected divergence between

the modulatory profiles across color- and motion-selective regions suggests there may be unique

spatial profiles for attentional modulation by feature relevance in color- and motion-selective

regions.

Finally, we compared the AMI at the stimulus location (Fig. 5D), which indexes the difference in

activation across feature-selective attention conditions, within and between color- and motion-

selective ROIs. The motion-selective ROIs TO1/TO2 showed a significantly positive AMI (such that

the stimulus representation was stronger during attend-motion than attend-color trials, $t(9) =$

3.34, p = 0.006, d_z = 1.06), and the color-selective ROIs hV4/VO1/VO2 showed a significantly

negative AMI ($t(9)$ = 4.18, p < 0.001, d_z = 1.32). Importantly, the AMIs derived from each set of

regions significantly differed ($t(9)$ = 4.5, p < 0.001, d_z = 1.42). Altogether, this result supports a

dissociable modulation of activation profiles across color- and motion-selective ROIs based on the

attended feature dimension, with task-relevant feature-selective ROIs more strongly encoding

stimulus locations when they are relevant for task performance.

Discussion

The aim of the current study was to establish the spatial profile of goal-driven feature-selective

attention within neural feature dimension maps (Fig. 1). We presented an identical colorful moving

dot stimulus on each trial and cued participants to attend to either the color or motion of that

stimulus (Fig. 2). Our design isolated the influence of observer goals because the goal-relevant

feature dimension could change on each trial, but the visual input was consistent across the entire

experiment. Region-level univariate methods (Fig. 4 & Appendix Fig. A1) showed unique

modulations for color- and motion-selective regions: motion regions were feature-selective, but

not modulated by attention directed to a peripheral stimulus as compared to fixation; color-

selective regions were modulated by covert spatial attention, but this modulation was not feature-

selective. However, by using an inverted encoding model (Fig. 3) to reconstruct spatial maps from

both motion- and color-selective regions, we were able to show that stimulus representations in

both regions were modulated by task demands and were strongest when attending to each

region's preferred feature dimension (Fig. 5 & Appendix Fig. A2). Interestingly, different spatial

profiles were observed for each feature-selective region: enhancement in color regions was

strongly *localized* to the stimulus relative to the opposite location, while enhancement in motion

regions was not as strictly constrained to the stimulus location, consistent with some degree of

global enhancement of spatial responses. Not only do these findings support the role of each

region as a 'neural feature dimension map' (Thayer & Sprague, 2023), they extend previous work

by characterizing the unique spatial profiles of attentional modulation that emerge in color- and

motion-selective regions.

Previous univariate fMRI studies average across voxels with variable spatial selectivity
(Kriegeskorte & Bandettini, 2007; Serences & Saproo, 2012), making it difficult to ascertain the
spatial profile of attentional modulations across visual field maps encoded by feature-selective
regions. However, the multivariate approach implemented here revealed that color and motion
regions have unique spatial prioritization profiles. Namely, neural responses in color regions
primarily differed across attention conditions at the stimulus location (Fig. 5B-D), consistent with
a local enhancement of stimulus processing at task-relevant locations (Fig. 1). Motion regions also
showed a stronger response at the stimulus location when motion was attended (Fig. 5B/D), along
with some degree of enhancement at the opposite, task-irrelevant location (Fig. 5B/C). This is
consistent with a global enhancement of motion processing across all visual field locations when
motion is made task-relevant (Fig. 1), similar to previous findings for attention to a specific motion
value (Saenz et al., 2002; Serences & Boynton, 2007). Characterizing these spatial profiles of
prioritization due to feature-selective and spatial attention was only possible by assessing
multivariate activation patterns in visual field coordinates using IEM.

Our fMRI results are consistent with participants deploying spatial attention to the stimulus
location on attend-color/attend-motion trials, with a stronger observed response at the stimulus
location in both regions when attending to either stimulus feature relative to fixation (Fig. 5B). If
neural prioritization was solely driven by spatial attention, then multivariate responses in feature-
selective regions would be equivalent between the attend color and attend motion conditions as
all features at the locus of spatial attention are enhanced (Eckstein, 2011; A. Treisman, 1985; A.
M. Treisman & Gelade, 1980). Rather, our AMI results indicate that activation profiles in both
color- and motion-selective regions were reweighted by attention to their preferred feature
dimension (Fig. 5D), suggesting that feature-selective attention acted in concert with spatial
attention to sculpt activation profiles across each neural feature dimension map to support
dynamic task demands (Maunsell & Treue, 2006; Scolari et al., 2014). Even though we observed
clear feature preferences, it is important to highlight that these effects could have been reduced
because attention can spread to irrelevant feature dimensions of an object (O'Craven et al., 1997,
1999; M. A. Schoenfeld et al., 2003, 2014), meaning that neural responses may have been
enhanced even when attending to a region's non-preferred feature dimension.

Feature-selective enhancement occurs when a feature value is relevant, resulting in an increased

response from all feature-selective neural populations, regardless of spatial selectivity, and drives

a spatially global enhancement that has been observed for color (McAdams & Maunsell, 2000;

Saenz et al., 2002; Zhang & Luck, 2009), motion (Martínez-Trujillo & Treue, 2002; Schneider, 2011;

M. Schoenfeld et al., 2007; Serences & Boynton, 2007; Treue & Trujillo, 1999), and orientation

(Ester et al., 2016; McAdams & Maunsell, 2000). Interestingly, we observed more global

enhancement in motion regions when attending to motion, but a local enhancement in color

regions when attending to color. Notably, global prioritization through feature-selective attention

is primarily observed when there are multiple stimuli at distinct locations, where one stimulus is

relevant and another is irrelevant to current goals, and is strongest at the irrelevant stimulus

location when the feature value matches the relevant stimulus feature (e.g., Saenz et al., 2002). If

feature-selective attention modulates responses across the visual field, then all neural populations

tuned to the relevant feature should have increased firing, even when there is no stimulation. For

example, Serences & Boynton (2007) found evidence for successful decoding of the attended

motion direction when motion was goal-relevant from voxels with no visual input.

There is evidence to suggest that color and motion have different spatial profiles when feature-

selective attention is deployed. For instance, Leonard and colleagues (2015) instructed

participants to identify color-defined targets in a central letter stream. When distractors matched

the relevant color and were far away from the target, they weakly captured attention, but capture

gradually increased as distractors were closer to the target. Similar interactions between spatial

attention and feature-selective attention for color have been observed in attention and visual

search tasks, where feature-specific modulations can be restricted to a single stimulus location

when cued to direct spatial attention to a hemifield (Andersen et al., 2011; Berggren & Eimer,

2020; Diepen et al., 2016; Dodwell et al., 2025). This indicates that feature-selective attention for

color is spatially dependent (van Es et al., 2018), diverging from the canonical characterization of

spatially global prioritization of relevant features. In contrast, motion directions are almost always

prioritized globally. Numerous studies have shown that when a specific motion direction is

relevant, enhancement occurs throughout the visual field (Boynton et al., 2006; Mendoza et al.,

2011; Treue & Trujillo, 1999), even when parametrically manipulating the distance between target

and distractor motion stimuli (Liu & Mance, 2011), and when spatial attention is directed to other

locations (Bosworth & Dobkins, 2002; Dobkins & Bosworth, 2001; Melcher et al., 2004; Talipski et

al., 2022). Motion processing can even be improved when spatial attention is directed away from

a motion stimulus (Motoyoshi et al., 2015). Furthermore, one fMRI study found increased

responses in both hemifields in motion selective regions when attending to motion but only

contralateral responses increases in color selective regions when attending to color (Stoppel et al.,

2007). Accordingly, the evidence overall appears to point to color and motion being prioritized

differently by feature- and spatial-based attention, with a greater emphasis for local

enhancements of representations of color and global enhancement of representations of motion.

Our results may provide unique insight into the neural mechanisms that result in different spatial

activation patterns for stimuli containing task-relevant color and motion information. The IEM

approach employed here allowed for a demonstration of global modulation by feature-selective

attention in motion-sensitive regions TO1/TO2, and so the absence of a similar observation for

color-sensitive regions hV4/VO1/VO2 offers compelling evidence for a primarily local modulation

of stimulus representations when color was task relevant. One reason why motion regions may

implement global modulations while color regions locally enhance responses is that vRFs in

TO1/TO2 are much larger than hV4/VO1/VO2 (Wandell & Winawer, 2015). Thus, motion regions

may necessarily enhance processing across large swaths of the visual field when motion is task

relevant, providing a distinct neural mechanism for the differences in spatial prioritization

(McMains et al., 2007; van Es et al., 2018). This difference in encoding properties between regions

may stem from the idiosyncrasies of color and motion as feature dimensions. By definition, to

compute motion direction, the start and end point of an item are used to generate a direction

vector (Albright & Stoner, 1995; Borst & Egelhaaf, 1989). Therefore, the global enhancement

observed in the current study may be due to the specialization of TO1/TO2 in processing motion,

a unique feature dimension which necessitates integration of information across distinct locations.

Notably, the current study employed different stimulus selectivity requirements than many other

studies investigating attention to features. There was no requirement to prioritize a single location

over others in our paradigm because only a single stimulus was ever presented. Additionally, our

participants were instructed to attend to an entire feature dimension (e.g., color), rather than a

specific feature value (e.g., orange). Other paradigms often include multiple stimuli and instruct

participants to attend a specific feature value (Boynton et al., 2006; Mendoza et al., 2011; Treue

& Trujillo, 1999; Andersen et al., 2011; Dodwell et al., 2025; van Diepen et al., 2016; Stormer &

Alvarez, 2014). It is possible that the profile of attentional modulation would change across neural

feature dimension maps when multiple stimuli are competing for attentional priority (Desimone

& Duncan, 1995; Harrison et al., 2024; Sprague, Itthipuripat, et al., 2018), or when a specific

feature value is cued as relevant.

Competing models make divergent predictions as to how task relevance of specific feature

dimensions impacts activation profiles over feature dimension maps within the context of visual

search. Thus, it is necessary to characterize activation profiles across the entire visual field (Fig. 3)

to finely assess properties of feature dimension maps and adjudicate between these models.

'Dimensional weighting' models claim relevance enhances activation across entire feature

dimension maps in a spatially global manner (Found & Müller, 1996; Liesefeld & Müller, 2019,

2023; Müller et al., 1995), which is consistent with our observations that attention to stimulus

motion enhanced activation across the entire spatial extent of TO1/TO2. Other models claim that

a spatial 'attentional window' restricts how much of the visual field is prioritized (Theeuwes, 2010,

2023), which is consistent with spatial activation profiles in hV4/VO1/VO2. As such, it appears

aspects of both models are supported by the current findings. However, taken together, results

from the current study have interesting parallels to the dimension weighting priority map model.

Dimensional weighting appears to occur more weakly when color is the relevant feature dimension

(Liesefeld, Liesefeld, Pollmann, et al., 2019). It has been argued that this is because color is

comprised of multiple dimensions (e.g., luminance, red-green, and blue-yellow in CIE *Lab* color

space). Rather than considering color to uniquely contain multiple dimensions, as motion is

similarly multi-dimensional (e.g., velocity, direction, and complex motion patterns), it may be that

color is unique for other reasons, including peculiarities of neural architectures supporting local

processing within individual feature dimension maps. Perhaps the local modulations we observed

in hV4/VO1/VO2 are the neural substrate of the finding that task-driven enhancement of color

acts on a more localized portion of the visual field (Theeuwes, 2010, 2023), whereas our

observation of more global enhancement in TO1/TO2 when motion is attended supports robust

dimensional weighting across all spatial locations that is typically observed (Liesefeld, Liesefeld, &

Müller, 2019; Liesefeld & Müller, 2019; Sauter et al., 2018).

Previous reports (McMains et al., 2007; Runeson et al., 2013) investigated how individual color-

and motion-selective regions are modulated by task demands. Our findings were remarkably

consistent with those found by McMains and colleagues (2007). Univariate analyses in color-

selective region hV4 showed spatial selectivity, while the motion-selective region was not spatially

selective. Note, that the spatial analysis conducted by McMains et al. was very coarse, as it was

only able to show that the hemisphere contralateral to the stimulus had a greater response than

the ipsilateral hemisphere. To assay the full spatial profile of activation relative to the stimulus

location, multivariate techniques like the one implemented in the current study must be used.

Importantly, our IEM approach allowed for a fully randomized stimulus position on each trial,

minimizing the impact of spatial expectation effects, and eliminating the impact of any

idiosyncrasies of cortical sampling of different visual field locations. Additional similarities between

our results and those of McMains et al (2007) are present as there was no significant modulation

based on attended feature dimension in hV4, such that there was no greater univariate response

when color was goal relevant compared to motion (Appendix Fig. 1). Also converging with their

results, in our study, the more anterior color-selective regions VO1 and VO2 (parallel to TEO as

described in their study; Brewer et al., 2005) showed the primary evidence for color-selective

attentional modulations, with statistically less-significant effects observed in hV4 (Appendix Figs.

1, 2). Based on this finding, it could be that these anterior regions are especially subject to

attentional modulations. This appears to be unique to when color is goal-relevant, as hV4

qualitatively responds just as strongly, if not more strongly, to task-irrelevant but salient color

stimuli than VO1/VO2 in a previous study (Thayer & Sprague, 2023). This intriguing functional

distinction between extracting salient locations in posterior regions and relevant locations in

anterior regions, which echoes similar sharp emergence of top-down cognitive signals in

extrastriate cortex during visual working memory (Mendoza-Halliday et al., 2014), should be

explored in future work spanning additional manipulations of visual stimuli and task demand

conditions.

In sum, we found that feature-selective retinotopic regions in visual cortex track the location of

stimuli throughout the visual field and showed a stronger response to those stimuli when each

region's preferred feature was goal-relevant. Regions had different spatial modulation profiles,

with color regions locally enhancing activation at the stimulus location, and motion regions

showing a global enhancement of activation. These findings emphasize the role of TO1/TO2 and

hV4/VO1/VO2 as feature dimension maps and specify how they prioritize information in the visual

field, which can be exploited to fully understand how we successfully navigate our visual

environment.

References

References

Albright, T. D., & Stoner, G. R. (1995). Visual motion perception. *Proceedings of the National*

Academy of Sciences, *92*(7), 2433–2440. https://doi.org/10.1073/pnas.92.7.2433

Amano, K., Wandell, B. A., & Dumoulin, S. O. (2009). Visual field maps, population receptive field

sizes, and visual field coverage in the human MT+ complex. *Journal of Neurophysiology*,

102(5), 2704–2718.

Andersen, S. K., Fuchs, S., & Müller, M. M. (2011). Effects of Feature-selective and Spatial Attention

at Different Stages of Visual Processing. *Journal of Cognitive Neuroscience*, *23*(1), 238–246.

https://doi.org/10.1162/jocn.2009.21328

Awh, E., Belopolsky, A. V., & Theeuwes, J. (2012). Top-down versus bottom-up attentional control:

A failed theoretical dichotomy. *Trends in Cognitive Sciences*, *16*(8), 437–443.

Bahle, B., Thayer, D. D., Mordkoff, J. T., & Hollingworth, A. (2019). The architecture of working

memory: Features from multiple remembered objects produce parallel, coactive guidance

of attention in visual search. *Journal of Experimental Psychology: General*.

Baker, D. H., Vilidaite, G., Lygo, F. A., Smith, A. K., Flack, T. R., Gouws, A. D., & Andrews, T. J. (2021).

Power contours: Optimising sample size and precision in experimental psychology and

human neuroscience. *Psychological Methods*, *26*, 295–314.

https://doi.org/10.1037/met0000337

Beauchamp, M. S., Cox, R. W., & Deyoe, E. A. (1997). Graded Effects of Spatial and Featural

Attention on Human Area MT and Associated Motion Processing Areas. *Journal of*

Neurophysiology, *78*(1), 516–520. https://doi.org/10.1152/jn.1997.78.1.516

Beck, D. M., & Kastner, S. (2005). Stimulus context modulates competition in human extrastriate

cortex. *Nature Neuroscience*, *8*(8), Article 8. https://doi.org/10.1038/nn1501

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing

under dependency. *The Annals of Statistics*, *29*(4), 1165–1188.

https://doi.org/10.1214/aos/1013699998

Berggren, N., & Eimer, M. (2020). Spatial filtering restricts the attentional window during both

singleton and feature-based visual search. *Attention, Perception, & Psychophysics*, *82*(5),

2360–2378. https://doi.org/10.3758/s13414-020-01977-5

Bichot, N. P., Rossi, A. F., & Desimone, R. (2005). Parallel and Serial Neural Mechanisms for Visual

Search in Macaque Area V4. *Science*, *308*(5721), 529–534.

https://doi.org/10.1126/science.1109676

Bisley, J. W., & Goldberg, M. E. (2003). Neuronal activity in the lateral intraparietal area and spatial

attention. *Science*, *299*(5603), 81–86.

Bisley, J. W., & Goldberg, M. E. (2006). Neural Correlates of Attention and Distractibility in the

Lateral Intraparietal Area. *Journal of Neurophysiology*, *95*(3), 1696–1717.

https://doi.org/10.1152/jn.00848.2005

Borst, A., & Egelhaaf, M. (1989). Principles of visual motion detection. *Trends in Neurosciences*,

12(8), 297–306. https://doi.org/10.1016/0166-2236(89)90010-6

Bosworth, R. G., & Dobkins, K. R. (2002). The Effects of Spatial Attention on Motion Processing in

Deaf Signers, Hearing Signers, and Hearing Nonsigners. *Brain and Cognition*, *49*(1), 152–

169. https://doi.org/10.1006/brcg.2001.1497

Boynton, G. M., Ciaramitaro, V. M., & Arman, A. C. (2006). Effects of feature-based attention on

the motion aftereffect at remote locations. *Vision Research*, *46*(18), 2968–2976.

https://doi.org/10.1016/j.visres.2006.03.003

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Brewer, A. A., Liu, J., Wade, A. R., & Wandell, B. A. (2005). Visual field maps and stimulus selectivity

in human ventral occipital cortex. *Nature Neuroscience*, *8*(8), 1102–1109.

Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, *51*(13), 1484–1525.

https://doi.org/10.1016/j.visres.2011.04.012

Chawla, D., Rees, G., & Friston, K. J. (1999). The physiological basis of attentional modulation in

extrastriate visual areas. *Nature Neuroscience*, *2*(7), 671–676.

https://doi.org/10.1038/10230

Cook, E. P., & Maunsell, J. H. R. (2002). Attentional Modulation of Behavioral Performance and

Neuronal Responses in Middle Temporal and Ventral Intraparietal Areas of Macaque

Monkey. *Journal of Neuroscience*, *22*(5), 1994–2004.

https://doi.org/10.1523/JNEUROSCI.22-05-01994.2002

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in

the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215. https://doi.org/10.1038/nrn755

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review

of Neuroscience*, *18*(1), 193–222.

Diepen, R. M. van, Miller, L. M., Mazaheri, A., & Geng, J. J. (2016). The Role of Alpha Activity in

Spatial and Feature-Based Attention. *eNeuro*, *3*(5).

https://doi.org/10.1523/ENEURO.0204-16.2016

Dobkins, K. R., & Bosworth, R. G. (2001). Effects of set-size and selective spatial attention on

motion processing. *Vision Research*, *41*(12), 1501–1517. https://doi.org/10.1016/S0042-

6989(01)00038-4

Dodwell, G., Nako, R., & Eimer, M. (2025). EEG evidence for spatial selectivity in feature-based

preparation for visual search. *Biological Psychology*, *196*, 109016.

https://doi.org/10.1016/j.biopsycho.2025.109016

Dumoulin, S. O., & Wandell, B. A. (2008). Population receptive field estimates in human visual

cortex. *Neuroimage*, *39*(2), 647–660.

Eckstein, M. P. (2011). Visual search: A retrospective. *Journal of Vision*, *11*(5), 14–14.

https://doi.org/10.1167/11.5.14

Ester, E. F., Sutterer, D. W., Serences, J. T., & Awh, E. (2016). Feature-Selective Attentional

Modulations in Human Frontoparietal Cortex. *Journal of Neuroscience*, *36*(31), 8188–8199.

https://doi.org/10.1523/JNEUROSCI.3935-15.2016

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*

Methods, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Folk, C. L., Remington, R. W., & Johnston, J. C. (1992). Involuntary covert orienting is contingent

on attentional control settings. *Journal of Experimental Psychology: Human Perception and*

Performance, *18*, 1030–1044. https://doi.org/10.1037/0096-1523.18.4.1030

Found, A., & Müller, H. J. (1996). Searching for unknown feature targets on more than one

dimension: Investigating a "dimension-weighting" account. *Perception & Psychophysics*,

58(1), 88–101. https://doi.org/10.3758/BF03205479

Franz, V. H., & Loftus, G. R. (2012). Standard errors and confidence intervals in within-subjects

designs: Generalizing Loftus and Masson (1994) and avoiding the biases of alternative

accounts. *Psychonomic Bulletin & Review*, *19*(3), 395–404.

https://doi.org/10.3758/s13423-012-0230-1

Harrison, A., Thayer, D., & Sprague, T. (2024). Stimulus representations in neural priority maps are

equally enhanced by attention independent of the number of attended locations. *Journal*

of Vision, *24*(10), 1507. https://doi.org/10.1167/jov.24.10.1507

Huk, A. C., Dougherty, R. F., & Heeger, D. J. (2002). Retinotopy and functional subdivision of human

areas MT and MST. *Journal of Neuroscience*, *22*(16), 7195–7205.

Ikkai, A., & Curtis, C. E. (2008). Cortical Activity Time Locked to the Shift and Maintenance of Spatial

Attention. *Cerebral Cortex*, *18*(6), 1384–1394. https://doi.org/10.1093/cercor/bhm171

Itti, L., & Koch, C. (2001). Computational modelling of visual attention. *Nature Reviews*

Neuroscience, *2*(3), 194.

Kastner, S., De Weerd, P., Pinsk, M. A., Elizondo, M. I., Desimone, R., & Ungerleider, L. G. (2001).

Modulation of Sensory Suppression: Implications for Receptive Field Sizes in the Human

Visual Cortex. *Journal of Neurophysiology*, *86*(3), 1398–1411.

https://doi.org/10.1152/jn.2001.86.3.1398

Kastner, S., Pinsk, M. A., De Weerd, P., Desimone, R., & Ungerleider, L. G. (1999). Increased activity

in human visual cortex during directed attention in the absence of visual stimulation.

Neuron, *22*(4), 751–761.

Kay, K. N., Winawer, J., Mezer, A., & Wandell, B. A. (2013). Compressive spatial summation in

human visual cortex. *Journal of Neurophysiology*, *110*(2), 481–494.

Kriegeskorte, N., & Bandettini, P. (2007). Analyzing for information, not activation, to exploit high-

resolution fMRI. *NeuroImage*, *38*(4), 649–662.

https://doi.org/10.1016/j.neuroimage.2007.02.022

Leber, A. B., & Egeth, H. E. (2006). It's under control: Top-down search strategies can override

attentional capture. *Psychonomic Bulletin & Review*, *13*(1), 132–138.

https://doi.org/10.3758/BF03193824

Leonard, C. J., Balestreri, A., & Luck, S. J. (2015). Interactions between space-based and feature-

based attention. *Journal of Experimental Psychology: Human Perception and Performance*,

41(1), 11–16. https://doi.org/10.1037/xhp0000011

Liesefeld, H. R., Liesefeld, A. M., & Müller, H. J. (2019). Distractor-interference reduction is

dimensionally constrained. *Visual Cognition*, *27*, 247–259.

https://doi.org/10.1080/13506285.2018.1561568

Liesefeld, H. R., Liesefeld, A. M., Pollmann, S., & Müller, H. J. (2019). Biasing Allocations of Attention

via Selective Weighting of Saliency Signals: Behavioral and Neuroimaging Evidence for the

Dimension-Weighting Account. In T. Hodgson (Ed.), *Processes of Visuospatial Attention and*

Working Memory (pp. 87–113). Springer International Publishing.

https://doi.org/10.1007/7854_2018_75

Liesefeld, H. R., & Müller, H. J. (2019). Distractor handling via dimension weighting. *Current Opinion*

in Psychology, *29*, 160–167. https://doi.org/10.1016/j.copsyc.2019.03.003

Liesefeld, H. R., & Müller, H. J. (2023). *Target Salience and Search Modes: A Commentary on*

Theeuwes (2023) (1). *6*(1), Article 1. https://doi.org/10.5334/joc.279

Liu, T., & Mance, I. (2011). Constant spread of feature-based attention across the visual field.

Vision Research, *51*(1), 26–33. https://doi.org/10.1016/j.visres.2010.09.023

Luck, S. J., Gaspelin, N., Folk, C. L., Remington, R. W., & Theeuwes, J. (2020). Progress toward

resolving the attentional capture debate. *Visual Cognition*.

https://www.tandfonline.com/doi/full/10.1080/13506285.2020.1848949

Mackey, W. E., Winawer, J., & Curtis, C. E. (2017). Visual field map clusters in human frontoparietal

cortex. *Elife*, *6*, e22974.

Mante, V., Sussillo, D., Shenoy, K. V., & Newsome, W. T. (2013). Context-dependent computation

by recurrent dynamics in prefrontal cortex. *Nature*, *503*(7474), Article 7474.

https://doi.org/10.1038/nature12742

Martínez-Trujillo, J. C., & Treue, S. (2002). Attentional Modulation Strength in Cortical Area MT

Depends on Stimulus Contrast. *Neuron*, *35*(2), 365–370. https://doi.org/10.1016/S0896-

6273(02)00778-X

Maunsell, J. H., & Treue, S. (2006). Feature-based attention in visual cortex. *Trends in*

Neurosciences, *29*(6), 317–322.

Mazer, J. A., & Gallant, J. L. (2003). Goal-related activity in V4 during free viewing visual search.

Evidence for a ventral stream visual salience map. *Neuron*, *40*(6), 1241–1250.

https://doi.org/10.1016/s0896-6273(03)00764-5

McAdams, C. J., & Maunsell, J. H. R. (2000). Attention to Both Space and Feature Modulates

Neuronal Responses in Macaque Area V4. *Journal of Neurophysiology*, *83*(3), 1751–1755.

https://doi.org/10.1152/jn.2000.83.3.1751

McMains, S. A., Fehd, H. M., Emmanouil, T.-A., & Kastner, S. (2007). Mechanisms of feature-and

space-based attention: Response modulation and baseline increases. *Journal of*

Neurophysiology, *98*(4), 2110–2121.

Melcher, D., Crespi, S., Bruno, A., & Morrone, M. C. (2004). The role of attention in central and

peripheral motion integration. *Vision Research*, *44*(12), 1367–1374.

https://doi.org/10.1016/j.visres.2003.11.023

Mendoza, D., Schneiderman, M., Kaul, C., & Martinez-Trujillo, J. (2011). Combined effects of

feature-based working memory and feature-based attention on the perception of visual

motion direction. *Journal of Vision*, *11*(1), 11. https://doi.org/10.1167/11.1.11

Mendoza-Halliday, D., Torres, S., & Martinez-Trujillo, J. C. (2014). Sharp emergence of feature-

selective sustained activity along the dorsal visual pathway. *Nature Neuroscience*, *17*(9),

1255–1262. https://doi.org/10.1038/nn.3785

Moore, T., & Armstrong, K. M. (2003). Selective gating of visual signals by microstimulation of

frontal cortex. *Nature*, *421*(6921), 370–373.

Moran, J., & Desimone, R. (1985). Selective Attention Gates Visual Processing in the Extrastriate

Cortex. *Science*, *229*(4715), 782–784. https://doi.org/10.1126/science.4023713

Motoyoshi, I., Ishii, T., & Kamachi, M. G. (2015). Limited attention facilitates coherent motion

processing. *Journal of Vision*, *15*(13), 1. https://doi.org/10.1167/15.13.1

Müller, H. J., Heller, D., & Ziegler, J. (1995). Visual search for singleton feature targets within and

across feature dimensions. *Perception & Psychophysics*, *57*(1), 1–17.

https://doi.org/10.3758/BF03211845

O'Craven, K. M., Downing, P. E., & Kanwisher, N. (1999). fMRI evidence for objects as the units of

attentional selection. *Nature*, *401*(6753), 584–587.

O'Craven, K. M., Rosen, B. R., Kwong, K. K., Treisman, A., & Savoy, R. L. (1997). Voluntary Attention

Modulates fMRI Activity in Human MT–MST. *Neuron*, *18*(4), 591–598.

https://doi.org/10.1016/S0896-6273(00)80300-1

Ogawa, T., & Komatsu, H. (2004). Target Selection in Area V4 during a Multidimensional Visual

Search Task. *Journal of Neuroscience*, *24*(28), 6371–6382.

https://doi.org/10.1523/JNEUROSCI.0569-04.2004

Ogawa, T., & Komatsu, H. (2006). Neuronal dynamics of bottom-up and top-down processes in

area V4 of macaque monkeys performing a visual search. *Experimental Brain Research*,

173(1), 1–13. https://doi.org/10.1007/s00221-006-0362-5

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers

into movies. *Spatial Vision*, *10*(4), 437–442. https://doi.org/10.1163/156856897X00366

Poltoratski, S., Ling, S., McCormack, D., & Tong, F. (2017). Characterizing the effects of feature

salience and top-down attention in the early visual system. *Journal of Neurophysiology*,

118(1), 564–573. https://doi.org/10.1152/jn.00924.2016

Reynolds, J. H., & Desimone, R. (2003). Interacting Roles of Attention and Visual Salience in V4.

Neuron, *37*(5), 853–863. https://doi.org/10.1016/S0896-6273(03)00097-7

Runeson, E., Boynton, G. M., & Murray, S. O. (2013). Effects of task and attentional selection on

responses in human visual cortex. *Journal of Neurophysiology*, *109*(10), 2606–2617.

Saenz, M., Buracas, G. T., & Boynton, G. M. (2002). Global effects of feature-based attention in

human visual cortex. *Nature Neuroscience*, *5*(7), 631–632.

Sauter, M., Liesefeld, H. R., Zehetleitner, M., & Müller, H. J. (2018). Region-based shielding of visual

search from salient distractors: Target detection is impaired with same- but not different-

dimension distractors. *Attention, Perception, & Psychophysics*, *80*(3), 622–642.

https://doi.org/10.3758/s13414-017-1477-4

Schneider, K. A. (2011). Subcortical Mechanisms of Feature-Based Attention. *Journal of*

Neuroscience, *31*(23), 8643–8653. https://doi.org/10.1523/JNEUROSCI.6274-10.2011

Schoenfeld, M. A., Hopf, J.-M., Merkel, C., Heinze, H.-J., & Hillyard, S. A. (2014). Object-based

attention involves the sequential activation of feature-specific cortical modules. *Nature*

Neuroscience, *17*(4), 619–624. https://doi.org/10.1038/nn.3656

Schoenfeld, M. A., Tempelmann, C., Martinez, A., Hopf, J.-M., Sattler, C., Heinze, H.-J., & Hillyard,

S. A. (2003). Dynamics of feature binding during object-selective attention. *Proceedings of*

the National Academy of Sciences, *100*(20), 11806–11811.

https://doi.org/10.1073/pnas.1932820100

Schoenfeld, M., Hopf, J.-M., Martinez, A., Mai, H., Sattler, C., Gasde, A., Heinze, H.-J., & Hillyard, S.

(2007). Spatio-temporal Analysis of Feature-Based Attention. *Cerebral Cortex*, *17*(10),

2468–2477. https://doi.org/10.1093/cercor/bhl154

Scolari, M., Ester, E. F., & Serences, J. T. (2014). Feature- and Object-Based Attentional Modulation

in the Human Visual System. In A. C. (Kia) Nobre & S. Kastner (Eds.), *The Oxford Handbook*

of   Attention   (p.   0).   Oxford   University   Press.

https://doi.org/10.1093/oxfordhb/9780199675111.013.009

Serences, J. T., & Boynton, G. M. (2007). Feature-Based Attentional Modulations in the Absence

of   Direct   Visual   Stimulation.   *Neuron*,   *55*(2),   301–312.

https://doi.org/10.1016/j.neuron.2007.06.015

Serences, J. T., & Saproo, S. (2012). Computational advances towards linking BOLD and behavior.

Neuropsychologia, *50*(4), 435–446.

Serences, J. T., & Yantis, S. (2006). Selective visual attention and perceptual coherence. *Trends in*

Cognitive Sciences, *10*(1), 38–45.

Sprague, T. C., Adam, K. C., Foster, J. J., Rahmati, M., Sutterer, D. W., & Vo, V. A. (2018). Inverted

encoding models assay population-level stimulus representations, not single-unit neural

tuning. *eNeuro*, *5*(3).

Sprague, T. C., Boynton, G. M., & Serences, J. T. (2019). *Inverted encoding models estimate sensible*

channel responses for sensible models. https://doi.org/10.1101/642710

Sprague, T. C., Ester, E. F., & Serences, J. T. (2016). Restoring latent visual working memory

representations in human cortex. *Neuron*, *91*(3), 694–707.

Sprague, T. C., Itthipuripat, S., Vo, V. A., & Serences, J. T. (2018). Dissociable signatures of visual

salience and behavioral relevance across attentional priority maps in human cortex.

Journal of Neurophysiology, *119*(6), 2153–2165.

Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in the human

occipital, parietal and frontal cortices. *Nature Neuroscience*, *16*(12), 1879.

Stoppel, C. M., Boehler, C. N., Sabelhaus, C., Heinze, H.-J., Hopf, J. M., & Schoenfeld, M. A. (2007).

Neural mechanisms of spatial- and feature-based attention: A quantitative analysis. *Brain*

Research, *1181*, 51–60. https://doi.org/10.1016/j.brainres.2007.07.019

Störmer, V. S., & Alvarez, G. A. (2014). Feature-Based Attention Elicits Surround Suppression in

Feature Space. *Current Biology*, *24*(17), 1985–1988.

https://doi.org/10.1016/j.cub.2014.07.030

Swisher, J. D., Halko, M. A., Merabet, L. B., McMains, S. A., & Somers, D. C. (2007). Visual

Topography of Human Intraparietal Sulcus. *The Journal of Neuroscience*, *27*(20), 5326–

5337. https://doi.org/10.1523/JNEUROSCI.0991-07.2007

Talipski, L. A., Goodhew, S. C., & Edwards, M. (2022). No effect of spatial attention on the

processing of a motion ensemble: Evidence from Posner cueing. *Attention, Perception, &*

Psychophysics, *84*(6), 1845–1857. https://doi.org/10.3758/s13414-021-02392-0

Thayer, D. D., & Sprague, T. C. (2023). *Feature-specific salience maps in human cortex* (p.

2023.03.29.534828). bioRxiv. https://doi.org/10.1101/2023.03.29.534828

Theeuwes, J. (2010). Top–down and bottom–up control of visual selection. *Acta Psychologica*,

135(2), 77–99. https://doi.org/10.1016/j.actpsy.2010.02.006

Theeuwes, J. (2023). *The Attentional Capture Debate: When Can We Avoid Salient Distractors and*

When Not? (1). *6*(1), Article 1. https://doi.org/10.5334/joc.251

Treisman, A. (1985). Preattentive processing in vision. *Computer Vision, Graphics, and Image*

Processing, *31*(2), 156–177. https://doi.org/10.1016/S0734-189X(85)80004-9

Treisman, A. M., & Gelade, G. (1980). A feature-integration theory of attention. *Cognitive*

Psychology, *12*(1), 97–136.

Treue, S., & Trujillo, J. C. M. (1999). Feature-based attention influences motion processing gain in

macaque visual cortex. *Nature*, *399*(6736), 575–579.

van Es, D. M., Theeuwes, J., & Knapen, T. (2018). Spatial sampling in human visual cortex is

modulated   by   both   spatial   and   feature-based   attention. *eLife*,   *7*.

https://doi.org/10.7554/eLife.36928

Vo, V. A., Sprague, T. C., & Serences, J. T. (2017). Spatial tuning shifts increase the discriminability

and fidelity of population codes in visual cortex. *Journal of Neuroscience*, *37*(12), 3386–

3401.

Wandell, B. A., Dumoulin, S. O., & Brewer, A. A. (2007). Visual field maps in human cortex. *Neuron*,

56(2), 366–383.

Wandell, B. A., & Winawer, J. (2015). Computational neuroimaging and population receptive fields.

Trends in Cognitive Sciences, *19*(6), 349–357.

Winawer, J., & Witthoft, N. (2015). Human V4 and ventral occipital retinotopic maps. *Visual

Neuroscience*, *32*, E020. https://doi.org/10.1017/S0952523815000176

Witkowski, P. P., & Geng, J. J. (2023). Prefrontal Cortex Codes Representations of Target Identity

and   Feature   Uncertainty. *Journal   of   Neuroscience*,   *43*(50),   8769–8776.

https://doi.org/10.1523/JNEUROSCI.1117-23.2023

Wolfe, J. M. (1994). Guided search 2.0 a revised model of visual search. *Psychonomic Bulletin &

Review*, *1*(2), 202–238.

Wolfe, J. M., & Horowitz, T. S. (2004). What attributes guide the deployment of visual attention

and how do they do it? *Nature Reviews Neuroscience*, *5*(6), 495.

Wolfe, J. M., & Horowitz, T. S. (2017). Five factors that guide attention in visual search. *Nature*

Human Behaviour, *1*(3), 1–8. https://doi.org/10.1038/s41562-017-0058

Zhang, W., & Luck, S. J. (2009). Feature-based attention modulates feedforward visual processing.

Nature Neuroscience, *12*(1), 24–25. https://doi.org/10.1038/nn.2223

1007

1008

 <u>APPENDIX</u>

Figure A1: BOLD response across all voxels for individual regions of interest show selective attentional modulation based on feature preferences. A. Mean BOLD response for each attention condition from all individual retinotopic regions (feature-selective retinotopic regions hV4, VO1, VO2, TO1, and TO2, presented in aggregate in Fig. 4, are highlighted). Both individual motion-selective regions showed no response modulation when attending to either feature of the stimulus (color or motion) relative to when attending to fixation (all p's > 0.05). Each color-selective region was modulated by task demands, with a greater region-level response when attending to the color ($p < 0.05$) or the motion ($p < 0.05$) relative to attending to fixation. B. Attention modulation index for each individual feature-selective region. Both motion-selective regions had a positive response that differed from zero (TO1 and TO2: $p < 0.05$, uncorrected), indicating a stronger response when attending to motion. Color-selective regions VO1 and VO2 had a negative response that differed from zero (VO1 and VO2: $p < 0.05$, uncorrected), indicating a preference for color. However, hV4 did not differ from zero ($p > 0.05$). * indicate significant differences based on permuted paired or single sample t tests ($p < 0.05$), corrected for multiple comparisons via FDR across 13 ROIs. Gray asterisks indicate trends, defined based on p < 0.05 without correction. Error bars indicate within-subject SEM. Faint dots indicate individual subject means. Col: color, Mot: motion, Fix: fixation.

1010

1011

Figures 4, 5	Attend Fixation	Attend Color	Attend Motion	Fixation vs Color	Fixation vs Motion	Color vs Motion	AMI
V1	0.122	0.333	0.582	0.605	0.151	0.343	0.365
V2	**0.006**	0.138	0.176	*0.042*	**0.005**	0.308	0.286
V3	0.061	0.938	0.483	**0.006**	**0**	0.321	0.327
V3ab	0.249	0.041	**0.012**	*0.032*	**0.018**	0.848	0.874
hV4	0.054	**0**	**0**	**0.002**	**0.013**	0.212	0.224
VO1	**0.023**	**0**	**0**	**0**	**0**	**0.016**	*0.013*
VO2	*0.027*	**0**	**0**	**0**	**0.004**	*0.031*	*0.029*
LO1	**0**	**0**	**0**	0.234	0.073	0.781	0.792
LO2	**0**	**0**	**0**	0.204	0.106	0.980	0.974
TO1	**0**	**0**	**0**	0.090	0.234	**0.020**	*0.024*
TO2	**0**	**0**	**0**	*0.035*	0.771	**0.017**	*0.014*
TO1/TO2	**0**	**0**	**0**	0.067	0.598	**0.014**	**0.024**
hV4/VO1/VO2	**0.029**	**0**	**0**	**0**	**0**	0.055	0.061
IPS0/IPS1	**0**	**0**	**0**	0.506	0.617	0.783	0.788
IPS2/IPS3	**0**	**0**	**0**	0.758	0.302	0.536	0.555

Table 1. Statistics for univariate analyses. *P* values for univariate comparisons for individual attention conditions from zero and between specific attention conditions for each ROI. Comparisons include paired-sample *t* tests between the attend fixation, color, and motion conditions. One-sample *t* tests for individual attention conditions and AMI are also included. Even though the Color vs Motion and AMI analysis are effectively the same comparison, *p* values are slightly different due to the random shuffling procedure used to generate non-parametric distributions for statistical comparison (see statistical procedures above) and different number of comparisons when doing FDR corrections (78 for Color/Motion/Fixation comparisons; 13 for AMI). Bold denotes significance after FDR correction for all comparisons (*q* = 0.05). Significance before FDR corrections using α = 0.05 denoted with italics.

1012

1013

Figure A2: Reconstructed spatial maps for individual retinotopic ROIs. A. Reconstructions from all individual retinotopic ROIs, for the attend fixation, color, and motion conditions. Qualitatively, there is a strong response at the aligned location, particularly when attending to either the motion or color of the stimulus. **B.** We determined whether each region was modulated by task demands by assessing the strength of the stimulus representation for each stimulus condition as well as the opposite location. In all feature-selective regions, the neural response at the stimulus location was stronger when attending to either feature of the stimulus as compared to attend fixation trials, indicating that each region tracks task-relevance of the stimulus location. At the opposite location, responses were often strongest when attending to fixation. **C.** Difference in map activation between the stimulus and opposite locations for each stimulus condition. Responses were always greater when attending to either the color or motion of the stimulus for all regions (all p's < 0.05). **D.** Difference in map activation between the attend motion and attend color conditions. Responses reliably differed from zero for TO1, VO1, and VO2, where TO1 had stronger response when attending to motion and VO1 and VO2 had stronger responses when attending to color. * Significant

difference based on permuted paired-samples *t* test ($p < 0.05$), corrected for multiple comparisons via FDR. Gray asterisks indicate trends, defined as p < 0.05, uncorrected. Error bars indicate within-subject SEM. Faint dots indicate individual subject means.

1014

1015

Figures 6, 7	Stimulus Location			Opposite Location			Location Difference			
	Fixation vs Color	Fixation vs Motion	Color vs Motion	Fixation vs Color	Fixation vs Motion	Color vs Motion	Fixation vs Color	Fixation vs Motion	Color vs Motion	AMI
V1	**0.023**	**0.004**	0.856	**0**	**0**	0.79	**0.003**	**0**	0.778	0.853
V2	**0**	**0**	**0.036**	**0.011**	**0.003**	0.294	**0**	**0**	0.619	0.055
V3	**0**	**0**	0.114	**0.017**	**0**	0.241	**0**	**0**	0.324	0.11
V3ab	**0**	**0**	**0**	*0.044*	**0.006**	0.616	**0**	**0**	0.185	**0**
hV4	**0**	**0**	0.055	0.351	0.234	0.991	**0**	**0**	0.087	0.064
VO1	**0**	**0**	**0.023**	0.084	0.057	0.701	**0**	**0**	0.099	*0.024*
VO2	**0**	**0.006**	**0.002**	0.066	**0.003**	0.83	**0**	**0.002**	0.106	**0**
LO1	**0**	**0**	*0.048*	**0.002**	**0.003**	0.919	**0**	**0**	0.157	*0.047*
LO2	**0**	**0.008**	0.588	**0.015**	*0.037*	0.949	**0**	**0.007**	0.613	0.623
TO1	**0.007**	**0**	**0.042**	**0.005**	**0.002**	0.240	**0**	**0**	0.912	*0.024*
TO2	**0**	**0.005**	0.312	**0.011**	**0.008**	0.749	**0**	**0**	0.881	0.353
TO1/TO2	**0**	**0**	**0.011**	**0**	**0**	0.619	**0**	**0**	0.610	**0.006**
hV4/VO1/VO2	**0**	**0**	**0.005**	0.040	**0.008**	0.748	**0**	**0**	**0.019**	**0**
IPS0/IPS1	**0**	0.071	0.080	0.084	**0**	0.555	**0**	**0.005**	0.280	0.076
IPS2/IPS3	0.085	0.623	**0.004**	0.523	0.277	0.422	0.084	0.908	0.055	**0.005**

Table 2. Statistics for multivariate analyses. *P* values for IEM comparisons between specific attention conditions at certain locations in the visual field for each ROI. Comparisons include paired-sample *t* tests between the attend fixation, color, and motion conditions. One-sample *t* tests for AMI are also included. Even though the Color vs Motion at the stimulus location and AMI analyses are effectively the same comparison, *p* values are slightly different due to the random shuffling procedure used to generate non-parametric distributions for statistical comparison (see statistical procedures above) and different number of comparisons when doing FDR corrections (39 for Fixation/Color/Motion comparisons; 13 for AMI). Bold denotes significance after FDR correction for all comparisons (*q* = 0.05). Significance before FDR corrections using α = 0.05 denoted with italics.