Super compensatory substitutions restore protein function and flatten fitness landscapes

Wenhong Jiang[1,5†][*], Hengchuang Yin[2,5†], Saad S. Khan[3,5†], Yiparemu Abuduhaibaier[4] and Lucas B. Carey[5,6*]

[1] Key Laboratory of Fermentation Engineering (Ministry of Education), National 111 Center for Cellular Regulation and Molecular Pharmaceutics, Hubei University of Technology, Wuhan, Hubei, 430068, China

[2] Xinjiang Technical Institute of Physics and Chemistry, Chinese Academy of Science, Urumqi, China

[3] Departments of Bioengineering, Biology, Chemistry and Chemical Biology, Single Cell Proteomics Center, and Barnett Institute, Northeastern University, Boston, MA 02115, US

[4] Henan University School of Stomatology, Kaifeng, China

[5] Center for Quantitative Biology, Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[6] Pando Bioscience, Watertown, MA.

†. These authors have contributed equally to this work and share first authorship.
*. Corresponding author: Wenhong Jiang: jiangwh@pku.edu.cn ; Lucas B. Carey: lucas.carey@gmail.com

Classification

Major: Biological Sciences

Minor: Genetics

Keywords

compensatory mutations, fitness landscape, deep mutational scanning, evolution, protein stability

Significance

Compensatory mutations allow deleterious mutations to persist and even fix in populations, shaping protein evolution. Yet, comprehensive analyses of their prevalence and mechanisms have been lacking. Through a systematic analysis of large‐scale deep mutational scanning (DMS) datasets, we identified a class of super compensatory substitutions that enhance fitness across diverse genetic backgrounds. Our results reveal a previously unrecognized mode of compensatory effects that promote mutational robustness in evolving proteins.

Abstract

Compensatory mutations can mitigate the fitness costs of deleterious mutations, yet their distribution, structural contexts, and evolutionary roles remain poorly characterized. Here, we systematically analyzed a public deep mutational scanning (DMS) dataset comprising ~400,000 amino acid variants of imidazole glycerol-phosphate dehydratase (IGPD), with substitutions spanning the length of the protein. We found that roughly one third of otherwise nonfunctional missense variants can regain function through additional substitutions, predominantly at solvent exposed residues with weak local contacts. Notably, we identified a distinct class of super compensatory substitutions that enhance fitness across diverse genetic backgrounds and buffer mutational effects, effectively flattening the local fitness landscape. Comparative analyses of additional public DMS datasets across different proteins and assays revealed that super compensators recur across natural sequence–function maps, indicating a broadly represented mode of compensatory effects that sustains mutational persistence and contributes to protein evolutionary robustness.

Introduction

Most random amino acid mutations destabilize proteins and impair their function (1,2). Nevertheless, deleterious mutations can accumulate and persist within populations through multiple mechanisms which offset fitness costs. Epistatic interactions can mask fitness defects in specific genetic contexts; a mutation that is highly deleterious in one genetic background can become nearly neutral in the presence of another mutation, permitting the accumulation of otherwise deleterious variants (3-6). Proteins can also exhibit mutational robustness, maintaining structural stability and function despite acquiring destabilizing mutations (7, 8). At the organismal level, allelic redundancy can buffer fitness defects when deleterious mutations affect only a subset of gene copies (9, 10). Environmental pressures, such as antibiotic exposure, can render

otherwise deleterious mutations selectively advantageous, promoting their persistence in populations (11).

For deleterious nucleotide substitutions, fitness can sometimes be restored through direct reversion to the original sequence. However, when multiple mutations accumulate, or when selective pressures favor retention of the deleterious mutations, additional compensatory mutations provide an alternative route to fitness recovery. These mutations can restore organismal fitness to near wild-type (WT) or even WT levels (Fig. 1 *A*), a phenomenon documented across diverse biological systems (6, 11-13). Notably, compensatory mutations can occur at higher frequencies than direct reversions (14), particularly following antibiotic-induced population bottlenecks (15-18).

Intragenic compensatory mutations broadly operate through two mechanistic modes: global structural stabilization and pairwise interaction repair. Due to their effect on protein stability, global compensatory mutations have weak spatial constrains relative to destabilizing mutations, provided they contribute to the same folded energetic unit. This spatial flexibility enables broad-spectrum functional rescue across multiple mutations positions (18). In contrast, pairwise compensatory mechanisms typically involve residues in direct three-dimensional contact, forming active-site interfaces, α-helix packing cores, or allosteric communication pathways (19). As a result, local compensation is often highly context specific (20) and can even become destabilizing in the absence of the partnered deleterious mutations (14).

Previous investigations of compensatory evolution have largely relied on small-scale experiments examining the effects of limited sets of genetic variants on individual macromolecules, including proteins (11, 12, 21, 22), mRNA (23, 24), and tRNA (25). These studies demonstrated that compensatory mutations can restore fitness, but also reported cases where recovery appeared to rely primarily on direct reversion (26, 27). Because only a small fraction of sequence space (< 100 genotypes per study) was explored, findings varied substantially across studies (13, 28, 29). This variability and limited sampling have impeded the identification of robust predictors of compensatory potential and obscured broader principles governing the resolution of deleterious mutations at the macromolecular level.

To address this gap, we analyzed a large-scale deep mutational scanning (DMS) dataset comprising ~400,000 amino acid variants of the yeast *HIS3* gene, which encodes the histidine

biosynthesis enzyme imidazoleglycerol-phosphate dehydratase (IGPD/His3p) (30). This dataset

provides near-complete mutational coverage (>95% of residues) and includes ~30,000 variants

that impair growth in histidine-deficient media. At a scale more than three orders of magnitude

larger than previous studies, this dataset enables a systematic exploration of the protein level

principles governing compensatory evolution. Our analysis shows that compensatory potential

can be predicted, in part, from protein structural features. Most notably, we identify a distinct

class of *super-compensator* substitutions that enhance fitness across diverse genetic backgrounds

and buffer the fitness effects of mutations, effectively flattening fitness landscapes. The

prevalence of these super-compensators is further supported by analyses of several additional

DMS datasets. Together, these findings position compensatory substitutions, particularly super

compensators, as significant contributors to the shaping of evolutionary trajectories and

maintenance of genetic diversity in populations.

Results

A DMS Dataset to Identify Intragenic Compensation

To systematically examine the full spectrum of compensatory mutations across an entire protein,

we analyzed a large□scale deep mutational scanning (DMS) library of the *Saccharomyces*

cerevisiae HIS3 gene (30). The *HIS3* coding sequence was divided into twelve 20–30 residue

segments, each subjected to localized mutagenesis. Fitness of all variants was quantified through

pooled competitive growth assays and normalized such that genotypes carrying premature stop

codons were assigned a fitness of zero and wild-type WT a value of one (Materials and Methods).

After filter the data for quality (Material and Methods), the final dataset comprised 372,867

amino acid variant genotypes. Notably, many of these variant combinations correspond to

naturally occurring alleles observed across 21 extant yeast species (Fig.□S1*A, B*). Among them,

31,701 (8.5%) genotypes displayed a relative fitness of zero and served as the foundation for

subsequent analyses (Fig.□S1*C*).

In our data, we find that unfit (fitness = 0) genotypes can be rescued by additional substitutions

(Fig. 1 *B*). To quantify the extent of rescue, we defined rescuability as the average fitness gained

by unfit genotypes upon acquiring one or more additional substitutions (Fig. 1 *C*; Materials and

Methods). Using this metric, we find that only a small subset of unfit genotypes exhibit

appreciable rescuability (Fig. 1 *D*).

To control for errors in fitness estimation, we used nonsense genotypes containing an internal

stop codon as a negative control. Functional rescue of such genotypes is implausible and indeed

we observed little to no rescuability among them in our data. Using their rescuability distribution,

we estimated a false discovery rate of 5% for rescuability values ≥ 0.02 and 1% for values

≥ 0.04 (Fig. 1 *E*). Based on these calibrations, we classified an unfit genotype as rescuable

when its rescuability was ≥ 0.1, identifying 3,750 (11.8%) such genotypes. Notably, comparable

rescuability cutoffs were obtained across a range of definitions for fit and unfit genotypes (Fig.

S1*D*) as well as under alternative analytical procedures (Fig. S1*E*).

Figure 1

Fig. 1. Widespread compensatory substitutions in a deep mutational scanning library. (A)
Schematic demonstrating modes of rescue of loss-of function genotypes via compensatory
substitutions. (B) An example of compensatory substitutions in which a genotype with the fitness of
zero (blue) is four substitutions away from WT and for which we measured the fitness of 23 genotypes
with additional substitutions. The green edges connect to rescuing genotypes, and the red edges
connect to non-rescuing genotypes. (C) Fitness data from panel (B) illustrating computations for
rescuability. The dark blue dot is a rescuable unfit genotype while black dots are genotypes which
cannot be rescued. Error bars denote the standard deviation in the estimation of fitness. (D) The
distribution of rescuability of all unfit genotypes. (E) Comparison of rescuability distributions among
genotype classes, N denotes synonymous variants used in fitness estimation for amino acid genotype.
Purple, all missense genotypes; blue, missense genotypes carrying ≥ 10 synonymous variants;
green, nonsense genotypes (premature stop codons) serving as a low rescuability control.
148

Characterization of Structural Features Governing Compensation Potential

149
150

As nonfunctional variants span nearly the entire protein (Fig. S1*F*), we used the His3p library to
evaluate structural features influencing intragenic compensation. To quantify compensatory
potential structurally, we defined position-wise rescuability as the mean rescuability of all unfit
genotypes carrying substitutions at a given residue (Fig. 2 *A*). Position-wise rescuability varied
markedly across residues, with solvent-exposed, core-distal positions, which tend to be weakly
evolutionarily constrained, exhibiting significantly higher rescuability (Fig. 2 *B*), consistent with
previous observations (29).
158

Figure 2

160 **Fig. 2. Rescuability correlates with protein structural features.** (A) The rescuability of each
161 position in the His3p monomer was calculated as the average rescuability of all genotypes with that

position substituted. Error bars represent the standard deviation. (B) Position-wise rescuability correlates with protein structural features and evolutionary conservation. (C) Rescuable genotypes exhibit lower predicted ΔΔG_fold values than non‑rescuable ones (one‑tailed Wilcoxon rank‑sum test). (D) Most rescuing genotypes show lower ΔΔG_fold values than the corresponding rescuable genotypes. (E) Among genotypes with more than 50% of substitutions located in subunit‑interface regions, rescuable genotypes display lower ΔΔG_bind values than non‑rescuable genotypes (one‑tailed Wilcoxon rank‑sum test). (F) For genotypes with more than 50% substitutions in subunit interface regions, most rescuing genotypes have smaller ΔΔG_bind than rescuable genotypes.

To investigate the mechanistic relationship between rescuability and structural features, we examined predicted changes in Gibbs free energy change of folding (ΔΔG_fold) for 10,619 representative genotypes. Across all variants, rescuable genotypes exhibited lower ΔΔG_fold than non-rescuable ones, indicating that severe structural destabilization constrains compensatory potential (Fig. 2 *C*; Fig. S2*A*). Moreover, the addition of compensatory mutations partially mitigated destabilization, while restoration to wild-type stability was uncommon (Fig. 2 *D*). Compensatory stabilization was most pronounced for moderately destabilized variants (ΔΔG_fold < 40), suggesting that structural stabilization contributes significantly, but not exclusively, to intragenic compensation.

Because His3p functions as a homo‑24‑mer complex, we also assessed stability changes at subunit interfaces, where substitutions can influence both monomer folding and inter‑subunit binding (31-33). For genotypes with more than 50% of substitution in interface regions, predicted changes in Gibbs free energy change of binding (ΔΔG_bind) mirror the folding analysis: rescuable genotypes had smaller destabilizing shifts, and compensatory substitutions further mitigated losses in binding affinity (Fig. 2 *E* and *F*; Fig. S2*B*). Together, these results suggest that intragenic compensation enhances protein robustness by alleviating destabilizing effects on both monomeric structure and multimeric assembly.

Super Compensatory Substitutions Increase the Fitness of Diverse Variants

Given the observed structural enrichment of rescuability, we next sought to understand modes of compensation across different genetic backgrounds. Specifically, we asked whether compensatory substitutions act in a genotype specific manner or whether some substitutions can restore fitness across a broad range of deleterious genotypes.

Fig. 3. Super compensatory substitutions increase fitness across diverse genetic backgrounds.
(A) A schematic figure quantifies the compensatory ability of A110D in the 115E background. For substitution A110D in non-WT state 115E, paired genotypes differing only at position 110 are compared. Fitness differences (Δf) are computed and genotype pairs with $\Delta f \geq 0.5$ are considered compensatory. The resulting compensatory ratio is represented as a directed edge from 115E to 110D. (B) Directed network of compensatory interactions. Nodes represent non-WT amino acid states and directed edges compensatory effects of substitutions. Edge shading indicates fraction of substituted genotypes considered compensatory. (C) Compensatory ability of individual substitutions, defined as the sum of incoming edge weights in (B). Only substitutions from Segment 1 are shown. (D) Distribution of compensatory ability across all substitutions, showing that only a small subset exhibit high compensatory capacity. (E) Neural-network model predicting fitness via a latent variable (fitness potential), computed as the sum of amino acid specific fitness impact scores mapped to fitness by a

non-linear transformation. (F) An example illustrating modulation of fitness and fitness potential by
substitutions: A110D (red) increases fitness potential and fitness, whereas K115E (blue) decreases
both. (G) Substitutions with higher compensatory ability tend to exhibit higher fitness impact scores
relative to WT.

To quantify these modes of compensation, we represented intragenic rescue relationships using a
directed graph in which nodes correspond to amino-acid states and directed edges encode
compensatory effects. An edge from amino acid state A to B indicates that introducing
substitution B increases the fitness of genotypes carrying state A. Edge weights represent the
fraction of such genotypes rescued beyond a defined fitness threshold ($\geq$ 0.5; Fig. 3 *A-B*). The
overall compensatory capacity of a substitution was defined as the sum of its incoming edge
weights, quantifying the breadth of rescue across heterogeneous genetic contexts. Thus, the
network captures how well each substitution can rescue genotypes with distinct existing amino
acid states.

This analysis revealed a small subset of super compensatory substitutions—specific amino acid
changes (e.g., A110D) capable of increasing the fitness of genetically diverse variants
(Fig. 3 *C* and *D*; Fig. S3*A*). The corresponding non WT amino-acid states resulting from
these substitutions (e.g., 110D) are hereafter referred to as super compensators.

To investigate why certain substitutions emerge as super compensatory, we employed a published
neural network model that assigns each amino acid state a fitness impact score (30),
quantifying its contribution to a latent variable termed fitness potential. The model maps fitness
potential to predicted fitness via a sigmoidal transformation (Fig. 3 *E* and *F*; Fig. S4). Within
this framework, each genotype's fitness potential equals the sum of the fitness impact scores of its
constituent amino acid states, providing an interpretable representation linking amino acid
state specific effects to overall fitness. Remarkably, super compensators exhibited substantially
higher fitness impact scores than their WT counterparts (Fig. 3 *F* and *G*). These elevated scores
enable super compensatory substitutions to increase fitness potential—and thus predicted
fitness—across diverse genetic backgrounds, offering an explanation for their broad
compensatory action.

240 Super Compensators Buffer Substitution effects

241 To understand how super compensators achieve broad compensation, we examined their
242 influence on the fitness effects of other substitutions. Specifically, we compared the fitness

effects of identical substitutions introduced into wild‑type (WT) versus super compensator backgrounds. Remarkably, replacing a WT amino acid with a super compensator substantially attenuated the fitness effect of both deleterious and beneficial substitutions at other positions, indicating a strong buffering effect (Fig. 4 *A*).

Figure 4

Fig. 4. Super compensators buffer the fitness effects of both deleterious and beneficial substitutions. (A) Representative examples showing that super compensator 110D attenuates the fitness effects of both a deleterious (K115T) and beneficial (V114I) substitution. (B-D) A non-super compensator (110Q, x-axis) shows no systemic effect on substitution fitness compared with the WT amino acid state (110A, y-axis). Paired fitness effects were evaluated using relative (B) and absolute (C) fitness difference and skewness of the absolute differences (D). (E-G) A super compensator (110D, x-axis) reduces the fitness effects of substitutions relative to the WT state (110A, y-axis). Comparisons using relative (E) and absolute (F) fitness difference reveal buffering through collapse toward the vertical and produce a positively skewed distribution of fitness effect differences (G). (H) Conceptual model linking buffering to the sigmoidal relationship between fitness potential and predicted fitness. Non-WT amino acid states with positive fitness impact scores (eg., 110D) can shift

genotypes towards a high-fitness plateau where fitness is insensitive to additional substitutions. States with negative impact scores (eg., 115E) can shift genotypes off the plateau, increasing sensitivity. (I) Amino acid states with higher fitness impact scores relative to WT exhibit greater buffering capacity, quantified as increased positive skewness of fitness effect differences (Spearman R= 0.78, p-value = 0). (J) Non-WT amino acid states with higher compensatory capacity similarly exhibit stronger buffering, as indicated by increased positive skewness (Spearman R= 0.67, p-value = 0). (K) Example of local landscape smoothing by a super compensator. 93V reduces fitness variability among neighboring variants, effectively flattening the local fitness landscape. (L) Non-WT amino acid states with higher fitness impact scores relative to WT are associated with more stable protein structures, reflected by lower $\Delta\Delta G\_bind$ values.

To quantify this buffering effect, we computed the fitness difference (Δf) of each substitution with the WT (Fig. 4 *B*, y axis) and the non-WT amino acid state (Fig. 4 *B*, x axis). If substitution effects were comparable in both contexts, data points would distribute symmetrically around the diagonal. Instead, for super compensators such as 110D, the distribution collapsed along the vertical axis (Fig. 4 *E*), demonstrating substantial dampening of substitution effects. We further quantified this behavior by calculating the absolute deviation between the x and y axes (Fig. 4 *C*, *F*), and assessing the skewness of this distribution. Substitutions in super compensator backgrounds exhibited both reduced absolute fitness effects and greater skewness relative to non super compensator contexts (Fig. 4 *D*, *G*), confirming the enhanced buffering capacity of super compensators.

To identify the underlying cause, we revisited the sigmoidal relationship between fitness potential and predicted fitness defined by the neural network model. Most genotypes in our library reside on the high fitness plateau, where substantial variation in fitness potential produces only minor change in predicted fitness (Fig. 4 *H*). Beneficial substitutions that push genotypes further along this plateau have reduced sensitivity toward fitness deviations due to additional substitutions, yielding positively skewed fitness effect distributions. Conversely, deleterious substitutions drive genotypes off the plateau, increasing sensitivity and generating negatively skewed distributions (Fig. 4 *H*). Across all genotypes, we found that amino acid states with higher fitness impact scores than their WT counterparts exhibited higher skewness in the distribution of absolute fitness effect differences (Fig. 4 *I*). Similarly, substitutions with higher compensatory ability showed correspondingly increased skewness (Fig. 4 *J*). These observations indicate that super compensators buffer both deleterious and beneficial substitution effects by operating within a flattened region of the fitness landscape.

Whereas prior studies have emphasized the ruggedness of protein fitness landscapes,
characterized by numerous local peaks and valleys (34-37), our results show that super
compensators can locally smooth these landscape. The super compensator 93V provides a clear
example: in variants containing substitutions at positions 66 and 86, 93V substantially
reduces fitness variability among neighboring variants, effectively flattening the local fitness
surface (Fig. 4*K*).

Super Compensators Stabilize Inter-residue Interactions

Building on the role of stability restoration in compensation, we next evaluated structural features
of super compensatory substitutions. In general, super compensators significantly increased
overall protein stability (Fig. 4 *L*; Table S1). For instance, 110D forms a salt bridge with R112,
effectively stabilizing a neighboring loop region. In contrast, substitutions exhibiting lower
fitness impact scores than the WT, such as I19D, were predicted to disrupt a β sheet and its
adjacent helix (Table S1).

Figure 5

310
Fig. 5. The Super compensator 189A buffers the fitness effects of deleterious substitutions. (A)
Amino-acid composition of the constructed DMS library. (B) Genotypes with premature stop codons
exhibit similar fitness as those without in the presence of histidine and show reduced fitness in its
absence. (C) Super compensators increase the fitness of most genotypes carrying random mutations.
(D) The average fitness difference between genotypes with 189A and with 189S is higher than zero
(blue), and the fitness difference is more pronounced among deleterious genotypes (orange).

Notably, most super compensators are spatially distant from the catalytic sites (> 8 Å), with
none located within 4 Å of the substrate (Fig. S5). This spatial separation supports a
mechanism of global stabilization rather than direct modulation of active site interactions.

Super Compensators Enhance Organisms' tolerance to Deleterious Mutations

To experimentally verify the functional consequences of super compensation, we performed deep
mutational scanning to determine whether super compensators confer mutational robustness by
buffering deleterious substitutions. We designed the library to contain two fully randomized
flanking positions (NNK codons) surrounding a central position encoding either the WT or the
super compensator S189A amino acid (Fig. 5 *A*). Under histidine deficient conditions,
variants carrying premature stop codons displayed severely reduced fitness, which returned to
WT levels upon histidine supplementation (Fig. 5 *B*), confirming the reliability of our assay.

Introduction of the super compensator 189A resulted in a marked fitness increase for 77% of
all variants in the library (Fig. 5 *C*). The rescue effect was strongest among genotypes harboring
mildly deleterious substitutions (Fig. 5 *D*). Together, these results provide direct experimental
evidence that super compensators can increase mutational tolerance by buffering fitness loss
caused by otherwise deleterious mutations.

Super Compensators are Prevalent across Diverse Proteins

To determine whether super compensators represent a general phenomenon beyond His3p, we
applied the same analytical framework to published DMS datasets from other randomly
mutagenized proteins assayed using diverse fitness readouts (38). This analysis revealed that
super compensators are broadly distributed across proteins and similarly buffer the fitness effects
of mutation in independent systems (Fig. S6).

Notably, fitness was defined differently across datasets, yet our findings were robust to these
differences. In His3p fitness was quantified by yeast growth rate, whereas in another dataset it
was protein stability in a cell free system (39) and in the case of GFP, using fluorescence intensity
as a proxy for protein folding and activity (40). Together, these results support a general model
in which super compensators exert their effects, at least in part, by enhancing protein stability.

Discussion

Our analysis reveals the principles governing intragenic compensation, delineating both the characteristics of rescuable genotypes and compensatory substitutions. We found that the compensatory potential of non-functional variants is strongly correlated with protein structural context.: substitutions at solvent-exposed, weakly spatially constrained positions are more easily rescued. This finding clarifies why Tomala et al. (13) identified a limited set of compensatory substitutions for temperature-sensitive mutations, which often reside in structurally critical regions, consistent with independent evidence that some deleterious mutations are fundamentally uncompensatable (27). Consistent with Tomala et al.'s results, we observe that deleterious mutations causing modest structural destabilization can be rescued, reinforcing our observation that the destabilizing effect of substitution in folded and conserved positions is stronger than those in the solvent-exposed positions (31, 41). Beyond structural context, we also demonstrate that amino acid physicochemical properties can serve as predictors for fitness. Specifically, when the AAindex values—a set of quantitative descriptors of amino acid attributes—in mutant variants approximates those of the WT, fitness recovery is observed (Fig. S7). That is to say, compensation can occur when mutations bring the AAindex values closer to those of the WT.

Focusing on the properties of compensatory substitutions themselves, we identified a distinct class of super compensatory substitutions capable of increasing fitness of diverse genetic backgrounds. This phenomenon parallels the concept of cryptic mutations (42), which facilitate the accumulation of otherwise deleterious substitutions. Our experimental validation directly demonstrates this capacity: introducing super compensators reduced deleterious effects of mutations, thereby increasing mutational robustness. The existence of super compensators can thus provide an alternative explanation toward the mutational robustness of proteins. Previous work by Guo et al (43) and György Abrusán et al (44) found mutational robustness to vary across protein secondary structures. This aligns with our observation of increased compensation at solvent exposed positions and indicates that structure context shapes both the emergence and effectiveness of compensatory mechanisms.

The evolutionary implications of super compensators may extend beyond laboratory assays. If super compensators can increase the fitness of diverse variants, they should leave a detectable signature in natural sequences by enabling the retention of mildly deleterious alleles. Consistent with this expectation, comparative analysis of 335 His3p orthologs revealed that super compensators frequently co-occur with mildly deleterious amino acid states across extant yeast species (Fig. S8). Moreover, the strength of the super compensator (its fitness impact score) was

positively correlated with the severity of its partner deleterious amino acid states (Fig. S8*B*). These observations provide a plausible explanation for the sequence divergence observed among His3p orthologs and suggests that super compensators may have contributed to natural genetic variation.

These insights have practical implications for protein engineering. The ability of super compensators to confer mutational robustness suggests that predicting and incorporating them could serve as an effective strategy for designing proteins with enhanced stability and expanded functions. Accordingly, prioritizing stabilizing substitutions represents a valuable component for protein design, consistent with prevailing approaches that emphasize predicting stability-increasing mutations (45-48).

While the His3p DMS dataset is invaluable, its limitations warrant consideration. First, its basis in extant sequence backgrounds may underrepresent severely deleterious substitutions. Second, segmenting the gene precluded the detection of inter-segment compensatory effects. Nevertheless, to our knowledge, the His3p library remains one of the largest and most comprehensive resources that contains multi-site mutants, making it ideal for probing structure-compensation relationships. Furthermore, we are able to recapitulate our results regarding the stabilizing role of Super compensator across other large-scale DMS datasets, including those based on GFP fluorescence (40) and other stability-directed assays (39). This cross-validation underscores the generalizability of our conclusions beyond His3p.

In summary, this work deepens our understanding of how compensatory evolution is constrained by protein structure, reveals how super compensators promote mutational robustness and sequence diversity, and highlights how these principles can inform rational strategies for protein engineering.

Materials and Methods

Experimental Design

The experimental methodology and analytical procedures were already fully described in the previous work (30). We briefly summarize the key steps as follows: first, plasmids containing each HIS3 gene variant encoding IGPD with distinct amino acid substitutions were constructed and transformed into Δ*HIS3 S. cerevisiae*. Mixed populations of yeast strains carrying different *HIS3* variants were then co-cultured in bulk competition assays. Deleterious

substitutions in IGPD resulted in slow growth and decline of variants from the population over
time, neutral or beneficial substitutions enabled rapid growth and eventual dominance.

Through competitive growth analysis, variants with missense substitutions exhibiting growth
rates comparable to or lower than premature stop codon-containing variants were assigned a
fitness value of zero. In contrast, WT genotypes and the majority of single-substitution
variants demonstrated fitness values of 1. The experimental design included two biological
replicates with two technical replicates per biological replicate. Following rigorous quality
control, which excluded data with a replicate correlation of < 0.85, a further filter was applied to
ensure robust fitness assessment. Segment 9 exhibited low correlation between biological
replicates and was therefore excluded from this work (30). Specifically, amino acid genotypes
represented by fewer than three synonymous nucleotide variants were discarded. The resulting
dataset consisted of genotypes with an average of seven amino acid substitutions from the
reference sequence. The fitness of each genotype was assessed by averaging nine synonymous
variants. As these synonymous mutants exhibited little to no fitness difference (30), this internal
replication provided high confidence.

Calculation of Rescuability

For each unfit genotype, we identified all genotypes containing the same set of substitutions plus
one or more additional substitutions (hereafter referred to as the superset, total count N). Fitness
for each amino acid genotype was estimated using synonymous nucleotide variants, yielding a
mean fitness (S) and standard deviation (S.D.). The 95% confidence interval for fitness was
defined as $S \pm 1.96 \times S.D.$ A superset genotype A was considered to compensate for genotype B
when its lower confidence bound exceeded the upper confidence bound of B, i.e., $S_A - 1.96 S.D._A \geq$
$S_B + 1.96 S.D._B$. For each unfit genotype, we calculated fitness difference ($\Delta f = S_A - S_B$) for all
compensating superset genotypes (N1 out of N). Rescuability was then defined as the sum these
Δf values normalized by the total superset count :

$$Rescuability = \frac{\sum_1^{N1} \Delta f}{N}$$

As an independent validation, we used a non-parametric approach using synonymous variants.
For each candidate compensating genotype A and unfit genotype B, we compared the distribution
of fitness values of their respective synonymous variant sets (G_A and G_B) using a two sample
Kolmogorov-Smirnov test ($\alpha=0.05$). Genotype A was considered compensatory if the fitness
distribution of G_A was significantly higher than G_B. Fitness difference (Δf) and rescuability
values were then computed analogously to the primary method.

Calculation of Structural and Evolutionary Metrics

WCN Score The WCN calculation was performed using established methodology from Lin CP et al.(49), with the formula:

$$\text{WCN}_i = \sum_{i \neq j}^{N} \frac{1}{r2(ij)}$$

where r_{ij} denotes the distance between the C_α geometric center of residues i and residues j in IGPD (N total residues). The WCN value for each position was calculated by averaging measurements across 24 monomeric structures.

RSA Score The RSA score for each position of IGPD was derived from the DSSP file (generated using IGPD's PDB file) (50). The RSA values were normalized using the amino acid-specific surface area reference established by Matthew et al. (51).

ΔΔG Prediction The ΔΔG associated with structural stability or interface binding following substitutions was calculated using the standard Rosetta (53) protocol with default parameter settings.

Conservation Score Conservation scores were derived using sequence entropy analysis of IGPD homologs. Monomeric IGPD sequences from 21 extant species were aligned using T-coffee Expresso (52). For each alignment position X, sequence entropy was calculated as:

$$H(X) = \sum_{x=0}^{20} P(X) log2P(X)$$

Where P(X) represents the probability of each amino acid at position X, estimated from the multiple sequence alignment. The conservation score for position X is defined as:

$$C(X) = (1-H(X))/(1-gap\_fr)$$

Where gap_fr quantifies the gap occurrence rate at position X.

Association of Physicochemical Features with Fitness and Compensation

For each genotype, we quantified physicochemical distance from the wild-type sequence as the difference in AAindex values (AAindex distance) relative to WT across 553 AAindex descriptors. Analyses were performed within mutation sets, defined as genotypes carrying substitutions at the same positions but differing in amino-acid identity. For each mutation set and AAindex descriptor, we computed Spearman correlations (ρ) between AAindex distance and corresponding fitness across genotypes. Mutation sets were required to contain at least 82 genotypes, corresponding to the minimum sample size needed to detect modest correlation (ρ=0.3) with α = 0.05 as determined by an a priori power analysis using G*Power 3.1 (54). Correlation P values were adjusted for multiple testing across all descriptors and mutation sets using the Benjamini–Hochberg false discovery rate (FDR), correlations with FDR < 0.05 were considered significant. For visualization, non-significant correlations were set to zero.

 Directed-Graph Representation of Compensatory Interactions

For each substitution at position i (S_i) and each non-WT amino acid state at distinct position j

(N_j), we identified pairs of genotypes that only differed at position i: $G_1 = (S_i, N_j)$ and $G_2 = (WT_i, N_j)$. The

compensatory ability of S_i with respect to N_j was quantified as the fraction of such genotype pairs for which

$\Delta f = f(G_1) - f(G_2) > 0.5$. These relationships were represented as directed edges from S_i to N_j in a

network graph, with edge weights corresponding to the fraction of rescued genotype pairs. The

visualizations produced for Figure 3*B* and Figure S3*A* were produced using NetworkX (55) in

Python.

Quantification of Buffering of Substitutions Fitness Effects

To assess how super compensators modulate the magnitude of fitness effect of other substitutions,

we considered only substitutions that produced a nonzero fitness effect. The fitness effect of a

substitution was defined as the difference in fitness between two genotypes that differ only at a

single position (for example, G_1=00000, G_2=00100, where 0 denotes WT amino acid and 1

denotes a non-WT amino acid). For genotype quadruplet (G1=00000, G2=00100, G3=00001,

G4=00101) with fitness values f1, f2, f3, f4, we calculated $|\Delta f1|=|f2 - f1|$ and $|\Delta f2|=|f4 - f3|$.

Buffering of fitness effects by a non-WT amino acid state was quantified as the skewness of the

distribution of $|\Delta f1|$- $|\Delta f2|$, which captures the tendency to reduce the magnitude of fitness effects

of substitutions.

DMS Library Construction

To experimentally validate the predictions that super compensators can increase mutational

robustness, one representative super compensator was selected, that is S189A. The sites for

random mutagenesis were selected randomly, which are positions 181 and 202.

Plasmid Construction To construct the libraries, we first identified naturally occurring, or, when

necessary, inserted restriction sites that were in close proximity to the desired super compensators

sites and mutagenesis sites within the *S. cerevisiae HIS3* gene. Single-stranded oligonucleotides

that coded for the desired amino acid substitutions (including all synonymous nucleotide variants)

were synthesized with 15-25 bp homology arms flanking restriction sites.

Genomic DNA was extracted (Multi Sciences, MK010001) from the prototrophic S288C derived

FY4 strain. The *HIS3* locus (ATG – 250 bp, TAG + 300 bp) was then amplified using 500 ng of

the extracted DNA, primers 376, 379, and the Golden Star T6 DNA polymerase (Tsingke

Biotech, TSE101). The amplicon was PCR purified (Qiagen MinElute, 28004) and inserted

between the SacI and KpnI sites (destroying them) within the pGREG506 ADH1pr-MCM-

mCherry vectors (a gift from the Chao Tang's lab) via Gibson Assembly (Tiangen Easygeno,

VI201) to yield plasmid p142. After PCR, both amplicons were verified, purified, and assembled

into the vector backbone to yield plasmid p143.

Library Construction To construct the final libraries, 7.5 μg of p142 and p143 were digested for

4 hours at 37°C with 9 μL of KpnI, NheI (NEB, R3131S), and BstXI (NEB, R01131S) and EagI

(NEB, R3505S) respectively. Each reaction was then loaded on a 0.7% agarose gel, with 0.5X

TAE running buffer, gel purified (Magen, D2111), and concentrated (Tianmo Biotech, TD413).

Subsequently, the library ssOligos (primers 337, 338) were then assembled into 500 ng of the

digested vectors to a final concentration of 0.25 μM in 15 μL Gibson Assembly reactions (2 step

Gibson Assembly, 3 minutes at 37°C, 60 minutes at 50°C). Finally, the assembly reactions were

split into 5 transformations in chemically competent *E. Coli* (Transgen Biotech, CD101) and

spread onto LB+Amp plates, and incubated overnight (~14 hours) at 37°C. After incubation, the

colonies were verified (colony PCR, Sanger sequencing) and scraped off the plates (~55,000

colonies/library), mixed and the library plasmids were extracted (Tiangen Biotech, DP106),

quantified, and stored.

Yeast Transformation and Bulk Competition Competent yeast cells were prepared (Zymo

Research, T2001) using yeast strain Y120 (derived from UCC8363; MKT1-RM SAL1-RM

CAT5-RM MIP1-BY ura3Δ0 his3:KanMX). Subsequently, 5 μg of each plasmid was transformed

into ~52.5 x 10^6 cells and incubated at 30°C, 220 RPM for 50 hrs in 1L synthetic complete

dropout medium (SCD) lacking Uracil (-Ura).

After transformation (yielding ~750,000 colonies/library), ~3 x 10^8 cells per library were

inoculated into 1 L of SCD-Ura and grown at 30°C, 220 RPM for 6 hrs. After that, ~3 x 10^7 cells

per library were inoculated into 1 L SCD-His (Histidine) and ~2.1 x 10^7 cells into 700 mL SCD-

Ura and incubated at 30°C, 220 RPM. The competition was carried out for 120 hours with 12

hours between bottlenecks (2.1 x 10^7 cells per library transferred into fresh media), two samples

of 15 mL (technical replicates) were taken from each culture every 24 hours, centrifuged (3000 g,

5 mins) and stored (-20°C), the experiment from transformation to completion was carried out

twice (biological replicates).

Plasmid Extraction and Illumina Sequencing After competition, cell pellets from each time

point were thawed and plasmids extracted using the Tiangen Midiprep kit (DP106). Extracted

DNA (~10 ng) was labeled and amplified using staggered primers (primers 345-356) and Q5

DNA polymerase (NEB, M0492S) in 50 μL reactions (98°C for the 30 s; 98°C for 10 s, 55°C for

30 s and 72°C for 30 s for 17 cycles; 72°C for 2 min). The amplicons were purified (Qiagen

MinElute, 28004) and pooled samples were submitted to Genewiz company for Illumina library

construction and sequencing.

Data Processing Raw paired-end FASTQ files were merged using FLASH (56), demultiplexed

according to the unique primer pairs used to amplify each sample at specific timepoint, and reads

for each genotype were counted using starcode (57). Reads for each genotype were the sum of all

synonymous nucleotide genotypes and the sum across two technical replicates. Fitness for each

genotype was calculated using PyFitSeq (58), and the average fitness across both biological

replicates was used for further analysis.

Prevalence Analysis of Super Compensators

DMS datasets for 217 distinct proteins were retrieved from the ProteinGym database (38). The

datasets contain quantitative fitness measurement derived from diverse biochemical and cellular

readouts, including protein stability, enzymatic activity, and bacterial growth rate. The

established computational pipeline for identifying super compensators was then applied to these

datasets to detect potential super compensators and quantify the buffering ability against

substitutions.

Data, Materials, and Software Availability

557   All data and codes used for analysis and creating figures are available on Zenodo at

558   https://doi.org/10.5281/zenodo.17769277

Acknowledgments

We thank Dr. Chu Wang and Dr. Luhua Lai at Peking University for helpful discussions and

comments on the manuscript. This work is funded by China Postdoctoral Science Foundation

(2025M782617 to W.H.J.), the Natural Science Foundation of Hubei Province (JCZRQNB202600450

to W.H.J), Open Project Funding of the Key Laboratory of Fermentation Engineering (Ministry of

Education) (202509FE11 to W.H.J.), the Natural Science Foundation of Xinjiang Uygur Autonomous

Region (2025D01A136 to H.C.Y.). Additional funds to L.B.C. are provided by Peking-Tsinghua

Center for Life Sciences.

Author contributions: L.B.C. and W.H.J. designed the study; W.H.J., S.S.K., H.C.Y., Yiparemu and

L.B.C. carried out the experiments and analyzed data; W.H.J., H.C.Y., S.S.K. and L.B.C wrote the

manuscript. All the authors read and accepted the final version of the manuscript.

Competing interests: The authors declare no competing interest.

References

1. K. B. Zeldovich, P. Chen, E. I. Shakhnovich, Protein stability imposes limits on organism
complexity and speed of molecular evolution. *Proc Natl Acad Sci U S A* **104**, 16152-
16157 (2007).
2. G. Faure, E. V. Koonin, Universal distribution of mutational effects on protein stability,
uncoupling of protein robustness from sequence evolution and distinct evolutionary
modes of prokaryotic and eukaryotic proteins. *Physical Biology* **12** (2015).
3. S. Bershtein, M. Segal, R. Bekerman, N. Tokuriki, D. S. Tawfik, Robustness-epistasis link
shapes the fitness landscape of a randomly drifting protein. *Nature* **444**, 929-932 (2006).
4. N. E. Morton, J. F. Crow, H. J. Muller, AN ESTIMATE OF THE MUTATIONAL DAMAGE IN
MAN FROM DATA ON CONSANGUINEOUS MARRIAGES. *Proc Natl Acad Sci U S A* **42**, 855-
863 (1956).
5. G. I. Lang *et al.*, Pervasive genetic hitchhiking and clonal interference in forty evolving
yeast populations. *Nature* **500**, 571-574 (2013).
6. J. F. Storz, Compensatory mutations and epistasis for protein function. *Curr Opin Struct
Biol* **50**, 18-25 (2018).
7. J. D. Bloom *et al.*, Evolution favors protein mutational robustness in sufficiently large
populations. *BMC Biol* **5**, 29 (2007).
8. P. A. Lind, L. Arvidsson, O. G. Berg, D. I. Andersson, Variation in Mutational Robustness
between Different Proteins and the Predictability of Fitness Effects. *Mol Biol Evol* **34**,
408-418 (2017).
9. S. Pasek, J. L. Risler, P. Brézellec, The role of domain redundancy in genetic robustness
against null mutations. *J Mol Biol* **362**, 184-191 (2006).
10. Z. Gu *et al.*, Role of duplicate genes in genetic robustness against null mutations. *Nature*
421, 63-66 (2003).
11. S. Maisnier-Patin, D. I. Andersson, Adaptation to the deleterious effects of antimicrobial
drug resistance mutations by compensatory evolution. *Res Microbiol* **155**, 360-369
(2004).
12. J. D. Bloom, L. I. Gong, D. Baltimore, Permissive secondary mutations enable the
evolution of influenza oseltamivir resistance. *Science* **328**, 1272-1275 (2010).
13. K. Tomala, P. Zrebiec, D. L. Hartl, Limits to Compensatory Mutations: Insights from
Temperature-Sensitive Alleles. *Mol Biol Evol* **36**, 1874-1883 (2019).
14. A. Poon, L. Chao, The rate of compensatory mutation in the DNA bacteriophage
phiX174. *Genetics* **170**, 989-999 (2005).
15. N. Meftahi *et al.*, Evidence for the critical role of a secondary site rpoB mutation in the
compensatory evolution and successful transmission of an MDR tuberculosis outbreak
strain. *J Antimicrob Chemother* **71**, 324-332 (2016).

16.   I. Comas *et al.*, Whole-genome sequencing of rifampicin-resistant Mycobacterium
tuberculosis strains identifies compensatory mutations in RNA polymerase genes. *Nat*
Genet **44**, 106-110 (2011).
17.   A. Handel, R. R. Regoes, R. Antia, The role of compensatory mutations in the emergence
of drug resistance. *PLoS Comput Biol* **2**, e137 (2006).
18.   Q. Liu *et al.*, Have compensatory mutations facilitated the current epidemic of
multidrug-resistant tuberculosis? *Emerg Microbes Infect* **7**, 98 (2018).
19.   N. J. Ose, P. Campitelli, T. Modi, S. B. Ozkan, Second-order allosteric control as a
mechanism for compensatory mutations in B-cell translocation gene 2. *Protein Sci* **34**,
e70270 (2025).
20.   D. N. Ivankov, A. V. Finkelstein, F. A. Kondrashov, A structural perspective of
compensatory evolution. *Curr Opin Struct Biol* **26**, 104-112 (2014).
21.   D. K. Hanson, D. M. Tiede, S. L. Nance, C. H. Chang, M. Schiffer, Site-specific and
compensatory mutations imply unexpected pathways for proton delivery to the QB
binding site of the photosynthetic reaction center. *Proc Natl Acad Sci U S A* **90**, 8929-
8933 (1993).
22.   M. G. Mateu, A. R. Fersht, Mutually compensatory mutations during evolution of the
tetramerization domain of tumor suppressor p53 lead to impaired hetero-
oligomerization. *Proc Natl Acad Sci U S A* **96**, 3595-3599 (1999).
23.   W. Stephan, D. A. Kirby, RNA folding in Drosophila shows a distance effect for
compensatory fitness interactions. *Genetics* **135**, 97-103 (1993).
24.   D. A. Kirby, S. V. Muse, W. Stephan, Maintenance of pre-mRNA secondary structure by
epistatic selection. *Proc Natl Acad Sci U S A* **92**, 9047-9051 (1995).
25.   J. Domingo, G. Diss, B. Lehner, Pairwise and higher-order genetic interactions during the
evolution of a tRNA. *Nature* **558**, 117-121 (2018).
26.   M. Filteau *et al.*, Evolutionary rescue by compensatory mutations is constrained by
genomic and environmental backgrounds. *Mol Syst Biol* **11**, 832 (2015).
27.   A. Jakubowska, R. Korona, Lack of evolutionary conservation at positions important for
thermal stability in the yeast ODCase protein. *Mol Biol Evol* **26**, 1431-1434 (2009).
28.   A. F. Poon, L. Chao, Functional origins of fitness effect-sizes of compensatory mutations
in the DNA bacteriophage phiX174. *Evolution* **60**, 2032-2043 (2006).
29.   C. Ferrer-Costa, M. Orozco, X. de la Cruz, Characterization of compensated mutations in
terms of structural and physico-chemical properties. *J Mol Biol* **365**, 249-256 (2007).
30.   V. O. Pokusaeva *et al.*, An experimental assay of the interactions of amino acids from
orthologous sequences shaping a complex fitness landscape. *PLoS Genet* **15**, e1008079
(2019).
31.   N. Tokuriki, D. S. Tawfik, Stability effects of mutations and protein evolvability. *Curr Opin*
Struct Biol **19**, 596-604 (2009).
32.   A. Stein, D. M. Fowler, R. Hartmann-Petersen, K. Lindorff-Larsen, Biophysical and
Mechanistic Models for Disease-Causing Protein Variants. *Trends Biochem Sci* **44**, 575-
588 (2019).
33.   R. Casadio, M. Vassura, S. Tiwari, P. Fariselli, P. Luigi Martelli, Correlating disease-related
mutations to their effect on protein stability: a large-scale analysis of the human
proteome. *Hum Mutat* **32**, 1161-1170 (2011).
34.   A. S. Kondrashov, S. Sunyaev, F. A. Kondrashov, Dobzhansky–Muller incompatibilities in
protein evolution. *Proceedings of the National Academy of Sciences* **99**, 14878-14883
(2002).

35.   P. Bak, H. Flyvbjerg, B. Lautrup, Coevolution in a rugged fitness landscape. *Phys Rev A*
662   **46**, 6724-6730 (1992).

663   36.   J. Van Cleve, D. B. Weissman, Measuring ruggedness in fitness landscapes. *Proc Natl*
664   *Acad Sci U S A* **112**, 7345-7346 (2015).

665   37.   J. de Visser, S. F. Elena, I. Fragata, S. Matuszewski, The utility of fitness landscapes and
666   big data for predicting evolution. *Heredity (Edinb)* **121**, 401-405 (2018).

667   38.   P. Notin *et al.*, ProteinGym: Large-Scale Benchmarks for Protein Design and Fitness
668   Prediction. *bioRxiv* 10.1101/2023.12.07.570727 (2023).

669   39.   K. Tsuboyama *et al.*, Mega-scale experimental analysis of protein folding stability in
670   biology and design. *Nature* **620**, 434-444 (2023).

671   40.   L. Gonzalez Somermeyer *et al.*, Heterogeneity of the GFP fitness landscape and data-
672   driven protein design. *eLife* **11**, e75842 (2022).

673   41.   L. H. Greene *et al.*, Role of conserved residues in structure and stability: tryptophans of
674   human serum retinol-binding protein, a model for the lipocalin superfamily. *Protein Sci*
675   **10**, 2301-2316 (2001).

676   42.   J. Zheng, J. L. Payne, A. Wagner, Cryptic genetic variation accelerates evolution by
677   opening access to diverse adaptive peaks. *Science* **365**, 347-353 (2019).

678   43.   H. H. Guo, J. Choe, L. A. Loeb, Protein tolerance to random amino acid change. *Proc Natl*
679   *Acad Sci U S A* **101**, 9205-9210 (2004).

680   44.   G. Abrusán, J. A. Marsh, Alpha Helices Are More Robust to Mutations than Beta Strands.
681   *PLoS Comput Biol* **12**, e1005242 (2016).

682   45.   P. Popov *et al.*, Computational design of thermostabilizing point mutations for G
683   protein-coupled receptors. *Elife* **7** (2018).

684   46.   B. Borgo, J. J. Havranek, Automated selection of stabilizing mutations in designed and
685   natural proteins. *Proceedings of the National Academy of Sciences* **109**, 1494-1499
686   (2012).

687   47.   M. I. Zimmerman *et al.*, Prediction of New Stabilizing Mutations Based on Mechanistic
688   Insights from Markov State Models. *ACS Cent Sci* **3**, 1311-1321 (2017).

689   48.   B. Steipe, B. Schiller, A. Plückthun, S. Steinbacher, Sequence statistics reliably predict
690   stabilizing mutations in a protein domain. *J Mol Biol* **240**, 188-192 (1994).

691   49.   C. P. Lin *et al.*, Deriving protein dynamical properties from weighted protein contact
692   number. *Proteins* **72**, 929-935 (2008).

693   50.   P. Carter, C. A. Andersen, B. Rost, DSSPcont: Continuous secondary structure
694   assignments for proteins. *Nucleic Acids Res* **31**, 3293-3295 (2003).

695   51.   M. Z. Tien, A. G. Meyer, D. K. Sydykova, S. J. Spielman, C. O. Wilke, Maximum allowed
696   solvent accessibilites of residues in proteins. *PLoS One* **8**, e80635 (2013).

697   52.   C. Notredame, D. G. Higgins, J. Heringa, T-Coffee: A novel method for fast and accurate
698   multiple sequence alignment. *J Mol Biol* **302**, 205-217 (2000).

699   53.   S. Lyskov, J. J. Gray, The RosettaDock server for local protein-protein docking. *Nucleic*
700   *Acids Res* **36**, W233-238 (2008).

701   54.   F. Faul, E. Erdfelder, A. Buchner, A. G. Lang, Statistical power analyses using G*Power
702   3.1: tests for correlation and regression analyses. *Behav Res Methods* **41**, 1149-1160
703   (2009).

704   55.   A. A. Hagberg, L. A. National, L. Alamos, D. A. Schult, P. J. Swart, Exploring Network
705   Structure , Dynamics , and Function using NetworkX. (2008).

706   56.   T. Magoč, S. L. Salzberg, FLASH: fast length adjustment of short reads to improve
707   genome assemblies. *Bioinformatics* **27**, 2957-2963 (2011).

708   57.   E. Zorita, P. Cuscó, G. J. Filion, Starcode: sequence clustering based on all-pairs search.
709   *Bioinformatics* **31**, 1913-1919 (2015).
710   58.   F. Li, M. L. Salit, S. F. Levy, Unbiased Fitness Estimation of Pooled Barcode or Amplicon
711   Sequencing Studies. *Cell Syst* **7**, 521-525.e524 (2018).

Supplementary Figures and Tables

Figure S1

716
Fig. S1. The DMS library is derived from extant yeast sequences and spans nearly the full length
of His3p. (A) Position wise conservation and amino acid conservation across His3p. The red track
shows conservation score across 21 extant yeast species at each position of the His3p monomer. The
blue track shows the conservation score in the DMS library with heatmap below showing the
frequency of each amino acid in each position in the library. (B) Most genotypes in the library are
composed of extant amino acids. (C) Fitness distribution of all genotypes in the library, the blue bar
indicates percentage of unfit genotypes. (D) We defined "rescue" as a fitness increase from 0 to a
value significantly above 0, and calculated rescuability accordingly. The correlation between
rescuability values was then examined across a range of fitness thresholds used to define unfit and fit
genotypes. (E) Different rescuability calculation methods produce strong agreement of rescuability
estimates for unfit genotypes ($R^2 = 0.84$). (F) Unfit genotypes span nearly the entire length of His3p.
728

Figure S2

731
Fig. S2. Rescuable genotypes are more structurally stable than non-rescuable genotypes. (A)
Across varying numbers of substitutions, rescuable genotypes display higher predicted structural
stability than non-rescuable genotypes. (B) Among genotypes with more than 50% substitutions
in subunit interface regions, rescuable genotypes remain more structurally stable than non-
rescuable genotypes across varying numbers of substitutions.

Figure S3

739
Fig. S3. There exist several substitutions that increase fitness of diverse genotypes. (A)
Compensatory relationships of all pairs of substitutions and non-WT amino acid states at distinct
positions for all segments, excluding segment 9. (B) Compensatory capacity of each substitution
across all segments, excluding segment 9.

Figure S4

744
745 **Fig. S4. Fitness potential is a latent variable that can predict fitness.** Each panel illustrates
746 the non-linear mapping learned by a neural network model for each of the 12 His3p segments
747 analyzed, excluding segment 9.

Fig. S5. Distance from substituted position of each non-WT amino acid state to the substrate. Distance from each non-WT amino acid state to the substrate (Å, y-axis) was plotted against compensatory ability (x-axis) for all segments except segment 9. There is no significant correlation between the distance from each non-WT amino acid state to the substrate and the its compensatory ability.

756
757

Figure S6

Fig. S6. Super compensators identified across diverse proteins. (A) Distribution of compensatory ability for individual substitution across different proteins. Substitutions with prominent compensatory ability ($\geq$ 0.6) are labeled. The fitness metric used in each dataset (stability or fluorescence intensity) is indicated in the title of each panel. (B) Super compensators buffer the fitness effects of other substitutions. Each panel corresponds to the super compensator highlighted in panel A (which showed the highest compensatory ability for that protein). The percentage of substitutions exhibiting reduced fitness effects in the super compensator background is labeled.

Figure S7

770
Fig. S7. Amino acid physicochemical properties can predict fitness. The heatmap showing
Spearman correlations between fitness and physicochemical properties (AAIndex values)
distance from WT. Columns correspond to AAIndex descriptors and rows to distinct mutation
sets. The high spearman correlation indicts that move the AAindex values of variants closer to
WT can increase the fitness. The high negative spearman correlations indicate that moving the
AAIndex values of variants closer to those of WT increases fitness.
777

Figure S8

Fig. S8. Co occurring amino acid state pairs in extant His3p orthologs, across all segments excluding segments 9 and 10. (A) Co-occurrence network of amino acid state pairs across 355 extant yeast species. Nodes represent amino acid states and edges the co-occurrence of a pair. Edge shading reflects the fraction of species in which a given pair co-occurs. Non-extant amino acid states are denoted by lighter node shading. Pairs that co-occur in at least 10 species are labelled with white text. (B) Among pairs of non WT amino acid states observed in extant species, amino-acids states with lower fitness impact scores relative to WT (x-axis, binned) tend to co-occur with states exhibiting higher fitness-impact scores (y-axis).

794 **Table S1. Structural effects of super compensatory and non-compensatory substitutions.**
795 Predicted structural effects of 7 super compensators (highest fitness impact score relative to WT) and
796 five non compensatory substitutions (lowest fitness impact score relative to WT)**.**

Substitution	Effects	Super compensatory substitutions
S189A	Remarkably stabilize the helical structure	Yes
E169R	Hydrogen bonds may form between E165 and an adjacent side chain to stabilize the helix.	Yes
N137G	The conformation interval of R138 is enlarged, and the salt bridge between R138 and phosphate group in substrate is weakened	Yes
A110D	Form a hydrogen bond with R112 to stabilize loop conformation	Yes
E144N	To weaken its rejection with D178 and stabilize the β-sheet structure, and may also form hydrogen bonds with D178	Yes
S73A	Remarkably stabilize the helical structure	Yes
T93V	Beneficial for hydrophobic accumulation in helical structure	Yes
I19D	Destroy the beta sheet and helical stack	No
I19N	Destroy the beta sheet and helical stack	No
I200P	Destroy helix and hydrophobic accumulation	No
H67P	Destroy the helix, the hydrogen bond	No
F163I	Destroy the helix and the aromatic ring accumulation	No