Autoimmune regulator deficiency causes sterile epididymitis and impacts male fertility

through disruption of inorganic physiology

Soo Hyun Ahn,[1] Katrina Halgren,[2] Geoffrey Grzesiak,[1] Keith W. MacRenaris,[3] Aaron Sue,[3]

Huirong Xie,[4] Elena Demireva,[4] Thomas V. O'Halloran,[3,5,6*] and Margaret G. Petroff[1,6,7*]

*Corresponding Authors

Corresponding author email: petrof10@msu.edu

Affiliations

[1] Department of Pathobiology Diagnostic Investigation, College of Veterinary Medicine, Michigan

State University, East Lansing, MI, USA.

[2]Lyman Briggs College, Michigan State University, East Lansing, MI, USA

[3]Elemental Health Institute and Department of Microbiology Genetics and Immunology, Michigan

State University, East Lansing, MI, USA.

[4]Transgenic and Genome Editing Facility, Institute for Quantitative Health Science & Engineering,

Research Technology Support Facility, Michigan State University, East Lansing, MI, USA

[5]Department of Chemistry, Michigan State University, East Lansing, MI, USA

[6]Institute for Integrative Toxicology, Michigan State University, East Lansing, MI, USA.

[7]Department of Microbiology, Genomics, and Immunology, Michigan State University, East

Lansing, MI, USA

Funding: This research is supported by NIH grants R01 HD100832, R01 GM11484 and the

National Research Resource for Quantitative Mapping in the life Sciences, supported by the Office

of the Director, National Institutes of Health, and the National Institute for General Medical

Sciences under grant number P41 GM135018.

Running Title: Aire-deficiency disrupts inorganic physiology and leads to sterile

epididymitis

Abstract

Autoimmune regulator (AIRE), a transcription factor expressed by medullary thymic epithelial

cells, is required for shaping the self-antigen tolerant T cell receptor repertoire. Humans with

mutations in *AIRE* suffer from Autoimmune Polyglandular Syndrome Type 1 (APS-1). Among

many symptoms, men with APS-1 commonly experience testicular insufficiency and infertility,

but the mechanisms causing infertility are unknown. Using an *Aire*-deficient mouse model, we

demonstrate that male subfertility is caused by sterile epididymitis characterized by immune cell

infiltration and extensive fibrosis. In addition, we reveal that the presence of autoreactive immune

cells and inflammation in epididymides of *Aire*-deficient mice are required for iron (Fe) deposition

in the interstitium, which is brought on by macrophages. We further demonstrate that male

subfertility is associated with a decrease in metals zinc (Zn), copper (Cu), and selenium (Se) which

serve as cofactors in several antioxidant enzymes. We also show increase in DNA damage of

epididymal sperm of $Aire^{-/-}$ animals as a key contributing factor to subfertility. The absence of *Aire*

results in autoimmune attack of the epididymis leading to fibrosis, Fe deposition, and Cu, Zn and

Se imbalance, ultimately resulting in sperm DNA damage and subfertility. These results highlight

the requirement of *Aire* to promote immune tolerance throughout the epididymis, disruption of

which causes an imbalance of inorganic elements with resulting consequence on male fertility.

Key points:

Breakdown of epididymal self-tolerance promotes disruption of inorganic elements.

Autoimmunity causes interstitial fibrosis resulting in sperm DNA damage and subfertility.

Elevated interstitial iron and macrophages contribute to fibrosis.

Introduction

Infertility, a condition in which a couple is unable to achieve pregnancy after 12 months of unprotected intercourse, affects approximately 1 in 6 individuals worldwide, with lifetime prevalence ranging from 10.8 to 23.4% depending on global region.[1] Male factors are thought to account about 20% of cases, and contribute to as much as 60%.[2] Male infertility can stem from defective testicular function resulting in abnormal sperm concentration, motility, and morphology.[3] However, many cases of infertility are unexplained, occurring despite normal sperm parameters.[4] In these cases, underlying autoimmune disease may contribute by promoting inflammation of the reproductive organs, especially the epididymis,[5] which could perturb sperm maturation.

Autoimmunity, or loss of immune self-tolerance, results from inappropriate T cell reactivity towards self-antigens. Self-tolerance in T cells is established in the thymus through the expression of a transcription factor called Autoimmune Regulator (AIRE) by medullary thymic epithelial cells (mTECs). In these cells, AIRE regulates transcription of tissue-restricted antigens, which are then presented in the context of major histocompatibility complex; nascent autoreactive T cells that interact too strongly with the presented self-antigen are subsequently eliminated or directed into a regulatory phenotype.[6] In humans, mutation in the *AIRE* gene causes Autoimmune Polyglandular Syndrome Type 1 (APS-1), a rare autoimmune disease that manifests in a variety of symptoms including mucocutaneous candidiasis, hypoparathyroidism, adrenal insufficiency, gonadal insufficiency, and infertility.[7–9]

Using an Aire deficient (*Aire*[-/-]) mouse model, we previously documented CD3+ T cell infiltration

into the prostate gland, seminal vesicles, epididymis, and more rarely, the testis.[10] In addition,

while sperm production and morphology appeared normal, sperm from *Aire*[-/-] males were severely

impaired in their ability to fertilize eggs. *In vitro* fertilization using *Aire*[-/-] epididymal sperm

resulted in only 9% of embryos reaching the 2-cell stage, and none reaching the blastocyst stage.[10]

These results suggest that malfunction of sperm in *Aire*[-/-] males are caused during post-

spermatogenic maturation in the epididymis, possibly due to abnormalities in the epididymal

environment.

The epididymis is an elongated, tightly coiled tube that connects each testicle to the vas deferens.

In addition to storing and transporting sperm in its distal-most portion (cauda) prior to ejaculation,

the epididymis provides important factors that transform the immature testicular sperm into

mature, fertilization-competent sperm.[11,12] The epididymis is indispensable for final post-testicular

maturation of sperm; if they do not traverse the entire length of the epididymis, they remain

incapable of fertilization.[13,14] The rodent epididymis is divided into four anatomical regions: the

initial segment (most proximal to the testis), the caput (head), the corpus (body), and the cauda

(tail) (most distal to the testis, merging into the vas deferens). With their distinct transcriptomic

profiles[15] and luminal contents,[16] each segment has specific roles in facilitating sperm maturation.

The epididymis preserves sperm in a dormant state by tightly regulating the pH of the lumen.[17,18]

Epididymal epithelial cells also produce antioxidant enzymes, including glutathione peroxidases,

perioxiredoxins, and superoxide dismutases, for which essential metals serve as requisite

cofactors.[12,19–21 22,23] Thus, any disruption in this epididymal environment may cause defects in

sperm maturation and their fertilization competence, ultimately resulting in infertility.

Here, we demonstrate that deletion of *Aire* promotes targeting of the epididymis by autoreactive immune cells, resulting in epididymitis, interstitial fibrosis, dysregulation of inorganic physiology, and severely compromising fertility. Using a range of emerging methods to determine quantitative inorganic phenotypes,[24] we demonstrate that autoimmune-mediated epididymal damage results in interstitial fibrosis and is associated with iron deposition, depletion of essential antioxidant cofactors such as zinc (Zn), copper (Cu), and selenium (Se), and DNA damage of the sperm. By crossing *Aire*[-/-] mice to mice deficient in T and B cells through deficiency in Recombination Activating Gene 1 (Rag1), which plays a fundamental role in generating T and B cells, we also demonstrate the role of the adaptive immune system in compromising fertility in *Aire*[-/-] males and driving fibrinogenesis and inorganic physiology.

Materials and Methods

Animals

Experiments with animals complied with NIH's *Guide for the Care and Use of Laboratory Animals*,[25] and were approved by the Institutional Animal Care and Use Committee at Michigan State University. Mice were maintained on a 12h:12h light cycle with rodent chow (Teklad 2019; Inotiv), which is formulated to support gestation in transgenic and inbred animals, and water available *ad libitum*. Mice containing mutations in the autoimmune regulator (*Aire*) gene were generated on the Balb/cJ background (Jackson Laboratories) by the Michigan State University Transgenic and Genome Editing Facility using CRISPR/Cas9 targeting. A single guide RNA (gRNA) targeting exon 2 of the *Aire* gene (**ENSMUSG00000000731**), with protospacer and PAM sequence 5'- GAC GCT CCG TCT GAA GGA GA – AGG 3' (g158), was synthesized by IDT.

Ribonucleoprotein (RNP) complexes of Cas9 and gRNA were assembled and electroporated into

Balb/c zygotes to generate mutant mice as previously described.[26] Resulting edited founders

carrying 32 base-pair (bp), 44bp, or 86bp deletions were identified and backcrossed with Balb/c

mice to establish the transgenic lines. Preliminary findings showed profound male subfertility in

all three lines; data used in this manuscript were generated from the 32 bp line.

Aire$^{+/-}$ mice were crossed with *Rag1*-deficient animals to generate mice that were heterozygous for

both *Aire* and *Rag1* genes. Mice were genotyped by PCR analysis of tail biopsy genomic DNA

using the following primers: Aire F1: 5'-CAC ACC CTA TGC CCC CTA AC-3', Aire R1- 5`-

ACA TGG GCC GAC TTG TAT CC-3`; Rag1MutFwd: 5`-TGG ATG TGG AAT GTG TGC

GAG-3`; Rag1Common: 5`-TAC ACC TTG GCG ACG ACT CCT-3`, Rag1WTFwd: 5`-TCT

GGA CTT GCC TCC TCT GT-3`. *Aire*-deleted animals were confirmed for the absence of Aire

protein in the thymus by immunofluorescent staining, using wild type thymus as a control.

Fertility trial

6-week-old male *Aire*$^{+/+}$ (n=4) and *Aire*$^{-/-}$ (n=8) animals were paired with 6-week-old Balb/c virgin

females to assess fertility. These animals were either WT or heterozygous for the *Rag1* gene. To

assess the role of the adaptive immune system in fertility outcome of *Aire*$^{-/-}$ mice, additional

pairings of *Aire*$^{-/-}$*Rag1*$^{-/-}$ males (n=4) and *Aire*$^{+/+}$*;Rag1*$^{-/}$ males (n=3) were established. Mice

remained paired for 14.5 to 16 weeks, with the exception that two of the three *Aire*$^{+/+}$*;Rag1*$^{-/-}$ males

were paired for ~6 weeks. At the conclusion of the fertility trial, males were euthanized for

downstream analyses at 19 to 23 weeks of age; additional males were euthanized at 7-9 weeks.

Females were euthanized at the same time as the males, except when they were pregnant, in which

case they were euthanized following birth of the litter. Fertility rate was calculated as:

$$(number\ of\ litters \div number\ of\ days\ paired) \times 21$$

We chose 21 days as an expected length of mouse gestation; if dams produced a litter of pups

every 21 days, the resultant number would be close to 1.

Histology

Paraffin-embedded tissues were fixed in 4% paraformaldehyde at 4°C overnight, transferred to

70% ethanol, embedded in paraffin, and sectioned at 5 μm onto SuperFrost slides (Thermofisher

Scientific). Alternatively, tissues were fixed in 4% paraformaldehyde at 4°C overnight, transferred

to 30% sucrose, and embedded in OCT medium (Sakura) by freezing the medium in liquid-

nitrogen chilled isopentane. OCT-embedded samples were kept at -80°C until cryosectioning

(Leica Biosystems). For hematoxylin and eosin staining,[27] we used the Leica SelecTech staining

system consisting of Hematoxylin 560, Blue Buffer 8, Define, Alcoholic Eosin Y 515. Sections

were deparaffinized in Histoclear (National Diagnostics) and rehydrated in a gradient of ethanol

(100%, 95% and 80%) and water. Slides were stained in hematoxylin for 3 minutes, washed in

water for 1 minute, and immersed in 95% ethanol for 40 seconds followed by alcoholic eosin Y

for 12 seconds. Slides were then washed using two exchanges of 90% ethanol for 40 seconds and

dehydrated in 100% ethanol for 5 minutes. Finally, slides were washed in Histoclear for 5 minutes

and mounted using Cytoseal 60 (Fisher Scientific). For Masson's trichrome staining, which was

used to assess fibrosis, serially sectioned samples were submitted to the Investigative

Histopathology Core at Michigan State University. Using this stain, connective tissue stains blue,

nuclei stain purple/red, and cytoplasm stains pink/red.

Immunohistochemistry and Immunofluorescence

Paraffin-embedded epididymal tissue sections were immunohistochemically stained with rabbit

anti-CD8 (ab217344, Abcam), rabbit anti-CD4 (Abcam), rat anti-CD19 (6OMP31, Thermofisher

Scientific), rat anti-F4/80 (123102, Biolegend), or rabbit anti-αSMA (14395-1-AP, Proteintech)

antibodies. After deparaffinization and rehydration, we performed a heat-mediated antigen

retrieval step using Reveal Decloaker sodium citrate buffer (Biocare Medical), heated to 199°F for

20 minutes. After cooling to room temperature, samples were incubated with the SuperBlock™

blocking buffer (37515, ThermoFisher Scientific) for 10 minutes followed by 3% hydrogen

peroxide in methanol to quench endogenous peroxidase activity for 10 minutes. Sections were

washed with phosphate buffered saline (PBS) before incubating with the primary antibody.

Adjacent sections were used for isotype controls. Biotinylated anti-rabbit or anti-rat IgG secondary

antibody (Vector Laboratories) was applied, followed by Ready-To-Use Horseradish Peroxidase

and DAB (Vector Laboratories) for color development.

To quantify positively stained cells, slides were scanned using a Zeiss Axioscan 7 Microscope

Slide Scanner (Carl Zeiss AG). Three nonoverlapping, randomly chosen regions of interest (ROI;

350 µm × 950 µm) were from the caput, corpus, and cauda epididymides were captured using Zen

Microscopy Software (Carl Zeiss). In some animals, only two non-overlapping regions could be

imaged. Three independent blinded evaluators counted DAB+ cells from slides stained with anti-

CD4 and anti-CD8 antibodies. The non-parametric Wilcoxon's t-test was used to compare the

means of DAB+ cells between $Aire^{+/+}$ and $Aire^{-/-}$ samples. Intraclass Correlation Coefficient was

used to assess the strength of inter-evaluator agreement and was calculated using the "icc" R

package. The ICC for this experiment was 0.91, indicating a high similarity between values

measured by the three independent evaluators.

For immunofluorescence, OCT-embedded samples were sectioned at 5 μm, washed in PBS

for 10 minutes, blocked using SuperBlock Blocking Buffer, and incubated overnight with

antibodies against F4/80, CD163 (1:400; Abcam), and/or Ferritin (1:1000; F5012, Sigma Aldrich).

The following day, the slides were washed and incubated with secondary antibodies (AF488-anti-

rat or AF546-anti-rabbit IgG; 0.01mg/mL, ThermoFisher Scientific) and mounted using ProLong

Gold antifade reagent with DAPI (ThermoFisher Scientific).

Flow cytometry

For flow cytometeric analysis of lymphocytes, epididymides from 2 males were pooled into 1

sample, such that 1 biological replicate consisted of cells from 2 males and 4 epididymides. Once

excised, cauda epididymides were dissected from the rest of the epididymis and collected into

TYH media warmed to 37°C to allow sperm to swim out of the sample. Cauda epididymides were

further processed and analyzed separately from caput-corpus epididymides such that spatial

information could be maintained while ensuring sufficient cellular yield. Tissues were finely

minced using spring scissors, collected into gentleMACS™ M Tubes (Miltenyi Biotec), and

incubated in 5 mL of Miltenyi Tissue Dissociation Kit 1 (Miltenyi Biotec) for 15 minutes at 37°C,

using GentleMACS™ Dissociator (Miltenyi Biotec) for 1 minute per sample at room temperature.

The samples were returned to 37°C to incubate for a 15 additional minutes. The digest was filtered

through a 70 μm filter into 50 mL conical tubes containing 10 mL of PBS with 5% FBS to

neutralize the digestive enzyme. Each sample was topped up to 30 mL with 5% FBS and

centrifuged at 400g for 5 minutes. The cell pellet was washed with 10 mL of PBS/5%FBS buffer,

resuspended in 0.4 mL of PBS/5%FBS buffer, and aliquoted into 96 well plates for staining. All

cells were incubated with a viability stain (1:1000; Invitrogen™ Live/Dead™ Fixable Blue Dead

Cell Stain kit, ThermoFisher Scientific) for 30 minutes on ice, followed by staining with Mouse

BD FcBlock™ (1:100, 553142BD Pharmingen, Cat. No.) for 20 minutes on ice. Cells were

pelleted and resuspended with the primary antibody cocktail on ice for 30 minutes. The following

antibodies were used: BV421 anti-CD45 (1:200; 103133, 30-F11, Biolegend,), KIRAVIA Blue

520 anti-CD4 (1:100, clone GK1.5; Biolegend), BV785 anti-CD8 (1:100, 53-6.7, Biolegend), PE-

Dazzle 594 anti-CD44 (1:100, IM7, Biolegend), PE-Cy5 anti-CD69 (1:50; H1.2F3, Biolegend),

APC anti-CD25 (1:50; PC61, Biolegend), BV510 anti-B220 (1:100; RA3-6B2, Biolegend). After

incubation with primary antibodies, the cells were washed three times in 200 µL of flow staining

buffer, pelleting the cells between each wash. Lastly, cells were fixed using 1% paraformaldehyde

in PBS for 15 minutes on ice and stored at 4°C until analysis using Cytek® Aurora spectral flow

cytometer (Cytek® Biosciences,).

Gelatin standard preparation

Gelatin standards (blank, 1, 5, 10, 20, 50 parts per million (ppm)) (**Supplemental Table 1**) were

produced by mixing in IV-Stock-4 (Inorganic Ventures) to 10% porcine gelatin (G2500-500G,

Sigma Aldrich) and heated to 55°C. The blank contained no IV-Stock-4. 400 µL of the preparation

was pipetted directly onto the cryostat mounting chuck, left to freeze at -20°C for 4 minutes, and

sectioned at 20 µm thickness onto the SuperFrost glass slides (Thermofisher Scientific). A 50 µL

aliquot of each standard was aliquoted in a pre-weighed metal-free tube to determine elemental

concentrations using an Agilent 8900 Triple Quadrupole - Inductively Coupled Plasma - Mass

Spectrometer (ICP-QQQ-MS, Agilent) equipped with the Agilent SPS 4 Autosampler, integrate

sample introduction system (ISiS), x-lens, and micromist nebulizer. To each sample, 300 µL of

trace metal grade 70% nitric acid (Fisher chemical, #A509P212), was added to digest the sample

at 70 °C for 4 hours. Following digestion, 9.7 mL of ultrapure deionized water was added to each

sample. All samples were weighed prior to analysis on the ICP-QQQ-MS.

Spatial, quantitative mapping element distribution using LA-ICP-TOF-MS

Paraffin-embedded 5 µm tissue sections were mounted onto charged glass slides and laser ablated

using the Elemental Scientific Lasers BioImage 266 nm laser ablation system (Elemental Scientific

Lasers), which is equipped with an ultra-fast low dispersion TwoVol3 ablation chamber and a dual

concentric injector (DCI3). The aerosolized sample was transferred to the Tofwerk icpTOF S2

mass spectrometer (TOFWERK AG) where the elemental content was analyzed real time

according to mass/charge (m/z) ratio. Daily tuning was performed using NIST SRM612 glass

certified reference material (National Institute for Standards and Technology). High intensities for

^{140}Ce and ^{55}Mn were used to optimize torch alignment, lens voltages, and nebulizer gas flow while

maintaining low oxide formation based on the ^{232}Th^{16}O+/^{232}Th+ ratio (< 0.5). Parameters are listed

in **Supplemental Table 2**. Laser scanning was performed in a raster pattern with a laser repetition

rate of 100 Hz using a 10 µm circular spot size with laser fluence set to 15 J/cm^2. Ten lines of

gelatin standards were ablated using the same parameters but with the raster spacing set at 20 µm.

Scanning data was recorded using TofPilot 1.3.4.0 (TOFWERK AG) and saved in the open-source

hierarchical data format (HDF5). Post-acquisition data processing was performed with Tofware

v3.2.0, a data analysis package used as an add-on to IgorPro (Wavemetric Inc.). Imaging and mass

calibration was conducted using the Iolite version 4.8.6. Software (Elemental Scientific Lasers) .[28]

Elemental Composition Determination by ICP-QQQ-MS

Mouse epididymides separated into caput, corpus, and cauda were placed into a pre-weighed metal-free 15mL conical tubes (Labcon) to be analyzed using ICP-QQQ-MS (Agilent). Prior to analysis, each metal-free tube was weighed using an analytical balance pre- and post-addition of the tissue sample to ascertain the weight of the tissue. Tissues were digested in 300 µL of 70% nitric acid (Fisher, #A509P212) and incubated at 65°C overnight. The following day, 9.7 mL of ultrapure deionized water was added to bring the final nitric acid concentration to 3% by volume. Each tube was weighed again before analysis to ascertain the final weight of the sample. Standards were prepared using IV-Stock-4 (Inorganic Ventures) at 0.01, 0.1, 1, 10, and 100 parts per billion (ppb) using 3% nitric acid in metal-free 15 mL conical tubes. The element concentrations of the samples were calculated using the following equation:

$$ug/g\ (ppm) = (weight\ of\ the\ ICP\ sample\ (g)\ * \ (0.001$$

$$* \ ppb))\ /\ weight\ of\ the\ sample\ (g)$$

Comet assay

The assay was conducted using the CometAssay® Single Cell Gel Electrophoresis Assay kit (Catalog # 4250-050-K, R&D Systems). Cauda epididymides were collected from *Aire*$^{+/+}$ and *Aire*$^{-/-}$ males into a 1.5 mL tube filled with 250 µL of PBS, gently minced with spring scissors, and incubated for 15 minutes at 37°C to allow the sperm to swim out of the tissue. The supernatant was collected into a separate 1.5mL centrifuge tube and counted using a hemocytometer by mixing 5 µL of the suspension with 95 µL of MilliQ water. Fifty µL of the resuspended sperm (1 million cells/mL) were mixed with 500 µL of 1% LMAgarose, spread onto a glass slide, and placed at 4°C

for 30 minutes. The samples were then incubated with lysis buffer + Proteinase K (500 µg/mL) for

18 hours at 4°C. The following day, the slides were immersed in cold Neutral Electrophoresis

Buffer (0.5M Tris-HCL, 1.5M sodium acetate) and subjected to electrophoresis at 21V/cm for 45

minutes at 4°C. Next, the slides were immersed in DNA Precipitation Solution and immersed in

70% ethanol for 30 minutes each at room temperature. The slides were dried at 37°C and stained

with 100 uL of SYBR® Gold Nucleic Acid Gel stain (1:10000, S11494, ThermoFisher Scientific)

for 30 minutes at room temperature. Lastly, the slides were briefly washed in tap water, dried, and

imaged using a 488 nm filter (Zeiss Axioscan 7). To calculate the percentage of sperm with comet

tails, Fiji software was used to convert the images to 8-bit, and threshold was set between 65-70

range to 255, followed by the "Analyze Particle" function in Fiji, which counted the number of

sperm heads (particle size set to 10 - 250) present on the image. To calculate proportion of sperm

with fragmented DNA, the comet tails were manually counted from the same image, divided by

the total number of particles detected, and multiplied by 100.

Statistical analysis

R studio and the packages ggplot2, ggbeeswarm, readxl, ggpubr, ggsci, icc, and tidyverse were

used to generate graphs and conduct statistical analysis. Normal distribution of the data was tested

using the Shapiro-Wilks test. For comparison of the means between two independent, normally

distributed samples, Student's t-test was used; for more than two, one-way ANOVA followed by

Tukey's post-hoc test was used. For samples that failed Shapiro-Wilks normality test, Wilcoxon's

test was used to compare the means of two samples, and Kruskal-Wallis test for two or more

samples, followed by Dunn's post-hoc test. For the comet assay, a paired t-test was conducted. *P*-

values less than 0.05 were deemed statistically significant.

Results

***Aire*[-/-] males subfertility is dependent on the adaptive immune system**

We generated transgenic animals heterozygous for *Aire* and *Rag1* genes on the Balb/c background. The absence of *Rag1* gene with *Aire*-deficiency (*Aire*[-/-]) produces animals without adaptive T and B cells (*Aire*[-/-];*Rag1*[-/-]), allowing us to address the role of central tolerance to male reproductive tissues. Since we rederived the *Aire*[-/-] animals using CRISPR/Cas9, we validated the absence of Aire protein in the thymus using immunofluorescence **(Fig 1A)**. We also tested whether the *Aire*[-/-] animals recapitulated the development of autoimmunity and infertility, as previously demonstrated by us and others.[6,10,29] We observed immune cell infiltration in both reproductive and non-reproductive organs (**Fig. 1B**), confirming that our *Aire*[-/-] mouse model recapitulated established mouse models of APS-1.[6] Because we observed smaller testes with immune cell infiltration in about 14% of *Aire*[-/-] males examined,[10] we euthanized 10- and 20-week-old *Aire*[-/-] males to assess testicular histology. Hematoxylin and eosin staining showed normal seminiferous tubules with intact spermatogenesis, comparable to *Aire*[+/+] testes in a majority of our *Aire*[-/-] samples at 10 weeks of age (*Aire*[+/+]; n=4, *Aire*[-/-]; n=4) and 20 weeks of age (*Aire*[+/+]; n=9, *Aire*[-/-]; n=10, data not shown), suggesting that testes were not targeted by immune cells, or that the testes-blood-barrier prevented immune cell infiltration in these *Aire*[-/-] mice.

To test fertility, 6-week-old *Aire*[+/+] (n=3) or *Aire*[-/-] (n=7), which were either heterozygous or wild-type for the *Rag1* gene, were singly housed with virgin WT Balb/c females. Additionally, fertility of *Aire*[+/+];*Rag1*[-/-] (n=3) and *Aire*[-/-];*Rag1*[-/-] (n=4) males was evaluated. Wild type (*Aire*[+/+]*Rag1*[+]) males demonstrated robust fertility, with an average delivery rate of 0.85 litters per 21 days and 74 pups total over ~14 weeks. In contrast, the *Aire*[-/-];*Rag1*[+] males had an average delivery rate of 0.08 litters per 21 days, with only 17 total pups produced. The two litters produced by the seven

Aire$^{-/-}$ males occurred in the first 7 weeks of pairing, whereas *Aire*$^{+/+}$ males were able to sire litters

for the entire duration of pairing. *Aire*$^{-/-}$;*Rag1*$^{-/-}$ males showed fertility comparable to *Aire*$^{+/+}$ mice

(Fig. 1C), showing that adaptive immune cells are responsible for subfertility in *Aire*$^{-/-}$

***Aire*$^{-/-}$ epididymides show immune cell infiltration**

Because the testis of *Aire*$^{-/-}$ mice appeared normal, we focused our attention on the epididymis,

where the testicular sperm undergo maturation, gain fertilization competency, and are stored

before ejaculation. Histological analysis of 10 and 20-week-old *Aire*$^{-/-}$ epididymides revealed

thickening of the interstitium and immune cell infiltration consisting of CD4+ and CD8+ T cells,

and CD19+ B cells **(Fig. 1B, 2A).** To measure the degree of infiltration of T cells, we counted the

immunohistochemically stained cells, with attention to age-dependent variation and differences

between the epididymal caput (head), corpus (body), and cauda (tail) **(Fig. 2B)**. While CD4 and

CD8 cells in *Aire*$^{+/+}$ epididymides were found only sporadically, both cell types were frequently

found in *Aire*$^{-/-}$ mice in all three segments of *Aire*$^{-/-}$ animals in both young and old animals. No

differences between regions of epididymis within each strain were observed ($P>0.05$). Age-

dependent differences were not obvious, therefore data from young and aged animals were

combined. T cell infiltration of all three epididymal regions of *Aire*$^{-/-}$ mice were significantly

higher than those of *Aire*$^{+/+}$ mice (**Fig. 2C, D**). As expected, neither T nor B cells could be

identified in *Rag1*-deficient mice, regardless of absence or presence of *Aire* (not shown).

To characterize the activation status of the infiltrating epididymal T cells, we performed flow

cytometry, separating the caput and corpus from the cauda epididymides. Within genotype, no

differences were observed in the relative abundance of CD45hi immune cells in the different

regions of the epididymis ($P>0.05$). Between genotypes, the proportion of CD45hi cells among

total cells trended towards increasing in the corpus-caput, and significantly increased in the cauda

epididymis, of $Aire^{-/-}$ mice as compared to $Aire^{+/+}$ controls (**Fig. 3B**). Similarly, the proportion of

B cells remained similar within genotype, but increased in the cauda epididymis of $Aire^{-/-}$ mice

(**Fig. 3C**). Relative abundance of CD4+ and CD8+ T cells was higher in the epididymal caput-

corpus of $Aire^{-/-}$ mice than that of $Aire^{+/+}$ controls (**Fig. 3D, E**). In the cauda epididymis, CD4+ T

cells remained similar between knockout and control mice (**Fig. 3F**), but CD8+T cells increased

approximately 6-fold in $Aire^{-/-}$ mice **(Fig. 3G)**. Finally, the early activation marker CD69 was

elevated on CD8+ T cells of both caput-corpus (**Fig. 3E**) and cauda (**Fig. 3G**); in CD4+ T cells,

the CD69 expression appeared to increase but this did not reach statistical significance, possibly

due to insufficient statistical power (**Fig. 3D, F**). Nearly all CD4+ and CD8+ T cells were CD44^{high},

indicating prior antigen experience; levels of CD44, however, were unchanged between genotypes

(**Fig. 3D-G**). CD25 expression was also unchanged bewteen $Aire^{-/-}$ and $Aire^{+/+}$ mice, except for

CD4+ cells obtained from cauda epididymis (**Fig. 3F**). Few CD8+ T cells in either genotype

expressed IL-2Rα/activation marker CD25 (**Fig3. E, G**).

^{56}Fe is increased in $Aire^{-/-}$ epididymis and correlates with fibrosis

Thickening of the epididymal interstitium in $Aire^{-/-}$ mice (**Fig. 1B**) could indicate fibrotic change

of the tissue. To test this, tissues were stained with Masson's trichrome, which revealed increased

presence of collagen fibers in $Aire^{-/-}$ epididymides **(Fig. 4)**. We also probed for αSMA+

myofibroblast using immunohistochemistry, which are pertinent to fibrogenesis and scar

formation[30] (**Supplemental Fig. 1**). These cells were abundant in the interstitium of $Aire^{-/-}$

epididymis compared to $Aire^{+/+}$, indicating their potential involvement in fibrogenic

transformation of the interstitium in our *Aire*$^{-/-}$ animals. The epididymis of the age-matched *Aire*$^{-}$

$^{/-}$;*Rag1*$^{-/-}$ animals, which possess innate immune cells but lack mature T and B lymphocytes, did

not develop fibrosis (**Fig. 4, third column**), demonstrating that autoreactive adaptive immune cells

are required to promote fibrogenesis of the epididymis.

Dysfunction in iron metabolism and homeostasis can cause hemochromatosis and idiopathic

pulmonary fibrosis, both of which are characterized by extensive fibrotic changes of the organ.[31,32]

To determine whether iron accumulation contributed to fibrosis in *Aire*$^{-/-}$ epididymis, we dissected

the epididymis into its anatomically distinct regions (**Fig. 2B**) and used ICP-QQQ-MS for

elemental quantification. Using this method, we found that ^{56}Fe concentration (µg/g tissue)

regardless of age was elevated throughout the epididymis from *Aire*$^{-/-}$ males compared to *Aire*$^{+/+}$

and *Aire*$^{-/-}$;*Rag1*$^{-/-}$ (**Fig. 5A**). In *Aire*$^{+/+}$ epididymides, the mean concentration of ^{56}Fe was

comparable between caput, corpus, and cauda (21.5 ± 5.2 µg/g; 26.8 ± 10.1 µg/g; 29.8 ± 4.9µg/g,

respectively). However, in *Aire*$^{-/-}$ epididymides, concentrations were 3-4-fold higher (caput, 69.8

± 51.3 µg/g; corpus, 110.1 ± 35.7 µg/g; cauda, 85.2 µg/g ± 16.0 µg/g). The ^{56}Fe concentration in

each of the epididymal segments of the *Aire*$^{-/-}$;*Rag1*$^{-/-}$ tissues were comparable to *Aire*$^{+/+}$ tissues

and values were not significantly different (**Fig. 5A**). ^{56}Fe concentration in the caudal sperm

fraction was similar between *Aire*$^{+/+}$ and *Aire*$^{-/-}$, suggesting that increased ^{56}Fe is localized to the

tissue rather than the sperm (**Fig. 5B**).

To visualize the spatial distribution of iron in the epididymis, we used laser ablation – inductively

coupled plasma – time of flight – mass spectrometry (LA-ICP-TOF-MS) (**Fig. 5C**). Analogous to

the spatial single-cell RNA sequencing that illuminates transcriptomics of a sample on a per-pixel

basis, LA-ICP-TOF-MS can delineate the total elemental content of a sample and instantaneously

allow spatial visualization of all elements within the periodic table in the tissue sample. Using this

technique, we found that ^{56}Fe was especially concentrated within the interstitium of the corpus and

cauda regions of both the young and the aged *Aire$^{-/-}$* epididymides; this was not observed in either

Aire$^{+/+}$ or *Aire$^{-/-}$;Rag1$^{-/-}$* tissues **(Fig. 5C).**

F4/80+ macrophages are sources of ^{56}Fe in the interstitium

We next asked what cells the excess iron originates from in *Aire$^{-/-}$* epididymides. Because of known

associations between macrophages and iron in fibrosis of the lung,[33] we hypothesized that these

cells are a source of iron. We first used a macrophage-specific marker, F4/80, to characterize the

distribution of macrophages in the epididymis of *Aire$^{+/+}$*, *Aire$^{-/-}$* and *Aire$^{-/-}$;Rag1$^{-/-}$* animals using

immunohistochemistry (**Fig. 6A**). In the control tissues, F4/80+ macrophages were found as small,

densely stained cells in the interstitium of the epididymis. In *Aire$^{-/-}$* tissues, macrophages became

elongated and were found throughout the interstitium of *Aire$^{-/-}$* epididymis (**Fig. 6A),**

corresponding with the locations of fibrosis and iron accumulation (**Fig. 4, Fig. 5**). F4/80+

macrophages found within fibrotic regions of the epididymis were also CD163+ (**Fig. 6B**), a

marker commonly used to identify alternatively activated macrophages with profibrotic

transcriptomic signatures.[34] Collectively, the results suggest that macrophages with a pro-fibrotic

phenotype contribute to interstitial fibrosis in the *Aire$^{-/-}$* epididymis.

Due to the ability of macrophages to store iron in the form of ferritin and hemosiderin, we asked

whether iron co-localized with F4/80+ macrophages. Indeed, we found that ferritin, a 24-mer iron

storage protein,[35] often co-localized with F4/80-positive macrophages, particularly in Aire$^{-/-}$

animals (**Fig. 6C**). Interestingly, in wild type animals, ferritin was found in and F4/80 overlapped

less faithfully, suggesting heterogeneity of iron regulatory mechanisms in F4/80+ macrophages,

an/or the presence of iron-carrying non-macrophages such as eosinophils or neutrophils.[36,37] To

further confirm the asociation between iron and macrophages in Aire[-/-] mice, we labelled

epididymides with a primary antibody for F4/80 followed by a secondary [172]Yb-tagged antibody,

and in the same sections, produced elemental maps using LA-ICP-TOF-MS. The resulting images

were co-registered on a per-pixel bases and showed co-localization of [172]Yb-positive pixels with

the [56]Fe signature, strongly suggesting that macrophages are the source of excess intracellular iron

(**Fig. 6D**).

Excess iron in tissues could lead to reactive oxygen species through the Fenton reaction,[38–40] and

in turn could cause DNA damage. Therefore, we measured DNA fragmentation in sperm isolated

from caudal epididymis of *Aire*[+/+] or *Aire*[+/-] and *Aire*[-/-] males using the comet assay, which reveals

single- and double-stranded breaks in the DNA. We found an increased percentage of sperm

containing DNA damage, as revealed by "comet tails" in *Aire*[-/-] samples compared to controls (**Fig.**

7; *Aire*[+/+], $2.35 \pm 0.92\%$ vs. *Aire*[-/-], 7.98 ± 3.69).

Decreased concentrations of Cu, Zn, and Se also correlate with increased sperm DNA

fragmentation in *Aire*[-/-] animals

Oxidative stress due to increased reactive oxygen species (ROS) can result in DNA damage in

cells.[41,42] Because sperm are transcriptionally inert, they do not possess the ability for *de novo*

synthesis of antioxidants and thus are especially prone to oxidative damage. To protect the sperm

from such assault, the epididymal epithelial cells secrete antioxidant enzymes - glutathione

peroxidases (Gpxs) and peroxiredoxins - into the luminal compartment[12,19,43] for which essential

elements such as Cu, Zn, and Se are of core structural component. To test this, we measured the

concentration of metals from the caput, corpus, and cauda epididymis using ICP-QQQ-MS.

Compared to *Aire*[+/+] and *Aire*[-/-]*;Rag1*[-/-] samples, concentrations of Cu, Zu, and Se were reduced,

particularly in cauda epididymis of *Aire*[-/-] animals **(Supplemental Fig. 2, A-E).** Elemental analysis

by LA-ICP-TOF-MS corroborated the findings for ^{63}Cu and ^{64}Zn, which were strikingly reduced

in *Aire*[-/-] epididymis as compared to that of *Aire*[+/+] controls.

Discussion

Male-factor infertility, which accounts for about half of these cases, can stem from several

problems, including underlying autoimmune disease. In this instance, autoreactive lymphocytes

may target the generative sources of male germ cells – the testis and epididymis – perturbing sperm

maturation and negatively impacting fertility. We previously showed that male mice deficient in

Aire, the transcription factor critical for establishing self-antigen specific tolerance to developing

T cells in the thymus, suffered from subfertility with increased infiltration of CD3+ T cells into

the testis and generation of anti-testis and anti-epididymal antibodies.[10] Here, we report increased

numbers of activated T cells and macrophages within the epididymis, which was associated with

disruptions in inorganic physiology and sperm DNA damage.

We derived an *Aire*-deficient mouse model by targeting exon 2 of *Aire* using CRISPR/Cas9

technology. These mice recapitulated prior findings of *Aire* deficiency, in that both endocrine and

non-endocrine organs showed evidence of autoimmunity as well as severe male subfertility.

Importantly, the absence of adaptive immune cells, achieved by crossing Aire-deficient mice to

Rag1-deficient mice, completely restored fertility. This observation conclusively demonstrates that infertility in *Aire*[-/-] males is driven by the presence of autoreactive lymphocytes rather than a germ cell or reproductive tract intrinsic role for Aire.

In many animals, we found high numbers of CD4+ and CD8+ T cells, as well as B cells, in the epididymis of *Aire*[-/-] males, while we did not observe testicular abnormalities. This result suggests that infertility may arise from impaired epididymal function. Using immunohistochemistry, we found that T cell abundance varied between animals but also showed as age- and region-specific differences in the epididymis. Additionally, flow cytometry revealed increased proportions of activated (CD69+) CD8+ T cells in all segments of the epididymis of *Aire*[-/-] animals. While CD44 expression was high on nearly all T cells regardless of genotype, the late activation marker, CD25, was upregulated in only CD4+ T cells from caudal *Aire*[-/-] epididymis, possibly indicating a memory phenotype in this region. Together, these data suggest that Aire deficiency promotes activation of CD8+ and CD4+ T cells, potentially due to chronic interaction between the T cell receptor and self-antigens presented in the context of MHC class I or class II in the Aire[-/-] epididymal environment. Future experiments will include additional markers that will reveal memory and exhaustion phenotypes of these cells.

In addition to characterizing the immune cell infiltrates, we set out to understand the potential mechanism of fibrosis in autoimmune-mediated epididymitis by combining immunohistochemical staining and mass spectrometry mapping of the metal content using LA-ICP-TOF-MS and ICP-QQQ-MS. While epididymal fibrosis is a hallmark feature of inflamed epididymis due to infection or experimentally induced autoimmune epididymitis,[44,45] our findings of interstitial iron overload as well as ferritin-positive interstitial macrophages are reminiscent of hemochromatosis and

bleomycin-induced idiopathic pulmonary fibrosis where the associations between iron,

macrophages, and fibrosis are well-established.[31,33,46–48] It should be recognized that, although

macrophages likely account for most F4/80+ cells, other cells express this marker (eg, a subset of

dendritic cells[49], and eosinophils[36]) and thus may also be responsible for carrying iron.

Conversely, ferritin+/F480-negative cells may also account for excess iron in the Aire[-/-]

epididymides. Dysregulation of iron chemistry can promote oxidative damage through Fenton

reaction and increase hydroxyl radicals that readily cause cellular DNA damage, which may

instigate fragmentation of sperm DNA and fibrosis of the interstitium through the wound healing

process. While the sequence of events – whether accumulation of iron promotes fibrosis such as

that observed in the lung[48] or vice versa[33] – is unknown, we surmise that the autoimmune injury

to the epididymis activates F4/80+ interstitial macrophages evidenced by their morphological

changes, which by working together with myofibroblasts, promotes fibrosis of the epididymal

interstitium.[50]

We show that autoimmune-mediated epididymitis decreases the concentrations of essential metals

in the epididymis, including Cu, Zn, and Se. These findings were surprising because imbalance of

these elements in seminal fluid is associated with sperm abnormality and infertility in humans,[51,52]

To protect the sperm from oxidative damage, the epididymal epithelial cells express a wide variety

of antioxidant enzymes including Cu-Zn superoxide dismutases (including SOD1 and SOD3[53]),

glutathione peroxidases (Gpxs[11,54–56]), and thioredoxin-glutathione reductase (i.e. TXNRD3[57]).

These enzymes have been shown to be released into the lumen in the form of epididymosomes by

the epithelial cells.[58] Deficiency of these essential cofactors may confer deleterious effect on the

activity of key antioxidant enzymes including SODs and selenoproteins TXNRD3, Gpx1, 3, and

4, exposing sperm to reactive oxygen species-mediated oxidative stress and DNA damage.[59,60] We

speculate that decreases in Cu, Zn, and Se lead to defects in antioxidant enzyme production or

function, resulting in sperm DNA damage. While further studies are required to confirm levels of

antioxidant activity and production of reactive oxygen species, this model provides a plausible

explanation for why sperm from *Aire*[-/-] males fail to allow embryo progression to blastocyst stage

upon *in vitro* fertilization.[60,61]

In summary, we demonstrate that Aire deficiency and resultant autoimmune attack to the

epididymis disrupts inorganic physiology of the epididymis, in particular accumulation of Fe in

the interstitial space, and decreases in Zn, Se, Cu, which may in turn promote increased ROS

production and damage to sperm DNA. Interstitial macrophages as well as myofibroblasts likely

play a significant role in the fibrogenesis using similar mechanisms found in the liver and the

lung.[62] Based on the results from *Aire*[-/-]*;Rag1*[-/-] animals, we demonstrate that fibrogenesis, Fe

accumulation, and Zn, Cu and Se depletion in the interstitium require the presence of adaptive

immune cells. Future studies will focus on further characterizing these immune population in the

Aire[-/-] epididymis to delineate how they participate in the fibrogenic transformation of the

epididymis and promote imbalance of essential metals that are critical to maintain male fertility.

Further studies are necessary to understand the mechanisms that contribute to decreases in essential

metals such as Se, Cu, and Zn in autoimmune-mediated subfertility.

Acknowledgements

We thank Dr. Lindsay Moritz and Dr. Sue Hammoud at University of Michigan for the provision

of the comet assay protocol. Metal analysis using ICP-QQQ-MS and LA-ICP-TOF-MS was

performed at the Quantitative Bio Element Analysis and Mapping Center (QBEAM) located at

Michigan State University, East Lansing, MI, USA. Figures were created using BioRender.

536
537

References

1. *Infertility Prevalence Estimates, 1990–2021. Geneva: World Health Organization*.; 2023.

2. Practice Committee of the American Society for Reproductive Medicine. Diagnostic
evaluation of the infertile male: a committee opinion. *Fertil Steril*. 2015;103(3):e18-25.

3. Sharma A, Minhas S, Dhillo WS, Jayasena CN. Male infertility due to testicular disorders. *J*
Clin Endocrinol Metab. 2021;106(2):e442-e459.

4. Hamada A, Esteves SC, Agarwal A. Unexplained male infertility. *Hum Androl*. 2011;1(1):2-
16.

5. Finelli R, Leisegang K, Finocchi F, De Masi S, Agarwal A, Damiani G. The impact of
autoimmune systemic inflammation and associated medications on male reproductive health
in patients with chronic rheumatological, dermatological, and gastroenterological diseases: A
systematic review. *Am J Reprod Immunol*. 2021;85(5):e13389.

6. Anderson MS, Venanzi ES, Klein L, et al. Projection of an immunological self shadow within
the thymus by the aire protein. *Science*. 2002;298(5597):1395-1401.

7. Ahonen P, Myllärniemi S, Sipilä I, Perheentupa J. Clinical variation of autoimmune
polyendocrinopathy–candidiasis–ectodermal dystrophy (APECED) in a series of 68 patients.
N Engl J Med. 1990;322(26):1829-1836.

8. Eisenbarth GS, Gottlieb PA. Autoimmune polyendocrine syndromes. *N Engl J Med*.
2004;350(20):2068-2079.

9. Michels AW, Gottlieb PA. Autoimmune polyglandular syndromes. *Nat Rev Endocrinol*.
2010;6(5):270-277.

10. Warren BD, Ahn SH, Brittain KS, et al. Multiple lesions contribute to infertility in males
lacking autoimmune regulator. *Am J Pathol*. 2021;191(9):1592-1609.

11. Hinton BT, Palladino MA, Rudolph D, Lan ZJ, Labus JC. The role of the epididymis in the
protection of spermatozoa. *Curr Top Dev Biol*. 1996;33:61-102.

12. O'Flaherty C. Orchestrating the antioxidant defenses in the epididymis. *Andrology*.
2019;7(5):662-668.

13. Gervasi MG, Visconti PE. Molecular changes and signaling events occurring in spermatozoa
during epididymal maturation. *Andrology*. 2017;5(2):204-218.

14. Fournier-Delpech S, Colas G, Courot M. The first cleavage of tubal sheep eggs after fertilization with epididymal or ejaculated spermatozoa. *C R Seances Acad Sci III*. 1981;292(7):515-517.

15. Rinaldi VD, Donnard E, Gellatly K, et al. An atlas of cell types in the mouse epididymis and vas deferens. *Elife*. 2020;9.

16. Dacheux JL, Gatti JL, Dacheux F. Contribution of epididymal secretory proteins for spermatozoa maturation. *Microsc Res Tech*. 2003;61(1):7-17.

17. Da Silva N, Shum WWC, Breton S. Regulation of vacuolar proton pumping ATPase-dependent luminal acidification in the epididymis. *Asian J Androl*. 2007;9(4):476-482.

18. Carr DW, Usselman MC, Acott TS. Effects of pH, lactate, and viscoelastic drag on sperm motility: A species Comparison1. *Biol Reprod*. 1985;33(3):588-595.

19. Ingold I, Aichler M, Yefremova E, et al. Expression of a catalytically inactive mutant form of glutathione peroxidase 4 (Gpx4) confers a dominant-negative effect in male fertility. *J Biol Chem*. 2015;290(23):14668-14678.

20. Chabory E, Damon C, Lenoir A, et al. Epididymis seleno-independent glutathione peroxidase 5 maintains sperm DNA integrity in mice. *J Clin Invest*. Published online June 22, 2009.

21. Dou Q, Turanov AA, Mariotti M, et al. Selenoprotein TXNRD3 supports male fertility via the redox regulation of spermatogenesis. *J Biol Chem*. 2022;298(8):102183.

22. Bettger WJ. Zinc and selenium, site-specific versus general antioxidation. *Can J Physiol Pharmacol*. 1993;71(9):721-724.

23. Hamilton LE, Oko R, Miranda-Vizuete A, Sutovsky P. Sperm redox system equilibrium: Implications for fertilization and male fertility. *Adv Exp Med Biol*. 2022;1358:345-367.

24. Zee DZ, MacRenaris KW, O'Halloran TV. Quantitative imaging approaches to understanding biological processing of metal ions. *Curr Opin Chem Biol*. 2022;69(102152):102152.

25. Albus U. Guide for the Care and Use of Laboratory Animals (8th edn). *Lab Anim*. 2012;46(3):267-268.

26. Ding D, Liu J, Dong K, et al. PNLDC1 is essential for piRNA 3′ end trimming and transposon silencing during spermatogenesis in mice. *Nat Commun*. 2017;8(1).

27. Croy A, Yamada AT, Demayo FJ, Adamson SL. *The Guide to Investigation of Mouse Pregnancy*. Elsevier Gezondheidszorg; 2018.

28. Paton C, Hellstrom J, Paul B, Woodhead J, Hergt J. Iolite: Freeware for the visualisation and processing of mass spectrometric data. *J Anal At Spectrom*. 2011;26(12):2508.

29. Jasti S, Warren BD, McGinnis LK, Kinsey WH, Petroff BK, Petroff MG. The autoimmune regulator prevents premature reproductive senescence in female mice. *Biol Reprod*. 2012;86(4):110.

30. Tai Y, Woods EL, Dally J, et al. Myofibroblasts: Function, formation, and scope of molecular
therapies for skin fibrosis. *Biomolecules*. 2021;11(8):1095.

31. Mehta KJ, Farnaud SJ, Sharp PA. Iron and liver fibrosis: Mechanistic and clinical aspects.
World J Gastroenterol. 2019;25(5):521-538.

32. Kowdley KV. Iron overload in patients with chronic liver disease. *Gastroenterol Hepatol (N
Y)*. 2016;12(11):695-698.

33. Ali MK, Kim RY, Brown AC, et al. Critical role for iron accumulation in the pathogenesis of
fibrotic lung disease. *J Pathol*. 2020;251(1):49-62.

34. Wendisch D, Dietrich O, Mari T, et al. SARS-CoV-2 infection triggers profibrotic
macrophage responses and lung fibrosis. *Cell*. 2021;184(26):6243-6261.e27.

35. Munro HN, Linder MC. Ferritin: structure, biosynthesis, and role in iron metabolism. *Physiol
Rev*. 1978;58(2):317-396.

36. McGarry MP, Stewart CC. Murine eosinophil granulocytes bind the murine macrophage-
monocyte specific monoclonal antibody F4/80. *J Leukoc Biol*. 1991;50(5):471-478.

37. Sasmono RT, Ehrnsperger A, Cronau SL, et al. Mouse neutrophilic granulocytes express
mRNA encoding the macrophage colony-stimulating factor receptor (CSF-1R) as well as
many other macrophage-specific transcripts and can transdifferentiate into macrophages in
vitro in response to CSF-1. *J Leukoc Biol*. 2007;82(1):111-123.

38. Bansal AK, Bilaspuri GS. Effect of ferrous sulphate and ascorbic acid on motility, viability
and lipid peroxidation of crossbred cattle bull spermatozoa. *Animal*. 2008;2(1):100-104.

39. Aitken RJ, Wingate JK, De Iuliis GN, McLaughlin EA. Analysis of lipid peroxidation in
human spermatozoa using BODIPY C11. *Mol Hum Reprod*. 2007;13(4):203-211.

40. Papanikolaou G, Pantopoulos K. Iron metabolism and toxicity. *Toxicol Appl Pharmacol*.
2005;202(2):199-211.

41. Gonzalez-Hunt CP, Wadhwa M, Sanders LH. DNA damage by oxidative stress: Measurement
strategies for two genomes. *Curr Opin Toxicol*. 2018;7:87-94.

42. Loft S. Oxidative DNA damage in human sperm influences time to pregnancy. *Hum Reprod*.
2003;18(6):1265-1272.

43. Chabory E, Damon C, Lenoir A, et al. Epididymis seleno-independent glutathione peroxidase
5 maintains sperm DNA integrity in mice. *J Clin Invest*. 2009;119(7):2074-2085.

44. Wijayarathna R, Pasalic A, Nicolas N, et al. Region-specific immune responses to
autoimmune epididymitis in the murine reproductive tract. *Cell Tissue Res*. 2020;381(2):351-
360.

45. Pleuger C, Ai D, Hoppe ML, et al. The regional distribution of resident immune cells shapes
distinct immunological environments along the murine epididymis. *Elife*. 2022;11.

46. Pei Z, Qin Y, Fu X, et al. Inhibition of ferroptosis and iron accumulation alleviates pulmonary fibrosis in a bleomycin model. *Redox Biol*. 2022;57(102509):102509.

47. Misharin AV, Morales-Nebreda L, Reyfman PA, et al. Monocyte-derived alveolar macrophages drive lung fibrosis and persist in the lung over the life span. *J Exp Med*. 2017;214(8):2387-2404.

48. Maus M, López-Polo V, Mateo L, et al. Iron accumulation drives fibrosis, senescence and the senescence-associated secretory phenotype. *Nat Metab*. 2023;5(12):2111-2130.

49. Sun X, Jones HP, Dobbs N, Bodhankar S, Simecka JW. Dendritic cells are the major antigen presenting cells in inflammatory lesions of murine Mycoplasma respiratory disease. *PLoS One*. 2013;8(2):e55984.

50. Adler M, Mayo A, Zhou X, et al. Principles of cell circuits for tissue repair and fibrosis. *iScience*. 2020;23(2):100841.

51. Bleau G, Lemarbre J, Faucher G, Roberts KD, Chapdelaine A. Semen selenium and human fertility. *Fertil Steril*. 1984;42(6):890-894.

52. Nenkova G, Petrov L, Alexandrova A. Role of trace elements for oxidative status and quality of human sperm. *Balkan Med J*. 2017;34(4):343.

53. Alvarez JG, Storey BT. Role of superoxide dismutase in protecting rabbit spermatozoa from O2 toxicity due to lipid peroxidation. *Biol Reprod*. 1983;28(5):1129-1136.

54. Schwaab V, Faure J, Dufaure JP, Drevet JR. GPx3: the plasma-type glutathione peroxidase is expressed under androgenic control in the mouse epididymis and vas deferens. *Mol Reprod Dev*. 1998;51(4):362-372.

55. Noblanc A, Kocer A, Chabory E, et al. Glutathione peroxidases at work on epididymal spermatozoa: An example of the dual effect of reactive oxygen species on mammalian male fertilizing ability. *J Androl*. 2011;32(6):641-650.

56. Drevet JR. The antioxidant glutathione peroxidase family and spermatozoa: A complex story. *Mol Cell Endocrinol*. 2006;250(1-2):70-79.

57. Wang H, Dou Q, Jeong KJ, Choi J, Gladyshev VN, Chung JJ. Redox regulation by TXNRD3 during epididymal maturation underlies capacitation-associated mitochondrial activity and sperm motility in mice. *J Biol Chem*. 2022;298(7):102077.

58. Barrachina F, Ottino K, Elizagaray ML, et al. Regulatory T cells play a crucial role in maintaining sperm tolerance and male fertility. *Proc Natl Acad Sci U S A*. 2023;120(37).

59. Lopes S M Sc, Sun JG PhD, Jurisicova A PhD, Meriano J B Sc, Casper RF MD. Sperm deoxyribonucleic acid fragmentation is increased in poor-quality semen samples and correlates with failed fertilization in intracytoplasmic sperm injection. *Fertil Steril*. 1998;69(3):528-532.

60. Seli E, Gardner DK, Schoolcraft WB, Moffatt O, Sakkas D. Extent of nuclear DNA damage
in ejaculated spermatozoa impacts on blastocyst development after in vitro fertilization. *Fertil*
Steril. 2004;82(2):378-383.

61. Lawler T, Parlato L, Warren Andersen S. Racial disparities in colorectal cancer
clinicopathological and molecular tumor characteristics: a systematic review. *Cancer Causes*
Control. Published online September 9, 2023.

62. Chung S, Overstreet JM, Li Y, et al. TGF-β promotes fibrosis after severe acute kidney injury
by enhancing renal macrophage infiltration. *JCI Insight*. 2018;3(21).

 Figure Legends

Figure 1. Aire-deficient (*Aire*^{-/-}) males develop autoimmunity towards non-reproductive and

reproductive organs and are subfertile. (A) Immunofluorescence of Aire in *Aire*$^{-/-}$ and *Aire*$^{+/+}$

thymi. AF488 (green), Aire; AF568 (red), Cytokeratin 14 (K14, mTEC marker); DAPI (blue),

nuclei. Asterisks highlight areas in which additional Aire$^+$ cells can be seen outside the inset. Note

the absence of the characteristic intranuclear speckles in the *Aire*$^{-/-}$ thymus. Scale bar = 10μm. (B)

Representative hematoxylin and eosin-stained lung, salivary gland, prostate, and cauda epididymis

sections from 20 week-old *Aire*$^{-/-}$ and *Aire*$^{+/+}$ males. Lung: n=3 per genotype; prostate: *Aire*$^{-/-}$,

n=11/*Aire*$^{+/+}$, n=7; caudal epididymis: n=11 per genotype. Arrows point to lymphoid aggregates.

Scale bars = 100μm. (C) Fertility outcomes of *Aire*$^{-/-}$ and *Aire*$^{+/+}$ males on the *Rag1*-sufficient or -

deficient background. ANOVA followed by Tukey's post hoc test. ns: *P*>0.05, *PP < 0.0001.

Figure 2. Immune cell infiltrates consisting of CD4+ T cells, CD8+ T cells and CD19+ B cells

occur early and persist with age in *Aire*^{-/-} epididymides. (A) Immunohistochemical staining for

CD4+ T cells, CD8+ T cells and CD19+ B cells in the epididymis of 19-20 week old *Aire*$^{+/+}$ and

A*Aire*$^{-/-}$ epididymides. CD4+ and CD8+ T cells are also found sporadically in the *Aire*$^{+/+}$

epididymis (arrows). *Aire*$^{+/+}$(19-20 weeks: n=5)*,* and *Aire*$^{-/-}$ (19-20 weeks: n=7), Scale bar = 50

μm. (B) Photomicrograph of the three main anatomical regions of the epididymis (Caput, H;

Corpus; B; Cauda; T). (C, D) Manually enumerated CD4+ (*Aire*$^{+/+}$: n=8; *Aire*$^{-/-}$: n=9) and CD8+

(*Aire*$^{+/+}$: n=10, *Aire*$^{-/-}$ n=12) T cells in the epididymis. Red dots denote 19-20 week old animals;

black dots denote 7-9 week old animals. Statistical Analysis: Wilcoxon non-parametric t-test. ns:

p>0.05, *PP<0.01.

Figure 3. Flow cytometric analysis of CD4+ and CD8+ T cells from *Aire$^{-/-}$* and *Aire$^{+/+}$* epididymides. (A) Gating strategy for flow cytometric assessment of lymphocyte population in the epididymis. FSC-A: forward scatter area, SSC-A: side scatter area, FSC-H: forward scatter height. Epididymides were separated into Caput-Corpus and Cauda, digested into single cell suspensions, and analyzed by flow cytometry for the expression of CD25, CD44, and CD69. (B) Relative abundance of CD45hi cells in *Aire$^{+/+}$* and *Aire$^{-/-}$* epididymides; (C) Relative abundance of CD45hi B220+ cells in the epididymis; (D, F) Relative abundance of CD45hi CD4+T, followed by CD4+CD25+ CD4+CD44+, and CD4+CD69+ T cells in the epididymis; (E,G) Relative abundance of CD45hi CD8+ T cells, followed by CD8+CD25+, CD8+CD44+, and CD8+CD69+ T cells in the epididymis. Animals analyzed were 19-20 weeks old. Wilcoxon t-test, ns: *P*>0.05, **P*< 0.01.

Figure 4. Fibrotic changes in epididymides of *Aire$^{-/-}$* males. Masson's trichrome staining reveals extensive fibrosis (blue staining) in the interstitium of *Aire$^{-/-}$* epididymides of 9-10 weeks and 19-20-week-old males compared to that of *Aire$^{+/+}$* and *Aire$^{-/-}$;Rag1$^{-/-}$* controls. Shown are the cauda and portions of the corpus epididymis. Scale bar = 1000µm.

Figure 5. ^{56}Fe concentration is increased in the *Aire$^{-/-}$* epididymis and localizes to the interstitium. (A) ICP-QQQ-MS analysis of epididymal segments. Red = 7-9 wks, Black = 19-20 wks. **(B)** ^{56}Fe measurement in sperm using ICP-QQQ-MS. **(C)** LA-ICP-TOF-MS reveals the spatial distribution of ^{56}Fe to the interstitium of the *Aire$^{+/+}$* (n=3 for each age), *Aire$^{-/-}$* (n=4 for each age), and *Aire$^{-/-}$;Rag1$^{-/-}$* (n=4 for 9-10wks, n=3 for 19-20wks) epididymis. Statistical analysis: Kruskal-Wallis One-Way ANOVA followed by Dunn-Bonferroni post-hoc was conducted to compare the means of each group for (A). Wilcoxon-test for (B). *****P*<0.00005, ns = not significant.

Figure 6. F4/80+ macrophages carry intracellular ferritin-associated iron in the interstitium of *Aire*[-/-] epididymis. (A) Representative immunohistochemical staining for F4/80+ macrophages in cauda epididymides of *Aire*[-/-] compared to *Aire*[+/+] and *Aire*[-/-];*Rag1*[-/-] mice. *Aire*[+/+](7-10 weeks: n=4, 19-20 weeks: n=5), *Aire*[-/-] (7-10 weeks: n= 4, 19-20 weeks: n=7), and *Aire*[-/-];*Rag1*[-/-] (7-10 weeks: n=4, 19-20 weeks: n=3) Scale bar = 100 μm. **(B)** Immunofluorescence staining for F4/80+ (Alexa Fluor 488, green) and CD163+ (Alexa Fluor 546, red) macrophages in *Aire*[+/+] and *Aire*[-/-] epididymal tail. Scale bar = 50 μm. **(C)** Dual immunofluorescence staining for F4/80 (Alexa Fluor 488, green) and ferritin (Alexa Fluor 546, red) in *Aire*[+/+] and *Aire*[-/-] cauda epididymis. Scale bar = 100 μm. **((D)** Representative LA-ICP-TOF-MS images of *Aire*[-/-] epididymis stained with F4/80 primary antibody followed by a secondary antibody tagged to a lanthanide ([172]Yb) revealing colocalization of [56]Fe with F4/80+ macrophages. Scale bar = 1000μm.

Figure 7. Comet assay reveal increased DNA fragmentation in caudal sperm from *Aire*[-/-] males. Epididymal tail sperm from *Aire*[+/+] or *Aire*[+/-] and *Aire*[-/-] males aged between 7 to 20 weeks were processed for comet analysis. Scale bar = 1000μm *p = 0.048, paired t-test.

Supplementary Table 1. Elemental concentration in the gelatin standards determined by ICP-QQQ-MS.

Supplementary Table 2. LA-ICP-TOF-MS parameters.

Supplementary Figure 1. αSMA+ myofibroblasts abound in the interstitium of *Aire*[-/-] cauda epididymis in comparison to *Aire*[+/+] epididymis of similar age. Scale bar = 200 μm for top and bottom row; 50 μm for middle row

Supplementary Figure 2. Concentrations of ^{63}Cu and ^{64}Zn are decreased in *Aire$^{-/-}$*

epididymis. (A-C) ICP-QQQ-MS analysis of individual segments of the epididymis show

regional decreases in ^{63}Cu, ^{64}Zn, and ^{78}Se in the *Aire$^{-/-}$* epididymis compared to *Aire$^{+/+}$* or *Aire$^-$*

$^{/-}$;Rag1$^{-/-}$. Red = 7-9 weeks old, Black = 19-20 weeks old. Elemental distribution of ^{63}Cu

Statistical analysis: Kruskal-Wallis One-Way ANOVA followed by Dunn-Bonferroni post-hoc

was conducted to compare the means of each group. *p**(D)** and ^{64}Zn **(E)** in 20-week-old *Aire$^{+/+}$*

and *Aire$^{-/-}$* epididymis using LA-ICP-TOF-MS; Se abundance was below the detection limit of

this technology. Scale bar = 1000 μm.

A

anti-Aire
anti-K14
DAPI

Inset

Aire$^{+/+}$

Aire$^{-/-}$

B

	Aire$^{+/+}$	*Aire*$^{-/-}$
Lung		
Salivary Gland		
Prostate Gland		
Caudal epididymis		

C

Litters/21 days

Aire	+/+	+/+	-/-	-/-
Rag1 +/+ ; +/-	+/+	-/-	+/+	-/-

Figure 1. **Aire-deficient (*Aire^-/-^*) males develop autoimmunity towards non-reproductive and reproductive organs and are subfertile. (A)** Immunofluorescence of Aire in *Aire^-/-^* and *Aire^+/+^* thymi. AF488 (green), Aire; AF568 (red), Cytokeratin 14 (K14, mTEC marker); DAPI (blue), nuclei. Asterisks highlight areas in which additional Aire$^+$ cells can be seen outside the inset. Note the absence of the characteristic intranuclear speckles in the *Aire^-/-^* thymus. Scale bar = 10µm. **(B)** Representative hematoxylin and eosin-stained lung, salivary gland, prostate, and cauda epididymis sections from 20 week-old *Aire^-/-^* and *Aire^+/+^* males. Lung: n=3 per genotype; prostate: *Aire^-/-^*, n=11/*Aire^+/+^*, n=7; caudal epididymis:n=11 per genotype. Arrows point to lymphoid aggregates. Scale bars = 100 µm. **(C)** Fertility outcomes of *Aire^-/-^* and *Aire^+/+^* males on the *Rag1*-sufficient or -deficient background. ANOVA followed by Tukey's post hoc test. ns: *P*>0.05, **P*<0.05, *****P* < 0.0001.

Figure 2. Immune cell infiltrates consisting of CD4+ T cells, CD8+ T cells and CD19+ B cells occur early and persist with age in *Aire$^{-/-}$* epididymides. (A) Immunohistochemical staining for CD4+ T cells, CD8+ T cells and CD19+ B cells in the epididymis of 19-20 week old *Aire$^{+/+}$* and A*ire$^{-/-}$* epididymides. CD4+ and CD8+ T cells are also found sporadically in the *Aire$^{+/+}$* epididymis (arrows). *Aire$^{+/+}$*(19-20 weeks: n=5) ,and *Aire$^{-/-}$* (19-20 weeks: n=7), Scale bar = 50 μm. **(B)** Photomicrograph of the three main anatomical regions of the epididymis (Caput, H; Corpus; B; Cauda; T). **(C, D)** Manually enumerated CD4+ (*Aire$^{+/+}$*: n=8; *Aire$^{-/-}$*: n=9) and CD8+ (*Aire$^{+/+}$*: n=10, *Aire$^{-/-}$*: n=12) T cells in the epididymis. Red dots denote 19-20 week old animals; black dots denote 7-9 week old animals. Statistical Analysis: Wilcoxon non-parametric t-test. ns: p>0.05, *P<0.05, **P<0.01.

Figure 3. Flow cytometric analysis of CD4+ and CD8+ T cells from *Aire*$^{-/-}$ and *Aire*$^{+/+}$ epididymides. (A) Gating strategy for flow cytometric assessment of lymphocyte population in the epididymis. FSC-A: forward scatter area, SSC-A: side scatter area, FSC-H: forward scatter height. Epididymides were separated into Caput-Corpus and Cauda, digested into single cell suspensions, and analyzed by flow cytometry for the expression of CD25, CD44, and CD69. **(B)** Relative abundance of CD45hi cells in *Aire*$^{+/+}$ and *Aire*$^{-/-}$ epididymides; **(C)** Relative abundance of CD45hi B220+ cells in the epididymis; **(D, F)** Relative abundance of CD45hi CD4+T, followed by CD4+CD25+ CD4+CD44+, and CD4+CD69+ T cells in the epididymis; **(E,G)** Relative abundance of CD45hi CD8+ T cells, followed by CD8+CD25+, CD8+CD44+, and CD8+CD69+ T cells in the epididymis. Wilcoxon t-test, ns: *P*>0.05, **P*<0.05, **p < 0.01.

Figure 4. Masson's trichrome staining reveals extensive fibrosis (blue staining) in the interstitium of *Aire^-/-* epididymides of 9-10 weeks and 19-20-week-old males compared to that of *Aire^+/+* and *Aire^-/-;Rag1^-/-* controls. Shown are the cauda and portions of the corpus epididymis. Scale bar = 1000 μm.

Figure 5. ^{56}Fe concentration is increased in the *Aire$^{-/-}$* epididymis and localizes to the interstitium. (A) ICP-QQQ-MS analysis of epididymal segments. Red = 7-9 wks, Black = 19-20 wks. **(B)** ^{56}Fe measurement in sperm using ICP-QQQ-MS. **(C)** LA-ICP-TOF-MS reveals the spatial distribution of ^{56}Fe to the interstitium of the *Aire$^{+/+}$* (n=3 for each age), *Aire$^{-/-}$* (n=4 for each age), and *Aire$^{-/-}$;Rag1$^{-/-}$* (n=4 for 9-10wks, n=3 for 19-20wks) epididymis. Statistical analysis: Kruskal-Wallis One-Way ANOVA followed by Dunn-Bonferroni post-hoc was conducted to compare the means of each group for (A). Wilcoxon-test for (B). ****P<0.00005, ns = not significant.

Figure 6. F4/80+ macrophages carry intracellular ferritin-associated iron in the interstitium of *Aire*[-/-] epididymis. (A) Representative immunohistochemical staining for F4/80+ macrophages in cauda epididymides of *Aire*[-/-] compared to *Aire*[+/+] and *Aire*[-/-];*Rag1*[-/-] mice. *Aire*[+/+](7-10 weeks: n=4, 19-20 weeks: n=5), *Aire*[-/-] (7-10 weeks: n= 4, 19-20 weeks: n=7), and *Aire*[-/-];*Rag1*[-/-] (7-10 weeks: n=4, 19-20 weeks: n=3) Scale bar = 100 μm. **(B)** Immunofluorescence staining for F4/80+ (Alexa Fluor 488, green) and CD163+ (Alexa Fluor 546, red) macrophages in *Aire*[+/+] and *Aire*[-/-] epididymal tail. Scale bar = 50 μm. **(C)** Dual immunofluorescence staining for F4/80 (Alexa Fluor 488, green) and ferritin (Alexa Fluor 546, red) in *Aire*[+/+] and *Aire*[-/-] cauda epididymis. Scale bar = 100 μm. **(D)** Representative LA-ICP-TOF-MS images of *Aire*[-/-] epididymis stained with F4/80 primary antibody followed by a secondary antibody tagged to a lanthanide (^{172}Yb) revealing colocalization of ^{56}Fe with F4/80+ macrophages. Scale bar = 1000 μm.

Figure 7. Comet assay reveal increased DNA fragmentation in caudal sperm from *Aire*$^{-/-}$ **males.** Epididymal tail sperm from *Aire*$^{+/+}$ or *Aire*$^{+/-}$ and *Aire*$^{-/-}$ males aged between 7 to 20 weeks were processed for comet analysis. Scale bar = 1000 μm *p = 0.048, paired t-test.

Supplemental Figure 1. αSMA+ myofibroblasts abound in the interstitium of *Aire*-/- cauda epididymis in comparison to *Aire*+/+ epididymis of similar age. Scale bar = 200 μm for top and bottom row; 50 μm for middle row

Supplementary Figure 2. Concentrations of ^{63}Cu and ^{64}Zn are decreased in *Aire*$^{-/-}$ epididymis. (A-C) ICP-QQQ-MS analysis of individual segments of the epididymis show regional decreases in ^{63}Cu, ^{64}Zn, and ^{78}Se in the *Aire*$^{-/-}$ epididymis compared to *Aire*$^{+/+}$ or *Aire*$^{-/-}$;*Rag1*$^{-/-}$. Red = 7-9 weeks old, Black = 19-20 weeks old. Elemental distribution of ^{63}Cu Statistical analysis: Kruskal-Wallis One-Way ANOVA followed by Dunn-Bonferroni post-hoc was conducted to compare the means of each group. *p<0.05, **p<0.005, ***p<0.0005, ****p<0.00005. **(D)** and ^{64}Zn **(E)** in 20-week-old *Aire*$^{+/+}$ and *Aire*$^{-/-}$ epididymis using LA-ICP-TOF-MS; Se abundance was below the detection limit of this technology. Scale bar = 1000 μm.

Supplementary Table 1. Elemental concentration in the gelatin standards

Sample Name	^{24}Mg	^{44}Ca	^{55}Mn	^{57}Fe	^{58}Ni	^{63}Cu	^{64}Zn
Gelatin Blank	2.106	0.916	0	0.679	0.013	0.030	0.096
Gelatin 50 ppm	53.32	44.85	55.40	56.99	56.55	56.93	55.02
Gelatin 20 ppm	26.62	21.17	26.39	26.97	27.03	27.64	26.34
Gelatin 10 ppm	14.38	11.01	13.66	14.14	14.01	14.13	13.08
Gelatin 5 ppm	8.512	6.498	7.535	8.074	7.409	7.821	7.243
Gelatin 1 ppm	3.460	2.113	1.681	2.564	1.768	1.832	1.776

Supplementary Table 2. LA-ICP-TOF-MS parameters

ICP-MS Paramters (Tofwerk S2)		
Parameter	**Unit**	**Value**
RF Power	W	1550
Sampling Depth	mm	4.9
Cone Material	Nickel	-
Cone Insert (STD)	mm	3.5
Plasma Gas Flow	L/min	14.0
Auxillary Gas Flow	L/min	8.0
Nebulizer Gas Flow	L/min	0.95-1.01
Measurement Mode	CCT Mode	-
CCT Gas Flow (100% He)	mL/min	5
CCT Focus lens	V	-19.5
CCT Entry Lens	V	-180
CCT Mass	V	250
CCT Bias	V	10
CCT Exit Lens	V	-200
Time-of-Flight Parameters (Tofwerk S2)		
Parameter	**Unit**	**Value**
m/z range	amu	14 - 256
Resolution	m/Δm	1000
ODG Settings	ms	24
Notch Bias	V	-80
Notch (40 amu)	V	1.6
Notch (36 amu)	V	1
Notch (28 amu)	V	2
Notch (15.8 amu)	V	2
Laser Ablation Parameters (ESL Bioimage 266 nm)		
Parameter	**Unit**	**Value**
Spot Size	μm	10
Interline Distance (y)	μm	10
Overlap (x)	μm	0
Repetition Rate	Hz	100
Laser Power	%	60
Laser Fluence	J/cm^2	15

Sample Energy	mJ	0.010-0.015
Imaging Cup Flow Rate (He)	mL/min	200
Imaging Chamber Flow Rate (He)	mL/min	250
PEEK Tubing I.D.	mm	0.75