Epigenetic phase variation in the gut microbiome enhances bacterial adaptation

Mi Ni[1], Yu Fan[1], Yujie Liu[1*], Yangmei Li[1*], Wanjin Qiao[1*], Lauren E. Davey[2,3*], Xue-Song Zhang[4*], Magdalena Ksiezarek[1], Edward Mead[1], Alan Touracheau[1], Wenyan Jiang[1], Martin J. Blaser[4], Raphael H. Valdivia[2], and Gang Fang[1#]

[1]*Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.*
[2]*Department of Integrative Immunobiology, Duke Microbiome Center, Duke University School of Medicine, Durham, NC, USA.*
[3]*Department of Biochemistry and Microbiology, University of Victoria, Victoria, BC, Canada.*
[4]*Center for Advanced Biotechnology and Medicine, Rutgers University, New Brunswick, NJ, USA.*

* These authors contributed equally to this work.

[#] *Address correspondence to* gang.fang@mssm.edu

Abstract

The human gut microbiome within the gastrointestinal tract continuously adapts to variations in diet, medications, and host physiology. A central strategy for genetic adaptation is epigenetic phase variation (ePV) mediated by bacterial DNA methylation, which can regulate gene expression, enhance clonal heterogeneity, and enable a single bacterial strain to exhibit variable phenotypic states. Genome-wide and site-specific ePV have been well characterized in human pathogens' antigenic variation and virulence factor production. However, the role of ePV in facilitating adaptation within the human microbiome remains poorly understood. Here, we comprehensively cataloged genome-wide and site-specific ePV in human infant and adult gut microbiomes. First, using long-read metagenomic sequencing, we detected genome-wide ePV mediated by complex structural variations of DNA methyltransferases, highlighting the ones associated with antibiotics or fecal microbiota transplantation. Second, we analyzed an extensive collection of public short-read metagenomic sequencing datasets, uncovering a greater prevalence of genome-wide ePV in the human gut microbiome. Third, we quantitatively detected site-specific ePVs using single-molecule methylation analysis to identify dynamic variations associated with antibiotic treatment or probiotic engraftment. Finally, we performed an

in-depth assessment of an *Akkermansia muciniphila* isolate from an infant, highlighting that ePV can regulate gene expression and enhance the bacterial adaptive capacity by employing a bet-hedging strategy to increase tolerance to differing antibiotics. Our findings indicate that epigenetic modifications are a common and broad strategy used by bacteria in the human gut to adapt to their environment.

Introduction

The human gut microbiome is continuously subjected to a myriad of changes including those due to changes in diet, medications, and physiological states.[1-3] In response to these alterations, the gut microbiome shows a remarkable adaptive capacity.[4-6] Characterizing this adaptive capacity is crucial for understanding the dynamic relationship between the gut microbiome and host physiology, especially in the context of human health and disease. While many microbiome studies are focused on profiles of microbial species and the relevant functions across different human diseases,[4,5,7-11] great progress has also been made with regard to the molecular mechanisms that underpin the microbiome's adaptive capacity including point mutations,[12-14] copy number variation,[15] structural variation (SV),[16,17] gene amplification,[18,19] invertible promoters[20] and intragenic inversions.[21] These genomic variations provide a single bacterial strain with the plasticity of expressing alternative, heritable and reversible phenotypic states, including phase variation (PV).[22,23] This mechanism increases the chances of bacterial survival and adaptation by producing a genetically diverse population that can withstand multiple stresses.[20,21]

Another mechanism of bacterial adaptation involves DNA methylation, which can regulate gene expression, enhance clonal heterogeneity and mediate epigenetic phase variation

(ePV, intra-strain epigenetic variation that leads to phenotypic differences, **Fig. 1A**).[24-27] Bacterial DNA contains three primary types of DNA methylation: N6-methyladenine (6mA), N4-methylcytosine (4mC) and 5-methylcytosine (5mC).[28] These DNA modifications are catalyzed by DNA methyltransferases (MTases) that are either part of restriction-modification systems or orphan enzymes without cognate restriction genes.[24,28] Phase variation driven by DNA methylation has been characterized in multiple human pathogens, where it facilitates processes such as antigenic variation and virulence factor production,[29-31] which can impact how the bacterium interacts with its environment and its host.[27,32-34]

ePV characterized across diverse human pathogens can be categorized into two types: genome-wide or site-specific.[35-38] Genome-wide ePV is usually mediated by genetic phase variation affecting an MTase or its corresponding specificity unit (which determines the recognition specificity, or sequence motif, of an MTase). In a diverse population, DNA MTase phase variation results in genome-wide activity of the MTase that alternates between two states, leading to varying levels of methylation across the population, typically approaching either ~100% or ~0% methylation of its target motif (**Fig. 1B**). Phase variation of the specificity unit associated with a DNA MTase can create two or more alternative genome-wide recognition motifs (**Fig. 1B**). Both types of genome-wide ePV can affect the expression of multiple downstream genes via epigenetic regulation, which provide greater flexibility than single-gene phase variation in creating transcriptional heterogeneity (e.g. *Streptococcus pneumoniae,*[39] *Haemophilus influenzae,*[40] *Helicobacter pylori,*[41] *etc.*) and selective advantages to bacterial pathogens (e.g. *Mycobacterium tuberculosis*[42]). In contrast to genome-wide ePV, site-specific ePV involves a small number (one or few) methylation sites and is usually driven by the competitive binding between a DNA MTase and other DNA binding proteins such as a transcription factor (TF) (**Fig. 1C**). Two well-characterized examples are the ePV of the *pap* operon[43] and the *agn43* gene[44] in *Escherichia coli*, where the 6mA MTase targeting 5'-GATC-3'

sites compete with Lrp and OxyR, respectively. These site-specific epigenetic states are inheritable and reversible and have the advantage of mediating phase variation without altered genomic sequences, thereby mitigating mutation costs, for example, associated with antibiotic resistance.[45] Both genome-wide and site-specific ePV facilitate the manifestation of diverse phenotypic states within a single bacterial strain population, enhancing the ability to adapt to changing conditions.[46]

DNA methylation is highly abundant and diverse among the species constituting the human gut microbiome,[47] yet the role of ePV and its impact on bacterial adaptation in the human gut remains unexplored. In this study, we aimed to address three major questions: (i) How prevalent are genome-wide and site-specific ePV in the human gut microbiome? (ii) What are the functions of genes regulated by ePV? (iii) In the context of a host, how do environmental (e.g. antibiotics) and individual-specific (e.g. fecal microbiota transplantation, FMT) selective pressures modulate ePV?

To address these questions, we systematically investigated the occurrence and implications of ePV in the human infant and adult gut microbiome (**Fig. 1D-E**). Using long-read metagenomic and meta-epigenomic analysis, we detected genome-wide ePVs mediated by complex SVs in DNA MTases and their specificity units. We observed intra-strain ePVs associated with antibiotic use within individuals, as well as between individuals in FMT. Furthermore, we built on the extensive collection of publicly available short-read metagenomic sequencing datasets (2,345 unique samples) and estimated the widespread prevalence of genome-wide ePVs mediated by SVs in DNA MTases and their specificity units across both healthy and disease-afflicted cohorts. We identified site-specific ePVs associated with probiotic engraftment and antibiotic treatment using single-molecule quantitative methylation analysis.

Importantly, we observed that an *Akkermansia muciniphila* strain in an infant gut microbiome

110 had an ePV detected following a course of amoxicillin (AMX); we recapitulated this shift *in vitro*

and functionally characterized how this ePV enhances the adaptive capacity of the *A.*

muciniphila strain in response to different antibiotics classes (**Fig. 1E**).

Results

115 **Genome-wide epigenetic phase variations in the gut microbiome**

We developed methods for joint metagenomic and meta-epigenomic analysis using

long-read DNA sequencing (**Fig. S1A**) and applied them to publicly available long-read

metagenomic sequencing data as well as new data we generated for this study (**Fig. S1B,**

Table S1). We focused on detecting SVs involving DNA MTases and/or their corresponding

120 specificity units, including invertible promoters, simple sequence repeats, and complex SVs

spanning one or multiple coding regions (**Methods**). These SVs have been linked to genome-

wide ePV in bacterial pathogens[39-42] but not in the gut microbiome. While short-read DNA

sequencing can detect invertible promoter PVs and simple repeats, long-read DNA sequencing

can resolve complex SVs by assigning individual long reads to a different possible genome

125 rearrangement built from each reference genome (**Fig. S1B, Methods**).[21,48-50] We used long-

read mapping tool (pbalign / NGMLR[50]) with additional quality controls to ensure read

homogeneity at genomic regions flanking the SVs that reflect intra-strain variation rather than

multiple strains (**Methods**). For example, from the microbiome sample from a healthy adult ,[51]

we detected a complex SV within an MTase in *Phocaeicola vulgatus* (formerly *B. vulgatus*),

130 mediated by three paired/matching inverted repeats (IRs), resulting in three possible genotypes

(g1-g3; **Fig. 2A**). Individual long reads can be unambiguously assigned to each of the three

genotypes, facilitating genotype-specific quantitation of each (**Fig. 2A**). Across the 221 long-

read metagenomic samples analyzed, we detected 174 SVs involving DNA MTase and/or

specificity units (**Table S2**), which encode 2-8 alternative genotypes based on long-read

135   mapping analysis (**Fig. 2B**, **Fig. S1C**). One of these SVs was detected from *Gemmiger*

qucibialis [within class Clostridia] in the context of an FMT trio (donor: D1; donor 5 years: D1.5y;

and recipient post-FMT: R1-postFMT; **Fig. 2C**). Specifically, this SV has two alternative

genotypes (g1 and g2) which demonstrated different patterns when colonizing donor and

recipient. While g1 is the dominant genotype in the *G. qucibialis* strain in the FMT donor sample

140   (very stable over a five-year time window, **Table S3**), g2 becomes the dominant genotype in the

recipient one year after FMT (**Fig. 2C**, **Methods**), suggesting differing host selective pressures

among the FMT donor and recipient. [See the IGV screenshot showing the ePV results from the

same strain (**Fig. S2**). Additionally, our other long-term manuscript provides detailed methods

for tracking strains, eliminating confounding factors from multiple strains, and validating pure

145   strains through culture analysis using matched fecal samples.[52]].

Because PacBio sequencing data allows direct detection of 6mA and 4mC, *de novo*

methylation motif and quantitative methylation analysis at single molecule resolution

(**Methods**),[28,53,54] using meta-epigenomic analysis, we were able to directly assess ePVs

150   mediated by the SVs. We focused on nine fecal samples (from three infants, ages 1-3) enrolled

in the Early Childhood Antibiotics and the Microbiome (ECAM) study.[3] The three infants were

selected based on their exposure to AMX treatment, with fecal samples available before and

after administration of the antibiotics, which had substantial impact on microbiome composition

in each case (**Fig. 2D-2F**). To illustrate how the SV of an MTase and specificity unit mediated

155   genome-wide ePV, we describe one example detected from a *Bacteroides xylanisolvens* strain

in Infant 1. Two pairs of IRs in a Type I restriction modification system (RMS) create four

possible genotypes via genome rearrangement, with four expected methylation recognition

motifs (**Fig. 2G**). For these four genotypes, *de novo* methylation analysis resolved specific

methylation motifs, and ePVs estimated from methylation analysis correlate with the genotype

frequencies estimated from long-read mapping (**Fig. 2G**). Across the three infants, we observed

bacterial strains whose methylation motifs were stable over time (**Fig. 2H-2J**), as well as those

with genome-wide ePV in at least one methylation motif following exposure to AMX (**Fig. 2H-**

2J), which suggest possible antibiotics-induced selective pressure of genome-wide ePVs.

Although long-read sequencing is very helpful for metagenomic analysis, most existing

microbiome studies were based on short-read sequencing platforms. Short-read DNA

sequencing have limited ability to resolve complex SVs, they are reliable for detecting relatively

simpler SVs (e.g. with only two alternative genotypes). Building on PhaseFinder,[20] we adapted a

pipeline for detecting putative ePVs from short-read metagenomic shotgun sequencing data

(**Fig. 3A-3B; Methods**) and applied it to 2,345 unique metagenomic samples from nine studies

including healthy people (e.g. the Human Microbiome Project[55]) and additional cohorts (**Table**

S4). Across 1,308 unique species, we detected 2,556 putative genome-wide ePVs with at least

one alternative genotype based on SVs in MTases or specificity units (**Fig. 3C**). Four orders

(Clostridiales, Bacteroidales, Enterobacterales, Bifidobacteriales) were enriched for ePVs (**Fig.**

3C; Fig. S3A-S3B), although we observed broad prevalence of potential ePV (IRs in MTase or

specificity units, with neighboring invertase) across many other orders (**Fig. 3C**). Among the

detected putative ePVs, some are associated with the use of antibiotics. For example, a *B.*

uniformis strain from an infant cohort[56] has a putative ePV with two alternative MTase

genotypes, which had clear changes upon using AMX, across four infants (**Fig. 3A, 3D**). The

MTase gene and SV involved in this ePV was detected in 23 of 153 *B. uniforms* isolates

available at NCBI, with each of these 23 strains exhibiting one of the two ePV genotypes (**Fig.**

3E).

Site-specific epigenetic phase variation in the gut microbiome

185   Differing from the genome-wide ePVs resulting from changes in MTase activity or

specificity, a site-specific ePV involves only one or several methylation sites (up to 75) and is

usually driven by the site-specific binding competition between a DNA MTase and another DNA

binding protein such as a TF (**Fig. 4A,**). For a specific sequence motif site recognized by a DNA

MTase in a bacterial strain, only some bacterial cells are methylated while the remainder are not

190   (**Fig 4A**, left panel), and the methylation percentage can be estimated for each site (**Fig 4A**,

right panel). We built on previous work for PacBio DNA sequencing-based single-molecule

methylation detection,[28,57-59] now with enhanced quality control to ensure site-specific ePV

reflects variation within a single strain rather than a mixture of different strains (**Methods**).

Independent validation confirmed that the pipeline can reliably quantify the extent of partial

195   methylation (**Fig. S4A**).

        We applied this analysis pipeline to the nine infant fecal samples and the FMT trio

described above, and an additional sample for each infant, discovering. 885 site-specific ePVs

with < 50% site-specific methylation across 32 strains (representing 27 unique species),

200   including some motif sites that were mostly (<10%) unmethylated (boxplot in **Fig. 4B**). The

number of these partial methylated sites per genome differs across various orders (line plot

highlighted in red, **Fig. 4B**). The partial methylation sites were classified as either intergenic or

intra-genic based on their positions relative to protein-coding genes. Of the 885 site-specific

ePVs, 174 (19.7%) are in intergenic regions (**Fig. S4B**). This significant enrichment of partially

205   methylated sites in intergenic regions (Chi-square $p < 10e-5$) is consistent with the proposed

mechanism of site-specific competitive binding between MTases and transcription factors (or

repressors),[43,44] although intragenic DNA methylation might also be associated with the

regulation of gene expression in bacteria. A site-specific ePV from a CAG-41 sp900066215

strain, is within the promoter region of a gene encoding an alkaline shock protein (Asp23) (**Fig.**

210   **S4C**). In summary, genes with site-specific ePVs are enriched for functional categories related

to metabolic pathways, cellular processes, and environmental processing including several genes associated with host interactions, antibiotic resistance, and biofilm formation (**Fig. 4C**).

To determine whether site-specific ePVs are associated with the selective pressure of environmental factors, we examined 12 ECAM cohort fecal samples from infants with AMX exposures as described. We detected ePV sites with >40% differences in their methylation status between two consecutive sampling time points. In Infant 3 (**Fig. 4D**), we found three site-specific ePVs in *Bacteroides ovatus* (RMS were annotated in **Fig. 4D and Fig. S4D**) whose methylation levels significantly increased upon the infant's treatment with AMX (**Fig. 4E-4F**). These include the essential sigma factor σ^{70}, which regulates genes involved in basic cellular processes including protein synthesis, energy production, and cell wall biosynthesis.[60] The susC/susD family outer membrane proteins, which serve as phase-variable sites, have been identified in various *Bacteroides* species such as *B. fragilis* and *B. thetaiotaomicron*.[61,62] Similarly, the *phoB* gene, which encodes a response regulator protein essential for the PhoBR two-component signal transduction system, exhibits site-specific ePVs. This system regulates the expression of multiple genes under phosphate-limiting conditions[63] and plays a critical role in biofilm formation and environmental stress response.[64] The presence of site-specific ePVs in these genes likely contributes to transcriptional heterogeneity, providing additional adaptive capacity under AMX exposure. Similarly, we examined the FMT fecal samples as described above. Between donor and post-FMT recipient samples **(Fig. 4G)**, we detected ePV sites with differentially methylation ≥ 50% (**Fig. 4H**). In donor 1, we found four site-specific ePVs in *Bacteroides plebeius* whose methylation levels had major decreases 5 years after FMT (**Fig. 4H-4I, Fig. S4E**). One of the four sites is located in the *MBL* gene encoding a β-lactamase, which are emerging as the most notable resistance determinants in Enterobacteriaceae[65,66], while another site is located between *susD* and *TonB*, both were reported to be phase variable sites in several *Bacteroides* species.[52,61,62]

To determine whether site-specific ePVs may be affected by selective pressure in the gut environment differing from *in vitro* culturse, we fed a commercially available probiotic capsule [Glucose Control, Pendulum] with five strains (*A. muciniphila*, *Bifidobacterium longum*, *Clostridium beijerinckii*, *Clostridium butyricum*, and *Anaerobutyricum hallii*) to Germ-free B6 mice and collected fecal samples at day 7 (**Fig. 4J**). PacBio metagenomic sequencing data showed that *A. muciniphila* stably colonized all four tested mice (**Fig. S4F**). Site-specific methylation analysis found 38 ePVs (across 15 unique RA6mATTY sites) in the probiotic *A. muciniphila* strain with >50% differential methylation between the inoculum and the fecal samples (**Fig. 4J-4L; Fig. S5A**); Four of the 16 genes flanking the 15 unique RA6mATTY sites exhibited fitness cost based on INSeq data,[67] including genes related to bacteria's interaction with the host environment(*pmp10*; **Fig 4L**)[68,69] and stress responses (*prmA, PAP*). Taking *prmA* as an example, we found it flanked by a pair of ePV sites (validated by restriction digestion assay; **Fig. S5B**), for which inversion would enhance the methylation level of the promoter (**Fig. 4L**). The orientations of these site-specific ePVs were associated with the successful engraftment of the probiotic *A. muciniphila* strain in the gut environment.

Genes, functions and selective advantage mediated by a genome-wide ePV in an *A. muciniphila* strain

We next focused on a specific strain of *A. muciniphila,* a promising next-generation probiotic associated with several positive health outcomes and drug responses[70-74] to identify the genes, functions, and selective advantages that are mediated by a genome-wide ePV. This species was selected for its prevalent genome-wide and site-specific ePVs (**Fig. 2I, 4K-4L**) and its significant response to AMX exposure observed in an infant fecal sample (**Fig. 5A**). This

strain encodes a Type I RMS with three nested IRs, creating eight different genotypes (g1-g8) and six unique methylation sequence recognition motifs (**Fig. 5A**). Upon the use of AMX, the subpopulation of g2 increased dramatically from 11.1% to 26.5% (**Fig. 5A; Methods**).

265 Consequently, there was a genome-wide ePV in methylation analysis, specifically the genome-wide loss of 6mA at GG6mAN$_9$TGAA sites and genome-wide gain of 6mA at G6mACN$_7$RTTC sites (**Fig. S5C**).

This finding suggests that that ePV may enhance the adaption of *A. muciniphila* to
270 antibiotic exposure. To address that question, we first isolated the *A. muciniphila* strain (**Methods**) and selected a clone (hereafter referred to as Akk_ECAM) that is dominated by the g1 genotype representing the same ePV state as in the infant's fecal sample before AMX exposure (**Fig. S5D**). After ten passages of sub-cultures in liquid media with or without AMX (**Fig. 5B; Fig. S5E; Methods**). Consistent with the observations made from the infant's fecal
275 samples, with the passages, we observed a gradual population-level change from g1 to g2 ePV genotype (**Fig. 5C**) along with genome-wide methylation changes from GG6mAN$_9$TGAA to G6mACN$_7$RTTC (**Fig. 5D**). However, in two different clones, we did not observe the same changes in genome-wide ePV suggesting that *A. muciniphila* adapt AMX through multiple mechanisms.[75] Because single SNP have been associated with commensal fitness during
280 inflammation,[76] we characterized this ePV in a large number (n=46) of g1 clones (**Fig. 5E**). Consistent with our initial observations, there was variability among these 46 clones in terms of how the ePV state changed during the nine passages in presence of AMX: three changed to g2, one changed to g3 and seven changed to g4 (**Fig. 5F**); with the remaining clones continued as g1. With the initial abundance of the non-g1 ePV state < 0.1% at the beginning of the passage
285 that 11 of 46 clones had changed in their ePV states is statistically significant (Fisher's exact test $p = 0.0037$; **Fig. 5G**). In contrast, with control media (without AMX), all the 46 clones remained at g1 after the passages. This difference provide evidence that this genome-wide ePV

may enhance the tolerance of the *A. muciniphila* strain to AMX. Because g3 preserves the same methylation motif as g1 (GG6mAN$_9$TGAA) while g2/g4 both lose the methylation motif (**Fig. 5A**), we hypothesize that the loss of GG6mAN$_9$TGAA methylation in g2 and g4 influence the expression of *A. muciniphila* genes that increases its tolerance to AMX.

To test this hypothesis, we selected totally 11 clones with preserved (g1, g3) or lost (g2, g4) GG6mAN$_9$TGAA methylation and profiled their transcripts by RNA-seq after the final two passages in the presence of AMX (**Fig. S6A**). The RNA-seq data analysis showed that the *A. muciniphila* transcriptional signatures with two ePV states cluster separately in a principal component analysis (**Fig. 5H**). Differential gene expression analysis identified a single gene [Amuc_0914 homolog, herein named *mucC*, for *muciniphila* gene from cupin family] that is significantly (negative binomial test, $p = 3.76E-07$) and consistently up-regulated when GG6mAN$_9$TGAA methylation is lost in g2 and g4 clones (**Fig. 5I, Fig. S6B**). Next, we asked whether its expression was methylation-specific; we found the promoter of *mucC* contains a GG6mN$_9$TGAA methylation site between its -35 region and translation initiation site (**Fig. 5J**). We also performed whole genome shotgun sequencing on colonies that had changes in ePV states with AMX treatment and found certain genetic variations across the clones but none associated with the AMX treatment (**Fig. S6C-S6D**). These finds identify methylation-mediated regulation of *mucC* expression.

The *mucC* gene is highly conserved across *A. muciniphila* strains (**Fig. S6E, Table S5**) encoding a small protein characterized by a cupin fold,[77] which consists of a conserved β-barrel structure (AlphaFold prediction,[78,79] **Fig. S6F**). MucC homologs are also found in other bacterial species (**Fig. S6G**). In *A. muciniphila*, *mucC* is immediately adjacent to Ribonuclease R (rnr), a member of the RNase II superfamily (**Fig. S6H**) that hydrolyze RNA in the 3' - 5' direction.

RNase R is associated with selective mRNA degradation, particularly of mRNAs that lack a stop codon in bacteria[80] and are expressed under stress conditions.[81] Because the genetic tools of *A. muciniphila* are limited[67], we expressed *mucC in trans* in *E. coli* cells (pZE21- *mucC*) and confirmed its protein expression by mass spectrometry (**Methods; Fig. S7A-S7D**). By comparing *mucC* + *E. coli* (pZE21- *mucC*) with negative controls (empty plasmid, pZE21), *mucC* expression conferred significantly greater tolerance to AMX based on their MICs and survival rate (**Fig.5K**).

Finally, because ePV enhances non-genetic clonal heterogeneity and empowers the adaptive capacity of single cells in a population,[38-42] we hypothesized that while the ePV state with high *mucC* expression enhances tolerance to AMX, other conditions may favor an ePV state with low *mucC* expression. High basal level of *mucC* expression doesn't create a fitness cost *in vitro*, as we found that the removal of AMX didn't select for the g1 ePV state with lower *mucC* expression (**Fig. S7E**). However, after testing additional stressors, we discovered that ciprofloxacin selects the g1 ePV state with lower *mucC* expression (**Fig. 5L, Table S7**), demonstrating dynamic adaptation depending on a local stress.

Overall, our findings indicate that a complex ePV detected in an *A. muciniphila* strain that originated from an infant gut microbiome enhances clonal heterogeneity via regulating *mucC* gene expression with several ePV states associated with high *mucC* expression selected by AMX and low *mucC* expression selected by ciprofloxacin. The reversible selection by two different antibiotics highlights the advantage of ePV to empower the same *A. muciniphila* strain to epigenetically adapt to different stress conditions. Noteworthy, this evolutionary strategy, where a small subpopulation generates ePV heterogeneity sufficient to increase population-

level adaptation across various stress conditions, underscores the adaptive advantage at the population level rather than individual cells.

Discussion

Our study highlights the prevalence of ePV across diverse bacterial species within the gut microbiota, underscoring the role of epigenetic mechanisms in enhancing clonal heterogeneity and enabling strains to adapt to environmental stressors such as antibiotic

345   treatments or when colonizing new hosts.

We leverage long-read metagenomic data to simultaneously detect complex SVs and DNA methylation, providing a detailed view of the epigenetic landscape of bacteria within gastrointestinal tract. Furthermore, our analysis of an extensive collection of short-read

350   metagenomic data has highlighted SVs near MTase and specificity units, suggesting the broad presence of ePV in healthy and diseased human cohorts. As long-read sequencing is more often used in microbiome studies, we anticipate that our findings will open new dimensions to characterize how gut bacteria adapt to various stress conditions and host interactions using epigenetic mechanisms. The genes associated with ePV in our datasets suggest that the

355   biological consequences of ePV are broad, impacting microbe-host interactions, antibiotic resistance, and the metabolic capacities of gut bacteria **(Figs. 2-4)**.

Compared to genetic phase variations, ePV offers several advantages in enhancing clonal heterogeneity. The reversibility of ePV, without changing DNA sequence enables

360   individual bacterial strains to adapt dynamically to diverse stresses. Furthermore, genome-wide

ePV can simultaneously regulate multiple genes, which provides coordinated activation or inhibition of multiple functionally related genes.[29,32] As shown in bacterial human pathogens, regulation by global epigenetic modification highlights an intermediate state toward long-term adaptation to stress conditions and interactions with the human immune system.[75,82,83] The

365   diverse functions of the genes associated with genome-wide and site-specific ePVs suggest that ePVs may regulate many important functions (e.g. antibiotic resistance, bacteria-host interaction, fitness, etc) of gut bacteria to enhance their adaptation and resilience in the changing environment.

370   The ePV characterized in the *A. muciniphila* strain regulated differential expression of the *mucC* gene, upon exposure to two common antibiotics, AMX and ciprofloxacin **(Fig. 5)**, is a great example of a new epigenetic bet-hedging mechanism of commensal bacteria. Our results indicate that in the human gut, ePVs may help bacterial populations regain heterogeneity after bottlenecks encountered during colonization of a new host or severe perturbations due to

375   antibiotic exposures. These ePV-driven regulatory mechanisms underscore new opportunities for targeted epigenetic interventions to improve desired functions of beneficial bacteria. For example, by manipulating ePV, we may strategically enhance the resilience and functional capabilities of beneficial bacteria, which might improve the success rates of probiotic engraftment and the efficacy of treatments for microbiota-associated conditions.

        Despite strong evidence that the regulation of *mucC* expression by ePV enhances tolerance of *A. muciniphila* to AMX, we noted that this is not the only mechanism to adapt to antibiotic stress. As reported in other studies, genetic mutations can also confer increased fitness in other bacterial species. For instance, a subpopulation shift in specific clones has been

385   reported in *B. thetaiotaomicron*, which was associated with resistance to oxidative stress

through alterations in the sequence of the IctA protein.[76] While implicating different pathways, our finding highlights that ePV is one way to enhance clonal heterogeneity and should be considered within a systemic view that includes other genetic, transcriptional, and translational forms of regulations.[15,20,21,55]

Figure captions

Figure 1. An integrated framework for epigenetic phase variation (ePV) detection and function validation from the human gut microbiome

395 **(A)** A specific bacterial strain may have sub-populations that exhibit heterogeneous methylation patterns, leading to distinct gene expression and phenotypes. This may enhance the ability of a single strain to adapt to different stress conditions. **(B)** Schematic diagram of genome-wide ePV events, mediated by the structural variation located within restriction-modification system (RMS) or methyltransferase (MTase) genes. **(C)** Schematic of site-specific ePV events, resulting from

400 the competition between MTase and transcription factors (TFs). **(D)** Prevalence and phylogenetic distribution of ePV were explored across various healthy and clinical-related cohorts. Phylogenetic tree is color-coded at phylum level. **(E)** Workflow for functional analysis of genes involved in ePV events. Bacterial strains carrying the ePVs were isolated from fecal sample. An *in vitro* passage assay was performed under conditions mimicking the *in vivo*

405 environment, followed by whole-genome shotgun and targeted DNA sequencing. RMS genotypes were quantified and associated with treatment conditions. RNA sequencing was conducted to profile transcriptomic differences between genotypes. Differentially expressed genes (DEGs) were overexpressed further in native strains or model species, and phenotypic tests were carried out to confirm whether the altered phenotype was due to the expression of

410 the gene of interest (GOI).

Figure 2. Long-read metagenomic sequencing deciphers complex ePV events

(**A**) An ePV observed in a *Phocaecola vulgatus* strain from a healthy adult gut microbiome, mediated by three paired/matching inverted repeats (IRs), resulting in three distinct RMS

415 genotypes (g1-g3). The alignment plots show raw reads aligned with their corresponding reference genomes. The proportion of each genotype is displayed on the right side of the read alignment chart. (**B**) Additional ePV events detected in various bacterial species. Gene diagrams illustrate the rearrangement of RMS gene fragments, while pie charts show the

proportion of each RMS variant. (**C**) ePV events identified in an FMT cohort. The top panel illustrates the FMT sample collection from Recipient 1 (see **Table S3**). The bottom panel shows an ePV in a *Gemmiger qucibialis* strain, detected in both donor and post-FMT samples. (**D-F**) Metagenomes analysis of fecal samples from three infants in the ECAM cohort. The top panel presents the timeline of sample collection and amoxicillin (AMX) exposure. The bottom panel displays species-level relative abundance profiles, including only species with a sequencing depth > 10X. (**G**) An ePV event observed in a *Bacteroides xylanisolvens* strain, mediated by two pair of IR features, resulting in four RMS genotypes (g1-g4). RMS genotype was profiled in pre- and post-AMX treatment samples, alongside the methylome (m1-m4). AMX treatment inhibited the population of g2 (highlighted in purple) while promoting the growth of g1 (highlighted grey). Simultaneously, the methylome shift from m2 to m1, mirroring the genotype changes. (**H-J**) Additional methylation changes and ePV events in different bacterial species in Infant 1 (**H**), Infant 2 (**I**) and Infant 3 (**J**). The methylation profiles were generated based on the single-site methylation analysis (see Methods) and manually curated. (**A-C, G**) M: MTase represented by green arrows; S1, specific subunit 1 represented by red arrows; S2, specific subunit 2 represented by yellow arrows. The pink and black rectangles above the gene schematic diagram are IRs.

Figure 3. Genome-wide ePV events are widespread across different cohorts and associated with clinical treatments

(**A**) An ePV observed in a *Bacteroides uniforms* strain from an adult gut microbiome. Due to read length limitation, short-read data can only detect simple ePV (or "inverton", as defined in a previous study), which are mediated by a pair of IR features and result in two distinct RMS genotypes (g1, g2). The proportion of each genotype is shown to the right of the gene diagram. The read alignment chart illustrates how paired-end (PE) reads were used to detect the inverton-like ePV. (**B**) Additional inverton-like ePVs detected across various bacterial species. Gene diagrams show the rearrangement of RMS gene fragments, while pie charts depict the proportion of each RMS variant. (**C**) Phylogenetic tree of inverton-like ePVs. Potential ePV (based solely on the reference genome) and confirmed ePV (supported by both the reference genome and metagenomic read) are grouped into three categories based on the RMS genotype (**Fig. S3**) and are projected onto the phylogenetic tree. ePV enriched orders are highlighted according to the colors scheme in the legend. (**D**) Response of *B. uniforms* ePV to AMX treatment. The *B. uniforms* ePV mentioned in **Fig. 3A** was detected in four infants from the DIABIMMUNE cohort, both pre- and post-AMX treatment. The RMS genotype shifted from g1 to g2 in response to treatment. (**E**) Prevalence analysis of the aforementioned *B. uniforms* ePV.

Among 153 complete *B. uniforms* genomes from NCBI, 23 carry this ePV. The bar plot on the right shows the genotype distribution across each isolate. **(A, B, E)** M: methyltransferase represented by green arrows; S1, specific subunit 1 represented by red arrows; S2, specific subunit 2 represented by yellow arrows. The pink rectangles above the gene schematic are IRs.

Figure 4. Site-specific ePV events are widespread across different cohorts and associated with bacterial adaptation to environmental stress

(**A**) For a specific bacterial strain, only a subset of cells are methylated, while the remainder are not (left panel). The schematic in the middle panel depicts a transcription factor (TF) competing with an MTase at a specific site. The ePV site (highlighted within the dashed rectangle) was identified from multiple molecules using single-molecule methylation analysis. The right panel illustrates the deconvolution results obtained from this single-site methylation analysis. (**B**) Summary of all partial methylation sites detected in this study. Values were normalized based on the strain number within the order (diamond points highlighted in blue). Methylation fraction values within the order were also determined (box plot). (**C**) Function categories of the genes affected by ePV sites. Heatmap showing the number of genes regulated by ePV, with host strain taxonomy (order) as column label and gene function category (pathway) as row labels. The color scale represents the number of genes, indicating the occurrence of each functional category per strain. (**D**) The top panel presents the time line of AMX administration and sample collection for Infant 3. The bottom panel shows the involved methylation-related genes in a *B. ovatus* strain from Infant 3. (**E**) Distribution of partially methylated G6mATC sites across the *B. ovatus* genome. The genome consists of three contigs, randomly ordered along the x axis. Three of the partially methylated sites were labeled with numbers. (**F**) Changes in partially methylated G6mATC sites in response to AMX treatment. The motif sites were examined on both strands, and only sites that showed partial methylation on both strands were considered final partially methylated sites (the same criteria were applied to all other partially methylated sites in Fig. **4I,L**). The gene diagram shows the location of each site within the affected gene. **(E, F)** Pre- (black) and post-AMX (red) methylations were highlighted. (**G**) Sample collection information for Recipient 2 from FMT cohort. (**H**) Distribution of partially methylated G6mATC sites across the *Bacteroides plebeius* genome. Four of the partially methylated sites were labeled with numbers. (**I**) Change in partially methylated G6mATC sites in response to FMT. The gene diagram shows the location of each site within the affected gene. **(H,I)** Pre- (black) and post-FMT (red) methylations were highlighted. (**J**) Design and sample collection information for Akk-fed mouse assay. (**K**) Distribution of partial methylated RA6mATTY sites across the *A.*

muciniphila genome. Three of the partially methylated sites were labeled with numbers. (**L**) Changes in partially methylated RA6mATTY sites from capsule to *in vivo* colonization (mouse stool). The gene diagram shows the location of each site within the affected gene. **(K, L)** Capsule (black) and *in vivo* (red) methylations were highlighted.

Figure 5. Characterization of a complex genome-wide ePV and its role in *Akkermansia* tolerance to AMX

(**A**) An ePV event observed in an *A. muciniphila* strain from Infant 2 in the ECAM cohort, mediated by three pairs of IR features, resulting in eight RMS genotypes (g1-g8). RMS genotype was profiled in pre- and post-AMX treatment samples, along with the methylation motifs. AMX treatment inhibited the population of g1 (highlighted in grey) while promoting the growth of g2 (highlighted in purple) or other non g1 populations. (**B**) Experimental design of the *in vitro* passage assay. *A. muciniphila* was isolated from the fecal sample of Infant 2 (hereafter referred to as Akk_ECAM). After recovery on the BHI agar plates, a culture from single colony was subjected to 2 mg / L AMX treatment for 10 passages. Cultures without AMX served as negative controls. (**C**) RMS genotype profiling using targeted sequencing. In response to AMX treatment, the RMS genotype shifted from g1 dominance to g2 dominance, whereas g1 remained dominant through passage 10 in the absence of AMX. (**D**) Methylation motif shift in the Akk_ECAM strain. Genomic DNA was extracted from colony 3 of the first batch *in vitro* passage assay. Whole genome shotgun sequencing was performed using Pacbio Sequel II platform. Following single-molecule methylation analysis, the methylation level of the two motifs of interest was investigated. This is consistent with the *in vivo* findings, where AMX treatment decreased in the g1 methylation subpopulation (highly methylated at GG6mAN$_7$TGAA) and increased the non-g1 methylation subpopulation (highly methylated at G6mACN$_7$RTTC). **(E)** Experimental design of the second batch of the *in vitro* passage assay. After recovery on BHI agar plates, 46 colonies were subjected to 2 mg / L AMX treatment for nine passages. Culture without AMX served as a negative control. (**F**) RMS genotype profiling with targeted sequencing. The RMS genotype shifted from g1 dominance to non-g1 (g2-g4) dominance in response to AMX treatment, while g1 remained dominant through passage 9 in the absence of AMX treatment. (**G**) Summary of the 2nd batch *in vitro* passage assay. Genotypes were determined by targeted sequencing and grouped into two categories, g1 and non-g1 (g2-g4). Fisher's exact test was used to assess the statistical significance of the difference in proportions between pre- and post-AMX treatment. The results indicate a significant association ($p = 0.0037$), suggesting that g2-g4 genotypes were selected by AMX treatment. (**H**) PCA analysis of the RNA-seq samples. Under AMX treatment, g2/g4 samples were clearly separated from g1/g3

samples. (**I**) DEGs were identified by comparing g1/g3 and g2/g4 samples under AMX treatment. Thresholds for DEGs were $\log_2$ fold changes > 1 or < -1, with an adjusted p value < 0.05. *mucC* was the only upregulated gene (highlighted in red). *tRNA-Phe* was the only downregulated gene (highlighted in green). (**J**) The promoter elements and methylation motif (GG6mAN$_9$TGAA) were represented in the diagram. The fraction of g1 methylation motif in the promoter decreased with AMX treatment. (**K**) AMX tolerance of *E. coli* and overexpression of the *mucC* gene. The *mucC* gene was codon-optimized, synthesized and fused with a 6X HIS tag at the N-terminus. It was inserted into a medium-copy plasmid with a constitutive promoter, and AMX tolerance was assessed by using the ETEST and CFU counting methods respectively. Each strain was tested in triplicate. (**L**) RMS reversibility assay. g2/g4 dominant strains from the second batch of the *in vitro* assay were treated under various conditions. The point plot shows the targeted sequencing results, and the zoomed-in bar plot indicates that ciprofloxacin treatment can reverse non-g1 (g2/g4) genotypes back to g1 genotype. Each condition was tested in two replicates.

Data availability

All sequencing data generated in this study have been submitted to NCBI SRA with the accession number PRJNA1207694, and will be made publicly available upon publication.

Acknowledgments

We thank the Icahn School of Medicine at Mount Sinai colleagues Jeremiah J. Faith and Ilaria Mogno for helping with the characterization of the adult FMT fecal samples. This work was supported by grant no. R35 GM139655 (G.F.) from the National Institutes of Health and seed fund (G.F.) from the Icahn School of Medicine at Mount Sinai. L.E.D. lab is supported by grant no. RGPIN-2022-03915. This work was also supported in part through the computational resources and staff expertise provided by the Department of Scientific Computing at the Icahn School of Medicine at Mount Sinai.

Author contributions

M.N. and G.F. designed the methods. M.N. performed most of the computational analysis. Y.F., M.K. and A.T. assisted with computational analysis. M.N., Y.Liu, Y. Li, W.Q., L.E.D., X-S.Z, M.K. and E.A.M. performed experiments. M.N., Y.F., Y.Liu, Y. Li, W.Q., L.E.D., X-S.Z, M.K., E.A.M., W.J., M.J.B., R.H.V. and G.F. contributed to data interpretation. M.N. and G.F. wrote the manuscript with input and comments from all coauthors. G.F. conceived and supervised the project.

Declaration of interests

R.H.V. is a co-founder of Bloom Science (San Diego, CA)

References

1. Nandi, T., Lee, I.R., Ghosh, T., Ng, A.H.Q., Chng, K.R., Li, C., Tan, Y.H., Lye, D., Barkham, T., Chen, S., et al. (2018). Gut microbiome recovery after antibiotic usage is mediated by specific bacterial species. bioRxiv. 10.1101/350470.
2. Cox, Laura M., Yamanishi, S., Sohn, J., Alekseyenko, Alexander V., Leung, Jacqueline M., Cho, I., Kim, Sungheon G., Li, H., Gao, Z., Mahana, D., et al. (2014). Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. Cell *158*, 705-721. 10.1016/j.cell.2014.05.052.
3. Bokulich, N.A., Chung, J., Battaglia, T., Henderson, N., Jay, M., Li, H., D. Lieber, A., Wu, F., Perez-Perez, G.I., Chen, Y., et al. (2016). Antibiotics, birth mode, and diet shape microbiome maturation during early life. Sci. Transl. Med. *8*, 343ra382-343ra382. 10.1126/scitranslmed.aad7121.
4. Wirbel, J., Pyl, P.T., Kartal, E., Zych, K., Kashani, A., Milanese, A., Fleck, J.S., Voigt, A.Y., Palleja, A., Ponnudurai, R., et al. (2019). Meta-analysis of fecal metagenomes reveals global microbial signatures that are specific for colorectal cancer. Nat. Med. *25*, 679-689. 10.1038/s41591-019-0406-6.
5. Franzosa, E.A., Sirota-Madi, A., Avila-Pacheco, J., Fornelos, N., Haiser, H.J., Reinker, S., Vatanen, T., Hall, A.B., Mallick, H., McIver, L.J., et al. (2019). Gut microbiome structure and metabolic activity in inflammatory bowel disease. Nat. Microbiol. *4*, 293-305. 10.1038/s41564-018-0306-4.
6. Nagy-Szakal, D., and Kellermayer, R. (2011). The remarkable capacity for gut microbial and host interactions. Gut Microbes *2*, 178-182. 10.4161/gmic.2.3.16107.
7. Demir, M., Lang, S., Hartmann, P., Duan, Y., Martin, A., Miyamoto, Y., Bondareva, M., Zhang, X., Wang, Y., Kasper, P., et al. (2022). The fecal mycobiome in non-alcoholic fatty liver disease. J. Hepatol. *76*, 788-799. 10.1016/j.jhep.2021.11.029.

8. Cammann, D., Lu, Y.M., Cummings, M.J., Zhang, M.L., Cue, J.M., Do, J., Ebersole, J., Chen, X.N., Oh, E.C., Cummings, J.L., and Chen, J.C. (2023). Genetic correlations between Alzheimer's disease and gut microbiome genera. Sci. Rep. *13*, 5258. 10.1038/s41598-023-31730-5.

9. Ferreiro, A.L., Choi, J., Ryou, J., Newcomer, E.P., Thompson, R., Bollinger, R.M., Hall-Moore, C., Ndao, I.M., Sax, L., Benzinger, T.L.S., et al. (2023). Gut microbiome composition may be an indicator of preclinical Alzheimer's disease. Sci. Transl. Med. *15*, eabo2984. 10.1126/scitranslmed.abo2984.

10. Smith, M., Dai, A.Q., Ghilardi, G., Amelsberg, K.V., Devlin, S.M., Pajarillo, R., Slingerland, J.B., Beghi, S., Herrera, P.S., Giardina, P., et al. (2022). Gut microbiome correlates of response and toxicity following anti-CD19 CAR T cell therapy. Nat. Med. *28*, 713-+. 10.1038/s41591-022-01702-9.

11. Valles-Colomer, M., Blanco-Miguez, A., Manghi, P., Asnicar, F., Dubois, L., Golzato, D., Pinto, F., Cumbo, F., Huang, K.D., Manara, S., et al. (2023). The person-to-person transmission landscape of the gut and oral microbiomes. Nature *614*, 125-135. 10.1038/s41586-022-05620-1.

12. Zhu, X., Qin, J.Y., Tan, C.Y., and Ning, K. (2021). The seasonal changes of the gut microbiome of the population living in traditional lifestyles are represented by characteristic species-level and functional-level SNP enrichment patterns. BMC Genomics *22*, 83. 10.1186/s12864-021-07372-0.

13. Ma, C.C., Wasti, S., Huang, S., Zhang, Z., Mishra, R., Jiang, S.M., You, Z.K., Wu, Y.X., Chang, H.B., Wang, Y.Y., et al. (2020). The gut microbiome stability is altered by probiotic ingestion and improved by the continuous supplementation of galactooligosaccharide. Gut Microbes *12*. 10.1080/19490976.2020.1785252.

14. Zahavi, L., Lavon, A., Reicher, L., Shoer, S., Godneva, A., Leviatan, S., Rein, M., Weissbrod, O., Weinberger, A., and Segal, E. (2023). Bacterial SNPs in the human gut microbiome associate with host BMI. Nat. Med. 10.1038/s41591-023-02599-8.

15. Greenblum, S., Carr, R., and Borenstein, E. (2015). Extensive strain-level copy-number variation across human gut microbiome species. Cell *160*, 583-594. 10.1016/j.cell.2014.12.038.

16. Zeevi, D., Korem, T., Godneva, A., Bar, N., Kurilshikov, A., Lotan-Pompan, M., Weinberger, A., Fu, J., Wijmenga, C., Zhernakova, A., and Segal, E. (2019). Structural variation in the gut microbiome associates with host health. Nature *568*, 43-48. 10.1038/s41586-019-1065-y.

17. Zhernakova, D.V., Wang, D.M., Liu, L., Andreu-Sánchez, S., Zhang, Y., Ruiz-Moreno, A.J., Peng, H.R., Plomp, N., Del Castillo-Izquierdo, A., Gacesa, R., et al. (2024). Host genetic regulation of human gut microbial structural variation. Nature *625*. 10.1038/s41586-023-06893-w.

18. Xu, J., Mahowald, M.A., Ley, R.E., Lozupone, C.A., Hamady, M., Martens, E.C., Henrissat, B., Coutinho, P.M., Minx, P., Latreille, P., et al. (2007). Evolution of symbiotic bacteria in the distal human intestine. PLoS Biol. *5*, 1574-1586, e156. 10.1371/journal.pbio.0050156.

19. Levan, S.R., Stamnes, K.A., Lin, D.L., Panzer, A.R., Fukui, E., McCauley, K., Fujimura, K.E., McKean, M., Ownby, D.R., Zoratti, E.M., et al. (2019). Elevated faecal 12,13-diHOME concentration in neonates at high risk for asthma is produced by gut bacteria and impedes immune tolerance. Nat. Microbiol. *4*, 1851-1861. 10.1038/s41564-019-0498-2.

20. Jiang, X., Hall, A.B., Arthur, T.D., Plichta, D.R., Covington, C.T., Poyet, M., Crothers, J., Moses, P.L., Tolonen, A.C., Vlamakis, H., et al. (2019). Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host adaptation in the gut. Science *363*, 181-187. 10.1126/science.aau5238.

21. Chanin, R.B., West, P.T., Wirbel, J., Gill, M.O., Green, G.Z.M., Park, R.M., Enright, N., Miklos, A.M., Hickey, A.S., Brooks, E.F., et al. (2024). Intragenic DNA inversions expand bacterial coding capacity. Nature. 10.1038/s41586-024-07970-4.

22. Jonsson, A.B., Nyberg, G., and Normark, S. (1991). Phase variation of gonococcal pili by frameshift mutation in pilC, a novel gene for pilus assembly. EMBO J. *10*, 477-488-488. 10.1002/j.1460-2075.1991.tb07970.x.

23. Segal, E., Billyard, E., So, M., Storzbach, S., and Meyer, T.F. (1985). Role of chromosomal rearrangement in *N. gonorrhoeae* pilus phase variation. Cell *40*, 293-300. 10.1016/0092-8674(85)90143-6.

24. Sánchez-Romero, M.A., and Casadesús, J. (2020). The bacterial epigenome. Nat. Rev. Microbiol. *18*, 7-20. 10.1038/s41579-019-0286-2.

25. Gopalan-Nair, R., Coissac, A., Legrand, L., Lopez-Roques, C., Pécrix, Y., Vandecasteele, C., Bouchez, O., Barlet, X., Lanois, A., Givaudan, A., et al. (2024). Changes in DNA methylation contribute to rapid adaptation in bacterial plant pathogen evolution. PLoS Biol. *22*, e3002792. 10.1371/journal.pbio.3002792.

26. Forni, D., Pozzoli, U., Mozzi, A., Cagliani, R., and Sironi, M. (2024). Depletion of CpG dinucleotides in bacterial genomes may represent an adaptation to high temperatures. NAR genom. bioinform. *6*, lqae088. 10.1093/nargab/lqae088.

27. Atack, J.M., Tan, A., Bakaletz, L.O., Jennings, M.P., and Seib, K.L. (2018). Phasevarions of bacterial pathogens: Methylomics sheds new light on old enemies. Trends Microbiol. *26*, 715-726. 10.1016/j.tim.2018.01.008.

28. Beaulaurier, J., Schadt, E.E., and Fang, G. (2019). Deciphering bacterial epigenomes using modern sequencing technologies. Nat. Rev. Genet. *20*, 157-172. 10.1038/s41576-018-0081-3.

29. Manso, A.S., Chai, M.H., Atack, J.M., Furi, L., De Ste Croix, M., Haigh, R., Trappetti, C., Ogunniyi, A.D., Shewell, L.K., Boitano, M., et al. (2014). A random six-phase switch regulates pneumococcal virulence via global epigenetic changes. Nat. Commun. *5*, 5055. 10.1038/ncomms6055.

30. Oliveira, P.H., Ribis, J.W., Garrett, E.M., Trzilova, D., Kim, A., Sekulovic, O., Mead, E.A., Pak, T., Zhu, S., Deikus, G., et al. (2020). Epigenomic characterization of *Clostridioides difficile* finds a conserved DNA methyltransferase that mediates sporulation and pathogenesis. Nat. Microbiol. *5*, 166-180. 10.1038/s41564-019-0613-4.

31. Nahar, N., Tram, G., Jen, F.E.-C., Phillips, Z.N., Weinert, Lucy A., Bossé, Janine T., Jabbari, Jafar S., Gouil, Q., Du, Mei R.M., Ritchie, Matthew E., et al. (2023). *Actinobacillus pleuropneumoniae* encodes multiple phase-variable DNA methyltransferases that control distinct phasevarions. Nucleic Acids Res. 10.1093/nar/gkad091.

32. Srikhanta, Y.N., Fox, K.L., and Jennings, M.P. (2010). The phasevarion: phase variation of type III DNA methyltransferases controls coordinated switching in multiple genes. Nat. Rev. Microbiol. *8*, 196-206. 10.1038/nrmicro2283.

33. Ben-Assa, N., Coyne, M.J., Fomenkov, A., Livny, J., Robins, William P., Muniesa, M., Carey, V., Carasso, S., Gefen, T., Jofre, J., et al. (2020). Analysis of a phase-variable restriction modification system of the human gut symbiont *Bacteroides fragilis*. Nucleic Acids Res. 10.1093/nar/gkaa824.

34. Huang, X.T., Wang, J.J., Li, J., Liu, Y.N., Liu, X., Li, Z.Y., Kurniyati, K., Deng, Y.J., Wang, G.L., Ralph, J.D., et al. (2020). Prevalence of phase variable epigenetic invertons among host-associated bacteria. Nucleic Acids Res. *48*, 11468-11485. 10.1093/nar/gkaa907.

35. Brunet, Y.R., Bernard, C.S., Gavioli, M., Lloubès, R., and Cascales, E. (2011). An epigenetic switch involving overlapping fur and DNA methylation optimizes expression of a type VI secretion gene cluster. PLoS Genet. *7*, e1002205. 10.1371/journal.pgen.1002205.

36. Lim, H.N., and van Oudenaarden, A. (2007). A multistep epigenetic switch enables the stable inheritance of DNA methylation states. Nat. Genet. *39*, 269-275. 10.1038/ng1956.

37. Jen, F.E.C., Scott, A.L., Tan, A., Seib, K.L., and Jennings, M.P. (2020). Random switching of the ModA11 type III DNA methyltransferase of *Neisseria meningitidis* regulates entner-doudoroff aldolase expression by a methylation change in the *eda* promoter region. J. Mol. Biol. *432*, 5835-5842. 10.1016/j.jmb.2020.08.024.

38. Seib, K.L., Srikhanta, Y.N., Atack, J.M., and Jennings, M.P. (2020). Epigenetic regulation of virulence and immunoevasion by phase-variable restriction-modification systems in bacterial pathogens. In Annual Review of Microbiology, Vol 74, 2020, S. Gottesman, ed. pp. 655-671. 10.1146/annurev-micro-090817-062346.

39. Li, J., Li, J.W., Feng, Z.X., Wang, J.J., An, H.R., Liu, Y.N., Wang, Y., Wang, K.L., Zhang, X.G., Miao, Z., et al. (2016). Epigenetic switch driven by DNA inversions dictates phase variation in *Streptococcus pneumoniae*. PLoS Path. *12*, e1005762. 10.1371/journal.ppat.1005762.

40. De Bolle, X., Bayliss, C.D., Field, D., Van De Ven, T., Saunders, N.J., Hood, D.W., and Moxon, E.R. (2000). The length of a tetranucleotide repeat tract in *Haemophilus influenzae* determines the phase variation rate of a gene with homology to type III DNA methyltransferases. Mol. Microbiol. *35*, 211-222. 10.1046/j.1365-2958.2000.01701.x.

41. de Vries, N., Duinsbergen, D., Kuipers Ernst, J., Pot Raymond, G.J., Wiesenekker, P., Penn Charles, W., van Vliet Arnoud, H.M., Vandenbroucke-Grauls Christina, M.J.E., and Kusters Johannes, G. (2002). Transcriptional phase variation of a type iii restriction-modification system in *Helicobacter pylori*. J. Bacteriol. *184*, 6615-6623. 10.1128/jb.184.23.6615-6624.2002.

42. Li, H.-c., Chen, T., Yu, L., Guo, H.-x., Chen, L., Chen, Y.-h., Chen, M., Zhao, J., Yan, H.-m., Zhou, L., and Wang, W. (2020). Genome-wide DNA methylation and transcriptome and proteome changes in *Mycobacterium tuberculosis* with para-aminosalicylic acid resistance. Chem. Biol. Drug Des. *95*, 104-112. 10.1111/cbdd.13625.

43. Braaten, B.A., Blyn, L.B., Skinner, B.S., and Low, D.A. (1991). Evidence for a methylation-blocking factor (mbf) locus involved in pap pilus expression and phase variation in *Escherichia coli*. J. Bacteriol. *173*, 1789-1800. 10.1128/jb.173.5.1789-1800.1991.

44. Kaminska, R., and van der Woude, M.W. (2010). Establishing and maintaining sequestration of dam target sites for phase variation of *agn43* in *Escherichia coli*. J. Bacteriol. *192*, 1937-1945. 10.1128/jb.01629-09.

45. Ghosh, D., Veeraraghavan, B., Elangovan, R., and Vivekanandan, P. (2020). Antibiotic Resistance and Epigenetics: More to It than Meets the Eye. Antimicrob. Agents Chemother. *64*, 10.1128/aac.02225-02219.

46. Carasso, S., Zaatry, R., Hajjo, H., Kadosh-Kariti, D., Ben-Assa, N., Naddaf, R., Mandelbaum, N., Pressman, S., Chowers, Y., Gefen, T., et al. (2024). Inflammation and bacteriophages affect DNA inversion states and functionality of the gut microbiota. Cell Host Microbe. 10.1016/j.chom.2024.02.003.

47. Won, C., and Yim, S.S. (2024). Emerging methylation-based approaches in microbiome engineering. Biotechnology for Biofuels and Bioproducts *17*, 96. 10.1186/s13068-024-02529-x.

48. English, A.C., Salerno, W.J., and Reid, J.G. (2014). PBHoney: identifying genomic variants via long-read discordance and interrupted mapping. BMC Bioinformatics *15*, 180. 10.1186/1471-2105-15-180.

49. Cretu Stancu, M., van Roosmalen, M.J., Renkens, I., Nieboer, M.M., Middelkamp, S., de Ligt, J., Pregno, G., Giachino, D., Mandrile, G., Espejo Valle-Inclan, J., et al. (2017). Mapping and phasing of structural variation in patient genomes using nanopore sequencing. Nat. Commun. *8*, 1326. 10.1038/s41467-017-01343-4.

50. Sedlazeck, F.J., Rescheneder, P., Smolka, M., Fang, H., Nattestad, M., von Haeseler, A., and Schatz, M.C. (2018). Accurate detection of complex structural variations using single-molecule sequencing. Nat. Methods *15*, 461-468. 10.1038/s41592-018-0001-7.

51. Chen, L., Zhao, N., Cao, J., Liu, X., Xu, J., Ma, Y., Yu, Y., Zhang, X., Zhang, W., Guan, X., et al. (2022). Short- and long-read metagenomics expand individualized structural variations in gut microbiomes. Nat. Commun. *13*, 3175. 10.1038/s41467-022-30857-9.

52. Fan, Y., Ni, M., Aggarwala, V., Mead, E.A., Ksiezarek, M., Kamm, M.A., Borody, T.J., Paramsothy, S., Kaakoush, N.O., Grinspan, A., et al. (2024). Long-read metagenomics empowers precise tracking of bacterial strains and their genomic changes after fecal microbiota transplantation. bioRxiv. 10.1101/2024.09.30.615906.

53. Beaulaurier, J., Zhang, X.-S., Zhu, S., Sebra, R., Rosenbluh, C., Deikus, G., Shen, N., Munera, D., Waldor, M.K., Chess, A., et al. (2015). Single molecule-level detection and long read-based phasing of epigenetic variations in bacterial methylomes. Nat. Commun. *6*, 7438. 10.1038/ncomms8438.

54. Kong, Y., Cao, L., Deikus, G., Fan, Y., Mead, E.A., Lai, W., Zhang, Y., Yong, R., Sebra, R., Wang, H., et al. (2022). Critical assessment of DNA adenine methylation in eukaryotes using quantitative deconvolution. Science *375*, 515-522. 10.1126/science.abe7489.

55. Huttenhower, C., Gevers, D., Knight, R., Abubucker, S., Badger, J.H., Chinwalla, A.T., Creasy, H.H., Earl, A.M., FitzGerald, M.G., Fulton, R.S., et al. (2012). Structure, function and diversity of the healthy human microbiome. Nature *486*, 207-214. 10.1038/nature11234.

56. Yassour, M., Vatanen, T., Siljander, H., Hämäläinen, A.-M., Härkönen, T., Ryhänen, S.J., Franzosa, E.A., Vlamakis, H., Huttenhower, C., Gevers, D., et al. (2016). Natural history of the infant gut microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. Sci. Transl. Med. *8*, 343ra381-343ra381. 10.1126/scitranslmed.aad0917.

57. Zhu, S., Beaulaurier, J., Deikus, G., Wu, T.P., Strahl, M., Hao, Z., Luo, G., Gregory, J.A., Chess, A., He, C., et al. (2018). Mapping and characterizing N6-methyladenine in eukaryotic genomes using single-molecule real-time sequencing. Genome Res. 10.1101/gr.231068.117.

58. Beaulaurier, J., Zhu, S., Deikus, G., Mogno, I., Zhang, X.-S., Davis-Richardson, A., Canepa, R., Triplett, E.W., Faith, J.J., Sebra, R., et al. (2018). Metagenomic binning and association of plasmids with bacterial host genomes using DNA methylation. Nat. Biotechnol. *36*, 61-69. 10.1038/nbt.4037.

59. Kong, Y., Mead, E.A., and Fang, G. (2023). Navigating the pitfalls of mapping DNA and RNA modifications. Nat. Rev. Genet. *24*, 363-381. 10.1038/s41576-022-00559-5.

60. Park, J., and Wang, H.H. (2021). Systematic dissection of σ^{70} sequence diversity and function in bacteria. Cell Reports *36*, 109590. 10.1016/j.celrep.2021.109590.

61. Nakayama-Imaohji, H., Hirakawa, H., Ichimura, M., Wakimoto, S., Kuhara, S., Hayashi, T., and Kuwahara, T. (2009). Identification of the site-specific DNA invertase responsible for the phase variation of susC/susD family outer membrane proteins in *Bacteroides fragilis*. J. Bacteriol. *191*, 6003-6011. 10.1128/jb.00687-09.

62. Joglekar, P., Sonnenburg, E.D., Higginbottom, S.K., Earle, K.A., Morland, C., Shapiro-Ward, S., Bolam, D.N., and Sonnenburg, J.L. (2018). Genetic variation of the susC/susD homologs from a polysaccharide utilization locus underlies divergent fructan specificities and functional adaptation in *Bacteroidesthetaiotaomicron* strains. Msphere *3*, e00185-18. 10.1128/mSphereDirect.00185-18.

63. Sultan, S.Z., Silva, A.J., and Benitez, J.A. (2010). The PhoB regulatory system modulates biofilm formation and stress response in El Tor biotype *Vibrio cholerae*. FEMS Microbiol. Lett. *302*, 22-31. 10.1111/j.1574-6968.2009.01837.x.

64. Cai, X.H., Yang, S.Y., Peng, Y.H., Tan, K.N., Xu, P., Wu, Z.H., Kwan, K.Y., and Jian, J.C. (2024). Regulation of PhoB on biofilm formation and hemolysin gene *hlyA* and *ciaR* of Streptococcus agalactiae. Vet. Microbiol. *289*, 109961. 10.1016/j.vetmic.2023.109961.

65. Logan, L.K., and Bonomo, R.A. (2016). Metallo-β-Lactamase (MBL)-producing Enterobacteriaceae in United States children. OFID *3*, ofw090. 10.1093/ofid/ofw090.

66. Cherak, Z., Loucif, L., Moussi, A., Bendjama, E., Benbouza, A., and Rolain, J.M. (2022). Emergence of metallo-β-lactamases and OXA-48 carbapenemase producing gram-negative bacteria in hospital wastewater in Algeria: A potential dissemination pathway into the environment. Microb. Drug Resist. *28*, 23-30. 10.1089/mdr.2020.0617.

67. Beck, L.C., Masi, A.C., Young, G.R., Vatanen, T., Lamb, C.A., Smith, R., Coxhead, J., Butler, A., Marsland, B.J., Embleton, N.D., et al. (2022). Strain-specific impacts of probiotics are a significant driver of gut microbiome development in very preterm infants. Nat. Microbiol. *7*, 1525-1535. 10.1038/s41564-022-01213-w.

68. Vandahl, B.B., Pedersen, A.S., Gevaert, K., Holm, A., Vandekerckhove, J., Christiansen, G., and Birkelund, S. (2002). The expression, processing and localization of polymorphic membrane proteins in Chlamydia pneumoniae strain CWL029. BMC Microbiol. *2*, 36. 10.1186/1471-2180-2-36.

69. Pedersen, A.S., Christiansen, G., and Birkelund, S. (2001). Differential expression of Pmp10 in cell culture infected with Chlamydia pneumoniae CWL029. FEMS Microbiol. Lett. *203*, 153-159. 10.1111/j.1574-6968.2001.tb10834.x.

70. Ansaldo, E., Slayden, L.C., Ching, K.L., Koch, M.A., Wolf, N.K., Plichta, D.R., Brown, E.M., Graham, D.B., Xavier, R.J., Moon, J.J., and Barton, G.M. (2019). *Akkermansia muciniphila* induces intestinal adaptive immune responses during homeostasis. Science *364*, 1179-1184. doi:10.1126/science.aaw7479.

71. Bae, M., Cassilly, C.D., Liu, X., Park, S.-M., Tusi, B.K., Chen, X., Kwon, J., Filipčík, P., Bolze, A.S., Liu, Z., et al. (2022). Akkermansia muciniphila phospholipid induces homeostatic immune responses. Nature *608*, 168-173. 10.1038/s41586-022-04985-7.

72. Derosa, L., Routy, B., Thomas, A.M., Iebba, V., Zalcman, G., Friard, S., Mazieres, J., Audigier-Valette, C., Moro-Sibilot, D., Goldwasser, F., et al. (2022). Intestinal *Akkermansia muciniphila* predicts clinical response to PD-1 blockade in patients with advanced non-small-cell lung cancer. Nat. Med. *28*, 315-324. 10.1038/s41591-021-01655-5.

73. Yoon, H.S., Cho, C.H., Yun, M.S., Jang, S.J., You, H.J., Kim, J.-h., Han, D., Cha, K.H., Moon, S.H., Lee, K., et al. (2021). *Akkermansia muciniphila* secretes a glucagon-like peptide-1-inducing protein that improves glucose homeostasis and ameliorates metabolic disease in mice. Nat. Microbiol. *6*, 563-573. 10.1038/s41564-021-00880-5.

74. Zheng, Y., Wang, T., Tu, X., Huang, Y., Zhang, H., Tan, D., Jiang, W., Cai, S., Zhao, P., Song, R., et al. (2019). Gut microbiome affects the response to anti-PD-1 immunotherapy in patients with hepatocellular carcinoma. JITC *7*, 193. 10.1186/s40425-019-0650-9.

75. Motta, S.S., Cluzel, P., and Aldana, M. (2015). Adaptive resistance in bacteria requires epigenetic inheritance, genetic noise, and cost of efflux pumps. PLoS One *10*, e0118464. 10.1371/journal.pone.0118464.

76. Tawk, C., Lim, B., Bencivenga-Barry, N.A., Lees, H.J., Ramos, R.J.F., Cross, J., and Goodman, A.L. (2023). Infection leaves a genetic and functional mark on the gut population of a commensal bacterium. Cell Host Microbe *31*, 811-826.e816. 10.1016/j.chom.2023.04.005.

77. Dunwell, J.M., Purvis, A., and Khuri, S. (2004). Cupins: the most functionally diverse protein superfamily? Phytochemistry *65*, 7-17. 10.1016/j.phytochem.2003.08.016.

78. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure prediction of

biomolecular interactions with AlphaFold 3. Nature *630*, 493-500. 10.1038/s41586-024-
07487-w.

79.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596*, 583-589. 10.1038/s41586-021-
03819-2.

80.   Cheng, Z.-F., and Deutscher, M.P. (2005). An important role for RNase R in mRNA
decay. Mol. Cell *17*, 313-318. 10.1016/j.molcel.2004.11.048.

81.   Chen, C., and Deutscher, M.P. (2005). Elevation of rnase r in response to multiple stress
conditions. J. Biol. Chem. *280*, 34393-34396. 10.1074/jbc.C500333200.

82.   Bryant, J.M., Brown, K.P., Burbaud, S., Everall, I., Belardinelli, J.M., Rodriguez-Rincon,
D., Grogono, D.M., Peterson, C.M., Verma, D., Evans, I.E., et al. (2021). Stepwise
pathogenic evolution of *Mycobacterium abscessus*. Science *372*, eabb8699.
doi:10.1126/science.abb8699.

83.   Gopalan-Nair, R., Coissac, A., Legrand, L., Lopez-Roques, C., Pécrix, Y.,
Vandecasteele, C., Bouchez, O., Barlet, X., Lanois, A., Givaudan, A., et al. (2024).
Changes in DNA methylation contribute to rapid adaptation in bacterial plant pathogen
evolution. PLoS Biol. *22*, e3002792. 10.1371/journal.pbio.3002792.

Materials and methods

Strains, media and growth conditions

Escherichia coli str. K-12 substr. MG1655 was used for heterologous expression of the gene of interest.
E. coli strains were cultured in LB broth (LB broth Miller, Becton) or LB agar (LB agar Miller, Becton) at
37°C. When grown in liquid medium, the cultures were aerated with shaking at 220 rpm, unless
otherwise specified.

Akkermansia muciniphila strains employed in this work include Akk_ECAM and Akk_probiotics. The
Akk_ECAM strain was isolated in this study (details provided below), while the Akk_probiotics strain was
derived from a commercial probiotic product (Glucose Control, Pendulum). All *Akkermansia* strains were
routinely grown statically under an anaerobic condition (5% hydrogen, 5% carbon dioxide, and 90%
nitrogen) using a type C anaerobic chamber with Indicator (#7250000, Coy laboratory) in BHI media
(Brain Heart Infusion Agar, Gibco) supplemented with 0.25% v/v porcine gastric mucin (M1778, Sigma-
Aldrich). For the selection of amoxicillin (AMX)-tolerant *Akkermansia*, amoxicillin sodium salt was used at
2 mg / L.

Stool specimen collection

ECAM samples were obtained from a previous study with an Institutional Review Board–approved study
which was conducted in healthy, pregnant mothers in New York City from 2011 to 2014.[1] FMT samples
were obtained from a previous study with Institutional Review Board approval.[2]

Isolation of Akkermansia strains from infant stool

The *A. muciniphila* strain Akk_ECAM was isolated using a modified version of a previously described protocol[3] based on growth in mucin medium (3 mM KH_2PO_4, 3 mM Na_2HPO_4, 5.6 mM NH_4Cl, 1 mM $MgCl_2$, 1 mM $Na_2S \cdot 9H_2O$, 47 mM $NaHCO_3$, 1 mM $CaCl_2$, 40 mM HCl, trace elements and vitamins, and 0.25% porcine gastric mucin). To enrich for *A. muciniphila*, a loopful of stool sample (~20 mg) was added into 2 ml of mucin medium and incubated at 37°C for 48 h under anaerobic conditions. The culture was passaged in mucin media three times by inoculating 100 μl into 2 ml of mucin medium and incubating until turbid. After three sequential passages, a suspension sample was streaked on BHI agar supplemented with 0.25% mucin and incubated for one week. Single colonies were re-streaked on BHI to purify, and the resulting purified colonies were then grown in 2 ml of mucin medium. To identify the isolated bacteria, DNA was isolated from 1 ml of culture (DNeasy Blood and Tissue Kit, Qiagen), and a portion of the 16S rRNA gene was amplified by PCR and sequenced (see primer sequences listed in **Table S6**).

DNA extraction

Extraction of DNA from fecal samples

Fecal DNA was extracted with a QIAamp PowerFecal Pro DNA kit (#51804, Qiagen) according to the manufacturer's instructions. Briefly, 50-100 mg of fecal sample was first homogenized by beads-beating using a Retsch mixer mill (MM400, Verder Scientific Inc., Newtown, PA) for 10 min at 25 Hz speed (5 min each side) and then subjected to column-based extraction steps as described in the kit protocol.

Extraction of DNA from bacterial cells

Genomic DNA was extracted from bacterial cells using either a Wizard Genomic DNA Purification Kit (A1120, Promega) or a Qiagen DNeasy Blood and Tissue kit (#69504, Qiagen) following the manufacturer's instruction. The DNA was eluted in 100 μL elution buffer and stored at -20 °C.

RNA extraction

The bacterial cell pellets were collected from 0.5 ml of cell cultures grown in BHI media supplemented with 0.25% mucin after 48h of incubation and saved in -80 °C. Total RNA was extracted with an RNeasy Mini kit (#74104, Qiagen). Briefly, cells were first lysed in RLT buffer (from the kit) with the addition of

10 µl/ml β-mercaptoethanol. The lysate was further processed by adding the content of a PowerBead tube (glass, 0.1 mm; Qiagen) and homogenizing the mixture for 5 min at a speed of 30 Hz using a Retsch mixer mill. Subsequent steps followed the manufacturer's instructions. Contaminating genomic DNA was depleted using on-column DNase treatments with DNase (#79254, Qiagen). RNA quality was assessed with a Qubit 4 Fluorometer (Invitrogen) using the Qubit 1× dsDNA HS Assay Kit to detect DNA contamination and the Qubit RNA HS Assay Kit to measure total RNA. Ribosomal RNA was depleted using NEBNext rRNA Depletion Kit (Bacteria) (#E7850, NEB). The quality of total RNA was checked using an Agilent 2100 Bioanalyzer (#5067-4626, Agilent Technologies USA).

Real-time quantitative RT-PCR

One microgram of total RNA was converted to cDNA using SuperScript™ VILO™ MasterMix kit (#11755050, Invitrogen). Gene-specific primer pairs are provided in **Table S6**. SYBR-based RT-qPCR was performed in 384-well plates (#4306737; Applied Biosystems) and analyzed by a qPCR system (QuantStudio 6 Flex, Thermo Fisher Scientific). The thermal cycling protocol was as follows: 10 min at 50°C, 1 min at 90°C, followed by 40 cycles of 30 s at 60°C and 1 min at 90°C, with a final melt curve stage. SYBR signal was normalized to the ROX reference dye. Cycle of quantification (Cq) values were automatically determined using QuantStudio Design & Analysis Software (v1.3). Based on the melting curves, the primers were specifically bound to the target genome. All the samples were run in replicates technical replicates.

Library preparation and sequencing

Oxford Nanopore whole genome sequencing

gDNA samples were sheared to an average length of 10 kb using g-Tubes (#520079, Covaris) and purified with 0.4 × AMPure XP beads (A63882, Beckman Coulter Inc.) to remove short fragments. Native libraries were prepared using Oxford Nanopore Technologies (ONT) ligation sequencing kit and native barcoding kit (SQK-LSK109, and SQK-NBD112-24, respectively). PCR barcoding was performed using the Native Barcoding Expansion 1-12 kit (EXP-NBD104) and/or 13-24 kit (EXP-NBD114) for sample multiplexing. Whole genome amplification (WGA) were conducted with the REPLI-g Mini kit (#150023, Qiagen) using 10 ng of input DNA and 16 h of incubation. WGA samples were treated with T7 endonuclease I (M0302S, NEB) to maximize sequencing yield, as per ONT documentation. Native or amplified gDNA samples were sequenced on R9.4.1 and R10.4.1 MinION flow cells using MinION MK1B or MK1C instruments. Basecalling was performed with Guppy v.5.0.7 (ONT) in GPU mode using the dna_r9.4.1_450bps_sup.cfg model for R9.4 data or Dorado v0.4.1 (ONT) with dna_r10.4.1_e8.2_400bps_sup.cfg model for R10.4 data.

ONT targeted sequencing

The RMS region was amplified using specific forward and reverse primers (**Table S6**) to generate a 6.6-kb amplicon, which included sequences enable sample barcoding. PCR was performed in a 50 μl reactions containing 250 ng of gDNA and 0.7 μM of primers. Specific barcodes were assigned to the PCR products using the PCR Barcoding Expansion 1–96 kit (EXP-PBC096, ONT). A second-round PCR was performed with 0.5 nM purified PCR product, 1 × LongAMP PCR buffer, 2 mM of $MgCl_2$, 0.3 mM of dNTPs and 5 units of LongAmp Taq DNA polymerase (NEB) in a 50 μl reaction. The thermal cycling protocol was 95°C for 3 min, followed by 15 cycles of 95°C for 15 s, 62°C for 15 s, 65°C for 7 min, with a final extension at 65°C for 7 min. Amplicons were sequenced on R9.4.1 / R10.4.1 Flongle flow cells.

PacBio WGS sequencing

The SMRTbell library was prepared according to the manufacturer's instructions. DNA was sheared to ~5,000 bp using a Covaris LE220 Focused-ultrasonicator (#500569, Covaris). The samples underwent damage and end repair, followed by ligation of single-stranded barcoded adaptors. Sequencing primer was annealed to the SMRTbells and incubated with templates for 1 hour at 20 ºC using a primer-to-template ratio of 20:1. The Sequel 2.1 polymerase was bound to the annealed complex at a polymerase to SMRTbell ratio of 30:1 and incubated for 4 h at 30 °C. The binding complex was cleaned, diluted to the final loading concentration (125-175 pM), and held at 4 °C until diffusion-based loading on the Sequel II instrument (PacBio). Sequencing run used 8M sequencing chips, with 30 h movies recorded following the manufacturer's guidelines. Most metagenome samples except for some FMT samples, were sequenced on the latest Revio instrument (PacBio). Circular consensus sequencing (CCS) reads were generated using CCS v.3.4.1 (https://github.com/PacificBiosciences/ccs) with default settings.

Illumina RNA sequencing

RNA-seq library construction for Illumina sequencing was performed using the NEBNext Ultra II RNA Library Prep Kit for Illumina (#E7770, NEB). Library quality and size were assessed using an Agilent 4200 TapeStation system (Agilent Technologies, Inc.) and quantified with a Qubit 4 Fluorimeter using Qubit DNA HS Assay Kit (Invitrogen). All samples were equitably pooled, size-selected, and sequenced on an Illumina HiSeq 2500 platform, generating paired-end reads of 150 base pairs (bp) each.

Phylogenetic analysis of HMP reference genomes

The bacterial genomics sequences and proteome files were downloaded from the NIH Human Microbiome Project (HMP) legacy data browser website (https://www.hmpdacc.org/hmp/HMRGD/). A phylogenetic tree was constructed using PhyloPhlAn v3.0.67[4] with proteome files as input and high--

diversity configuration. The tree was visualized and edited using the interactive tree of life (iTOL) software (https://itol.embl.de).

De novo assembly of metagenomes

ONT read metagenomic assembly and taxonomic annotation

Basecalled reads were processed using metaFlye tool v2.9[5] with the parameters ---nano-raw --meta --keep-haplotypes --iterations 2. Meta-assemblies were divided into species-level bins using the MaxBin

985 2.0 software.[6] Strainberry v1.1[7] was used to phase haplotypes and separate strain level contigs. Contigs were subsequently binned using the "binning" module of metaWRAP v1.3.2.[8] The quality of the reconstructed MAGs was evaluated using the 'bin_refinement' module of metaWRAP v1.3.2, and taxonomic assignment was performed using GTDB-tk v3.8.[9]

SMRT read metagenomic assembly and taxonomic annotation

990 HiFi sequencing reads were assembled using hifiasm_meta 0.13-r308 (https://github.com/xfengnefx/hifiasm-meta) with default parameters, focusing on primary contigs. To maximize MAGs recovery, the HiFi-MAG-Pipeline (https://github.com/PacificBiosciences/pb-metagenomics-tools/tree/master/HiFi-MAG-Pipeline) was deployed, increasing both the total number of MAGs and single-contig circular MAGs.

995 ### Hybrid assembly

Nanopore-based long reads and Illumina paired-end short reads were used for a hybrid assembly with MUFFIN,[10] producing high-quality and complete bins.

Gene annotation and RMS prediction

1000 Gene annotation for bacterial genomes and MAGs was performed using Prokka v1.14.6.[11] RMS identification followed previously established methods.[12] Curated reference protein sequences for types I, II, IIC, and III RMS, and type IV REases were downloaded from the REBASE[13] "gold standards" dataset (accessed November 2016). A dedicated BLAST database was created for the specific gene (hsdS) of Type I RMS using 39,729 HsdS proteins from REBASE. The TRD domain in hsdS genes were identified

1005 using HMMER v3.3 (http://hmmer.org) with a profile built from 31 TRD domains.

Differential-expression analysis of RNA-seq data

Read quality was assessed using FastQC v.0.11.9 (http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Adapter sequences and low-quality reads were removed using Trimmomatic v.0.36[14] with the following parameters: PE, -phred33,

 ILLUMINACLIP:<adapters.fa>:2:30:10:8: keepBothReads, SLIDINGWINDOW:4:15, LEADING:20,
TRAILING:20, and MINLEN:50. rRNA sequences were then filtered out using SortMeRNA v.2.1b [15] with
the SILVA 16S/23S[16] and the Rfam 5S[17] rRNA database. Non-rRNA reads were aligned to the reference
genome using BWA-MEM v.0.7.17-r1198.[18] Read assignment was performed with FeatureCounts
v.2.0.1,[19] excluding multi-mapping and multi-overlapping reads. Normalization and differential expression

 testing were conducted using DESeq2 v.1.18.1,[20] with differentially expressed genes defined as those
with an FDR-corrected $p < 0.05$ and $|log2[FC]| > 1$.

Gene set enrichment analysis

Gene function was annotated by eggmapper v2.1.6 against the defaults eggNOG database.[21] GO and
KEGG enrichment analyses were performed using the R package clusterProfiler v.3.18.1. [22] KEGG

 pathways and modules were filtered for UniRef90 v.2019_06 entries, downloaded from Uniprot website.[23]
Enrichments at different taxonomic levels were calculated and visualized as heatmap using
ComplexHeatmap package v.2.6.2.[24]

Genome-wide epigenetic phase variation (ePV) detection in long-read data

 Identification of simple sequence repeat (SSR) in MAGs
Assembled MAGs were analyzed to identify SSR signatures. FASTA files containing all the assembled
sequences were input into the MISA Perl script
(https://github.com/cfljam/SSR_marker_design/tree/master). The following minimum repeat thresholds for
microsatellites were specified: (unit size/minimum number of repeats): (1/10) (2/6) (3/5) (4/5) (5/5) (6/5).

 The maximum length of spacer sequence was set to 100 bp in the MISA setup file.

Structural variation (SV) detection
Basecalled ONT reads were aligned to the de novo assembled MAGs with NGMLR v 0.2.6.[25] Hifi PacBio
reads were aligned to the de novo assemblies with pbalign (SMRTlink v10,
https://downloads.pacbcloud.com/public/software/installers/smrtlink_10.1.0.119588.zip). The resulting

 alignment files were processed with sniffles v2.0.7[26] for structural variant calling. Detected SVs were
mapped to their corresponding genome locations, and SVs overlapping predicted RMS regions / orphan
MTases (within a 2,000 bp window) were defined as potential ePVs. The ePV list was refined through
manually curation.
Quantification of ePV with raw reads

 To overcome the limitations of SV detection tools in quantifying read counts assigned to genomic variants
in complex rearrangement regions, we developed an in-house pipeline: (1) predicted all possible genomic
variants base on the reference genome and the read alignment; (2) create an *in silico* reference for each
variants and merged them; (3) realigned reads to the new reference genomes; (4) applied strict filtered

thresholds, including "MAPQ 60 & mg > 99" and "-max softsclip 300", (5) counted the reads uniquely
1045 mapped to each genotype.

Site-specific ePV detection for PacBio data

Preprocess of PacBio raw data

PacBio sequencing data contained both DNA sequence and kinetic data, which were used to detect
1050 single-molecule methylation signal. Raw SMRT Sequel II data was processed into CCS reads and
demultiplexed using SMRTlink v10.

Genome-wide aggregated methylation analysis

Subreads and CCS reads were aligned to MAGs using pbalign (SMRTlink v10) with default parameters,
and alignments were separated by MAG. The IPD ratio was calculated with ipdSummary (SMRTlink v10)
1055 for each MAGs. De novo methylation motifs were identified using motifMaker (SMRTlink v10) with default
parameters. All of the above analyses aggregate the CCS data based on the MAG genomes, as opposed
to single-molecule analysis, which is based on each individual CCS reads.

Single molecule level methylation analysis

Subreads were split based on CCS read ID and mapped to their corresponding CCS reads using pbalign
1060 with default parameters. Alignments were filtered with strict thresholds, including "MAPQ 60 & mg > 99"
and "-max softsclip 300". Single-molecule IPD ratio was calculated with ipdSummary. For Revio data, raw
data lacked subread information and require an in-house scripts for single-molecule analysis (released
the detailed scripts on Zenodo).

Identification of single site ePV

1065 IPD ratio files from single-molecule methylation signals were processed in R v4.0.3. Low
methylation sites (< 50%) were identified on both strands, and only sites with low methylation on
both strands were considered partially methylated sites. The methylation variation sites were
visualized across MAGs, with loci and potentially affected genes highlighted in a gene diagram
chart.

1070

Genome-wide ePV detection in Illumina short-read data

For short-read metagenomic data, ePVs exclusively referred to inverton, as defined in a
previous study.[27] Invertons across the metagenomic samples were identified using the
PhaseFinder[27] pipeline, with HMP reference genomes (https://hmpdacc.org/hmp/HMRGD/) as

1075 reference. Invertons overlapping predicted RMS regions / orphan MTase (within a 2,000 bp window) were classified as potential ePVs.

Protein extraction, purification and identification

Protein preparation
E. coli MG1655 cells were harvested by centrifugation at 5,000 × g for 10 minutes at 4°C. The
1080 supernatant was collected in 50 ml aliquots. The cell pellet was resuspended in lysis buffer containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 2 mM tris (2-carboxyethyl)phosphine (TCEP, #68957, Sigma-Aldrich) and a protein inhibitor cocktail (P8340, Sigma-Aldrich). The suspension was lysed by sonication on ice (15-second pulses, 10 times, with 2-min intervals). The lysate was clarified by centrifugation at 12,000 × g for 30 min at 4°C to remove cell debris. The supernatant from cell culture was concentrated to
1085 1 ml using an Amicon Ultra Centrifugal Filter unit (3 kDa, Millipore) and subjected to sonication as described above. The resulting crude protein solution was purified using a Capturem His-Tagged Purification kit (#635719, TaKaRa Bio Inc.) following the manufacturer's instructions.

SDS-PAGE
Protein samples were mixed with 4× Sodium dodecyl sulfate (SDS) loading buffer (Bio-Rad Laboratories,
1090 Inc) and heated at 98°C for 5 min. SDS-polyacrylamide gel electrophoresis (SDS-PAGE) was performed using Mini-PROTEIN TGX Gels (4–20%, Bio-Rad Laboratories, Inc). Gels were stained with Coomassie Brilliant Blue R-250, and protein bands were visualized using a gel documentation system (Bio-Rad Laboratories, Inc).

Intact Protein Mass Spectrometry
1095 For intact protein mass analysis, purified protein samples were buffer exchanged into 1 M ammonium acetate solution using a 3 kDa Amicon Ultra Centrifugal Filter unit (Millipore). Qualitative analysis of recombinant protein was performed with a 1260 HPLC system coupled to a 6545 XT Q-TOF MS instrument operated in positive ESI mode (Agilent Technologies, Inc.).

The acquired spectra were extracted and deconvoluted with Bioconfirm A 2.0 software (Agilent
1100 Technologies) with the default parameters, except that the mass range was set to be 10 kDa to 100 kDa.

In vitro passage assay of *A. muciniphila*

AMX selection assay
1105 An *A. muciniphila* isolate with the g1 RMS genotype was selected as the starting material and cultured on BHI agar plates supplemented with 0.25% mucin for three days. 46 single colonies were picked and subcultured in 3 ml BHI broth supplemented with 0.25% mucin for two additional passages. The above subcultures were diluted to an initial OD_{600} of 0.01 and a 50 µl aliquot was inoculated in 150 µl BHI liquid

medium supplemented with 0.25% mucin, and with or without 2 mg / L AMX. The subculture process was repeated for nine passages. Cells from each passage were collected and stored. gDNA extraction and targeted sequencing were performed as described above.

RMS Reversible assay

To assess the conditions under which the RMS genotype could revert from non-g1 to g1, *A. muciniphila* colonies with the non-g1 RMS genotype from the AMX selection assay were stressed with various microbiological mediums, antibiotics, and atmospheric conditions (see detailed design and conditions in **Table S7**). RMS genotypes were determined across the assay using targeted sequencing.

Eexpression of the *mucC* gene in *E. coli*

To overexpress the *mucC* gene, we synthesize the gene with codons optimized for expression in *E. coli*. The gene was inserted into a median-copy-number plasmid containing a constitutive expression promoter. The recombination plasmid was then transformed into *E. coli*. Overnight cell cultures were diluted 1:100 into fresh LB medium containing different concentrations of AMX. The growth curve of each group of cells (OD_{600}) was monitored using a microplate spectrophotometer (SpectraMax M5, USA).

AMX sensitivity test

AMX test strips (AES, Liofilchem) were placed on the surface of a BHI agar plate supplemented with 0.25% mucin and inoculated with a 500 µl bacterial suspension. The plates were incubated at 37°C for 48 h under anaerobic conditions. The MIC was determined by the intersection of the lower part of the ellipse-shaped growth inhibition area with the test strip.

Akkermansia-fed Mouse model

A probiotic supplement including *Clostridium beijerinckii* WB-STR-0005, *Clostribium butyricum* WB-STR-0006, *A. muciniphila* WB-STR-0001 (introduced earlier as Akk_probiotics), *Anaerobutyricum hallii* WB-STR-0008, and *Bifidobacterium infantis* 100 (Pendulum, USA) was administered to healthy male germ-free (GF) C57BL/6J mice (JAX stock #000664) (n=4). 12-week-old GF mice were fed 10 mg probiotic supplement via Mini Capsule Kit (Braintree Scientific XKITM) for the first two days. Fecal samples were collected at day 7.

Quantification and statistical analysis

Statistical analyses were performed using R (v.4.0.3). Statistical significance was determined using unpaired two-tailed Mann–Whitney–Wilcoxon test, unpaired two-tailed Welch's t-test, and unpaired Student's t-test, as indicated in figure legends. Data are shown as mean $\pm$ s.d. $P < 0.05$ was considered statistically significant. Information regarding the number of samples and statistical tests are provided in the figure legends.

References

1. Bokulich, N.A., Chung, J., Battaglia, T., Henderson, N., Jay, M., Li, H., D. Lieber, A., Wu, F., Perez-Perez, G.I., Chen, Y., et al. (2016). Antibiotics, birth mode, and diet shape microbiome maturation during early life. Sci. Transl. Med. *8*, 343ra382-343ra382. 10.1126/scitranslmed.aad7121.
2. Aggarwala, V., Mogno, I., Li, Z., Yang, C., Britton, G.J., Chen-Liaw, A., Mitcham, J., Bongers, G., Gevers, D., Clemente, J.C., et al. (2021). Precise quantification of bacterial strains after fecal microbiota transplantation delineates long-term engraftment and explains outcomes. Nat. Microbiol. *6*, 1309-1318. 10.1038/s41564-021-00966-0.
3. Derrien, M., Vaughan, E.E., Plugge, C.M., and de Vos, W.M. (2004). *Akkermansia muciniphila* gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. Int. J. Syst. Evol. Microbiol. *54*, 1469-1476. 10.1099/ijs.0.02873-0.
4. Asnicar, F., Thomas, A.M., Beghini, F., Mengoni, C., Manara, S., Manghi, P., Zhu, Q., Bolzan, M., Cumbo, F., May, U., et al. (2020). Precise phylogenetic analysis of microbial isolates and genomes from metagenomes using PhyloPhlAn 3.0. Nat. Commun. *11*, 2500. 10.1038/s41467-020-16366-7.
5. Kolmogorov, M., Bickhart, D.M., Behsaz, B., Gurevich, A., Rayko, M., Shin, S.B., Kuhn, K., Yuan, J., Polevikov, E., Smith, T.P.L., and Pevzner, P.A. (2020). metaFlye: scalable long-read metagenome assembly using repeat graphs. Nat. Methods *17*, 1103-1110. 10.1038/s41592-020-00971-x.
6. Wu, Y.-W., Simmons, B.A., and Singer, S.W. (2015). MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics *32*, 605-607. 10.1093/bioinformatics/btv638.
7. Vicedomini, R., Quince, C., Darling, A.E., and Chikhi, R. (2021). Strainberry: automated strain separation in low-complexity metagenomes using long reads. Nat. Commun. *12*, 4485. 10.1038/s41467-021-24515-9.
8. Uritskiy, G.V., DiRuggiero, J., and Taylor, J. (2018). MetaWRAP—a flexible pipeline for genome-resolved metagenomic data analysis. Microbiome *6*, 158. 10.1186/s40168-018-0541-1.
9. Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., and Parks, D.H. (2022). GTDB-Tk v2: memory friendly classification with the genome taxonomy database. Bioinformatics *38*, 5315-5316. 10.1093/bioinformatics/btac672.
10. Van Damme, R., Hölzer, M., Viehweger, A., Müller, B., Bongcam-Rudloff, E., and Brandt, C. (2021). Metagenomics workflow for hybrid assembly, differential coverage binning, metatranscriptomics and pathway analysis (MUFFIN). PLoS Comp. Biol. *17*, e1008716. 10.1371/journal.pcbi.1008716.
11. Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. Bioinformatics *30*, 2068-2069. 10.1093/bioinformatics/btu153.
12. Oliveira, P.H., Ribis, J.W., Garrett, E.M., Trzilova, D., Kim, A., Sekulovic, O., Mead, E.A., Pak, T., Zhu, S., Deikus, G., et al. (2020). Epigenomic characterization of *Clostridioides difficile* finds a

conserved DNA methyltransferase that mediates sporulation and pathogenesis. Nat. Microbiol. *5*, 166-180. 10.1038/s41564-019-0613-4.

13. Roberts, R.J., Vincze, T., Posfai, J., and Macelis, D. (2014). REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. Nucleic Acids Res. *43*, D298-D299. 10.1093/nar/gku1046.

14. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

15. Kopylova, E., Noé, L., and Touzet, H. (2012). SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. Bioinformatics *28*, 3211-3217.

16. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F.O. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res. *41*, D590-D596.

17. Griffiths-Jones, S., Bateman, A., Marshall, M., Khanna, A., and Eddy, S.R. (2003). Rfam: an RNA family database. Nucleic Acids Res. *31*, 439-441.

18. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics *25*, 1754-1760.

19. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

20. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 1-21.

21. Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. (2021). Eggnog-mapper v2: Functional annotation, orthology assignments, and domain prediction at the metagenomic scale. Mol. Biol. Evol. *38*, 5825-5829. 10.1093/molbev/msab293.

22. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. OMICS *16*, 284-287. 10.1089/omi.2011.0118.

23. Consortium, T.U. (2022). UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res. *51*, D523-D531. 10.1093/nar/gkac1052.

24. Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics *32*, 2847-2849. 10.1093/bioinformatics/btw313.

25. Sedlazeck, F.J., Rescheneder, P., Smolka, M., Fang, H., Nattestad, M., von Haeseler, A., and Schatz, M.C. (2018). Accurate detection of complex structural variations using single-molecule sequencing. Nat. Methods *15*, 461-468. 10.1038/s41592-018-0001-7.

26. Smolka, M., Paulin, L.F., Grochowski, C.M., Horner, D.W., Mahmoud, M., Behera, S., Kalef-Ezra, E., Gandhi, M., Hong, K., Pehlivan, D., et al. (2024). Detection of mosaic and population-level structural variants with Sniffles2. Nat. Biotechnol. 10.1038/s41587-023-02024-y.

27. Jiang, X., Hall, A.B., Arthur, T.D., Plichta, D.R., Covington, C.T., Poyet, M., Crothers, J., Moses, P.L., Tolonen, A.C., Vlamakis, H., et al. (2019). Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host adaptation in the gut. Science *363*, 181-187. 10.1126/science.aau5238.

Fig. 1

A

Epigenetic phase variation (ePV)

A bacterial strain

Methylome A
Methylome B

Transcriptome A
Transcriptome B

Phenotype A
Phenotype B

Condition A

Condition B

B

Genome-wide ePV

Active MTase

MTase off → Inactive MTase

MTase motif (A → B) → Active MTase

C

Site-specific ePV

Active MTase

TF competing

Active MTase

Active MTase
Inactive MTase
TF Transcriptional factor

Specific subunit A
Specific subunit B
Methylation

Sequence motif A
Sequence motif B

D

Prevalence of ePV in the human microbiome (new data + public data)

Diverse cohorts

HMP
Infant
Antibiotics
Diseases
FMT

E

Function characterization of an ePV

In vitro passage assay

DNA sequencing

gDNA + amplicons

Gene function validation

GOI

RNA sequencing

Total RNA → cDNA

Fig. 2

Fig. 3

A

Bacteroides uniformis Type II MTase

genotype (g1)
(g2)

g1: 48.7%
g2: 51.3%

Raw reads assignment (IGV)

2,391 bp

g1

g2

B

Bacteroides finegoldii Type I RMS

g1
g2

g1: 38.2%
g2: 61.8%

Klebsiella michiganensis Type II MTase

g1
g2

g1: 25.7%
g2: 74.3%

Bacteroides fragilis Type III MTase

g1
g2

g1: 1%
g2: 99%

M — MTase   Other genes   S1 — Specific gene 1
S2 — Specific gene 2   Inverted repeat (IR)   Promoter

C

① Genome-wide ePV
② Candidate genome-wide ePV

◁ Type I
○ Type II
□ Type III

Order   Clostridiales   Enterobacterales   Bacteroidales   Bifidobacteriales

D

Infant 4   AMX →   Infant 5   AMX →

Infant 6   AMX →   Infant 7   AMX →

E

B. uniformis isolates (153) → Hit genomes (23)

9   14

Count

g1   g2

Fig. 4

Fig. 5