1 **CellSP: Module discovery and visualization for subcellular spatial transcriptomics**
2 **data**
3

4 Bhavay Aggarwal[1], Saurabh Sinha[1, 2]*.
5
6 [1] The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology, Atlanta,
7 GA, 30332, USA
8 [2] H. Milton Stewart School of Industrial & Systems Engineering, Georgia Institute of Technology, Atlanta,
9 GA, 30318, USA
10
11
12 *Correspondence: saurabh.sinha@bme.gatech.edu

ABSTRACT

Spatial transcriptomics has enabled the study of mRNA distributions within cells, a key aspect of cellular function. However, there is a dearth of tools that can identify and interpret functionally relevant spatial patterns of subcellular transcript distribution. To address this, we present CellSP, a computational framework for identifying, visualizing, and characterizing consistent subcellular spatial patterns of mRNA. CellSP introduces the concept of "gene-cell modules", which are gene sets with non-random subcellular transcript distributions in many cells. It provides intuitive visualizations of the captured patterns and offers functional insights into each discovered module. We demonstrate that CellSP reliably identifies functionally significant modules across diverse tissues and technologies. We use the tool to discover subcellular spatial phenomena related to myelination, axonogenesis and synapse formation in the mouse brain. We find immune response-related modules that change between kidney cancer and healthy samples, and myelination-related modules specific to mouse models of Alzheimer's Disease.

INTRODUCTION

Spatial transcriptomics (ST) technologies and analytical tools available today offer unprecedented views of gene expression patterns within the intricate tapestry of tissues [1] [2] [3]. At the cutting edge of this technology are single-molecule resolution assays [4] [5] [6], which provide a detailed view of transcript distributions within individual cells, promising to transform our understanding of mRNA localization and its relationship with cellular functions. This relationship has been documented in the literature through anecdotal examples showing that RNA localization to specific subcellular regions may underlie efficient spatial organization of cellular processes [7], rapid local translation in response to external stimuli [8], maintenance of cell polarity [9] [10], facilitation of cell migration [11], coordination of developmental patterning [12], etc. Thus, there is growing recognition of the need to expand our understanding of subcellular RNA localization [13], and the time is ripe for this pursuit, supported by the state-of-the-art subcellular spatial transcriptomics tools.

While experimental techniques have advanced rapidly – enabling, for instance the mapping of individual transcripts of hundreds to thousands of genes in thousands of cells [14] [15] [16] – analytical methods have lagged behind. Available ST-related tools are mostly designed for analysis at the single-cell resolution [17]. More recently, methods targeting subcellular phenomena have emerged [13] [18] [19] [20] that can highlight subcellular localization patterns involving single genes or gene pairs, in individual cells. For example, a gene may be annotated as having transcripts localized to cell edges by the BENTO tool [13] or having a "radial" distribution in a cell by the SPRAWL software [19]. Similarly, a gene pair may be found to be significantly colocalized by the InSTAnT toolkit [18], in an individual cell or across many cells. However, when applied to a typical subcellular ST dataset, these tools identify tens of thousands of statistically significant subcellular spatial patterns involving individual genes or gene pairs in individual cells. This overwhelming volume of results imposes a significant interpretive burden, complicating the transition from statistical findings to meaningful and actionable biological insights. The goal of this work is to bridge this gap with an analytical method that identifies the most salient subcellular spatial patterns in an ST data set, along with biological interpretations of those patterns.

An important lesson from decades of transcriptomics data analysis is the concept of gene "module" – a set of genes that share some common pattern of expression. Systematic identification of gene modules [21] followed by statistical enrichment tests [22] is a popular approach to distil a large number of statistical observations into a more compact set of systems-level insights. Inspired by this paradigm, we sought a method to define gene modules with shared subcellular spatial patterns. One simple strategy is to determine if a gene exhibits a specific

subcellular localization (e.g., "cell edge" according to BENTO) in many cells, and group together
all genes exhibiting the same frequent localization into a module. This strategy is overly
permissive, since two genes annotated with the same frequent localization pattern may not exhibit
that common pattern in the same cells. We therefore conceptualized a "gene-cell" module as a
set of genes that exhibit the same subcellular pattern in the same cells. However, requiring all
genes of a module to exhibit the same subcellular pattern in all module cells is overly restrictive,
and will result in fragmented modules. To address this, we adopted a biclustering technique for
discovering gene-cell modules from larger collections of genes and gene pairs identified by
subcellular pattern discovery tools. This algorithm forms the core of the software presented here,
called "CellSP".
73
CellSP analyzes single-molecule resolution ST data, identifies significant subcellular spatial
distribution patterns at the gene level, and distills them into a compact list of gene-cell modules
that typically comprise tens of genes and hundreds of cells. It provides specialized techniques for
visualizing such modules and their defining spatial patterns. It uses gene set enrichment analysis
to describe the genes comprising the module. Additionally, it uses machine learning classification
to distinguish module-associated from other cells in the tissue based on their transcriptomic
profiles, identifying genes and biological properties (beyond cell type) that characterize module
cells.
82
We used CellSP to analyze ST data sets from four different studies generated using two different
technologies. We present the results of these analyses and provide guidance on interpreting the
discovered modules. We identify modules associated with myelination in mouse brain tissues
across various technologies and find these modules to exhibit differences linked to Alzheimer's
disease. The reported differences are not necessarily detectable through traditional gene
expression analyses. We also find modules related to cell-cell adhesion and axonogenesis, and
a closer examination of these modules sheds light on interactions among neighboring cells. Use
of CellSP on a cancer data set reveals immune-response related gene-cell modules representing
subcellular phenomena specific to cancer versus healthy tissue. In summary, CellSP establishes
a robust framework for the systematic detection, visualization and comparison of subcellular
transcriptomic patterns, coupled with a statistical characterization of their associated biological
functions.
95
96
97 **RESULTS**
98
Overview of CellSP
CellSP is a tool for analysis of spatial transcriptomics (ST) data at single-molecule resolution
(**Figure 1A**). It identifies "gene-cell modules", where a module is defined as a set of genes
exhibiting a specific subcellular spatial distribution pattern (of their transcripts) in a set of cells
(We use the terms "gene-cell module" and "module" interchangeably.) Modules are recovered
based on statistical criteria and represent the occurrence of interesting, persistent subcellular
spatial phenomena in the tissue. There are three main steps involved:
106
Step 1 – Subcellular pattern discovery. Here, the statistical tools SPRAWL [19] and InSTAnT [18]
are used to rigorously identify subcellular spatial patterns involving individual genes (SPRAWL)
or gene pairs (InSTAnT), in each cell (**Figure 1B**). SPRAWL identifies four types of subcellular
patterns – peripheral, radial, punctate and central – describing the distribution of a gene's
transcripts within the cell, while InSTAnT tests if transcripts of a gene pair tend to be proximal to
each other more often than expected by chance. Patterns of each type are stored in a matrix

whose columns represent genes or gene pairs and rows represent cells (**Figure 1C**, left), with
entries representing strength of pattern occurrence. Five such matrices are produced, one for
each of the four spatial patterns from SPRAWL analysis and one from InSTAnT analysis.
116
Step 2 – Module discovery. Next, a statistical tool called LAS (Large Average Submatrices) [23]
is used to analyze each pattern annotation matrix and identify "biclusters", i.e., a subset of rows
and columns with a large average value (**Figure 1C**, middle). Each bicluster represents a set of
genes or gene pairs that exhibit the same type of subcellular pattern in the same set of cells, with
statistical significance estimated by a Bonferroni-based score (**Methods**). Biclusters identified in
this manner may overlap in terms of rows (cells) and columns (genes/gene pairs), yielding many
redundant modules. To address this, CellSP deploys an iterative module-coalescing process
where pairs of modules comprising similar sets of cells and genes are combined into larger
modules (**Methods**, **Figure 1C**, right). The module discovery step utilizes a parallel computing
implementation to accelerate runtime and facilitate the analysis of large datasets.
127
Step 3 – Module characterization. To aid biological interpretation, CellSP reports shared
properties of the genes and cells of each discovered module (**Figure 1E**). Genes are
characterized using Gene Ontology (GO) enrichment tests, while cells are characterized by their
cell type composition if such information is available. To provide a more precise characterization
of a module's cells, CellSP trains a machine learning classifier to discriminate those cells from all
other cells, using the expression levels of all genes other than the module genes. (Here, it uses
a reference scRNA-seq data set, if available, to impute levels of genes missing in the ST data set;
see **Methods**.) Genes that are highly predictive in this task are then subjected to GO enrichment
tests, furnishing hypotheses about biological processes and pathways that are active specifically
in the module cells.
138
User experience: The CellSP tool ingests a single-molecule resolution ST data set in the popular
"AnnData" format with cell delineations (**Methods**) and returns a listing of gene-cell modules
identified for all five pattern types (peripheral, radial, punctate, central, colocalization). The
detailed report of each module includes the type of subcellular pattern, identities of genes, number
of cells, GO terms characterizing the module genes, cell type composition and GO terms
associated with module cells. The latter two are provided depending on the availability of cell type
annotations and reference scRNA-seq data (**Methods**), both of which are optional.
146
Visualization: CellSP introduces new techniques for intuitive visualization of subcellular
phenomena represented by modules. Here, we outline these visualization methods in general
terms, specific examples are provided in subsequent sections. For modules defined by gene pair
colocalization, it creates a heatmap showing that colocalization frequencies are higher for module
genes versus other genes, and in module cells versus other cells (**Figure 1D**, right). Modules of
genes that localize in "central" or "peripheral" pattern are depicted using a density plot of transcript
distribution at varying radial locations in an idealized circular cell, averaged over all module cells
(**Figure 1D**, middle). Subcellular patterns of types "radial" and "punctate" are depicted using a
density plot binned by sectors of an idealized circular cell, aggregated over all module cells by
aligning their densest sector (**Figure 1D**, left). In each of these visualizations, a side-by-side visual
comparison with non-module genes is provided to underscore the specificity of the pattern to the
module genes. Additional visual aids provided in the report include sample cells illustrating the
subcellular phenomenon directly, "REVIGO" [24] plots of GO term enrichments, and Uniform
Manifold Approximation and Projection (UMAP) [25] as well as spatial plots of module cells.
161
CellSP reveals myelination- and axonogenesis-related subcellular spatial phenomena in
preoptic area of mouse hypothalamus

In its first demonstrative application, we used CellSP to analyze a MERFISH data set comprising
5,149 cells in the hypothalamic preoptic area (POA) of a mouse brain [26], identifying 38 modules
spanning the five pattern types and each module comprising 9 genes and 138 cells on average
(**Figure S1A, Data S1**). Each gene appears in less than four modules on average (**Figure S1B**)
and each cell appears in less than three modules on average (**Figure S1C**), indicating a small
degree of overlap among the modules. The modules comprise genes enriched in a range of
biological processes and cellular components, especially cytoskeleton and cellular structure
regulation, neuronal function and development, and metabolic processes (**Data S1**). Note that the
underlying subcellular pattern discovery tool InSTAnT reports 358 significant gene pairs (p-value
< 1e-3) that colocalize in 35 cells on average, while SPRAWL reports 134 genes having a
significant pattern (score threshold = 0.5) in 351 cells on average across the four patterns. The
resulting compendium of significant statistical findings reported by these tools, tens of thousands
in number, creates a substantial interpretive burden. CellSP effectively distills these patterns into
a manageable set of modules, streamlining the crucial step of biological interpretation.
178
We next illustrate the format in which CellSP reports its discovered modules (**Figure S2**) and how
specific biological insights can be gleaned from these. The module "M0_I" consists of the four
genes Sgk1 (serum/glucocorticoid regulated kinase 1), Ttyh2 (tweety family member 2), Ermn
(ermin, ERM-like protein) and Ndrg1 (N-myc downstream regulated 1), and a set of 332 cells
(**Figure 2A**). It was identified using InSTAnT in the pattern discovery step (Step 1, see above),
which means that these genes exhibit statistically significant colocalization in the module cells, a
selection of which are shown in **Figure 2B**. Notably, the colocalization is observed in varying
subcellular regions such as the nucleus, nuclear periphery, cytoplasm and cell membrane. (See
Figure S1E for more examples.) **Figure 2C** illustrates how CellSP helps visualize a colocalization
module, highlighting the specificity of the phenomenon to the module's genes and cells. The
module cells are mostly mature oligodendrocytes (67%, **Figure 2D**) and the module genes are
enriched for GO terms such as the molecular function "cytoskeletal protein binding" (p-value =
0.001), the cellular component "myelin sheath" (p-value 0.004) and the biological process
"supramolecular fiber organization" (p-value = 0.02) (**Figure 2E**). (Note that cytoskeletal protein
binding and supramolecular fiber organization play important roles in formation and maintenance
of myelin sheath [27] [28].) Moreover, according to CellSP's classification-based characterization
(Step 3, see above), the module cells are marked by genes enriched for "ensheathment of
neurons" (p-value = E-12) (**Figure 2F**). Thus, CellSP analysis suggests a connection between
subcellular colocalization of the module and the myelination process in mature oligodendrocytes.
A possible interpretation is that the mRNAs of these four genes are captured in the process of
being transported to the myelin sheath for local translation, as has been recorded for at least two
other myelin biogenesis-related genes, viz., Mbp (Myelin Basic Protein) [29] and Mobp (Myelin
Oligodendrocyte Basic Protein) [30]. Interestingly, another CellSP-reported module – "M11_S" –
represents a punctate subcellular pattern (**Figure S1D**) involving three of the four above genes,
viz., Sgk1, Ttyh2, Ermn, and another myelination-related gene, Gjc3 (gap junction protein gamma
3) that has been shown to localize in myelin sheaths [31].
205
An example of a module discovered using SPRAWL in the pattern annotation step is "M0_S"
(**Figure 3A**), consisting of four genes – Fn1 (fibronectin 1), Slco1a4 (solute carrier organic anion
transporter family member 1a4), Rgs5 (regulator of G protein signaling 5), Sema3c (semaphorin
3C) – distributed in a peripheral pattern (**Figure 3B**) in 221 cells. **Figure 3C** illustrates how CellSP
helps visualize such a module, revealing the greater tendency of this module's genes to have
peripheral localization, compared with background genes. The module genes are enriched in
regulation of axonogenesis (**Figure 3E**) [32] [33] [34], while the module cells, mostly neurons
(50% inhibitory and 30% excitatory, **Figure 3D**) are marked by genes associated with cell
adhesion (**Figure 3F**), a key process during axonogenesis. The peripheral localization pattern

defining the module suggests local translation of proteins that localize in neurites (cell delineations
mostly capture somata [35]) or extracellular matrix (ECM). Consistent with this speculation, RGS5
protein is known to exhibit strong synaptic localization [36] [37], while FN1 protein is an
extracellular matrix component [38], supporting the possibility of local translation. Considering the
module's statistical association with cell-adhesion functions, we examined the transcript
distribution in adjacent pairs of cells and observed that the peripheral localization tends to be at
the facing boundaries (**Figure 3G-I**, **Figure S3**), a phenomenon not seen in non-module cells.
Putting together the different facets of this module's characterization by CellSP, we hypothesize
that the four genes are locally translated into proteins that localize at soma boundaries and
perform adhesion-related functions. This may be the case if the module cells, mostly neurons, are
actively forming synapses, undergoing neurite outgrowth, etc. [39], which is plausible considering
that the preoptic region profiled through these data is an important site of developing neurites
[40].
228
Subcellular spatial patterns recur across cell types and regions of brain
We next analyzed a large ST data set that profiles a whole mouse brain using Xenium technology
[41]. This profiles a panel of 248 genes in 162,033 cells grouped into 50 clusters based on their
transcriptomes (**Figure S4**)**.** We used CellSP to analyze each cell cluster separately, as large
expression differences across clusters may confound the discovery of subcellular pattern
modules, and because we sought to test module reproducibility across varying cellular contexts.
CellSP identified ~22 modules per cell cluster, each module comprising 8 genes and 68 cells on
average (**Data S2**). Some of the prominent biological characterizations (GO terms) reported for
these modules included synaptic processes, myelination, hormone receptor activity, cellular
responses to external stimuli, extracellular components, secretory processes and inter-cellular
communication.
240
Closer inspection of these results revealed that modules with certain biological characterizations
recur in multiple cell clusters (cell types and/or regions) of the brain. For instance, we observed
36 modules, across 18 different clusters (**Data S3, Figure S6**) to be associated with myelination
process, i.e., CellSP reported the GO term "ensheathment of neurons" as enriched (nominal p-
value < 0.01) in module genes as well as in marker genes of module cells, for each of these
modules. There is substantial overlap in their constituent genes (**Figure 4C**), with the three most
frequently recurring constituents being the myelin-associated genes Gjc3, Sox10 (SRY-box
transcription factor 10) and Opalin (oligodendrocytic myelin paranodal and inner loop protein) [42]
[43] [44]. Gjc3 is known to be expressed in myelinating glial cells [31], Sox10 is a transcription
factor responsible for the activation of several myelin-specific genes [45] and Opalin is known to
promote oligodendrocyte differentiation and axon myelination [46]. We confirmed that the
inclusion of a gene in a module is not a mere reflection of its expression levels in module cells
(**Figure S5A**), though higher expression does play a role (**Figure S5B**), in that a gene that is less
expressed in a cell is less likely to have its subcellular pattern, if any, detected by a statistical
procedure. These myelination-associated modules comprise 3493 cells overall, which are
observed across clusters with diverse transcriptomic profiles (**Figure 4A**) and in different brain
regions (**Figure 4B**). The shared biological properties of these modules are further elucidated by
the numerous other GO terms enrichments common to them (**Figure 4D,E**), including
mesenchymal stem cell (MSC) differentiation and oligodendrocyte differentiation. MSCs have the
potential to differentiate to or stimulate the maturation of oligodendrocytes [47], the primary
myelinating cells.
262
Another prominent theme in the whole brain CellSP results was the repeated finding of modules
characterized by "GABA-ergic synapse", a GO subcellular localization term. Twenty modules
across 14 different cell clusters were reported to have this annotation enriched (p-value < 0.0001)

in their constituent genes as well as in expression markers of their member cells (**Figure S7A,B**).
These modules were observed in different expression contexts (**Figure 4F**) and physical regions
of the brain (**Figure 4G**), and to exhibit different types of subcellular patterns (**Figure S9, Data**
S6). Several genes appear repeatedly as members of these modules, with the most frequent ones
being Gad1 (glutamate decarboxylase 1), Gad2 (glutamate decarboxylase 2) and Rab3b (RAB3B,
member RAS oncogene family) (**Figure S7C, Figure S8**), which play critical roles in maintaining
basal levels of GABA (Gamma-aminobutyric acid), as well as synthesis and vesicle release of
GABA in an activity-dependent manner at inhibitory synapses [48] [49] [50]. There is evidence
suggesting that GAD2 and RAB3B proteins are localized in presynaptic terminals [49] [51], hinting
at an explanation for the subcellular colocalization of their transcripts.
276
CellSP-detected modules can be specific to biological conditions
The analyses above show us that gene-cell modules recovered by CellSP reveal subcellular
transcriptomic phenomena related to biological processes such as myelination and
axonogenesis. This raises the natural question: do these modules also reveal subcellular
phenomena that vary under different tissue conditions? To investigate this, we used CellSP on
data sets comprising case-control pairs of tissues. The first such analysis involved Xenium data
on a kidney cancer (papillary renal cell carcinoma) FFPE sample (**Figure S10A**) and a healthy
kidney sample [52]. As in the whole-brain analysis above, we first clustered the cells of each
sample based on their transcriptomes and analyzed each cell cluster with CellSP, identifying 11
modules in the cancer sample and 15 modules in the healthy sample (**Data S5, S6**). The cancer
modules were mostly (10/11) found in two of the 18 cell clusters (**Figure 5A, Figure S10B**) –
Cluster 6 and Cluster 9. Six of these cancer modules comprise genes and cells characterized by
CellSP as being immune response-related, with enrichment of GO terms such as "immune
response", "defense response", "T cell activation", "positive regulation of leukocyte proliferation",
etc. (**Data S5**).
292
Examination of the 15 modules in the healthy sample (**Data S6**) suggested that the immune
response-related modules found in the cancer sample were specific to it and were not present in
the healthy sample. To pursue this observation objectively, we used CellSP visualization routines
to contrast a module's spatial pattern in the module cells (from the cancer sample) versus cells in
the other (healthy) tissue sample. As condition-specific gene expression can be a confounder for
such analysis, this analysis used a subset of healthy tissue cells that matched the cancer module
cells in the expression levels of module genes (**Methods**). **Figures 5B-D** show the cancer-versus-
healthy tissue comparison for three of the seven cancer-associated immune-response modules.
For instance, module C6_M5_S (**Figure 5B**) comprises a set of 31 genes that are localized
centrally in a subset of 22 cells (**Figure S10C**) and was detected in Cluster 6 of the cancer sample
(**Figure 5A**). The genes are enriched for the GO term "defense response" (p-value 8.9E-08). As
shown in **Figure 5B** (right), the same set of genes lacks this centrally localized distribution of
transcripts in expression-matched cells from the healthy sample. A second example is the module
C6_M0_S (**Figure 5C**), a set of 24 genes, also enriched in "defense response" (p-value 3.3E-04),
that exhibit radial subcellular localization in a set of 18 cells (**Figure S10D**) of Cluster 6. CellSP
visualization (**Figure 5C,** left) confirms that the spatial pattern is significantly different between
module genes and other genes when examining the module cells. The adjacent panel (**Figure**
5C, right) shows that this prominent contrast is not seen in expression-matched cells from healthy
tissue, supporting the cancer-specificity of this module. The third cancer-specific module we
highlight here is C6_M3_I (**Figure 5D**), a set of seven genes enriched in the term "T cell
activation", that exhibit significant subcellular colocalization in 66 cells (**Figure S10E**) of Cluster
6. The strength of its spatial pattern in cancer versus healthy tissue is visualized in **Figure 5D**,
where the top left quadrant of the heatmap indicates that the module genes have a stronger
tendency for pairwise colocalization in the module (cancer) cells compared to expression-

matched non-module cells from the healthy sample. (The other quadrants, especially the bottom-
right, show that this contrast is not observed for a random subset of non-module genes.) In
summary, CellSP discovers immune-response related gene-cell modules representing significant
subcellular patterns that are specific to cancer versus healthy tissue.
321
CellSP detects subcellular patterns associated with Alzheimer's Disease in a mouse
model
We next used CellSP to identify gene-cell modules in brains of mouse models of Alzheimer's
Disease (AD). For this, we analyzed Xenium data on a coronal section of one hemisphere of
TgCRND8 transgenic male mouse brain at three time points (pathological progression stages),
as well as wild-type (WT) mouse brain at similar time points [53]. Following the same procedure
as above, we clustered cells in each brain sample and used CellSP to find significant modules in
each cell cluster, recovering ~11 gene-cell modules per cluster on average (**Figure S11**, **Data**
S7). To contextualize these modules with the AD genetics literature, we focused on six genes
from the Xenium gene panel implicated in AD risk [54] (**Figure 5E**). We observed the gene Picalm
(phosphatidylinositol binding clathrin assembly protein) to appear in many modules in the AD
mice, especially in the early and middle time points, significantly more frequently than in WT mice
(t-test p-value 0.047) (**Figure S12**). The Picalm protein is responsible for recruiting clathrin and
adaptor protein-2 (AP-2) to the plasma membrane [55] as part of clathrin-mediated endocytosis,
and contributes to clearance of amyloid-β (Aβ) at the blood brain barrier [56]. It is linked to
processes that are disrupted in AD and is also genetically associated with the disease [57]. We
thus speculate that the observed subcellular patterns involving this gene are reflections of the
gene's dysfunction in AD mice. We observed another AD risk gene – Trem2 (triggering receptor
expressed on myeloid cells 2) – to feature in many CellSP modules (12) (**Figure S13**) in the late-
stage AD mouse brain, significantly more than in the other five samples from WT or AD mice. This
late stage is associated with increase in AD-associated microglial population [58], and Trem2 is
primarily expressed in microglia, where it functions as a transmembrane receptor [59] enabling
the progression to a mature disease-associated microglia phenotype [58] and is involved in the
response to Aβ plaques [60]. Variants in Trem2 gene have been identified as a significant risk
factor for late-onset AD [61], suggesting again that the observed gene-cell modules reveal
subcellular phenomena reflecting its dysfunction. The four other AD risk genes were found in few
modules overall (**Figure 5E**) and these modules were of similar counts in AD vs. WT mice. We
also note that the above AD-associated changes involving Picalm and Trem2 expression,
observed at the subcellular spatial level, would not have been detected at the level of overall
cellular expression (**Figure S14**).
352
We next examined the CellSP-detected modules in AD and WT brains with a focus on myelination.
Given the reported connection between myelin damage and AD pathology [62]; and building on
our earlier identification of myelination-related gene-cell modules in the mouse brain, we
investigated whether such modules have any unique characteristics in one phenotypic group
compared to the other. We concentrated on all identified modules associated with the GO term
"ensheathment of neurons" (nominal p-value < 0.01) and, to identify any characteristics that
differentiate these modules between AD and WT, we examined the frequency of myelination-
related genes co-occurring in the same module (**Figure 5F**). While most gene pairs show similar
co-occurrence statistics between the two groups, the pairs Olig2 (oligodendrocyte transcription
factor 2)-Plp1 (proteolipid protein 1) and Olig2-Mobp have a significantly higher co-occurrence in
WT brain compared to AD brain (proportion test p-value < 0.05). (Also see **Data S8**). To follow up
on these intriguing observations involving Olig2, a gene essential to maturation of
oligodendrocytes (the cells responsible for producing myelin), we next identified all genes that co-
occur with Olig2 in a group-specific manner (proportion test p-value < 0.05). (The previous
analysis was limited to eight select myelination-related genes, including Olig2.) We found 10 such

genes (**Figure 5G**, blue bars), nine of which co-occur with Olig2 preferentially in gene-cell
modules of the WT brains, and consist of the above-noted genes Plp1, Mobp as well as other
genes linked to myelination, including Opalin and Cnp (2',3'-cyclic nucleotide 3'
phosphodiesterase) [46] [63] [64] [65]. This inter-group difference in module composition is most
in time points 1 and 2 (**Data S9**) which correspond to periods of continued myelination in the mice
brain [66]. These observations suggest that the integrity and functionality of subcellular modules
found in WT brains may be impaired in AD brains, reflecting a broader dysfunction in myelination
processes in AD [67]. In summary, our myelination-focused examination of CellSP modules points
to a potential significance of Olig2 in Alzheimer's disease pathology, highlighting a shift from
normal myelination functions in healthy brains.
378
The specific relationships and insights uncovered here are a glimpse into the dynamics of the
subcellular transcriptome during AD progression, not merely a reflection of differential co-
expression of genes at the whole cell level. To substantiate this, we repeated the module
discovery using co-expression analysis with the popular WGCNA tool [21]. Modules were
detected for each cell cluster in each of the six brain samples, as above, except that these were
gene co-expression modules rather than CellSP-identified gene-cell modules. These co-
expression modules do not show any preferential co-occurrence of Olig2 with any of genes
identified above except one – Olig2-Gatm (glycine amidinotransferase) (p-value < 0.05, **Figure
5G**, orange bars). This analysis underscores the ability of CellSP modules to reveal subcellular
pattern changes between phenotypic conditions, that may not be discernible at the cellular level.
389
390

DISCUSSION
With the increasing popularity of spatial transcriptomics (ST) techniques of single-molecule
resolution [68] [69] [70] new tools have been proposed to identify genes and gene pairs that exhibit
interesting transcript distribution patterns within cells [18] [19] [13]. However, there remain
fundamental and practical challenges not addressed by these analysis tools. A practical problem
is that they annotate subcellular patterns such as gene-gene colocalization or gene localization
preferences in individual cells, leading to very large compendia of significant patterns that are
difficult to sift through for actionable insights. A conceptual limitation is that while they can
effectively shortlist genes that merit further exploration on account of their subcellular
distributions, they do not highlight subsets of cells where interesting spatial phenomena manifest
and could point to special functional properties of those cells. The simple approach of examining
every cell that exhibits a statistically significant spatial pattern is impractical, as a very large
number of cells harboring diverse patterns of varying functional origins are enumerated in this
way. On the other hand, examining all cells where a particular gene (or pair) has the same
subcellular distribution pattern will demand that we repeat such examination for a large number
of genes, again leading to a substantial interpretive burden.
407

The key innovation of CellSP is the idea of a "gene-cell module", which is one solution to the
above shortcomings of existing tools. Such a module, if found to be statistically significant, draws
our attention to a subset of cells that share a subcellular spatial pattern involving the same subset
of genes, suggesting a functional commonality among those cells. Furthermore, CellSP
automatically searches for that functional characterization by using machine learning to identify
gene markers of the member cells of the module, and reporting biological properties enriched in
the marker genes. This is how our analysis of MERFISH data on hypothalamic preoptic area
uncovered a collection of 332 cells, mostly mature oligodendrocytes, that appear to be involved
in myelination, and another subset of 221 cells, mostly neurons, that are characterized by
axonogenesis-related functions. These discoveries would not have been possible with existing

tools. In identifying a gene-cell module, CellSP not only highlights an interesting subset of cells,
it simultaneously draws our attention to a set of genes that share an interesting transcript
distribution pattern in those cells, prompting functional hypotheses involving those spatial
patterns. Such gene set discovery is a standard technique of transcriptomics analysis, e.g., as
co-expression module identification [21] and a CellSP module extends this time-tested concept
to subcellular colocalization of genes. It was this functionality that led to the identification of the
four-gene module comprising myelination-related genes (Sgk1, Ttyh2, Ermn and Ndrg1) that
colocalize with each other in many mature oligodendrocytes, and another module comprising
axonogenesis-related genes (Fn1, Slco1a4, Rgs5, Sema3c) whose transcripts localize at the
periphery in the same collection of cells, raising the possibility of localized translation of these
genes as part of cell adhesion processes.
429
We note that the InSTAnT toolkit [18] provides module discovery functionality, where a module
has a different statistical interpretation from that adopted in CellSP – it is a subset of genes that
colocalize with each other significantly frequently across the entire population of cells. We
compared CellSP to the gene module discovery methods implemented in InSTAnT (**Note S1**) and
found that the former offers significant advantages in terms of biological interpretation and ease
of use. Importantly, CellSP-reported modules include four additional types of subcellular spatial
patterns (radial, peripheral, punctate and central patterns discovered by SPRAWL [19]) and are
accompanied by innovative visualizations and machine learning-based characterization of
module cells, features that are crucial to the scientific discovery process involving functional
subcellular patterns.
440
We recognize that certain aspects of CellSP can be further improved in future versions. For
example, the heuristic nature of the LAS search algorithm and our module merging step may
result in variability across different runs, and alternative search algorithms should be explored to
address this. As another example, the scheme for visualizing subcellular idealize cells as unit
circles, which may not accurately depict elongated or irregularly shaped cells; techniques for
average shape construction [71] may be fruitful for future work in this direction. Finally, the
comparison of gene-cell modules between conditions is not implemented as a statistical tool in
CellSP and was performed by us through downstream processing of its results; such a tool for
"differential module discovery" will be another important goal of future work.
450
By analyzing data sets from diverse organs and tissues, such as whole brain, specific brain
regions, kidney, etc., under various biological conditions such as Alzheimer's disease models or
kidney cancer, and different technologies such as MERFISH and Xenium, we demonstrated how
CellSP can be used to explore large and high-dimensional ST data and extract new insights into
subcellular transcript distribution. The discovered patterns are suggestive of co-transportation or
co-localization of RNAs that participate in similar biological functions [72], and also support the
phenomenon of local translation [73] which requires the mRNA to be transported to precise
locations within the cell where they are translated into proteins. Furthermore, we showed how
gene-cell modules can be compared across conditions to discover subcellular changes
associated with disease states, which are not recover-able using traditional differential expression
analysis. Overall, CellSP offers a powerful new approach to subcellular spatial transcriptomics
data analysis.
463
464
465
466 **METHODS**
467

CellSP module discovery algorithm

CellSP identifies persistent subcellular spatial phenomena in the tissue by defining "gene-cell modules" as sets of genes with specific subcellular transcript spatial patterns across cells. This process is performed in two steps -

Subcellular pattern discovery. We utilize the existing tools InSTAnT [18] and SPRAWL [19] for subcellular pattern detection in individual cells. InSTAnT detects gene pair colocalization in single cells, employing the "Proximal Pairs" (PP) test to calculate colocalization p-values for each gene pair in every cell. It also assigns a global colocalization p-value to each gene pair, via the "Conditional Poisson Binomial" (CPB) test. We use a user-tunable threshold of $1e^{-3}$ on this p-value to limit the set of candidate gene pairs. A matrix M_I is constructed to store the PP test p-values with dimensions $n_{cells} \times n_{genepairs}$ where $n_{genepairs}$ is the number of candidate gene pairs. We then perform a negative log transformation on this matrix.

SPRAWL detects subcellular mRNA localization patterns and classifies them into four categories - peripheral, central, radial, and punctate. Each gene receives a statistical significance score (on a scale of -1 to 1) representing the strength of the spatial pattern in each cell. These scores are stored in four separate matrices $M_{peripheral}$, $M_{central}$, M_{radial}, $M_{punctate}$ each with dimensions $n_{cells} \times n_{genes}$.

Module discovery. Each entry of the matrices obtained from InSTAnT and SPRAWL represents the strength of pattern occurrence for each gene/gene-pair in each cell. To aggregate these patterns across genes and cells, we use the LAS [23] algorithm to perform biclustering on each of these matrices. The LAS algorithm iteratively searches for a submatrix (bicluster) with significantly high average value, removes the influence of selected submatrix from the original data matrix, and repeats the process until the desired number of submatrices are retrieved or no significant submatrices are found. The LAS significance score for a $k \times l$ submatrix U within a data matrix X of dimensions $m \times n$ is defined as

$$S(U) = -\log\left[\binom{m}{k}\binom{n}{l}\Phi\left(-\tau\sqrt{kl}\right)\right]$$

where $\Phi(\cdot)$ denotes the cumulative distribution function (CDF) of the standard normal distribution, and τ denotes the average value of the submatrix U.

The reported submatrices are what we call "gene-cell modules" or "modules". Due to the heuristic nature of the search, it is possible that a reported module is part of a larger module with a similarly high average value, but this larger module is not discovered. To mitigate this issue across reported modules, we implement a two-step module expansion process:

1. Cell Expansion – For each module, we iterate through all cells and add a cell to the module if the cell's average value across the module's genes exceeds the original module's average value.
2. Module Mergers – Iterating over modules in descending order of significance score, we calculate the overlap coefficient for the genes of each module pair. The overlap coefficient is defined as

$$Overlap(A, B) = \frac{|A \cap B|}{min(|A|, |B|)}$$

514
where A and B denote the gene sets of the two modules. If the overlap coefficient exceeds
0.667, then we consider merging the module pair by creating a new module that comprises
the union of genes/gene pairs of the original pair and the union of cells of the original pair.
If the new module's significance score surpasses the module significance threshold, the
merged module is retained, and the original pair of modules is removed. This process is
repeated iteratively, prioritizing the merger of highly significant modules first, until no
further mergers are possible. This merger step is useful to reduce the number of modules
detected and combine modules of similar composition (high overlap in constituent genes)
into one. This step is made optional.
524
525   The module discovery process (including module expansion) is conducted independently for
526   modules derived from each of the five matrices $M_I, M_{peripheral}, M_{central}, M_{radial}, M_{punctate}$ that
527   represent different kinds of subcellular patterns.
528
In the final reported list of gene-cell modules, two modules may comprise overlapping sets of cells
and genes, allowing the same cells and genes to be part of multiple modules (**Figure 1**). Modules
detected from the SPRAWL-derived matrices $M_{peripheral}, M_{central}, M_{radial}, M_{punctate}$ are
reported as one group (their module identifiers have the suffix "_S" and a unique numeric prefix),
while modules detected from the InSTAnT-derived matrix M_I are reported as another group
(identifiers with suffix "_I" and a unique numeric prefix).
535
536   **Module Characterization**
537   To investigate factors (other than cell types) that underlie or are associated with subcellular spatial
538   phenomena, CellSP performs a two-level characterization of the detected modules using gene
539   set enrichment analysis and predictive modeling.
540
541   Module Gene Characterization: The biological functions associated with module genes are
542   characterized using PantherDB [74] for gene enrichment analysis. For each module, the module
543   genes are used as the query gene set, while the gene panel of the ST assay serves as the
544   background set. This analysis identifies pathways, biological processes, and molecular functions
545   enriched in module genes.
546
Module Cell Characterization: To characterize the cells in each module, CellSP trains a classifier
to distinguish between module cells (positive set) and all other cells (negative set). Each cell's
gene expression profile (total transcript count of each gene in that cell) is used as the cell's feature
vector, i.e., this is not a spatially resolved featurization. Since an ST assay of single-molecule
resolution typically profiles a limited number of genes, the number of features is relatively modest.
To address this, CellSP uses the Tangram tool [75] to impute the expression of additional genes
in each cell of the ST data using scRNA-seq data of the same tissue, if available. The imputation
process expands the gene set significantly, increasing the number of features from a few
hundreds (in the original ST data) to several thousands (as available in scRNA-seq data). We
construct an extended gene expression matrix by retaining the original expression values for
genes present in the ST panel and incorporating the imputed expression values for genes absent
from the panel. The extended gene expression matrix is used to train a Random Forest classifier
of module cells. The top 20 most informative genes are identified based on their feature
importance, which is quantified using SHAP [76] [77]. These genes, along with any other genes
that exhibit a high Pearson correlation (r > 0.98) with them, are designated as "marker genes" of
the module cells. These genes are then subjected to gene set characterization using PantherDB

to identify pathways, biological processes, and molecular functions that characterize the module's
cells.
565
Module Pattern Visualization
567   To help visualize modules defined by the five types of subcellular spatial patterns (four types
568   identified by SPRAWL and the colocalization pattern identified by InSTAnT), we developed three
569   complementary plotting techniques.
570
SPRAWL detects localization patterns (peripheral, central, radial or punctate) for each gene in
each cell. To aggregate these spatial localization patterns across many cells, CellSP transforms
each cell into a uniform representation within a unit circle. This transformation is achieved using
the smallest enclosing circle algorithm to identify the smallest circle enclosing all transcripts in the
cell. The circle is then centered at the origin and scaled to have a unit radius. For "central" and
"peripheral" patterns, the unit circle is divided into C (default value of 5) concentric rings, and the
proportion of module gene transcripts within each ring is calculated. The transcript abundance in
a ring is then averaged across all cells of the module, and the ring is displayed with color shade
depicting the average abundance. A similar visualization is constructed for all non-module cells
and the two plots are placed side-by-side. For modules displaying "central" patterns, we expect
higher abundance in the innermost rings, while "peripheral" patterns are anticipated to exhibit
higher abundance in the outermost rings.
583
For modules defined by "punctate" or "radial" patterns, the unit circle representing an idealized
view of a cell is divided into S (defaults to 10) sectors. Gene transcript density is computed in each
sector for module genes. To account for the directional variability of these patterns, each cell's
idealized circle view is rotated such that the sector with the maximum density aligned with 0°.
Following this alignment, densities are aggregated across cells by calculating mean densities for
each sector. This results in a density distribution across sectors, with the highest density in the
first sector by construction. This calculation is then repeated for non-module genes, and the
density distributions (across sectors) for module genes and non-module genes are plotted
overlaid on each other, in different colors. In this framework, module genes are expected to
concentrate primarily in the first sector, while non-module genes are anticipated to exhibit more
uniform distributions across sectors, despite having the highest density in the first sector.
595
InSTAnT detects colocalized gene pairs as those whose transcripts are in close proximity (within
distance d) of each other. To visualize such patterns aggregated over cells, CellSP first constructs
a d-radius neighborhood graph of transcripts for each module cell (where d equals the distance
threshold parameter used in InSTAnT runs), calculates the number of neighboring transcript pairs
for each pair of module genes and averages these counts across all module cells, thus obtaining
a "proximity score" for that gene pair aggregating information across module cells. This process
is then repeated over non-module cells to obtain a proximity score of the same gene pair but now
aggregating information from non-module cells. The proximity enrichment score of the gene pair
is then defined as the log ratio of proximity scores from module cells and non-module cells. High
positive values of the proximity enrichment score indicate that the gene pair exhibits a greater
degree of colocalization in module cells compared to non-module cells. To add further contrast,
a randomly selected set of non-module ("control") genes, equal in count to the module genes, is
included in the visualization: a heatmap is constructed whose rows and columns represent
module genes (top half of rows and left half of columns) and the selected control genes (bottom
half of rows and right half of columns), and values depict proximity enrichment scores of gene
pairs. The upper-left quadrant corresponds to pairs of module genes, the lower-right quadrant to
pairs of control genes, and the remaining two quadrants correspond to a module gene paired with
a non-module gene.

Gene Set Enrichment Visualization

CellSP provides the ability to visualize the gene set enrichment reports generated by PantherDB
using the Revigo tool [24]. Revigo summarizes lists of GO terms by clustering them based on
their semantic similarity, identifying representative terms. These terms are visualized as circles in
a scatterplot, where the circle size reflects the gene set size and the color intensity indicates
statistical significance, with darker colors representing higher significance. This visualization
highlights the importance, similarity, and uniqueness of terms, making it easier to interpret
enrichment results.
622

We used a special type of Revigo plots to aggregate gene set enrichment reports across multiple
detected modules (**Figure 4D,E**; this is not a standard functionality of CellSP). We first collected
all GO terms with a module association at a p-value below 0.01, across the modules, along with
their p-values. This aggregated list of significant terms was then processed through Revigo, which
calculated the principal component analysis (PCA) values for each term. Using these PCA
coordinates, we generated a 2D histogram to visualize the aggregated GO terms across modules.
The PCA values determined the positions of the bins in the plot and the number of modules each
term was associated significantly with was represented by the color intensity of the corresponding
bin. This visualization highlights the most commonly enriched biological themes across modules
while preserving the semantic relationships among GO terms.
634

Cross-Condition Module Comparison

Human Kidney Dataset: To compare the detected modules between conditions in the Human
Kidney dataset, we assessed whether the subcellular patterns of modules from the cancer
condition were present in the control tissue. We first performed min-max normalization of gene
expression values for both the control and cancer datasets independently. For a gene-cell module
detected in the cancer dataset, we calculated the total gene expression of module genes in each
module cell and determined their minimum P_{min} and maximum P_{max} as percentiles of total gene
expression in the population of all cells in the cancer dataset. Using these percentiles, we
identified cells from the control dataset that have total gene expression (of module genes)
between P_{min} and P_{max} percentile of the population of control cells. Thus, we identified a subset
of the control cells whose total expression is similar to the module cells (which belong to the
cancer data set). Subcellular patterns of module genes were then examined in these control cells
and compared to the patterns in the module cells from the cancer dataset.
647

Mouse Alzheimer's Disease Dataset: To identify changes in functionally similar (myelination-
related) modules across conditions (AD model mice and WT mice), we asked whether two sets
of modules (one set from each condition) differ in their gene composition. One way to answer this
is based on how many modules in each condition include a specific gene, and whether the gene's
frequency of inclusion is different between conditions. This approach was used for **Figure 5E**. A
complementary approach is based on counting how many modules in each condition include a
specific gene pair and comparing the proportion of such modules between conditions using a one-
sided proportion test [78]. The test statistic for the proportion test is given by

$$z = \frac{\widehat{p_1} - \widehat{p_2}}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{n_1}+\frac{1}{n_2}\right)}}$$

658 where $\widehat{p_1}$ is the proportion for group 1, $\widehat{p_2}$ is the proportion for group 2, $\hat{p}$ is the pooled proportion
659 and n_1, n_2 are the sample sizes for group 1 and 2 respectively.
660

Comparison with traditional analysis methods

To compare insights generated by our approach against traditional methods, we performed differential expression analysis and co-expression network analysis on the datasets. For differential expression analysis, we utilized Scanpy [79] to apply the Wilcoxon rank-sum test, comparing gene expression between the two conditions under investigation. This provided insights into genes with significantly altered expression levels across conditions.

For co-expression network analysis, we employed WGCNA [21] using the PyWGCNA framework [80]. Similar to our approach using CellSP, WGCNA was run independently for each cell cluster in the dataset. WGCNA assigns a module identity (class label) to each gene within a cluster based on its co-expression patterns. We used these module identities to calculate the frequency with which a gene pair is assigned to the same WGCNA module across the dataset. These frequencies were then compared between the conditions using a one-sided proportion test to assess condition-specific co-expression relationships.

CellSP User Guide

CellSP provides flexible and tunable parameters that allow users to adapt analyses to their specific datasets and objectives. For InSTAnT, the distance threshold (d) is typically set to approximately 5% of the average cell diameter. The significance threshold (α) for the CPB p-values is set to $1e^{-5}$ by default. If this results in too few significant gene pairs (< 250), we recommend relaxing the threshold to $1e^{-3}$ or lower. To manage computational complexity, an additional parameter, K, allows users to select the top K gene pairs for further analysis. For SPRAWL, we adapted the original scripts into CellSP to enable parallelization and use the default parameters. Similarly, LAS scripts from the implementation available in biclustlib [81] were parallelized and integrated into CellSP. LAS includes two primary adjustable parameters: N and RS. The parameter N determines the number of modules to search and is set to a default value of 10. We recommend keeping this parameter between 5 to 50. The parameter RS specifies the number of randomized searches performed to enhance the quality of the biclusters. By default, RS is set to 50,000; however, for larger datasets (e.g., > 200 genes or > 10,000 cells), we recommend reducing this value to balance computational demands while maintaining reasonable bicluster quality. Visualization in CellSP is also highly customizable. Users can fine-tune the representation of subcellular patterns using parameters such as the number of concentric circles (C), the number of sectors (S), and the distance threshold (d). These settings provide flexibility in highlighting key spatial patterns. Details for the specific parameters used in each analysis can be found in **Data S10**. On a dataset containing 10,000 cells and 248 genes, CellSP completes its analysis in approximately 3 hours. Additional runtime details can be found in **Note S2** and **Data S13.**

DATA AVAILABILITY

The MERFISH dataset for the Mouse Preoptic Hypothalamus region [26] was obtained through direct communication with Dr. Jeffrey Moffitt. After excluding cells labeled with ambiguous cell types, the dataset comprises 5,149 cells distributed across 9 distinct cell types, with a gene panel of 135 genes.

The Xenium datasets are publicly available at 10x Genomics Datasets. We used the "Fresh Frozen Mouse Brain for Xenium Explorer Demo" [41], "Human Kidney Preview Data" [52], and the "Xenium In Situ Analysis of Alzheimer's Disease Mouse Model Brain Coronal Sections from One Hemisphere Over a Time Course" [53] datasets for the mouse brain, human kidney, and

Alzheimer's disease experiments, respectively. The Mouse Brain dataset contains 162,033 cells grouped into 50 clusters based on the expression profiles of 248 genes. The Human Kidney dataset comprises 56,509 cells grouped into 19 clusters in the cancer tissue and 97,546 cells grouped into 21 clusters in the control tissue. The gene panel includes 377 genes. The Mouse Alzheimer's dataset includes six tissue samples from two conditions—Wild Type and Alzheimer's (TgCRND8 mouse model)—at three timepoints (2.5 months, 5.7 months, and 13+ months). The Alzheimer's samples contain 53,908, 58,681, and 61,435 cells across the timepoints, while the Wild Type samples contain 58,230, 58,685, and 59,933 cells, with a gene panel of 347 genes.

CODE AVAILABILITY

CellSP is open source and available at https://github.com/bhavaygg/CellSP.

ACKNOWLEDGEMENTS

We thank Abhishek Ojha for his help in designing the statistical tests. This research utilized resources from PACE at the Georgia Institute of Technology, Atlanta, USA. Funding: This work was supported by the National Institutes of Health (R35GM131819 to S.S.), and Georgia Institute of Technology (Wallace H. Coulter Distinguished Faculty Chair: S.S.)

731 **REFERENCES**

[1] K. R. Maynard, L. Collado-Torres, L. M. Weber, C. Uytingco, B. K. Barry, S. R. Williams, J. L. Catallini, M. N. Tran, Z. Besich, M. Tippani and others, "Transcriptome-scale spatial gene expression in the human dorsolateral prefrontal cortex," *Nature neuroscience,* vol. 24, p. 425–436, 2021.

[2] B. Watson, B. Paul, L. Amir-Zilberstein, A. Segerstolpe, R. Rahman, A. Shih, J. Deguine, R. Xavier, J. R. Moffitt and A. C. Mullen, "Spatial transcriptomics of healthy and fibrotic human liver at single-cell resolution," *bioRxiv,* p. 2024–02, 2024.

[3] P. Cadinu, K. N. Sivanathan, A. Misra, R. J. Xu, D. Mangani, E. Yang, J. M. Rone, K. Tooley, Y.-C. Kye, L. Bod and others, "Charting the cellular biogeography in colitis reveals fibroblast trajectories and coordinated spatial remodeling," *Cell,* vol. 187, p. 2010–2028, 2024.

[4] H. Shi, Y. He, Y. Zhou, J. Huang, K. Maher, B. Wang, Z. Tang, S. Luo, P. Tan, M. Wu and others, "Spatial atlas of the mouse central nervous system at molecular resolution," *Nature,* vol. 622, p. 552–561, 2023.

[5] E. N. Farah, R. K. Hu, C. Kern, Q. Zhang, T.-Y. Lu, Q. Ma, S. Tran, B. Zhang, D. Carlin, A. Monell and others, "Spatially organized cellular communities form the developing human heart," *Nature,* vol. 627, p. 854–864, 2024.

[6] M. Zhang, X. Pan, W. Jung, A. R. Halpern, S. W. Eichhorn, Z. Lei, L. Cohen, K. A. Smith, B. Tasic, Z. Yao and others, "Molecularly defined and spatially resolved cell atlas of the whole mouse brain," *Nature,* vol. 624, p. 343–354, 2023.

[7] C. E. Holt and S. L. Bullock, "Subcellular mRNA localization in animal cells and why it matters.," *Science (New York, N.Y.),* vol. 326, no. 5957, p. 1212–1216, November 2009.

[8] D. Wang, Z. Zhang, Y. Jiang, Z. Mao, D. Wang, H. Lin and D. Xu, "DM3Loc: multi-label mRNA subcellular localization prediction and analysis based on multi-head self-attention mechanism," *Nucleic acids research,* vol. 49, p. e46–e46, 2021.

[9] T. Schuster, A. Amoah, A. Vollmer, G. Marka, J. Niemann, M. Saçma, V. Sakk, K. Soller, M. Vogel, A. Grigoryan and others, "Quantitative determination of the spatial distribution of components in single cells with CellDetail," *Nature Communications,* vol. 15, p. 10250, 2024.

[10] J. Barr, K. V. Yakovlev, Y. Shidlovskii and P. Schedl, "Establishing and maintaining cell polarity with mRNA localization in Drosophila.," *BioEssays : news and reviews in molecular, cellular and developmental biology,* vol. 38, no. 3, p. 244–253, March 2016.

[11] M. Dermit, M. Dodel, F. C. Y. Lee, M. S. Azman, H. Schwenzer, J. L. Jones, S. P. Blagden, J. Ule and F. K. Mardakheh, "Subcellular mRNA Localization Regulates Ribosome Biogenesis in Migrating Cells.," *Developmental cell,* vol. 55, no. 3, p. 298–313.e10, November 2020.

[12] R. M. Parton, A. Davidson, I. Davis and T. T. Weil, "Subcellular mRNA localisation at a glance," *Journal of cell science,* vol. 127, p. 2127–2133, 2014.

[13] C. K. Mah, N. Ahmed, N. A. Lopez, D. C. Lam, A. Pong, A. Monell, C. Kern, Y. Han, G. Prasad, A. J. Cesnik and others, "Bento: a toolkit for subcellular analysis of spatial transcriptomics data," *Genome Biology,* vol. 25, p. 82, 2024.

[14] K. H. Chen, A. N. Boettiger, J. R. Moffitt, S. Wang and X. Zhuang, "Spatially resolved, highly multiplexed RNA profiling in single cells," *Science,* vol. 348, p. aaa6090, 2015.

[15] C.-H. L. Eng, M. Lawson, Q. Zhu, R. Dries, N. Koulena, Y. Takei, J. Yun, C. Cronin, C. Karp, G.-C. Yuan and others, "Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH+," *Nature,* vol. 568, p. 235–239, 2019.

[16] X. Wang, W. E. Allen, M. A. Wright, E. L. Sylwestrak, N. Samusik, S. Vesuna, K. Evans, C. Liu, C. Ramakrishnan, J. Liu, G. P. Nolan, F.-A. Bava and K. Deisseroth, "Three-dimensional intact-tissue sequencing of single-cell transcriptional states.," *Science (New York, N.Y.),* vol. 361, no. 6400, July 2018.

[17] K. Dong and S. Zhang, "Deciphering spatial domains from spatially resolved transcriptomics with an adaptive graph attention auto-encoder," *Nature communications,* vol. 13, p. 1739, 2022.

[18] A. Kumar, A. W. Schrader, B. Aggarwal, A. E. Boroojeny, M. Asadian, J. Lee, Y. J. Song, S. D. Zhao, H.-S. Han and S. Sinha, "Intracellular spatial transcriptomic analysis toolkit (InSTAnT)," *Nature communications,* vol. 15, p. 7794, 2024.

[19] R. Bierman, J. M. Dave, D. M. Greif and J. Salzman, "Statistical analysis supports pervasive RNA subcellular localization and alternative 3'UTR regulation," *eLife,* vol. 12, p. RP87517, 2024.

[20] F. C. Walter, O. Stegle and B. Velten, "FISHFactor: a probabilistic factor model for spatial transcriptomics data with subcellular resolution," *Bioinformatics,* vol. 39, p. btad183, 2023.

[21] P. Langfelder and S. Horvath, "WGCNA: an R package for weighted correlation network analysis," *BMC bioinformatics,* vol. 9, p. 1–13, 2008.

[22] A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander and others, "Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles," *Proceedings of the National Academy of Sciences,* vol. 102, p. 15545–15550, 2005.

[23] A. A. Shabalin, V. J. Weigman, C. M. Perou and A. B. Nobel, "Finding large average submatrices in high dimensional data," *The Annals of Applied Statistics,* p. 985–1012, 2009.

[24] F. Supek, M. Bošnjak, N. Škunca and T. Šmuc, "REVIGO summarizes and visualizes long lists of gene ontology terms," *PloS one,* vol. 6, p. e21800, 2011.

[25] L. McInnes, J. Healy and J. Melville, "Umap: Uniform manifold approximation and projection for dimension reduction," *arXiv preprint arXiv:1802.03426,* 2018.

[26] J. R. Moffitt, D. Bambah-Mukku, S. W. Eichhorn, E. Vaughn, K. Shekhar, J. D. Perez, N. D. Rubinstein, J. Hao, A. Regev, C. Dulac and others, "Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic region," *Science,* vol. 362, p. eaau5324, 2018.

[27] H. Wang, A. Tewari, S. Einheber, J. L. Salzer and C. V. Melendez-Vasquez, "Myosin II has distinct functions in PNS and CNS myelin sheath formation.," *The Journal of cell biology,* vol. 182, no. 6, p. 1171–1184, September 2008.

[28] H. Inouye, J. Liu, L. Makowski, M. Palmisano, M. Burghammer, C. Riekel and D. A. Kirschner, "Myelin organization in the nodal, paranodal, and juxtaparanodal regions revealed by scanning x-ray microdiffraction," *PLoS One,* vol. 9, p. e100592, 2014.

[29] L. S. Laursen, C. W. Chan and C. ffrench-Constant, "Translation of myelin basic protein mRNA in oligodendrocytes is regulated by integrin activation and hnRNP-K," *Journal of Cell Biology,* vol. 192, p. 797–811, February 2011.

[30] R. Gould and S. Brady, "Identifying mRNAs Residing in Myelinating Oligodendrocyte Processes as a Basis for Understanding Internode Autonomy.," *Life (Basel, Switzerland),* vol. 13, no. 4, April 2023.

[31] B. M. Altevogt, K. A. Kleopa, F. R. Postma, S. S. Scherer and D. L. Paul, "Connexin29 is uniquely distributed within myelinating glial cells of the central and peripheral nervous systems.," *The Journal of neuroscience : the official journal of the Society for Neuroscience,* vol. 22, no. 15, p. 6458–6470, August 2002.

[32] E. Koncina, L. Roth, B. Gonthier and D. Bagnard, "Role of semaphorins during axon growth and guidance," *Axon growth and guidance,* p. 50–64, 2007.

[33] C. Liu, Q. Hu, J. Jing, Y. Zhang, J. Jin, L. Zhang, L. Mu, Y. Liu, B. Sun, T. Zhang, Q. Kong, G. Wang, D. Wang, Y. Zhang, X. Liu, W. Zhao, J. Wang, T. Feng and H. Li, "Regulator of G protein signaling 5 (RGS5) inhibits sonic hedgehog function in mouse cortical neurons.," *Molecular and cellular neurosciences,* vol. 83, p. 65–73, September 2017.

[34] D. A. Tonge, H. T. de Burgh, R. Docherty, M. J. Humphries, S. E. Craig and J. Pizzey, "Fibronectin supports neurite outgrowth and axonal regeneration of adult brain neurons in vitro.," *Brain research,* vol. 1453, p. 8–16, May 2012.

[35] V. Petukhov, R. J. Xu, R. A. Soldatov, P. Cadinu, K. Khodosevich, J. R. Moffitt and P. V. Kharchenko, "Cell segmentation in imaging-based spatial transcriptomics," *Nature Biotechnology,* vol. 40, p. 345–354, October 2021.

[36] P. J. Thul, L. Åkesson, M. Wiking, D. Mahdessian, A. Geladaki, H. Ait Blal, T. Alm, A. Asplund, L. Björk, L. M. Breckels, A. Bäckström, F. Danielsson, L. Fagerberg, J. Fall, L. Gatto, C. Gnann, S. Hober, M. Hjelmare, F. Johansson, S. Lee, C. Lindskog, J. Mulder, C. M. Mulvey, P. Nilsson, P. Oksvold, J. Rockberg, R. Schutten, J. M. Schwenk, Å. Sivertsson, E. Sjöstedt, M. Skogs, C. Stadler, D. P. Sullivan, H. Tegel, C. Winsnes, C. Zhang, M. Zwahlen, A. Mardinoglu, F. Pontén, K. von Feilitzen, K. S. Lilley, M. Uhlén and E. Lundberg, "A subcellular map of the human proteome.," *Science (New York, N.Y.),* vol. 356, no. 6340, May 2017.

[37] The Human Protein Atlas, *RGS5.*

[38] J. Wang, L. Yin and Z. Chen, "Neuroprotective role of fibronectin in neuron-glial extrasynaptic transmission.," *Neural regeneration research,* vol. 8, no. 4, p. 376–382, February 2013.

[39] H. Togashi, T. Sakisaka and Y. Takai, "Cell adhesion molecules in the central nervous system.," *Cell adhesion & migration,* vol. 3, no. 1, p. 29–35, 2009.

[40] J. Biran, M. Tahor, E. Wircer and G. Levkowitz, "Role of developmental factors in hypothalamic function.," *Frontiers in neuroanatomy,* vol. 9, p. 47, 2015.

[41] 10x Genomics, "Fresh Frozen Mouse Brain Replicates," *In Situ Gene Expression dataset analyzed using Xenium Onboard Analysis 1.0.2,* 2023, Jan 22.

[42] C. K. Abrams, "Diseases of connexins expressed in myelinating glia.," *Neuroscience letters,* vol. 695, p. 91–99, March 2019.

[43] T. Turnescu, J. Arter, S. Reiprich, E. R. Tamm, A. Waisman and M. Wegner, "Sox8 and Sox10 jointly maintain myelin gene expression in oligodendrocytes," *Glia,* vol. 66, pp. 279-294, 2018.

[44] R. Tang, J. Vargas-Medrano, E. Ramos, P. Thompson and B. Gadad, "Gene Expression Analysis of CCL2, MOBP and OPALIN in Major Depressive Disorder and Suicidality (P5-6.005)," *Neurology,* vol. 98, p. 3266, 2022.

[45] J. Hornig, F. Fröb, M. R. Vogl, I. Hermans-Borgmeyer, E. R. Tamm and M. Wegner, "The transcription factors Sox10 and Myrf define an essential regulatory network module in differentiating oligodendrocytes," *PLoS genetics,* vol. 9, p. e1003907, 2013.

[46] O. de Faria Jr, A. S. Dhaunchak, Y. Kamen, A. D. Roth, T. Kuhlmann, D. R. Colman and T. E. Kennedy, "TMEM10 promotes oligodendrocyte differentiation and is expressed by oligodendrocytes in human remyelinating multiple sclerosis plaques," *Scientific reports,* vol. 9, p. 3606, 2019.

[47] S. George, M. R. Hamblin and H. Abrahamse, "Differentiation of mesenchymal stem cells to neuroglia: in the context of cell signalling," *Stem cell reviews and reports,* vol. 15, p. 814–826, 2019.

[48] M. S. Dicken, A. R. Hughes and S. T. Hentges, "Gad1 mRNA as a reliable indicator of altered GABA release from orexigenic neurons in the hypothalamus.," *The European journal of neuroscience,* vol. 42, no. 9, p. 2644–2653, November 2015.

[49] Z. Z. Pan, "Transcriptional control of Gad2.," *Transcription,* vol. 3, no. 2, p. 68–72, 2012.

[50] O. M. Schlüter, J. Basu, T. C. Südhof and C. Rosenmund, "Rab3 superprimes synaptic vesicles for release: implications for short-term synaptic plasticity.," *The Journal of neuroscience : the official journal of the Society for Neuroscience,* vol. 26, no. 4, p. 1239–1246, January 2006.

[51] T. Tsetsenis, T. J. Younts, C. Q. Chiu, P. S. Kaeser, P. E. Castillo and T. C. Südhof, "Rab3B protein is required for long-term depression of hippocampal inhibitory synapses and for normal reversal learning," *Proceedings of the National Academy of Sciences,* vol. 108, p. 14300–14305, 2011.

[52] 10x Genomics, "Human Kidney Preview Data (Xenium Human Multi-Tissue and Cancer Panel)," *In Situ Gene Expression dataset analyzed using Xenium Onboard Analysis 1.5.0,* 2023, July 17.

[53] 10x Genomics, "Xenium In Situ Analysis of Alzheimer's Disease Mouse Model Brain Coronal Sections from One Hemisphere Over a Time Course," *In Situ Gene Expression dataset analyzed using Xenium Onboard Analysis 1.4.0,* 2023, July 13.

[54] G. Novikova, M. Kapoor, J. Tcw, E. M. Abud, A. G. Efthymiou, S. X. Chen, H. Cheng, J. F. Fullard, J. Bendl, Y. Liu and others, "Integration of Alzheimer's disease genetics and myeloid genomics identifies disease risk regulatory elements and genes," *Nature communications,* vol. 12, p. 1610, 2021.

[55] S. Baig, S. A. Joseph, H. Tayler, R. Abraham, M. J. Owen, J. Williams, P. G. Kehoe and S. Love, "Distribution and expression of picalm in Alzheimer disease.," *Journal of neuropathology and experimental neurology,* vol. 69, no. 10, p. 1071–1077, October 2010.

[56] W. Xu, L. Tan and J.-T. Yu, "The Role of PICALM in Alzheimer's Disease.," *Molecular neurobiology,* vol. 52, no. 1, p. 399–413, August 2015.

[57] W. Xu, H.-F. Wang, L. Tan, M.-S. Tan, C.-C. Tan, X.-C. Zhu, D. Miao, W.-J. Yu, T. Jiang, L. Tan and others, "The impact of PICALM genetic variations on reserve capacity of posterior cingulate in AD continuum," *Scientific reports,* vol. 6, p. 24480, 2016.

[58] 10x Genomics, *Exploring Alzheimer's-like pathology at subcellular resolution using Xenium In Situ.*

[59] J. Yang, Z. Fu, X. Zhang, M. Xiong, L. Meng and Z. Zhang, "TREM2 ectodomain and its soluble form in Alzheimer's disease.," *Journal of neuroinflammation,* vol. 17, no. 1, p. 204, July 2020.

[60] L.-F. Lue, C. T. Schmitz, G. Serrano, L. I. Sue, T. G. Beach and D. G. Walker, "TREM2 Protein Expression Changes Correlate with Alzheimer's Disease Neurodegenerative Pathologies in Post-Mortem Temporal Cortices.," *Brain pathology (Zurich, Switzerland),* vol. 25, no. 4, p. 469–480, July 2015.

[61] Q. Qin, Z. Teng, C. Liu, Q. Li, Y. Yin and Y. Tang, "TREM2, microglia, and Alzheimer's disease.," *Mechanisms of ageing and development,* vol. 195, p. 111438, April 2021.

[62] E. Papuć and K. Rejdak, "The role of myelin damage in Alzheimer's disease pathology.," *Archives of medical science : AMS,* vol. 16, no. 2, p. 345–351, 2020.

[63] S. Ruskamo, A. Raasakka, J. S. Pedersen, A. Martel, K. Škubník, T. Darwish, L. Porcar and P. Kursula, "Human myelin proteolipid protein structure and lipid bilayer stacking," *Cellular and Molecular Life Sciences,* vol. 79, p. 419, 2022.

[64] P. Montague, A. S. McCallion, R. W. Davies and I. R. Griffiths, "Myelin-associated oligodendrocytic basic protein: a family of abundant CNS myelin proteins in search of a function," *Developmental neuroscience,* vol. 28, p. 479–487, 2006.

[65] M. Gravel, B. Trapp, J. Peterson and P. E. Braun, "CNP in myelination: Overexpression alters oligodendrocyte morphogenesis," *Cell Biology and Pathology of Myelin: Evolving Biological Concepts and Therapeutic Approaches,* p. 75–82, 1997.

[66] R. R. Sturrock, "Myelination of the mouse corpus callosum," *Neuropathology and applied neurobiology,* vol. 6, p. 415–420, 1980.

[67] M. Ota, N. Sato, Y. Kimura, Y. Shigemoto, H. Kunugi and H. Matsuda, "Changes of myelin organization in patients with Alzheimer's disease shown by q-space

myelin map imaging," *Dementia and Geriatric Cognitive Disorders Extra,* vol. 9, p. 24–33, 2019.

[68] R. R. Stickels, E. Murray, P. Kumar, J. Li, J. L. Marshall, D. J. Di Bella, P. Arlotta, E. Z. Macosko and F. Chen, "Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2," *Nature biotechnology,* vol. 39, p. 313–319, 2021.

[69] H. Zeng, J. Huang, H. Zhou, W. J. Meilandt, B. Dejanovic, Y. Zhou, C. J. Bohlen, S.-H. Lee, J. Ren, A. Liu and others, "Integrative in situ mapping of single-cell transcriptional states and tissue histopathology in a mouse model of Alzheimer's disease," *Nature Neuroscience,* vol. 26, p. 430–446, 2023.

[70] Y. Kim, W. Cheng, C.-S. Cho, Y. Hwang, Y. Si, A. Park, M. Schrank, J.-E. Hsu, A. Anacleto, J. Xi and others, "Seq-Scope: repurposing Illumina sequencing flow cells for high-resolution spatial transcriptomics," *Nature protocols,* p. 1–47, 2024.

[71] I. L. Dryden and K. V. Mardia, "Statistical Shape Analysis.,(Wiley: New York, NY.)," 1998.

[72] P. J. Santangelo, N. Nitin and G. Bao, "Direct visualization of mRNA colocalization with mitochondria in living cells using molecular beacons," *Journal of Biomedical Optics,* vol. 10, p. 044025–044025, 2005.

[73] S. Das, M. Vera, V. Gandin, R. H. Singer and E. Tutucci, "Intracellular mRNA transport and localized translation," *Nature reviews Molecular cell biology,* vol. 22, p. 483–504, 2021.

[74] H. Mi, D. Ebert, A. Muruganujan, C. Mills, L.-P. Albou, T. Mushayamaha and P. D. Thomas, "PANTHER version 16: a revised family classification, tree-based classification tool, enhancer regions and extensive API," *Nucleic acids research,* vol. 49, p. D394–D403, 2021.

[75] T. Biancalani, G. Scalia, L. Buffoni, R. Avasthi, Z. Lu, A. Sanger, N. Tokcan, C. R. Vanderburg, Å. Segerstolpe, M. Zhang and others, "Deep learning and alignment of spatially resolved single-cell transcriptomes with Tangram," *Nature methods,* vol. 18, p. 1352–1362, 2021.

[76] S. M. Lundberg and S.-I. Lee, "A Unified Approach to Interpreting Model Predictions," in *Advances in Neural Information Processing Systems 30*, I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan and R. Garnett, Eds., Curran Associates, Inc., 2017, p. 4765–4774.

[77] S. M. Lundberg, G. Erion, H. Chen, A. DeGrave, J. M. Prutkin, B. Nair, R. Katz, J. Himmelfarb, N. Bansal and S.-I. Lee, "From local explanations to global understanding with explainable AI for trees," *Nature Machine Intelligence,* vol. 2, pp. 2522-5839, 2020.

[78] S. Seabold and J. Perktold, "statsmodels: Econometric and statistical modeling with python," in *9th Python in Science Conference*, 2010.

[79] F. A. Wolf, P. Angerer and F. J. Theis, "SCANPY: large-scale single-cell gene expression data analysis," *Genome biology,* vol. 19, p. 1–5, 2018.

[80] N. Rezaie, F. Reese and A. Mortazavi, "PyWGCNA: a Python package for weighted gene co-expression network analysis," *Bioinformatics,* vol. 39, p. btad415, 2023.

[81] V. A. Padilha and R. J. G. B. Campello, "A systematic comparative evaluation of biclustering techniques," *BMC bioinformatics,* vol. 18, p. 1–25, 2017.

Figures

Figure 1: Schematic of CellSP. (A) CellSP analyzes single-molecule resolution spatial
transcriptomics data from a tissue; such data provides subcellular locations of individual
transcripts of a set of genes within each delineated cell. **(B)** Subcellular Pattern Discovery. CellSP
utilizes existing tools InSTAnT and SPRAWL for the discovery of subcellular spatial patterns of
genes within each cell. Types of patterns detected include gene-gene colocalization reported by
InSTAnT and four types of subcellular localization preferences (peripheral, punctate, central,
radial) reported by SPRAWL. **(C)** Module Discovery. The next step identifies spatial patterns
involving multiple genes and cells. Patterns identified in the previous step (represented as
matrices of cells x genes or gene-pairs) are subjected to a biclustering analysis that identifies a
subset of rows and columns with a large average value, representing a "gene-cell module": a set
of genes or gene-pairs exhibiting the same subcellular pattern within the same cells. Gene-cell
modules that overlap in a significant number of genes (or gene pairs) are coalesced to reduce
redundancy. This step outputs a set of gene-cell modules with varying number of cells and genes.
(D) CellSP provides intuitive visualizations of gene-cell modules, which often span tens to
hundreds of cells making direct inspection impractical. CellSP visualizations are tailored to the
type of pattern defining the module and summarize the strength of subcellular spatial pattern
involving module-associated genes and cells while contrasting them with background genes and
cells. **(E)** Module Characterization. Module-associated genes are subjected to Gene Ontology
(GO) analysis to gain functional insights into detected gene-cell modules. Alternatively, a Random
Forest classifier is trained to differentiate module cells from non-module cells and the most
predictive genes, called cell-marker genes, are subjected to GO analysis. To address the limited
size of gene panels in subcellular spatial transcriptomic datasets, CellSP employs Tangram to
impute gene expression from single-cell RNA-seq datasets of the same tissue, enabling more
comprehensive analysis. Results of GO analysis are visualized using REVIGO plots.

Figure 2: **CellSP reveals myelination-related subcellular spatial phenomena in mouse**
brain. (A) Summary table of an example gene-cell module, "M0_I", discovered by CellSP from
MERFISH data on hypothalamic preoptic area in mouse brain. (The suffix "_I" indicates that it was
discovered by aggregating single-cell level colocalization patterns reported by InSTAnT.) "GO
Genes" shows the Gene Ontology (GO)-based characterization of the four module genes. "GO
Cells" shows the GO-based characterization of the genes whose expression levels are most
discriminative of the 332 module cells versus remaining cells. **(B)** Direct visualization of six of the
332 module cells, with blue boundaries indicating automatically delineated cell periphery, red
boundaries indicating nuclear periphery, each dot representing a transcript, colored in green for
module genes and in gray for other genes. **(C)** CellSP visualization of module M0_I. The heatmap
shows pairwise colocalization propensity of gene pairs in module cells compared to non-module
cells, as a log ratio, with warmer colors (more positive values) indicating specificity of the
colocalization to the module cells. The first half of rows and columns corresponds to module genes
while the second half corresponds to a random subset of non-module genes, so the contrast
between the upper left quadrant (more warm color cells) and other quadrants underscores that
the phenomenon is specific to the module gene pairs. **(D)** UMAP visualization of all cells in the
sample, with colors indicating cell types and module cells shown in black. The module mostly
comprises mature oligodendrocytes (ODM). **(E)** REVIGO plot of biological processes enriched in
module genes, showing a prominent association with myelination-related terms. **(F)** REVIGO plot
of biological processes associated with gene markers of module cells, showing that these cells
are also characterized by genes enriched for myelination-related terms.

Figure 3: CellSP reveals axonogenesis related subcellular spatial phenomena in mouse brain. (A) Summary table of an example gene-cell module, "M0_S", discovered by CellSP from MERFISH data on hypothalamic preoptic area in mouse brain. (The suffix "_S" indicates that it was discovered by aggregating single-cell level colocalization patterns reported by SPRAWL.) **(B)** Direct visualization of six of the 221 module cells, with each dot representing a transcript, colored in green for module genes and in gray for other genes. **(C)** CellSP visualization of module M0_S. The subcellular space is idealized as a circle with concentric rings representing varying distances from the "center" (see Methods) and color intensity representing transcript abundance of a gene set in a ring, averaged over the module cells. The upper circle depicts this information for module genes and the lower circle represents all other genes for contrast, which in this case highlights that the module gene transcripts are enriched in peripheral regions (outermost ring). This type of visualization is used for any gene-cell module with peripheral or central pattern. **(D)** UMAP visualization of all cells in the tissue, with colors indicating cell types and module cells shown in black. The module mostly comprises neurons. **(E)** REVIGO plot of biological processes associated with module genes, highlighting enrichment for axonogenesis-related terms. **(F)** REVIGO plot of biological processes associated with marker genes of module cells, showing that these cells are characterized by genes enriched for cell adhesion-related terms. **(G-H)** Spatial plot

802   of all cells in tissue (H) with module cells highlighted in red, along with detailed visualization of a
803   subset of module cells in close spatial proximity (G,I). In these detailed views, module cells are
804   depicted with a stronger boundary, while their fill color corresponds to their cell type. Transcripts
805   of module genes are highlighted in red and are localized at the contact boundaries between
806   module cells.

Figure 4: **CellSP uncovers myelination and GABA-ergic synapse related modules across cell types and regions of mouse brain. (A)** UMAP visualization of all cells in a Xenium data set of whole mouse brain, with cells of 36 myelination-related gene-cell modules detected by CellSP shown in color. (Cells of each module are shown in a different color.) This demonstrates that CellSP captures myelination-related subcellular patterns across cells with diverse gene expression profiles, indicating that these patterns are not merely a reflection of overall gene expression levels. **(B)** Spatial plot of all cells in the whole brain data set, with cells of myelination-related modules (same as in A) shown in different colors, while cells not in any of these modules are shown in grey. **(C)** Matrix showing gene composition of each of the 36 myelination-related modules (rows); blue indicates a gene's inclusion in a module. Genes Sox10, Gjc3, and Opalin appear in all or most modules, while four others appear in a majority of modules. **(D)** REVIGO plot of biological processes associated with module genes, aggregated over all 36 modules. Color intensity of a bin in this plot represents the number of significant module associations of one or more GO terms that map to that bin in the semantic space computed by REVIGO. **(E)** REVIGO plot of biological processes associated with gene markers of module cells, aggregated over all the 36 modules. **(F)** UMAP visualization of all cells in data set, with cells in any of the 20 discovered GABA-ergic synapse-related modules shown in color. (Cells of each module are shown in a different color.) **(G)** Spatial plot of all cells in the sample, with cells from all 20 GABA-ergic synapse-related modules shown in different colors, while cells not in any of these modules are shown in grey.

828
Figure 5: **CellSP detects differences in subcellular spatial patterns between tissue**
conditions. (A) Spatial plot of a Kidney PRCC (kidney cancer) tissue (Xenium data), highlighting
cluster 6 cells in blue, cluster 9 cells in green, and cells of any of the 6 CellSP-detected immune-
response related gene-cell modules in red. **(B, C, D)** CellSP visualization of three selected
immune system related gene-cell modules discovered in the cancer sample. (B) shows module
R6_M5_S (central pattern) in left panel while right panel is a visualization of the subcellular
distribution of the module genes in expression-matched cells from the healthy kidney sample
("control"). Module genes are more uniformly localized towards the cell's "center" in the cancer
sample than in healthy sample. (C) shows module R6_M0_S (radial pattern) in left panel and the
right panel depicts the subcellular distribution of module genes in cells from the healthy sample.
The cancer sample (left panel) shows a strong clear difference between module genes and non-
module genes in terms of radial spread of transcripts, while this contrast is insignificant in the
healthy sample (right panel). (D) shows module R6_M3_I, defined by a colocalization pattern. The
colocalization visualization has been modified to provide contrast with control cells rather than
non-module cancer cells. The module genes are much more proximal to each other in cancer
module cells than in random cells from the healthy sample. **(E)** Gene-cell modules detected in
mouse hemibrain Xenium samples from three TgCRND8 mice (AD model) at different time points
and three wildtype (WT) mice at similar time points were examined for inclusion of six AD-related
genes (columns). Shown are the number of detected modules in each of the six samples (rows)
that include a gene. Notably, Picalm and Trem2 have more modules detected in AD samples than
in WT, consistent with their known roles in Alzheimer's Disease-related processes. **(F)** Chord
diagram depicting the number of modules detected in AD samples (bottom) or WT samples (top)
that include a pair of genes, shown for genes annotated with function "ensheathment of neurons".
The gene pairs "Olig2-Plp1" and "Olig2-Mobp" (shown in black) exhibit a significant difference
between AD and WT samples. **(G)** Blue bars show z-scores computed from a proportion test
comparing AD and WT samples in terms the fraction of CellSP modules that include Olig2 and
another gene (x-axis labels), shown only for partner genes where the proportion test yields a
significant p-value. Orange bars show similarly calculated z-scores for modules defined by cell-
level co-expression, as computed by WGCNA.