Corticosteroid resistance is predetermined by early immune response dynamics at acute Graft-versus-Host disease onset

One Sentence Summary (150 characters): Steroid-resistant acute graft versus host disease is set at disease onset as evidenced by specific transcriptomic signature, cell-to-cell crosstalk and trajectories.

Sophie Le Grand[1], Yannick Marie[2], Delphine Bouteiller[2], Émilie Robert[3], Régis Peffault de Latour[3,4], Gérard Socié[1,4], Nicolas Vallet[5]*, David Michonneau[1,4]*

Affiliations:

[1]INSERM UMR 1342, Translational Immunology in Immunotherapy and Hematology, IHU Leukemia Institute Paris Saint Louis, Paris Cité University, F-75010, Paris, France

[2]Genotyping and Sequencing Facility, Paris Brain Institute-ICM, Hôpital de la Pitié-Salpêtrière, CNRS UMR 7225, INSERM U1127, Sorbonne Université UM75, CS21414, Paris, France.

[3]Cryostem Consortium, Technopôle de Château Gombert, Marseille, France

[4]Hematology Transplantation, Saint Louis Hospital, Assistance Publique des Hôpitaux de Paris, 75010 Paris, France

[5]Hematology and Cell Therapy department, University Hospital of Tours, Inserm U1069 N2COx

*co-last authors

Corresponding author:

David Michonneau

Saint Louis hospital, hematology and transplantation

1 avenue Claude Vellefaux

75010 PARIS, FRANCE

david.michonneau@aphp.fr

Abstract

Steroid-resistant acute graft versus host disease (SR-aGVHD) is the leading life-threatening complication following allogeneic hematopoietic stem cell transplantation. Novel therapeutics development is impeded by scares knowledge on biological pathways leading to steroid resistance at time of aGVHD diagnosis. To gain insight into our understanding on circulating immune cell subsets and functions at time of aGVHD, a single cell deep phenotyping and transcriptome analysis was performed on peripheral blood mononuclear cells from patients with aGVHD before steroid treatment or without aGVHD. We aimed at identifying biological patterns associated with steroid resistance at early onset of aGVHD. First, circulating immune cell subsets were associated with increased incidence of aGVHD, but not with steroid sensitivity. Then, pathway analysis and inferred ligand/receptor interactions revealed major functional divergences between steroid-sensitive (SS-) and SR-aGVHD, including enrichment of TNFα activation in SR-GVHD, as well as TNF/TNFR, CCL3, CCL4 and IL18 signaling, and decreased interferon α and γ signaling pathways, suggesting that steroid resistance in an intrinsic property of immune cells before any treatment. To go deeper into the understanding of mechanisms at play during SR-aGVHD, we modeled immune trajectories within CD8$^+$ T cells and evidenced specific direct transition, from an early naive state to a highly activated one. By contrast, SS-aGVHD involved specific gene signatures across multiple intermediate differentiation stages during cell-to-cell transitions. These findings provide evidence that steroid resistance is driven by intrinsic mechanisms already present at the onset of alloimmune response, that may serve as potential new therapeutic targets.

1 INTRODUCTION

Allogeneic hematopoietic stem cell transplantation (HSCT) is a major treatment for

hematological malignancies. Its therapeutic effect is based on alloreactivity of donor T cells

against tumor cells. However, donor T cells also react against healthy host cells, which may

lead to graft-versus-host disease (GVHD). Acute GVHD (aGVHD) may involve skin, gut, or

liver tissue. Its pathophysiology is initiated after tissue damage from high dose alkylating agent

and radiation based-conditioning regimens leading to recognition of non-self-major and minor

histocompatibility antigens (MHC and mHag(*1*)) and an alloreactive immune response,

exacerbated by altered tissue repair mechanisms and microbiome dysbiosis(*2*). Despite first

line treatment with prompt corticosteroid initiation(*3, 4*), more than 40% of patients are steroid

resistant (SR-aGVHD)(*5, 6*). Steroid resistance is defined by a progressive disease after three

days of standard dose of corticosteroids, an absence of clinical response after 7 days, an

absence of complete remission after two to four weeks, or recurrence while steroids are

stopped or tapered. Although ruxolitinib improves symptoms for 60% of SR-GVHD patients(*7,*

8), their prognosis is very dismal(*3, 7, 9*).

While biomarkers were recently developed to improve aGVHD diagnosis and risk

stratification(*10–14*), response to steroid is still unpredictable and biological mechanisms

underlying steroid resistance remains only partly understood, even in the few experimental

transplantation models of SR–GVHD described(*15–17*). In humans, specific whole gene

expression analysis at diagnosis of lower intestinal acute SR-GVHD identified an increased

repair-associated gene (amphiregulin and the aryl hydrocarbon receptor) and M2

macrophages pathways(*18*), suggesting that steroid resistance in not restricted to the T cells

response as initially suspected. Indeed, neutrophils participate in tissue damage at GVHD

onset and triggering, through reactive oxygen species production(*19*) and promote T cell

expansion(*20*) and differentiation to Th17 cells(*21, 22*). As corticosteroids increase neutrophils

count and development, it has been hypothesized that corticosteroids may trigger a positive

feedback loop between pathogenic T cells and neutrophils in SR-GVHD context(*23*).

Furthermore, neutrophil-predominant noncanonical inflammation pathways were recently

identified in gut-SR-GVHD biopsies, notably ubiquitin specific peptidase 17–like family of

genes(24). While gut microbiota's role has been extensively studied in GVHD, it is not well

understood in SR-GVHD. Butyrate, a short-chain-fatty-acid product from *Blautia* bacteria is

associated with better long terms outcomes in aGVHD(25–27), but one study found a negative

association between butyrate production and SR-GVHD, due to the ability of butyrate to impair

human colonic stem cells from recovering epithelial monolayer(28). Other studies identified

SR-GVHD biomarkers such as IL-22(15), but with controversial potential(29, 30). Most of these

studies rely on murine models(15–17, 23, 31), and there is currently no clear overview of

immune subsets associated with steroid resistance in human GVHD.

We hypothesize that reshaping of immune system might be initiated early in

alloimmune responses, therefore predisposing to corticosteroids resistance. To describe

immune subsets and their respective function, cellular indexing of transcriptomes and epitopes

sequencing (CITE-Seq) was performed on peripheral blood mononuclear cells (PBMC)

samples collected at the onset of aGVHD before first line treatment, or in patient without GVHD

at day 100 after HSCT. Whole landscape of circulating immune cells at early onset of aGVHD

and its association with GVHD severity was uncovered. Deep analysis of transcriptomic

profiles revealed that steroid resistance is an intrinsic property of the immune response,

characterized by specific enriched biological pathways, specific cell-to-cell crosstalk through

ligand-receptor interactions and a specific immune trajectory involving most immune subsets.

Altogether, our results highlight that steroid-resistant GVHD is an early characteristic of allo-

immune response before any immunosuppressive treatment and pave the way to more

targeted first-line therapy to improve allogeneic HSCT outcomes.

RESULTS

Cohort's characteristics

Samples from 53 patients were retrieved from the prospective CRYOSTEM multicenter collection. PBMC were collected before corticosteroids initiation for patients diagnosed with aGVHD, and at day 100 post allo-HSCT for non-GVHD patients. Hematological malignancies, HSCT procedure, clinical and GVHD characteristics were balanced between groups, except for graft source, delay between sample and transplantation date (**Table S1)**. Among the 21 aGVHD patients, 16 (76%) were steroid sensitive (SS-GVHD) and 5 (24%) were steroid refractory (SR-GVHD), with a median delay of 24 days [13-96] and 17 days [11-70] between transplantation and sample, respectively (p=0.3) (**Figure 1A**). For validation and control purposes, 6 samples from healthy blood donor were used. After samples preparation for CITE-seq experiments, sequencing, quality control filtering and data processing acquisition, more than 300,000 single cells were available for immune profiling and transcriptome analysis (**Figure 1B**). Unsupervised clustering resulted in the identification of 40 immune cell subsets (**Figure 1C and S1**), characterized by their surface antigens expression (**Figure 1D**).

Immune signature and increased proliferation associate with aGVHD onset but not with response to steroids

Compared to non-aGVHD patients, classical monocytes $CD73^+$ were significantly enriched in aGVHD patients, whereas NK and B cell subsets were significantly decreased (**Figure 1E**). To encompass the compositional structure of immune subsets frequencies, principal coordinates analysis (PCoA) was used. Four distinct immune clusters were identified, thereafter called immunotypes (**Figure 2A**). aGVHD patient's distribution was significantly different among immunotypes (p=0.001, **Table S2**, **Figure 2B**), as well as the GVHD cumulative incidence (**Figure 2C,** Immunotype 4 GVHD incidence = 72% at day 100, *vs* immunotype 1 = 55%, *vs* immunotype 2 = 38%, *vs* immunotype 4 = 0%, global log-rank p=0.005). GVHD-free and

relapse-free survival (GRFS) was significantly associated with immunotypes (p=0.0048,

Figure S2), however there were no significant differences in non-relapse mortality, cumulative

incidence of relapse or of chronic GVHD (**Figure S2**), and no association with aGVHD organ

involvement (**Table S2**). Association with other clinical variables is depicted in **Figure S3**.

Immunotype 1 and 4 included a large number of patients with aGVHD (**Figure 2B–C**)

and were characterized by an overall reduction in immune cell heterogeneity, illustrated with

an imbalanced immune subsets representation (**Figure 2D**). Especially, immunotypes 1 and 4

presented a high display of classical monocytes $CD11b^{bright}$ (IgM^+ or IgM^-) compared to other

immunotypes. Immunotype 2 and 3 were characterized by a balanced distribution across

different immune cell populations, and a lower classical monocytes $CD11b^{bright}$ cells

representation. Remarkably, immunotype 3 was characterized by an absence of patients with

aGVHD, a decrease in classical monocytes $CD11b^{bright}$ and in other monocytes populations,

whereas effector T $CD4^+$ and $CD8^+$ were predominantly increased.

To further investigate whether the over-representation may encompass a high cell

division rate at time of aGVHD diagnosis, we determined cell cycle phase scoring within each

immune cell population. Immune populations were predominantly in S phase (**Figure 3A**),

reflecting increase cell proliferation at aGVHD onset.

No significant differences among immune cells abundances, immunotypes and

cell-cycle scoring was observed with respect to corticosteroid resistance (**Figure 2B**, **Table**

S2, Figure S4 and S5). In summary, immune cells subset clearly identified patients without

GVHD but hardly discriminated SS and SR-GVHD. We thus hypothesized that cell intrinsic

biological functions may predispose to SR-GVHD.

Steroid resistance is associated with enrichment of TNFα and metabolic pathways at aGVHD onset

We first considered the biological pathways enriched at the onset of aGVHD by comparison with patients without GVHD (**Figure S6**). When considering aGVHD as a whole, we observe overactivation of IFNα and γ pathways in all immune cells, and a decrease in TNFα signaling, especially in CD8$^+$ T cells and myeloid cells.

To study which biological process may correlate with subsequent SS- or SR-aGVHD, we then performed a second enrichment analysis in patients with SS- or SR-aGVHD. Compared to SS-aGVHD, inflammation pathways were enriched in SR-aGVHD patients within all immune cells, particularly TNFα and hypoxia pathways (**Figure 3B**). STAT5 and MTORC pathways were particularly upregulated in exhausted effector memory CD4$^+$ and CD8$^+$ T cells, in naïve CD8$^+$ T cells and in dendritic immune cells. The IL6/JAK STAT pathway was also highly enriched in naive CD8$^+$ T cells and classical CD11b^{bright} monocytes. Genes associated with metabolism pathways were overexpressed, especially oxidative phosphorylation in naïve CD4$^+$ T cells, double negative T cells and myeloid cells. Conversely, there was a decreased activation of the IFNα and γ responses within all immune subsets, except NK cells.

Cell to cell crosstalk during SR-GVHD specifically involves IL1, CCL3, CCL4, TNFα and IL18 signaling

To address whether cell biological processes involved in SR-aGVHD may be involved in cell-to-cell interactions leading to early steroid-resistance, we examined intercellular communication using a computational inferring method(*32*). When focusing on CD8$^+$ T cells as receiver, 20 ligands were identified as prioritized ligands, and their expression on sender cells is shown in **Figure 4A**. IL1B, CCL3 and CCL4 ligands were particularly overexpressed in

SR-aGVHD condition (**Figure 4B**). Regulatory potential of each ligand with predicted targets

genes in shown in **Figure 4C**.

Predicted ligands and targets interaction were then analyzed for CD4 regulatory T cells

(**Figure S7A**), NK cells (**Figure S7B**), CD4[+]T cells (**Figure S8A**), and double negative T cells

(**Figure S8B**). An overview of all these interactions is depicted as a circus plot (**Figure 4D**)

based on biologically confirmed interactions previously designated as *bona fide* interactions

(**Figure S9A**). TNF ligands-to-cell signaling appeared to be overexpressed in most sender

cells in SR-aGVHD patients, as well as CCL3, CCL4 and IL18 signaling. ADAM17, which is

involved in TNF activation(*33*), was overexpressed in all senders and interacted with TNFR,

ITGAL and ITGB2 receptors. Interestingly, IFN/IFNR interactions was enhanced across all

receivers in SR-aGVHD but absent in SS-aGVHD.

Conversely, when comparing patients with aGVHD to those without aGVHD, immune

cells crosstalk appeared different to that disclosed in the SS- and SR–aGVHD setting

(**Figure S9B**). Consistent with enrichment analysis, TNF/TNF-R interaction was decreased

among DN T cells, NK cells and T regulatory lymphocytes. TGFB1/TGFBR interactions, which

are considered as being associated with an immunoregulatory role, were significantly

increased in patient with no aGVHD compared to aGVHD patients, especially within CD8[+] T

cells. BTLA/TNFRSF14 interaction is another pathway previously identified in animal models

as a regulator of alloimmune response(*34*, *35*), and was overexpressed in patients without

GVHD.

Specific trajectories were associated with SR-GVHD

As biological processes and cell-to-cell interactions targeting CD8[+] T cells were associated

with SR-aGVHD, we hypothesized that immune cells subsets from SR-aGVHD and SS-

aGVHD patients may exert specific CD8[+] T cell trajectories. To address this hypothesis,

trajectories within CD8[+] T cell were defined on a new cell clustering based on mRNA

expression (**Figure S10**). Two trajectories were identified for CD8⁺ T cell: (i) lineage 1

described in **Figure 5A-B** and (ii) lineage 2 described in **Figure S11**.

Frequency of cells in lineage 1 slightly increased over pseudotime in SS-aGVHD

patients through multiple transition states starting from naïve double positive T cell (cluster 27)

to effector memory CD8⁺ T cell expressing ICOS (cluster 2) **(Figure 5A-B).** By contrast,

SR-aGVHD patients were characterized by a direct transition with very few cells in the

transition subsets (cluster 15, 9 and 10), most immune subset being grouped in cluster 27 at

early pseudotime or in cluster 2 in late pseudotime. The same pattern was observed in the

second CD8⁺ T cell trajectory, with a transition from naïve double positive T cell (cluster 27) to

CD8⁺ TEMRA cells expressing ICOS and PD1 (cluster 25) (**Figure S11A-B**).

To highlight different gene expression pattern in SS-aGVHD and SR-aGVHD we first

studied individual genes whose expression varied significantly along pseudotime and across

transition state (**Figure 5C, S11C, Table S3-S10**). Among the most significant genes involved

in transition states, we identified CCL4, CCL5, IFI6, LAG3, and TIGIT (**Table S11 and S12**).

SR-aGVHD patients exhibited earlier and higher CCL4 and LAG3 expression in pseudotime

(**Figure 5D and S11D**). Consistently with pathway enrichment analysis, IFI6 expression,

indicative of IFNα activation, was higher in SS-aGVHD patients, particularly in early-

differentiated CD8⁺ T cells.

To study overall biological processes, we used hierarchical clustering to identify

modules of correlated gene among the top 500 expressed gene along pseudotime lineages.

This approach identified 5 modules of genes involved in cell-to-cell transition (**Figure 5C**).

Over-representation analysis (ORA) from Gene Ontology knowledgebase within each cell

subset highlighted the main biological pathways involved at each transition state (**Figure 5E**).

SR-aGVHD was notably associated with an early enrichment of pathways associated with cell

activation and high transcriptional and translational activity in naïve double positive T cell

(**Figure S11E**). A detailed analysis of pathways enriched in each module revealed striking

differences in the dynamics of T cell activation during SR-GVHD (trajectory 1 in **Figure S12**

and S13, and trajectory 2 in **Figure S14 and S15**).

Overall, these results reveal that even before starting corticosteroids, aGVHD is already

characterized by different trajectories and pattern of gene activation between transition states,

that can define a specific dynamic of alloimmune response during steroid resistance.

DISCUSSION

Despite recent advances in the management of aGVHD, steroid-resistant treatment remains

a challenge. This study is the first to describe immune subsets and transcriptional profiles in

human at time of aGVHD. Early identification of patients at risk of developing steroid resistance

is crucial for rapid and tailored therapeutic intervention. Up to now, a comprehensive

understanding of the circulating immune cells mechanisms associated with steroid resistance

has not been addressed. Our results highlight that steroid resistance is characterized at the

onset of aGVHD, by a specific transcriptomic profile that reveal biological pathways, cell-to-cell

interactions and trajectories that differ from steroid sensitive patients.

Immunotypes and cell cycle scoring enabled the identification of specific patterns

associated with aGVHD compared to non-aGVHD patients. Immunotype 4, characterized by

poor immune cell diversity and by classical CD11$^+$ monocytes abundance was associated with

worsened aGVHD free survival. This observation is consistent with previous studies identifying

monocytes and macrophages as fully parts of the immune response at aGVHD onset (*18*, *36–*

39).

However, cell subsets distribution did not allow to distinguish SS- and SR-aGVHD.

When focusing on immune cells functions rather than distinct phenotypes, we found specific

patterns associated with SR-GVHD, especially higher TNFα pathway activation. This was

strengthened with inferred interaction analyses which revealed an increased TNF/TNFR

signaling, notably in CD8$^+$ T cells. Although this *in silico* prediction would need to be confirmed

in vivo, these results might suggest that early anti-TNF therapies could be beneficial in patients

with the corticosteroid resistance profile we described herein. This is supported by a previous

clinical phase 2 trial testing a combined therapy of anti-TNF and corticosteroids, with increased

response rate (69% vs 33%; P < 0.001) without increased infections risk(*40*).

Our results suggest that the alloimmune response responsible for GVHD is already

different at its initiation in patients who will be steroid-resistant. Overall, the integration of our

data shows that steroid resistance could result from the cytokine production profile by myeloid

populations, especially monocytes producing CCL3, CCL4, IL1, and IL18, which stimulate the

activation of naive CD8$^+$ T cells and induce direct differentiation into effector T cells (**Figure 6**).

These results are consistent with published data on the role of myeloid populations(*19*, *20*)

and inflammasome activation(*41*) in the initiation of acute GVHD in mice, or of M2 macrophage

in the gut of patients with SR-GVHD(*18*). Additionally, previous data had also suggested that

the effect of IL-18 differs depending on whether GVHD is mainly mediated by CD8$^+$ or

CD4$^+$ T cells in murine models(*42*). Our data suggest that, in humans, this effector phase

depends on the activation profile of CD8$^+$ T cells, and that IL-18 may worsen the severity of

CD8-mediated GVHD by increasing steroid resistance. IL18 is a proinflammatory cytokine

which can promote interferon γ production(*43*), and its serum level in post-HSCT patients was

associated with aGVHD severity (r = 0.861, P < 0.001)(*44*). Furthermore, our results highlight

the key role of TNFα production by myeloid cells in SR-GVHD on CD4+ T activation that could

amplify the alloimmune response. These results confirm data obtained in animal models and

show that myeloid subsets may not only participate in the initiation of GVHD(*19*) but that their

response profile could already initiate the first mechanisms of steroid resistance.

Interferon have been described for their crucial role in GVHD development(*45–47*). We

found that IFNα and γ pathways were enriched at time of aGVHD and before corticosteroid

treatment. However, SR-aGVHD was associated with lower activation of IFNα and γ pathways

in most immune subsets, except for naïve CD8 T cells and non-classical monocytes. This could

explain why in SR-aGVHD, INFG/IFNGR interactions increased cell-to-cell crosstalk between

CD8+ T cells, monocytes and CD4+ T cells. In addition, it has been suggested previously that

IFNγ production could contribute to increase the severity of digestive GVHD(*48, 49*) and that

IFNλ blockade could improve gut GVHD severity(*50*).

Finally, trajectory inferences highlighted specific layouts associated with SR-aGVHD.

Cells were particularly abundant in low and high pseudotime in SR-GVHD patients, while there

was a progressive increase along pseudotime in SS-GVHD patients **(Figure 6)**. It suggests

that cells in SR-GVHD patients quickly differentiate to terminally activated states. This was

consistent with, higher and earlier gene expression of LAG3 and TIGIT in SR-GVHD patients.

Consistent with results from cell-cell interactions analysis, CCL4 and CCL5 were expressed at

higher levels and earlier in SR-GVHD patients, as also described in animal model(*51*).

Because samples were collected before cortico-steroid initiation, our work sheds light

on early signature pathways leading to steroid resistance. These pathways appeared distinct

from those involved in the general inflammatory response of acute GVHD and could be

targeted as early as steroid initiation. Transcriptomic analysis offered a powerful approach to

unravel immune cells functionality, combined with precise phenotypic population identification,

allowing for the identification of cellular heterogeneity in SR-GVHD immune response. The

complexity of the immune response in the context of GVHD was already investigated at the

single cell level, through T cell subset analysis following TCR clonotype mapping(*52*). This

elegant approach, revealed that each patient possessed a highly intricate repertoire, with

different tissues exhibiting distinct clonal populations, varying degrees of repertoire overlap

between sites, and diverse numbers of dominant clones shared across multiple tissues (*52*).

More recently, a similar approach highlighted the impact of the tissue microenvironment on

maintaining T-cell activation and driving divergent immune responses in peripheral organs

(*53*). Our results rely on PBMCs samples and hence, provide a complementary perspective on

acute GVHD, that uncover biological pathways associated with steroid resistance,

irrespectively of targeted organs. Studies based directly on blood samples are essential in

clinical practice, as they provide an easy access to potential predictors that could guide

therapeutic management.

Our study offers new insights into the immune complexity, presenting a broader

perspective of the immune response. Since our samples were all collected at onset of GVHD,

the sampling time varied between patients, which could impact the immune response,

particularly cytokine secretion. Therefore, while our study underscores the need for validation

in a larger cohort with careful matching of sampling times, it could also serve as a robust basis

for future studies involving a broader patient population. Furthermore, the use of models would

require additional validation *in vivo*. Finally, additional studies focusing on immune infiltrates

in target organs of GVHD would certainly help to complete our understanding of acute GVHD

pathophysiology, and especially of mechanisms associated with corticosteroid resistance.

Overall, these findings provide evidence that corticosteroid resistance is an intrinsic

mechanism already present at the onset of alloimmune response, highlighting specific patterns

that may serve as potential new therapeutic targets.

MATERIALS AND METHODS

Cohort

This study was conducted with annotated samples from patients and healthy donors provided by the CRYOSTEM Consortium and the SFGM-TC (Société Francophone de Greffe de Moelle et de Thérapie Cellulaire). Patients were included in the CRYOSTEM cohort after validation by the scientific committee (study number CS 18-01). CRYOSTEM cohort and collection were approved by IRB Sud-Méditerranée 1 (reference number DC-2014-2312 and DC-2019-3425) and the Commission Nationale de l'Informatique et des Libertés for data protection (reference number nz70243374i n°912120). All patients gave their written consent for clinical research and sample collection. This non-interventional research study with no additional clinical procedure was carried out in accordance with the Declaration of Helsinki. Data analyses were carried out using an anonymized database. Healthy donor's PBMC were isolated from residual blood after apheresis provided by Etablissement Français du Sang (18/EFS/032).

CITE Seq experiment

After thawing cells in a 50% fetal calf serum (FCS)/RPMI medium, 10x Genomics Feature Barcoding recommendations were followed. Cells were washed twice using a 10 mL solution of the same media. 500 000 cells were transferred in a 0.04% PBS/bovine serum albumin (BSA) medium and washed. Cells were then suspended in a 1% PBS/BSA solution before incubation with oligonucleotide-coupled antibodies TotalSeq-B (panel described in **Table S13**) for 60 min on ice. Cells were washed 4 times (4°C, RCF 300, 5 minutes) and suspended to obtain a concentration of 1 000 cells/µL. 10 000 cells were subjected in Gel Bead Emulsion using Chromium 10x Genomics controller, according to manufacturer guidelines. Chromium™ Next GEM Single Cell 3' Kit v3.1 (10X Genomics, cat. 1000268) and 3' Feature Barcode Kit (10X Genomics, cat. 1000262) were used to prepare reagents.

To perform single cell RNA sequencing after cDNA amplification, the concentration of

each sample was measured using Tapestation 2200 (Agilent). To prepare the cDNA libraries

for 10x Genomics Chromium controller, we used the single-cell 3′ v3.1 kit (Dual Index) with

Feature Barcode technology for Cell Surface Protein, following manufacturer guidelines. QC

libraries were performed using Tapestation 2200 (Agilent). Libraries were equimolar pooled to

obtain at least 20 000 read pairs per cell and 5 000 read pairs for feature barcode part after

sequencing on Illumina Novaseq 6000 (100 cycles cartridge). The input number of cells was

estimated at 10 000 cells/samples. FASTQ files were obtained with bcl2fastq (Illumina).

Data quality control

Alignment, filtering, barcode counting, UMI counting were performed with count function from

10x Genomics Cell Ranger (version 6.0) using GRCh38 genome assembly as reference data.

Then aggr function was used to aggregate results without normalization. A R environment

(version 4.3.1) was subsequently used for the following analyses. Briefly, using Seurat R

package(*54*), dead cells were excluded if mitochondrial RNA was above 18%; singlet were

identified as droplets with RNA feature between 200 and 4 700; RNA and cell surface antigen

counts were normalized and scaled using Seurat R package normalization and scaling function

for linear transformation. 2 000 highly variable RNA features were identified.

Immune cell clustering

Using a principal component analysis method on the scaled data, cells were clustered

according to cell surface antigen expression. We used UMAP for cluster visualization. Seurat

R package was then used to calculate cell cycle phase based on canonical transcripts, and

regressing these out of the data.

Enrichment pathways analysis

Fast pre-ranked gene set enrichment analysis was performed using FGSEA package(*55*) with "hallmarks" gene reference data set from Molecular signature database(*56*). Pathways were compared between GVHD and non-GVHD patients, and between SR-GVHD and SS-GVHD patients. Gene expression comparison between GVHD and non GVHD patients, and between SS- and SR-GVHD patients was provided by differential expression analysis. Enrichment scores associated with adjusted p-value < 0.05 were selected for dot plot visualization.

NicheNet Analysis

Ligand-receptors interaction was computed using NicheNet package(*32*). Sender cells were identified as myeloid cells, B cells, CD4$^+$ conventional T cells and regulatory T cells. Receiver cells were identified as CD4$^+$ and CD8$^+$ T lymphocytes, NK cells and regulatory T cells. Three pooled human gene expression data of interacting cells(*57–59*) were used as input and combined with a prior model that integrates existing knowledge on ligand-to-target signaling paths to perform ligand activity analyses. Circos plot visualization integrated *bona fide* interactions of each receiver, and increased or decreased interactions were integrated from "ligand differential expression heatmap" NicheNet output.

Trajectory inferences

Trajectory were based on RNA expression clustering (**Figure S10**). SingleCellExperiment object was subset by immune populations (CD8+ T cells, CD4+ T cells, NK cells, B cells and myeloid cells) before pseudotime calculation using slingshot package(*60*). TradSeq package was then use for downstream analysis(*61*). Briefly, associationTest R function evidenced genes whose expression is associated with pseudotime. Since pseudotime is a continuous value, it was divided into 50 categories for graphical representation with heatmap. Only the 500 most expressed gene in SS- and SR-GVHD were selected for heatmap visualization

(**Table S3-S6**). Each point on X axis corresponded to a cell cluster with similar pseudotimes.

The immune populations represented on top of the heatmap corresponded to the immune cell

population most abundantly represented within the cell cluster (**Table S7-S10**). Gene markers

specific to low pseudotime or high pseudotime were found thanks to starVsEndTest command.

Expression along pseudotime in each condition was plotted for four genes from the top 100

starVsEndTest output **(Figure 5, Tables S11 and S12)**. To explore the pathways associated

with gene expression of the top 500 most expressed genes along pseudotime from each of

the 5 modules of the heatmap, we used over-representation analysis using enrichGO

command from clusterProfiler R package(*62*). Only pathways associated with adjusted

pval < 0.05 from "molecular functions" and "biological process" of the Gene Ontology

knowledgbase(*63, 64*) were selected for dot plots visualization.

List of Supplementary Materials

Figure S1 to S15

Fig. S1: Immune populations in each group of patients

Fig. S2: Immunotype survival analysis

Fig. S3: Immunotype and clinical variables

Fig. S4: Immune cell abundance in SS- and SR-GVHD patients

Fig. S5: Cell cycle in SS- and SR-GVHD patients and enrichment pathways in aGVHD

Fig. S6: Pathway enrichments in aGVHD patients

Fig. S7: Interactions combined plots in regulatory T cells and NK cells

Fig. S8: Interactions combined plots in CD4$^+$ T cells and double negative T cells

Fig. S9: Bona fide interactions in all receivers

Fig. S10: Clustering based on RNA expression

Fig. S11: CD8$^+$ T cells lineage 2 trajectory analysis

Fig. S12: Over representation analysis in lineage 1 SR-GVHD patients

Fig. S13: Over representation analysis in lineage 1 SS-GVHD patients

Fig. S14: Over representation analysis in lineage 2 SR-GVHD patients

Fig. S15: Over representation analysis in lineage 2 SS-GVHD patients

Table S1 to S2

Table S1: Patients' characteristics

Table S2: Characteristics according to immunotypes

Supplementary data files S3 to S13

Data file S3: Top 500 genes expression matrix in lineage 1 SR-GVHD patients

Data file S4: Top 500 genes expression matrix in lineage 1 SS-GVHD patients

Data file S5: Top 500 genes expression matrix in lineage 2 SR-GVHD patients

Data file S6: Top 500 genes expression matrix in lineage 2 SS-GVHD patients

387 Data file S7: Immune population mostly represented along pseudotimes in lineage 1 SR-GVHD

388 patients

389 Data file S8: Immune population mostly represented along pseudotimes in lineage 1 SS-GVHD

390 patients

391 Data file S9: Immune population mostly represented along pseudotimes in lineage 1

392 2 SR-GVHD patients

393 Data file S10: Immune population mostly represented along pseudotimes in lineage 2 SS-

394 GVHD patients

395 Data file S11: StartvsEnd output in lineage 1

396 Data file S12: StartvsEnd output in lineage 2

397 Data file S13: Total-Seq-B antibody panel

REFERENCES

1.   N. Cieri, N. Hookeri, K. Stromhaug, L. Li, J. Keating, P. Díaz-Fernández, V. Gómez-García de
Soria, J. Stevens, R. Kfuri-Rubens, Y. Shao, K. A. Kooshesh, K. Powell, H. Ji, G. M. Hernandez, J.
Abelin, S. Klaeger, C. Forman, K. R. Clauser, S. Sarkizova, D. A. Braun, L. Penter, H. T. Kim, W. J.
Lane, G. Oliveira, L. S. Kean, S. Li, K. J. Livak, S. A. Carr, D. B. Keskin, C. Muñoz-Calleja, V. T. Ho,
J. Ritz, R. J. Soiffer, D. Neuberg, C. Stewart, G. Getz, C. J. Wu, Systematic identification of minor
histocompatibility antigens predicts outcomes of allogeneic hematopoietic cell transplantation. *Nat*
Biotechnol, 1–12 (2024).
2.   R. Zeiser, B. R. Blazar, Acute Graft-versus-Host Disease — Biologic Process, Prevention, and
Therapy. *N Engl J Med* **377**, 2167–2179 (2017).
3.   A. C. Harris, R. Young, S. Devine, W. J. Hogan, F. Ayuk, U. Bunworasate, C.
Chanswangphuwana, Y. A. Efebera, E. Holler, M. Litzow, R. Ordemann, M. Qayed, A. S. Renteria, R.
Reshef, M. Wölfl, Y.-B. Chen, S. Goldstein, M. Jagasia, F. Locatelli, S. Mielke, D. Porter, T. Schechter,
Z. Shekhovtsova, J. L. M. Ferrara, J. E. Levine, International, Multicenter Standardization of Acute
Graft-versus-Host Disease Clinical Data Collection: A Report from the Mount Sinai Acute GVHD
International Consortium. *Biol Blood Marrow Transplant* **22**, 4–10 (2016).
4.   L. Axt, A. Naumann, J. Toennies, S. P. Haen, W. Vogel, D. Schneidawind, S. Wirths, R.
Moehle, C. Faul, L. Kanz, S. Axt, W. A. Bethge, Retrospective single center analysis of outcome, risk
factors and therapy in steroid refractory graft-versus-host disease after allogeneic hematopoietic cell
transplantation. *Bone Marrow Transplant* **54**, 1805–1814 (2019).
5.   P. J. Martin, J. D. Rizzo, J. R. Wingard, K. Ballen, P. T. Curtin, C. Cutler, M. R. Litzow, Y.
Nieto, B. N. Savani, J. R. Schriber, P. J. Shaughnessy, D. A. Wall, P. A. Carpenter, First- and second-
line systemic treatment of acute graft-versus-host disease: recommendations of the American Society
of Blood and Marrow Transplantation. *Biol Blood Marrow Transplant* **18**, 1150–1163 (2012).
6.   J. R. Westin, R. M. Saliba, M. De Lima, A. Alousi, C. Hosing, M. H. Qazilbash, I. F. Khouri,
E. J. Shpall, P. Anderlini, G. Rondon, B. S. Andersson, R. Champlin, D. R. Couriel, Steroid-Refractory
Acute GVHD: Predictors and Outcomes. *Adv Hematol* **2011**, 601953 (2011).
7.   R. Zeiser, N. von Bubnoff, J. Butler, M. Mohty, D. Niederwieser, R. Or, J. Szer, E. M. Wagner,
T. Zuckerman, B. Mahuzier, J. Xu, C. Wilke, K. K. Gandhi, G. Socié, REACH2 Trial Group, Ruxolitinib
for Glucocorticoid-Refractory Acute Graft-versus-Host Disease. *N Engl J Med* **382**, 1800–1810 (2020).
8.   O. Penack, M. Marchetti, M. Aljurf, M. Arat, F. Bonifazi, R. F. Duarte, S. Giebel, H. Greinix,
M. D. Hazenberg, N. Kröger, S. Mielke, M. Mohty, A. Nagler, J. Passweg, F. Patriarca, T. Ruutu, H.
Schoemans, C. Solano, R. Vrhovac, D. Wolff, R. Zeiser, A. Sureda, Z. Peric, Prophylaxis and
management of graft-versus-host disease after stem-cell transplantation for haematological
malignancies: updated consensus recommendations of the European Society for Blood and Marrow
Transplantation. *Lancet Haematol* **11**, e147–e159 (2024).
9.   P. J. Martin, G. Schoch, L. Fisher, V. Byers, C. Anasetti, F. R. Appelbaum, P. G. Beatty, K.
Doney, G. B. McDonald, J. E. Sanders, A retrospective analysis of therapy for acute graft-versus-host
disease: initial treatment. *Blood* **76**, 1464–1472 (1990).
10.   J. L. M. Ferrara, A. C. Harris, J. K. Greenson, T. M. Braun, E. Holler, T. Teshima, J. E. Levine,
S. W. J. Choi, E. Huber, K. Landfried, K. Akashi, M. Vander Lugt, P. Reddy, A. Chin, Q. Zhang, S.
Hanash, S. Paczesny, Regenerating islet-derived 3-alpha is a biomarker of gastrointestinal graft-versus-
host disease. *Blood* **118**, 6702–6708 (2011).
11.   M. T. Vander Lugt, T. M. Braun, S. Hanash, J. Ritz, V. T. Ho, J. H. Antin, Q. Zhang, C.-H.
Wong, H. Wang, A. Chin, A. Gomez, A. C. Harris, J. E. Levine, S. W. Choi, D. Couriel, P. Reddy, J. L.
M. Ferrara, S. Paczesny, ST2 as a marker for risk of therapy-resistant graft-versus-host disease and
death. *N Engl J Med* **369**, 529–539 (2013).
12.   M. J. Hartwell, U. Özbek, E. Holler, A. S. Renteria, H. Major-Monfried, P. Reddy, M. Aziz, W.
J. Hogan, F. Ayuk, Y. A. Efebera, E. O. Hexner, U. Bunworasate, M. Qayed, R. Ordemann, M. Wölfl,
S. Mielke, A. Pawarode, Y.-B. Chen, S. Devine, A. C. Harris, M. Jagasia, C. L. Kitko, M. R. Litzow,
N. Kröger, F. Locatelli, G. Morales, R. Nakamura, R. Reshef, W. Rösler, D. Weber, K. Wudhikarn, G.
A. Yanik, J. E. Levine, J. L. Ferrara, An early-biomarker algorithm predicts lethal graft-versus-host

disease and survival. *JCI Insight* **2**, e89798 (2017).

13. H. Major-Monfried, A. S. Renteria, A. Pawarode, P. Reddy, F. Ayuk, E. Holler, Y. A. Efebera,
W. J. Hogan, M. Wölfl, M. Qayed, E. O. Hexner, K. Wudhikarn, R. Ordemann, R. Young, J. Shah, M.
J. Hartwell, M. S. Chaudhry, M. Aziz, A. Etra, G. A. Yanik, N. Kröger, D. Weber, Y.-B. Chen, R.
Nakamura, W. Rösler, C. L. Kitko, A. C. Harris, M. Pulsipher, R. Reshef, S. Kowalyk, G. Morales, I.
Torres, U. Özbek, J. L. M. Ferrara, J. E. Levine, MAGIC biomarkers predict long-term outcomes for
steroid-resistant acute GVHD. *Blood* **131**, 2846–2855 (2018).

14. G. Socié, D. Niederwieser, N. von Bubnoff, M. Mohty, J. Szer, R. Or, J. Garrett, A. Prahallad,
C. Wilke, R. Zeiser, Prognostic value of blood biomarkers in steroid-refractory or steroid-dependent
acute graft-versus-host disease: a REACH2 analysis. *Blood* **141**, 2771–2779 (2023).

15. Q. Song, X. Wang, X. Wu, T. H. Kang, H. Qin, D. Zhao, R. R. Jenq, M. R. M. van den Brink,
A. D. Riggs, P. J. Martin, Y.-Z. Chen, D. Zeng, IL-22-dependent dysbiosis and mononuclear phagocyte
depletion contribute to steroid-resistant gut graft-versus-host disease in mice. *Nat Commun* **12**, 805
(2021).

16. V. Arnhold, W. Y. Chang, S. A. Jansen, G. Thangavelu, M. Calafiore, P. Vinci, Y.-Y. Fu, T.
Ito, S. Takashima, A. Egorova, J. Kuttiyara, A. Perlstein, M. van Hoesel, C. Liu, B. R. Blazar, C. A.
Lindemans, A. M. Hanash, Corticosteroids impair epithelial regeneration in immune-mediated intestinal
damage. *J Clin Invest* **134**, e155880 (2024).

17. T. Toubai, C. Rossi, I. Tawara, C. Liu, C. Zajac, K. Oravecz-Wilson, D. Peltier, Y. Sun, H.
Fujiwara, S.-R. Wu, M. Riwes, I. Henig, S. Kim, P. Reddy, Murine Models of Steroid Refractory Graft-
versus-Host Disease. *Sci Rep* **8**, 12475 (2018).

18. S. G. Holtan, A. Shabaneh, B. C. Betts, A. Rashidi, M. L. MacMillan, C. Ustun, K. Amin, B. P.
Vaughn, J. Howard, A. Khoruts, M. Arora, T. E. DeFor, D. Johnson, B. R. Blazar, D. J. Weisdorf, J.
Wang, Stress responses, M2 macrophages, and a distinct microbial signature in fatal intestinal acute
graft-versus-host disease. *JCI Insight* **5**, e129762, 129762 (2019).

19. L. Schwab, L. Goroncy, S. Palaniyandi, S. Gautam, A. Triantafyllopoulou, A. Mocsai, W.
Reichardt, F. J. Karlsson, S. V. Radhakrishnan, K. Hanke, A. Schmitt-Graeff, M. Freudenberg, F. D.
von Loewenich, P. Wolf, F. Leonhardt, N. Baxan, D. Pfeifer, O. Schmah, A. Schönle, S. F. Martin, R.
Mertelsmann, J. Duyster, J. Finke, M. Prinz, P. Henneke, H. Häcker, G. C. Hildebrandt, G. Häcker, R.
Zeiser, Neutrophil granulocytes recruited upon translocation of intestinal bacteria enhance graft-versus-
host disease via tissue damage. *Nat Med* **20**, 648–654 (2014).

20. J. Hülsdünker, K. J. Ottmüller, H. P. Neeff, M. Koyama, Z. Gao, O. S. Thomas, M. Follo, A.
Al-Ahmad, G. Prinz, S. Duquesne, H. Dierbach, S. Kirschnek, T. Lämmermann, M. J. Blaser, B. T. Fife,
B. R. Blazar, A. Beilhack, G. R. Hill, G. Häcker, R. Zeiser, Neutrophils provide cellular communication
between ileum and mesenteric lymph nodes at graft-versus-host disease onset. *Blood* **131**, 1858–1869
(2018).

21. M. Pelletier, L. Maggi, A. Micheletti, E. Lazzeri, N. Tamassia, C. Costantini, L. Cosmi, C.
Lunardi, F. Annunziato, S. Romagnani, M. A. Cassatella, Evidence for a cross-talk between human
neutrophils and Th17 cells. *Blood* **115**, 335–343 (2010).

22. D. Minns, K. J. Smith, V. Alessandrini, G. Hardisty, L. Melrose, L. Jackson-Jones, A. S.
MacDonald, D. J. Davidson, E. Gwyer Findlay, The neutrophil antimicrobial peptide cathelicidin
promotes Th17 differentiation. *Nat Commun* **12**, 1285 (2021).

23. Q. Song, U. Nasri, D. Zeng, Steroid-Refractory Gut Graft-Versus-Host Disease: What We Have
Learned From Basic Immunology and Experimental Mouse Model. *Front Immunol* **13**, 844271 (2022).

24. B. K. Patel, M. J. Raabe, E. R. Lang, Y. Song, C. Lu, V. Deshpande, L. T. Nieman, M. J. Aryee,
Y.-B. Chen, D. T. Ting, Z. DeFilipp, Spatial transcriptomics reveals distinct tissue niches linked with
steroid responsiveness in acute gastrointestinal GVHD. *Blood* **142**, 1831–1844 (2023).

25. R. R. Jenq, Y. Taur, S. M. Devlin, D. M. Ponce, J. D. Goldberg, K. F. Ahr, E. R. Littmann, L.
Ling, A. C. Gobourne, L. C. Miller, M. D. Docampo, J. U. Peled, N. Arpaia, J. R. Cross, T. K. Peets,
M. A. Lumish, Y. Shono, J. A. Dudakov, H. Poeck, A. M. Hanash, J. N. Barker, M.-A. Perales, S. A.
Giralt, E. G. Pamer, M. R. M. van den Brink, Intestinal Blautia Is Associated with Reduced Death from
Graft-versus-Host Disease. *Biol Blood Marrow Transplant* **21**, 1373–1383 (2015).

26. Y. Furusawa, Y. Obata, S. Fukuda, T. A. Endo, G. Nakato, D. Takahashi, Y. Nakanishi, C.
Uetake, K. Kato, T. Kato, M. Takahashi, N. N. Fukuda, S. Murakami, E. Miyauchi, S. Hino, K. Atarashi,

S. Onawa, Y. Fujimura, T. Lockett, J. M. Clarke, D. L. Topping, M. Tomita, S. Hori, O. Ohara, T. Morita, H. Koseki, J. Kikuchi, K. Honda, K. Hase, H. Ohno, Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature* **504**, 446–450 (2013).

27. N. D. Mathewson, R. Jenq, A. V. Mathew, M. Koenigsknecht, A. Hanash, T. Toubai, K. Oravecz-Wilson, S.-R. Wu, Y. Sun, C. Rossi, H. Fujiwara, J. Byun, Y. Shono, C. Lindemans, M. Calafiore, T. M. Schmidt, K. Honda, V. B. Young, S. Pennathur, M. van den Brink, P. Reddy, Gut microbiome-derived metabolites modulate intestinal epithelial cell damage and mitigate graft-versus-host disease. *Nat Immunol* **17**, 505–513 (2016).

28. J. L. Golob, M. M. DeMeules, T. Loeffelholz, Z. Z. Quinn, M. K. Dame, S. S. Silvestri, M. C. Wu, T. M. Schmidt, T. L. Fiedler, M. J. Hoostal, M. Mielcarek, J. Spence, S. A. Pergam, D. N. Fredricks, Butyrogenic bacteria after acute graft-versus-host disease (GVHD) are associated with the development of steroid-refractory GVHD. *Blood Adv* **3**, 2866–2869 (2019).

29. M. Couturier, B. Lamarthée, J. Arbez, J.-C. Renauld, C. Bossard, F. Malard, F. Bonnefoy, M. Mohty, S. Perruche, P. Tiberghien, P. Saas, B. Gaugler, IL-22 deficiency in donor T cells attenuates murine acute graft-versus-host disease mortality while sparing the graft-versus-leukemia effect. *Leukemia* **27**, 1527–1537 (2013).

30. B. Lamarthée, F. Malard, C. Gamonet, C. Bossard, M. Couturier, J.-C. Renauld, M. Mohty, P. Saas, B. Gaugler, Donor interleukin-22 and host type I interferon signaling pathway participate in intestinal graft-versus-host disease via STAT1 activation and CXCL10. *Mucosal Immunol* **9**, 309–321 (2016).

31. M. A. Schroeder, J. F. DiPersio, Mouse models of graft-versus-host disease: advances and limitations. *Disease Models & Mechanisms* **4**, 318–333 (2011).

32. R. Browaeys, W. Saelens, Y. Saeys, NicheNet: modeling intercellular communication by linking ligands to target genes. *Nat Methods* **17**, 159–162 (2020).

33. D. R. Edwards, M. M. Handsley, C. J. Pennington, The ADAM metalloproteinases. *Mol Aspects Med* **29**, 258–289 (2008).

34. J. C. Albring, M. M. Sandau, A. S. Rapaport, B. T. Edelson, A. Satpathy, M. Mashayekhi, S. K. Lathrop, C.-S. Hsieh, M. Stelljes, M. Colonna, T. L. Murphy, K. M. Murphy, Targeting of B and T lymphocyte associated (BTLA) prevents graft-versus-host disease without global immunosuppression. *J Exp Med* **207**, 2551–2559 (2010).

35. Y. Sakoda, J.-J. Park, Y. Zhao, A. Kuramasu, D. Geng, Y. Liu, E. Davila, K. Tamada, Dichotomous regulation of GVHD through bidirectional functions of the BTLA-HVEM pathway. *Blood* **117**, 2506–2514 (2011).

36. F. Martins, E. Planet, D. Marino, M. Ansari, D. Trono, Single-cell transcriptome analysis reveals atypical monocytes circulating ahead of acute graft-versus-host disease clinical onset. *J Leukoc Biol*, qiae229 (2024).

37. H. Sun, L. Wu, X. Zhao, Y. Huo, P. Dong, A. Pang, Y. Zheng, Y. Han, S. Ma, E. Jiang, F. Dong, T. Cheng, S. Hao, Monocytes as an early risk factor for acute graft-versus-host disease after allogeneic hematopoietic stem cell transplantation. *Front Immunol* **15**, 1433091 (2024).

38. L. Jardine, U. Cytlak, M. Gunawan, G. Reynolds, K. Green, X.-N. Wang, S. Pagan, M. Paramitha, C. A. Lamb, A. K. Long, E. Hurst, S. Nair, G. H. Jackson, A. Publicover, V. Bigley, M. Haniffa, A. J. Simpson, M. Collin, Donor monocyte-derived macrophages promote human acute graft-versus-host disease. *J Clin Invest* **130**, 4574–4586 (2020).

39. K. Seno, M. Yasunaga, H. Kajiya, K. Izaki-Hagio, H. Morita, M. Yoneda, T. Hirofuji, J. Ohno, Dynamics of M1 macrophages in oral mucosal lesions during the development of acute graft-versus-host disease in rats. *Clin Exp Immunol* **190**, 315–327 (2017).

40. J. E. Levine, S. Paczesny, S. Mineishi, T. Braun, S. W. Choi, R. J. Hutchinson, D. Jones, Y. Khaled, C. L. Kitko, D. Bickley, O. Krijanovski, P. Reddy, G. Yanik, J. L. M. Ferrara, Etanercept plus methylprednisolone as initial therapy for acute graft-versus-host disease. *Blood* **111**, 2470–2475 (2008).

41. D. Jankovic, J. Ganesan, M. Bscheider, N. Stickel, F. C. Weber, G. Guarda, M. Follo, D. Pfeifer, A. Tardivel, K. Ludigs, A. Bouazzaoui, K. Kerl, J. C. Fischer, T. Haas, A. Schmitt-Gräff, A. Manoharan, L. Müller, J. Finke, S. F. Martin, O. Gorka, C. Peschel, J. Ruland, M. Idzko, J. Duyster, E. Holler, L. E. French, H. Poeck, E. Contassot, R. Zeiser, The Nlrp3 inflammasome regulates acute graft-versus-host disease. *J Exp Med* **210**, 1899–1910 (2013).

42. C.-K. Min, Y. Maeda, K. Lowler, C. Liu, S. Clouthier, D. Lofthus, E. Weisiger, J. L. M. Ferrara,
P. Reddy, Paradoxical effects of interleukin-18 on the severity of acute graft-versus-host disease
mediated by CD4+ and CD8+ T-cell subsets after experimental allogeneic bone marrow transplantation.
Blood **104**, 3393–3399 (2004).

43. H. Okamura, H. Tsutsi, T. Komatsu, M. Yutsudo, A. Hakura, T. Tanimoto, K. Torigoe, T.
Okura, Y. Nukada, K. Hattori, Cloning of a new cytokine that induces IFN-gamma production by T
cells. *Nature* **378**, 88–91 (1995).

44. Y. Fujimori, H. Takatsuka, Y. Takemoto, H. Hara, H. Okamura, K. Nakanishi, E. Kakishita,
Elevated interleukin (IL)-18 levels during acute graft-versus-host disease after allogeneic bone marrow
transplantation. *Br J Haematol* **109**, 652–657 (2000).

45. R. J. Robb, G. R. Hill, The interferon-dependent orchestration of innate and adaptive immunity
after transplantation. *Blood* **119**, 5351–5358 (2012).

46. S. Wang, T. Cheng, X. Chen, C. Zeng, W. Qin, Y. Xu, IFN-γ induces acute graft-versus-host
disease by promoting HMGB1-mediated nuclear-to-cytoplasm translocation and autophagic
degradation of p53. *Clin Sci (Lond)* **138**, 1287–1304 (2024).

47. F. T. Hakim, S. Memon, P. Jin, M. M. Imanguli, H. Wang, N. Rehman, X.-Y. Yan, J. Rose, J.
W. Mays, S. Dhamala, V. Kapoor, W. Telford, J. Dickinson, S. Davis, D. Halverson, H. B. Naik, K.
Baird, D. Fowler, D. Stroncek, E. W. Cowen, S. Z. Pavletic, R. E. Gress, Upregulation of IFN-Inducible
and Damage-Response Pathways in Chronic Graft-versus-Host Disease. *J Immunol* **197**, 3490–3503
(2016).

48. T. Ara, D. Hashimoto, E. Hayase, C. Noizat, R. Kikuchi, Y. Hasegawa, K. Matsuda, S. Ono, Y.
Matsuno, K. Ebata, R. Ogasawara, S. Takahashi, H. Ohigashi, E. Yokoyama, K. Matsuo, J. Sugita, M.
Onozawa, R. Okumura, K. Takeda, T. Teshima, Intestinal goblet cells protect against GVHD after
allogeneic stem cell transplantation via Lypd8. *Science Translational Medicine* **12**, eaaw0720 (2020).

49. Q. Song, U. Nasri, D. Zeng, Steroid-Refractory Gut Graft-Versus-Host Disease: What We Have
Learned From Basic Immunology and Experimental Mouse Model. *Front. Immunol.* **13** (2022).

50. A. S. Henden, M. Koyama, R. J. Robb, A. Forero, R. D. Kuns, K. Chang, K. S. Ensbey, A.
Varelias, S. H. Kazakoff, N. Waddell, A. D. Clouston, R. Giri, J. Begun, B. R. Blazar, M. A. Degli-
Esposti, S. V. Kotenko, S. W. Lane, K. L. Bowerman, R. Savan, P. Hugenholtz, K. H. Gartlan, G. R.
Hill, IFN-λ therapy prevents severe gastrointestinal graft-versus-host disease. *Blood* **138**, 722–737
(2021).

51. S. W. Choi, G. C. Hildebrandt, K. M. Olkiewicz, D. A. Hanauer, M. N. Chaudhary, I. A. Silva,
C. E. Rogers, D. T. Deurloo, J. M. Fisher, C. Liu, D. Adams, S. W. Chensue, K. R. Cooke, CCR1/CCL5
(RANTES) receptor-ligand interactions modulate allogeneic T-cell responses and graft-versus-host
disease following stem-cell transplantation. *Blood* **110**, 3447–3455 (2007).

52. S. DeWolf, Y. Elhanati, K. Nichols, N. R. Waters, C. L. Nguyen, J. B. Slingerland, N.
Rodriguez, O. Lyudovyk, P. A. Giardina, A. I. Kousa, H. Andrlová, N. Ceglia, T. Fei, R. Kappagantula,
Y. Li, N. Aleynick, P. Baez, R. Murali, A. Hayashi, N. Lee, B. Gipson, M. Rangesa, Z. Katsamakis, A.
Dai, A. G. Blouin, M. Arcila, I. Masilionis, R. Chaligne, D. M. Ponce, H. J. Landau, I. Politikos, R.
Tamari, A. M. Hanash, R. R. Jenq, S. A. Giralt, K. A. Markey, Y. Zhang, M.-A. Perales, N. D. Socci,
B. D. Greenbaum, C. A. Iacobuzio-Donahue, T. J. Hollmann, M. R. M. van den Brink, J. U. Peled,
Tissue-specific features of the T cell repertoire after allogeneic hematopoietic cell transplantation in
human and mouse. *Sci Transl Med* **15**, eabq0476 (2023).

53. K. I. Omdahl, R. S. Bermea, R. Fleming, K. Kimler, J. Kaminski, L. P. Hariri, A. Ly, X. Rui, L.
Cagnin, J. Lane, U. Gerdemann, B. R. Blazar, V. Tkachev, L. S. Kean, Organ-specific
microenvironments drive divergent T cell evolution in acute graft-versus-host disease. *Sci Transl Med*
17, eads1298 (2025).

54. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk,
C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T.
Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert, R.
Satija, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

55. G. Korotkevich, V. Sukhov, N. Budin, B. Shpak, M. N. Artyomov, A. Sergushichev, Fast gene
set enrichment analysis. bioRxiv [Preprint] (2021). https://doi.org/10.1101/060012.

56. A. Liberzon, C. Birger, H. Thorvaldsdóttir, M. Ghandi, J. P. Mesirov, P. Tamayo, The Molecular

Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417–425 (2015).

57. J. A. Ramilowski, T. Goldberg, J. Harshbarger, E. Kloppmann, M. Lizio, V. P. Satagopam, M. Itoh, H. Kawaji, P. Carninci, B. Rost, A. R. R. Forrest, A draft network of ligand-receptor-mediated multicellular signalling in human. *Nat Commun* **6**, 7866 (2015).

58. A. J. Pawson, J. L. Sharman, H. E. Benson, E. Faccenda, S. P. H. Alexander, O. P. Buneman, A. P. Davenport, J. C. McGrath, J. A. Peters, C. Southan, M. Spedding, W. Yu, A. J. Harmar, NC-IUPHAR, The IUPHAR/BPS Guide to PHARMACOLOGY: an expert-driven knowledgebase of drug targets and their ligands. *Nucleic Acids Res* **42**, D1098-1106 (2014).

59. A. D. Rouillard, G. W. Gundersen, N. F. Fernandez, Z. Wang, C. D. Monteiro, M. G. McDermott, A. Ma'ayan, The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016**, baw100 (2016).

60. K. Street, D. Risso, R. B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, S. Dudoit, Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19**, 477 (2018).

61. K. Van den Berge, H. Roux de Bézieux, K. Street, W. Saelens, R. Cannoodt, Y. Saeys, S. Dudoit, L. Clement, Trajectory-based differential expression analysis for single-cell sequencing data. *Nat Commun* **11**, 1201 (2020).

62. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

63. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, G. Sherlock, Gene Ontology: tool for the unification of biology. *Nat Genet* **25**, 25–29 (2000).

64. The Gene Ontology Consortium, S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry, H. J. Drabkin, D. Ebert, M. Feuermann, P. Gaudet, N. L. Harris, D. P. Hill, R. Lee, H. Mi, S. Moxon, C. J. Mungall, A. Muruganugan, T. Mushayahama, P. W. Sternberg, P. D. Thomas, K. Van Auken, J. Ramsey, D. A. Siegele, R. L. Chisholm, P. Fey, M. C. Aspromonte, M. V. Nugnes, F. Quaglia, S. Tosatto, M. Giglio, S. Nadendla, G. Antonazzo, H. Attrill, G. dos Santos, S. Marygold, V. Strelets, C. J. Tabone, J. Thurmond, P. Zhou, S. H. Ahmed, P. Asanitthong, D. Luna Buitrago, M. N. Erdol, M. C. Gage, M. Ali Kadhum, K. Y. C. Li, M. Long, A. Michalak, A. Pesala, A. Pritazahra, S. C. C. Saverimuttu, R. Su, K. E. Thurlow, R. C. Lovering, C. Logie, S. Oliferenko, J. Blake, K. Christie, L. Corbani, M. E. Dolan, H. J. Drabkin, D. P. Hill, L. Ni, D. Sitnikov, C. Smith, A. Cuzick, J. Seager, L. Cooper, J. Elser, P. Jaiswal, P. Gupta, P. Jaiswal, S. Naithani, M. Lera-Ramirez, K. Rutherford, V. Wood, J. L. De Pons, M. R. Dwinell, G. T. Hayman, M. L. Kaldunski, A. E. Kwitek, S. J. F. Laulederkind, M. A. Tutaj, M. Vedi, S.-J. Wang, P. D'Eustachio, L. Aimo, K. Axelsen, A. Bridge, N. Hyka-Nouspikel, A. Morgat, S. A. Aleksander, J. M. Cherry, S. R. Engel, K. Karra, S. R. Miyasato, R. S. Nash, M. S. Skrzypek, S. Weng, E. D. Wong, E. Bakker, T. Z. Berardini, L. Reiser, A. Auchincloss, K. Axelsen, G. Argoud-Puy, M.-C. Blatter, E. Boutet, L. Breuza, A. Bridge, C. Casals-Casas, E. Coudert, A. Estreicher, M. Livia Famiglietti, M. Feuermann, A. Gos, N. Gruaz-Gumowski, C. Hulo, N. Hyka-Nouspikel, F. Jungo, P. Le Mercier, D. Lieberherr, P. Masson, A. Morgat, I. Pedruzzi, L. Pourcel, S. Poux, C. Rivoire, S. Sundaram, A. Bateman, E. Bowler-Barnett, H. Bye-A-Jee, P. Denny, A. Ignatchenko, R. Ishtiaq, A. Lock, Y. Lussi, M. Magrane, M. J. Martin, S. Orchard, P. Raposo, E. Speretta, N. Tyagi, K. Warner, R. Zaru, A. D. Diehl, R. Lee, J. Chan, S. Diamantakis, D. Raciti, M. Zarowiecki, M. Fisher, C. James-Zorn, V. Ponferrada, A. Zorn, S. Ramachandran, L. Ruzicka, M. Westerfield, The Gene Ontology knowledgebase in 2023. *Genetics* **224**, iyad031 (2023).

Acknowledgments: The authors thank the patients who agreed to participate in this study by providing their blood samples. We also thank the iGenSeq platform team at ICM for sequencing the PBMC samples. We are grateful to the CRYOSTEM Consortium and the SFGM-TC for collecting patient samples and clinical data. **Funding:** This project was funded by la Fédération Leucémie Espoir and Fondation Maladies Rare (HPN-AM). **Author contributions:** Conceptualization: SLG, DM; Methodology: SLG, NV, DM; Samples processing: SLG, DB, YM, EB; Investigation: SLG, NV, DM; Visualization: SLG, GS, RPD, NV, DM; Funding acquisition: DM; Project administration: EB, DM, GS, RPD; Supervision: NV, DM; Writing – original draft: SLG, NV, DM; Draft revision: SLG, YM, DB, EB, RPF, GS, NV, DM. **Competing interests:** DM received research grants from Novartis and Sanofi, and consulting fees from Sanofi, Incyte, Novartis, Jazz Pharmaceuticals, CSL Behring and Mallinckrodt. Other authors declared that they have no competing interests. **Data availability:** Raw data are available on GEO public repository: GSE229733. Source code to reproduce the analyses, figures, and tables described in this manuscript are provided in a Git repository: https://gitlab.com/SophieLG/gvh_transcriptomic and long term preserved in Software Heritage:

https://archive.softwareheritage.org/browse/origin/directory/?origin_url=https://gitlab.com/SophieLG/gvh_transcriptomic×tamp=2024-12-09T14:03:56.753000%2B00:00

Figure 1: Overview of patient and cell distribution, cluster identification, and relative

abundance of immune cell populations.

A: Experimental design: HSCT patients (n=53) with (n=21) and without aGVHD (n=32) and

healthy donor as control group (n=6) were analyzed. PBMC were collected at day 100 post

HSCT for patients without aGVHD, and before steroid initiation for patients with aGVHD.

B: Number of cells included in the analysis after CITE-Seq processing. **C:** Uniform Manifold

Approximation and Projection (UMAP) after principal component analysis for immune cells

clustering, according to surface markers. **D:** Heatmap depicting immune cluster's identification

through surface antigen expression. Three irrelevant populations were removed and are

shown in **Figure S1**. **E**: Volcano plot exhibiting immune cells abundance in GVHD or

non-GVHD condition. Y axis reflects que -log 10(adjusted p-value) from differential expression

output, using non-parametric Mann-Whitney-Wilcoxon rank sum test. X axis reflects the

log(ratio) of median abundance in aGVHD condition, over non-aGVHD condition. Abbreviation:

CITE-Seq: Cellular indexing of transcriptomes and epitopes sequencing. aGVHD: acute graft

versus host disease. HSCT: allogeneic hematopoietic stem cell transplantation. PBMC:

Peripheral Blood Mononuclear Cells. SS-aGVHD: steroid sensitive acute graft versus host

disease. SR-aGVHD: steroid resistant acute graft versus host disease.

Figure 2: Immunotypes distinguished different immune patterns associated with aGVHD incidence.

A: Principal coordinates analysis (PCoA) revealed 4 different immune clusters (immunotypes)

according to cell population composition. **B:** Distribution of aGVHD patients among

immunotypes (left panel) and corticosteroid sensitivity (right panel). Fisher test was used for

statistical analysis. **C:** GVHD cumulative incidence with death as competing risk, estimated by

Kaplan-Meir method, Log rank test was used for statistical analysis. Immunotype 4 cumulative

incidence at day 100 = 72%, Immunotype 3 cumulative incidence at day 100 = 0%,

Immunotype 2 cumulative incidence at day 100 = 38%, Immunotype 1 cumulative incidence at

day 100 = 55%, log-rank p-value = 0.005. **D:** Bar plots showing top 15 contributing immune

subsets to immunotypes. Contribution was determined with permutational multivariate analysis

of variance (PERMANOVA).

Figure 3: Cellular function analysis identified distinct features associated with

SR-GVHD.

A: Volcano plots exhibiting cell cycle analysis after cell cycle phases scoring. X axis represents

log ratio of median number of cells in each phase in aGVHD patients over non GVHD patients.

Y axis represents log adjusted p values. Immune population significantly increased or

decreased are marked in red. Mann-Whitney-Wilcoxon tests were performed for statistical

analysis, and p values were adjusted using Benjamini and Hochberg method. **B:** Pathways

enrichment analysis comparing SR and SS-aGVHD, using gene set enrichment method and

"hallmarks" gene reference data set. Only significant values with adjusted p-value < 0.05 are

shown. NES: normalized enrichment score. SS-aGVHD: steroid sensitive acute graft versus

host disease. SR-aGVHD: steroid resistant acute graft versus host disease.

Figure 4: Cell-cell interactions with CD8+ T cells as receiver revealed specific patterns

associated with SR-aGVHD

NicheNet Output when focusing on CD8+ T cells as receiver is depicted on panel A, B and C.

A: Expression of top-ranked ligands on sender cells. Prioritized ligands identified are exhibited

on Y axis, and sender cells on X axis. The color determines the average expression and the

point size the percent of senders expressing the ligand. **B:** Relative expression of ligands on

sender cells. Expression is exhibited in log fold change (red: increase expression in

SR-aGVHD patients, Blue: decreased expression in SR-aGVHD patients). **C:** Ligand–target

matrix exhibiting the regulatory potential of interactions. On the X axis is shown the active

targets genes of the top-ranked ligands. Heatmap visualization showing which top-ranked

ligands are predicted to have regulated the expression of which differentially expressed genes.

D: Circos plots depicting interactions between senders and all receiver cells. Ligands are

shown on the left side of the circos plot, and targets on the right side. Only *bona fide*

interactions were selected (**Figure S9A**). Links exhibits positive log fold change (left panel,

increased in SR-aGVHD patients) or negative log fold change (right panel, increased in

SS-aGVHD patients) from "ligand differential expression heatmap" NicheNet's output. The

thickness of the link represents the numbers of sender populations involved in the interaction.

SR-aGVHD: Steroid resistant acute graft versus host disease. SS-aGVHD: Steroid sensitive

acute graft versus host disease.

Figure 5: SR-aGVHD is associated with specific trajectory in CD8⁺ T cells

A: Density plot for number of cells according to lineage 1 pseudotime visualization. SR-aGVHD cell abundance was marked in low and high pseudotimes, while SS-aGVHD was associated with a continuous increase in cell abundance along differentiation. **B:** Lineage 1 trajectory on RNA clustering. **C:** Heatmap depicting gene expression among the top 500 most expressed genes in SS-aGVHD patient (top panel) and in SR-aGVHD patients (low panel), along pseudotime. The color at the top of each heatmap represents the immune population in which the gene is expressed. **D:** Expression along pseudotime of four of the top 100 genes, whose expression varied the most between the start and end points of a lineage. **E:** Over-representation analysis to identify pathways associated with genes expression from each module of the heatmaps. Only pathways associated with adjusted pval < 0.05 from "molecular functions" and "biological process" of the Gene Ontology knowledgbase(*63, 64*) were selected for dot plots visualization. SR-aGVHD: Steroid resistant acute graft versus host disease. SS-aGVHD: Steroid sensitive acute graft versus host disease.

Figure 6: Steroid resistance is driven by biological pathways and specific networks of

interaction

Steroid-resistant aGVHD results of cytokine production by myeloid populations, especially

classical monocytes producing CCL3, CCL4, and IL18, which stimulate the activation of naive

CD8$^+$ T cells and induce direct differentiation into effector T cells. By contrast, steroid-sensitive

aGVHD is mainly driven by IL15 and IL12 production from classical monocytes and B cells,

and multiple transition states, from naïve CD8$^+$ T cells to activated EM CD8$^+$ T cells. aGVHD:

acute graft versus host disease, CM CD8$^+$ T cells: Central Memory CD8$^+$ T cells, EM CD8$^+$ T

cells: Effector Memory CD8$^+$ T cells, TEMRA CD8$^+$ T cells: Terminally differenciated Effector

Memory CD8$^+$ T cells