Corticosteroid resistance is predetermined by early immune response dynamics at acute Graft-Versus-Host disease onset

One Sentence Summary (150 characters): Steroid-resistant acute graft versus host disease is set at disease onset as evidenced by specific transcriptomic signature, cell-to-cell crosstalk and trajectories.

Sophie Le Grand[1], Yannick Marie[2], Delphine Bouteiller[2], Emilie Robert[3], Régis Peffault de Latour[3,4], Gérard Socié[1,4], Nicolas Vallet[5]*, David Michonneau[1,4]*

Affiliations:

[1]INSERM UMR 1342, Translational Immunology in Immunotherapy and Hematology, Paris Cité University, F-75010, Paris, France

[2]Genotyping and Sequencing Facility, Paris Brain Institute-ICM, Hôpital de la Pitié-Salpêtrière, CNRS UMR 7225, INSERM U1127, Sorbonne Université UM75, CS21414, Paris, France.

[3]Cryostem Consortium.

[4]Hematology Transplantation, Saint Louis Hospital, Assistance Publique des Hôpitaux de Paris, 75010 Paris, France

[5]Hematology and Cell Therapy department, University Hospital of Tours, Inserm U1069 N2COx

*co-last authors

Corresponding author:

David Michonneau

Saint Louis hospital, hematology and transplantation

1 avenue Claude Vellefaux

75010 PARIS, FRANCE

david.michonneau@aphp.fr

Abstract

Corticosteroid-resistant acute graft versus host disease (SR-aGVHD) is the leading life-threatening complication following allogeneic hematopoietic stem cell transplantation. Novel therapeutics development is impeded by scares knowledge on biological pathways leading to steroid resistance at time of aGVHD diagnosis. To gain insight into our understanding on circulating immune cell subsets and functions at time of aGVHD and before steroid initiation, a single cell deep phenotyping and transcriptome analysis was performed on peripheral blood mononuclear cells from aGVHD patients before first line treatment, and from patients without aGVHD. We first explored immune cell subsets as a global signature, called immunotype. One immunotype enriched in monocytes expressing CD11b was associated with increased incidence of aGVHD (72% at day 100, log-rank p=0.005), but immunotypes were not associated with steroid sensitivity. Pathway analysis and inferred ligand/receptor interactions revealed major divergence between steroid sensitive and SR-aGVHD, including enrichment of TNFα activation in SR-GVHD, as well as TNF/TNFR, CCL3, CCL4 and IL18 signaling. Interferon α and γ pathways activation were decreased in SR-aGVHD patients. Finally, trajectory inferences within CD8$^+$ T cells evidenced specific transition during SR-aGVHD, from an early differentiated state to a highly activated and differentiated one, without slight differentiation across intermediate stages, involving specific gene signature during state transition. These findings provide evidence that corticosteroid resistance is an intrinsic mechanism already present at the onset of alloimmune response. This comprehensive overview of pathways underlying corticosteroid resistance highlights specific patterns that may serve as potential new therapeutic targets.

1 INTRODUCTION

Allogeneic hematopoietic stem cell transplantation (HSCT) is a major treatment for

hematological malignancies. Its therapeutic effect is based on alloreactivity of donor T cells

against tumor cells. However, donor T cells also react against healthy host cells, which may

lead to graft-versus-host disease (GVHD). Acute GVHD (aGVHD) may involve skin, gut, or

liver tissue. Its pathophysiology is initiated after tissue damage from high dose alkylating agent

and radiation based-conditioning regimens leading to recognition of non-self-major and minor

histocompatibility antigens (MHC and mHag(*1*)) and an alloreactive immune response,

exacerbated by altered tissue repair mechanisms and microbiome dysbiosis(*2*). Despite first

line treatment with prompt corticosteroid initiation(*3, 4*), more than 40% of patients are steroid

resistant (SR-aGVHD)(*5–7*). Steroid resistance is defined by an absence of clinical response

after 7 days of standard dose of corticosteroids, a progressive disease after 3 days,

progressive disease with new organ involvement or recurrence while steroids are stopped or

tapered. Although ruxolitinib improves symptoms for 60% of SR-GVHD patients(*8*), their

prognosis is very dismal(*3, 9, 10*).

While biomarkers were recently developed to improve aGVHD diagnosis and risk

stratification(*11–15*), response to steroid is still unpredictable and biological mechanisms

underlying steroid resistance remains only partly understood, even in the few experimental

transplantation models of SR–GVHD described(*16–18*). In humans, specific whole gene

expression analysis at diagnosis of lower intestinal acute SR-GVHD identified an increased

repair-associated gene (amphiregulin and the aryl hydrocarbon receptor) and M2

macrophages pathways(*19*), suggesting that steroid resistance in not restricted to the T cells

response as initially suspected. Indeed, neutrophils participate in tissue damage at GVHD

onset and triggering, through reactive oxygen species production(*20*) and promote T cell

expansion(*21*) and differentiation to Th17 cells(*22, 23*). As corticosteroids increase neutrophils

count and development, it has been hypothesized that corticosteroids may trigger a positive

feedback loop between pathogenic T cells and neutrophils in SR-GVHD context(*24*).

Furthermore, neutrophil-predominant noncanonical inflammation pathways were recently

identified in gut-SR-GVHD biopsies, notably ubiquitin specific peptidase 17–like family of

genes(25). While gut microbiota's role has been extensively studied in GVHD, it is not well

understood in SR-GVHD. Butyrate, a short-chain-fatty-acid product from *Blautia* bacteria is

associated with better long terms outcomes in aGVHD(26–28), but one study found a negative

association between butyrate production and SR-GVHD, due to the ability of butyrate to impair

human colonic stem cells from recovering epithelial monolayer(29). Other studies identified

SR-GVHD biomarkers such as IL-22(16), but with controversial potential(30, 31). Most of these

studies rely on murine models(16–18, 32), and there is currently no clear overview of immune

subsets associated with steroid resistance in human GVHD.

We hypothesize that reshaping of immune system might be initiated early in

alloimmune responses, therefore predisposing to corticosteroids refractoriness. To describe

immune subsets and their respective function, cellular indexing of transcriptomes and epitopes

sequencing (CITE-Seq) was performed on peripheral blood mononuclear cells (PBMC)

samples collected at the onset of aGVHD before first line treatment, or in patient without GVHD

at day 100 after HSCT. Whole landscape of circulating immune cells at early onset of aGVHD

and its association with GVHD severity was uncovered. Deep analysis of transcriptomic

profiles revealed that steroid resistance is an intrinsic property of the immune response,

characterized by specific enriched biological pathways, specific cell-to-cell crosstalk through

ligand-receptor interactions and a specific immune trajectory involving most immune subsets.

Altogether, our results highlight that steroid-resistant GVHD is an early characteristic of allo-

immune response before any immunosuppressive treatment and pave the way to more

targeted first-line therapy to improve allogeneic HSCT outcomes.

RESULTS

Cohort's characteristics

Samples from 53 patients were retrieved from the prospective CRYOSTEM multicenter collection. PBMC were collected before corticosteroids initiation for patients diagnosed with aGVHD, and at day 100 post allo-HSCT for non-GVHD patients. Hematological malignancies, HSCT procedure and clinical characteristics were balanced between groups (**Table S1)**. Among the 21 aGVHD patients, 16 (76%) were steroid sensitive (SS-GVHD) and 5 (24%) were steroid refractory (SR-GVHD) (**Figure 1A**). For validation and control purposes, 6 samples from healthy blood donor were used. After samples preparation for CITE-seq experiments, sequencing, quality control filtering and data processing acquisition, more than 300,000 single cells were available for immune profiling and transcriptome analysis (**Figure 1B**). Unsupervised clustering resulted in the identification of 40 immune cell subsets (**Figure 1C and S1**), characterized by their surface antigens expression (**Figure 1D**).

Immune signature and increased proliferation associate with aGVHD onset but not with response to steroids

Compared to non-aGVHD patients, classical monocytes CD73$^+$ were significantly enriched in aGVHD patients, whereas NK and B cell subsets were significantly decreased (**Figure 1E**). To encompass the compositional structure of immune subsets frequencies, principal coordinates analysis (PCoA) was used. Four distinct immune clusters were identified, thereafter called immunotypes (**Figure 2A**). aGVHD patient's distribution was significantly different among immunotypes (p=0.001, **Table S2**, **Figure 2B**), as well as the GVHD cumulative incidence (**Figure 2C,** Immunotype 4 GVHD incidence = 72% at day 100, *vs* immunotype 1 = 55%, *vs* immunotype 2 = 38%, *vs* immunotype 4 = 0%, global log-rank p=0.005). GVHD-free and relapse-free survival (GRFS) was significantly associated with immunotypes (p=0.0048, **Figure S2**), however there were no significant differences in non-relapse mortality and

cumulative incidence of relapse (**Figure S2**). Association with other clinical variables is

depicted in **Figure S3**.

Immunotype 1 and 4 included a large number of patients with aGVHD (**Figure 2B–C**),

and were characterized by an overall reduction in immune cell heterogeneity, illustrated with

an imbalanced immune subsets representation (**Figure 2D**). Especially, immunotypes 1 and 4

presented a high display of classical monocytes CD11b^{bright} (IgM$^+$ or IgM$^-$) compared to other

immunotypes. Immunotype 2 and 3 were characterized by a balanced distribution across

different immune cell populations, and a lower classical monocytes CD11b^{bright} cells

representation. Remarkably, immunotype 3 was characterized by an absence of patients with

aGVHD, a decrease in classical monocytes CD11b^{bright} and in other monocytes populations,

whereas effector T CD4$^+$ and CD8$^+$ were predominantly increased.

To further investigate whether the over-representation may encompass a high cell

division rate at time of aGVHD diagnosis, we determined cell cycle phase scoring within each

immune cell population. Immune populations were predominantly in S phase (**Figure 3A**),

reflecting increase cell proliferation at aGVHD onset.

No significant differences among immune cells abundances, immunotypes and

cell-cycle scoring was observed with respect to corticosteroid resistance (**Figure 2B**, **Table**

S2, Figure S4 and S5). In summary, immune cells subset clearly identified patients without

GVHD but hardly discriminated SS and SR-GVHD. We thus hypothesized that cell intrinsic

biological functions may predispose to SR-GVHD.

Steroid resistance is associated with enrichment of TNFα and metabolic pathways at

aGVHD onset

We first considered the biological pathways enriched at the onset of aGVHD by

comparison with patients without GVHD (**Figure S6**). When considering aGVHD as a whole,

we observe overactivation of IFNα and γ pathways in all immune cells, and a decrease in

TNFα signaling, especially in CD8$^+$ T cells and myeloid cells.

To study which biological process may correlate with subsequent SS- or SR-aGVHD,

we then performed a second enrichment analysis in patients with SS- or SR-aGVHD.

Compared to SS-aGVHD, inflammation pathways were enriched in SR-aGVHD patients within

all immune cells; particularly TNFα and hypoxia pathways (**Figure 3B**). STAT5 and MTORC

pathways were particularly upregulated in exhausted effector memory CD4$^+$ and CD8$^+$ T cells,

in naïve CD8$^+$ T cells and in dendritic immune cells. The IL6/JAK STAT pathway was also

highly enriched in naive CD8$^+$ T cells and classical CD11b^{bright} monocytes.

Genes associated with metabolism pathways were overexpressed, especially oxidative

phosphorylation in naïve CD4$^+$ T cells, double negative T cells and myeloid cells. Conversely,

there was a decreased activation of the IFNα and γ responses within all immune subsets,

except NK cells.

Cell to cell crosstalk during SR-GVHD specifically involves IL1, CCL3, CCL4, TNFα and

IL18 signaling

To address whether cell biological processes involved in SR-aGVHD may be involved in

cell-to-cell interactions leading to early steroid-resistance, we examined intercellular

communication using a computational inferring method(*33*). When focusing on CD8$^+$ T cells

as receiver, 20 ligands were identified as prioritized ligands, and their expression on sender

cells is shown in **Figure 4A**. IL1B, CCL3 and CCL4 ligands were particularly overexpressed in

SR-aGVHD condition (**Figure 4B**). Regulatory potential of each ligand with predicted targets

genes in shown in **Figure 4C**.

Predicted ligands and targets interaction were then analyzed for CD4 regulatory T cells

(**Figure S7A**), NK cells (**Figure S7B**), CD4$^+$T cells (**Figure S8A**), and double negative T cells

(**Figure S8B**). An overview of all these interactions is depicted as a circus plot (**Figure 4D**)

based on biologically confirmed interactions previously designated as *bona fide* interactions

(**Figure S9A**). TNF ligands-to-cell signaling appeared to be overexpressed in most sender

cells in SR-aGVHD patients, as well as CCL3, CCL4 and IL18. ADAM17, which is involved in

TNF activation(*34*), was overexpressed in all senders and interacted with TNFR, ITGAL and

ITGB2 receptors. Interestingly, IFN/IFNR interactions was enhanced across all receivers in

SR-aGVHD but absent in SS-aGVHD.

Conversely, when comparing patients with aGVHD to those without aGVHD, immune

cells crosstalk appeared different to that disclosed in the SS- and SR–aGVHD setting

(**Figure S9B**). Consistent with enrichment analysis, TNF/TNF-R interaction was decreased

among DN T cells, NK cells and T regulatory lymphocytes. TGFB1/TGFBR interactions, which

are considered as being associated with an immunoregulatory role, were significantly

increased in patient with no aGVHD compared to aGVHD patients. Especially within CD8[+] T

cells, as well as BTLA/TNFRSF14 interaction, a pathway previously identified in animal models

as a regulator of alloimmune response(*35, 36*).

Specific trajectories were associated with SR-GVHD

As biological processes and cell-to-cell interactions targeting CD8[+] T cell were associated with

SR-aGVHD, we hypothesized that immune cells subsets from SR-aGVHD and SS-aGVHD

patients may exert specific CD8[+] T cell trajectories. To address this hypothesis, trajectories

within CD8[+] T cell were defined on a new cell clustering based on mRNA expression

(**Figure S10**). Two trajectories were identified: (i) lineage 1 described in **Figure 5A-B** and (ii)

lineage 2 described in **Figure S11**.

Frequency of cells in lineage 1 slightly increased over pseudotime in SS-aGVHD

patients through multiple transition states starting from naïve double positive T cell (cluster 27)

to effector memory CD8[+] T cell expressing ICOS (cluster 2) **(Figure 5A-B).** By contrast,

SR-aGVHD patients were characterized by a direct transition with very few cells in the

transition subsets (cluster 15, 9 and 10), most immune subset being grouped in cluster 27 at

early pseudotime or in cluster 2 in late pseudotime. The same pattern was observed in the

second CD8[+] T cell trajectory, with a transition from naïve double positive T cell (cluster 27) to

CD8[+] TEMRA cells expressing ICOS and PD1 (cluster 25) (**Figure S11A-B**).

To highlight different gene expression pattern in SS-aGVHD and SR-aGVHD we first

studied individual genes whose expression varied significantly along pseudotime and across

transition state (**Figure 5C, S11C, Table S3-S10**). Among the most significant genes involved

in transition states, we identified CCL4, CCL5, IFI6, LAG3, and TIGIT (**Table S11 and S12**).

SR-aGVHD patients exhibited earlier and higher CCL4 and LAG3 expression in pseudotime

(**Figure 5D and S11D**). Consistently with pathway enrichment analysis, IFI6 expression,

indicative of IFNα activation, was higher in SS-aGVHD patients, particularly in early-

differentiated CD8[+] T cells.

To study overall biological processes, we used hierarchical clustering to identify

modules of correlated gene among the top 500 expressed gene along pseudotime lineages.

This approach identified 5 modules of genes involved in cell-to-cell transition (**Figure 5C**).

Over-representation analysis (ORA) from Gene Ontology knowledgebase within each cell

subset highlighted the main biological pathways involved at each transition state (**Figure 5E**).

SR-aGVHD was notably associated with an early enrichment of pathways associated with cell

activation and high transcriptional and translational activity in naïve double positive T cell

(**Figure S11E**). A detailed analysis of pathways enriched in each module revealed striking

differences in the dynamics of T cell activation during SR-GVHD (trajectory 1 in **Figure S12**

and S13, and trajectory 2 in **Figure S14 and S15**).

Overall, these results reveal that even before starting corticosteroids, aGVHD is already

characterized by different trajectories and pattern of gene activation between transition states,

that can define a specific dynamic of alloimmune response during steroid resistance.

DISCUSSION

Despite recent advances in the management of aGVHD, steroid-resistant treatment remains a challenge. This study is the first to describe immune subsets and transcriptional profiles in human at time of aGVHD. Early identification of patients at risk of developing steroid resistance is crucial for rapid and tailored therapeutic intervention. Up to now, a comprehensive understanding of the circulating immune cells mechanisms associated with steroid resistance has not been addressed. Our results highlight that steroid resistance is characterized at the onset of aGVHD, by a specific transcriptomic profile that reveal biological pathways, cell-to-cell interactions and trajectories that differ from steroid sensitive patients.

Immunotypes and cell cycle scoring enabled the identification of specific patterns associated with aGVHD compared to non-aGVHD patients. Immunotype 4, characterized by poor immune cell diversity and by classical CD11[+] monocytes abundance was associated with worsened aGVHD free survival. This observation is consistent with previous studies identifying monocytes and macrophages as fully parts of the immune response at aGVHD onset (*19, 37–40*).

However, cell subsets distribution did not allow to distinguish SS- and SR-aGVHD. When focusing on immune cells functions rather than distinct phenotypes, we found specific patterns associated with SR-GVHD, especially higher TNFα pathway activation. This was strengthened with inferred interaction analyses which revealed an increased TNF/TNFR interaction, notably in CD8[+] T cells. Although this *in silico* prediction would need to be confirmed *in vivo*, these results might suggest that early anti-TNF therapies could be beneficial in patients with the corticosteroid resistance profile we described herein. This is supported by previous clinical phase 2 trial testing a combined therapy of anti-TNF and corticosteroids, with increased response rate (69% vs 33%; P < 0.001) without increased infections risk(*41*).

Interferon have been described for their crucial role in aGVHD development (*42, 43*). We found that IFNα and γ pathways were activated at time of aGVHD and before corticosteroid treatment. Unexpectedly, SR-aGVHD was associated with lower activation of IFNα and γ

pathways, and INFG/IFNGR interactions were increased in SR-aGVHD setting. This suggests

that the IFN signals might be blocked after receptor activation. Moreover, IFI6, a protein-coding

gene for IFNα, is expressed earlier and at higher levels during T cell differentiation in

SS-aGVHD.

Cross-talk analysis revealed numerous interactions specifically associated with

corticosteroids resistance, especially with cytokines/cytokines-receptors such as IL18, CCL4,

CCL5. Increased in CCL4 and CCL5 interactions within CD8[+] T cells populations suggest the

recruitment of effectors to inflammatory sites. IL18 is a proinflammatory cytokine which can

promote interferon γ production(*44*), and was already described as an important player in

aGVHD pathogenesis(*45*), as serum levels in post-HSCT patients were associated with

aGVHD severity (r = 0.861, P < 0.001)(*45*).

Finally, trajectory inferences highlighted specific layouts associated with SR-aGVHD.

Cells were particularly abundant in low and high pseudotime in SR-GVHD patients, while there

was a progressive increase along pseudotime in SS-GVHD patients. It suggests that cells in

SR-GVHD patients quickly differentiate to terminally activated states. This was consistent with,

higher and earlier gene expression of LAG3 and TIGIT in SR-GVHD patients. Consistent with

results from cell-cell interactions analysis, CCL4 and CCL5 were expressed at higher levels

and earlier in SR-GVHD patients, as already described in animal model(*46*).

Because samples were collected before cortico-steroid initiation, our work sheds light

on early signature pathways leading to corticosteroid resistance. These pathways appeared

distinct from those involved in the general inflammatory response of acute GVHD and could

be targeted as early as steroid initiation. Transcriptomic analysis offered a powerful approach

to unravel immune cells functionality, combined with precise phenotypic population

identification, allowing for the identification of cellular heterogeneity in SR-GVHD immune

response. The complexity of the immune response in the context of GVHD was already

investigated at the single cell level, through T cell subset analysis following TCR clonotype

mapping(*47*). This elegant approach, though limited by patient's number, revealed that each

patient possessed a highly intricate repertoire, with different tissues exhibiting distinct clonal

populations, varying degrees of repertoire overlap between sites, and diverse numbers of

dominant clones shared across multiple tissues(*47*). Another study investigated the role of

Friend virus leukemia integration 1 in aGVHD context and found that this virus was able to

activate and differentiate alloreactive CD4+ T cell(*48*).

Our study offers new insights into the immune complexity, presenting a broader

perspective of the immune response. Our findings are partly limited by the use of models that

would require additional validation *in vivo*. Moreover, additional studies focusing on immune

infiltrates in target organs of GvHD would certainly help to complete our understanding of acute

GVHD pathophysiology, and especially of mechanisms associated with corticosteroid

resistance.

Overall, these findings provide evidence that corticosteroid resistance is an intrinsic

mechanism already present at the onset of alloimmune response, highlighting specific patterns

that may serve as potential new therapeutic targets.

MATERIALS AND METHODS

Cohort

This study was conducted with samples from patients and healthy donors. Patients were

included in the CRYSOTEM cohort after validation by the scientific committee (study number

CS-1801). Cryostem consortium was approved by IRB Sud-Mediterranée 1 (reference number

AC-2011-1420) and the Commission National Informatique et Liberté for data protection

(reference number nz70243374i n°912120). All patients gave their written consent for clinical

research and sample collection. This non-interventional research study with no additional

clinical procedure was carried out in accordance with the Declaration of Helsinki. Data

analyses were carried out using an anonymised database. Healthy donor's PBMC were

isolated from residual blood after apheresis provided by Etablissement Français du Sang

(18/EFS/032).

CITE Seq experiment

After thawing cells in a 50% fetal calf serum (FCS)/RPMI medium, 10x Genomics Feature

Barcoding recommendations were followed. Cells were washed twice using a 10 mL solution

of the same media. 500 000 cells were transferred in a 0.04% PBS/bovine serum albumin

(BSA) medium and washed. Cells were then suspended in a 1% PBS/BSA solution before

incubation with oligonucleotide-coupled antibodies TotalSeq-B (panel described in **Table S13**)

for 60 min on ice. Cells were washed 4 times (4°C, RCF 300, 5 minutes) and suspended in

order to obtain a concentration of 1 000 cells/µL. 10 000 cells were subjected in Gel Bead

Emulsion using Chromium 10x Genomics controller, according to manufacturer guidelines.

Chromium™ Next GEM Single Cell 3' Kit v3.1 (10X Genomics, cat. 1000268) and 3' Feature

Barcode Kit (10X Genomics, cat. 1000262) were used to prepare reagents.

To perform single cell RNA sequencing after cDNA amplification, the concentration of

each sample was measured using Tapestation 2200 (Agilent). To prepare the cDNA libraries

for 10x Genomics Chromium controller, we used the single-cell 3′ v3.1 kit (Dual Index) with

Feature Barcode technology for Cell Surface Protein, following manufacturer guidelines. QC

libraries were performed using Tapestation 2200 (Agilent). Libraries were equimolar pooled to

obtain at least 20 000 read pairs per cell and 5 000 read pairs for feature barcode part after

sequencing on Illumina Novaseq 6000 (100 cycles cartridge). The input number of cells was

estimated at 10 000 cells/samples. FASTQ files were obtained with bcl2fastq (Illumina).

Data quality control

Alignment, filtering, barcode counting, UMI counting were performed with count function from

10x Genomics Cell Ranger (version 6.0) using GRCh38 genome assembly as reference data.

Then aggr function was used to aggregate results without normalization. A R environment

(version 4.3.1) was subsequently used for the following analyses. Briefly, using Seurat R

package(*49*), dead cells were excluded if mitochondrial RNA was above 18%; singlet were

identified as droplets with RNA feature between 200 and 4 700; RNA and cell surface antigen

counts were normalized and scaled using Seurat R package normalisation and scaling function

for linear transformation. 2 000 highly variable RNA features were identified.

Immune cell clusterization

Using a principal component analysis method on the scaled data, cells were clustered

according to cell surface antigen expression. We used UMAP for cluster visualization. Seurat

R package was then used to calculate cell cycle phase based on canonical transcripts, and

regressing these out of the data.

Enrichment pathways analysis

Fast preranked gene set enrichment analysis was performed using FGSEA package(*50*) with

"hallmarks" gene reference data set from Molecular signature database(*51*). Pathways were

compared between GVHD and non-GVHD patients, and between SR-GVHD and SS-GVHD

patients. Gene expression comparison between GVHD and non GVHD patients, and between

SS- and SR-GVHD patients was provided by differential expression analysis. Enrichment

scores associated with adjusted p-value < 0.05 were selected for dot plot visualisation.

NicheNet Analysis

Ligand-receptors interaction was computed using NicheNet package(*33*). Sender cells were

identified as myeloid cells, B cells, CD4[+] conventional T cells and regulatory T cells. Receiver

cells were identified as CD4[+] and CD8[+] T lymphocytes, NK cells and regulatory T cells. Three

pooled human gene expression data of interacting cells(*52–54*) were used as input and

combined with a prior model that integrates existing knowledge on ligand-to-target signalling

paths to perform ligand activity analyses. Circos plot visualisation integrated *bona fide*

interactions of each receiver, and increased or decreased interactions were integrated from

"ligand differential expression heatmap" NicheNet output.

Trajectory inferences

Trajectory were based on RNA expression clusterisation (**Figure S10**). SingleCellExperiment

object was subset by immune populations (CD8+ T cells, CD4+ T cells, NK cells, B cells and

myeloid cells) before pseudotime calculation using slingshot package(*55*). TradSeq package

was then use for downstream analysis(*56*). Briefly, associationTest R function evidenced

genes whose expression is associated with pseudotime. Since pseudotime is a continuous

value, it was divided into 50 categories for graphical representation with heatmap. Only the

500 most expressed gene in SS- and SR-GVHD were selected for heatmap visualization

(**Table S3-S6**). Each point on X axis corresponded to a cell cluster with similar pseudotimes.

The immune populations represented on top of the heatmap corresponded to the immune cell

population most abundantly represented within the cell cluster (**Table S7-S10**). Gene markers

specific to low pseudotime or high pseudotime were found thanks to starVsEndTest command.

Expression along pseudotime in each condition was plotted for four genes from the top 100

starVsEndTest output **(Figure 5, Tables S11 and S12)**. To explore the pathways associated

with gene expression of the top 500 most expressed genes along pseudotime from each of

the 5 modules of the heatmap, we used over-representation analysis using enrichGO

command from clusterProfiler R package(*57*). Only pathways associated with adjusted

pval < 0.05 from "molecular functions" and "biological process" of the Gene Ontology

knowledgbase(*58*, *59*) were selected for dot plots visualization.

List of Supplementary Materials

REFERENCES

1. N. Cieri, N. Hookeri, K. Stromhaug, L. Li, J. Keating, P. Díaz-Fernández, V. Gómez-García de Soria, J. Stevens, R. Kfuri-Rubens, Y. Shao, K. A. Kooshesh, K. Powell, H. Ji, G. M. Hernandez, J. Abelin, S. Klaeger, C. Forman, K. R. Clauser, S. Sarkizova, D. A. Braun, L. Penter, H. T. Kim, W. J. Lane, G. Oliveira, L. S. Kean, S. Li, K. J. Livak, S. A. Carr, D. B. Keskin, C. Muñoz-Calleja, V. T. Ho, J. Ritz, R. J. Soiffer, D. Neuberg, C. Stewart, G. Getz, C. J. Wu, Systematic identification of minor histocompatibility antigens predicts outcomes of allogeneic hematopoietic cell transplantation. *Nat Biotechnol*, 1–12 (2024).

2. R. Zeiser, B. R. Blazar, Acute Graft-versus-Host Disease — Biologic Process, Prevention, and Therapy. *N Engl J Med* **377**, 2167–2179 (2017).

3. A. C. Harris, R. Young, S. Devine, W. J. Hogan, F. Ayuk, U. Bunworasate, C. Chanswangphuwana, Y. A. Efebera, E. Holler, M. Litzow, R. Ordemann, M. Qayed, A. S. Renteria, R. Reshef, M. Wölfl, Y.-B. Chen, S. Goldstein, M. Jagasia, F. Locatelli, S. Mielke, D. Porter, T. Schechter, Z. Shekhovtsova, J. L. M. Ferrara, J. E. Levine, International, Multicenter Standardization of Acute Graft-versus-Host Disease Clinical Data Collection: A Report from the Mount Sinai Acute GVHD International Consortium. *Biol Blood Marrow Transplant* **22**, 4–10 (2016).

4. L. Axt, A. Naumann, J. Toennies, S. P. Haen, W. Vogel, D. Schneidawind, S. Wirths, R. Moehle, C. Faul, L. Kanz, S. Axt, W. A. Bethge, Retrospective single center analysis of outcome, risk factors and therapy in steroid refractory graft-versus-host disease after allogeneic hematopoietic cell transplantation. *Bone Marrow Transplant* **54**, 1805–1814 (2019).

5. O. Penack, M. Marchetti, M. Aljurf, M. Arat, F. Bonifazi, R. F. Duarte, S. Giebel, H. Greinix, M. D. Hazenberg, N. Kröger, S. Mielke, M. Mohty, A. Nagler, J. Passweg, F. Patriarca, T. Ruutu, H. Schoemans, C. Solano, R. Vrhovac, D. Wolff, R. Zeiser, A. Sureda, Z. Peric, Prophylaxis and management of graft-versus-host disease after stem-cell transplantation

for haematological malignancies: updated consensus recommendations of the European Society for Blood and Marrow Transplantation. *Lancet Haematol* **11**, e147–e159 (2024).

6. P. J. Martin, J. D. Rizzo, J. R. Wingard, K. Ballen, P. T. Curtin, C. Cutler, M. R. Litzow, Y. Nieto, B. N. Savani, J. R. Schriber, P. J. Shaughnessy, D. A. Wall, P. A. Carpenter, First- and second-line systemic treatment of acute graft-versus-host disease: recommendations of the American Society of Blood and Marrow Transplantation. *Biol Blood Marrow Transplant* **18**, 1150–1163 (2012).

7. J. R. Westin, R. M. Saliba, M. De Lima, A. Alousi, C. Hosing, M. H. Qazilbash, I. F. Khouri, E. J. Shpall, P. Anderlini, G. Rondon, B. S. Andersson, R. Champlin, D. R. Couriel, Steroid-Refractory Acute GVHD: Predictors and Outcomes. *Adv Hematol* **2011**, 601953 (2011).

8. R. Zeiser, N. von Bubnoff, J. Butler, M. Mohty, D. Niederwieser, R. Or, J. Szer, E. M. Wagner, T. Zuckerman, B. Mahuzier, J. Xu, C. Wilke, K. K. Gandhi, G. Socié, REACH2 Trial Group, Ruxolitinib for Glucocorticoid-Refractory Acute Graft-versus-Host Disease. *N Engl J Med* **382**, 1800–1810 (2020).

9. P. J. Martin, G. Schoch, L. Fisher, V. Byers, C. Anasetti, F. R. Appelbaum, P. G. Beatty, K. Doney, G. B. McDonald, J. E. Sanders, A retrospective analysis of therapy for acute graft-versus-host disease: initial treatment. *Blood* **76**, 1464–1472 (1990).

10. R. Zeiser, N. von Bubnoff, J. Butler, M. Mohty, D. Niederwieser, R. Or, J. Szer, E. M. Wagner, T. Zuckerman, B. Mahuzier, J. Xu, C. Wilke, K. K. Gandhi, G. Socié, Ruxolitinib for Glucocorticoid-Refractory Acute Graft-versus-Host Disease. *N Engl J Med* **382**, 1800–1810 (2020).

11. J. L. M. Ferrara, A. C. Harris, J. K. Greenson, T. M. Braun, E. Holler, T. Teshima, J. E. Levine, S. W. J. Choi, E. Huber, K. Landfried, K. Akashi, M. Vander Lugt, P. Reddy, A. Chin, Q. Zhang, S. Hanash, S. Paczesny, Regenerating islet-derived 3-alpha is a biomarker of gastrointestinal graft-versus-host disease. *Blood* **118**, 6702–6708 (2011).

12. M. T. Vander Lugt, T. M. Braun, S. Hanash, J. Ritz, V. T. Ho, J. H. Antin, Q. Zhang, C.-H. Wong, H. Wang, A. Chin, A. Gomez, A. C. Harris, J. E. Levine, S. W. Choi, D. Couriel,

P. Reddy, J. L. M. Ferrara, S. Paczesny, ST2 as a marker for risk of therapy-resistant graft-versus-host disease and death. *N Engl J Med* **369**, 529–539 (2013).

13.   M. J. Hartwell, U. Özbek, E. Holler, A. S. Renteria, H. Major-Monfried, P. Reddy, M. Aziz, W. J. Hogan, F. Ayuk, Y. A. Efebera, E. O. Hexner, U. Bunworasate, M. Qayed, R. Ordemann, M. Wölfl, S. Mielke, A. Pawarode, Y.-B. Chen, S. Devine, A. C. Harris, M. Jagasia, C. L. Kitko, M. R. Litzow, N. Kröger, F. Locatelli, G. Morales, R. Nakamura, R. Reshef, W. Rösler, D. Weber, K. Wudhikarn, G. A. Yanik, J. E. Levine, J. L. Ferrara, An early-biomarker algorithm predicts lethal graft-versus-host disease and survival. *JCI Insight* **2**, e89798 (2017).

14.   H. Major-Monfried, A. S. Renteria, A. Pawarode, P. Reddy, F. Ayuk, E. Holler, Y. A. Efebera, W. J. Hogan, M. Wölfl, M. Qayed, E. O. Hexner, K. Wudhikarn, R. Ordemann, R. Young, J. Shah, M. J. Hartwell, M. S. Chaudhry, M. Aziz, A. Etra, G. A. Yanik, N. Kröger, D. Weber, Y.-B. Chen, R. Nakamura, W. Rösler, C. L. Kitko, A. C. Harris, M. Pulsipher, R. Reshef, S. Kowalyk, G. Morales, I. Torres, U. Özbek, J. L. M. Ferrara, J. E. Levine, MAGIC biomarkers predict long-term outcomes for steroid-resistant acute GVHD. *Blood* **131**, 2846–2855 (2018).

15.   G. Socié, D. Niederwieser, N. von Bubnoff, M. Mohty, J. Szer, R. Or, J. Garrett, A. Prahallad, C. Wilke, R. Zeiser, Prognostic value of blood biomarkers in steroid-refractory or steroid-dependent acute graft-versus-host disease: a REACH2 analysis. *Blood* **141**, 2771–2779 (2023).

16.   Q. Song, X. Wang, X. Wu, T. H. Kang, H. Qin, D. Zhao, R. R. Jenq, M. R. M. van den Brink, A. D. Riggs, P. J. Martin, Y.-Z. Chen, D. Zeng, IL-22-dependent dysbiosis and mononuclear phagocyte depletion contribute to steroid-resistant gut graft-versus-host disease in mice. *Nat Commun* **12**, 805 (2021).

17.   V. Arnhold, W. Y. Chang, S. A. Jansen, G. Thangavelu, M. Calafiore, P. Vinci, Y.-Y. Fu, T. Ito, S. Takashima, A. Egorova, J. Kuttiyara, A. Perlstein, M. van Hoesel, C. Liu, B. R.

Blazar, C. A. Lindemans, A. M. Hanash, Corticosteroids impair epithelial regeneration in

immune-mediated intestinal damage. *J Clin Invest* **134**, e155880 (2024).

18.   T. Toubai, C. Rossi, I. Tawara, C. Liu, C. Zajac, K. Oravecz-Wilson, D. Peltier, Y. Sun,

H. Fujiwara, S.-R. Wu, M. Riwes, I. Henig, S. Kim, P. Reddy, Murine Models of Steroid

Refractory Graft-versus-Host Disease. *Sci Rep* **8**, 12475 (2018).

19.   S. G. Holtan, A. Shabaneh, B. C. Betts, A. Rashidi, M. L. MacMillan, C. Ustun, K. Amin,

B. P. Vaughn, J. Howard, A. Khoruts, M. Arora, T. E. DeFor, D. Johnson, B. R. Blazar, D.

J. Weisdorf, J. Wang, Stress responses, M2 macrophages, and a distinct microbial

signature in fatal intestinal acute graft-versus-host disease. *JCI Insight* **5**, e129762, 129762

(2019).

20.   L. Schwab, L. Goroncy, S. Palaniyandi, S. Gautam, A. Triantafyllopoulou, A. Mocsai,

W. Reichardt, F. J. Karlsson, S. V. Radhakrishnan, K. Hanke, A. Schmitt-Graeff, M.

Freudenberg, F. D. von Loewenich, P. Wolf, F. Leonhardt, N. Baxan, D. Pfeifer, O. Schmah,

A. Schönle, S. F. Martin, R. Mertelsmann, J. Duyster, J. Finke, M. Prinz, P. Henneke, H.

Häcker, G. C. Hildebrandt, G. Häcker, R. Zeiser, Neutrophil granulocytes recruited upon

translocation of intestinal bacteria enhance graft-versus-host disease via tissue damage.

Nat Med **20**, 648–654 (2014).

21.   J. Hülsdünker, K. J. Ottmüller, H. P. Neeff, M. Koyama, Z. Gao, O. S. Thomas, M. Follo,

A. Al-Ahmad, G. Prinz, S. Duquesne, H. Dierbach, S. Kirschnek, T. Lämmermann, M. J.

Blaser, B. T. Fife, B. R. Blazar, A. Beilhack, G. R. Hill, G. Häcker, R. Zeiser, Neutrophils

provide cellular communication between ileum and mesenteric lymph nodes at graft-versus-

host disease onset. *Blood* **131**, 1858–1869 (2018).

22.   M. Pelletier, L. Maggi, A. Micheletti, E. Lazzeri, N. Tamassia, C. Costantini, L. Cosmi,

C. Lunardi, F. Annunziato, S. Romagnani, M. A. Cassatella, Evidence for a cross-talk

between human neutrophils and Th17 cells. *Blood* **115**, 335–343 (2010).

23.   D. Minns, K. J. Smith, V. Alessandrini, G. Hardisty, L. Melrose, L. Jackson-Jones, A. S.

MacDonald, D. J. Davidson, E. Gwyer Findlay, The neutrophil antimicrobial peptide

cathelicidin promotes Th17 differentiation. *Nat Commun* **12**, 1285 (2021).

24.   Q. Song, U. Nasri, D. Zeng, Steroid-Refractory Gut Graft-Versus-Host Disease: What

We Have Learned From Basic Immunology and Experimental Mouse Model. *Front Immunol*

13, 844271 (2022).

25.   B. K. Patel, M. J. Raabe, E. R. Lang, Y. Song, C. Lu, V. Deshpande, L. T. Nieman, M.

J. Aryee, Y.-B. Chen, D. T. Ting, Z. DeFilipp, Spatial transcriptomics reveals distinct tissue

niches linked with steroid responsiveness in acute gastrointestinal GVHD. *Blood* **142**, 1831–

1844 (2023).

26.   R. R. Jenq, Y. Taur, S. M. Devlin, D. M. Ponce, J. D. Goldberg, K. F. Ahr, E. R. Littmann,

L. Ling, A. C. Gobourne, L. C. Miller, M. D. Docampo, J. U. Peled, N. Arpaia, J. R. Cross,

T. K. Peets, M. A. Lumish, Y. Shono, J. A. Dudakov, H. Poeck, A. M. Hanash, J. N. Barker,

M.-A. Perales, S. A. Giralt, E. G. Pamer, M. R. M. van den Brink, Intestinal Blautia Is

Associated with Reduced Death from Graft-versus-Host Disease. *Biol Blood Marrow*

Transplant **21**, 1373–1383 (2015).

27.   Y. Furusawa, Y. Obata, S. Fukuda, T. A. Endo, G. Nakato, D. Takahashi, Y. Nakanishi,

C. Uetake, K. Kato, T. Kato, M. Takahashi, N. N. Fukuda, S. Murakami, E. Miyauchi, S.

Hino, K. Atarashi, S. Onawa, Y. Fujimura, T. Lockett, J. M. Clarke, D. L. Topping, M. Tomita,

S. Hori, O. Ohara, T. Morita, H. Koseki, J. Kikuchi, K. Honda, K. Hase, H. Ohno, Commensal

microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature*

504, 446–450 (2013).

28.   N. D. Mathewson, R. Jenq, A. V. Mathew, M. Koenigsknecht, A. Hanash, T. Toubai, K.

Oravecz-Wilson, S.-R. Wu, Y. Sun, C. Rossi, H. Fujiwara, J. Byun, Y. Shono, C. Lindemans,

M. Calafiore, T. M. Schmidt, K. Honda, V. B. Young, S. Pennathur, M. van den Brink, P.

Reddy, Gut microbiome-derived metabolites modulate intestinal epithelial cell damage and

mitigate graft-versus-host disease. *Nat Immunol* **17**, 505–513 (2016).

29.   J. L. Golob, M. M. DeMeules, T. Loeffelholz, Z. Z. Quinn, M. K. Dame, S. S. Silvestri,

M. C. Wu, T. M. Schmidt, T. L. Fiedler, M. J. Hoostal, M. Mielcarek, J. Spence, S. A. Pergam,

D. N. Fredricks, Butyrogenic bacteria after acute graft-versus-host disease (GVHD) are

associated with the development of steroid-refractory GVHD. *Blood Adv* **3**, 2866–2869

(2019).

30.   M. Couturier, B. Lamarthée, J. Arbez, J.-C. Renauld, C. Bossard, F. Malard, F.

Bonnefoy, M. Mohty, S. Perruche, P. Tiberghien, P. Saas, B. Gaugler, IL-22 deficiency in

donor T cells attenuates murine acute graft-versus-host disease mortality while sparing the

graft-versus-leukemia effect. *Leukemia* **27**, 1527–1537 (2013).

31.   B. Lamarthée, F. Malard, C. Gamonet, C. Bossard, M. Couturier, J.-C. Renauld, M.

Mohty, P. Saas, B. Gaugler, Donor interleukin-22 and host type I interferon signaling

pathway participate in intestinal graft-versus-host disease via STAT1 activation and

CXCL10. *Mucosal Immunol* **9**, 309–321 (2016).

32.   M. A. Schroeder, J. F. DiPersio, Mouse models of graft-versus-host disease: advances

and limitations. *Disease Models & Mechanisms* **4**, 318–333 (2011).

33.   R. Browaeys, W. Saelens, Y. Saeys, NicheNet: modeling intercellular communication

by linking ligands to target genes. *Nat Methods* **17**, 159–162 (2020).

34.   D. R. Edwards, M. M. Handsley, C. J. Pennington, The ADAM metalloproteinases. *Mol*

Aspects Med **29**, 258–289 (2008).

35.   J. C. Albring, M. M. Sandau, A. S. Rapaport, B. T. Edelson, A. Satpathy, M.

Mashayekhi, S. K. Lathrop, C.-S. Hsieh, M. Stelljes, M. Colonna, T. L. Murphy, K. M.

Murphy, Targeting of B and T lymphocyte associated (BTLA) prevents graft-versus-host

disease without global immunosuppression. *J Exp Med* **207**, 2551–2559 (2010).

36.   Y. Sakoda, J.-J. Park, Y. Zhao, A. Kuramasu, D. Geng, Y. Liu, E. Davila, K. Tamada,

Dichotomous regulation of GVHD through bidirectional functions of the BTLA-HVEM

pathway. *Blood* **117**, 2506–2514 (2011).

37.   F. Martins, E. Planet, D. Marino, M. Ansari, D. Trono, Single-cell transcriptome analysis

531   reveals atypical monocytes circulating ahead of acute graft-versus-host disease clinical

532   onset. *J Leukoc Biol*, qiae229 (2024).

533   38.   H. Sun, L. Wu, X. Zhao, Y. Huo, P. Dong, A. Pang, Y. Zheng, Y. Han, S. Ma, E. Jiang,

534   F. Dong, T. Cheng, S. Hao, Monocytes as an early risk factor for acute graft-versus-host

535   disease after allogeneic hematopoietic stem cell transplantation. *Front Immunol* **15**,

536   1433091 (2024).

537   39.   L. Jardine, U. Cytlak, M. Gunawan, G. Reynolds, K. Green, X.-N. Wang, S. Pagan, M.

538   Paramitha, C. A. Lamb, A. K. Long, E. Hurst, S. Nair, G. H. Jackson, A. Publicover, V.

539   Bigley, M. Haniffa, A. J. Simpson, M. Collin, Donor monocyte-derived macrophages

540   promote human acute graft-versus-host disease. *J Clin Invest* **130**, 4574–4586 (2020).

541   40.   K. Seno, M. Yasunaga, H. Kajiya, K. Izaki-Hagio, H. Morita, M. Yoneda, T. Hirofuji, J.

542   Ohno, Dynamics of M1 macrophages in oral mucosal lesions during the development of

543   acute graft-versus-host disease in rats. *Clin Exp Immunol* **190**, 315–327 (2017).

544   41.   J. E. Levine, S. Paczesny, S. Mineishi, T. Braun, S. W. Choi, R. J. Hutchinson, D.

545   Jones, Y. Khaled, C. L. Kitko, D. Bickley, O. Krijanovski, P. Reddy, G. Yanik, J. L. M.

546   Ferrara, Etanercept plus methylprednisolone as initial therapy for acute graft-versus-host

547   disease. *Blood* **111**, 2470–2475 (2008).

548   42.   R. J. Robb, G. R. Hill, The interferon-dependent orchestration of innate and adaptive

549   immunity after transplantation. *Blood* **119**, 5351–5358 (2012).

550   43.   S. Wang, T. Cheng, X. Chen, C. Zeng, W. Qin, Y. Xu, IFN-γ induces acute graft-versus-

551   host disease by promoting HMGB1-mediated nuclear-to-cytoplasm translocation and

552   autophagic degradation of p53. *Clin Sci (Lond)* **138**, 1287–1304 (2024).

553   44.   H. Okamura, H. Tsutsi, T. Komatsu, M. Yutsudo, A. Hakura, T. Tanimoto, K. Torigoe,

554   T. Okura, Y. Nukada, K. Hattori, Cloning of a new cytokine that induces IFN-gamma

555   production by T cells. *Nature* **378**, 88–91 (1995).

45. Y. Fujimori, H. Takatsuka, Y. Takemoto, H. Hara, H. Okamura, K. Nakanishi, E.

Kakishita, Elevated interleukin (IL)-18 levels during acute graft-versus-host disease after

allogeneic bone marrow transplantation. *Br J Haematol* **109**, 652–657 (2000).

46. S. W. Choi, G. C. Hildebrandt, K. M. Olkiewicz, D. A. Hanauer, M. N. Chaudhary, I. A.

Silva, C. E. Rogers, D. T. Deurloo, J. M. Fisher, C. Liu, D. Adams, S. W. Chensue, K. R.

Cooke, CCR1/CCL5 (RANTES) receptor-ligand interactions modulate allogeneic T-cell

responses and graft-versus-host disease following stem-cell transplantation. *Blood* **110**,

3447–3455 (2007).

47. S. DeWolf, Y. Elhanati, K. Nichols, N. R. Waters, C. L. Nguyen, J. B. Slingerland, N.

Rodriguez, O. Lyudovyk, P. A. Giardina, A. I. Kousa, H. Andrlová, N. Ceglia, T. Fei, R.

Kappagantula, Y. Li, N. Aleynick, P. Baez, R. Murali, A. Hayashi, N. Lee, B. Gipson, M.

Rangesa, Z. Katsamakis, A. Dai, A. G. Blouin, M. Arcila, I. Masilionis, R. Chaligne, D. M.

Ponce, H. J. Landau, I. Politikos, R. Tamari, A. M. Hanash, R. R. Jenq, S. A. Giralt, K. A.

Markey, Y. Zhang, M.-A. Perales, N. D. Socci, B. D. Greenbaum, C. A. Iacobuzio-Donahue,

T. J. Hollmann, M. R. M. van den Brink, J. U. Peled, Tissue-specific features of the T cell

repertoire after allogeneic hematopoietic cell transplantation in human and mouse. *Sci*

Transl Med **15**, eabq0476 (2023).

48. S. D. Schutt, Y. Wu, A. Kharel, D. Bastian, H.-J. Choi, M. Hanief Sofi, C. Mealer, B.

McDaniel Mims, H. Nguyen, C. Liu, K. Helke, W. Cui, X. Zhang, Y. Ben-David, X.-Z. Yu,

The druggable transcription factor Fli-1 regulates T cell immunity and tolerance in graft-

versus-host disease. *J Clin Invest* **132**, e143950 (2022).

49. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A.

J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain,

A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish,

R. Gottardo, P. Smibert, R. Satija, Integrated analysis of multimodal single-cell data. *Cell*

184, 3573-3587.e29 (2021).

50.   G. Korotkevich, V. Sukhov, N. Budin, B. Shpak, M. N. Artyomov, A. Sergushichev, Fast

gene set enrichment analysis. bioRxiv [Preprint] (2021). https://doi.org/10.1101/060012.

51.   A. Liberzon, C. Birger, H. Thorvaldsdóttir, M. Ghandi, J. P. Mesirov, P. Tamayo, The

Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417–

425 (2015).

52.   J. A. Ramilowski, T. Goldberg, J. Harshbarger, E. Kloppmann, M. Lizio, V. P.

Satagopam, M. Itoh, H. Kawaji, P. Carninci, B. Rost, A. R. R. Forrest, A draft network of

ligand-receptor-mediated multicellular signalling in human. *Nat Commun* **6**, 7866 (2015).

53.   A. J. Pawson, J. L. Sharman, H. E. Benson, E. Faccenda, S. P. H. Alexander, O. P.

Buneman, A. P. Davenport, J. C. McGrath, J. A. Peters, C. Southan, M. Spedding, W. Yu,

A. J. Harmar, NC-IUPHAR, The IUPHAR/BPS Guide to PHARMACOLOGY: an expert-

driven knowledgebase of drug targets and their ligands. *Nucleic Acids Res* **42**, D1098-1106

(2014).

54.   A. D. Rouillard, G. W. Gundersen, N. F. Fernandez, Z. Wang, C. D. Monteiro, M. G.

McDermott, A. Ma'ayan, The harmonizome: a collection of processed datasets gathered to

serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016**, baw100

(2016).

55.   K. Street, D. Risso, R. B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, S. Dudoit,

Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC*

Genomics **19**, 477 (2018).

56.   K. Van den Berge, H. Roux de Bézieux, K. Street, W. Saelens, R. Cannoodt, Y. Saeys,

S. Dudoit, L. Clement, Trajectory-based differential expression analysis for single-cell

sequencing data. *Nat Commun* **11**, 1201 (2020).

57.   G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

58.   M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis,

K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis,

S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, G. Sherlock, Gene

Ontology: tool for the unification of biology. *Nat Genet* **25**, 25–29 (2000).

59.   The Gene Ontology Consortium, S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry,

H. J. Drabkin, D. Ebert, M. Feuermann, P. Gaudet, N. L. Harris, D. P. Hill, R. Lee, H. Mi, S.

Moxon, C. J. Mungall, A. Muruganugan, T. Mushayahama, P. W. Sternberg, P. D. Thomas,

K. Van Auken, J. Ramsey, D. A. Siegele, R. L. Chisholm, P. Fey, M. C. Aspromonte, M. V.

Nugnes, F. Quaglia, S. Tosatto, M. Giglio, S. Nadendla, G. Antonazzo, H. Attrill, G. dos

Santos, S. Marygold, V. Strelets, C. J. Tabone, J. Thurmond, P. Zhou, S. H. Ahmed, P.

Asanitthong, D. Luna Buitrago, M. N. Erdol, M. C. Gage, M. Ali Kadhum, K. Y. C. Li, M.

Long, A. Michalak, A. Pesala, A. Pritazahra, S. C. C. Saverimuttu, R. Su, K. E. Thurlow, R.

C. Lovering, C. Logie, S. Oliferenko, J. Blake, K. Christie, L. Corbani, M. E. Dolan, H. J.

Drabkin, D. P. Hill, L. Ni, D. Sitnikov, C. Smith, A. Cuzick, J. Seager, L. Cooper, J. Elser, P.

Jaiswal, P. Gupta, P. Jaiswal, S. Naithani, M. Lera-Ramirez, K. Rutherford, V. Wood, J. L.

De Pons, M. R. Dwinell, G. T. Hayman, M. L. Kaldunski, A. E. Kwitek, S. J. F. Laulederkind,

M. A. Tutaj, M. Vedi, S.-J. Wang, P. D'Eustachio, L. Aimo, K. Axelsen, A. Bridge, N. Hyka-

Nouspikel, A. Morgat, S. A. Aleksander, J. M. Cherry, S. R. Engel, K. Karra, S. R. Miyasato,

R. S. Nash, M. S. Skrzypek, S. Weng, E. D. Wong, E. Bakker, T. Z. Berardini, L. Reiser, A.

Auchincloss, K. Axelsen, G. Argoud-Puy, M.-C. Blatter, E. Boutet, L. Breuza, A. Bridge, C.

Casals-Casas, E. Coudert, A. Estreicher, M. Livia Famiglietti, M. Feuermann, A. Gos, N.

Gruaz-Gumowski, C. Hulo, N. Hyka-Nouspikel, F. Jungo, P. Le Mercier, D. Lieberherr, P.

Masson, A. Morgat, I. Pedruzzi, L. Pourcel, S. Poux, C. Rivoire, S. Sundaram, A. Bateman,

E. Bowler-Barnett, H. Bye-A-Jee, P. Denny, A. Ignatchenko, R. Ishtiaq, A. Lock, Y. Lussi,

M. Magrane, M. J. Martin, S. Orchard, P. Raposo, E. Speretta, N. Tyagi, K. Warner, R. Zaru,

A. D. Diehl, R. Lee, J. Chan, S. Diamantakis, D. Raciti, M. Zarowiecki, M. Fisher, C. James-

Zorn, V. Ponferrada, A. Zorn, S. Ramachandran, L. Ruzicka, M. Westerfield, The Gene

Ontology knowledgebase in 2023. *Genetics* **224**, iyad031 (2023).

Acknowledgments:

The authors thank the patients who agreed to participate in this study by providing their blood samples. We also thank the iGenSeq platform at ICM for sequencing the PBMC samples. We are grateful to the CRYOSTEM biobank for collecting patient samples and clinical data.

Funding:

This project was funded by la Fédération Leucémie Espoir and Fondation Maladies Rare (HPN-AM).

Author contributions:

Conceptualization: SLG, DM

Methodology: SLG, NV, DM

Samples processing: SLG, DB, YM, EB

Investigation: SLG, NV, DM

Visualization: SLG, GS, RPD, NV, DM

Funding acquisition: DM

Project administration: EB, DM, GS, RPD

Supervision: NV, DM

Writing – original draft: SLG, NV, DM

Draft revision: SLG, YM, DB, EB, RPF, GS, NV, DM

Competing interests:

Authors declare that they have no competing interests.

Data availability

Raw data are available on GEO public repository: GSE229733. A private reviewer access has been provided with the following secured token: ynmbkwaknfebdwb

Source code to reproduce the analyses, figures, and tables described in this manuscript are provided in a Git repository: https://gitlab.com/SophieLG/gvh_transcriptomic and long term preserved in Software Heritage:

662   https://archive.softwareheritage.org/browse/origin/directory/?origin_url=https://gitlab.com/Sop

663   hieLG/gvh_transcriptomic×tamp=2024-12-09T14:03:56.753000%2B00:00

 Figures legends:

Figure 1: Patients and cells characterization.

A: Experimental design: HSCT patients (n=53) with (n=21) and without aGVHD (n=32) and

healthy donor as control group (n=6) were analyzed. PBMC were collected at day 100 post

HSCT for patients without aGVHD, and before steroid initiation for patients with aGVHD.

B: Number of cells included in the analysis after CITE-Seq processing. **C:** Uniform Manifold

Approximation and Projection (UMAP) after principal component analysis for immune cells

clusterisation, according to surface markers. **D:** Heatmap depicting immune cluster's

identification through surface antigen expression. Three irrelevant populations were removed

and are shown in **Figure S1**. **E**: Volcano plot exhibiting immune cells abundance in GVHD or

non-GVHD condition. Y axis reflects que -log 10(adjusted p-value) from differential expression

output, using non-parametric Mann-Whitney-Wilcoxon rank sum test. X axis reflects the

log(ratio) of median abundance in aGVHD condition, over non-aGVHD condition. Abbreviation:

CITE-Seq: Cellular indexing of transcriptomes and epitopes sequencing. aGVHD: acute graft

versus host disease. HSCT: allogeneic hematopoietic stem cell transplantation. PBMC:

Peripheral Blood Mononuclear Cells. SS-aGVHD: steroid sensitive acute graft versus host

disease. SR-aGVHD: steroid resistant acute graft versus host disease.

Figure 2: Immunotypes distinguished different immune patterns associated with

aGVHD incidence.

A: Principal coordinates analysis (PCoA) revealed 4 different immune clusters (immunotypes)

according to cell population composition. **B:** Distribution of aGVHD patients among

immunotypes (left panel) and corticosteroid sensitivity (right panel). Fisher test was used for

statistical analysis. **C:** GVHD cumulative incidence with death as competing risk, estimated by

Kaplan-Meir method, Logrank test was used for statistical analysis. Immunotype 4 cumulative

incidence at day 100 = 72%, Immunotype 3 cumulative incidence at day 100 = 0%,

Immunotype 2 cumulative incidence at day 100 = 38%, Immunotype 1 cumulative incidence at

day 100 = 55%, log-rank p-value = 0.005. **D:** Bar plots showing top 15 contributing immune

subsets to immunotypes. Contribution was determined with permutational multivariate analysis

of variance (PERMANOVA).

Figure 3: Cell cycle scoring and enrichment pathways

A: Volcano plots exhibiting cell cycle analysis after cell cycle phases scoring. X axis represents

log ratio of median number of cells in each phase in aGVHD patients over non GVHD patients.

Y axis represents log adjusted p values. Immune population significantly increased or

decreased are marked in red. Mann-Whitney-Wilcoxon tests were performed for statistical

analysis, and p values were adjusted using Benjamini and Hochberg method. **B:** Pathways

enrichment analysis comparing SR and SS-aGVHD, using gene set enrichment method and

"hallmarks" gene reference data set. Only significant values with adjusted p-value < 0.05 are

shown. NES: normalized enrichment score. SS-aGVHD: steroid sensitive acute graft versus

host disease. SR-aGVHD: steroid resistant acute graft versus host disease.

Figure 4: Cell-cell interactions with CD8+ T cells as receiver revealed specific patterns

associated with SR-aGVHD

NicheNet Output when focusing on CD8+ T cells as receiver is depicted on panel A, B and C.

A: Expression of top-ranked ligands on sender cells. Prioritized ligands identified are exhibited

on Y axis, and sender cells on X axis. The color determines the average expression and the

point size the percent of senders expressing the ligand. **B:** Relative expression of ligands on

sender cells. Expression is exhibited in log fold change (red: increase expression in

SR-aGVHD patients, Blue: decreased expression in SR-aGVHD patients). **C:** Ligand–target

matrix exhibiting the regulatory potential of interactions. On the X axis is shown the active

targets genes of the top-ranked ligands. Heatmap visualization showing which top-ranked

ligands are predicted to have regulated the expression of which differentially expressed genes.

D: Circos plots depicting interactions between senders and all receiver cells. Ligands are

shown on the left side of the circos plot, and targets on the right side. Only *bona fide*

interactions were selected (**Figure S8**). Links exhibits positive log fold change (left panel,

increased in SR-aGVHD patients) or negative log fold change (right panel, increased in

SS-aGVHD patients) from "ligand differential expression heatmap" NicheNet's output. The

thickness of the link represents the numbers of sender populations involved in the interaction.

SR-aGVHD: Steroid resistant acute graft versus host disease. SS-aGVHD: Steroid sensitive

acute graft versus host disease.

Figure 5: SR-aGVHD is associated with specific trajectory in CD8$^+$ T cells

A: Density plot for number of cells according to lineage 1 pseudotime visualization. SR-aGVHD

cell abundance was marked in low and high pseudotimes, while SS-aGVHD was associated

with a continuous increase in cell abundance along differentiation. **B:** Lineage 1 trajectory on

RNA clusterization. **C:** Heatmap depicting gene expression among the top 500 most expressed

genes in SS-aGVHD patient (top panel) and in SR-aGVHD patients (low panel), along

pseudotime. The color at the top of each heatmap represents the immune population in which

the gene is expressed. **D:** Expression along pseudotime of four of the top 100 genes, whose

expression varied the most between the start and end points of a lineage.

E: Over-representation analysis to identify pathways associated with genes expression from

each module of the heatmaps. Only pathways associated with adjusted pval < 0.05 from

"molecular functions" and "biological process" of the Gene Ontology knowledgbase(*58*, *59*)

were selected for dot plots visualisation. SR-aGVHD: Steroid resistant acute graft versus host

disease. SS-aGVHD: Steroid sensitive acute graft versus host disease.

Fig. S1: Immune populations in each group of patients:

HSCT: hematopoietic Stem Cell Transplantation. GVHD: Graft Vetsus Host Disease.

Fig. S2: Immunotype survival analysis:

Survival analyses according to immunotypes

Fig. S3: Immunotype and clinical variables:

Clinical parameters and immunotypes.

Fig. S4: Immune cell abundance in SS- and SR-GVHD patients:

Volcano plot exhibiting immune cells abundance in SR-GVHD and SS-GVHD condition. Y axis reflects que -log 10(adjusted p-value) from differential expression output, using non-parametric Wilcoxon rank sum test. X axis reflects the log(ratio) of median abundance in SR-GVHD condition, over SS-GVHD condition. Abbreviation: GVHD: graft versus host disease. HSCT: allogeneic hematopoietic stem cell transplantation. SS-GVHD: Steroid Sensitive acute graft versus host disease. SR-GVHD: Steroid resistant acute graft versus host disease.

Fig. S5: Cell cycle in SS- and SR-GVHD patients and enrichment pathways in aGVHD:

Volcano plots exhibiting cell cycle analysis after cell cycle phases scoring. X axis represents log ratio of median number of cells in each phase in SR-GVHD patients over SS-GVHD patients. Y axis represents log adjusted p values. Immune population significantly increased or decreased are marked in red. Mann-Whitney-Wilcoxon tests were performed for statistical analysis, and p values were adjusted using Benjamini and Hochberg method.

Fig. S6: Pathway enrichments in aGVHD patients:

Pathways enrichment analysis comparing aGVHD and no GVHD, using gene set enrichment method and "hallmarks" gene reference data set. Only significant values with adjusted

p-value < 0.05 are shown. NES: normalized enrichment score. SS-GVHD: Steroid Sensitive

acute graft versus host disease. SR-GVHD: Steroid resistant acute graft versus host disease.

Fig. S7: Interactions combined plots in regulatory T cells and NK cells:

A: Combined plots of cell-cell interactions analysis, with T regulators lymphocytes as receiver.

B: Combined plots of cell-cell interactions analysis, with NK cells as receiver.

Fig. S8: Interactions combined plots in CD4$^+$T cells and double negative T cells

A: Combined plots of cell-cell interactions analysis, with CD4$^+$ T lymphocytes as receiver.

B: Combined plots of cell-cell interactions analysis, with double negative T cells as receiver.

Fig. S9: Bona fide interactions in all receivers:

A: Known interactions (*bona fide*) with prioritized ligands of each receiver populations.

B: Increased and decreased interactions in patients with aGVHD, according to *bona fide*

interactions.

Fig. S10: Clustering based on RNA expression

A: Clusters based on mRNA expression. Immune cells were identified using protein surface

expression after clustering. **B:** Immune cells on UMAP in GVHD patients, no GVHD patients

and healthy donors.

Fig. S11: CD8$^+$ T cells lineage 2 trajectory analysis:

A: Density plot for number of cells according to lineage 2 pseudotime visualization. SR-GVHD

cell abundance was marked in low and high pseudotimes, while SS-GVHD was associated

with a continuous increase in cell abundance along differentiation. **B:** Lineage 2 trajectory on

RNA clusterisation. **C:** Heatmap of RNA expression of the 500 most expressed gene in

SS-GVHD patient (top panel) and in SR-GVHD patients (low panel), along pseudotime. The

color at the top of each heatmap represents the immune population in which the gene is

expressed. **D:** Expression along pseudotime of four of the top 100 genes, whose expression

varied the most between the start and end points of a lineage. **E:** Over-representation analysis

to identify pathways associated with genes expression from each module of the heatmaps.

Only pathways associated with adjusted pval < 0.05 from "molecular functions" and "biological

process" of the Gene Ontology knowledgbase(*58*, *59*) were selected for dot plots visualisation.

SR–GVHD: Steroid resistant graft versus host disease. SS–GVHD: Steroid sensitive graft

versus host disease.

Fig. S12: Over representation analysis in lineage 1 SR-GVHD patients:

Over-representation analysis in lineage 1 and SR-GVHD patients to identify pathways

associated with genes expression from each module of the heatmaps. Only pathways

associated with adjusted pval < 0.05 from "molecular functions" and "biological process" of the

Gene Ontology knowledgbase(*58*, *59*) were selected for dot plots visualisation. SR–

GVHD: Steroid resistant graft versus host disease. SS–GVHD: Steroid sensitive graft versus

host disease.

Fig. S13: Over representation analysis in lineage 1 SS-GVHD patients:

Over-representation analysis in lineage 1 and SS-GVHD patients to identify pathways

associated with genes expression from each module of the heatmaps. Only pathways

associated with adjusted pval < 0.05 from "molecular functions" and "biological process" of the

Gene Ontology knowledgbase(*58*, *59*) were selected for dot plots visualisation. SR–

GVHD: Steroid resistant graft versus host disease. SS–GVHD: Steroid sensitive graft versus

host disease.

Fig. S14: Over representation analysis in lineage 2 SR-GVHD patients:

Over-representation analysis in lineage 2 and SR-GVHD patients to identify pathways

associated with genes expression from each module of the heatmaps. Only pathways

associated with adjusted pval < 0.05 from "molecular functions" and "biological process" of the

Gene Ontology knowledgbase(*58*, *59*) were selected for dot plots visualisation. SR–

GVHD: Steroid resistant graft versus host disease. SS–GVHD: Steroid sensitive graft versus

host disease.

Fig. S15: Over representation analysis in lineage 2 SS-GVHD patients:

Over-representation analysis in lineage 2 and SS-GVHD patients to identify pathways

associated with genes expression from each module of the heatmaps. Only pathways

associated with adjusted pval < 0.05 from "molecular functions" and "biological process" of the

Gene Ontology knowledgbase(*58*, *59*) were selected for dot plots visualisation. SR–

GVHD: Steroid resistant graft versus host disease. SS–GVHD: Steroid sensitive graft versus

host disease.

Table S1: Patients' characteristics

[1]Wilcoxon rank sum exact test; Pearson's Chi-squared test; Fisher's exact test; Wilcoxon rank

sum test

ALL: Acute Lymphoblastic leukemia, AML: Acute Myeloblastic Leukemia, CML: Chronic

Myeloid Leukemia, CR: Complete Remission, CMV: Cytomegalovirus, GVHD: graft versus

host disease, HSCT : Hematopoietic Stem Cell Transplantation, MDS: Myedlodysplastic

syndrome, NA: Not applicable, NHL: Non Hodgkin Lymphoma

Table S2: Characteristics according to immunotypes:

[1]n (%); Median(Range)

[2]Fisher's exact test; Kruskal-Wallis rank sum test

ALL: Acute Lymphoblastic leukemia, AML: Acute Myeloblastic Leukemia, CML: Chronic

Myeloid Leukemia, CR: Complete Remission, CMV: Cytomegalovirus, GVHD: graft versus

host disease, HSCT : Hematopoietic Stem Cell Transplantation, MDS: Myedlodysplastic

syndrome, NA: Not applicable, NHL: Non Hodgkin Lymphoma

Table S3: Top 500 genes expression matrix in lineage 1 SR-GVHD patients

Table S4: Top 500 genes expression matrix in lineage 1 SS-GVHD patients

Table S5: Top 500 genes expression matrix in lineage 2 SR-GVHD patients

Table S6: Top 500 genes expression matrix in lineage 2 SS-GVHD patients

Table S7: Immune population mostly represented along pseudotimes in lineage 1 SR-
GVHD patients

Table S8: Immune population mostly represented along pseudotimes in lineage 1 SS-
GVHD patients

Table S9: Immune population mostly represented along pseudotimes in lineage 1
2 SR-GVHD patients

Table S10: Immune population mostly represented along pseudotimes in lineage 2 SS-
GVHD patients

Table S11: StartvsEnd output in lineage 1

Table S12: StartvsEnd output in lineage 2

Table S13: Total-Seq-B antibody panel

Figure 1

A allo-HSCT n=53 — Diagnosis of acute GVHD n=21 — D+100 Patients without acute GVHD n=32 — Steroid regimen — Steroid-sensitive (SS) n=16 — Steroid-refractory (SR) n=5

B CITE-seq → Cluster on cell surface antigens → Cell subsets analyses, Transcriptomic analyses, Trajectory inferences, Cell-to-cell interactions

n cells: 73 917, 95 236, 131 331, 27 290, 64 946

- Healthy controls
- Acute GVHD before steroid regimen
- No acute GVHD
- Steroid-refractory
- Steroid-sensitive

C T cells, NK cells, Monocytes, B cells; UMAP1, UMAP2

D Scaled expression 0 / 0.5 / 1

Clusters (left labels):
- 26 - Non switched mem B cells
- 9 - Naive B cells
- 29 - MAIT cells
- 40 - ILC2
- 4 - CD4+ T cells th22 CTLA4low PD1low
- 27 - Regulatory T cells
- 13 - EM CD4+ T cells ICOS+ CTLA4+ PD1+
- 12 - Naive CD8+ T cells
- 8 - Naive CD4+ T cells
- 23 - CD8+ T cells TC2 activated CTLA4low
- 30 - SCM CD8+ T cells TCRva72+
- 24 - Double negative th17 T cells CCR6+
- 19 - CM CD4+ T cells TH1 CTLA4bright
- 21 - EM CD8+ T cells CTLA4bright
- 16 - Stem cell memory CD8+ T cells
- 1 - EMRA CD8+ CTLA4+ 2B4+ T cells
- 11 - EM CD8+ CTLA4+ 2B4+ T cells
- 37 - Plasmocytes
- 25 - Plasmacytoïd dendritic cells
- 10 - Classical monocytes CD73+
- 15 - NK cells CD56bright CD16-
- 7 - NK cells CD56dim NKp30bright NKp46+
- 18 - NK cells CD56- NKp46low NKp30+
- 36 - NK cells CD56dim NKp46+ NKp30+
- 6 - Non classical monocytes
- 33 - Myeloïds macrophages
- 22 - Class. monocytes CD73+ CD39+
- 31 - Classical monocytes CD39low
- 35 - Classical monocytes IgM+ CD11c+
- 2 - Classical monocytes KLRG1+
- 0 - Classical monocytes CD11bbright
- 5 - Classical monocytes NKp46+
- 32 - Intermediate monocytes
- 28 - Dendritic cells
- 39 - Classical monocytes
- 20 - Class. monocytes CD11blow CD11clow
- 3 - Classical monocytes IgM+ CD11bbright
- 34 - Mature LB1

Markers (bottom): CTLA4, KLRG1, CCR7, CCR6, CD73, CXCR3, NKG2D, 2B4, NKp30, PD1, CD95, CCR5, CD39, CD11b, CD14, HLA-DR, CD11, CD127, CD27, CD3, ICOS, CD28, CD4, CD45RA, CD8, CDp46, CD56, CD16, CD161, TCRva72, CD25, CCR4, CD38, KIR2DL1, CD123, CXCR5, CD24, CD20, CD19, IgM, IgD

% cell in HD

E −log(adjusted p value)

- SCM CD8+ T cells TCRva72+
- EM CD8+ T cells CTLA4bright
- CM CD4+ T cells TH1 CTLA4bright
- Classical monocytes CD73+
- Naive B cells
- NK cells CD56- NKp46low NKp30+
- NK cells CD56dim NKp46+ NKp30+
- Non switched mem B cells
- NK cells CD56dim NKp30bright NKp46+

Decreased in aGVHD — Increased in aGVHD

- Adj p-value < 0.05
- Adj p-value > 0.05

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Patients with aGVHD

Patients without aGVHD

Healthy donors

- 0 - Classical monocytes CD11b^{bright}
- 1 - EMRA CD8$^+$ CTLA4$^+$ 2B4$^+$ T cells
- 2 - Classical monocytes KLRG1$^+$
- 3 - Classical monocytes IgM$^+$ CD11b^{bright}
- 4 - CD4$^+$ T cells th22 CTLA4^{low} PD1^{low}
- 5 - Classical monocytes NKp46$^+$
- 6 - Non classical monocytes
- 7 - NK cells CD56^{dim} NKp30^{bright} NKp46$^+$
- 8 - Naive CD4$^+$ T cells
- 9 - Naive B cells
- 10 - Classical monocytes CD73$^+$
- 11 - EM CD8$^+$ CTLA4$^+$ 2B4$^+$ T cells
- 12 - Naive CD8$^+$ T cells
- 13 - EM CD4$^+$ T cells ICOS$^+$ CTLA4$^+$ PD1$^+$
- 14 - Irrelevant population
- 15 - NK cells CD56^{bright} CD16$^-$
- 16 - Stem cell memory CD8$^+$ T cells
- 17 - Irrelevant population
- 18 - NK cells CD56$^-$ NKp46^{low} NKp30$^+$
- 19 - CM CD4$^+$ T cells TH1 CTLA4^{bright}
- 20 - Class. monocytes CD11b^{low} CD11c^{low}

- 21 - EM CD8$^+$ T cells CTLA4^{bright}
- 22 - Class. monocytes CD73$^+$ CD39$^+$
- 23 - CD8$^+$ T cells TC2 activated CTLA4^{low}
- 24 - Double negative th17 T cells CCR6$^+$
- 25 - Plasmacytoïd dendritic cells
- 26 - Non switched mem B cells
- 27 - Regulatory T cells
- 28 - Dendritic cells
- 29 - MAIT cells
- 30 - SCM CD8$^+$ T cells TCRva72$^+$
- 31 - Classical monocytes CD39^{low}
- 32 - Intermediate monocytes
- 33 - Myeloïds macrophages
- 34 - Mature LB1
- 35 - Classical monocytes IgM$^+$ CD11c$^+$
- 36 - NK cells CD56^{dim} NKp46$^+$ NKp30$^+$
- 37 - Plasmocytes
- 17 - Irrelevant population
- 39 - Classical monocytes
- 40 - ILC2

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

A

T regulators lymphocytesas receivers

B

NK cells as receivers

Figure S8

CD4+ T cells as receiver

Double negative T cells as receiver

Figure S9

A

CD4⁺ T cells as receiver

CD8⁺ T cells as receiver

T regulators lymphocytes as receiver

Double negative T cells as receiver

NK cells as receiver

B

Enriched interactions in aGVHD

Enriched interactions in absence of aGVHD

Figure S10

A

Row labels (top to bottom):
- 4. CM CD4+ T cells th22
- 1. EM CD4+ T cells th9
- 27. Naive double+ T cells
- 3. Regulatory T cells ICOS+
- 34. Regulatory T cells CCR5+
- 25. CD8+ TEMRA cells ICOS^{bright}PD1+
- 8. Unconventional T cells
- 33. CD8+ TEMRA CTLA4+
- 15. CM CD8+ T cells exhausted
- 9. CD8+ TEMRA 2B4+
- 10. CD8+ TEMRA 2B4+ CTLA4+
- 2. EM CD8+ T cells ICOS+
- 14. Non switched B cells
- 21. Activated B cells CD73-
- 36. Activated B cells CXCR5-
- 11. NK cells CD56^{dim} CD16^{low}
- 22. NK cells CD56^{dim} CD16^{bright}
- 17. NK cells CD56^{bright} CD16^{dim}
- 40. Plasmocytes
- 31. Dendritic cells
- Ireelevant population
- Irrelevant population
- 23. Clas. monocytes CD73+ CCR6+
- Irrelevant population
- 32. Class. monocytes CD73+CD11b^{bright}
- Irrelevant population
- Irrelevant population
- 6. Int. monocytes CD11c^{bright}
- 29. Int. monocytes 2B4+
- 16. Clas. monocytes KLRG1+
- 13. Clas. monocytes CD39^{high} CTLA4+
- 0. Clas. monocytes PD1^{high} KLRG1^{low}
- 30. Clas. monocytes PD1^{low}
- 5. Clas. monocytes CD25+
- 35. Clas. monocytes NKG2D+
- 24. Clas. monocytes CD11b^{bright}
- 7. Clas. monocytes PD1^{bright}
- 18. Clas. monocytes CCR6lowKLRG1^{low}
- 12. Clas. monocytes CCR4+
- 19. Clas. monocytes CCR6-
- 38. Transitional B cells
- 26. Clas. monocytes PD1+IgM+

Column labels: KLRG1, CCR7, CTLA4, CCR6, CD95, PD1, 2B4, NKG2D, CCR5, CD39, CD14, CD11b, HLA-DR, CD11c, CD161, TCRVa72, CD3, CD8, CD25, ICOS, CCR4, CD27, CD4, CD28, CD127, CD45RA, CXCR3, CD73, IgM, CD24, CD20, IgD, CD19, CXCR5, KID2DL1, CD16, NKp30, NKp46, CD56, CD38, CD123

%cell in HD

B

GVHD+ patients | GVHD- patients | Healthy donors

Legend: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41

UMAP_1

UMAP_2

Figure S11

Figure S12

Lineage 1 and SR-GVHD patients

Figure S13

Lineage 1 and SS-GVHD patients

Figure S14

Lineage 2 and SR-GVHD patients

Figure S15

Lineage 2 and SS GVHD patients