Cis-regulatory chromatin contacts form de novo in the absence of loop extrusion

Nicholas G. Aboreden[1,2]*, Han Zhao[3]*, Fengnian Shan[3,4], Fuhai Liu[3], Haoyue Zhang[3]**,
Gerd A. Blobel[1,2]**

[1]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[2]Division of Hematology, The Children's Hospital of Philadelphia, Philadelphia, PA, USA

[3]Institute of Molecular Physiology, Shenzhen Bay Laboratory, Shenzhen, China

[4]South China University of Technology, Guangzhou, China

*These authors contributed equally to this work

**Correspondence:

zhang_adam@szbl.ac.cn (H.Z.)

blobel@chop.edu (G.A.B)

<u>SUMMARY</u>

NIPBL promotes chromatin loop extrusion by the cohesin complex until it stalls at
convergently oriented CTCF sites, leading to the formation of structural loops. However,
to what extent loop extrusion contributes to the establishment vs maintenance of *cis*-
regulatory element (CRE) connectivity is poorly understood. Here, we explored the de
novo establishment of chromatin folding patterns at the mitosis-to-G1-phase transition
upon acute NIPBL loss. NIPBL depletion primarily impaired the formation of cohesin-
mediated structural loops with NIPBL dependence being proportional to loop length. In
contrast, the majority of CRE loops were established independently of loop extrusion
regardless of length. However, NIPBL depletion slowed the re-formation of CRE loops
with weak enhancers. Transcription of genes at NIPBL-independent loop anchors was
activated normally in the absence of NIPBL. In sum, establishment of most regulatory
contacts and gene transcription following mitotic exit is independent of loop extrusion.

<u>INTRODUCTION</u>

Mammalian genomes are organized in a nonrandom, multi-layered manner within the
nucleus. Spatial organization is intimately connected to gene regulation as, for example,
distal transcriptional enhancers are in physical proximity with their promoters[1,2]. An
important player in the establishment and maintenance of genome structure is the

cohesin complex, which actively extrudes the chromatid until it encounters convergently
oriented CTCF sites, leading to the formation of looped contacts, also referred to as
structural loops[3-6]. These loops give rise to topologically associating domains (TADs),
which promote regulatory interactions within their boundaries and disfavor contacts
across them[7-11].

TADs and structural loops are dynamic entities, persisting for mere minutes and are
thus subject to constant turnover[12,13]. Cohesin activity is modulated by factors such as
WAPL/PDS5[14-18], which facilitate the release of cohesin from chromatin, and NIPBL, a
cohesin activator. NIPBL was identified in Drosophila as a facilitator of long-range
enhancer-promoter communication[19] and is required for cohesin-mediated loop
extrusion[14,20]. While NIPBL has been thought of as a cohesin loader onto chromatin [21-
25], recent evidence suggests an additional role, in which NIPBL primarily enhances the
processivity of cohesin extrusion[15,26-30]. In-vitro biochemical experiments demonstrated
that cohesin's ATPase activity is greatly stimulated by NIPBL, providing a mechanistic
explanation for how NIPBL enhances cohesin processivity[27,28].

Despite the established roles of CTCF and cohesin in shaping genome architecture, the
contribution of the CTCF/cohesin machinery to cis-regulatory element (CRE)
connectivity is limited. The transcription of most genes and the interaction between
numerous CREs persist following acute CTCF or cohesin depletion[18,31-36], pointing at
the existence of CTCF/cohesin-independent mechanism that connect CREs. However,
to what extent cohesin-mediated loop extrusion is necessary for the initial establishment
of CRE interactions versus their maintenance, remains to be explored more deeply.

Mitosis is characterized by a dramatic reorganization of chromatin architecture, with
most measurable features such as compartments, TADs, and chromatin loops being
transiently disrupted, only to be rapidly re-established during G1 entry[37-41]. This unique
transition period offers a powerful opportunity to study the establishment of architectural
features and the factors involved.

The study of cohesin in chromatin architecture during mitotic exit is complicated by its
essential role in maintaining sister chromatid cohesion in pro-metaphase. Therefore, to
explore the role of loop extrusion in establishing chromatin architecture without
disrupting mitosis we depleted NIPBL. Specifically, we performed Hi-C in cells that were
depleted of NIPBL during mitosis, followed by synchronous release into G1. We found
that most, but not all, cohesin-dependent structural loops failed to form. The
dependence of structural loops on NIPBL was associated with loop length - longer loops
were more sensitive to NIPBL depletion than shorter ones. Importantly, most CRE loops
and gene transcription were re-established normally. Hence, loop extrusion only plays a
minor role in establishing CRE connectivity implying alternative mechanisms, such as
transcription (co-)factor clustering by which CREs are formed in the 3D space.

RESULTS

NIPBL supports the formation of TADs and stripes

To test the requirement of NIPBL for the de novo formation of chromatin architecture
genome-wide, we tagged endogenous NIPBL bi-allelically with an 5-Ph-IAA-inducible
degron[42,43] and a HALO tag in the G1E-ER4[44] erythroblast cell line using CRISPR-
mediated gene editing. We introduced an osTIR2-expressing construct into the cells to
allow for 5-Ph-IAA-inducible NIPBL degradation. NIPBL was virtually completely
degraded upon 4 hrs of 5-Ph-IAA treatment, as measured by western blot in cell lysates
(Fig. 1a). To investigate the impact of NIPBL loss specifically during the mitosis-to-G1
transition, a period when chromatin architecture is rebuilt, we treated NIPBL-AID cells
with 5-Ph-IAA for 4 hours during nocodazole-induced prometaphase arrest and
continued the 5-Ph-IAA treatment upon nocodazole release into G1. To analyze the
temporal dynamics of chromatin architecture formation upon NIPBL depletion, we
collected 3 timepoints during G1 entry; two in early G1 (45min and 1hr) and one in late
G1 (4hr) (Fig. 1b). 2N cells were purified based on DNA content via DAPI staining to
ensure appropriate release into G1, followed by in situ Hi-C.

We began testing the role of NIPBL in establishing chromatin architecture by analyzing
the re-formation of TADs. Using the hicFindTads function from HiCExplorer[45], we
identified a total of 2,257 TADs in control cells in late G1 (4 hours). NIPBL loss globally
impaired TAD re-formation (Fig.1c). However, residual TADs exhibited a progressive
strengthening as cells advanced through G1, possibly due to NIPBL-independent
extrusion by cohesin. To validate that the NIPBL-dependent TADs are indeed cohesin-
dependent structures, we analyzed Hi-C data from asynchronously growing (mostly
interphase) G1E-ER4 cells bearing AID tagged SMC3 (cohesin) alleles, treated with 5-
Ph-IAA for 4 hours[46]. SMC3 depletion led to robust disruption of these TADs, confirming
that the domains in question are indeed cohesin-mediated.

To independently assess NIPBL's influence on cohesin-mediated loop extrusion, we
analyzed the formation of "stripes" or "flares" in NIPBL-depleted cells using the Stripenn
algorithm[47]. Stripes, which appear as horizontal or vertical lines on Hi-C maps,
represent loop extrusion intermediates[6,36,48]. We found that stripes were virtually absent
at early G1 timepoints following mitotic NIPBL depletion and began to appear by late
G1, although they remained less intense and shorter than those observed in control
cells (Fig. 1d,e). Control cells exhibited robust stripe formation in early G1, reflecting
strong cohesin activity upon immediate entry into G1. Thus, mitotic NIPBL depletion
impairs cohesin activity, potentially by reducing its extrusion rate and/or efficiency.

Mitotic NIPBL loss strengthens compartmentalization

Prior reports suggest that cohesin-mediated loop extrusion counteracts genome
compartmentalization[20,49,50]. To test whether NIPBL loss enhances the segregation of A
and B-type compartments during mitotic exit, we measured genome-wide
compartmentalization as cells entered G1 in the presence or absence of NIPBL. NIPBL
loss strengthened compartmentalization as cells progressed into G1. This was
evidenced by a sharper distinction between A and B compartments and increased intra-
compartment versus inter-compartment interactions (Extended Data Fig. 1c-e).

CRE loops can form independently of loop extrusion

CRE loops can persist upon acute depletion of CTCF or cohesin in interphase,
suggesting that alternative mechanisms maintain CRE connectivity[34,51-53]. However,
whether cohesin-mediated extrusion is required for the initial establishment of CRE
loops upon mitotic exit remains to be directly tested. Indeed, it has been proposed that
cohesin may be more essential for the establishment of de novo loops/gene expression
profiles during cell state transitions than it is for maintaining them[54]. We identified loops
using the cooltools.dots function in untreated 4hr G1 and 5-Ph-IAA-treated 4hr G1
samples. We quantified loop strength for each loop in each condition by measuring the
observed contact frequency within the center (peak) pixel divided by a locally adjusted
expected value. We then assigned a log2FC value comparing NIPBL replete and
depleted conditions for each loop and used a log2FC cutoff of -/+ 0.5 to define
differential loops.

We classified loops into three categories based on our prior annotations[41]: 1) structural
loops – CTCF/RAD21(cohesin) peaks at both anchors and CREs present at one or no
anchor, 2) CRE loops – CREs at both anchors and CTCF/RAD21 peaks at one or no
anchor, 3) dual function loops – CREs at both anchors and CTCF/RAD21 peaks at both
anchors (Fig. 2a). "Anchors" we defined as 20kb windows around the centers of the
contacts. NIPBL depletion predominantly impaired the formation of structural and dual-
function loops. Specifically, at late G1, 56% of structural loops failed to achieve maximal
intensity while most of the rest were delayed in their formation (Fig. 2b-d). Similar
results were observed for dual function loops. In stark contrast, 76% of CRE loops
formed at similar rates and achieved normal intensities in the absence of NIPBL.

If an extrusion process is involved in long range contact formation, loop length and
dependence on NIPBL are expected to correlate. Indeed, NIPBL's influence on cohesin-
dependent structural loops was proportional to loop length (longer structural loops
tended to be more sensitive to NIPBL depletion than shorter ones) (Fig. 2e,f).
Interestingly, this trend did not exist for CRE loops which could be robust to NIPBL loss
across length scales (Fig. 2e). Thus, CRE loops can be established independently of
NIPBL/cohesin-mediated loop extrusion.

Normal activation of most genes in G1 in the absence of NIPBL

To test whether loop extrusion is required to establish transcription, we measured
nascent transcription by TT-seq in cells depleted of NIPBL during mitosis and released
into G1 phase for either 45 minutes or 4 hours. NIPBL loss resulted in relatively modest
gene expression changes, with 187 genes differentially expressed at the 45-minute
timepoint and 188 at 4 hours (log2FC cutoff 1/-1, Padj < 0.05) (Fig. 3a). 93 differentially
expressed genes (DEGs) were shared between both timepoints (Fig. 3b). Genes known
to be activated in early G1 such as Fos and Jun were not differentially expressed at
either timepoint, suggesting that NIPBL depletion does not interfere with mitotic release
and/or G1 entry.

Of note, NIPBL-dependent genes did not overlap with NIPBL-dependent CRE loop
anchors (not shown). In contrast, these genes were enriched at architectural stripes,
TAD boundaries, and NIPBL-dependent dual-function loop anchors (Fig. 3c-d). Thus,
the small subset of NIPBL-dependent genes likely relies on structural loops (including
dual function loops) to promote CRE proximity, explaining their dependence on NIBPL.

One caveat to this analysis is that the genome undergoes a global spike in
transcriptional activity during the M-G1 phase transition[55]. This spike may mask NIPBL-
dependent changes in gene expression. Nevertheless, our data suggest that
transcription is re-established mostly normally, despite loop extrusion inhibition.

LDB1- and YY1-dependent loops can form in the absence of NIPBL

What mechanisms drive CRE connectivity in the absence of NIPBL? To this end, we
focused on LDB1 and YY1- two architectural factors previously shown to have cohesin-
independent roles[52,53,56]. Using Micro-C datasets generated in asynchronous LDB1-AID
and YY1-AID G1E-ER4 cells, we tested whether loops dependent on LDB1 or YY1 were
affected by mitotic NIPBL depletion. Focusing on loops with CREs in both anchors (this
includes dual-function and CRE loop categories), we found that the majority of loops
dependent on either LDB1 or YY1 were able to form in the absence of NIPBL (Fig. 4a,b
and Extended Data Fig. 2a,b). However, a subset of LDB1/YY1 loops were NIPBL-
dependent and tended to be closely flanked by encompassing structural loops
(consistent with previous reports of cohesin-dependent CRE loops[51,53] Fig. 4c and
Extended Data Fig. 2c). Additionally, another subset of LDB1/YY1 loops exhibited
delayed re-formation upon NIPBL depletion, but eventually recovered in late G1. Thus,
NIPBL is dispensable for the establishment of the majority of LDB1/YY1-mediated
regulatory interactions but may facilitate the timely formation of a subset of such loops.

An important technical consideration is that loop strength was calculated by dividing the
observed contacts within the loop's peak pixel by a locally adjusted expected value.
Since NIPBL depletion disrupts most TADs, global background contact frequencies are
lower (Extended Data Fig. 2d). Regardless, the strength (observed over locally-adjusted
expected signal) of LDB1/YY1-dependent/NIPBL-independent loops remains

unchanged or even increases. Measured increases upon NIPBL depletion may be
genuine or simply result from the peak pixels appearing stronger over the reduced local
background.

To further test whether YY1 and LDB1 mediated interactions can form in the absence of
NIPBL, we used an independent approach by generating pairs of YY1, LDB1, or RAD21
(as a control) ChIP-seq peaks. We created all possible pairs of peaks within a maximum
distance of 500kb from each other and averaged the contact frequency between these
pairs in the presence and absence of NIPBL (4hr post nocodazole release). As
expected, contacts between RAD21 sites were weakened by NIPBL depletion. Contacts
between LDB1 and YY1 peaks were retained to a greater extent, even though
background contact frequencies surrounding the "loop" center were reduced (Extended
Data Fig. 2e). This indicates that LDB1 and YY1-driven interactions can be established
independently of NIPBL/cohesin, providing an alternative mechanism of loop formation.

NIPBL Facilitates Rapid Formation of Loops Involving Weak CREs During Mitotic
Exit

CRE proximity to structural loops can render CRE loops dependent on NIPBL (Fig. 4c
and Extended Data Fig. 2c). However, prior work also suggested that CRE loops
involving weak enhancers tend to rely more on support by structural loops than those
involving stronger enhancers[51]. To investigate whether CRE activity is associated with
loop dependence on NIPBL, we grouped all asynchronous CRE loops into three
categories based on H3K27ac signal intensity at their anchors: top 25%, middle 50%,
and bottom 25%. Loops with the lowest H3K27ac levels were most affected by NIPBL
depletion and were delayed in their formation, while loops with the highest H3K27ac
levels remained largely unaffected and formed quickly during mitotic exit (Fig. 4d).
Under NIPBL replete conditions, CRE loops of all three groups were established at
comparable rates. Thus, weak enhancers tend to rely on extrusion to quickly establish
loops during mitotic exit even though many of these still eventually form in late G1.

DISCUSSION

The mitosis-to-G1 phase transition offers a unique opportunity to study the mechanisms
driving chromatin architecture formation. However, disrupting cohesin during the
mitosis-to-G1 phase transition is complicated due to its essential role in sister chromatid
cohesion and segregation. This challenge has precluded direct assessment of the role
of cohesin-mediated loop extrusion in re-establishing chromatin architecture. To
circumvent this challenge, we used mitotic degradation of NIPBL, a key activator of
cohesin. This approach allowed us to selectively impair cohesin's extrusion activity
during mitotic exit, providing valuable insights into NIPBL's role in regulating cohesin
and the necessity of loop extrusion for re-establishing chromatin architecture.

Long range structural loops are dependent on NIPBL while shorter ones can form but with delayed kinetics. The relationships of loop size and kinetics are consistent with a model in which NIPBL enhances cohesin's extrusion processivity, allowing it to extrude further and faster. This is consistent with in vitro experiments demonstrating an extrusion promoting function of NIPBL[27,28]. We surmise that the processivity effects are not simply the result of reduced cohesin loading in the absence of NIPBL. In an independent study, direct depletion of cohesin to the same degree as that observed upon NIBPL depletion had little effect on loop extrusion (Elphège Nora, personal communication). Therefore, our in vivo data are consistent with NIPBL's ability to enhance cohesin extrusion, rather than merely serving as a cohesin loader onto chromatin.

Our findings demonstrate that NIPBL is dispensable for the formation of many CRE loops, regardless of their length. This was true even for CRE loops spanning similar distances as structural loops which were dependent on NIPBL. Thus, CRE loops can form independently of cohesin-mediated extrusion and structural loop formation. Consistent with this, NIPBL depletion had minimal effects on transcription reactivation during mitotic exit, with only a small subset of genes being affected. Therefore, the majority of regulatory interactions likely form via alternative mechanisms, such as transcription (co)factor oligomerization/clustering.

For example, we showed that that NIPBL is largely dispensable for most LDB1- and YY1-driven interactions. Furthermore, the loops mediated by these two architectural proteins do not overlap significantly, suggesting that LDB1 and YY1 operate through distinct extrusion-independent mechanisms[53]. However, LDB1- and YY1-dependent loops represent only a subset of all CRE interactions[52,53], suggesting that additional (unidentified) factors are involved.

We found that the levels of H3K27ac at loop anchors were predicative of loop sensitivity to NIPBL depletion. In those cases in which loops with among weak CRE did form, they did so with delayed kinetics. Conversely, loops involving highly active enhancers exhibited similar formation kinetics in the presence and absence of NIPBL. This differential dependency upon NIPBL could result from lower transcription (co)factor concentrations at weak CREs, rendering them more reliant on loop extrusion to quickly facilitate long-range interactions.

In sum, by leveraging mitotic NIPBL degradation and synchronous release into G1, we directly tested the role of loop extrusion in the de novo establishment of chromatin architecture. While NIPBL is required for TAD formation, most regulatory interactions form independently. These findings reveal distinct extrusion-dependent and -independent mechanisms that organize chromatin architecture, uncoupling loop extrusion from affinity-based mechanisms in the formation of regulatory connectivity.

Fig. 1: NIPBL is required for domain and stripe formation after mitosis.

a Western Blot in asynchronous NIPBL-AID G1E-ER4 clones upon 4hr of 5-Ph-IAA
treatment. **b** Schematic of NIPBL depletion time course during G1 entry. **c** Pile-up plots
showing average signals for TADs identified in untreated 4h release condition. **d** Pile-up
plots showing average signals for stripes identified in untreated 4h release condition. **e.**
Stripe length for stripes identified in untreated 4h release condition compared to those
identified in 5-Ph-IAA-treated 4h release condition.

Fig. 2: Chromatin loops display distinct responses to NIPBL loss.

a. Schematic of loop stratification strategy. **b.** Differential loop analysis between
untreated and 5-Ph-IAA-treated 4h release conditions. (Top) scatter plots showing
individual loop responses to NIPBL depletion. (Bottom) APA plots showing average
signals for weakened, unchanged, and strengthened loops. **c.** Fraction of strengthened,
unchanged, and weakened loops that are classified as CRE, structural or dual-function.
d. Temporal dynamics of loop strength for loops that achieve normal intensities by late
G1 upon NIPBL depletion. **e.** Change in loop strength due to either NIPBL depletion (4h
nocodazole release) or SMC3 depletion (asynchronous) for structural and CRE loops.
Loops stratified by their length. **f.** Hi-C contact matrices of representative region
containing structural loops with variable reliance upon NIPBL. (Left) NIPBL-AID at 4h
release condition. (Right) SMC3-AID in asynchronous cells before/after 4h 5-Ph-IAA
treatment.

Fig. 3: Modest transcriptional changes upon NIPBL loss.

a. Volcano plots showing differential gene expression upon mitotic NIPBL depletion and
release into G1 for 45min (left) and 4h (right). **b.** Overlap of differentially expressed
genes between 45min and 4h timepoints. **c.** Enrichment of differently expressed genes
(4h timepoint) at stripe anchors, TAD boundaries and NIPBL-dependent dual-function
loop anchors. **d.** Hi-C contact matrices in the presence and absence of NIPBL (4h G1
release) of representative region containing NIPBL-dependent stripe anchored on
NIPBL-dependent gene.

Fig. 4: LDB1-dependent loops can form after mitosis in the absence of NIPBL.

a. (Top) Loop strength values during G1 entry for LDB1-dependent CRE loops classified
as NIPBL independent, delayed, and NIPBL dependent. (Bottom) Hi-C contact matrices
of representative regions containing LDB1-dependent loops that are independent,
delayed, and dependent on NIPBL. **b.** APA plots showing average signal for LDB1-
dependent CRE loops in the presence and absence of NIPBL (4h G1 condition). **c.**
Distance of LDB1-dependent CRE loops to encompassing structural loops. Only loops

with identified structural loops encompassing them are shown. **d**. Loop strength (right)
and observed contacts (left) for all asynchronous CRE loops during G1 entry in the
presence and absence of NIPBL. Loops are stratified by the average H3K27ac signal at
their anchors.

<u>EXTENDED DATA FIGURE LEGENDS</u>

Extended data Fig. 1: Hi-C replicate correlation and compartment analyses.

a. Heatmap showing Pearson correlation coefficients for Hi-C samples based on
eigenvector 1 values. **b**. (top) pile-up plots showing average TAD formation during G1
entry for each NIPBL-AID clone. (bottom) Pile-up plots showing average stripe
formation (normalized by stripe length) for each NIPBL-AID clone. **c.** (left) saddle plots
showing compartment strength during G1 entry at all timepoints. (right) quantification of
compartment strength. **d.** Pile-up plots showing average contact frequencies for A
compartments flanked by B compartments. **e.** Hi-C contact matrices showing
representative region exhibiting increased compartmentalization upon NIPBL depletion.

Extended data Fig. 2: LDB1- and YY1-dependent loops can form in the absence of
NIPBL.

a. YY1-dependent CRE loop strength during G1 entry in the presence and absence of
NIPBL. **b.** APA plots showing average signal for YY1-dependent CRE loops in the
presence and absence of NIPBL (4h G1 condition). **c.** Distance of YY1-dependent CRE
loops to encompassing structural loops. Only loops with identified structural loops
encompassing them are shown. **d.** Observed contacts for LDB1- and YY1-dependent
loops during G1 entry in the presence and absence of NIPBL. Loops stratified by NIPBL
independent, delayed, and NIPBL dependent. These are the same loops from Extended
Data Fig. 2a, and Fig. 4a. **e.** Paired chip-seq analysis. Asynchronous RAD21, LDB1,
and YY1 peaks were paired to create all possible pairs of peaks within a maximum
distance of 500kb. APA plots showing contact frequencies at late G1 (4h) in the
presence and absence of NIPBL for each set of peak pairs.

<u>DATA AVAILABILITY</u>

Hi-C and TT-seq datasets generated in this study are deposited in the GEO database
(GSE256073). External data from prior studies listed here: LDB1, CTCF, RAD21, YY1,
and H3K27ac ChIP-seq (GSE254377), LDB1 asynchronous Micro-C (GSE254373),
YY1 asynchronous Micro-C (GSE247254).

<u>METHODS</u>

Nocodazole arrest/release and 5-Ph-IAA treatment

The G1E-ER4:Wapl^{dTag}/Nipbl^{mAID} cells expressing OsTIR2 were treated with nocodazole
(200ng/ml) for 8 hours at a density of 0.5 million/ml. To degrade Nipbl during mitotic
arrest, 5-Ph-IAA (1uM) was added into the culture during the last 4 hours of nocodazole
arrest. To acquire post mitotic samples without Nipbl, the cells were were released from
nocodazole and resuspended in warm medium containing 5-Ph-IAA (1uM). The cells
were allowed to enter G1 phase for 45 minutes, 1 hour or 4 hours respectively. Cells
without 5-Ph-IAA treatment were collected in parallel as controls.

Cell sorting

For *in-situ* Hi-C experiments, control and 5-Ph-IAA treated samples were acquired as
described previously [51]. Briefly, at the time of collection, cells were pelleted, washed and
re-suspended in 1 x PBS. Cells were crosslinked with 2% formaldehyde for 10min at RT
and subsequently quenched with 1M (final concentration) glycine for 5min at RT. Cells
were pelleted and re-suspended in 1 x FACS sorting buffer (1 x PBS, 2% FBS, 2mM
EDTA and 0.02% NaN3) containing 20ng/ml DAPI at a density of about 50-100 million
cells/ml. G1 phase cells were sorted based on DAPI signal (2N). Sorted cells were
snap-frozen and stored at -80℃.

***In-situ* Hi-C.**

In-situ Hi-C was performed as previously described [57]. Briefly, purified cells (0.1-0.2
million) were lysed in 1ml cold Cell Lysis Buffer (10mM Tris pH 8, 10mM NaCl, 0.2%
NP-40) for 20min on ice. Subsequently, nuclei were washed pelleted and washed once
with cold 1.4×NEB buffer 3.1 (NEB Cat#B7203S). Nuclei were then incubated in 25ul
1.4× NEB buffer 3.1 containing 0.1% SDS for 10min at 65℃. SDS was then quenched
by 1% (final concentration) Triton X-100. Chromatin digest was performed by treatment
with 25U DpnII restriction enzyme for overnight at 37℃. After overnight digestion,
another stroke of DpnII (25U) was added into the system to further digest the genome
for 4 hours at 37℃. DpnII heat inactivation was performed at 65℃ for 20min. Digested
chromatin was then repaired with 6.6U of DNA Polymerase I, Large (Klenow) fragment
(NEB, Cat#M0210) in the presence of dCTP, dGTP, dTTP (Diamond Cat#B110049,
B110050, B110051) and Biotin-14-dATP (active motif, Catalog No: 14138). Blunt end
chromatin was then ligated with 2000 U of T4 DNA ligase (NEB, Cat#M0202M) for 4
hours at 16℃ and 4 hours at room temperature. Chromatin was then reverse-
crosslinked overnight at 65℃ in the presence of 1% SDS and proteinase K (G-clone
Cat#EZ0970). DNase-free RNase A (Vazyme Cat#DE111-01) was added to the system
for 30min at 37℃ to remove RNA. DNA was then extracted by phenol-chloroform
extraction, precipitated, and dissolved in 130☐l of nuclease free water. DNA was then
sonicated to fragments of 200-300bp using a QSonica Q800R3 sonicator (40%
amplitude, 15s ON and 15s OFF, 13min sonication). Fragmented DNA was purified
through 1.8x VAHTS DNA Clean Beads (Vazyme Cat#N411). Biotin labeled DNA

fragments were pulled down by Dynabeads MyOne Streptavidin C1 beads (Thermo
Fisher Scientific, Cat#65002). DNA capturing streptavidin beads were washed twice
with 1 × B&W buffer (5mM Tris-HCl, pH 7.5; 0.5mM EDTA; 1M NaCl) and subject to
bead library preparation using VAHTS Universal DNA Library Prep Kit for MGI (Vazyme,
Cat#NDM607-02) based on the manufacture's protocol. Adaptor ligated DNA library was
eluted in 0.1% SDS by incubating at 98℃ for 10min. Finally, the library was amplified for
6-8 cycles on a thermal cycler, using the VAHTS® HiFi Amplification Mix. PCR products
were then purified with 1.8x VAHTS DNA Clean Beads and sequenced on MGI
DNBSEQ-T7 sequencing platform.

<u>Quantification and data analysis</u>

Loop calling and differential loop analysis

.hic files were converted to cool files using hic2cool (v0.8.3). Cooltools.dots was used
on merged contact maps for each treatment condition to identify loops. We first
identified loops using 10kb and 25kb resolutions. We used the following parameters to
call dots: max_loci_separation=2_000_000, clustering_radius=20_000,
lambda_bin_fdr=0.05, n_lambda_bins=50. We utilized rounded donut, vertical,
horizontal and rounded lowleft kernels to define dots (pixels) that were enriched relative
to local backgrounds.

We created a consensus loop list by: 1- merging loops identified in the untreated (4hr
nocodazole release) and 5-Ph-IAA-treated (4hr nocodazole release) conditions for 10kb
and 25kb resolutions, 2- merging untreated/5-Ph-IAA loops from each resolution,
retaining the highest (smallest) resolution coordinates if a loop was called at both
resolutions. We then calculated loop strength at all consensus loops for untreated and
5-Ph-IAA-treated Hi-C contact matrices by calculating the observed/locally-adjusted
expected value. We calculated the locally-adjusted expected value by multiplying the
expected value at the peak pixel by the sum of the observed contacts in the rounded
donut region divided by the sum of the expected contacts in the rounded donut region.
Loop strength was calculated for each loop using the resolution at which the loop was
originally identified. Loops with NA, 0, or infinite strength values were removed to filter
out those in sparse regions in either untreated/5-Ph-IAA-treated conditions. We then
calculated a log2FC for each loop (5-Ph-IAA/untreated) and set a log2FC cutoff of -/+
0.5 to define weakened/strengthened loops upon NIPBL depletion.

Stratifying loops based on ChIP-seq peaks and CRE annotations

We stratified loops into different categories based on whether their anchors overlapped
(by at least 1bp) CTCF/cohesin ChIP-seq peaks and/or CREs. We used a list of
annotated CREs from our prior study[41] based on H3K27ac ChIP-seq and distance to
TSSs. Briefly, enhancers were defined as H3K27ac-enriched regions that were located

at least 1kb away from a TSS. Promoters were defined as H3K27ac-enriched regions
located within 1kb of a TSS. A list of asynchronous CTCF/cohesin co-occupied sites
were used from another prior study[53]. Loops were defined as structural loops:
CTCF/cohesin co-occupied sites in both anchors and CREs in 1 or 0 anchors, CRE
loops: CREs in both anchors, CTCF/cohesin co-occupied sites in 1 or 0 anchors, and
dual-function loops: CTCF/cohesin co-occupied sites in both anchors and CREs in both
anchors.

Quantifying asynchronous Micro-C loop strength

411
We used a list of asynchronous Micro-C loops from our prior study that had CREs in
both anchors (this included dual-function and CRE categories)[53]. We calculated the
dependence of each loop on LDB1[53], YY1[52] or NIPBL (this study) by utilizing contact
maps (Micro-C for LDB1/YY1, Hi-C for NIPBL) generated in the presence and absence
of each factor. We quantified loop strength for each loop as previously described using
10kb contact maps for each dataset. We filtered out loops that were in sparse regions in
any dataset by removing NA, 0 or infinite strength values. We further filtered for loops
with a minimum loop strength value of 1.2 in all datasets. Importantly, this conservative
strategy resulted in fewer loops that were dependent on either LDB1 or YY1 than we
previously reported. This is due to the fact that some LDB1/YY1-dependent loops may
be in sparse regions in the NIPBL datasets or not present/weak in late G1. Additionally,
short-range LDB1/YY1 dependent loops may be "missed" due to quantifying loop
strength at 10kb resolution.

We defined LDB1/YY1-dependent loops as those exhibiting a loop strength log2 fold
change (log2FC) of less than -0.5, corresponding to an approximate 30% reduction in
loop strength following LDB1 or YY1 depletion. To asses their responses to NIPBL
depletion, we then classified LDB/YY1 dependent loops into three categories based on
loop strength comparisons between untreated and 5-Ph-IAA-treated conditions (-
/+NIPBL) at 45min, 1hr and 4hr post-nocodazole release. NIPBL-independent loops-
loops with log2FC values greater than -0.5 at all time points, indicating minimal
influence of NIPBL. NIPBL-transiently affected loops-loops with log2FC values less than
-0.5 at either the 45min or 1hr time points, but greater than -0.5 at the 4hr time point,
suggesting a temporary reduction in loop strength due to NIPBL depletion. NIPBL-
dependent loops-loops with log2FC values less than -0.5 at the 4hr timepoint, indicating
that ultimately these loops could not form in the absence of NIPBL to the same degree
as in the untreated condition.

Identifying and Visualizing TADs and Stripes

438
439 We identified TADs using 10kb resolution contact maps using the HiCFindTADs function
440 from HiCExplorer with default parameters. We generated Pileups of all TADs identified
441 in the untreated 4hr nocodazole release condition using the python package Coolpup.

442 We identified stripes using 10kb resolution contact maps using Stripenn[47], and
443 generated pileups of all stripes from untreated 4hr nocodazole release with Coolpup.

A/B compartment assignment and saddle plotting

445 We used cooltools (v0.7.1) to compute cis eigenvector values on 100kb binned Hi-C
446 matrices from all NIPBL-AID treatment conditions. A compartments were identified as
447 those with positive EV1 values, and B compartments as those with negative EV1
448 values. This assignment was based on GC content. We generated saddle plots using
449 the cooltools.saddle function to show all AA, BB, AB, and BA interactions genome-wide.

TT-seq differential gene analysis

Strand-spec TT-seq reads were quantified in gene bodies using deepTools v3.5.5. We
quantified raw, mapped reads in positively-stranded genes using reads from the forward
strand and vice versa for negatively-stranded genes. We used the longest transcript of
Refseq annotated, mm9 genes. We then used DESEQ2 to perform differential
expression analysis.

1. Vermunt, M.W., Zhang, D., and Blobel, G.A. (2019). The interdependence of
gene-regulatory elements and the 3D genome. J Cell Biol *218*, 12-26.
10.1083/jcb.201809040.
2. Panigrahi, A., and O'Malley, B.W. (2021). Mechanisms of enhancer action: the
known and the unknown. Genome Biol *22*, 108. 10.1186/s13059-021-02322-1.
3. Rao, S.S., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D.,
Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., and Aiden,
E.L. (2014). A 3D map of the human genome at kilobase resolution reveals
principles of chromatin looping. Cell *159*, 1665-1680. 10.1016/j.cell.2014.11.021.
4. Hansen, A.S. (2020). CTCF as a boundary factor for cohesin-mediated loop
extrusion: evidence for a multi-step mechanism. Nucleus *11*, 132-148.
10.1080/19491034.2020.1782024.
5. de Wit, E., and Nora, E.P. (2023). New insights into genome folding by loop
extrusion from inducible degron technologies. Nat Rev Genet *24*, 73-85.
10.1038/s41576-022-00530-4.
6. Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny,
L.A. (2016). Formation of Chromosomal Domains by Loop Extrusion. Cell Rep
15, 2038-2049. 10.1016/j.celrep.2016.04.085.
7. Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and
Ren, B. (2012). Topological domains in mammalian genomes identified by
analysis of chromatin interactions. Nature *485*, 376-380. 10.1038/nature11082.
8. Nora, E.P., Lajoie, B.R., Schulz, E.G., Giorgetti, L., Okamoto, I., Servant, N.,
Piolot, T., van Berkum, N.L., Meisig, J., Sedat, J., et al. (2012). Spatial
partitioning of the regulatory landscape of the X-inactivation centre. Nature *485*,
381-385. 10.1038/nature11049.

9. Hou, C., Li, L., Qin, Z.S., and Corces, V.G. (2012). Gene density, transcription,
and insulators contribute to the partition of the Drosophila genome into physical
domains. Mol Cell *48*, 471-484. 10.1016/j.molcel.2012.08.031.
10. Sexton, T., Yaffe, E., Kenigsberg, E., Bantignies, F., Leblanc, B., Hoichman, M.,
Parrinello, H., Tanay, A., and Cavalli, G. (2012). Three-dimensional folding and
functional organization principles of the Drosophila genome. Cell *148*, 458-472.
10.1016/j.cell.2012.01.010.
11. Beagan, J.A., and Phillips-Cremins, J.E. (2020). On the existence and
functionality of topologically associating domains. Nat Genet *52*, 8-16.
10.1038/s41588-019-0561-1.
12. Gabriele, M., Brandao, H.B., Grosse-Holz, S., Jha, A., Dailey, G.M., Cattoglio, C.,
Hsieh, T.S., Mirny, L., Zechner, C., and Hansen, A.S. (2022). Dynamics of CTCF-
and cohesin-mediated chromatin looping revealed by live-cell imaging. Science
376, 496-501. 10.1126/science.abn6583.
13. Sabaté, T., Lelandais, B., Robert, M.-C., Szalay, M., Tinevez, J.-Y., Bertrand, E.,
and Zimmer, C. (2024). Universal dynamics of cohesin-mediated loop extrusion.
bioRxiv.
14. Haarhuis, J.H.I., van der Weide, R.H., Blomen, V.A., Yanez-Cuna, J.O.,
Amendola, M., van Ruiten, M.S., Krijger, P.H.L., Teunissen, H., Medema, R.H.,
van Steensel, B., et al. (2017). The Cohesin Release Factor WAPL Restricts
Chromatin Loop Extension. Cell *169*, 693-707 e614. 10.1016/j.cell.2017.04.013.
15. Davidson, I.F., and Peters, J.M. (2021). Genome folding through loop extrusion
by SMC complexes. Nat Rev Mol Cell Biol *22*, 445-464. 10.1038/s41580-021-
00349-7.
16. Tedeschi, A., Wutz, G., Huet, S., Jaritz, M., Wuensche, A., Schirghuber, E.,
Davidson, I.F., Tang, W., Cisneros, D.A., Bhaskara, V., et al. (2013). Wapl is an
essential regulator of chromatin structure and chromosome segregation. Nature
501, 564-568. 10.1038/nature12471.
17. Ouyang, Z., Zheng, G., Tomchick, D.R., Luo, X., and Yu, H. (2016). Structural
Basis and IP6 Requirement for Pds5-Dependent Cohesin Dynamics. Mol Cell *62*,
248-259. 10.1016/j.molcel.2016.02.033.
18. Wutz, G., Varnai, C., Nagasaka, K., Cisneros, D.A., Stocsits, R.R., Tang, W.,
Schoenfelder, S., Jessberger, G., Muhar, M., Hossain, M.J., et al. (2017).
Topologically associating domains and chromatin loops depend on cohesin and
are regulated by CTCF, WAPL, and PDS5 proteins. EMBO J *36*, 3573-3599.
10.15252/embj.201798004.
19. Rollins, R.A., Korom, M., Aulner, N., Martens, A., and Dorsett, D. (2004).
Drosophila nipped-B protein supports sister chromatid cohesion and opposes the
stromalin/Scc3 cohesion factor to facilitate long-range activation of the cut gene.
Mol Cell Biol *24*, 3100-3111. 10.1128/MCB.24.8.3100-3111.2004.
20. Schwarzer, W., Abdennur, N., Goloborodko, A., Pekowska, A., Fudenberg, G.,
Loe-Mie, Y., Fonseca, N.A., Huber, W., Haering, C.H., Mirny, L., and Spitz, F.
(2017). Two independent modes of chromatin organization revealed by cohesin
removal. Nature *551*, 51-56. 10.1038/nature24281.

21.   Gillespie, P.J., and Hirano, T. (2004). Scc2 couples replication licensing to sister
527   chromatid cohesion in Xenopus egg extracts. Curr Biol *14*, 1598-1603.
528   10.1016/j.cub.2004.07.053.
529   22.   Watrin, E., Schleiffer, A., Tanaka, K., Eisenhaber, F., Nasmyth, K., and Peters,
530   J.M. (2006). Human Scc4 is required for cohesin binding to chromatin, sister-
531   chromatid cohesion, and mitotic progression. Curr Biol *16*, 863-874.
532   10.1016/j.cub.2006.03.049.
533   23.   Murayama, Y., and Uhlmann, F. (2014). Biochemical reconstitution of topological
534   DNA binding by the cohesin ring. Nature *505*, 367-371. 10.1038/nature12867.
535   24.   Kurokawa, Y., and Murayama, Y. (2020). DNA Binding by the Mis4(Scc2) Loader
536   Promotes Topological DNA Entrapment by the Cohesin Ring. Cell Rep *33*,
537   108357. 10.1016/j.celrep.2020.108357.
538   25.   Gutierrez-Escribano, P., Newton, M.D., Llauro, A., Huber, J., Tanasie, L., Davy, J.,
539   Aly, I., Aramayo, R., Montoya, A., Kramer, H., et al. (2019). A conserved ATP- and
540   Scc2/4-dependent activity for cohesin in tethering DNA molecules. Sci Adv *5*,
541   eaay6804. 10.1126/sciadv.aay6804.
542   26.   Alonso-Gil, D., and Losada, A. (2023). NIPBL and cohesin: new take on a classic
543   tale. Trends Cell Biol *33*, 860-871. 10.1016/j.tcb.2023.03.006.
544   27.   Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.M. (2019).
545   DNA loop extrusion by human cohesin. Science *366*, 1338-1345.
546   10.1126/science.aaz3418.
547   28.   Kim, Y., Shi, Z., Zhang, H., Finkelstein, I.J., and Yu, H. (2019). Human cohesin
548   compacts DNA by loop extrusion. Science *366*, 1345-1349.
549   10.1126/science.aaz4475.
550   29.   Alonso-Gil, D., Cuadrado, A., Gimenez-Llorente, D., Rodriguez-Corsino, M., and
551   Losada, A. (2023). Different NIPBL requirements of cohesin-STAG1 and cohesin-
552   STAG2. Nat Commun *14*, 1326. 10.1038/s41467-023-36900-7.
553   30.   Petela, N.J., Gligoris, T.G., Metson, J., Lee, B.G., Voulgaris, M., Hu, B., Kikuchi,
554   S., Chapard, C., Chen, W., Rajendra, E., et al. (2018). Scc2 Is a Potent Activator
555   of Cohesin's ATPase that Promotes Loading by Binding Scc1 without Pds5. Mol
556   Cell *70*, 1134-1148 e1137. 10.1016/j.molcel.2018.05.022.
557   31.   Thiecke, M.J., Wutz, G., Muhar, M., Tang, W., Bevan, S., Malysheva, V., Stocsits,
558   R., Neumann, T., Zuber, J., Fraser, P., et al. (2020). Cohesin-Dependent and -
559   Independent Mechanisms Mediate Chromosomal Contacts between Promoters
560   and Enhancers. Cell Rep *32*, 107929. 10.1016/j.celrep.2020.107929.
561   32.   Rao, S.S.P., Huang, S.C., Glenn St Hilaire, B., Engreitz, J.M., Perez, E.M.,
562   Kieffer-Kwon, K.R., Sanborn, A.L., Johnstone, S.E., Bascom, G.D., Bochkov, I.D.,
563   et al. (2017). Cohesin Loss Eliminates All Loop Domains. Cell *171*, 305-320
564   e324. 10.1016/j.cell.2017.09.026.
565   33.   Nora, E.P., Goloborodko, A., Valton, A.L., Gibcus, J.H., Uebersohn, A., Abdennur,
566   N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted Degradation of
567   CTCF Decouples Local Insulation of Chromosome Domains from Genomic
568   Compartmentalization. Cell *169*, 930-944 e922. 10.1016/j.cell.2017.05.004.
569   34.   Hsieh, T.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Darzacq, X., and
570   Tjian, R. (2022). Enhancer-promoter interactions and transcription are largely

maintained upon acute loss of CTCF, cohesin, WAPL or YY1. Nat Genet *54*,
1919-1932. 10.1038/s41588-022-01223-8.

35. Rhodes, J.D.P., Feldmann, A., Hernandez-Rodriguez, B., Diaz, N., Brown, J.M.,
Fursova, N.A., Blackledge, N.P., Prathapan, P., Dobrinic, P., Huseyin, M.K., et al.
(2020). Cohesin Disrupts Polycomb-Dependent Chromosome Interactions in
Embryonic Stem Cells. Cell Rep *30*, 820-835 e810.
10.1016/j.celrep.2019.12.057.

36. Vian, L., Pekowska, A., Rao, S.S.P., Kieffer-Kwon, K.R., Jung, S., Baranello, L.,
Huang, S.C., El Khattabi, L., Dose, M., Pruett, N., et al. (2018). The Energetics
and Physiological Impact of Cohesin Extrusion. Cell *175*, 292-294.
10.1016/j.cell.2018.09.002.

37. Naumova, N., Imakaev, M., Fudenberg, G., Zhan, Y., Lajoie, B.R., Mirny, L.A.,
and Dekker, J. (2013). Organization of the mitotic chromosome. Science *342*,
948-953. 10.1126/science.1236083.

38. Gibcus, J.H., Samejima, K., Goloborodko, A., Samejima, I., Naumova, N.,
Nuebler, J., Kanemaki, M.T., Xie, L., Paulson, J.R., Earnshaw, W.C., et al. (2018).
A pathway for mitotic chromosome formation. Science *359*.
10.1126/science.aao6135.

39. Ito, K., and Zaret, K.S. (2022). Maintaining Transcriptional Specificity Through
Mitosis. Annu Rev Genomics Hum Genet *23*, 53-71. 10.1146/annurev-genom-
121321-094603.

40. Kadauke, S., and Blobel, G.A. (2013). Mitotic bookmarking by transcription
factors. Epigenetics Chromatin *6*, 6. 10.1186/1756-8935-6-6.

41. Zhang, H., Emerson, D.J., Gilgenast, T.G., Titus, K.R., Lan, Y., Huang, P., Zhang,
D., Wang, H., Keller, C.A., Giardine, B., et al. (2019). Chromatin structure
dynamics during the mitosis-to-G1 phase transition. Nature *576*, 158-162.
10.1038/s41586-019-1778-y.

42. Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., and Kanemaki, M.
(2009). An 5-Ph-IAA-based degron system for the rapid depletion of proteins in
nonplant cells. Nat Methods *6*, 917-922. 10.1038/nmeth.1401.

43. Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid Protein
Depletion in Human Cells by 5-Ph-IAA-Inducible Degron Tagging with Short
Homology Donors. Cell Rep *15*, 210-218. 10.1016/j.celrep.2016.03.001.

44. Weiss, M.J., Yu, C., and Orkin, S.H. (1997). Erythroid-cell-specific properties of
transcription factor GATA-1 revealed by phenotypic rescue of a gene-targeted
cell line. Mol Cell Biol *17*, 1642-1651. 10.1128/MCB.17.3.1642.

45. Ramirez, F., Bhardwaj, V., Arrigoni, L., Lam, K.C., Gruning, B.A., Villaveces, J.,
Habermann, B., Akhtar, A., and Manke, T. (2018). High-resolution TADs reveal
DNA sequences underlying genome organization in flies. Nat Commun *9*, 189.
10.1038/s41467-017-02525-w.

46. Zhao, H., Lin, Y., Lin, E., Liu, F., Shu, L., Jing, D., Wang, B., Wang, M., Shan, F.,
Zhang, L., et al. (2024). Genome folding principles uncovered in condensin-
depleted mitotic chromosomes. Nat Genet *56*, 1213-1224. 10.1038/s41588-024-
01759-x.

47.   Yoon, S., Chandra, A., and Vahedi, G. (2022). Stripenn detects architectural
616   stripes from chromatin conformation data using computer vision. Nat Commun
617   *13*, 1602. 10.1038/s41467-022-29258-9.
618   48.   Barrington, C., Georgopoulou, D., Pezic, D., Varsally, W., Herrero, J., and Hadjur,
619   S. (2019). Enhancer accessibility and CTCF occupancy underlie asymmetric TAD
620   architecture and cell type specific genome topology. Nat Commun *10*, 2908.
621   10.1038/s41467-019-10725-9.
622   49.   Liu, Y., and Dekker, J. (2022). CTCF-CTCF loops and intra-TAD interactions
623   show differential dependence on cohesin ring integrity. Nat Cell Biol *24*, 1516-
624   1527. 10.1038/s41556-022-00992-y.
625   50.   Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N., and Mirny, L.A. (2018).
626   Chromatin organization by an interplay of loop extrusion and compartmental
627   segregation. Proc Natl Acad Sci U S A *115*, E6697-E6706.
628   10.1073/pnas.1717730115.
629   51.   Zhang, H., Lam, J., Zhang, D., Lan, Y., Vermunt, M.W., Keller, C.A., Giardine, B.,
630   Hardison, R.C., and Blobel, G.A. (2021). CTCF and transcription influence
631   chromatin structure re-configuration after mitosis. Nat Commun *12*, 5157.
632   10.1038/s41467-021-25418-5.
633   52.   Lam, J.C., Aboreden, N.G., Midla, S.C., Wang, S., Huang, A., Keller, C.A.,
634   Giardine, B., Henderson, K.A., Hardison, R.C., Zhang, H., and Blobel, G.A.
635   (2024). YY1-controlled regulatory connectivity and transcription are influenced by
636   the cell cycle. Nat Genet *56*, 1938-1952. 10.1038/s41588-024-01871-y.
637   53.   Aboreden, N.G., Lam, J.C., Goel, V.Y., Wang, S., Wang, X., Midla, S.C., Quijano,
638   A., Keller, C.A., Giardine, B.M., Hardison, R.C., et al. (2024). LDB1 establishes
639   multi-enhancer networks to regulate gene expression. Mol Cell.
640   10.1016/j.molcel.2024.11.037.
641   54.   Cuartero, S., Weiss, F.D., Dharmalingam, G., Guo, Y., Ing-Simmons, E., Masella,
642   S., Robles-Rebollo, I., Xiao, X., Wang, Y.F., Barozzi, I., et al. (2018). Control of
643   inducible gene expression links cohesin to hematopoietic progenitor self-renewal
644   and differentiation. Nat Immunol *19*, 932-941. 10.1038/s41590-018-0184-1.
645   55.   Hsiung, C.C., Bartman, C.R., Huang, P., Ginart, P., Stonestrom, A.J., Keller, C.A.,
646   Face, C., Jahn, K.S., Evans, P., Sankaranarayanan, L., et al. (2016). A
647   hyperactive transcriptional state marks genome reactivation at the mitosis-G1
648   transition. Genes Dev *30*, 1423-1439. 10.1101/gad.280859.116.
649   56.   Krivega, I., and Dean, A. (2017). LDB1-mediated enhancer looping can be
650   established independent of mediator and cohesin. Nucleic Acids Res *45*, 8255-
651   8268. 10.1093/nar/gkx433.
652   57.   Zhao, H., Lin, Y., Lin, E., Liu, F., Shu, L., Jing, D., Wang, B., Wang, M., Shan, F.,
653   Zhang, L., et al. (2024). Genome folding principles uncovered in condensin-
654   depleted mitotic chromosomes. Nat Genet. 10.1038/s41588-024-01759-x.

Figure 1

a

b NIPBL depletion timecourse

c all TADs identified in untreated NIPBL-AID 4h condition

d all stripes identified in untreated NIPBL-AID 4h condition

e stripe length NIPBL-AID (4h)

P = 4.16 x 10⁻⁶

Figure 2

a loop stratification

structural
n=3,202

CRE
CTCF
cohesin

CRE
n=1,381

dual-function
n=1,037

b

structural
CRE
dual-function

loop strength
untreated 4h release vs 5-Ph-IAA 4h release

log₂ 5-Ph-IAA loop strength
log₂ untreated loop strength

structural loops
strengthened 127
unchanged 1,283
weakened 1,792

CRE loops
strengthened 324
unchanged 721
weakened 336

dual-function loops
strengthened 163
unchanged 461
weakened 413

weakened +NIPBL −NIPBL
unchanged +NIPBL −NIPBL
strengthened +NIPBL −NIPBL

log2 (obs/exp) 2.5 0

c proportion of differential loops at late G1 (4hr)

fraction of loops

loop type

loop categories
CRE
dual-function
structural

strengthened   unchanged   weakened

d Temporal dynamics of loops that achieve normal intensities by late G1 (4hr)

Structural loops
Untreated
5-Ph-IAA
Loop Strength
Prometa 45min 1h 4h

dual-function loops
Untreated
5-Ph-IAA
Loop Strength
Prometa 45min 1h 4h

CRE loops
Untreated
5-Ph-IAA
Loop Strength
Prometa 45min 1h 4h

e Loop length distribution

Structural loops
ΔNIPBL (G1 4h)
ΔSMC3 (async)
loop strength (log2FC)

CRE loops
ΔNIPBL (G1 4h)
ΔSMC3 (async)
loop strength (log2FC)

100kb 200kb 300kb 400kb 500kb 600kb 700kb 800kb 900kb 1Mb >1Mb
increasing loop length
100kb bins

f

ΔNIPBL (G1)
+NIPBL
weakened structural loops
unchanged structural loop
−NIPBL
0.01 0

ΔSMC3 (async)
+SMC3
−SMC3
0.01 0

CTCF 1.52 0
RAD21 1.12 0

CTCF 1.52 0
RAD21 1.12 0

Figure 3

a

TT-seq NIPBL depletion (45min)

Strongly downregulated (Log2FC < -1, Padj < 0.05): 85
Weakly downregulated (Log2FC -0.5 to -1, Padj < 0.05):183
Weakly upregulated (Log2FC 0.5 to 1, Padj < 0.05):131
Strongly upregulated (Log2FC > 1, Padj < 0.05):102

TT-seq NIPBL depletion (4hr)

Strongly downregulated (Log2FC < -1, Padj < 0.05): 107
Weakly downregulated (Log2FC -0.5 to -1, Padj < 0.05):135
Weakly upregulated (Log2FC 0.5 to 1, Padj < 0.05):75
Strongly upregulated (Log2FC > 1, Padj < 0.05):81

b

DEGs
log2FC<-1/or>1, Padj < 0.05

45min 94 93 95 4hr

c

Enrichment of DEGs at 3D genomic features (4hr timepoint)

Stripes

TAD boundaries

NIPBL-dependent
dual-function loop anchors

d

Dapk2 locus
(sustained downregulated)

+NIPBL

stripe

-NIPBL

Dapk2

CTCF

RAD21

Figure 4

a

LDB1 dependent CRE loops

NIPBL independent n=145

delayed n=113

NIPBL dependent n=96

Untreated - - -
5-Ph-IAA - - -

Loop Strength

Prometa   45min   1h   4h

chr8:67,363,443-69,099,866

chr10:16,800,000-17,500,100

chr13:51,878,822-52,362,777

Prometa

LDB1 (async)
CTCF (async)
RAD21 (async)
H3K27ac (async)

+NIPBL
1h
-NIPBL

+NIPBL
4h
-NIPBL

b

ΔNIPBL 4h

Structural loops | LDB1 dependent CRE loops
NIPBL independent | NIPBL dependent

+NIPBL

-NIPBL

log2FC

log2 (obs/exp)

log2 auxin/untreated

c

distance of LDB1 dependent CRE loops to encompassing structural loop

nearest distance to encompassing structural loop anchor (kb)

NIPBL independent   delayed   NIPBL dependent

Structural loop
CRE loop
CRE   CRE
Nearest distance (kb)

d

all asynchronous CRE loops

25% highest H3K27ac signal
middle 50% H3K27ac
25% lowest H3K27ac signal

loop strength

Untreated

Median loop Strength

Prometa   45min   1h   4h

5-Ph-IAA

Median loop Strength

Prometa   45min   1h   4h

observed contacts

Untreated

Median observed contacts

Prometa   45min   1h   4h

5-Ph-IAA

Median observed contacts

Prometa   45min   1h   4h

Extended Data Fig.1

a

Pearson correlation
(eigenvector 1, 100kb binned)

clone2 untreated 4h release
clone1 untreated 4h release
clone2 +aux 4h release
clone1 +aux 4h release
clone2 +aux 1h release
clone1 +aux 1h release
clone2 +aux 45min release
clone1 +aux 45min release
clone2 prometaphase
clone1 prometaphase

clone1 prometaphase
clone2 prometaphse
clone1 +aux 45min release
clone2 +aux 45min release
clone1 +aux 1h release
clone2 +aux 1h release
clone1 +aux 4h release
clone2 +aux 4h release
clone1 untreated 4h release
clone2 untreated 4h release

Pearson correlation
1
0

b

domains

-5-Ph-IAA
prometaphase   45min   1h   4h

+5-Ph-IAA
45min   1h   4h

Clone1

Clone2

1.8
1
0.57

stripes
normalized by stripe length

-5-Ph-IAA
prometaphase   45min   1h   4h

+5-Ph-IAA
45min   1h   4h

Clone1

Clone2

1.8
1
0.57

c

compartment saddle plots

prometaphase

BB   BA

EV1

AB   AA

EV1

untreated 4min release   untreated 1h release   untreated 4h release

+5-Ph-IAA 45min release   +5-Ph-IAA 1h release   +5-Ph-IAA 4h release

2
0.5

saddle strength

(AA + BB) / (AB + BA)

+5-Ph-IAA 4h release

untreated 4h release

extent

d

A compartments flanked by B compartments

untreated 4h release   +5-Ph-IAA 4h release

B
A
B

1.8
1
0.57

e

chr1:32.5Mb-42.5Mb
25kb resolution

0.0015
0

untreated 4h release

+5-Ph-IAA 4h release

chr1:34.9Mb-37.6Mb
10kb resolution

0.001
0

untreated

+5-Ph-IAA

EV1 untreated   2.5 / -1.5
EV1 5-Ph-IAA   2.5 / -1.5
A compartments
B compartments

Extended Data Fig.2

a

NIPBL independent
n=80

Untreated
5-Ph-IAA

Loop Strength

Prometa 45min 1h 4h

NIPBL transiently dependent
n=56

Untreated
5-Ph-IAA

Loop Strength

Prometa 45min 1h 4h

NIPBL dependent
n=68

Untreated
5-Ph-IAA

Loop Strength

Prometa 45min 1h 4h

b ΔNIPBL 4h

YY1 dependent CRE loops

NIPBL independent / NIPBL dependent

+NIPBL — log2 (obs/exp) 2

-NIPBL — log2 (obs/exp) 2

log2FC — log2 (auxin/untreated) 0 / -0.8

c

distance of
YY1 dependent CRE loops to
encompassing structural loop

0.068
* 0.612

nearest distance to encompassing structural loop anchor (kb)

NIPBL independent
NIPBL transient dependent
NIPBL dependent

Structural loop
CRE loop
CRE CRE
Nearest distance (kb)

d Observed contacts (absolute contact frequencies between loop anchors)

LDB1 dependent CRE loops

NIPBL independent
n=145

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

NIPBL transiently dependent
n=113

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

NIPBL dependent
n=96

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

YY1 dependent CRE loops

NIPBL independent
n=80

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

NIPBL transiently dependent
n=56

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

NIPBL dependent
n=68

Untreated
5-Ph-IAA

Observed contacts

Prometa 45min 1h 4h

e

Paired ChIP-seq peaks
(500kb max distance)

RAD21
n=434,110

LDB1
n=55,870

YY1
n=83,409

+NIPBL — log2 (obs/exp) 0.8

-NIPBL — log2 (obs/exp) 0.8