Smooth pursuit eye movements contribute to long-latency reflex modulation in the lower extremity

Oindrila Sinha[1], Matteo Bertucco[3], Isaac Kurtzer[4], Tarkeshwar Singh[1,2]

[1]Department of Kinesiology, The Pennsylvania State University, University Park, PA – 16802, USA
[2]Neuroscience Institute, The Pennsylvania State University, University Park, PA – 16802, USA
[3]Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy
[4]Department of Biomedical Sciences, College of Osteopathic Medicine, New York Institute of Technology, New York City, NY, USA

Running title: Smooth pursuit modulates lower limb long-latency reflex

Keywords: perturbation, posture stabilization, collision, EMG, center-of-pressure

Abstract word count: 218
Manuscript word count (Text + Figure captions + References): 3,693

ABSTRACT

Somatosensory mediated reactions play a fundamental role in adapting to environmental changes, particularly through long-latency responses (LLRs). We investigated how smooth pursuit eye movements (SPEM), which are slow eye movements used to track moving objects, influence LLRs of the upper and lower limb during mechanical interactions with moving objects. Seventeen participants stabilized their limb in anticipation of a collision with a virtual object approaching at 25 cm/s while standing. This task occurred while subjects either visually pursued the object or fixated a central location. Mechanical perturbations were applied at two time points: approximately 200ms and 60ms before the anticipated collision. On a random subset of trials, the robot applied a mechanical perturbation either 200ms (early) or 60ms (late) before the anticipated collision. As in previous studies LLRs were observed in leg muscles to upper limb displacement. Moreover, the leg LLRs were modulated by gaze, being larger during pursuit than fixation but only during late perturbations. This timing-specific modulation aligns with previous reports of policy transitions in feedback control roughly 60ms before impact. Upper limb LLRs were not impacted by gaze indicating a prioritization on postural control circuits. This work extends our understanding of the neural mechanisms underlying sensorimotor integration and highlights the sophisticated nature of the human motor control system in coordinating eye movements with whole-body postural responses.

NEW and NOTEWORTHY

We show for the first time that smooth pursuit eye movements contribute to modulation of long-latency reflexes in the lower limb during virtual object interactions in upright stance. Smooth pursuit eye movements are used to track moving objects. These findings suggest a neurophysiological link between control of eye movements and posture stabilization.

INTRODUCTION

Our sensorimotor system is well adapted to the complex and changing environment we inhabit. This competence is exemplified by a waiter receiving an unexpected bump and stabilizing their tray based on its content, the physical surroundings, and even shifting their weight to remain upright. Controlled experiments show that such sophistication is expressed as early as 50-60 ms post-perturbation and termed the long-latency reflex (LLR). LLRs exhibit coordinated responses across multiple effectors, including upper and lower extremities (1, 2). Further, LLRs are also modulated by task goals and experimental contexts (3, 4) and are influenced by inputs from multiple sensory modalities, including vision (5–7).

The modulatory impact of vision on somato-sensory evoked LLRs reflects the funneling of information from frontoparietal cortical areas onto motor cortical circuits which generate the LLR. In addition to modulation by static visual information, such as target size (6) and location (8), LLRs are scaled by the magnitude and direction of continuous motion signals, specifically the coherence of random-dot kinematograms as used in studies of decision-making studies (7). The key cortical substrate for motion processing is middle temporal complex (MT+) and its impact on LLRs is especially notable since MT+ plays a crucial role in smooth pursuit eye movements (SPEM) used to track moving objects (9, 10). During SPEM, an efference copy of the extraretinal signal associated with eye movements has been hypothesized to contribute to motion perception (12–14), to guide upper limbs actions such as interceptive hand movements (15), and anticipatory posture stabilization during interactions with moving objects (16, 17). The broad influence of SPEM on limb motor control and the linkage of MT+ to LLR modulation suggests an important but untested possibility: LLRs may be modulated by gaze behavior.

To test this hypothesis, we examined healthy individuals as they stood upright, countered a moving virtual object with their arm, and adopted different gaze behaviors: freely pursuing the moving object with their eyes or fixating a central location. On a random subset of trials, we applied mechanical perturbations to the arm to evoke LLRs of the upper and lower limbs. We applied mechanical perturbations either ~200 ms or ~60 ms prior to the impending collision as previous work on the role of vision for ball catching showed that stretch reflexes are modulated ~60 ms before collision to increase limb impedance (18, 19). We found that the leg LLRs, but not arm LLRs, were modulated by gaze behavior immediately before the impeding collision.

METHODS

Participants

Seventeen right-handed participants (age: 23.3 ± 1.2 years; 9 males, 8 females), with no history of neurological or musculoskeletal conditions, were recruited. Participants provided written informed consent and were compensated ($10/h). The study was approved by Penn State's Institutional Review Board.

Apparatus

Participants interacted with virtual stimuli using a Kinarm Endpoint robot (Kinarm, ON, Canada) integrated with an SR EyeLink 1000 Remote eye-tracker (SR Research, Canada). Visual stimuli were projected (120 Hz) onto a semi-transparent mirror via a VPixx display (VPixx Technologies, Canada). Direct vision of the hand was occluded (Fig. 1A). Participants stood upright on a force plate (Bertec, USA), head tilted ~30° forward, while manipulating the robot

handle. EMG data were recorded wirelessly (Trigno, Delsys Inc, USA) from bilateral tibialis anterior muscles and right biceps and triceps brachii. EMG data and center of pressure were sampled at 1,000 Hz. Hand forces were measured using an ATI-Mini40 force transducer (ATI Automation, USA) and also sampled at 1,000 Hz. The monocular eye-tracker, mounted 80 cm from participants, recorded left eye movements at 500 Hz (accuracy: 0.25-0.5°, data loss: 1.6 ± 0.9%, (20)) and was calibrated using Kinarm's calibration routine. Experiments were conducted in dim light. Participants held only the robot handle and kept their forehead slightly away from the head support to prevent receiving additional somatosensory cues.

Experimental task

Participants maintained their hand position within a virtual target (1×1 cm rectangle) while standing (Fig. 1A). Each trial began with cursor positioning at a start location, followed by fixation on a cross (see Fig. 1B). After the cross turned off (250-750 ms random delay), a virtual object appeared and moved toward the participant's hand position at 25 cm/s. In 75% of the trials, when the object collided with the hand, the robot applied a 10 N force impulse in the posterior direction (10 ms rise and fall times, 90 ms plateau) to simulate a collision (see Fig. 1C). Participants were instructed to maintain hand position within the target while resisting the force impulse during collision. In the remaining 25% of trials, mechanical perturbations (10N, 10 ms rise time) were applied before anticipated object collision, either 200 ms (Early) or 60 ms (Late) ms before contact. Perturbations were delivered as 10 N step forces in the anterior-posterior direction. The order of the trials was randomized within a block.

Experimental conditions

We used two gaze conditions in this experiment (central fixation, FC; and smooth pursuit eye movements, SPEM). In the smooth pursuit (SPEM) condition, participants were instructed to pursue the object all the way till contact. In the Fixation (FC) condition, the visual fixation point was positioned 5 cm anterior to the start position along the body midline (see Fig. 1D).

The experiment consisted of 10 blocks of 32 trials for each gaze condition (640 trials). We focused our analysis only on those perturbations that were directed toward the body (PerturbToward), comparing responses between central fixation (FC) and pursuit (SPEM) conditions.

Study design

Participants began the study with 16 EMG normalization trials. In these trials, they encountered a 10N static background force directed either toward them (8 trials) or away from them (8 trials). They were required to maintain arm position against this force, and the collected EMG signals were later used for normalization purposes. After this, they performed 10 blocks of FC and 10 blocks of SPEM gaze conditions. The order of the gaze conditions was randomized across participants.

Data recording and analyses

EMG signals were band-pass filtered (20-450 Hz), rectified, and smoothed using a 10-point moving average (for plotting). After baseline subtraction, signals were normalized to root mean square (RMS) values from normalization trials. For each gaze and experimental condition, trials were ensemble-averaged before quantifying response intervals. The long-latency feedback response (LLR) window for the upper limb was defined as 50-100 ms post onset of perturbation. For the lower extremity, the LLR window was extended to 75-120 ms to account for conduction

delays (2). The voluntary response windows were 100-150 ms for the upper extremity and 120-150 ms for the lower extremity. The mean of EMG signals within these windows was compared across gaze conditions.

Figure 1: Experimental setup, protocols, and gaze data. (A) Participants stood at a KINARM endpoint robot with force plate. (B) Trials began with hand positioning in a virtual target, followed by fixation cross and moving object presentation (25 cm/s). (C) Force profiles: collision trials (75%) delivered 90 ms trapezoidal forces; perturbation trials (25%) applied step forces either early (Early, ~200 ms pre-contact) or late (Late, ~60 ms pre-contact) in anterior-posterior directions. (D) Participants completed 10 randomized blocks each of smooth pursuit (SPEM) and central fixation (FC) conditions. (E) Single trial example (left panel) and mean gaze position (pursuit: red, fixation: blue) and object position (green) along the y-axis for all participants (right panel). (F) Pursuit percentage during fixation (FC) and pursuit (SPEM) conditions show that participants tracked moving objects with SPEM during pursuit condition.

Gaze data were preprocessed following established protocols (21). Point-of-regard (POR) data were low-pass filtered (15 Hz), converted to spherical coordinates, and transformed to ocular kinematics. Angular gaze speed was computed using a Savitzky-Golay filter (6th-order, 27-frame window). Gaze events (saccades, fixations, smooth pursuits) were classified using a threshold-based algorithm. Figure 1E shows gaze and target position along the Y-axis for two exemplary trials in the left panel and the mean across participants in the right panel. Kinetic and kinematic data were low-pass filtered at 50 Hz and 15 Hz, respectively, using a double-pass, zero-lag, third order filter. Center of pressure (COP) in the x (mediolateral)- and y (anterior/posterior)-directions was calculated with the following equations:

$$COP_X = \frac{-M_Y + F_Xd}{F_Z}$$

$$COP_Y = \frac{M_X - F_Yd}{F_Z}$$

Where F_x, F_y, and F_z represent ground reaction forces in the x-, y-, and z-directions, respectively, and M_x and M_y denote moments about the x- and y-axes. 'd' signifies the distance from the top of the force plate to its center.

Statistics

Data were tested for normality, homoscedasticity, and sphericity. If any of the assumptions were not met, data were log-transformed. A repeated measures ANOVA examined effects of Gaze Condition (FC, SPEM) and perturbation timing (Early, Late) on EMG amplitudes, center-of-pressure excursions, and hand force signals, with participants as random effects. Effect sizes were quantified using generalized η^2. Post-hoc comparisons were done using Holm-corrected pairwise tests ($\alpha = 0.05$).

RESULTS

To test our hypothesis, we constrained the gaze in the Fixation (FC) condition and allowed participants to freely pursue the moving object in the Pursuit (SPEM) conditions. Eye-tracking data confirmed successful gaze manipulation between conditions. During Fixation (FC), smooth pursuit duration averaged $0.0 \pm 0.1\%$, compared to $87.5 \pm 1.3\%$ during Pursuit (SPEM) blocks (see Fig. 1F). RM-ANOVA revealed a significant main effect of gaze condition [$F(1,33) = 22$, $P < 0.001$, $\eta^2 = 0.9$].

Pre-perturbation EMG amplitudes were comparable between gaze conditions across all recorded muscles. This baseline similarity was critical for comparing feedback responses, as short-latency response amplitudes typically scale with pre-perturbation activation, unlike long-latency responses (22, 23). Although short-latency responses were not our primary focus, controlled baseline activity enabled valid comparison of feedback responses. The pre-perturbation and pre-collision anticipatory EMG activity is shown in the supplementary material (Figs S1 and S2).

Constraining eye movements did not affect LLR amplitudes in the triceps brachii
Long-latency feedback responses in triceps brachii showed no significant gaze-dependent modulation during either the early or the late perturbations [Early: $F(1,31) = 2.3$, $p = 0.7$, $\eta^2 = 0.01$; Late: $F(1,31) = 6.1$, $p = 0.4$, $\eta^2 = 0.08$]. This suggests that extraretinal signals from smooth pursuit eye movements did not influence upper limb LLR responses, see Figure 2A for a representative participant data and Figure 2B for group-averaged responses. Responses are only shown for perturbations which stretched the triceps.

Figure 2: Long-latency reflexes in triceps brachii. The top panel shows the direction and timing of the perturbation applied by the robot. Left plot shows early (Early) and right plot shows late (Late) perturbations. (A) Representative single-participant data showing long-latency reflex (LLR, orange) and voluntary response (purple) windows. (B) Group-averaged EMG (n=17). No significant gaze-dependent modulation observed in either response window.

Constraining eye movements significantly decreased the amplitude of LLR in the lower limb muscles

LLR responses in bilateral tibialis anterior muscles showed gaze-dependent modulation, particularly during perturbations soon before the anticipated contact (Late perturbation). Figure 3A-B illustrates a representative participant's data (upper panels) and the group average (lower panels) for left and right tibialis anterior. The center of pressure adjustments (COP_y) remained unaffected (Supplementary Materials, Figure S3), suggesting selective influence of extraretinal signals on long-latency feedback responses, but not on overall metrics of posture stabilization.

Figure 3: Gaze-dependent modulation of lower limb reflexes. The top panel shows the direction and timing of the perturbation applied by the robot. EMG activity during early (Early, left) and late (Late, right) perturbations. (A) Left tibialis anterior: Single-participant example (top) and group average (n=17, bottom) showing gaze-dependent modulation in long-latency (orange) and voluntary (purple) windows. (B) Right tibialis anterior: Similar modulation pattern in single-participant (top) and group data (bottom).

Mean long-latency reflex responses) in tibialis anterior muscles were log-transformed and compared across the two gaze conditions at both perturbation times. Right tibialis anterior showed significantly larger long-latency feedback responses during pursuit compared to fixation [$F_{(1,31)} = 20$, $P < 0.05$, $\eta^2 = 0.2$; Fig. 4B]. During late perturbations, the differences between the gaze conditions in the left tibialis muscles approached significance [$F_{(1,31)} = 8.2$, $P = 0.1$, $\eta^2 = 0.03$; Fig. 4A]. These results indicate timing-specific modulation of postural reflexes by extraretinal signals, particularly in the right leg.

Figure 4: Long-latency reflex modulation. Box plots showing LLR amplitude (log-transformed EMG, 75-120 ms post-perturbation) during early (EARLY, left plots) and late perturbations (Late, right plots) for (A) left and (B) right tibialis anterior muscles. Gaze conditions: pursuit (SPEM) vs. fixation (FC). *P < 0.05.

DISCUSSION

Our findings replicated and extended multiple research domains. First, we demonstrated that upper limb perturbations elicited long-latency reflexes (LLRs) in the lower limb, consistent with previous observations in postural and limb perturbation studies (2, 24) and the broader framework of multi-segment integration in rapid feedback control (3). Second, we identified gaze-dependent modulation of leg LLRs, with enhanced responses during tracking of approaching objects with smooth pursuit eye movements (SPEM) compared to central target fixation. This gaze dependency aligns with extensive research on SPEM's role in motion processing (25) and interception movements (15), where SPEM demonstrates performance advantages and is preferentially selected in free-choice paradigms (16). Our previous work established that smooth pursuit eye movements (SPEM) lead to different patterns of anticipatory upper limb control when a seated subject must exert a matching force pulse against a colliding virtual object. Specifically, SPEM temporally advances response initiation and amplifies the anticipatory force magnitude compared to central fixation (16). Notably, peripheral fixation yields response patterns similar to SPEM, reflecting enhanced motion sensitivity in peripheral vision (17). The observed differential effects of SPEM versus central fixation on reactive compensation to mechanical perturbations complement our earlier findings on anticipatory compensation, suggesting substantial mechanistic overlap and supporting broader theoretical frameworks (26).

The gaze-dependency of leg LLRs manifested specifically during the period closest to anticipated collision, rather than earlier epochs. This temporal specificity indicates that the effect is not attributable to general SPEM-related upregulation, which was constant throughout, nor to global visual differences previously documented in vision versus no-vision LLR paradigms, where responses were enhanced in the no-vision condition (27). Instead, the observed modulation was epoch-specific, aligning with evidence for temporal adjustments in somatosensory processing (28), reflex engagement (5), and control strategies (29) immediately preceding mechanical interactions. The persistence of this pattern under visual occlusion remains unexplored and is an obvious next step.

Our findings likely reflect the integrative function of the middle temporal complex (MT+) in processing both retinal and extraretinal motion signals during smooth pursuit eye movements (30). MT+ engagement has been demonstrated in both small-object tracking and large-field motion processing tasks, such as random dot kinematograms (31), with its causal role in motion processing evidenced by post-ablation performance deficits (32). Building on Selen et al.'s (7) work that long-latency feedback response gains scale with decision time, we propose that MT+ motion signals may modulate premotor regions either directly or via parietal areas associated with perceptual evidence accumulation (33).

Our results strengthen evidence for MT+'s modulatory role in LLR gain control through integrated extraretinal and retinal signal processing and subsequent PMd signaling. However, our experimental design's temporal resolution, limited to two discrete perturbation timepoints, precluded investigation of continuous MT+ and premotor dynamics. Additionally, unlike random-dot motion paradigms requiring ongoing evidence accumulation, our task demanded discrete sensorimotor responses. Despite these methodological distinctions, both paradigms implicate MT+ in LLR gain modulation, suggesting a conserved role in sensorimotor integration.

The selective modulation of lower extremity feedback responses (Figs 3 & 4) was unexpected but aligns with established literature demonstrating context-dependent differential control of upper and lower limbs, such as during fixed-support versus surface perturbation conditions (34). This limb-specific modulation suggests preferential gaze-dependent postural control in the lower

extremities during standing. We hypothesize that similar gaze-dependent modulation would manifest in upper limb LLRs during seated conditions.

Taken together, the gaze-dependent modulation of LLRs was restricted to lower extremity muscles, suggesting a hierarchical organization of postural control during upper limb tasks. This selective modulation may reflect the prioritization of postural stability (35), where SPEM-associated extraretinal signals preferentially influence postural control circuits. The anatomical basis for this modulation could involve known projections from MT+ or superior colliculus—key nodes in the SPEM network—to the pontine nuclei and reticular formation, respectively. However, the precise neural mechanisms underlying this selective modulation and the relative contribution of these pathways to postural stability require further investigation.

References

1. **Kurtzer I**, **Pruszynski JA**, **Scott SH**. Long-latency responses during reaching account for the mechanical interaction between the shoulder and elbow joints. *Journal of Neurophysiology* 102: 3004–3015, 2009.

2. **Lowrey CR**, **Nashed JY**, **Scott SH**. Rapid and flexible whole body postural responses are evoked from perturbations to the upper limb during goal-directed reaching. *Journal of Neurophysiology* 117: 1070–1083, 2017.

3. **Kurtzer IL**, **Pruszynski JA**, **Scott SH**. Long-latency reflexes of the human arm reflect an internal model of limb dynamics. *Current Biology* 18: 449–453, 2008.

4. **Pruszynski JA**, **Omrani M**, **Scott SH**. Goal-dependent modulation of fast feedback responses in primary motor cortex. *Journal of Neuroscience* 34: 4608–4617, 2014.

5. **Lacquaniti F**, **Borghese N**, **Carrozzo M**. Transient reversal of the stretch reflex in human arm muscles. *Journal of Neurophysiology* 66: 939–954, 1991.

6. **Nashed JY**, **Crevecoeur F**, **Scott SH**. Influence of the behavioral goal and environmental obstacles on rapid feedback responses. *Journal of Neurophysiology* 108: 999–1009, 2012.

7. **Selen LPJ**, **Shadlen MN**, **Wolpert DM**. Deliberation in the motor system: reflex gains track evolving evidence leading to a decision. *Journal of Neuroscience* 32: 2276–2286, 2012.

8. **Ito S**, **Gomi H**. Visually-updated hand state estimates modulate the proprioceptive reflex independently of motor task requirements. *eLife* 9: e52380, 2020. doi: 10.7554/eLife.52380.

9. **Komatsu H**, **Wurtz RH**. Relation of cortical areas MT and MST to pursuit eye movements. I. Localization and visual properties of neurons. *Journal of Neurophysiology* 60: 580–603, 1988.

10. **Newsome WT**, **Wurtz RH**, **Komatsu H**. Relation of cortical areas MT and MST to pursuit eye movements. II. Differentiation of retinal from extraretinal inputs. *Journal of Neurophysiology* 60: 604–620, 1988.

11. **Dukelow SP**, **DeSouza JF**, **Culham JC**, **van den Berg AV**, **Menon RS**, **Vilis T**. Distinguishing subregions of the human MT+ complex using visual fields and pursuit eye movements. *Journal of Neurophysiology* 86: 1991–2000, 2001.

12. **Dichgans J**, **Wist E**, **Diener H**, **Brandt T**. The Aubert-Fleischl phenomenon: A temporal frequency effect on perceived velocity in afferent motion perception. *Experimental Brain Research* 23: 529–533, 1975.

13. **Gegenfurtner KR**, **Xing D**, **Scott BH**, **Hawken MJ**. A comparison of pursuit eye movement and perceptual performance in speed discrimination. *Journal of Vision* 3: 19–19, 2003.

14. **Spering M**, **Gegenfurtner KR**. Contextual effects on motion perception and smooth pursuit eye movements. *Brain Research* 1225: 76–85, 2008.

15. **Fooken J**, **Kreyenmeier P**, **Spering M**. The role of eye movements in manual interception: A mini-review. *Vision Research* 183: 81–90, 2021. doi: 10.1016/j.visres.2021.02.007.

16. **Sinha O**, **Madarshahian S**, **Gómez-Granados A**, **Paine ML**, **Kurtzer I**, **Singh T**. Smooth pursuit eye movements contribute to anticipatory force control during mechanical stopping of moving objects. *Journal of Neurophysiology* 129: 1293–1309, 2023. doi: 10.1152/jn.00075.2023.

17. **Sinha O**, **Rosenquist T**, **Fedorshak A**, **Kpankpa J**, **Albenze E**, **Bonnet C**, **Bertucco M**, **Kurtzer I**, **Singh T**. Predictive posture stabilization before contact with moving objects: equivalence of smooth pursuit tracking and peripheral vision. *Journal of Neurophysiology* 132: 695–709, 2024. doi: 10.1152/jn.00158.2024.

18. **Lacquaniti F**, **Borghese N**, **Carrozzo M**. Internal models of limb geometry in the control of hand compliance. *Journal of Neuroscience* 12: 1750–1762, 1992.

19. **Lacquaniti F**, **Carrozzo M**, **Borghese NA**. Time-varying mechanical behavior of multijointed arm in man. *Journal of Neurophysiology* 69: 1443–1464, 1993.

20. **Dunn MJ**, **Alexander RG**, **Amiebenomo OM**, **Arblaster G**, **Atan D**, **Erichsen JT**, **Ettinger U**, **Giardini ME**, **Gilchrist ID**, **Hamilton R**, **Hessels RS**, **Hodgins S**, **Hooge ITC**, **Jackson BS**, **Lee H**, **Macknik SL**, **Martinez-Conde S**, **McIlreavy L**, **Muratori LM**, **Niehorster DC**, **Nyström M**, **Otero-Millan J**, **Schlüssel MM**, **Self JE**, **Singh T**, **Smyrnis N**, **Sprenger A**. Minimal reporting guideline for research involving eye tracking (2023 edition). .

21. **Singh T**, **Perry CM**, **Herter TM**. A geometric method for computing ocular kinematics and classifying gaze events using monocular remote eye tracking in a robotic environment. *Journal of NeuroEngineering and Rehabilitation* 13: 10, 2016. doi: 10.1186/s12984-015-0107-4.

22. **Pruszynski JA**, **Kurtzer I**, **Lillicrap TP**, **Scott SH**. Temporal evolution of "automatic gain-scaling." *Journal of Neurophysiology* 102: 992–1003, 2009.

23. **Nicolozakes CP**, **Sohn MH**, **Baillargeon EM**, **Lipps DB**, **Perreault EJ**. Stretch reflex gain scaling at the shoulder varies with synergistic muscle activity. *Journal of neurophysiology* 128: 1244–1257, 2022.

24. **Marsden CD**, **Merton PA**, **Morton HB**. Human postural responses. *Brain* 104: 513–534, 1981.

25. **de la Malla C**, **Smeets JB**, **Brenner E**. Pursuing a target with one's eyes helps judge its velocity. *Perception* 51: 919–922, 2022.

26. **Scott SH**. A functional taxonomy of bottom-up sensory feedback processing for motor actions. *Trends in Neurosciences* 39: 512–526, 2016.

27. **Lacquaniti F**, **Maioli C**. Adaptation to suppression of visual information during catching. *Journal of Neuroscience* 9: 149–159, 1989. doi: 10.1523/jneurosci.09-01-00149.1989.

28. **Voudouris D**, **Fiehler K**. Dynamic temporal modulation of somatosensory processing during reaching. *Scientific Reports* 11: 1928, 2021.

29. **Lacquaniti F**, **Maioli C**. The role of preparation in tuning anticipatory and reflex responses during catching. *Journal of Neuroscience* 9: 134–148, 1989. doi: 10.1523/jneurosci.09-01-00134.1989.

30. **Dunkley BT**, **Freeman TC**, **Muthukumaraswamy SD**, **Singh KD**. Cortical oscillatory changes in human middle temporal cortex underlying smooth pursuit eye movements. *Human Brain Mapping* 34: 837–851, 2013.

31. **Maunsell JH**, **Van Essen DC**. Functional properties of neurons in middle temporal visual area of the macaque monkey. I. Selectivity for stimulus direction, speed, and orientation. *Journal of Neurophysiology* 49: 1127–1147, 1983.

32. **Newsome WT**, **Wurtz RH**, **Dursteler MR**, **Mikami A**. Deficits in visual motion processing following ibotenic acid lesions of the middle temporal visual area of the macaque monkey. *Journal of Neuroscience* 5: 825–840, 1985.

33. **Heekeren HR**, **Marrett S**, **Ungerleider LG**. The neural systems that mediate human perceptual decision making. *Nature Reviews Neuroscience* 9: 467–479, 2008.

34. **Nashner L**, **Cordo P**. Relation of automatic postural responses and reaction-time voluntary movements of human leg muscles. *Experimental Brain Research* 43: 395–405, 1981.

35. **Nardon M**, **Sinha O**, **Kpankpa J**, **Albenze E**, **Bonnet C**, **Bertucco M**, **Singh T**. Prioritized adjustments in posture stabilization and adaptive reaching during neuromuscular fatigue of lower-limb muscles. *Journal of Applied Physiology* 137: 629–645, 2024. doi: 10.1152/japplphysiol.00252.2024.