Genomic Resources for the Scuttle Fly *Megaselia abdita*: A Model Organism for
Comparative Developmental Studies in Flies

Running Title: Genomic resources for *M. abdita*

Authors:

Ayse Tenger-Trolander[1*] <ayse.trolander@gmail.com> (ORCiD 0000-0001-8319-0273)
Ezra Amiri[1] (ORCiD 0000-0002-6241-6656)
Valentino Gantz[2,3]
Chun Wai Kwan[1,4]
Sheri A Sanders[5]
Urs Schmidt-Ott[1*] <uschmid@uchicago.edu> (ORCiD 0000-0002-1351-9472)
* Authors for correspondence

Affiliations:
1. University of Chicago, Dept. of Organismal Biology and Anatomy, 1027 East 57th Street,
Chicago, Illinois 60637, USA
2. Section of Cell and Developmental Biology, University of California San Diego, La Jolla,
CA 92093, USA.
3. Pattern Biosciences, Inc. 681 Gateway Blvd, South San Francisco, CA 94080
4. Laboratory for Epithelial Morphogenesis, RIKEN Center for Biosystems Dynamics
Research, 2-2-3 Minatojima-minamimachi, Chuo-ku, Kobe, Hyogo 650-0047, Japan
5. Notre Dame University, 252 Galvin Life Science Center/Freimann Life Science Center,
Notre Dame, Indiana 46556, USA

Key Words: Genome assembly, genome annotation, transcriptome, non-traditional model
organism, evolutionary development, synteny analysis, orphan genes

Summary statement
We report a chromosome-level genome assembly and annotation and transcriptomes for an
emerging developmental model organism, the phorid fly Megaselia abdita, which is
phylogenetically intermediate between Drosophila and mosquitoes.

Abstract

The order Diptera (true flies) holds promise as a model taxon in evolutionary developmental
biology due to the inclusion of the model organism, *Drosophila melanogaster*, and the ability to
cost-effectively rear many species in laboratories. One of them, the scuttle fly *Megaselia abdita*
(Phoridae) has been used in evolutionary developmental biology for 30 years and is an
excellent phylogenetic intermediate between fruit flies and mosquitoes but remains
underdeveloped in genomic resources. Here, we present a *de novo* chromosome-level
assembly and annotation of *M. abdita* and transcriptomes of 9 embryonic and 4 postembryonic
stages. We also compare 9 stage-matched embryonic transcriptomes between *M. abdita* and *D.*
melanogaster. Our analysis of these resources reveals extensive chromosomal synteny with *D.*
melanogaster, 28 orphan genes with embryo-specific expression including a novel F-box LRR

gene in *M. abdita*, and conserved and diverged features of gene expression dynamics between
M. abdita and *D. melanogaster*. Collectively, our results provide a new reference for studying
the diversification of developmental processes in flies.
42
Introduction

Comparing related species is a powerful approach to understanding how mechanisms of
development evolve and diversify. The naturally occurring diversity of mechanisms and their
phylogenetic history can aid in revealing core principles and inform our understanding of
evolutionary transitions. Additionally, careful comparisons of multiple species in "model taxa" are
critical for determining the directionality of change and identifying mutations responsible for
important evolutionary shifts in developmental processes. Given that complex developmental
gene networks can enhance a population's permissiveness for the passive fixation of mutational
variants that open novel paths of adaptive evolution (Kimura and Ohta, 1974; Lynch, 2007a,
2007b), such mutations may not be adaptive and their significance as drivers of evolutionary
change might be overlooked.

While it is impractical to adapt a large set of closely related vertebrate model organisms for
laboratory studies, insects, in particular Diptera (true flies), offer a cost-effective alternative
(Grimaldi and Engel, 2005; Schmidt-Ott and Lynch, 2016; Wiegmann et al., 2011). Flies are
particularly appealing for the comparative study of developmental mechanisms because they
include a leading model organism in developmental biology, *Drosophila melanogaster*, and
many more species that are relatively easily cultured in laboratories. Developmental biologists
have introduced several new dipteran model organisms in recent years, including the
humpbacked fly, *Megaselia abdita*, which has been particularly useful for studying the evolution
of developmental mechanisms in dipteran embryos because of its technical advantages and
phylogenetic position (Rafiqi et al., 2011). It belongs to the large family Phoridae, also known as
scuttle flies (Disney, 1994; Li et al., 2024), and represents a lineage that separated from the
Drosophila lineage ca. 145 million years ago at the beginning of the Cyclorrhapha radiation,
roughly 100 million years into the dipteran radiation (Grimaldi and Engel, 2005). Developmental
biologists started using *M. abdita* as an experimental system in the 1990s to study the evolution
of axial pattern formation (Bullock et al., 2004; Crombach and Jaeger, 2021; Crombach et al.,
2016; Liu et al., 2018; Rohr et al., 1999; Schmidt-Ott et al., 1994; Stauber et al., 1999; Stauber
et al., 2000; Stauber et al., 2002; Wotton et al., 2015a; Wotton et al., 2015b; Wotton et al.,
2015c; Yoder and Carroll, 2006), the evolution of extraembryonic tissue (Caroti et al., 2018;
Fraire-Zamora et al., 2018; Horn et al., 2015; Kwan et al., 2016; Rafiqi et al., 2008; Rafiqi et al.,
2010; Rafiqi et al., 2012; Schmidt-Ott and Kwan, 2022; Stauber et al., 1999; Wotton, 2014;
Wotton et al., 2014), and other aspects of embryo development (Caroti et al., 2015; Dey et al.,
2023; Tanaka et al., 2015; Vicoso and Bachtrog, 2015). However, the limited availability of
genomic resources in *M. abdita* (Jimenez-Guri et al., 2013; Vicoso and Bachtrog, 2015) and the
Phoridae in general (Feng et al., 2020; Rasmussen and Noor, 2009; Zhong et al., 2016) has
limited the potential of this model organism by precluding genome-wide and epigenetic
experimental approaches.

Here we provide a *de novo* assembled and annotated chromosome-level genome for *M. abdita*,
alongside stage-specific transcriptomes across its life cycle. These resources provide an
excellent basis for genome-wide and epigenetic experimental approaches for an understudied
but phylogenetically important branch of the Diptera. They also establish synteny relationships
with the chromosomes of *D. melanogaster*, highlight conserved and divergent features of Hox
gene clusters, and provide insights into embryonic gene expression dynamics and orphan
genes. Collectively, our results will help to establish dipterans as a model taxon to study the
evolution of developmental mechanisms from gene regulation to neural networks and behavior.

Results and Discussion

Genome Assembly

We generated a *de novo* reference genome for *Megaselia abdita* using combined long read and
chromatin conformation capture methods obtained from several hundred embryos of a 10-
generation inbred line. Long-read, high-fidelity sequences were generated by PacBio (HiFi
PacBio reads), and chromatin conformation capture sequences were generated by Dovetail
Genomics' Omni-C method. After removing 33 scaffolds identified as contamination, the initial
draft assembly spanned 592.8 megabases (Mb) contained in 89 scaffolds with an N50 of
12.6 Mb. Using HiRise, a software designed to scaffold genome assemblies with proximity
ligation data (Putnam et al., 2016), the draft assembly was refined using the Omni-C reads
(Figure S1).

The final assembly is 592.8 Mb contained in 15 scaffolds with an N50 of 212.8 Mb (Table 1).
The genome size is comparable to a previous estimate of 562.7 Mb based on flow cytometry
data (Picard et al. 2012). We estimated heterozygosity at 0% - 0.24% (Figure S2A). We masked
67.9% of the genome, constituting repetitive elements (Figure S2B). The largest three scaffolds
correspond to the three chromosomes of *M. abdita* (Table S1). We aligned the remaining 12
scaffolds to the largest three and found that these scaffolds contain repetitive sequences that
cannot be correctly assembled into chromosome-level scaffolds.

Genome Annotation

We annotated the reference genome of *M. abdita* using evidence from RNAseq transcripts and
protein sequence databases, and a robust genome annotation pipeline (Figure 1). The process
involved mapping RNAseq data and assembling the transcriptome, mapping protein sequences
to the reference genome, and generating gene models using various prediction software. We
then created consensus gene models using EVidenceModeler, updated the models to include
UTRs and alternative isoforms, and filtered out transposable element models before functionally
annotating the genes with Eggnog-mapper (Haas et al., 2008; Huerta-Cepas et al., 2019;
Cantalapiedra et al., 2021). In the Materials and Methods section, we provide a detailed
walkthrough of this pipeline.

M. abdita's genome contains 11,934 protein-coding genes (Table 2). We assessed the quality
of the annotation with Benchmarking Universal Single-Copy Orthologs (BUSCO) (Simão et al.,

2015; Manni et al., 2021). BUSCO evaluates annotation completeness by looking for the
presence of highly conserved 'single-copy orthologs' across specific taxonomic groups. For
example, the BUSCO Eukaryota database expects 255 orthologs, while the Diptera database
expects 3,285. For the *M. abdita* genome annotation we found 93% eukaryotic and 88%
dipteran 'complete single copy orthologs', indicating a high-quality assembly (Table 2).

The genome size of *M. abdita* is significantly larger than that of *D. melanogaster* (592.8 Mb vs
139.5 Mb). We also found the mean gene length of protein-coding genes in *M. abdita* to be
significantly longer than that of *D. melanogaster*, ~19 kilobases (kb) vs ~6.9 kb respectively.
However, the lengths of transcripts and exons are remarkably consistent between the two
species indicating that *M. abdita's* larger genome size is partially attributable to longer introns
(Table S2).

Synteny analysis reveals significant collinearity between *M. abdita* and *D. melanogaster*
genomes, including HOX gene cluster arrangement

Analysis of chromosomal synteny can aid in the identification of orthologs and highlight genomic
regions with conserved regulatory potential. To investigate genome-wide synteny between *M.*
abdita and *D. melanogaster*, we performed a synteny analysis, identifying significant collinearity
between their genomes, including the arrangement of the split HOX gene cluster. We compared
the synteny and collinearity of *M. abdita* scaffolds with the *D. melanogaster* genome and
identified 387 collinear blocks encompassing 2,396 genes (Figure 2A). Large portions of *D.*
melanogaster chromosomes are syntenic with single scaffolds in *M. abdita*. Scaffold 1 of *M.*
abdita largely corresponds to chromosome arm 3L, chromosome 4, and chromosome arm 2R of
D. melanogaster, while scaffold 2 aligns with chromosome arm 2L and the distal portion of
chromosome arm 3R, and scaffold 3 with the proximal portion of chromosome arm 3R and the X
chromosome. The arrangement of the HOX genes in *M. abdita* mirrored that of *D.*
melanogaster, with distinct Antennapedia and Ultrabithorax complexes separated by 53,306 kb
in *M. abdita* (9,978 kb in *D. melanogaster*). Both complexes are located on Scaffold 2 of *M.*
abdita's genome and maintain the same gene order seen in *D. melanogaster*, including the
cuticle gene complex (Figure 2B). Additionally, the genes *zerknüllt* (*zen*) and *amalgam* have
undergone duplication in *M. abdita* (Figure 2B). *Zen* has experienced many duplications in
Diptera (Mulhair and Holland 2024). Since only the ~60 amino acid homeodomain of *zen* is
comparable between species, it is difficult to establish the relatedness of *M. abdita's zen-like* to
other *zen* genes and duplications by sequence alone.

M. abdita's *zen* gene is expressed in the serosa and has been characterized in detail (Caroti et
al., 2018; Kwan et al., 2016; Rafiqi et al., 2008; Rafiqi et al., 2010; Rafiqi et al., 2012; Stauber et
al., 1999). Consistent with these studies, we detected the *M. abdita's zen* transcript in
embryonic stages 5 - 15, coinciding with the time when the serosa is specified and maintained
(Figure S3). In contrast, the newly identified *zen-like* gene's expression was only detected at
stage 5 (Figure S3). Our findings underscore how a well-annotated genome can facilitate
precise comparisons of chromosomal architecture which will enable the identification of both
conserved regulatory landscapes and structural variation implicated in phenotypic diversity.

Major transition in transcriptional expression profile during germband extension and retraction

The extent to which gene expression is similar or different between stages and species provides a basis for identifying developmental windows of accelerated change, heterochronic shifts, and evolutionary divergence. To identify conserved features and differences between embryonic stages of *Megaselia abdita* and *Drosophila melanogaster*, we performed RNA-seq on single embryos from stages 1, 5, 8, 9, 10, 12, 13, 15, 16, and 17 for both species (Figure S4). We staged embryos based on morphology, corresponding to established staging schemes for each species (Campos-Ortega & Hartenstein, 1997; Wotton et al., 2014). We later excluded the *M. abdita* stage 16 embryo due to failed sequencing. Despite similar embryo sizes and uniform library preparation and sequencing conditions, *M. abdita* embryos had roughly twice the number of reads per embryo compared to *D. melanogaster* (Figure S5A) though the total number of genes with reads assigned was comparable between species with 10,373 in *D. melanogaster* to 9,999 in *M. abdita*. The number of genes expressed was similar across embryonic stages between species as well (Figure S5B). The number of reads mapping to genes doubled in *M. abdita* (8,941 vs. 3,863 reads/gene) consistent with its 2x read count. We used normalized read counts (transcript per million or TPM) to account for the global expression level differences, sample-to-sample variation, and differences in transcript length between genes. Our RNAseq data showed a major shift in transcriptional expression during germband retraction in both *M. abdita* and *D. melanogaster*. A multidimensional scaling (MDS) plot, which plotted samples based on the similarity of their top 500 most differentially expressed genes, revealed clustering of embryo transcriptomes before and after germband retraction (Figure 3A).

The transition from germband retraction to dorsal closure marks a shift from system-wide body plan development to nervous system-specific development in both *M. abdita* and *D. melanogaster*

To more closely look at the gene networks in both species, we identified the differentially expressed genes (DEGs) between developmental stages. Due to the absence of biological replicates for each embryo, we used a k-mean clustering approach to group samples and calculate a global dispersion estimate ($\sigma = 0.36$ for both *M. abdita* and *D. melanogaster*). Both species have high dispersion estimates but variance in expression was very similar in both datasets (Figure S6 & S7). Despite the high dispersion in data, we had sufficient power to detect DEGs with fold differences ±3 and p-value < 0.001.

We performed pairwise comparisons of each sequential stage: the zygote, containing maternal-deposited transcripts (Stage 1); cellularization, where maternal to zygotic transition (MZT) of gene expression occurs (Stage 5); gastrulation, germband extension & retraction (Stages 8-12); dorsal closure (Stages 13-16), and the final embryonic stage (stage 17). We detected 1,567 and 2,398 DEGs in *M. abdita* and *D.* in *melanogaster* respectively (Figure 3B). In approximately 1 hour of developmental time between stages 12 (germband retraction) and 13 (dorsal closure), both species showed a strikingly dynamic shift in gene expression. However, *D. melanogaster* had 1,463 DEGs compared to *M. abdita's* 564 DEGs at this same time point (Figure 3B). Conversely, *M. abdita* exhibited more dynamic gene expression earlier, between cellularization

and gastrulation (stages 5-8), with 418 DEGs compared to 123 in *D. melanogaster* (Figure 3B).
Between stages 8 and 12, there are markedly fewer DEGs in both species, corresponding with
the period of gastrulation and germband retraction (Figure 3B). This pause in the turning on and
off of genes is coincident with the phylotypic stage of development (Kalinka et al., 2010). It is
important to reiterate that this analysis highlights the most dynamic (largest changes in
expression) genes between sequential developmental stages – the total number of expressed
genes at any given stage was very similar between species (Figure S5B) - suggesting that *D.
melanogaster* exhibits sharper increases and decreases in expression than *M. abdita* after the
phylotypic phase ends while *M. abdita* seems to have evolved more dynamic expression before
the phylotypic phase begins.

To analyze the expression patterns of DEGs, we clustered them using DEGreports and plotted
the expression of each cluster (Figure S8) (Pantano 2024). The clusters confirmed that the most
dynamic transcriptional shifts occur during the transitions between stages 1 to 5, 5 to 8, and 12
to 13 for both species (Figure 3B). We then assessed whether the genes expressed in each
cluster were significantly enriched for any Biological Process Gene Ontology (BP GO) terms
using the Search Tool for the Retrieval of Interacting Genes/Proteins (String) (Szklarczyk et al.
2023) (Figure S8). *M. abdita*'s clusters 4 and 9 and *D. melanogaster's* clusters 6 and 10 were
significantly enriched for embryonic, body plan, and systems development terms (Figure 4A &
S9A) and contained many well-characterized developmental genes in Drosophila (Figure
S9B&C). Genes found in *M. abdita* cluster 1 and *D. melanogaster* clusters 2 and 11 exhibited a
pronounced increase in expression at stage 13 and were enriched for terms related to nervous
system, muscle, and cuticle development (Figure 4B). These findings highlight a conserved shift
toward nervous system development at the end of germband retraction in both species, while
also suggesting subtle differences in the timing and clustering of body plan-related gene
expression. In addition to their similarities, *D. melanogaster's* more extreme expression changes
between stages 12 and 13 (Figure 4 A-C) are notable because at these stages *Drosophila*
employs an evolutionarily novel mechanism involving a new tissue called the amnioserosa for
dorsal closure (Schmidt-Ott and Kwan, 2022) which may be contributing to the distinct
transcriptional dynamics observed between the species.

Our RNAseq analysis identified major transcriptional transitions between developmental stages
1 and 5, 5 and 8, and 12 and 13, revealing conserved and diverged patterns of gene expression
between *M. abdita* and *D. melanogaster*. This work was made possible by mapping the stage-
specific transcriptomes to the annotated genome, which allows us to link expression patterns to
specific genomic features.

**233 Investigation of 'orphan' genes exclusively expressed during embryogenesis in *M. abdita*
reveals novel F-Box LRR gene**

As the evolution of new genes within lineages and species is an important mechanism of
diversification of developmental mechanisms (Chen et al., 2010; Chen et al. 2013), we searched
M. abdita's genome for 'orphan' genes. Orphan genes are either highly diverged from known
sequences or represent newly evolved genes (Vakirlis et al., 2020; Xia et al., in press). In *M.
abdita's* genome annotation, eggnog-mapper was unable to assign orthologs to 1,049 gene

models. Further searches using *blastn* on the coding sequences and *blastp* on the predicted
protein sequences did not reveal any sequence similarity to other Dipterans. Approximately
8.7% of *M. abdita's* genes were 'orphan' genes.

We found that approximately 10% (109/1,049) of these orphan genes were expressed in
embryos, including 28 that were expressed *exclusively* during embryogenesis (Figure 5A). Most
of these genes (24) exhibited sharp expression peaks, with 14 peaking at stage 5
(cellularization), 4 at stage 12 (germband retraction), and smaller groups peaking at stages 13,
15, and 17. Four genes showed broader expression patterns, with two highly expressed during
stages 8-10 (germband elongation) and two across stages 5-12 (cellularization to germband
retraction).

To further characterize these 28 embryo-specific genes, we examined their open reading
frames and protein sequences (Table S3). We classified 21 as likely coding and of those we
further classified 9 as likely stable proteins (blue highlighted rows in Table S3). Additional
searches of InterProScan's database revealed no known protein domains or any domains
consistent with known transposases (Jones et al., 2014). We then used Alphafold2 to generate
protein structure models (Jumper et al., 2021). Two of these genes resulted in confident
structures (Table S3, pTM > 0.5). One of them could not be related to any known protein (Figure
5B); however, the second showed significant structural similarity to F-box leucine-rich repeat
(LRR) proteins (Figure 5C). F-box LRRs are components of the 'E3 ubiquitin ligase SCF
complex' which ubiquitinates targeted proteins for later degradation by the cell. Specifically, the
F-box domain binds to *Skp1* and the LRR domain binds to the target, the molecule slated for
ubiquitination and degradation.

We then compared our F-box LRR orphan protein sequence to known F-box-LRR genes in *D.*
melanogaster but found no obvious ortholog. We identified orthologs to *D. melanogaster* F-box
LRR genes *Ppa*, *Kdm2*, *Fbxl4*, *Fbxl7*, *Fbl6*, CG32085, CG9003, and CG8272. In total, we
identified 15 F-box LRR genes in *M. abdita (*8 with clear *D. melanogaster* orthologs, 6 with
orthologs in other dipteran species, and our orphan). One of the dipteran F-box LRR orthologs
in *M. abdita* (Scaffold 3, geneID #3336) had an identical expression profile (stage 5) to the
orphan F-box LRR.

F-box LRRs are known to ubiquinate important developmental signaling molecules in *D.*
melanogaster including the pair-rule gene *paired* (*prd*) which is bound by the F-box LRR protein
Partner of paired (*Ppa*) (Raj et al., 2000). *Ppa* is unusual in that its expression is patterned
rather than uniform as most F-Box LRR genes seem to be in *D. melanogaster* embryogenesis
(Das et al., 2002). Given that in *D. melanogaster*, the 12 best-known F-box LRR proteins (*Skp2*,
Ppa, *Kdm2*, *FipoQ*, *Fbxl4*, *Fbxl7*, *Fbl6*, CG32085, CG13766, CG12402, CG9003, CG8272) are
expressed throughout embryogenesis according to our both our RNAseq data and ENCODE
gene expression data, the stage-restricted expression of *M. abdita*'s orphan F-box LRR gene
and 13 other orphan genes with expression peaking before gastrulation might reflect previously
overlooked developmental differences between *M. abdita* and *D. melanogaster* at the
blastoderm stage.

280 **Genome browser and genomic analysis tools**

281 To improve accessibility and usability of the genomic data hosted on NCBI, we developed a
282 genome browser ecosystem centered on JBrowse2 (Diesh et al. 2023). This ecosystem is
283 available as a cloud image on NSF's Jetstream2 platform (image: *Megaselia abdita Genome*
284 *Resources*, Hancock et al. 2021, Boerner et al. 2023), with a portable Docker image currently in
285 development. The browser integrates tools for comprehensive genomic analysis, including
286 BLAST search (SequenceServer2.0, Priyam et al. 2019), CRISPR guide RNA design (modified
287 crisprDesigner, Beeber and Chain 2020), differential gene expression (DGE) analysis via R
288 Shiny (freecount, Brooks et al. 2024), and synteny mapping (ShinySyn, Xiao and Lam 2022).
289 Future updates will include expanded sgRNA profiling, Docker support for deployment on
290 commercial cloud platforms, and continuing optimization of workflows, ensuring this resource
291 remains a powerful and accessible tool for genomic research and education.

292 **Conclusions**

293 The scuttle fly *M. abdita* is an important non-traditional model organism with hitherto very limited
294 genomic resources. We have filled this gap by providing genomic and transcriptomic resources.
295 By assembling a chromosome-level genome and annotating it, we revealed substantial
296 chromosomal synteny with *Drosophila melanogaster* while uncovering many orphan genes.
297 Additionally, our comparative transcriptome analysis across embryogenesis highlights
298 conserved and divergent regulatory dynamics. The discovery of a novel F-box LRR gene,
299 expressed exclusively during embryogenesis, underscores the potential of *M. abdita* to reveal
300 new insights into the evolution of developmental gene networks. Ultimately, we hope the
301 addition of these resources will further comparative research of developmental mechanisms in
302 Diptera and continue the development of Diptera as a model taxon more broadly.

303 **Materials and Methods**

304 **Generation of inbred *M. abdita* line**
305 To generate single crosses of *M. abdita*, we collected pupae at the end of the pupal stage, when
306 pupae darken approximately 1–2 days before eclosion, and transferred them to 35 x 10 mm
307 petri dishes (Fisher Scientific, Cat. No. 50-820-644) until hatching. We monitored the plates
308 every morning and every two hours after to collect virgin females. We paired each virgin female
309 with a single male fly and allowed them to mate for two days in 35 x 10 mm petri dishes
310 containing a gel solution made from 2% agar in water. On the third day, we prepared egg-laying
311 vials by boiling 0.8 g of a fish food mixture—composed of 1 part spirulina flakes (Aquatic Eco-
312 Systems Inc., Cat. No. ZSF5) and 2 parts sinking powder (Aquatic Eco-Systems Inc., Cat. No.
313 F1A)—in 10 mL of a 0.8% agar solution (EMD Millipore, Cat. No. 1.01614). We added
314 approximately 1 mL of this solution to 10 x 75 mm culture tubes (Fisher Scientific, Cat. No. 14-
315 961-25) and allowed it to solidify. After cooling, we added 0.1 g of fish food on top using
316 weighing paper, followed by 200 µL of water. We used a cotton swab to compact the food and
317 clean any excess moisture from the sides of the tubes. We then transferred the mating pairs
318 from the agar plates to the culture tubes and plugged the tubes with rayon balls (TIDI, Cat. No.
319 969162). We established multiple single crosses and tracked them using a progressive

hierarchical code to identify the lineage. We conducted nine generations of sibling × sibling
single-pair matings across three separate parallel lineages (A, B, and C). At generation six, we
generated pooled crosses within each lineage to allow for the mixing of potentially lethal
recessive alleles that may have accumulated during the single-cross procedure. Following this,
we maintained the B lineage, as it exhibited higher overall fertility and health.

1.1 – 1.3 Genome Assembly

1.1 de novo library preparation, sequencing, and assembly

We generated a *de novo* reference genome for *M. abdita* with sequencing data from HiFi
PacBio reads and Dovetail's OmniC libraries (Cantata Bio). For HiFi PacBio sequencing, we
collected and snap-froze in liquid nitrogen ~600 dechorionated embryos (mostly stages 16 and
17) from a 10-generation inbred line of a previously established laboratory culture of *M. abdita*
Schmitz, 1959 (Schmidt-Ott et al., 1994). We sent these samples to Dovetail Genomics for HiFi
PacBio library preparation and sequencing. Library preparation circularizes fragments so they
can be read many times to generate a high-fidelity consensus sequence. Sequencing on the
SMRT (Single Molecule, Real-Time) nanofluidic chip generates the long reads inherent to
PacBio. PacBio generated 25.4 gigabase-pairs reads (42x coverage) from which Dovetail
generated a haplotype-resolved draft assembly using the Hifiasm assembler (Hifiasm1 v0.15.4-
r347 with default parameters) (Cheng et al. 2021). We used blobtools v1.1.1 to identify possible
contamination and removed 33 scaffolds from the assembly (Challis et al., 2020; Laetsch and
Blaxter, 2017). After filtering out haplotigs and contig overlaps with purge_dups v1.2.5, 89
scaffolds remained (Guan et al., 2020).

1.2 Omni-C library preparation and sequencing

We collected and snap-froze an additional ~600-700 dechorionated embryos and ~60 young
pupae in liquid nitrogen from the same inbred line of *M. abdita* for Omni-C Library preparation
and sequencing. Omni-C is a proprietary technology for long-range proximity ligation and
sequencing of genomic libraries which captures spatial information within the genome through
chromatin fixation and sequencing. Omni-C differs from other Hi-C preparations in that it digests
the chromatin with a sequence-independent endonuclease. This eliminates biases inherent to
competitive restriction enzyme-based approaches. Samples were treated with formaldehyde to
fix the chromatin and then digested with DNAse I. The resulting ends were then repaired, and
biotinylated bridge adapters were ligated to the ends. Subsequent steps involved proximity
ligation of adapter-ligated ends, reversal of formaldehyde-induced crosslinks, and DNA
purification. Non-internally ligated biotin residues were removed from the purified DNA.
Sequencing libraries were prepared with NEBNext Ultra enzymes and Illumina-compatible
adapters. Before PCR enrichment, biotin-containing fragments were isolated using streptavidin
beads. Sequencing was performed on the Illumina HiSeqX platform.

1.3 Scaffolding assembly with HiRise

Both the Hifiasm draft assembly and Dovetail OmniC sequencing reads were used as input for
HiRise, a software tailored for scaffolding genome assemblies with proximity ligation data
(Putnam et al., 2016). Based on spatial data from the chromatin conformation capture (Figure
S1), HiRise pinpoints regions where contigs are joined incorrectly (misjoins) in the initial
assembly and utilizes this spatial information to re-orient contigs and construct larger scaffolds.
The OmniC library sequences were aligned to the draft assembly using the Burrows-Wheeler
Aligner (Li and Durbin, 2009). HiRise-analyzed read pairs are mapped to the draft scaffolds to
develop a genomic distance likelihood model. This model is then used to identify and break
potential misjoins, score prospective joins, and execute joins surpassing a set threshold. These
scaffolds consist of sequentially arranged contigs separated by gaps. We used QUAST to
calculate %GC, N50, L50, and Ns per 100 kbp (Gurevich et al., 2014). We then repeat-masked
the genome with RepeatMasker (Smit et al., 2013). To estimate heterozygosity and sequencing
error rates, we calculated the frequency spectrum of canonical 21-mers using jellyfish, and input
the resulting histograms into GenomeScope (Marçais and Kingsford, 2011; Vurture et al., 2017).
Using NUCmer (mummer v3.23 software package), we aligned the non-chromosome size
scaffolds (4 – 15) back to the three chromosome-size scaffolds (1 - 3) and found repetitive
sequences present in scaffolds 4 – 15 (Kurtz et al., 2004; Marçais et al., 2018).

2.1 – 2.3 Genome Annotation

2.1 Repeat masking the genome

To generate a custom library for repeat masking, we ran RepeatModeler v2.0.4 on the genome
to find/model potential repeats (Smit and Hubley, 2008). We then used RepeatMasker's script
fambd.py v0.4.3 to extract Arthropoda records from the dfam database and combined these
sequences with the RepeatModeler output to use as the repeat library. We used RepeatMasker
v4.1.5 to soft-masked the genome with our custom library of repetitive low-complexity DNA and
transposable elements (Smit et al., 2013).

2.2 Data used as evidence of genome features

We downloaded available Illumina RNAseq data for *M. abdita* from NCBI which included paired
end reads from 3 adults and pooled embryos (Table S4). We generated RNAseq data for 9
precisely staged embryos, first and third instar larval stages, and a 1-day-old pupal stage (Table
S4). For protein evidence, we downloaded all Dipteran protein sequences from NCBI's RefSeq
database which included 2,122,027 sequences from 617 species (Sayer et al., 2022). We also
downloaded Uniprot's complete protein sequence file (UniProt Release 2023_04) which
contains 570,157 sequences from 14,509 species (The UniProt Consortium, 2023).

2.3 Annotation pipeline

We annotated the reference genome of *M. abdita* using evidence from all RNAseq transcripts
(Table S4), protein sequences, and gene prediction software (Figure 1). The choices of software
used in this pipeline are based on the methods section of VanKuren's 'Draft *Papilio alphenor*
assembly and annotation' (VanKuren 2023). First, we assembled RNA transcripts using two
transcript assemblers: Stringtie v2.2.1 and Trinity v2.15.1 (Pertea et al, 2015; Grabherr et al.,

2011; Haas et al., 2013). Trinity performs both genome-guided and *de novo* assembly; we
assembled transcripts with both methods. For Trinity and Stringtie's genome-guided
assemblies, we first mapped reads to the genome with 'Spliced Transcripts Alignment to a
Reference' software (STAR v2.7.10b) (Dobin et al., 2013). Next, we used the 'Program to
Assemble Spliced Alignments' (PASA v2.5.3) to identify gene structures from all three
assemblies (Haas et al., 2003). We predicted gene models directly from the PASA assemblies
using the PASA plug-in TransDecoder v5.7.1 which identifies candidate coding regions from
Trinity and StringTie assemblies. To create protein alignments, we used the software Exonerate
v2.2.0 which maps protein sequences to the genome (Slater and Birney, 2005). We used the
BRAKER3 pipeline (braker.pl v3.0.6) to predict gene models from mapped RNAseq reads and
protein data (Gabriel et al., 2023). BRAKER3 relies on the software Augustus and GeneMark to
predict gene models (Stanke et al., 2006; Brůna et al., 2020). We also generated our own *ab*
initio gene structure predictions, using GlimmerHMM v3.0.4 (Majoros et al., 2004). First, we
collected "hints" for training the *ab initio* predictors by extracting protein-coding hints from the
protein alignments using Augustus' exonerate2hints function, intron hints from mapped RNA
seq reads using Augustus' bam2hints function, and exon/intron hints from the PASA assemblies
using Augustus' bam2exonhints function. We additionally used the coding predictions from
Transdecoder to create training models and further refined those models using the
lib.selectTrainingModels function from Funannotate (Palmer and Stajich, 2019). This training
data was used to run GlimmerHMM with hints as guidance (Majoros et al., 2004). We then
provided the PASA assemblies, mapped protein data, TransDecoder predictions, *ab initio*
predictions, BRAKER3 predictions, and a file weighting each line of evidence to the software
Evidence Modeler v2.1.0. Evidence Modeler constructed the consensus gene structures which
were updated by PASA to add UTRs and identify alternative transcripts (Haas et al., 2008). We
removed gene structures that overlapped with RepeatMasker output using a Funannotate
function called "RemoveBadModels." We then used Funannotate v1.8.1 to identify annotations
that match known transposable elements (TEs) and repeat proteins using BLAST and updated
the annotation to remove remaining TEs. We used AGAT v1.4.1 to remove genes with an open
reading frame (ORF) < 100 amino acids in length and any associated gene structures (Dainat,
2022). We assigned gene names using eggNOG-mapper v2.1.12 which relies on orthology
predictions to functionally annotate genes (Cantalapiedra et al., 2021; Huerta-Cepas et al.,
2019).

428
Synteny Analysis
We used MCScanX (primary release) to compare the synteny of *M. abdita* and *D. melanogaster*
genomes (Wang et al., 2012). MCScanX identifies syntenic blocks based on a score given to
each gene pair. We set the match_score = 50 (default), match size = 5 (number of genes
required to constitute a syntenic block), gap_pentaly = 0 (no penalty for gaps), and max_gaps =
100. We used the output of this run to generate our synteny map. We used the circlize R
package to plot the results (Gu et al., 2014).

436
Embryo Staging for RNAseq

For precise embryo staging, embryos of the appropriate age were mounted on a microscope
slide under halocarbon 27 oil and observed in a Zeiss Axiophot compound microscope
equipped with a 10x objective and DIC (differential interference contrast) optics until they
reached the desired stage. They were photographed and immediately processed for RNA
extraction as previously described (Lott et al., 2014). We collected *M. abdita* and stage-matched
D. melanogaster embryos at stages 1, 5, 8, 9, 10, 12, 13, 15, 16, and 17. Photos of the
sequenced embryos can be found in Figure S4.
445
RNA isolation and Sequencing
We incubated each sample for 5-10 minutes at room temperature in TRIzol. We then froze each
sample in TRIzol at -80° C. We extracted total RNA using the TRIzol/Phenol-chloroform protocol
detailed in the appendix (Protocol 10) of 'Functional evolution of a morphogenetic gradient' by
Chun Wai Kwan. The University of Chicago genomics core facility constructed cDNA libraries
using the TruSeq kit with PCR (Illumina, CA, USA). The cDNA libraries were barcoded and
multiplexed for 100bp paired-end sequencing on one lane of a HiSeq Illumina 2000 sequencer.
453
Differential Gene Expression Analysis
M. abdita had ~2x the reads for each sample compared to *D. melanogaster* (Figure S5A). We
chose not to down sample *M. abdita* reads to match *D. melanogaster* to avoid losing power to
detect changes between genes within *M. abdita* developmental stages. We justified this by
looking for any evidence that the higher number of reads skewed the relationship between gene
expression and variance, but found that the relationship between mean expression and variance
in expression is similar in both species without subsetting the data (Figure S6). Additionally, the
average, median, and variance in gene expression (TPM) were very similar across development
in both species (Figure S7).
463
We aligned *M. abdita* RNAseq reads to the reference genome generated in this publication and
D. melanogaster RNAseq reads to *D. melanogaster*'s genome (Accession: GCF_000001215.4).
We used Subread v2.0.5 to align reads (Liao et al., 2013). 90-99% of reads mapped to the
genome for each sample. We input the aligned bam files into Subread's featureCount function
which assigns the reads to a genomic feature from the annotation file (gff). At this point, we
performed the analysis in Rstudio using the free and open source statistical language R and
various R packages including EdgeR, tidyr, dplyr, and ggplot2 (R Core Team, 2023; Robinson
et al., 2009; Posit Team, 2024; Wickham et al. 2016; Wickham et al., 2023; Wickham et al.,
2024). We used EdgeR to assess gene expression differences between samples. We filtered
out lowly expressed genes and normalized data by library size (TMM normalization) for both
species. We calculated counts per million (CPM) to normalize the difference in raw reads
between samples and then calculated transcript per million (TPM) to account for differences in
gene length. We calculated gene length as the coding sequence length for each gene's longest
isoform. All TPM expression data are plotted as $\log_2(\text{TPM} + 1)$.
478
479 As our RNA-seq samples do not include biological replicates, we estimated the squared
480 coefficient of variation (BCV) using k-means clustering of the samples. We began by calculating
481 a distance matrix for the samples and extracting the first four eigenvectors, which together

explained >95% of the variance. To determine the optimal number of clusters (k), we calculated
the within-cluster sum of squared errors (WSS) and Silhouette scores for k values ranging from
1 to 8, selecting k = 5 based on these metrics. For *M. abdita*, clustering with k = 5 grouped the
embryonic stages as follows: Group 1 (stage 1), Group 2 (stage 5), Group 3 (stages 8–12),
Group 4 (stages 13–15), and Group 5 (stage 17). For *D. melanogaster*, the clusters were:
Group 1 (stage 1), Group 2 (stages 5 and 8), Group 3 (stages 9–12), Group 4 (stages 13–16),
and Group 5 (stage 17). Using these groupings, we estimated the dispersion as σ=0.36 for both
M. abdita and *D. melanogaster.* These estimates were applied globally. Differential expression
analysis was performed using EdgeR's exactTest() function, comparing gene expression
between pairwise embryonic stages rather than the k-means groups. Genes were considered
differentially expressed if they met the criteria: fold change < -3 or > 3 and p-value < 0.001.
We then took all differentially expressed genes (DEGs) and input that expression data into
DEGreports pattern() function. DEGreports clusters DEGs based on expression profile similarity
(Pantano 2024). We used these clusters for gene ontology enrichment analysis described
below.
496
Gene Ontology Enrichment Analysis
We used the STRING database v12.0 (Search Tool for the Retrieval of Interacting
Genes/Proteins) to perform enrichment analysis of Biological Process Gene Ontology (GO)
terms for our clustered DEG lists (Szklarczyk et al., 2023). STRING compares input gene sets
to a reference genome to identify networks of interacting genes and enrichments in biological
processes. Since *M. abdita* is not available in STRING, we first used NCBI's Blast tool to select
the top *D. melanogaster* ortholog match/hit for each *M. abdita* gene (Altschul et al., 1990). Using
FlyBase's batch download tool, we retrieved the corresponding FlyBase IDs, which STRING
accepts as input (Öztürk-Çolak et al., 2024). STRING performed the enrichment analysis
identifying the Biological Process GO terms significantly associated with each developmental
stage. STRING measures enrichment based on the strength of enrichment
(Log_{10}(observed/expected)), false discovery rate (p-values corrected for multiple testing with
Benjamini-Hochberg), and the signal (weighted harmonic mean between observed/expected
ratio and -log(FDR).
511
'Orphan' gene identification
We identified genes from *M. abdita*'s annotation file for which EggNOG-mapper could not assign
an ortholog. Next, we assessed the expression of these genes in our RNA-seq data. To identify
genes with exclusively embryonic expression, we focused on those with a CPM > 1 in at least
one of the nine embryonic stages and a CPM < 1 in pupal, larval, and adult stages. We
validated these genes further by analyzing their ORFs using NCBI's 'Open reading frame
finder'. We used CPC2 to evaluate the nucleic acid sequences to assess coding potential (Kang
et al., 2017). Finally, we performed sequence similarity searches using NCBI's blastn and blastp
tools, querying both nucleotide and protein sequences against the entire NCBI database as well
as against Dipteran-specific sequences (Altschul et al., 1990; Sayer et al., 2022).
522
Protein structure prediction and structural similarity search

We used Protparam to assess protein stability, aliphatic index, and hydropathicity of predicted
proteins (Gasteiger et al., 2005). We then searched all predicted protein sequences against
InterProScan to look for any missed domains, specifically to look for evidence of transposable
elements that were not discovered with blast searches (Jones et al., 2014). We used the
AlphaFold2.ipynb provided by ColabFold v1.5.5 to predict protein structures (Jumper et al.,
2021; Mirdita et al., 2022;). If the predicted protein had a predicted template modeling (pTM)
score > 0.5 we then uploaded the structure to FoldSeek's website and searched the available
databases (AlphaFold/Proteome, AlphaFold/Swiss-Prot, AlphaFold/UniProt50, BFMD) for
proteins with similar structure (van Kempen et al., 2024; Varadi et al., 2022; Varadi et al., 2024).
We used Protein Imager to generate publication-quality images of protein structures (Tomasello
et al., 2020).

Acknowledgments

We thank Lily Shiue and Qianyu Jin at Cantata Bio for assembling the genome. We thank
Nicholas VanKuren for his notes on genome annotation. We thank Yaikhomba Mutum for the
conversation on protein structure prediction and structural similarity searches. We thank Xiang-
Ru (Shannon) Xu for optimization of the *M. abdita* single cross procedure and the drawings in
figure S10.

JETSTREAM2: This work used Jetstream2 at Indiana University through allocation BIO220075
from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS)
program, which is supported by National Science Foundation grants #2138259, #2138286,
#2138307, #2137603, and #2138296.

GBCF: Bioinformatic work was supported in part by the Notre Dame University Genomics and
Bioinformatics Core Facility.

Competing interests

548   The authors declare no competing interests.

Funding

550   Research reported in this publication was supported by the National Institute of General Medical
551   Sciences of the National Institutes of Health under Award Number R01GM127366 and the
552   University of Chicago. The funders had no role in study design, data collection and analysis,
553   decision to publish, or preparation of the manuscript.

Data and Resource Availability

Genome: The annotated *Megaselia abdita* genome can be found in NCBI's Genome database
under BioProject Accession PRJNA1164289.
RNASeq data: All fastq files containing the raw sequencing reads for each sample have been
uploaded to NCBI's Sequence Read Archive (SRA) and can be found under the BioProject
Accession PRJNA1200075. Individual BioSample and SRA accession numbers can be found in

supplementary table S4 including those samples that were not generated in this study but used
as evidence for the annotation of the genome.
Genome Browser: Jetstream2 at Indiana University is a resource provider for NSF's ACCESS
program which aims to broaden access to super computing resources at no cost to researchers.
To access the *Megaselia abdita* genome browser and related tools, create an ACCESS ID, then
use this ID create an account and login to Jetstream2. You will apply for an 'allocation' of credits
which can be used on Jetstream2. Detailed instructions on the use of Jetstream2 can be found
at https://jetstream-cloud.org/get-started/index.html
568
References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local
alignment search tool. *Journal of Molecular Biology* 215, 403–410.

Beeber, D. and Chain, F. J. (2020). crispRdesignR: A Versatile Guide RNA Design
Package in R for CRISPR/Cas9 Applications. *J Genomics* **8**, 62–70.

Boerner, T. J., Deems, S., Furlani, T. R., Knuth, S. L. and Towns, J. (2023). ACCESS:
Advancing Innovation: NSF's Advanced Cyberinfrastructure Coordination Ecosystem:
Services & Support. In *Practice and Experience in Advanced Research Computing*
2023: Computing for the Common Good, pp. 173–176. New York, NY, USA:
Association for Computing Machinery.

Brooks, E. M., Sanders, S. A. and Pfrender, M. E. (2024). freeCount: A Coding Free
Framework for Guided Count Data Visualization and Analysis. In *Practice and*
Experience in Advanced Research Computing 2024: Human Powered Computing, pp.
1–4. New York, NY, USA: Association for Computing Machinery.

Brůna, T., Lomsadze, A. and Borodovsky, M. (2020). GeneMark-EP+: eukaryotic gene
prediction with self-training in the space of genes and proteins. *NAR Genomics and*
Bioinformatics **2**, lqaa026.

Bullock, S. L., Stauber, M., Prell, A., Hughes, J. R., Ish-Horowicz, D. and Schmidt-Ott,
U. (2004). Differential cytoplasmic mRNA localisation adjusts pair-rule transcription
factor activity to cytoarchitecture in dipteran evolution. *Development* **131**, 4251–4261.

Campos-Ortega, J. A. and Hartenstein, V. (1997). *The Embryonic Development of*
Drosophila melanogaster. Berlin, Heidelberg: Springer.

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. and Huerta-Cepas, J.
(2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and

Domain Prediction at the Metagenomic Scale. *Molecular Biology and Evolution* **38**,
5825–5829.

Caroti, F., Urbansky, S., Wosch, M. and Lemke, S. (2015). Germ line transformation and
in vivo labeling of nuclei in Diptera: report on *Megaselia abdita* (Phoridae) and
Chironomus riparius (Chironomidae). *Development Genes and Evolution* **225**, 179.

Caroti, F., González Avalos, E., Noeske, V., González Avalos, P., Kromm, D., Wosch,
M., Schütz, L., Hufnagel, L. and Lemke, S. (2018). Decoupling from yolk sac is
required for extraembryonic tissue spreading in the scuttle fly *Megaselia abdita*. *eLife*
7, e34616.

Challis, R., Richards, E., Rajan, J., Cochrane, G. and Blaxter, M. (2020). BlobToolKit –
Interactive Quality Assessment of Genome Assemblies. *G3: Genes|Genomes|Genetics*
10, 1361.

Chen, S., Zhang, Y. E. and Long, M. (2010). New Genes in Drosophila Quickly Become
Essential. *Science* **330**, 1682–1685.

Chen, S., Krinsky, B. H. and Long, M. (2013). New genes as drivers of phenotypic
evolution. *Nat Rev Genet* **14**, 645–660.

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. and Li, H. (2021). Haplotype-resolved
de novo assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170–
175.

Crombach, A. and Jaeger, J. (2012). Life's attractors : understanding developmental
systems through reverse engineering and in silico evolution. *Adv Exp Med Biol* **751**,
93–119.

Crombach, A., Wotton, K. R., Jiménez-Guri, E. and Jaeger, J. (2016). Gap Gene
Regulatory Dynamics Evolve along a Genotype Network. *Molecular Biology and*
Evolution **33**, 1293–1307.

Dainat J. 2022. Another Gtf/Gff Analysis Toolkit (AGAT): Resolve interoperability issues
and accomplish more with your annotations. Plant and Animal Genome XXIX
Conference. https://github.com/NBISweden/AGAT.

Das, T., Purkayastha-Mukherjee, C., D'Angelo, J. and Weir, M. (2002). A conserved F-
box gene with unusual transcript localization. Dev Genes Evol 212, 134–140.

Diesh, C., Stevens, G. J., Xie, P., De Jesus Martinez, T., Hershberg, E. A., Leung, A.,
Guo, E., Dider, S., Zhang, J., Bridge, C., et al. (2023). JBrowse 2: a modular genome
browser with views of synteny and structural variation. *Genome Biology* **24**, 74.

Dey, B., Kaul, V., Kale, G., Scorcelletti, M., Takeda, M., Wang, Y.-C. and Lemke, S.
(2023). Divergent evolutionary strategies preempt tissue collision in fly gastrulation.
2023.10.09.561568.

Disney, R. H. L. (1994). *Scuttle Flies: The Phoridae*. Dordrecht: Springer Netherlands.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M. and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15–21.

Feng, D., Li, J. and Liu, G. (2020). The complete mitochondrial genomes of two scuttle
flies, *Megaselia spiracularis* and *Dohrniphora cornuta* (Diptera: Phoridae).
Mitochondrial DNA B Resour **5**, 1208–1209.

Fraire-Zamora, J. J., Jaeger, J. and Solon, J. (2018). Two consecutive microtubule-
based epithelial seaming events mediate dorsal closure in the scuttle fly *Megaselia*
abdita. *eLife* **7**, e33807.

Gabriel, L., Brůna, T., Hoff, K. J., Ebel, M., Lomsadze, A., Borodovsky, M. and Stanke,
M. (2023). BRAKER3: Fully automated genome annotation using RNA-Seq and protein
evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. *bioRxiv*
2023.06.10.544449.

Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M., Appel, R. and
Bairoch, A. (2005). Protein Identification and Analysis Tools on the Expasy Server. In
The Proteomics Protocols Handbook, pp. 571–607.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I.,
Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length
transcriptome assembly from RNA-Seq data without a reference genome. *Nat*
Biotechnol **29**, 644–652.

Grimaldi, D. and Engel, M. (2005). *Evolution of Insects*. Cambridge University Press.

Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014)**.** Circlize implements and
enhances circular visualization in R. *Bioinformatics*, **30**, 2811-2812**.**

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y. and Durbin, R. (2020).
Identifying and removing haplotypic duplication in primary genome assemblies.
Bioinformatics **36**, 2896–2898.

Gurevich, A., Saveliev, V., Vyahhi, N. and Tesler, G. (2013). QUAST: quality assessment
tool for genome assemblies. *Bioinformatics* **29**, 1072–1075.

Haas, B. J., Delcher, A. L., Mount, S. M., Wortman, J. R., Smith Jr, R. K., Hannick, L. I.,
Maiti, R., Ronning, C. M., Rusch, D. B., Town, C. D., et al. (2003). Improving the
Arabidopsis genome annotation using maximal transcript alignment assemblies.
Nucleic Acids Research **31**, 5654–5666.

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell,
C. R. and Wortman, J. R. (2008). Automated eukaryotic gene structure annotation
using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome*
Biology **9**, R7.

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J.,
Couger, M. B., Eccles, D., Li, B., Lieber, M., et al. (2013). De novo transcript
sequence reconstruction from RNA-Seq: reference generation and analysis with
Trinity. *Nat Protoc* **8**, 10.1038/nprot.2013.084.

Hancock, D. Y., Fischer, J., Lowe, J. M., Snapp-Childs, W., Pierce, M., Marru, S.,
Coulter, J. E., Vaughn, M., Beck, B., Merchant, N., et al. (2021). Jetstream2:
Accelerating cloud computing via Jetstream. In *Practice and Experience in Advanced*
Research Computing 2021: Evolution Across All Dimensions, pp. 1–8. New York, NY,
USA: Association for Computing Machinery.

Horn, T., Hilbrant, M. and Panfilio, K. A. (2015). Evolution of epithelial morphogenesis:
phenotypic integration across multiple levels of biological organization. *Frontiers in*
Genetics **6**, 303.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K.,
Cook, H., Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., et al. (2019). eggNOG
5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. *Nucleic Acids Res* **47**, D309–D314.

Jiménez-Guri, E., Huerta-Cepas, J., Cozzuto, L., Wotton, K. R., Kang, H.,
Himmelbauer, H., Roma, G., Gabaldón, T. and Jaeger, J. (2013). Comparative
transcriptomics of early dipteran development. *BMC Genomics* **14**, 123.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., et al. (2014). InterProScan 5: genome-scale
protein function classification. *Bioinformatics* **30**, 1236–1240.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589.

Kalinka, A. T., Varga, K. M., Gerrard, D. T., Preibisch, S., Corcoran, D. L., Jarrells, J.,
Ohler, U., Bergman, C. M. and Tomancak, P. (2010). Gene expression divergence
recapitulates the developmental hourglass model. *Nature* **468**, 811–814.

Kang, Y.-J., Yang, D.-C., Kong, L., Hou, M., Meng, Y.-Q., Wei, L. and Gao, G. (2017).
CPC2: a fast and accurate coding potential calculator based on sequence intrinsic
features. *Nucleic Acids Research* **45**, W12–W16.

Kimura, M. and Ohta, T. (1974). On Some Principles Governing Molecular Evolution.
Proceedings of the National Academy of Sciences of the United States of America **71**,
2848.

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C. and
Salzberg, S. L. (2004). Versatile and open software for comparing large genomes.
Genome Biol **5**, R12.

Kwan, C. W., Gavin-Smyth, J., Ferguson, E. L. and Schmidt-Ott, U. (2016). Functional
evolution of a morphogenetic gradient. *eLife* **5**, e20894.

Kwan, C.W. (2017). Functional evolution of a morphogenetic gradient. [Doctoral
dissertation,The University of Chicago]. https://doi.org/10.6082/M1H41PH7

Li, H. and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–
Wheeler transform. *Bioinformatics* **25**, 1754–1760.

Li, X., Hash, J. M., Hartop, E., Yang, D., Smith, P. T. and Brown, B. V. (2024) A
molecular phylogeny of scuttle flies (Diptera: Phoridae) unveils extensive concordance
but intriguing divergences from morphological results. *Systematic Entomology*.

Liao, Y., Smyth, G. K. and Shi, W. (2013). The Subread aligner: fast, accurate and
scalable read mapping by seed-and-vote. *Nucleic Acids Research* **41**, e108.

Liu, Q., Onal, P., Datta, R. R., Rogers, J. M., Schmidt-Ott, U., Bulyk, M. L., Small, S.
and Thornton, J. W. (2018). Ancient mechanisms for the evolution of the bicoid
homeodomain's function in fly development. *eLife* **7**, e34594.

Lomsadze, A., Burns, P. D. and Borodovsky, M. (2014). Integration of mapped RNA-Seq
reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res*
42, e119.

Lott, S. E., Villalta, J. E., Zhou, Q., Bachtrog, D. and Eisen, M. B. (2014). Sex-Specific
Embryonic Gene Expression in Species with Newly Evolved Sex Chromosomes. *PLOS*
Genetics **10**, e1004159.

Lynch, M. (2007a). The frailty of adaptive hypotheses for the origins of organismal
complexity. *Proceedings of the National Academy of Sciences* **104**, 8597–8604.

Lynch, M. (2007b). *The origins of genome architecture*. Sunderland, MA: Sinauer
Associates, Inc.

Majoros, W. H., Pertea, M. and Salzberg, S. L. (2004). TigrScan and GlimmerHMM: two
open source ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–2879.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. and Zdobnov, E. M. (2021).
BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol*
Biol Evol **38**, 4647–4654.

Marçais, G. and Kingsford, C. (2011). A fast, lock-free approach for efficient parallel
counting of occurrences of k-mers. *Bioinformatics* **27**, 764–770.

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L. and Zimin, A.
(2018). MUMmer4: A fast and versatile genome alignment system. *PLOS*
Computational Biology **14**, e1005944.

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. and Steinegger, M.
(2022). ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682.

Mulhair, P. O. and Holland, P. W. H. (2024). Evolution of the insect Hox gene cluster:
Comparative analysis across 243 species. *Seminars in Cell & Developmental Biology*
152–153, 4–15.

Öztürk-Çolak, A., Marygold, S. J., Antonazzo, G., Attrill, H., Goutte-Gattat, D., Jenkins,
V. K., Matthews, B. B., Millburn, G., dos Santos, G., Tabone, C. J., et al. (2024).
FlyBase: updates to the Drosophila genes and genomes database. *Genetics* **227**,
iyad211.

Palmer, J. and Stajich, J. (2019). nextgenusfs/funannotate: funannotate v1.5.3.

Pantano, L. (2024). *DEGreport: Report of DEG analysis*. R package version 1.42.0,
http://lpantano.github.io/DEGreport/.

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T. and Salzberg,
S. L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-
seq reads. *Nat Biotechnol* **33**, 290–295.

Picard, C. J., Johnston, J. S. and Tarone, A. M. (2012). Genome Sizes of Forensically
Relevant Diptera. *Journal of Medical Entomology* **49**, 192–197.

Posit team. (2024). RStudio: Integrated development environment for R. Posit Software,
PBC, Boston, MA. <http://www.posit.co/>

Priyam, A., Woodcroft, B. J., Rai, V., Moghul, I., Munagala, A., Ter, F., Chowdhary, H.,
Pieniak, I., Maynard, L. J., Gibbins, M. A., et al. (2019). Sequenceserver: A Modern
Graphical User Interface for Custom BLAST Databases. *Mol Biol Evol* **36**, 2922–2924.

Putnam, N. H., O'Connell, B. L., Stites, J. C., Rice, B. J., Blanchette, M., Calef, R.,
Troll, C. J., Fields, A., Hartley, P. D., Sugnet, C. W., et al. (2016). Chromosome-
scale shotgun assembly using an in vitro method for long-range linkage. *Genome Res*
26, 342–350.

R Core Team. (2023) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rafiqi, Ab. M., Lemke, S., Ferguson, S., Stauber, M. and Schmidt-Ott, U. (2008).
Evolutionary origin of the amnioserosa in cyclorrhaphan flies correlates with spatial and
temporal expression changes of zen. *Proceedings of the National Academy of*
Sciences **105**, 234–239.

Rafiqi, A. M., Lemke, S. and Schmidt-Ott, U. (2010). Postgastrular zen expression is
required to develop distinct amniotic and serosal epithelia in the scuttle fly *Megaselia*.
Dev Biol **341**, 282–290.

Rafiqi, A. M., Lemke, S. and Schmidt-Ott, U. (2011). The scuttle fly *Megaselia abdita*
(Phoridae): a link between Drosophila and Mosquito development. *Cold Spring Harb*
Protoc **2011**, pdb.emo143.

Rafiqi, Ab. M., Park, C.-H., Kwan, C. W., Lemke, S. and Schmidt-Ott, U. (2012). BMP-
dependent serosa and amnion specification in the scuttle fly *Megaselia abdita*.
Development **139**, 3373–3382.

Raj, L., Vivekanand, P., Das, T. K., Badam, E., Fernandes, M., Jr, R. L. F., Brent, R.,
Appel, L. F., Hanes, S. D. and Weir, M. (2000). Targeted localized degradation of
Paired protein in Drosophila development. *Current Biology* **10**, 1265–1272.

Rasmussen, D. A. and Noor, M. A. (2009). What can you do with 0.1× genome coverage?
A case study based on a genome survey of the scuttle fly *Megaselia scalaris*
(Phoridae). *BMC Genomics* **10**, 382.

Robinson, M. D., McCarthy, D. J. and Smyth, G. K. (2009). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139.

Rohr, K. B., Tautz, D. and Sander, K. (1999). Segmentation gene expression in the
mothmidge Clogmia albipunctata (Diptera, Psychodidae) and other primitive dipterans.
Dev Gene Evol **209**, 145–154.

Sayers, E. W., Bolton, E. E., Brister, J. R., Canese, K., Chan, J., Comeau, D. C.,
Connor, R., Funk, K., Kelly, C., Kim, S., et al. (2021). Database resources of the
National Center for Biotechnology Information. *Nucleic Acids Res* **50**, D20–D26.

Schmidt-Ott, U., Sander, K. and Technau, G. M. (1994). Expression of engrailed in
embryos of a beetle and five dipteran species with special reference to the terminal
regions. *Rouxs Arch Dev Biol* **203**, 298–303.

Schmidt-Ott, U. and Lynch, J. A. (2016). Emerging developmental genetic model systems
in holometabolous insects. *Curr Opin Genet Dev* **39**, 116–128.

Schmidt-Ott, U. and Kwan, C. W. (2022). How two extraembryonic epithelia became one:
serosa and amnion features and functions of Drosophila's amnioserosa. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **377**, 20210265.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. and Zdobnov, E. M.
(2015). BUSCO: assessing genome assembly and annotation completeness with
single-copy orthologs. *Bioinformatics* **31**, 3210–3212.

Slater, G. S. C. and Birney, E. (2005). Automated generation of heuristics for biological
sequence comparison. *BMC Bioinformatics* **6**, 31.

Smit, A. and Hubley, R. (2008). RepeatModeler.

Smit, A., Hubley, R. and Green, P. (2013). RepeatMasker.

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S. and Morgenstern, B. (2006).
AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Research* **34**,
W435–W439.

Stauber, M., Jäckle, H. and Schmidt-Ott, U. (1999). The anterior determinant bicoid of
Drosophila is a derived Hox class 3 gene. *Proceedings of the National Academy of*
Sciences of the United States of America **96**, 3786.

Stauber, M., Taubert, H. and Schmidt-Ott, U. (2000). Function of bicoid and hunchback
homologs in the basal cyclorrhaphan fly *Megaselia* (Phoridae). *Proceedings of the*
National Academy of Sciences **97**, 10844–10849.

Stauber, M., Prell, A. and Schmidt-Ott, U. (2002). A single Hox3 gene with composite
bicoid and zerknüllt expression characteristics in non-Cyclorrhaphan flies. *Proceedings*
of the National Academy of Sciences **99**, 274–279.

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable,
A. L., Fang, T., Doncheva, N. T., Pyysalo, S., et al. (2023). The STRING database in
2023: protein-protein association networks and functional enrichment analyses for any
sequenced genome of interest. *Nucleic Acids Res* **51**, D638–D646.

Tanaka, K., Diekmann, Y., Hazbun, A., Hijazi, A., Vreede, B., Roch, F. and Sucena, É.
(2015). Multispecies Analysis of Expression Pattern Diversification in the Recently
Expanded Insect Ly6 Gene Family. *Mol Biol Evol* **32**, 1730–1747.

Tomasello, G., Armenia, I. and Molla, G. (2020). The Protein Imager: a full-featured
online molecular viewer interface with server-side HQ-rendering capabilities.
Bioinformatics **36**, 2909–2911.

The UniProt Consortium (2023). UniProt: the Universal Protein Knowledgebase in 2023.
Nucleic Acids Research **51**, D523–D531.

Vakirlis, N., Carvunis, A.-R. and McLysaght, A. (2020). Synteny-based analyses indicate
that sequence divergence is not the main source of orphan genes. *Elife* **9**, e53500.

van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L. M.,
Söding, J. and Steinegger, M. (2024). Fast and accurate protein structure search with
Foldseek. *Nat Biotechnol* **42**, 243–246.

VanKuren, N. (2023)**.** Draft *Papilio alphenor* assembly and annotation. Dryad.
https://doi.org/10.5061/dryad.n2z34tn2x

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan,
D., Stroe, O., Wood, G., Laydon, A., et al. (2022). AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space
with high-accuracy models. *Nucleic Acids Research* **50**, D439–D444.

Varadi, M., Bertoni, D., Magana, P., Paramval, U., Pidruchna, I., Radhakrishnan, M.,
Tsenkov, M., Nair, S., Mirdita, M., Yeo, J., et al. (2024). AlphaFold Protein Structure
Database in 2024: providing structure coverage for over 214 million protein sequences.
Nucleic Acids Research **52**, D368–D375.

Vicoso, B. and Bachtrog, D. (2015). Numerous transitions of sex chromosomes in
Diptera. *PLoS Biol* **13**, e1002078.

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski,
J. and Schatz, M. C. (2017). GenomeScope: fast reference-free genome profiling from
short reads. *Bioinformatics* **33**, 2202–2204.

Wang, Y., Tang, H., DeBarry, J. D., Tan, X., Li, J., Wang, X., Lee, T., Jin, H., Marler, B.,
Guo, H., et al. (2012). MCScanX: a toolkit for detection and evolutionary analysis of
gene synteny and collinearity. *Nucleic Acids Research* **40**, e49.

Wickham H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
York. ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.

Wickham. H., François, R., Henry, L., Müller, K., Vaughan, D. (2023). *dplyr: A Grammar*
of Data Manipulation. R package version 1.1.4, https://github.com/tidyverse/dplyr,
https://dplyr.tidyverse.org.

Wickham, H., Vaughan. D., and Girlich, M. (2024). *tidyr: Tidy Messy Data*. R package
version 1.3.1, https://github.com/tidyverse/tidyr, https://tidyr.tidyverse.org.

Wiegmann, B. M., Trautwein, M. D., Winkler, I. S., Barr, N. B., Kim, J.-W., Lambkin, C.,
Bertone, M. A., Cassel, B. K., Bayless, K. M., Heimberg, A. M., et al. (2011).
Episodic radiations in the fly tree of life. *Proceedings of the National Academy of*
Sciences **108**, 5690–5695.

Wotton, K. R. (2014). Heterochronic shifts in germband movements contribute to the rapid
embryonic development of the coffin fly *Megaselia scalaris*. *Arthropod Struct Dev* **43**,
589–594.

Wotton, K. R., Jiménez-Guri, E., Matheu, B. G. and Jaeger, J. (2014). A Staging
Scheme for the Development of the Scuttle Fly *Megaselia abdita*. *PLOS ONE* **9**,
e84421.

Wotton, K. R., Jiménez-Guri, E., Crombach, A., Janssens, H., Alcaine-Colet, A.,
Lemke, S., Schmidt-Ott, U. and Jaeger, J. (2015a). Quantitative system drift
compensates for altered maternal inputs to the gap gene network of the scuttle fly
Megaselia abdita. *eLife* **4**, e04785.

Wotton, K. R., Jiménez-Guri, E., Crombach, A., Cicin-Sain, D. and Jaeger, J. (2015b).
High-resolution gene expression data from blastoderm embryos of the scuttle fly
Megaselia abdita. *Sci Data* **2**, 150005.

Wotton, K. R., Jiménez-Guri, E. and Jaeger, J. (2015c). Maternal Co-ordinate Gene
Regulation and Axis Polarity in the Scuttle Fly *Megaselia abdita*. *PLOS Genetics* **11**,
e1005042.

Xia, S., Chen, J., Arsala, D., Emerson, J. and Long, M. (In press). The origin of new
genes is a general evolutionary process of functional innovation. *Nature Genetics*.

Xiao, Z. and Lam, H.-M. (2022). ShinySyn: a Shiny/R application for the interactive
visualization and integration of macro- and micro-synteny data. *Bioinformatics* **38**,
4406–4408.

Yoder, J. H. and Carroll, S. B. (2006). The evolution of abdominal reduction and the
recent origin of distinct Abdominal-B transcript classes in Diptera. *Evol Dev* **8**, 241–
251.

Zhong, M., Wang, X., Liu, Q., Luo, B., Wu, C. and Wen, J. (2016). The complete
mitochondrial genome of the scuttle fly, *Megaselia scalaris* (Diptera: Phoridae).
Mitochondrial DNA A DNA Mapp Seq Anal **27**, 182–184.

Figures and Tables

Table 1. *M. abdita* reference assembly statistics and quality metrics.

Reference Assembly Statistics	
Total length (bp)	592,824,975
Number of scaffolds	15
Scaffold N50 (bp)	212,802,314
Scaffold L50	2
# of Ns per 100kb	1.32
GC (%)	29.74%
Masked (%)	67.90%
Eukaroyotic BUSCO's recovered (%)	99%

Figure 1. Genome annotation pipeline. The starting input files are shown in black ovals and include the genome assembly along with two lines of evidence: RNAseq reads (fastq format) and protein sequences obtained from NCBI and UniProt. Software tools are represented in colored boxes: green indicates mapping software, blue indicates gene model generation

software, and purple indicates post-gene model processing and functional annotation tools.

Arrows pointing from a software box to unboxed text represent the output files generated by the

software. Arrows leading from unboxed text to a software box indicate input files used by the

software.

Table 2. Genome annotation statistics (left) and quality metrics (right) for *M. abdita*. The quality
metrics include the percentage of complete universal orthologs identified in the annotation
across five BUSCO datasets. The number of genes in each lineage-specific BUSCO database
is shown in parentheses.

M. abdita Genome Annotation Stasitics		Complete universal orthologs recovered in in *M. abdita* annotation (BUSCOs)	
Number of coding genes	11,934	Eukaryota (n = 255)	93.3%
Number of mrnas	20,560	Metazoa (n = 954)	90.0%
Number of mrnas with 3' & 5' UTR	17,607	Insecta (n = 1,367)	91.6%
Mean mRNAs/gene	1.7	Endopterygota (n = 2,124)	90.6%
Mean gene length (bp)	19,478	Diptera (n = 3,285)	88.0%

A)

B) ***M. abdita* HOX gene clusters**

Figure 2. A) Synteny analysis between *D. melanogaster* chromosomes and *M. abdita* scaffolds.
Lines represent groups of collinear genes, connecting their positions between *D. melanogaster*
chromosomes and *M. abdita* chromosome-sized scaffolds. Colors correspond to *D.*
melanogaster chromosomes. The ANT and UBX Hox gene complexes are highlighted by black
lines within the chromosomes. B) Visualization of *M. abdita* Hox gene clusters. The
Antennapedia (top) and Ultrabithorax (bottom) complexes are both located on Scaffold 2. Each
blue line indicates the genomic position and length of a gene.

Figure 3. A) Multidimensional scaling (MDS) plots based on the top 500 most differentially
expressed genes from single-embryo RNA-seq samples. The *M. abdita* data are shown on the
left (blue) and the *D. melanogaster* data on the right (yellow). Points represent individual
samples, and their colors correspond to groupings identified by *k*-means clustering. Arrows
indicate key developmental events: germband retraction begins and ends. B) Number of
differentially expressed genes (DEGs) between sequential developmental stages. Comparisons

(e.g., "1 vs 5") indicate the number of DEGs identified between stage 1 and stage 5. Yellow bars

represent *D. melanogaster*, and blue bars represent *M. abdita*.

Figure 4. A-C) Expression profiles of differentially expressed gene clusters during
embryogenesis and their enriched Biological Process Gene Ontology (BP GO) terms for *M.*
abdita (left - blue) and *D. melanogaster* (right - yellow). Grey lines show the expression profile of
the individual genes within the cluster and bolded lines represent the average expression profile
of the entire cluster. DEGreports generated cluster names (e.g., "Cluster 4") which are arbitrary
but retained here for continuity. Key embryonic developmental events: MZT + cellularization,
gastrulation + germband extension, and germband retraction + dorsal closure are highlighted in
yellow, blue, and red, respectively. Shared enriched BP GO terms for the boxed clusters are
listed on the right.

938
939
940 **Figure 5.** A) Expression profiles of orphan genes with expression limited to embryonic stages. E
941 indicates embryonic, L indicates larval, P indicates pupal, and A indicates adult (multiple
942 samples). Highlighted in pink and dark purple are two genes with high-confidence protein
943 structure predictions. All other 'orphan' gene expression profiles are shown in black.
944 B) AlphaFold-predicted protein structure for Scaffold_1_222469999.1571.

945 C) AlphaFold-predicted protein structure for Scaffold_3_155577901.980. Sequence similarity
946 searches suggest this gene encodes a novel F-box-LRR protein. In the structures, alpha helices
947 are highlighted in blue, and beta sheets are highlighted in yellow.