Authors:

Robert W. Burroughs
ORCID: 0000-0003-4384-3430
Department of Ecology and Evolution
Stony Brook University, Stony Brook, NY.
Robert.Burroughs@stonybrook.edu

Christopher J. Percival
ORCID: 0000-0002-8883-9737
Department of Anthropology
Stony Brook University, Stony Brook, NY.
Christopher.Percival@stonybrook.edu

Natasha S. Vitek
ORCID: 0000-0003-0587-8907
Department of Ecology and Evolution
Stony Brook University, Stony Brook, NY.
Natasha.Vitek@stonybrook.edu

Data availability statement: All mapped and analyzed proteomics data and phenotypic data analyzed, along with R-scripts and ProteoRE workflow are included as Supplemental Materials. These Supplemental Materials will be made available publicly upon publication via Dryad Digital Repository DOI: 10.5061/dryad.kh18932j0. Raw proteomics data will be made publicly available on the MassIVE database with Dataset Identifier: MSV000098600.

Funding statement: Author RWB was funded via an Institutional Research and Academic Career Development Award (IRACDA) made to Stony Brook University from the National Institute of General Medical Sciences of the National Institutes of Health [K12GM102778], and startup funding to NSV from Stony Brook University.

Experiments were paid for via startup funding to NSV from Stony Brook University

Conflict of interest disclosure: Authors have no conflicts of interest.

Ethics approval statement: Live animal experiments were conducted in accordance with guidelines and regulations from Stony Brook University Institutional Animal Care and Use Committee (IACUC). Research was conducted in accordance with plan and procedures in approved protocol SBU IACUC 2023-0014.

Patient consent statement: Not Applicable.

Permission to reproduce material from other sources: Not Applicable.

Clinical trial registration: Not Applicable.

Reduced Dietary Protein Induces Changes in the Dental Proteome

Robert W. Burroughs[1,2], Christopher J. Percival[3], Natasha S. Vitek[1]

[1] Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA.

[2]Center for Inclusive Education, Stony Brook University, Stony Brook, NY, USA.

[3]Department of Anthropology, Stony Brook University, Stony Brook, NY, USA.

Abstract

Experimental studies have demonstrated that nutritional changes during development

can result in phenotypic changes to mammalian cheek teeth. This developmental

plasticity of tooth morphology is an example of phenotypic plasticity. Because tooth

development occurs through complex interactions between manifold processes, there

are many potential mechanisms which can contribute to a tooth's norm of reaction.

Determining the identity of those mechanisms and the relative importance of each of

them is one of the main challenges to understanding phenotypic plasticity. Quantitative

proteomics combined with experimental studies allow for the identification of potential

molecular contributors to a plastic response through quantification of expressed gene

products. Here, we present the results of a quantitative proteomics analysis of mature

upper first molars in *Mus musculus* from a controlled feeding experiment. Pregnant and

nursing mothers were fed either a low-dietary protein (10%) treatment diet or control

(20%) diet. Low-dietary protein was not associated with reduced molar size or skull

length. However, expression of tooth-related proteins, immune system proteins, and

actin-based myosin proteins were significantly altered in our low-dietary protein

proteomics sample. The differential expression of immune proteins along with

systematic reduction in actin-based myosin protein expression are novel discoveries for

tooth proteomics studies. We propose that studies that aim to elucidate specific

mechanisms of molar phenotypic plasticity should prioritize investigations into the

relationships between IGF regulation and tooth development and actin-based myosin

expression and tooth development.

Research Highlights

A low-protein diet during development results in significantly altered protein expression

for odontogenetic and osteogenic proteins, immune system proteins, and actin-based

myosin proteins within *Mus musculus*, but does not alter skull length or molar size.

Introduction

Phenotypic plasticity, the differential expression of a phenotype, is often invoked as a

way that organisms respond to changing environments. Within lab and common garden

experiments, nutritional changes such as changes in the quantity of dietary protein

induce plastic changes to mammalian cheek tooth phenotypes (Patton & Brylski, 1987;

Paynter & Grainer, 1956; Shaw & Griffiths, 1963). Typically, these are changes in tooth

size, size and shape of tooth cusps, and timing of eruption (Holloway et al., 1961;

Patton & Brylski, 1987; Paynter & Grainer, 1956; Searle, 1954; Shaw & Griffiths, 1963).

This demonstrates that there is plasticity in dental development that can provide a short-

term, non-evolutionary response to changing environments (Levis & Pfennig, 2021).

However, measures of plastic phenotypic response to environmental change often do

not identify the molecular and genetic pathways underlying those changes. The use of

quantitative proteomics to quantify differences in protein expression could allow for the

identification and study of pathways that are altered by an environmental change (e.g.,

are plastic). Here, we present a study characterizing protein expression variation in a

sample of upper first molars (M1s) and phenotypic variation in lower first molars (lower

M1s) from lab mice fed a low-protein diet as part of a controlled feeding experiment.

This experiment was designed to induce developmental changes to dental traits and to

identify proteins and associated pathways that may underlie phenotypic plasticity. This

allows us to characterize those pathways most impacted by exposure to low dietary

protein during embryonic and early postnatal tooth development.

Proteomics

Proteomics is broadly the field focused on identifying, annotating, and quantifying

variation of proteins. Quantification of variation includes both protein sequence variation

and protein expression variation. Informatics approaches are applied to protein spectral

data collected via tandem liquid-chromatography mass-spectrometry (LC-MS/MS)

(Heck & Neely, 2020). Protein expression profiles are typically tissue- and

developmental stage-dependent and must be interpreted within the specific

spatiotemporal contexts in which samples were collected (Rebeaud et al., 2021).

An assemblage of proteins that is expressed within a specific tissue or structure is often

referred to as a 'proteome' (Sharma et al., 2020). For example, within the dental

proteome enamel and dentin forming proteins are found in high abundance (Sharma et

al., 2020). However, outside of dental tissues, these proteins are detected in low

abundance and only at certain developmental stages (Bansal et al., 2012; Ritchie,

2018).

If basic processes of tooth mineralization are impacted by dietary deficiency, we expect

a change in associated protein expression and a change in phenotype such as occlusal

pattern, size, and/or shape (Harjunmaa et al., 2014). It is possible that some aspects of

tooth mineralization are more easily perturbed by environmental change than others

and their associated pathways may more frequently underlie plastic changes.

Measuring the tooth proteome may allow us to identify potential candidates underlying

plastic changes in tooth phenotype.

Beyond stereotypical tooth-associated proteins, previous characterizations of the dental

proteome have suggested that teeth serve as reservoirs of more general patterns of

organismal protein expression during amelogenesis (tooth mineralization) (Froment et

al., 2021; Giovani et al., 2021; Green et al.; Sharma et al., 2020). For example, Green et

al. (2019) recovered immune system related proteins from enamel that was micro

sampled from near the enamel-dentin junction of mineralizing pig molars. These

immune proteins currently have no known function in amelogenesis. These proteins are

definitively present within enamel tissue, but they may originate in nearby salivary

glands that are in contact with enamel cell proliferation zones and regions of

mineralization at the enamel-dentin junction (Green et al., 2019; Jagr et al., 2019). If

mineralized teeth serve as an archive of broader organismal protein expression at the

time of mineralization, it is possible that tooth proteomic data could provide evidence of

more generalized responses to environmental perturbations.

This Study

To determine what pathways are altered by an impoverished diet, we conducted a

controlled feeding experiment and investigated protein expression in the cheek teeth.

The study presented here is part of a larger research program that will also investigate

the impacts of low dietary nutrition on a variety of tooth phenotypes. Pregnant and

nursing mouse dams were fed a low protein or control protein diet and effects were

measured in their offspring. Previous feeding studies on dietary protein quantity in rats

and mice suggested that a threshold of 10-12% (by weight) dietary protein reduction

from a control of 20-24% should induce phenotypically plastic changes to molar size,

skull length, and disrupt timing of molar development (Barbeito-Andrés et al., 2016

2016; Holloway et al., 1961; Miller & German, 1999; Paynter & Grainer, 1956; Paynter &

Grainer, 1961; Pucciarelli, 1980; Shaw & Griffiths, 1963).

With this context we anticipated that reduction in dietary protein would disrupt normal

tooth development and alter protein expression patterns during embryonic and early

postnatal development. We predicted a significant reduction in M1 size, consistent with

previous reports (e.g., (Holloway et al., 1961; Paynter & Grainer, 1956; Paynter &

Grainer, 1961; Shaw & Griffiths, 1963) and significant reduction in skull length based on

previous studies in rat and mouse (e.g., (Barbeito-Andrés et al., 2016; Lobe et al., 2006;

Miller & German, 1999; Pucciarelli, 1980). Previous studies suggested that halving

dietary protein would likely increase risk of infection and increase metabolic stress in

low protein mice (Giovani et al., 2021; Steward et al., 2023). Thus, we anticipated that

there might be additional proteomic signals related to stress or immune system function

if those protein signatures are preserved within dentition. Recovery of significant

differential expression of proteins because of a dietary change allowed us to identify

and rank the molecular pathways most likely to be perturbed by environmental

influences like dietary protein reduction.

Methods

Data Availability

All datasets, code, workflows, and raw phenotypic data reported are available as

Supplemental Materials on Dryad: Dryad DOI. Raw proteomics data are available on

MassIVE database by querying the following MassIVE identifier: MSV000098600.

Feeding Experiment

A breeding colony of inbred strain C57BL/6J (RRID: IMSR JAX:000664) mice was

established at the Division of Laboratory Animal Research at Stony Brook University, in

accordance with authorized IACUC protocol (SBU IACUC 2023-0014). Male and female

mice were acquired at 8 weeks of age and housed in residence to acclimate until

breeding began at 12 weeks of age. Males were placed with females for up to 72 hours

(3 day-night cycles) and females were checked daily for presence of a copulatory plug.

Once a plug was present or 72 hours had passed, the male and female were separated.

Females, regardless of plug presence, were then randomly assigned to control or

treatment diets to ensure that any developing embryos were on as consistent a diet as

possible. Mice assigned to the control group were fed a 20% raw protein diet (PicoLab

Rodent Diet 20, 5053). Mice assigned to treatment were fed a 10% raw protein diet

(Mod LabDiet 5053 with ~10% Protein Red, 5BQM). Prior to this assignment, all

specimens consumed the control diet.

If females were not pregnant, as evidenced by swollen abdomen after ~7 days post-

mating attempt, low-protein females were cycled back to the control protein (20%) diet.

They were re-acclimated to that diet for 14 days before additional mating was

attempted. To ensure that additional stress was not induced via single housing, pairs of

females were housed together during acclimation and pregnancy. Mating timings were

staggered to ensure that two females in the same cage would not give birth at the same

time. This allowed for the separation of females and their litters once they had given

birth.

Pregnant and nursing females in the treatment group were fed only the low-protein diet

for the remainder of their lifespan. Offspring were weaned at approximately ~21 days

postnatal (P21). Siblings were housed with their respective sex and fed their respective

diets until P28 to ensure full eruption of the third molars. Dams and offspring were

euthanized when the offspring were P28. Dams were not reused to avoid introducing

bias related to improvement of maternal care from first to second litter (Weber & Olsson,

2008). A total of 12 litters (7 treatment, 5 control) constituting 74 offspring (treatment

n=34, control n=30) from this feeding experiment were collected. Aside from the 74

collected offspring, there was postnatal attrition of one full litter of treatment pups (n=8),

where the mother declined to nurse the pups. The smallest pup from each of five litters

(treatment n=3; control n=2) also did not survive to weaning.

Sampling

For proteomics data we opportunistically sampled specimens from four litters of our

feeding experiment. In our sample, males were the more common sex, and thus we

selected by random, four male sibling pairs for destructive sampling. Selecting only

males also removed variation in protein expression that would be associated with

specimen sex within our small sample. The upper first molar (upper M1) was selected

for protein extraction because of its ease of extraction and because it and its lower jaw

counterpart are the first molars to develop and erupt (Figure 1A). Eight upper M1s

(treatment n=4; control n=4) were extracted from sibling pairs from four litters

immediately after euthanasia by dissecting away the maxillary gingiva and exposed

maxillary bone to reveal tooth roots. A blunt probe was used to lever the tooth out.

Excess tissue, including any root bundle, was removed with forceps. Extracted teeth

were then washed with 70% ETOH, wiped dry with a clean Kimwipe, and stored in new

cryotubes. Teeth were placed into a -80ºC freezer and maintained at -80ºC until

preparation for protein quantification.

For phenotypic data, we sampled the left lower first molar (lower M1) of specimens from

four treatment and four control litters. These litters include the four litters sampled for

proteomics. We measured the phenotype of 50 (treatment n=24; control n=26), where

24 were males and 26 were females. Three of the specimens sampled for proteomics (2

controls, 1 treatment) were excluded from the phenotypic data because they were held

out for future synchrotron scanning. Specimen data are included in Supplemental

Materials.

Phenotyping

We utilized µCT imaging to generate high-fidelity 3D surfaces of mouse specimens.

This approach allows us to utilize semi-automated phenotyping approaches. We batch

scanned specimens at the New York Institute of Technology Visualization Center on a

Bruker Skyscan 1173 at 75kV, 106µA, at a resolution of 19.938µm. Specimens were

decapitated post-euthanasia and ethafoam was inserted to separate the upper and

lower dentition. Heads were then wrapped in Kimwipes and stacked vertically in 50ml

Falcon tubes, with males and females grouped separately within the tube and one to

two litters of the same type (e.g., all control or all treatment) per tube. Litters were

separated by stacked ethafoam discs that form a visual barrier in reconstructed scan

data.

Post-processing of µCT data was done at Stony Brook University, by author RWB,

utilizing 3D Slicer (Fedorov et al., 2012)**.** Reconstructed batch scans were loaded into

Slicer and individual specimens were cropped from the batch scan volume. Minimum

thresholding was used to segment all bony elements within the scan. The Islands

segmentation tool was used to separate out the largest element within the segment, the

cranium. Once segmented, crania were converted to 3D models. It was necessary to

segment the entire tooth row, as opposed to isolate the lower M1, due to contact

between teeth. Minimum thresholding was again applied, but with increased minima

compared to bony elements to isolate the denser enamel. The Scissors tool was used

to circle the lower left tooth row and "Erase Outside" was selected to remove all

remaining elements within the segment. Finally, the Islands tool was again used to keep

only the largest element within the segment (the tooth row). Once segmented,

toothrows were converted to 3D models.

We utilized landmarks to measure phenotypic data on 3D models. To place landmarks,

we utilized Automated Landmarking through Point cloud Alignment and

Correspondence Analysis (ALPACA) implemented within the 3D Slicer SlicerMorph

extension (Porto et al., 2021; Rolfe et al., 2021). The strength of ALPACA lies in rapid

implementation and reduced user bias in landmark placement for landmark projection.

ALPACA accomplishes this with a point cloud-based deformation of a reference

landmark template and mesh to match non-reference specimens. Once the deformation

has occurred, landmarks from the reference specimen are projected onto the non-

reference specimen. Because this method utilizes a deformation-based approach it is

necessary that specimens be similar in shape (as they are in this intraspecific study of a

single tooth) and that placed landmarks be quality checked after running ALPACA. We

generated two reference templates: 1) a control cranium with two landmarks for

calculating straight-line skull length, 2) a control lower M1 with 25 landmarks

representing tooth outline/crown footprint (Figures 1B and 1C). After ALPACA was run,

a single author (RWB) checked all of ALPACAs landmark position predictions and

moved any landmarks that were not properly projected.

Analysis of landmark data was conducted in **R** with core R tools (RCoreTeam, 2023),

the **geomorph** (Adams et al., 2013) package, with visualization via the core **plot**

functions and **ggplot** package (Wickham, 2016). The landmark files and R script are in

Supplemental Materials. To test hypotheses of changes in skull-length we measured the

line between landmarks at the apex of the foramen magnum and the junction between

the maxillae (Figure 1B). A Welch's T-Test was calculated using the **t.test** function in R

to determine if there was a significant difference between treatment and control group

mean skull lengths. Because our prediction was that treatment specimens would be

smaller, we utilized a single-tail alternative hypothesis, implemented in R as "greater

than".

Crown footprint area of the left lower first molar (lower M1) was represented by 25

landmarks placed around the widest portion of the crown (Figure 1C). These landmarks

were used to estimate a convex hull that approximates the tooth crown area. We

compared mean crown area estimates between groups using Welch's T-Test, as

described for skull length.

The decision to phenotype the lower M1 while quantifying the proteome of upper M1

was based on sampling constraints. Both molars develop simultaneously, rendering

them as comparable as possible, while simultaneously accounting for the potential

reciprocal destructive sampling of each method collection approach. Previous studies

indicate that high power μCT has been shown to damage DNA and could possibly

degrade proteins (Immel et al., 2016) and our proteomics protocol required destroying

the entire upper M1. Choosing the lower molar for phenotyping also aids in future

phenotypic comparisons because lower molars are usually found in higher abundance

within paleontological samples, resulting in studies of phenotypic evolution and plasticity

often emphasizing measures of lower molars (Bell & Jass, 2011; Burroughs, 2019;

Procopio et al., 2018; Ungar, 2010). We chose an outline-based approach over

traditional straight-line measurements because crown area has been shown to be a

more sensitive tool for assessing size (Hopkins, 2018).

Protein analysis by LC-MS/MS

For protein analysis we first determined if our extraction and proposed sampling worked

by initially sampling only two treatment and two control specimens. After validating our

protocols, we sampled the remaining two treatment and two control specimens. Proteins

were isolated in 5% SDS, 100mM TEAB, 10mM DTT using a Precellys bead

homogenizer for two cycles and spun at 16,000 x G for 5 minutes, which pulverized the

tooth tissue. Supernatants were reduced at 55ºC for 30 minutes, and cysteines were

alkylated with 25mM iodoacetamide for 30 minutes at room temperature in the

dark. Samples were acidified with phosphoric acid, proteins were then precipitated with

90% methanol, 50mM TEAB, and bound to S-Trap solid phase cartridges (Zougman et

al., 2020). Protein precipitates were washed with 90% methanol, 50mM TEAB and

digested with trypsin (20μg, Sigma-Aldrich. Trypsin, TPCK-treated; #4352157) at 47ºC

for two hours. We ran two blank STRAP samples per analytical sample, no protein or

peptide was recovered from blank STRAP samples, except for a single keratin peptide

at m/z 1082. Precipitates were then sequentially eluted with 50mM TEAB, 0.2% formic

acid, and 50% acetonitrile, the 0.2% formic acid elution step was by centrifugation at

4000 x G for 1 minute each. Peptides were dried by vacuum centrifugation, desalted

using HLB SPE cartridges (Waters), dried and resuspended in 20µl 0.1% formic

acid/water.

Peptides were analyzed by C18 reverse phase LC-MS/MS (Thermo Nano1200) using a

2µl injection volume. HPLC C18 columns were prepared using a P-2000 CO2 laser

puller (Sutter Instruments) and silica tubing (100µm ID x 15 cm) and were self-packed

with 3u Reprosil resin. Peptides were separated using a flow rate of 300 nl/minute, and

a gradient elution step changing from 0.1% formic acid to 40% acetonitrile (CAN) over

90 minutes, followed by a 90% CAN wash and re-equilibration steps (buffer A 0.1%

formic acid/water, buffer B 90% acetonitrile/water with 0.1% formic acid). Parent peptide

mass and collision-induced fragment mass information were collected using an orbital

trap (Q-Exactive HF; Thermo) instrument followed by protein database searching using

Proteome Discoverer 2.4 (Thermo). Electrospray ionization was achieved using spray

voltage of ~2.3 kV. Information-dependent MS and MS/MS acquisitions were made

using a 50ms survey scan (m/z 375 – 1400) at 60,000 resolution, followed by 'top 20'

consecutive second product ion scans at 15,000 resolution. Peptide and spectra false

discovery rates were set to 0.01 and 0.05 FDR bins, with 10ppm MS and 0.05Da MS2

tolerances, allowing up to 2 missed tryptic cleavages, and a minimum peptide size of 6

amino acids. Variable modifications included M-oxidation, NQ-deamination, and ST

dehydration. Peptide-specific label free quantitation (mapping) was performed using

Proteome Discoverer 2.4 (SequestHT Percolator algorithm), linking peptides to

annotated mouse proteome (Uniprot mouse 16982 sequences; 4-2019, downloaded

October of 2023) and the MQ contaminant dataset for standardized protein

identification. Label free abundance estimations were performed (Sequest HT LFQ).

Raw data files are available on the MassIVE database using identifier MSV000098600.

Filtering

To assess if dietary changes resulted in significantly altered protein expression, we first

filtered our mapped proteomic data. Filtering was done in ProteoRE (Mehta et al.,

2023). Because spectral ionization can vary between LC-MS/MS analyses (e.g., (Muntel

et al., 2015; Z. Wang et al., 2024)) we initially treated our dataset as two distinct

datasets, representing the initial sampling (Group 1) and additional sampling (Group 2)

(Supplemental Materials). There was no expectation that Groups 1 and 2 were

substantially different from one another based on the experimental procedure: groups

differed only by the date of LC-MS/MS analysis. We chose a conservative approach of

filtering each dataset individually and concatenating the resulting filtered datasets,

ensuring that only proteins represented in both datasets were used for downstream

analyses and interpretations. Each sample group was initially filtered by excluding all

mapped proteins that had any one of the following criteria: A minimum false discovery

rate (q-value) greater than 0.05, representation by fewer than two peptides, or three or

fewer peptide spectral matches (PSMs). These criteria were selected based on

proteomics field standard practices (Al-Amrani et al., 2021).

Mapped Proteomics Analysis

To concatenate our results, a Venn analysis was performed to find the set of proteins

identified as unambiguously present in both groups, resulting in a combined group (CG)

dataset. This dataset reflected the shared mapped proteins between all eight (four

control, four treatment) specimens in our sample. Groups 1 and 2 were further filtered to

identify proteins with significant changes in protein expression between treatment and

control specimens. First, proteins where at least one treatment specimen in each group

had an abundance ratio adjusted p-value ≤0.05 were retained (n=135). Then, proteins

that had missing data for two or more specimen comparisons were discarded (n=15).

Those two datasets were subsequently concatenated via a Venn analysis to form a

combined group significant (CGSig) dataset. The result is that every protein in CGSig is

significantly differentially expressed in at least one treatment/control comparison in

Group 1 and at least one treatment/control comparison in Group 2.

Here, the abundance ratio is a normalized ratio of estimated protein abundance, based

on peak intensity, for one proteomics sample over another proteomics sample, in our

case a treatment specimen over a control specimen. Because protein abundances can

vary significantly in terms of magnitude between samples in label free quantification, the

estimated protein abundances are commonly transformed on a Log2 scale to normalize

(adjust) them (Liu & Zhang, 2021). To calculate the p-value for adjusted abundance

ratios p-values we used a two-tailed Students' T-Test. One result of this transformation

is that some calculated abundance ratios will effectively become -∞ or ∞ but are

represented in the dataset as values of -3.32 or 3.32. We calculated the average Log2

fold change reported in Table 1 by taking the mean of the estimated abundance ratios

from samples that had significant p-values, excluding significant samples that were

equal to -3.32 or 3.32. For some samples, calculated means were close to 0 and/or

removal of one outlier could change the sign (and thus interpretation) of the mean. In

those cases, we do not provide an average value, instead we provided the number of

specimens and their direction of fold change (Up/Down) (Table 1).

To identify protein functions, associated interactions, and general biological profiles

represented in CG and CGSig we performed Gene Ontology (GO) enrichment analysis

via the ClusterProfiler tool of ProteoRE and Pathway Enrichment Analysis via

REACTOME (Croft et al., 2011; Wu et al., 2021). For GO enrichment analyses, we

queried at two ontology levels, using cutoffs for p-value of 0.05 and q-value of 0.05.

Outputs for GO analyses were used to assign broad categorical function to proteins

(Table 1) based on Metabolic Functions, Cellular Component, and Biological Processes

categories (Figure 3).

For pathway enrichment analysis via REACTOME we queried the *Mus musculus*

REACTOME for the proteins found in the CGSig dataset. To calculate enrichment, the

number of entities (in our case proteins) identified as belonging within a specific

pathway are identified. Then the total number of entities (proteins) that could be

contained in that pathway is calculated and divided by the total number of entities

(proteins) from the organism (in our case *Mus*). The resulting 'Entity Ratio' is used to

correct for pathway size to determine which pathways are overrepresented compared to

a random distribution. A pathway is considered 'enriched' when the input from the

dataset (x) is a higher proportion of the total number of entities (n) than would be

expected by chance. The resulting probability estimate is represented by a p-value

calculated on a 95% confidence interval and pathways with Entities p-value ≤ 0.05 are

significantly enriched. Enriched pathways were then ranked based on the Entity Ratio,

which represents the percentage of all *Mus* proteins in that pathway with a significant

enrichment in our analysis (i.e., an Entity Ratio of 0.04 represents 4% of all *Mus*

proteins known in that pathway). Pathways with higher entity ratios are ranked higher

within Table 2.

Preliminary Test of Developmental Archive

Prior research and the design of our study allowed us to conduct a preliminary

investigation of whether enamel and dentin proteomes represent an archive of protein

expression during mineralization, rather than proteins expressed earlier in tooth

development or at the time of specimen collection. We queried the Mouse Gene

eXpression Database (GXD) to determine if a subset of proteins was expressed at

developmental stages prior to the onset of mineralization, during mineralization, or after

mineralization was complete. First, we investigated proteins known to be expressed

during tooth mineralization (Pandya et al., 2017). The study of Pandya et al. (2017)

reported 24 proteins present during tooth mineralization, we identified 20 of those 24

proteins in our CG dataset. We also investigated proteins related to

odontogenesis/osteogenesis, immune, and actin-based myosins from our CGSig

dataset. A protein's associated gene being expressed during mineralization, but not

earlier or later in time, would support the argument that tooth proteomes represent a

specific window of development during amelogenesis. The table containing the proteins

queried in the GXD is within Supplemental Materials.

Results

Phenotyping

We found no significant differences in size between treatment and control groups

(Table 3). Mean skull length for the treatment group was 19.20mm and control group

was 19.03mm (Figure 2A). Mean crown area for treatment group was 1.03mm^2 and

control group mean was 1.02mm^2 (Figure 2B).

Proteomics

Within G1, a total of 2189 unique proteins were mapped (Supplemental Materials). 1622

unique proteins were mapped for G2 (Supplemental Materials). The combined group

(CG) of proteins shared between G1 and G2 was 1469 unique proteins (Supplemental

Materials). Of the combined group, there were a total of 120 proteins with significant

differential expression (fold change) (CGSig; Table 1; Figure 4; Supplemental Materials

Figures 1-6).

Gene ontology profiling revealed that both CG and CGSig are comprised of sets of

proteins primarily associated with the Binding (in Molecular Function), the Cell generally

(as opposed to a specific cellular component), and the Biological Process of Metabolism

(Figure 3; Supplemental Materials). Pathway enrichment analysis via REACTOME

identified 387 biological pathways associated with significantly differentially expressed

proteins (Supplemental Materials). Of those 387, 34 were significantly enriched ($p \leq$

0.05) (Table 2). Approximately 47% of matched proteins are represented in the top 6

enriched pathways (Entity Ratio ≥ 0.01) (Table 2; Supplemental Materials).

Odontogenesis and Osteogenesis Proteins

A total of 8 proteins in CG were identified as belonging to biomineralization seven

proteins associated of them had significant differential expression (Log2 fold change)

between treatment and control groups (Table 1; Figure 4). In low-protein treatments, the

major enamel-forming protein, Amelogenin X (AMELX) had an average -1.52-fold

change in expression. One of the two major dentin-forming proteins, Dentin

sialophosphoprotein (DSPP) had a 1.77-fold increase in expression. The other dentin-

forming protein, Dentin matrix acidic phosphoprotein 1 (DMP1) did not have a clear

direction of differential expression; three treatment comparisons showed an increase in

protein expression, while two had a decrease in expression. For osteogenesis, Integrin

alpha-V (ITGAV) had -2.66-fold change and Tartrate-resistant acid phosphatase type 5

(ACP5) had -1.71-fold change. The remaining two osteogenic proteins Collagen alpha-

1(II) chain (COL2A1) and Galectin-7 (LGALS7) had variation that made interpretation

unclear. For COL2A1, one specimen had increased and two had decreased expression.

For LGALS7, three were increased and two decreased.

Inflammation and Immune Response

A total of 229 proteins were associated with Immune System Response. Nine proteins

associated with inflammation and immune system response had significant differential

expression (Log2 fold change) between treatment and control groups (Table 1; Figure

4). Two of the nine proteins, S100-A8 (S100A8) and Coagulation factor X (F10), had a -

1.27 and -0.23 fold change in treatment groups, respectively. Five of the nine proteins

had increases in expression: 0.92 for Platelet factor 4 (PF4), 1.10 for Galectin-3

(LGALS3), 1.50 for BPI fold-containing family A member 2 (BPIFA2), 2.30 for Protein

AMBP (AMBP), and 2.40 for Arginase-1 (ARG1). Two of the immune proteins,

Apoptosis-associated speck-like protein containing a CARD (PYCARD) and

Calmodulin-4 (CALM4), lacked consistent signal between pairwise specimen

comparisons. For PYCARD, one specimen showed increased expression and one

showed decreased expression. For CALM4, two specimens were increased in

expression and one decreased in expression.

Muscle Contraction

A total of 42 proteins were associated with muscle contraction 13 of those proteins had

significant differential expression. All of them had decreased expression for treatment

specimens (Table 1; Figure 4). Seven of the 13 proteins are actin-based myosins

(Myosins 1, 4, 7B, 8, Myosin Light chain 1, Tropomyosin 2, and Myosin regulatory light

chain 2), with the remaining six being actin-specific proteins (Troponin T, Actins Alpha 1

and Alpha 2, Titin, and Four and a half LIM domains protein 1).

Preliminary Test of Developmental Archive

Genes associated with two of 20 tooth proteins previously found during mineralization

(Pandya et al., 2017) are also found in tooth tissues prior to the onset of amelogenesis

based on a query of GXD. Those genes are *Alpl* and *Itgb1*. Both are found only in the

tooth developmental stage immediately prior to amelogenesis (TS21 associated with

embryonic days 12.5-14). The genes associated with the remaining 18 proteins are

found only in developmental stages associated with molar amelogenesis (TS22+,

embryonic day 15 through postnatal day 8). Querying genes associated with our

proteomic sample's differentially expressed immune and actin-based myosin proteins

within GXD revealed that immune genes are universally expressed within tooth tissues

during all reported stages until adulthood. Actin-based myosins were present from the

onset of mineralization forward. Two of the actin-based myosins (*Actn2* & *Fhl1*) were

reported as definitively absent from tooth tissues at E14.5 (i.e., just prior to

mineralization) (Visel et al., 2004). Additionally, one gene (*Myh1*) was present in

mineralization stages but was definitively absent from postnatal week 6-8 aged mouse

specimens (Freeman et al., 1998).

Discussion

Phenotyping

Our hypotheses were that specimens in the treatment group would have reduced skull

length and M1 size, results we did not recover. Both phenotypic results were

unexpected because many studies have reported variation in skull size associated with

nutritional quality in rats, mice, and pigs (e.g., (Barbeito-Andrés et al., 2016; Dickerson

& Hughes, 1972; Lobe et al., 2006; McCance et al., 1961; Miller & German, 1999;

Pucciarelli, 1980, 1981), and four previous studies reported variation in the size of the

M1 due to dietary changes (Paynter & Grainger, 1956; Holloway et al., 1961; Paynter &

Grainger, 1961; Shaw & Griffiths, 1963). Why then did we not recover similar signals?

First, we note that these previous studies sampled specimens at different ontogenetic

stages than we did, most sampled specimens that were older than ours (40-days

postnatal to 10-months postnatal), a single study (Barbeito-Andrés et al. 2016) sampled

specimens younger than ours (embryonic day 18 to postnatal day 1). Considering skull

length, mice grow continuously until they reach sexual maturity and rapidly between

birth and weaning. It is possible that any differences in skull length were simply not

manifested at the ontogenetic stage we sampled. This does not explain, however, the

lack of difference in M1 size reported previously, because M1s are determinate in

growth and cease to grow by postnatal day 8 (Ungar, 2010; Pandya et al, 2017). In

comparison to our study design using 10% dietary protein as the low-protein treatment,

some other studies used dietary protein quantity as low as 8% (e.g., Paynter &

Grainger, 1956; Holloway et al., 1961; Paynter & Grainger, 1961; Shaw & Griffiths,

1963; Miller & German 1999). This small additional reduction in dietary protein may be

sufficient to trigger effects. Finally, we recognize that different dental and cranial traits

are impacted by dietary protein more readily than others. Cranial traits have been

reported to be less impacted by reduced dietary protein than postcranial traits (e.g.,

McCance et al. 1961; Lobe et al. 2006) and M1 size may be less impacted than other

traits of the M1 (e.g., angle of cusps) (Paynter & Grainger, 1956; Holloway et al., 1961;

Paynter & Grainger, 1961).

At present we can only conclude that there is no difference in M1 size or skull length of

specimens from our groups. However, our proteomic results (discussed in detail below)

inform us that development of the tooth was perturbed in ways that likely impact other

aspects of tooth phenotype. For instance, changes in relative enamel or dentin

thickness would not be recovered by our estimates of size, because these are internal

structural changes to tooth composition. Future more comprehensive phenotyping

efforts may reveal significant differences.

Developmental Archive

Results from GXD queries support the argument that our tooth proteome dataset

primarily represents gene expression during mineralization and not prior to

mineralization. For example, genes for actin-based myosins of interest within our

dataset are not expressed in mouse teeth prior to the onset of amelogenesis. While it is

not possible to say whether the measured immune-system proteins in our proteomic

sample were expressed during mineralization or at the time of euthanasia, previous

studies (e.g., Green et al., 2019; Jagr et al., 2019) have identified immune and

inflammation related proteins incorporated in tooth tissues, either within enamel itself or

within other tissues of the enamel-dentin junction, such as salivary glands. In our case,

we recover several of the immune related proteins reported by those studies, including

S100A8 and CALM4, potentially supporting the idea that molar enamel represents an

archive of gene and protein expression during amelogenesis. This conclusion is further

supported by the fact that the gene for differentially expressed MYH1 in our dataset is

not expressed after mineralization, indicating our MYH1 signal likely represents protein

expression during mineralization rather than a later time point (Freeman et al. 1998).

Further, we would not expect expression of amelogenesis-specific genes or proteins

after the end of mineralization, because of the cessation of proliferation of ameloblasts

and lack of vascularization within the fully mineralized tooth (Alghadeer et al., 2023;

Nanci, 2007). While this hypothesis requires further directed experimental validation,

our results support the concept that the recovered tooth proteome represents a limited

window of development.

Odontogenic Proteins

We predicted that proteins associated with enamel and dentin formation would be

altered by our feeding experiment, and specifically that they would be decreased in

expression. The major enamel forming protein, AMELX, was significantly reduced in

expression for treatment specimens (Table 1; Figure 4). We anticipated that our dietary

protein reduction would result in decreased tooth size, associated with decreased

AMELX expression, because AMELX is a necessary component for the formation of

enamel. While changes in AMELX met our expectations, measures dentin forming

protein expression and lower M1 size did not.

In the case of DMP1, there is not a clear signal to interpret whether expression was

increased or decreased in our proteomics sample. This highlights the challenges of

drawing interpretations from proteomic data where individual variation can influence the

overall interpretation. This challenge is recognized by the field of quantitative

proteomics, but still represents an area where increased research efforts will be needed

(Al-Amrani et al., 2021; Chantada-Vazquez et al., 2022; Liu & Zhang, 2021; Steward et

al., 2023). A clearer interpretation of DMP1 expression, would be helpful for constructing

future hypotheses. For instance, decreased expression of DMP1 should lead to

decreased expression of DSPP and dentin hypomineralization, suggesting that DMP1

and DSPP expression contribute significantly to dentinogenesis imperfecta (Orsini et al.,

2014; Shi et al., 2020). Being able to robustly identify such patterns or, at minimum,

make supported interpretations based on the variable evidence, will enhance the utility

of future quantitative proteomic studies.

Our finding of increased DSPP expression seemed initially counterintuitive. However,

this result is supported by a recent study of protein expression in a hypomineralized

enamel defect found in humans (Mukhtar et al., 2022). In this study of hypomineralized

molars, the enamel defect impacts the first permanent molars of children and results in

a significant reduction in mineral density from normal teeth (Mukhtar et al., 2022). This

reduced density was associated with downregulation of AMELX, upregulation of DSPP,

but no reported differences in DMP1 expression (Mukhtar et al., 2022).

The causal mechanism for the pattern of increased DSPP expression is unknown. The

general role of DSPP is to control the conversion of dental pulp cells into odontoblasts

via binding with Integrin beta 6 (ITGB6) (Ritchie, 2018; Wan et al., 2016). Previous work

indicated that mice with either a DSPP heterozygous (DSPP$^{+/-}$) or DSPP knockout

(DSPP$^{-/-}$) genotypes experience dentin dysplasia and dentinogenesis imperfecta due to

haploinsufficiency of DSPP (Shi et al., 2020). Haploinsufficiency suggests that dentin is

impacted when DSPP expression is decreased but does not indicate what phenotype

results from elevated DSPP expression. Amelogenesis imperfecta enamel is typically

thin and chalky while dentin appears to be normally mineralized, suggesting that DSPP

overexpression does not result in dentin hypermineralization, but this has not been

experimentally validated and dentin structure was not reported by Mukhtar et al. (2022).

Our proteomic results appear consistent with protein expression patterns associated

with amelogenesis imperfecta and not dentinogenesis imperfecta, based on the shared

expression changes for AMELX and DSPP (Mukhtar et al., 2022; Orsini et al., 2014; Shi

et al., 2020). It is unlikely that both amelogenesis and dentinogenesis imperfecta are

simultaneously present within a single specimen's dentition. Only a single study has

reported compounded presence of amelogenesis and dentinogenesis imperfecta, which

occurred in an MSX2 knockout transgenic line (Aioub et al., 2007).

It is possible that the signal here is due to experimental design of this study, where

enamel and dentin are pooled together. The concern is that dentin constitutes a larger

amount of the overall tooth and could 'swamp' the proteomic signal. However, this

appears unlikely due to consistent signals of expression changes for many proteins not

related to dentin. In cases where a set of proteins 'swamp' the signal, it is due to

exceptionally high abundance of certain proteins, e.g., raw blood is dominated by ~20

proteins related to hemoglobin, swamping out other protein signals (Molloy et al., 2023).

In our case, recovery of 1469 proteins and 120 significantly different proteins associated

with numerous processes suggests that our signal is not driven by one tissue or

category of proteins within the pooled tissue sample.

Thus, we conclude that our measures of differentially expressed amelogenesis-

associated proteins indicate that the development of thinner and less mineralized dental

enamel could be a phenotypically plastic response to reduced dietary protein during

early development. Such a result would help explain why there are disparate signals

between proteomic and phenotypic results: while tooth size is not affected, tooth or

microstruture composition may be. To test this hypothesis, it is necessary to utilize

synchrotron scanning or destructive sampling in the form of histology. For now, it

remains to be tested if the mouse molars from our study have amelogenesis imperfecta.

Immune and Inflammation Proteins

Previous work on mapping protein expression across micro-sampled enamel sections

of pig molars had suggested that there was possibility of recovering immune and

inflammation related proteins from mineralized dentition (Green et al., 2019). In their

case, Green et al. (2019) were constructing a detailed map of proteomic expression

associated with amelogenesis. Their reported immune system proteins were localized

from enamel which came from along the enamel-dentin junction (Green et al., 2019).

Their study was not designed nor attempted to induce differential protein expression

based on experimental procedures. In our case, we were unsure if we would recover

immune and inflammation proteins because we created a tissue-averaged signal by

crushing and processing the entire M1. By finding these proteins in our proteomics

sample and recovering differential expression of them, we present a novel result of

immune response to an environmental change. Of the nine immune or inflammation

response proteins with significant fold change, five were increased in expression for

treatments relative to controls, two were reduced in expression, and two had mixed

interpretations (Table 1).

Seven of the nine proteins are primarily associated with neutrophil degranulation,

including transport and proliferation of neutrophils. Of these seven, five were increased

in expression. However, Calmodulin 4 (CALM4) had a mixed interpretation and

coagulation factor X (F10) was reduced in expression. Neutrophils function as critical,

but specialized, immune system response cells. Neutrophils contain granules of multiple

types (azurophilic, specific, ficolin-rich, tertiary, and secretory), that target specific

threats and/or regulate immune system response to specific infectious threats

(Eichelberger & Goldman, 2020; Othman et al., 2022 ; Yin & Heit, 2018).

The precise nature of what was being targeted by immune system activation is

unknown. However, the result of F10 being reduced in expression may provide some

insight into the potential infectious threat. Deficiency of F10 has been implicated as part

of an immune response to the common, antibiotic-resistant, bacterium *Acinetobacter*

baumannii (Choby et al., 2019). In those cases, F10 deficiency is indicative of an

increased abundance of neutrophils and macrophages (Choby et al., 2019). Thus,

though F10 is decreased in expression, relative to the increase in five other neutrophil

degranulation proteins, the combination of patterns supports a conclusion that our

treatment group had higher immune system response than our control group. Future

research efforts to systematically compare immune-related proteomic signatures from

mineralized structures to standardized health monitoring tools could prove fruitful.

We also recovered changes in PYCARD and S100A8 within our proteomics sample,

that are indicative of an inflammation response (Table 1). Studies have indicated that

PYCARD directly mitigates inflammation when upregulated and contributes to

inflammation when downregulated (Sartoretto et al., 2019; Wittmann et al., 2021). In our

sample, one treatment specimen showed a moderate increase in PYCARD expression

over controls, and one showed a moderate decrease (Table 1). Importantly, deviation in

either direction suggests that there is either an increased response to inflammation

(increased expression) or increased inflammation present (decreased expression)

(Sartoretto et al., 2019; Wittmann et al., 2021). Reduced expression is recovered for

S100A8, consistent with an ongoing physiological response to inflammation (S. Wang et

al., 2018). These patterns of differential expression may represent increased

inflammation response in some of the treatment specimens, as compared to control

specimens. Due to the limited nature of our data and potential variation in inflammation

response between specimens, these results should be interpreted cautiously.

Because of our controlled experimental design, we postulate that reduced dietary

protein is the cause of increased inflammation. A recent study investigating the impact

of low dietary protein during gestation indicates that intrauterine inflammation can occur

and result in increased inflammation present in the offspring of Syrian golden hamsters

(Mohammed et al., 2023). While Mohammed et al. (2023) focused on measuring

inflammation of the liver of the offspring, the connection between low dietary protein

during embryonic and postnatal development and increased inflammation was strongly

established. Our study and results further support this connection. Future studies should

aim to systematically investigate proteomic signals of mineralized structures along with

standardized inflammation panels. This would further establish the connection between

dietary protein, inflammation, and the signals archived within mineralized structures.

Muscle Contraction Proteins

All thirteen significantly modified Muscle Contraction pathway proteins were reduced in

expression. They were predominantly actin-based myosins, which play a critical role in

cellular movement and structure formation by acting as motor molecules (Guhathakurta

et al., 2018). Actin-based myosins are broadly implicated in the proper development of

many tissues, including dentition (Du et al., 2024; Guhathakurta et al., 2018; Luis &

Schnorrer, 2021). During dental development actin-based myosins contribute to the

proper formation of enamel rods by transporting ameloblasts (Duverger & Morasso,

2018). Lower expression of actin-based myosins, including some of those that are

differentially expressed in our study (e.g., Myosin-1, Myosin-4, Myosin-8, and Tropinin 1)

are associated with the syndromic form of amelogenesis imperfecta (Duverger &

Morasso, 2018). To date, no proteomic study of dentition has recovered differential

expression of actin-based myosins, but none of the previous dental proteomic studies

attempted to experimentally induce differential expression of these proteins. Future

investigations should center on understanding the distribution of actin-based myosins

within the tooth and the correlation between protein expression in teeth and other

tissues when dietary-protein is reduced.

Potential Pathways for Phenotypic Plasticity

We recovered 34 significantly enriched REACTOME pathways from our 120 significantly

differentially expressed proteins. Among those pathways, two of the top six most

enriched pathways, regulation of insulin-like growth factor and muscle contraction, have

proteins that play many roles during morphogenesis.

Insulin-like growth factor plays a role in a number of different developmental processes,

including odontogenesis, bone development, organ development, and brain

development (Baroncelli et al., 2017; Brown et al., 2017; Chen et al., 2009; Dodington et

al., 2021; Luo et al., 2020; Montivero et al., 2021; Oyanagi et al., 2019; Vassilakos et al.,

2019). Thus, regulation of IGF is a complex set of processes with many different factors

involved in regulating expression in different tissues. This includes regulation by thyroid

growth hormone (Yakar & Isaksson, 2016) and regulation via phosphorylation of binding

proteins to aid in transport, activation, and inhibition (Chrudinova et al., 2024; Dong et

al., 2022; Huttlin et al., 2010; Palma-Lara et al., 2023; Tagliabracci et al., 2015). Given

the importance of IGF to multiple developmental processes, regulation of IGF is a

probable candidate for a mechanism underlying phenotypic plasticity. Experimental

manipulation of IGF1 and IGF2 during odontogenesis has revealed systematic changes

to the size and number of cusps of developing molars (Oyanagi et al., 2019). The

connection between IGF1 and IGF2 gene expression and proteins associated with IGF

regulation, which includes Amelogenin X, is not currently well understood (Bansal et al.,

2012; Oyanagi et al., 2019; Pandya & Diekwisch, 2021). However, regulation of insulin-

like growth factor offers a promising pathway for future investigation, particularly in

elucidating responses of IGF gene expression and associated regulatory factors to

environmental perturbations, such as poor diet.

Similarly, our collection of proteins from the muscle contraction pathway are primarily

actin-based myosins. The function of myosins as motor proteins is well documented, but

the specific nature of how myosins interact during odontogenesis is poorly understood

(Du et al., 2024; Guhathakurta et al., 2018). Previous proteomic studies of teeth have

not reported differential expression of actin-based myosins, making this result

unexpected. Simultaneously, those studies were not attempting to experimentally

induce plastic responses like our study. The connection of actin-based myosins with

body size and potentially with amelogenesis imperfecta suggests that future

investigations may be fruitful. These future studies would hopefully confirm the results

we recovered and further explore the relationship between protein expression and

phenotypic plasticity.

Conclusions

Proteomic expression results supported our prediction that halving dietary protein during

embryogenesis and early postnatal development would alter expression of proteins

recovered from mouse molars. Specifically, we identified 120 differentially expressed

proteins associated with a reduction of dietary protein during embryonic and early

postnatal development. Changes in dietary protein have been proposed to result in

phenotypically plastic changes in molar size and skull length. Our study did not recover

significant differences between normal and low protein groups for either a measure of

molar size or skull length. The discordance between positive proteomic results and

negative phenotypic results suggests that phenotypic differences may be more nuanced

that initially predicted.

We recovered significant changes in proteins associated with dental development, that

are primarily within the pathway associated with regulation of Insulin-Like Growth

Factor. The connection between IGF and dental proteins remains to be further

investigated, but changes in expression in this pathway could directly influence tooth

structure. We also identified systematically reduced expression for proteins in the

Muscle contraction pathway, specifically actin-based myosins, a novel discovery for

tooth-derived proteomics data. Actin-based myosins are broadly implicated in vertebrate

development and are correlated with dental and general body development. Taken

together, it is possible that we perturbed tooth phenotypes, such as enamel or dentin

thickness or density, that are not captured by our measures of craniodental size.

Fortunately, the study presented here is part of a larger research program aimed at

more comprehensive phenotyping efforts. The results of this study will allow us to focus

on investigating phenotypic hypotheses derived from our proteomics results (e.g.,

testing for amelogenesis imperfecta). As well as continuing to investigate other aspects

of craniodental morphology, to include assessment of size of the second and third

molars and shape of all three molars, along with cranial traits such as depth and width

of the cranium. Inclusion of these more comprehensive phenotypic data may recover

differences between groups that were not recovered from the data reported here.

Our controlled feeding experiment also induced increased immune system activation

and inflammation response, as evidenced by increased expression of proteins in the

Neutrophil degranulation pathway. While this result does not directly inform us about

phenotypic plasticity, that we could potentially to derive such information from a fully

mineralized teeth could prove useful for studying the biology of deceased or extinct

organisms. New studies aimed at microsampling enamel, dentin, and along the enamel-

dentin junction to determine both the source of these signals and which tissues contain

them are necessary.

Finally, we propose that proteomic quantification of non-experimental organisms will

prove fruitful and predict that dietary changes in wild settings will change enamel and

dentin formation by altering aspects of the IGF regulatory pathway and potentially

expression of actin-based myosins. Future research efforts then should focus on

elucidating the connection between IGF gene expression and enamel and dentin protein

expression; determining the role of actin-based myosins in tooth and skeletal

development; and comparing the immune and inflammation signals in proteomic data to

classically used tools (e.g., blood panels and cortisol screening). With a combination of

phenotyping and proteomic investigations it will become clearer precisely what traits are

more likely to be perturbed, and what sections of these pathways are causal to those

changes. In turn, this will enhance the utility of quantitative proteomics for investigating

organismal biology more broadly.

783
784 **Acknowledgments**
785
We thank the Stony Brook Division of Laboratory Animal Research (DLAR), particularly

Laurie Levine, for animal husbandry and management. We thank Dr. John Haley of the

Stony Brook University Proteomics Core Facility, for sample preparation and initial post-

processing of proteomics data. We thank the New York Institute of Technology

Visualization Center staff, Dr. Brian Beatty and Kelsi Hurdle, for providing technical

support in scanning and reconstruction of µCT scans. We thank the Stony Brook

University Center for Inclusive Education staff: Lisa Ospitale, Diana Champney, and

Erica Valdez, and Stony Brook IRACDA PIs, Dr. Carol Carter and Karian Wright, for

financial and administrative support.

Author RWB was funded via an Institutional Research and Academic Career

Development Award (IRACDA) made to Stony Brook University from the National

Institute of General Medical Sciences of the National Institutes of Health

[K12GM102778], and startup funding to NSV from Stony Brook University. The content

803
804
805
806 **Cited**
807

Adams, D. C., Otárola-Castillo, E., & Paradis, E. (2013). geomorph: anrpackage for the
collection and analysis of geometric morphometric shape data. *Methods in Ecology*
and Evolution, 4(4), 393-399. doi:10.1111/2041-210x.12035
Aioub, M., Lezot, F., Molla, M., Castaneda, B., Robert, B., Goubin, G., . . . Berdal, A. (2007).
Msx2 -/- transgenic mice develop compound amelogenesis imperfecta,
dentinogenesis imperfecta and periodental osteopetrosis. *Bone, 41*(5), 851-859.
doi:10.1016/j.bone.2007.07.023
Al-Amrani, S., Al-Jabri, Z., Al-Zaabi, A., Alshekaili, J., & Al-Khabori, M. (2021). Proteomics:
Concepts and applications in human medicine. *World J Biol Chem, 12*(5), 57-69.
doi:10.4331/wjbc.v12.i5.57
Alghadeer, A., Hanson-Drury, S., Patni, A. P., Ehnes, D. D., Zhao, Y. T., Li, Z., . . . Ruohola-
Baker, H. (2023). Single-cell census of human tooth development enables
generation of human enamel. *Dev Cell, 58*(20), 2163-2180 e2169.
doi:10.1016/j.devcel.2023.07.013
Bansal, A. K., Shetty, D. C., Bindal, R., & Pathak, A. (2012). Amelogenin: A novel protein with
diverse applications in genetic and molecular profiling. *J Oral Maxillofac Pathol,*
16(3), 395-399. doi:10.4103/0973-029X.102495
Barbeito-Andrés, J., Gonzalez, P., & Hallgrímsson, B. (2016). Prenatal development of skull
and brain in a mouse model of growth restriction. *Revista Argentina de Antropología*
Biológica, 18(1). doi:10.17139/raab.2016.0018.01.05
Baroncelli, L., Cenni, M. C., Melani, R., Deidda, G., Landi, S., Narducci, R., . . . Berardi, N.
(2017). Early IGF-1 primes visual cortex maturation and accelerates developmental
switch between NKCC1 and KCC2 chloride transporters in enriched animals.
Neuropharmacology, 113(Pt A), 167-177. doi:10.1016/j.neuropharm.2016.02.034
Bell, C. J., & Jass, C. N. (2011). Polyphyly, paraphyly, provinciality, and the promise of
intercontinental correlation - Charles Repenning's contributions to the study of
arvicoline rodent evolution and biochronology. *Palaeontologia Electronica, 14*(3),
15.
Brown, S. M., Peters, R., & Lawrence, A. B. (2017). Up-regulation of IGF-1 in the frontal
cortex of piglets exposed to an environmentally enriched arena. *Physiol Behav, 173*,
285-292. doi:10.1016/j.physbeh.2017.02.030

Burroughs, R. W. (2019). Phylogenetic and Developmental Constraints Dictate the Number
of Cusps on Molars in Rodents. *Sci Rep, 9*(1), 10902. doi:10.1038/s41598-019-
47469-x
Chantada-Vazquez, M. D. P., Bravo, S. B., Barbosa-Gouveia, S., Alvarez, J. V., & Couce, M. L.
(2022). Proteomics in Inherited Metabolic Disorders. *Int J Mol Sci, 23*(23).
doi:10.3390/ijms232314744
Chen, J. H., Martin-Gronert, M. S., Tarry-Adkins, J., & Ozanne, S. E. (2009). Maternal protein
restriction affects postnatal growth and the expression of key proteins involved in
lifespan regulation in mice. *PLoS One, 4*(3), e4950.
doi:10.1371/journal.pone.0004950
Choby, J. E., Monteith, A. J., Himmel, L. E., Margaritis, P., Shirey-Rice, J. K., Pruijssers, A., . . .
Skaar, E. P. (2019). A Phenome-Wide Association Study Uncovers a Pathological
Role of Coagulation Factor X during Acinetobacter baumannii Infection. *Infect*
Immun, 87(5). doi:10.1128/IAI.00031-19
Chrudinova, M., Kirk, N. S., Chuard, A., Venugopal, H., Zhang, F., Lubos, M., . . . Altindis, E.
(2024). A viral insulin-like peptide inhibits IGF-1 receptor phosphorylation and
regulates IGF1R gene expression. *Mol Metab, 80*, 101863.
doi:10.1016/j.molmet.2023.101863
Croft, D., O'Kelly, G., Wu, G., Haw, R., Gillespie, M., Matthews, L., . . . Stein, L. (2011).
Reactome: a database of reactions, pathways and biological processes. *Nucleic*
Acids Res, 39(Database issue), D691-697. doi:10.1093/nar/gkq1018
Dickerson, J. W. T., & Hughes, P. C. R. (1972). Growth of the rat skeleton after severe
nutritional intrauterine and post-natal retardation. *Resuscitation, 1*(2), 163-170.
doi:10.1016/0300-9572(72)90019-6
Dodington, D. W., Yumol, J. L., Yang, J., Pollock-Tahiri, E., Sivasubramaniyam, T., Sacco, S.
M., . . . Woo, M. (2021). JAK2-IGF1 axis in osteoclasts regulates postnatal growth in
mice. *JCI Insight, 6*(5). doi:10.1172/jci.insight.137045
Dong, C., Lamichhane, B., Yamazaki, H., Vasquez, B., Wang, J., Zhang, Y., . . . Wang, X.
(2022). The phosphorylation of serine(55) in enamelin is essential for murine
amelogenesis. *Matrix Biol, 111*, 245-263. doi:10.1016/j.matbio.2022.07.001
Du, W., Verma, A., Ye, Q., Du, W., Lin, S., Yamanaka, A., . . . Hu, J. K. (2024). Myosin II
mediates Shh signals to shape dental epithelia via control of cell adhesion and
movement. *PLoS Genet, 20*(6), e1011326. doi:10.1371/journal.pgen.1011326
Duverger, O., & Morasso, M. I. (2018). Pleiotropic function of DLX3 in amelogenesis: from
regulating pH and keratin expression to controlling enamel rod decussation.
Connect Tissue Res, 59(sup1), 30-34. doi:10.1080/03008207.2017.1408602
Eichelberger, K. R., & Goldman, W. E. (2020). Manipulating neutrophil degranulation as a
bacterial virulence strategy. *PLoS Pathog, 16*(12), e1009054.
doi:10.1371/journal.ppat.1009054
Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., Fillion-Robin, J. C., Pujol, S., . . .
Kikinis, R. (2012). 3D Slicer as an image computing platform for the Quantitative
Imaging Network. *Magn Reson Imaging, 30*(9), 1323-1341.
doi:10.1016/j.mri.2012.05.001

Freeman, T., Dixon, A., Campbell, E., Tait, T., Richardson, P., Rice, K., . . . Bentley, D. (1998).
883   *Expression Mapping of Mouse Genes*.

884   Froment, C., Zanolli, C., Hourset, M., Mouton-Barbosa, E., Moreira, A., Burlet-Schiltz, O., &
885   Mollereau, C. (2021). Protein sequence comparison of human and non-human
886   primate tooth proteomes. *J Proteomics, 231*, 104045.
887   doi:10.1016/j.jprot.2020.104045

888   Giovani, P. A., Martins, L., Salmon, C. R., Mofatto, L. S., Leme, A. F. P., Puppin-Rontani, R.
889   M., . . . Kantovitz, K. R. (2021). Comparative proteomic analysis of dental cementum
890   from deciduous and permanent teeth. *J Periodontal Res, 56*(1), 173-185.
891   doi:10.1111/jre.12808

892   Green, D. R., Schulte, F., Lee, K. H., Pugach, M. K., Hardt, M., & Bidlack, F. B. (2019).
893   Mapping the Tooth Enamel Proteome and Amelogenin Phosphorylation Onto
894   Mineralizing Porcine Tooth Crowns. *Front Physiol, 10*, 925.
895   doi:10.3389/fphys.2019.00925

896   Guhathakurta, P., Prochniewicz, E., & Thomas, D. D. (2018). Actin-Myosin Interaction:
897   Structure, Function and Drug Discovery. *Int J Mol Sci, 19*(9).
898   doi:10.3390/ijms19092628

899   Harjunmaa, E., Seidel, K., Hakkinen, T., Renvoise, E., Corfe, I. J., Kallonen, A., . . . Jernvall, J.
900   (2014). Replaying evolutionary transitions from the dental fossil record. *Nature,*
901   *512*(7512), 44-48. doi:10.1038/nature13613

902   Heck, M., & Neely, B. A. (2020). Proteomics in Non-model Organisms: A New Analytical
903   Frontier. *J Proteome Res, 19*(9), 3595-3606. doi:10.1021/acs.jproteome.0c00448
904

905   Holloway, P. J., Shaw, J. H., & Swenney, E. A. (1961). EFFECTS OF VARIOUS
906   SUCROSE:CASEIN RATIOS IN PURIFIED DIETS ON THE TEETH AND SUPPORTING
907   STRUCTURES OF RATS. *Arch Oral Biol, 3*, 185-200.
908

909   Hopkins, S. S. B. (2018). Estimation of Body Size in Fossil Mammals. In D. A. Croft, D. F. Su,
910   & S. W. Simpson (Eds.), *Methods in Paleoecology: Reconstructing Cenozoic*
911   *Terrestrial Environments and Ecological Communities* (pp. 7-22). Cham: Springer
912   International Publishing.

913   Huttlin, E. L., Jedrychowski, M. P., Elias, J. E., Goswami, T., Rad, R., Beausoleil, S. A., . . .
914   Gygi, S. P. (2010). A tissue-specific atlas of mouse protein phosphorylation and
915   expression. *Cell, 143*(7), 1174-1189. doi:10.1016/j.cell.2010.12.001

916   Immel, A., Le Cabec, A., Bonazzi, M., Herbig, A., Temming, H., Schuenemann, V. J., . . .
917   Krause, J. (2016). Effect of X-ray irradiation on ancient DNA in sub-fossil bones -
918   Guidelines for safe X-ray imaging. *Sci Rep, 6*, 32969. doi:10.1038/srep32969

919   Jagr, M., Ergang, P., Pataridis, S., Kolrosova, M., Bartos, M., & Miksik, I. (2019). Proteomic
920   analysis of dentin-enamel junction and adjacent protein-containing enamel matrix
921   layer of healthy human molar teeth. *Eur J Oral Sci, 127*(2), 112-121.
922   doi:10.1111/eos.12594

923   Levis, N. A., & Pfennig, D. W. (2021). Innovation and Diversification Via Plasticity-Led
924   Evolution. In *Phenotypic Plasticity & Evolution* (pp. 211-240).

Liu, C., & Zhang, R. (2021). Biomineral proteomics: A tool for multiple disciplinary studies. *J*
Proteomics, 238, 104171. doi:10.1016/j.jprot.2021.104171
Lobe, S. L., Bernstein, M. C., & German, R. Z. (2006). Life-long protein malnutrition in the rat
(Rattus norvegicus) results in altered patterns of craniofacial growth and smaller
individuals. *J Anat, 208*(6), 795-812. doi:10.1111/j.1469-7580.2006.00565.x
Luis, N. M., & Schnorrer, F. (2021). Mechanobiology of muscle and myofibril
morphogenesis. *Cells Dev, 168*, 203760. doi:10.1016/j.cdev.2021.203760
Luo, Y., Liu, Z., Luo, S., Wang, X., & Tao, L. (2020). The developmental and experience-
dependent expression of IGF-2 in mice visual cortex. *Neurosci Lett, 721*, 134828.
doi:10.1016/j.neulet.2020.134828
McCance, R. A., Ford, E. H., & Brown, W. A. (1961). Severe undernutrition in growing and
adult animals. 7. Development of the skull, jaws and teeth in pigs. *Br J Nutr, 15*, 213-
224. doi:10.1079/bjn19610026
Mehta, S., Bernt, M., Chambers, M., Fahrner, M., Foll, M. C., Gruening, B., . . . Griffin, T. J.
(2023). A Galaxy of informatics resources for MS-based proteomics. *Expert Rev*
Proteomics, 20(11), 251-266. doi:10.1080/14789450.2023.2265062
Miller, J. P., & German, R. Z. (1999). Protein Malnutrition Affects the Growth Trajectories of
the Craniofacial Skeleton in Rats. *Journal of Nutrition, 129*, 2061-2069.
Molloy, M. P., Hill, C., McKay, M. J., & Herbert, B. R. (2023). Proteome Analysis of Whole
Blood Collected by Volumetric Absorptive Microsampling. In D. W. Greening & R. J.
Simpson (Eds.), *Serum/Plasma Proteomics: Methods and Protocols* (pp. 173-179).
New York, NY: Springer US.
Montivero, A. J., Ghersi, M. S., Silvero, C. M., Artur de la Villarmois, E., Catalan-Figueroa, J.,
Herrera, M., . . . Perez, M. F. (2021). Early IGF-1 Gene Therapy Prevented Oxidative
Stress and Cognitive Deficits Induced by Traumatic Brain Injury. *Front Pharmacol,*
12, 672392. doi:10.3389/fphar.2021.672392
Mukhtar, U., Goyal, A., Luthra-Guptasarma, M., Gauba, K., Kapur, A., & Thakur, A. K. (2022).
Label-free quantitative proteomics reveals molecular correlates of altered
biomechanical properties in molar incisor hypomineralization (MIH): an in vitro
study. *Eur Arch Paediatr Dent, 23*(1), 179-191. doi:10.1007/s40368-021-00687-2
Muntel, J., Boswell, S. A., Tang, S., Ahmed, S., Wapinski, I., Foley, G., . . . Springer, M. (2015).
Abundance-based classifier for the prediction of mass spectrometric peptide
detectability upon enrichment (PPA). *Mol Cell Proteomics, 14*(2), 430-440.
doi:10.1074/mcp.M114.044321
Nanci, A. (2007). Development of the Tooth and Its Supporting Tissues. In *Ten Cate's Oral*
Histology: Development, Structure, and Function (pp. 79-107). St. Louis: Elsevier.
Orsini, G., Majorana, A., Mazzoni, A., Putignano, A., Falconi, M., Polimeni, A., & Breschi, L.
(2014). Immunocytochemical detection of dentin matrix proteins in primary teeth
from patients with dentinogenesis imperfecta associated with osteogenesis
imperfecta. *Eur J Histochem, 58*(4), 2405. doi:10.4081/ejh.2014.2405
Othman, A., Sekheri, M., & Filep, J. G. (2022). Roles of neutrophil granule proteins in
orchestrating inflammation and immunity. *FEBS J, 289*(14), 3932-3953.
doi:10.1111/febs.15803

Oyanagi, T., Takeshita, N., Hara, M., Ikeda, E., Chida, T., Seki, D., . . . Takano-Yamamoto, T.
(2019). Insulin-like growth factor 1 modulates bioengineered tooth morphogenesis.
Sci Rep, 9(1), 368. doi:10.1038/s41598-018-36863-6

Palma-Lara, I., Garcia Alonso-Themann, P., Perez-Duran, J., Godinez-Aguilar, R., Bonilla-
Delgado, J., Gomez-Archila, D., . . . Palacios-Reyes, C. (2023). Potential Role of
Protein Kinase FAM20C on the Brain in Raine Syndrome, an In Silico Analysis. *Int J*
Mol Sci, 24(10). doi:10.3390/ijms24108904

Pandya, M., & Diekwisch, T. G. H. (2021). Amelogenesis: Transformation of a protein-
mineral matrix into tooth enamel. *J Struct Biol, 213*(4), 107809.
doi:10.1016/j.jsb.2021.107809

Pandya, M., Liu, H., Dangaria, S. J., Zhu, W., Li, L. L., Pan, S., . . . Diekwisch, T. G. H. (2017).
Integrative Temporo-Spatial, Mineralogic, Spectroscopic, and Proteomic Analysis of
Postnatal Enamel Development in Teeth with Limited Growth. *Front Physiol, 8*, 793.
doi:10.3389/fphys.2017.00793

Patton, J. L., & Brylski, P. V. (1987). Pocket Gophers in Alfalfa Fields: Causes and
Consequences of Habitat-Related Body Size Variation. *The American Naturalist,*
130(4), 493-506.

Paynter, K. J., & Grainer, R. M. (1956). The relation of nutrition to the morphology and size of
rat molar teeth. *journal of the Canadian Dental Association, 22*(9), 519-531.

Paynter, K. J., & Grainer, R. M. (1961). Influence of Nutrition and Genetics on Morphology
and Caries Susceptibility. *Journal of the American Medical Association, 177*(5), 306-
309.

Porto, A., Rolfe, S., & Maga, A. M. (2021). ALPACA: A fast and accurate computer vision
approach for automated landmarking of three-dimensional biological structures.
Methods Ecol Evol, 12(11), 2129-2144. doi:10.1111/2041-210X.13689

Procopio, N., Chamberlain, A. T., & Buckley, M. (2018). Exploring Biological and Geological
Age-related Changes through Variations in Intra- and Intertooth Proteomes of
Ancient Dentine. *J Proteome Res, 17*(3), 1000-1013.
doi:10.1021/acs.jproteome.7b00648

Pucciarelli, H. M. (1980). The effects of race, sex, and nutrition on craniofacial
differeniation in rats. A multivariate analysis. *Am J Phys Anthropol, 53*(3), 359-368.
doi:10.1002/ajpa.1330530307

Pucciarelli, H. M. (1981). Growth of the functional components of the rat skull and its
alteration by nutritional effects. A multivariate analysis. *Am J Phys Anthropol, 56*(1),
33-41. doi:10.1002/ajpa.1330560104

RCoreTeam. (2023). R: A Language and Environment for Statistical Computing. In. Vienna,
Austria: R Foundation for Statistical Computing.

Rebeaud, M. E., Mallik, S., Goloubinoff, P., & Tawfik, D. S. (2021). On the evolution of
chaperones and cochaperones and the expansion of proteomes across the Tree of
Life. *Proc Natl Acad Sci U S A, 118*(21). doi:10.1073/pnas.2020885118

Ritchie, H. (2018). The functional significance of dentin sialoprotein-phosphophoryn and
dentin sialoprotein. *Int J Oral Sci, 10*(4), 31. doi:10.1038/s41368-018-0035-9

Rolfe, S., Pieper, S., Porto, A., Diamond, K., Winchester, J., Shan, S., . . . Maga, A. M. (2021). SlicerMorph: An open and extensible platform to retrieve, visualize and analyze 3D morphology. *Methods Ecol Evol, 12*(10), 1816-1825. doi:10.1111/2041-210x.13669

Sartoretto, S., Gemini-Piperni, S., da Silva, R. A., Calasans, M. D., Rucci, N., Pires Dos Santos, T. M., . . . Zambuzzi, W. F. (2019). Apoptosis-associated speck-like protein containing a caspase-1 recruitment domain (ASC) contributes to osteoblast differentiation and osteogenesis. *J Cell Physiol, 234*(4), 4140-4153. doi:10.1002/jcp.27226

Searle, A. G. (1954). The Influence of Diet on Variation within Pure Lines. In *Genetical Studies on the Skeleton of the Mouse* (pp. 413-424).

Sharma, V., Rastogi, S., Kumar Bhati, K., Srinivasan, A., Roychoudhury, A., Nikolajeff, F., & Kumar, S. (2020). Mapping the Inorganic and Proteomic Differences among Different Types of Human Teeth: A Preliminary Compositional Insight. *Biomolecules, 10*(11). doi:10.3390/biom10111540

Shaw, J. H., & Griffiths, D. (1963). Dental Abnormalities in Rats Attributable to Protein Deficiency During Reproduction. *Journal of Nutrition, 80*, 123-141.

Shi, C., Ma, N., Zhang, W., Ye, J., Shi, H., Xiang, D., . . . Liu, Q. (2020). Haploinsufficiency of Dspp Gene Causes Dentin Dysplasia Type II in Mice. *Frontiers in Physiology, 11*. doi:10.3389/fphys.2020.593626

Steward, K. F., Refai, M., Dyer, W. E., Copie, V., Lachowiec, J., & Bothner, B. (2023). Acute stress reduces population-level metabolic and proteomic variation. *BMC Bioinformatics, 24*(1), 87. doi:10.1186/s12859-023-05185-4

Tagliabracci, V. S., Wiley, S. E., Guo, X., Kinch, L. N., Durrant, E., Wen, J., . . . Dixon, J. E. (2015). A Single Kinase Generates the Majority of the Secreted Phosphoproteome. *Cell, 161*(7), 1619-1632. doi:10.1016/j.cell.2015.05.028

Ungar, P. S. (2010). *Mammal Teeth: Origin, Evolution, and Diversity*. Baltimore: Johns Hopkins University Press.

Vassilakos, G., Lei, H., Yang, Y., Puglise, J., Matheny, M., Durzynska, J., . . . Barton, E. R. (2019). Deletion of muscle IGF-I transiently impairs growth and progressively disrupts glucose homeostasis in male mice. *FASEB J, 33*(1), 181-194. doi:10.1096/fj.201800459R

Visel, A., Thaller, C., & Eichele, G. (2004). GenePaint.org: an atlas of gene expression patterns in the mouse embryo. *Nucleic Acids Res, 32*(Database issue), D552-556. doi:10.1093/nar/gkh029

Wan, C., Yuan, G., Luo, D., Zhang, L., Lin, H., Liu, H., . . . Chen, Z. (2016). The Dentin Sialoprotein (DSP) Domain Regulates Dental Mesenchymal Cell Differentiation through a Novel Surface Receptor. *Sci Rep, 6*, 29666. doi:10.1038/srep29666

Wang, S., Song, R., Wang, Z., Jing, Z., Wang, S., & Ma, J. (2018). S100A8/A9 in Inflammation. *Front Immunol, 9*, 1298. doi:10.3389/fimmu.2018.01298

Wang, Z., Liu, P. K., & Li, L. (2024). A Tutorial Review of Labeling Methods in Mass Spectrometry-Based Quantitative Proteomics. *ACS Meas Sci Au, 4*(4), 315-337. doi:10.1021/acsmeasuresciau.4c00007

Weber, E. M., & Olsson, I. A. S. (2008). Maternal behaviour in Mus musculus sp.: An
ethological review. *Applied Animal Behaviour Science, 114*(1-2), 1-22.
doi:10.1016/j.applanim.2008.06.006
Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-
Verlag.
Wittmann, N., Behrendt, A. K., Mishra, N., Bossaller, L., & Meyer-Bahlburg, A. (2021).
Instructions for Flow Cytometric Detection of ASC Specks as a Readout of
Inflammasome Activation in Human Blood. *Cells, 10*(11).
doi:10.3390/cells10112880
Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., . . . Yu, G. (2021). clusterProfiler 4.0: A
universal enrichment tool for interpreting omics data. *Innovation (Camb), 2*(3),
100141. doi:10.1016/j.xinn.2021.100141
Yakar, S., & Isaksson, O. (2016). Regulation of skeletal growth and mineral acquisition by
the GH/IGF-1 axis: Lessons from mouse models. *Growth Horm IGF Res, 28*, 26-42.
doi:10.1016/j.ghir.2015.09.004
Yin, C., & Heit, B. (2018). Armed for destruction: formation, function and trafficking of
neutrophil granules. *Cell Tissue Res, 371*(3), 455-471. doi:10.1007/s00441-017-
2731-8
Zougman, A., Wilson, J. P., & Banks, R. E. (2020). A simple serum depletion method for
proteomics analysis. *Biotechniques, 69*(2), 148-151. doi:10.2144/btn-2020-0017
1072

Graphical Abstract:

Altered Developmental Pathways

Significantly Altered Protein Expression

Low dietary protein during early development

Altered Molar Size

Following Page

Figure 1: Models of skulls of treatment specimens used for proteomic sampling and processed from µCT scans. A: Lateral and Ventral view of skull, showing where right upper M1 was removed, M2 and M3 are still visible. Light blue shows M1 still in place, Fuschia dots indicate where roots for M1 once inserted. B: Dorsal view of skull, opacity adjusted to 50% to show location of landmarks for straight-line skull estimate. C: Scheme of 25 landmarks placed on left lower m1 used to assess crown area and shape between treatment and control specimens.

A

20mm　　　20mm

B

20mm

C

1.5mm

Figure 2: Results of phenotypic investigation. A: Estimates of straight line skull length.
B: Estimates of crown area.

Following Page

Figure 3: Most enriched pathways from Gene Ontology (GO) analysis of CGSig. A: Biological Processes, B: Cellular Components, C: Metabolic Function

A CGSig: GO: Biological Process

B CGSig: GO: Cellular Components

C CGSig: GO: Metabolic Function

Figure 4: Volcano of specimen proteomics data exemplified by selected comparison from each group. A: From Group 1 Treatment 2 over Control 1. B: From Group 2 Treatment 3 over Control 4. Blue dots show Adjusted Abundance Ratio P-values ≤ 0.05. Labeled dots are proteins related to osteogenesis, odontogenesis, inflammation and immune response, and muscle contraction. Volcano plots of remaining 6 comparisons are provided in Supplemental Materials.

Table 1: List of proteins with significant differential expression between low dietary protein (10%) treatment and normal dietary protein (20%) control groups. Average Log2 fold change calculated by averaging of specimens with significant change of abundance ratio (SI 1). Up=Significant specimens have positive Log2 fold change meaning increased expression of that protein. Down = Specimens have negative Log2 fold change meaning decreased expression of that protein. Proteins with blank averages are where a consistent signal was not identified due to individual variation. The number of specimens which could be individually interpreted as Up (U) or Down (D) is provided. Data are from SI 1.

Uniprot Accession	Protein and Gene Name	GO: Process	Avg. Log2	Up/ Down
P63277	Amelogenin, X isoform GN=Amelx	Odontogenesis	-1.52	Down
O55188	Dentin matrix acidic phosphoprotein 1 GN=Dmp1	Odontogenesis		3U/2D
P97399	Dentin sialophosphoprotein GN=Dspp	Odontogenesis	1.77	Up
P43406	Integrin alpha-V GN=Itgav	Osteogenesis	-2.66	Down
Q05117	Tartrate-resistant acid phosphatase type 5 GN=Acp5	Osteogenesis	-1.71	Down
P28481	Collagen alpha-1(II) chain GN=Col2a1	Osteogenesis		1U/2D
O54974	Galectin-7 GN=Lgals7	Osteogenesis		3U/2D
P27005	Protein S100-A8 GN=S100a8	Inflammation and Immune Response	-1.27	Down
O88947	Coagulation factor X GN=F10	Inflammation and Immune Response	-0.23	Down
Q9EPB4	Apoptosis-associated speck-like protein containing a CARD GN=Pycard	Inflammation and Immune Response		1U/1D
Q9JM83	Calmodulin-4 GN=Calm4	Inflammation and Immune Response		2U/1D
Q9Z126	Platelet factor 4 GN=Pf4	Inflammation and Immune Response	0.92	Up
P16110	Galectin-3 GN=Lgals3	Inflammation and Immune Response	1.10	Up
P07743	BPI fold-containing family A member 2 GN=Bpifa2	Inflammation and Immune Response	1.50	Up
Q07456	Protein AMBP GN=Ambp	Inflammation and Immune Response	2.30	Up
Q61176	Arginase-1 GN=Arg1	Inflammation and Immune Response	2.40	Up
Q9JJZ2	Tubulin alpha-8 chain GN=Tuba8	Cytoskeleton Formation	-2.29	Down
Q8R1Q8	Cytoplasmic dynein 1 light intermediate chain 1 GN=Dync1li1	Cytoskeleton Formation	-1.59	Down
Q9CRB9	MICOS complex subunit Mic19 GN=Chchd3	Cytoskeleton Formation	-1.17	Down
Q91ZU6	Dystonin GN=Dst	Cytoskeleton Formation	-0.56	Down
P08032	Spectrin alpha chain, erythrocytic 1 GN=Spta1	Cytoskeleton Formation		1U/1D
Q61554	Fibrillin-1 GN=Fbn1	Cytoskeleton Formation	0.09	Up
P15508	Spectrin beta chain, erythrocytic GN=Sptb	Cytoskeleton Formation		1U/1D
P55292	Desmocollin-2 GN=Dsc2	Cytoskeleton Formation		2U/1D
P48193	Protein 4.1 GN=Epb41	Cytoskeleton Formation		2U/1D
P10637	Microtubule-associated protein tau GN=Mapt	Cytoskeleton Formation	1.07	Up
O35902	Desmoglein-3 GN=Dsg3	Cytoskeleton Formation	1.20	Up
Q7TSF1	Desmoglein-1-beta GN=Dsg1b	Cytoskeleton Formation	1.22	Up
P97350	Plakophilin-1 GN=Pkp1	Cytoskeleton Formation	1.58	Up
Q02257	Junction plakoglobin GN=Jup	Cytoskeleton Formation	1.65	Up
E9Q557	Desmoplakin GN=Dsp	Cytoskeleton Formation	1.80	Up
G5E8K5	Ankyrin-3 GN=Ank3	Cytoskeleton Formation	1.87	Up
Q3UV17	Keratin, type II cytoskeletal 2 oral GN=Krt76	Development of Cornified Envelope		3U/2D
A1L317	Keratin, type I cytoskeletal 24 GN=Krt24	Development of Cornified Envelope	0.76	Up
P04104	Keratin, type II cytoskeletal 1 GN=Krt1	Development of Cornified Envelope	1.26	Up
Q61781	Keratin, type I cytoskeletal 14 GN=Krt14	Development of Cornified Envelope	1.46	Up
P97347	Repetin GN=Rptn	Development of Cornified Envelope	1.47	Up
Q61414	Keratin, type I cytoskeletal 15 GN=Krt15	Development of Cornified Envelope	1.51	Up
Q09PK2	Retroviral-like aspartic protease 1 GN=Asprv1	Development of Cornified Envelope	1.70	Up
Q62266	Cornifin-A GN=Sprr1a	Development of Cornified Envelope	1.75	Up
A6BLY7	Keratin, type I cytoskeletal 28 GN=Krt28	Development of Cornified Envelope	1.89	Up
Q62267	Cornifin-B GN=Sprr1b	Development of Cornified Envelope	2.93	Up
A2AQP0	Myosin-7B GN=Myh7b	Muscle Contraction	-2.37	Down
Q9QZ47	Troponin T, fast skeletal muscle GN=Tnnt3	Muscle Contraction	-2.34	Down

Q5SX39	Myosin-4 GN=Myh4	Muscle Contraction	-2.25	Down
P13542	Myosin-8 GN=Myh8	Muscle Contraction	-2.24	Down
Q5SX40	Myosin-1 GN=Myh1	Muscle Contraction	-1.91	Down
P05977	Myosin light chain 1/3, skeletal muscle isoform GN=Myl1	Muscle Contraction	-1.88	Down
P68134	Actin, alpha skeletal muscle GN=Acta1	Muscle Contraction	-1.86	Down
Q3SX28	TREMBL:Q3SX28;Q5KR48 (Bos taurus) Tropomyosin 2	Muscle Contraction	-1.61	Down
P13412	Troponin I, fast skeletal muscle GN=Tnni2	Muscle Contraction	-1.59	Down
P97457	Myosin regulatory light chain 2, skeletal muscle isoform GN=Mylpf	Muscle Contraction	-1.42	Down
P97447	Four and a half LIM domains protein 1 GN=Fhl1	Muscle Contraction	-1.06	Down
A2ASS6	Titin GN=Ttn	Muscle Contraction	-0.80	Down
Q9JI91	Alpha-actinin-2 GN=Actn2	Muscle Contraction	-0.48	Down
P07310	Creatine kinase M-type GN=Ckm	Phosphorylation	-2.03	Down
P28650	Adenylosuccinate synthetase isozyme 1 GN=Adssl1	Phosphorylation	-1.24	Down
Q9R0Y5	Adenylate kinase isoenzyme 1 GN=Ak1	Phosphorylation	-0.71	Down
P29699	Alpha-2-HS-glycoprotein GN=Ahsg	Phosphorylation		2U/1D
Q9R0H0	Peroxisomal acyl-coenzyme A oxidase 1 GN=Acox1	Regulation	-2.12	Down
P63330	Serine/threonine-protein phosphatase 2A GN=Ppp2ca	Regulation	-1.78	Down
Q9EST5	Acidic leucine-rich nuclear phosphoprotein 32 B GN=Anp32b	Regulation	-1.18	Down
P07759	Serine protease inhibitor A3K GN=Serpina3k	Regulation	-1.17	Down
P28665	Murinoglobulin-1 GN=Mug1	Regulation	-0.88	Down
Q9Z2N8	Actin-like protein 6A GN=Actl6a	Regulation	-0.62	Down
O55042	Alpha-synuclein GN=Snca	Regulation	0.29	Up
Q5FWK3	Rho GTPase-activating protein 1 GN=Arhgap1	Regulation		1U/1D
Q9D5V5	Cullin-5 GN=Cul5	Regulation	0.38	Down
Q6P253	Dermokine GN=Dmkn	Regulation	0.89	Up
Q08189	Protein-glutamine gamma-glutamyltransferase E GN=Tgm3	Regulation	1.04	Up
Q60604	Adseverin GN=Scin	Regulation	1.32	Up
Q9D2Q8	Protein S100-A14 GN=S100a14	Regulation	1.45	Up
Q8CGR5	Kallikrein-14 GN=Klk14	Regulation	1.55	Up
P21570	Angiogenin GN=Ang	Regulation	2.42	Up
Q62203	Splicing factor 3A subunit 2 GN=Sf3a2	Transcription and Repair	-0.21	Down
Q9R0Q6	Actin-related protein 2/3 complex subunit 1A GN=Arpc1a	Transcription and Repair	2.09	Up
P32848	Parvalbumin alpha GN=Pvalb	Binding	-2.24	Down
P62631	Elongation factor 1-alpha 2 GN=Eef1a2	Binding	-2.01	Down
O35744	Chitinase-like protein 3 GN=Chil3	Binding	-1.82	Down
P31786	Acyl-CoA-binding protein GN=Dbi	Binding	-1.72	Down
P81117	Nucleobindin-2 GN=Nucb2	Binding	-1.09	Down
O70456	14-3-3 protein sigma GN=Sfn	Binding	1.09	Up
P63024	Vesicle-associated membrane protein 3 GN=Vamp3	Transport	-3.07	Down
Q9JLM8	Serine/threonine-protein kinase DCLK1 GN=Dclk1	Transport	-2.40	Down
P18872	Guanine nucleotide-binding protein G(o) subunit alpha GN=Gnao1	Transport	-2.35	Down
Q8VEH3	ADP-ribosylation factor-like protein 8A GN=Arl8a	Transport	-2.18	Down
P61028	Ras-related protein Rab-8B GN=Rab8b	Transport	-2.07	Down
P04247	Myoglobin GN=Mb	Transport	-1.71	Down
P02089	Hemoglobin subunit beta-2 GN=Hbb-b2	Transport	-1.51	Down

P04117	Fatty acid-binding protein, adipocyte GN=Fabp4	Transport	-1.16	Down
P02088	Hemoglobin subunit beta-1 GN=Hbb-b1	Transport		1U/1D
P04919	Band 3 anion transport protein GN=Slc4a1	Transport		1U/1D
P97449	Aminopeptidase N GN=Anpep	Transport		1U/1D
P14602	Heat shock protein beta-1 GN=Hspb1	Transport	1.44	Up
P24668	Cation-dependent mannose-6-phosphate receptor GN=M6pr	Transport	2.37	Up
Q8VHI3	GDP-fucose protein O-fucosyltransferase 2 GN=Pofut2	Metabolism	-2.35	Down
Q5SWU9	Acetyl-CoA carboxylase 1 GN=Acaca	Metabolism	-2.03	Down
Q9D2R0	Acetoacetyl-CoA synthetase GN=Aacs	Metabolism	-1.93	Down
O70503	Very-long-chain 3-oxoacyl-CoA reductase GN=Hsd17b12	Metabolism	-1.87	Down
P19096	Fatty acid synthase GN=Fasn	Metabolism	-1.83	Down
P21550	Beta-enolase GN=Eno3	Metabolism	-1.77	Down
O70250	Phosphoglycerate mutase 2 GN=Pgam2	Metabolism	-1.75	Down
Q8R429	Sarcoplasmic/endoplasmic reticulum calcium ATPase 1 GN=Atp2a1	Metabolism	-1.73	Down
Q9EQ06	Estradiol 17-beta-dehydrogenase 11 GN=Hsd17b11	Metabolism	-1.38	Down
P13634	Carbonic anhydrase 1 GN=Ca1	Metabolism	-0.92	Down
Q9WUB3	Glycogen phosphorylase, muscle form GN=Pygm	Metabolism	-0.87	Down
Q9CQT1	Methylthioribose-1-phosphate isomerase GN=Mri1	Metabolism	-0.25	Down
P00920	Carbonic anhydrase 2 GN=Ca2	Metabolism		1U/1D
O89020	Afamin GN=Afm	Metabolism		1U/1D
O09131	Glutathione S-transferase omega-1 GN=Gsto1	Metabolism	1.22	Up
P97370	Sodium/potassium-transporting ATPase subunit beta-3 GN=Atp1b3	Metabolism	1.22	Up
P47739	Aldehyde dehydrogenase, dimeric ZDP-preferring GN=Aldh3a1	Metabolism	1.48	Up
Q9QYJ0	DnaJ homolog subfamily A member 2 GN=Dnaja2	Metabolism	2.00	Up
Q9R099	Transducin beta-like protein 2 GN=Tbl2	Unknown	-1.52	Down
Q00898	Alpha-1-antitrypsin 1-5 GN=Serpina1e	Unknown	-1.25	Down
Q9D7B7	Probable glutathione peroxidase 8 GN=Gpx8	Unknown		1U/2D
P00687	Alpha-amylase 1 GN=Amy1	Unknown		3U/2D
Q8CIT9	Suprabasin GN=Sbsn	Unknown	1.28	Up
Q9Z331	Keratin, type II cytoskeletal 6B GN=Krt6b	Unknown	1.30	Up
Q8CF02	Protein FAM25C GN=Fam25c	Unknown	1.81	Up

Table 2: Significantly enriched REACTOME pathways derived from CGSig dataset (120 proteins). Table is ordered from highest Entities Ratio to lowest. Entities Ratio is the number of proteins from Pathway Name/Total Number of Proteins for *Mus musculus* in REACTOME database. Entities pValue is hypothesis test of chances of the number of proteins found being within pathway compared to random significant pValue = enriched pathway. Accession numbers of proteins from CGSig which belong to a pathway, not an inclusive list of all proteins known within the pathway.

REACTOME Pathway identifier	Pathway name	Proteins Found in Pathway	Total Number of Proteins in Pathway	#Reactions found	#Reactions total	Entities Ratio	Entities pValue	Uniprot Accession of Proteins Found In Pathway
R-MMU-1266738	Developmental Biology	19	573	22	218	0.0520	0.00001468838	P97350; P04104; Q8CGR5; Q61414; Q3UV17; P15508; P97347; P28481; Q9R0Q6; Q61781; P43406; P08032; Q02257; Q9Z331; O35902; P55292; E9Q557; Q9JJZ2; A6BLY7
R-MMU-6798695	Neutrophil degranulation	14	532	8	10	0.0483	0.00202335301	P29699; P97350; P04104; P97449; Q61176; P43406; P16110; Q8VEH3; Q02257; Q9JM83; Q9EPB4; E9Q557; Q8R1Q8; P27005
R-MMU-6805567	Keratinization	13	191	14	26	0.0173	0.00000019904	P97350; P04104; Q8CGR5; Q61414; Q3UV17; P97347; Q61781; Q02257; Q9Z331; O35902; P55292; E9Q557; A6BLY7
R-MMU-397014	Muscle contraction	9	180	12	43	0.0163	0.00017018443	Q9JI91; Q8R429; Q9QZ47; P05977; P68134; P97457; P13412; P97370; P13542
R-MMU-5357801	Programmed Cell Death	5	151	6	155	0.0137	0.02528751686	P97350; O70456; P10637; O35902; E9Q557
R-MMU-381426	Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	4	123	1	6	0.0112	0.04619853375	P29699; Q61554; O55188; P63277
R-MMU-8957275	Post-translational protein phosphorylation	4	117	1	1	0.0106	0.03970547416	P29699; Q61554; O55188; P63277
R-MMU-109581	Apoptosis	5	110	6	110	0.0100	0.00733981442	P97350; O70456; P10637; O35902; E9Q557
R-MMU-1169410	Antiviral mechanism by IFN-stimulated genes	4	101	3	44	0.0092	0.02511274198	P63330; P10637; O55042; Q9JJZ2
R-MMU-6807878	COPI-mediated anterograde transport	4	100	6	12	0.0091	0.02433051178	P08032; P15508; Q9JJZ2; Q8R1Q8
R-MMU-1500931	Cell-Cell communication	4	100	5	56	0.0091	0.02433051178	Q91ZU6; P21570; Q02257; Q61781
R-MMU-6809371	Formation of the cornified envelope	13	98	7	19	0.0089	0.00000000008	P97350; P04104; Q8CGR5; Q61414; Q3UV17; P97347; Q61781; Q02257; Q9Z331; O35902; P55292; E9Q557; A6BLY7
R-MMU-373760	L1CAM interactions	4	76	3	24	0.0069	0.00988125906	P43406; P08032; P15508; Q9JJZ2
R-MMU-446728	Cell junction organization	4	76	5	40	0.0069	0.00988125906	Q91ZU6; P21570; Q02257; Q61781
R-MMU-9013106	RHOC GTPase cycle	3	75	2	6	0.0068	0.04930012225	P63024; Q02257; Q5FWK3
R-MMU-9013026	RHOB GTPase cycle	3	74	2	6	0.0067	0.04771882744	P63024; Q02257; Q5FWK3
R-MMU-9833482	PKR-mediated signaling	4	71	3	28	0.0064	0.00784286893	P63330; P10637; O55042; Q9JJZ2
R-MMU-3371497	HSP90 chaperone cycle for steroid hormone receptors (SHR) in the presence of ligand	3	71	4	22	0.0064	0.04312926881	Q9QYJ0; Q9JJZ2; Q8R1Q8
R-MMU-75105	Fatty acyl-CoA biosynthesis	3	70	4	20	0.0064	0.04165134733	P19096; Q5SWU9; O70503
R-MMU-9013406	RHOQ GTPase cycle	3	62	2	5	0.0056	0.03078079105	P63024; Q02257; Q5FWK3
R-MMU-3000178	ECM proteoglycans	4	52	7	14	0.0047	0.00264050200	O55188; P43406; P97399; P28481
R-MMU-75153	Apoptotic execution phase	4	50	4	53	0.0045	0.00229488049	P97350; P10637; O35902; E9Q557
R-MMU-375165	NCAM signaling for neurite out-growth	3	45	4	8	0.0041	0.01342809454	P08032; P15508; P28481
R-MMU-390522	Striated Muscle Contraction	6	39	4	4	0.0035	0.00000509269	Q9JI91; Q9QZ47; P05977; P68134; P13412; P13542
R-MMU-111465	Apoptotic cleavage of cellular proteins	4	38	4	38	0.0034	0.00084596405	P97350; P10637; O35902; E9Q557
R-MMU-1237044	Erythrocytes take up carbon dioxide and release oxygen	3	24	3	8	0.0022	0.00240137363	P00920; P13634; P04919
R-MMU-1480926	O2/CO2 exchange in erythrocytes	3	24	5	14	0.0022	0.00240137363	P00920; P13634; P04919
R-MMU-140875	Common Pathway of Fibrin Clot Formation	2	24	6	29	0.0022	0.02860143761	O88947; Q9Z126
R-MMU-1475029	Reversible hydration of carbon dioxide	2	17	2	8	0.0015	0.01507296362	P00920; P13634
R-MMU-1247673	Erythrocytes take up oxygen and release carbon dioxide	3	16	2	6	0.0015	0.00075788188	P00920; P13634; P04919
R-MMU-445095	Interaction between L1 and Ankyrins	2	12	1	1	0.0011	0.00778004526	P08032; P15508
R-MMU-351906	Apoptotic cleavage of cell adhesion proteins	3	11	3	10	0.0010	0.00025621707	P97350; O35902; E9Q557
R-MMU-446107	Type I hemidesmosome assembly	2	9	2	4	0.0008	0.00447020414	Q91ZU6; Q61781
R-MMU-5660668	CLEC7A/inflammasome pathway	1	4	2	3	0.0004	0.04264489669	Q9EPB4

Table 3: Welch's Two Sample T-Tests to test for significant difference in mean between treatment and control group crown area and skull length.

Crown area by Group			
	t = -1.1575	df = 47.474	p-value = 0.8736
95% Confidence Intervals	-0.02914971	Infinity	
Mean Control	1.021672mm^2		
Mean Treatment	1.033573mm^2		
Skull Length by Group			
	t = -86423	df = 41.17	p-value = 0.8038
95% Confidence Intervals	-0.5085681	Infinity	
Mean Control	19.02755mm		
Mean Treatment	19.20011mm		

Supplemental Figures:

Volcano Plots from Burroughs et al. (In Review) Tooth Proteomics

Low-dietary protein (10%) sample specimen over control (20% protein) sample specimen. Blue dots show adjusted p-value is ≤ 0.05. Labeled proteins refer to Table 1 in Burroughs et al. (In Review).

1: Treatment 1 Over Control 1

2: Treatment 1 over Control 2

3: Treatment 2 over Control 2

4: Treatment 3 over Control 3

5: Treatment 4 over Control 3

6: Treatment 4 over Control 4