1 # The first complete mitochondrial genome of the endangered
2 # spotted eagle ray *Aetobatus ocellatus*

3
4
5

6 Alan Marín[1]
14 [1]Laboratorio de Genética, Fisiología y Reproducción, Facultad de Ciencias, Universidad
15 Nacional del Santa, Chimbote, Perú

29 CONTACT: Alan Marín amarin@uns.edu.pe

Abstract

The complete mitochondrial genome of the endangered spotted eagle ray *Aetobatus ocellatus*

was determined for the first time using Next Generation Sequencing (NGS) reads mined from

the Sequence Read Archive (SRA) of the GenBank database (BioSample SAMN31811701,

collection site: Bohol Island, Philippines). The spotted eagle ray mitogenome (GenBank

accession pending) has 20,217 bp in length displaying a typical vertebrate mitogenome

organization (13 protein-coding genes, 22 tRNA genes, 2 rRNA genes, and a control region).

A phylogenetic analysis based on the nucleotide sequence information of 13 protein-coding

genes from different related species confirmed the species status of *A. ocellatus* as an Indo-

West Pacific species, which was formerly described as *A. narinari* whose current accepted

distribution is restricted to the Atlantic Ocean.

54
55
56
57 Keywords: Aetobatidae, mitogenome, phylogenetics, Next Generation Sequencing

Eagle rays of the genus *Aetobatus* (Myliobatiformes: Aetobatidae) suffer from strong

overfishing and slow recovery from overexploitation due to their low regeneration rate, slow

growth, and late maturity (Yamaguchi et al. 2021), which has led to a continued population

decline throughout their distribution ranges (Dulvy et al. 2021; Finucci et al. 2024) and

subsequent categorization of all of its members as Endangered or Vulnerable within the IUCN

red list (IUCN 2024). The classification of Aetobatidae species is particularly susceptible to

taxonomic confusion due to the interspecific overlap of their morphological features (White et

al. 2013) and during the last decade a number of taxonomic changes were introduced with the

aid of genetic markers, including the resurrection of old synonyms and description of new

species (White 2010; White and Moore 2013; White et al. 2013).

Herein, I report the first complete mitochondrial genome of the endangered spotted eagle ray

Aetobatus ocellatus, which has only recently been resurrected as the valid name for the Indo-

West Pacific member of the *narinari* complex (White et al. 2010). The *A. ocellatus* sample was

collected in Bohol Island, Philippines (BioSample: SAMN31811701) and sequenced by the

Smithsonian National Museum of Natural History in their "NOAA Genome Skimming of

Marine animals" project (BioProject accession: PRJNA720393). Paired-end FASTQ genomic

reads (2.2 Gbp within 14.6 million reads, Platform: Illumina NovaSeq 6000) were downloaded

from the public GenBank SRA repository (SRA accession: SRS15852991), trimmed with

BBDuk (14.5 million reads remained), and merged with BBMerge (4.1 million reads remained)

implemented in Geneious Prime v. 2021.1.1 (Biomatters Ltd., Auckland, New Zealand).

Unmerged (22,239) and merged (16,553) reads were independently assembled against the only

complete available mitogenome reference sequence of *Aetobatus* prior to this study (*A.

flagellum,* 20,201 bp, GenBank accession NC_022837) using the "Map to Reference" tool in

Geneious Prime. Protein-coding (PC), transport RNA (tRNA), and ribosomal RNA (rRNA)

genes were annotated using the "Annotate and Predict" feature of Geneious Prime by

comparing the novel *A. ocellatus* mitogenome to other complete mitogenomes of closely

related species. The position of tRNAs was verified using tRNAscan-SE (Chan and Lowe

2019).

The 20,217 bp consensus mitogenome of *A. ocellatus* (GenBank accession pending) has a

typical fish mitogenome organization (Winn et al. 2024) with 13 PC, two rRNA, 22 tRNA

genes, and one putative control region, showing an overall nucleotide composition of A: 31.4%,

C: 27.1%, G: 15.5%, and T: 26%. All PCGs use the typical ATG start codon except for the

COI which utilizes the alternative GTG start codon.

The Bayesian phylogenetic analysis was performed using MrBayes 3.2 (Ronquist &

Huelsenbeck 2003) and consisted of a concatenated matrix of 13 PCGs (11,460 bp) from 15

mitogenomes of closely related species retrieved from the GenBank database. Substitution

saturation in single codon positions from each PCG was analyzed using DAMBE 5 (Xia 2013).

The software jModelTest 2 (Darriba et al. 2012) was used to find the best-fit model of evolution

for each PCG. Figure 1 shows the Bayesian phylogenetic tree in which all samples were

grouped into 4 distinct family clades according to their taxonomy. Aetobatidae species formed

a basal monophyletic clade with robust statistical support. *A. ocellatus* was clustered together

with *A. narinari* into a discrete subclade (338 variable sites, genetic divergence 3.03%),

reflecting their low estimated divergence age (from 2 to 5 mya, Sales et al. 2019). The genetic

material analyzed herein (BioSample SAMN31811701) came from an organism collected in

the Indo-West Pacific region (Bohol Island, Philippines) originally identified as *A. narinari*.

However, the phylogenetic results obtained herein indicate that BioSample SAMN31811701

belongs to *A. ocellatus*, supporting its species status as an Indo-West Pacific species (White et

al. 2010), while *A. narinari* is restricted to the Atlantic region (Sales et al. 2019).

Figure 1. Bayesian phylogenetic tree inferred from 13 concatenated mitochondrial protein-

coding genes of Aetobatidae, Dasyatidae, Mobulidae, and Rhinopteridae species. The sequence

matrix (11,460 nt) used in the phylogenetic analysis consisted of unambiguously aligned

regions of the first, second, and third codon positions. The position of *Aetobatus ocellatus* is

highlighted in bold. *Pteroplatytrygon violacea* and *Dasyatis pastinaca* (Dasyatidae) were used

as outgroups. Posterior probabilities are shown at each node. GenBank accession numbers for

each species are shown in parentheses.

References

- Chan, P.P., & Lowe, T.M. (2019). tRNAscan-SE: Searching for tRNA Genes in
Genomic Sequences. Methods Mol Biol. 1962:1-14.

- Darriba, D., Taboada, G.L., Doallo, & R., Posada D. (2012). jModelTest 2: more
models, new heuristics and parallel computing. Nat Methods. 9(8): 772–772.

- Dulvy, N.K., Carlson, J., Charvet, P., Ajemian, M.J., Bassos-Hull, K, Blanco-Parra,
MP, Chartrain, E., Derrick, D., Dia, M., Diop, M., Doherty, P., Dossa, J., De Bruyne,
G., Herman, K., Leurs, G.H.L., Mejía-Falla, P.A., Navia, A.F., Pacoureau, N., Pérez
Jiménez, J.C., Pires, J.D., Seidu, I., Soares, A.-L., Tamo, A., VanderWright, W.J. &
Williams, A.B. (2021). *Aetobatus narinari* (amended version of 2021 assessment). The
IUCN   Red   List   of   Threatened   Species 2021:
e.T42564343A201613657. https://dx.doi.org/10.2305/IUCN.UK.2021-
2.RLTS.T42564343A201613657.en. Accessed on 10 January 2025.

- Finucci,   B.,   Rigby,   C.L.,   Armstrong,   A.O.   &   Rezaie-Atagholipour,
M. (2024). *Aetobatus ocellatus*. The IUCN Red List of Threatened Species 2024:
e.T42566169A124549514. https://dx.doi.org/10.2305/IUCN.UK.2024-
1.RLTS.T42566169A124549514.en. Accessed on 09 January 2025.

- IUCN. The IUCN Red List of Threatened Species (2024). *Aetobatus* spp. status search:
https://www.iucnredlist.org/search?taxonomies=104020849&searchType=species
Accessed on 09 January 2025.

- Ronquist, F., & Huelsenbeck, J.P. (2003). MrBayes 3: Bayesian phylogenetic inference
under mixed models. Bioinformatics. 19(12), 1572-1574.

- Sales, J.B.L., de Oliveira, C.N., dos Santos, W.C.R., Rotundo, M.M., Ferreira, Y.,
Ready, J., ... & da Silva Rodrigues-Filho, L.F. (2019). Phylogeography of eagle rays of
the genus Aetobatus: *Aetobatus narinari* is restricted to the continental western Atlantic
Ocean. Hydrobiologia. 836, 169-183.

- White, W.T. (2010). Clarification of *Aetobatus ocellatus* (Kuhl, 1823) as a valid species,
and a comparison with *Aetobatus narinari* (Euphrasen, 1790) (Rajiformes:
Myliobatidae). CSIRO Mar Atmos Res. 32, 141.

- White, W.T., Furumitsu, K., & Yamaguchi, A. (2013). A new species of eagle ray
Aetobatus narutobiei from the Northwest Pacific: an example of the critical role
taxonomy plays in fisheries and ecological sciences. PLoS One. 8(12), e83785.

-    White, W.T., & Moore, A.B. (2013). Redescription of *Aetobatus flagellum* (Bloch &
163   Schneider, 1801), an endangered eagle ray (Myliobatoidea: Myliobatidae) from the
164   indo–West Pacific. Zootaxa 3752(1), 199-213.

165   -   Winn, J.C., Maduna, S.N., & Bester-van der Merwe, A.E. (2024). A comprehensive
166   phylogenomic study unveils evolutionary patterns and challenges in the mitochondrial
167   genomes of Carcharhiniformes: A focus on Triakidae. Genomics, 116(1), 110771.

168   -   Xia, X. (2013). DAMBE5: a comprehensive software package for data analysis in
169   molecular biology and evolution. Mol Biol Evol. 30(7): 1720–1728.

170   -   Yamaguchi, A., Furumitsu, K., & Wyffels, J. (2021). Reproductive biology and
171   embryonic diapause as a survival strategy for the East Asian endemic eagle ray
172   *Aetobatus narutobiei*. Front Mar Sci. 8, 768701.