Title: Novel Insights into T-Cell Exhaustion and Cancer Biomarkers in PDAC Using ScRNA-Seq

Abstract

One of the aggressive and lethal cancers, pancreatic ductal adenocarcinoma (PDAC) is characterised by poor prognosis and resistance to conventional treatments. Moreover, tumor immune microenvironment (TIME) plays a crucial role in the progression and therapeutic resistance of PDAC. It is associated with T-cells exhaustion, leading to the progressive loss of T-cell functions with impaired ability to kill tumor cells. Therefore, this study employed single cell RNA sequencing (scRNA-seq) analysis and identified upregulated genes of cancer cells in two groups ("cancer cells_vs_all-PDAC" and "cancer-PDAC_vs_all-normal"), while of T-cells, including CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells were identified between conditions (PDAC_vs_Normal) with their respective cell types. Subsequently, common and unique markers of cancer cells from both groups were identified, resulting in three sub-groups (common in both groups, unique to "cancer cells_vs_all-PDAC" and unique to "cancer-PDAC_vs_all-normal" group). Moreover, the top-10 enriched Reactome pathways of cancer cells (common and unique) and T-cells were identified; and the genes implicated in those pathways were selected to perform PPI analysis, eventually revealing the top-10 hub-genes of each group (common and unique) of cancer cells (*GAPDH, AKT1, EGFR, CS, RHOA, TPI1, SDHA, TFRC, FASN, HIF1A, H4C6, MYC, H3C12, DDX21, USP7, RFC4, APEX1, CDK9, H2BC9, NOP2, FN1, COL1A1, COL1A2, COL3A1, COL5A2, COL6A1, COL5A1, BGN, COL6A2,* and *FBN1*) and T-cells (*ACTB, AKT1, CD4, CTNNB1, FN1, H3-3B, HIF1A, HSP90AA1, HSP90AB1, HSPA8, IFNG, ITGB1, JUN, MAPK3, MMP9, NFKB1, TP53, UBB,* and *UBC*). Furthermore, the gene expression validation of cancer cell and T-cells hub-genes was performed using GEPIA2 and TISCH2, while overall survival analysis of cancer cell hub-genes was performed using GEPIA2. Conclusively, this study unravelled a total of 16 novel markers (*CS, H4C6, H3C12, H2BC9, TP53, FN1, MMP9, ITGB1, HSP90AB1, AKT1, ACTB, MAPK3, UBC, UBB, CTNNB1, and H3-3B*) of cancer and T-cells, indicating that these might be implicated in the pathways leading to T-cell exhaustion, rendering growth, proliferation, and survival of cancer cells. This study provides the groundwork for future research into the immune landscape of PDAC, particularly T-cell exhaustion. However, further clinical studies are needed to validate these novel markers as potential therapeutic targets in PDAC patients.

Keywords: PDAC, T-cell exhaustion, Immune landscape of PDAC, PDAC TIME, PDAC heterogeneity

Introduction

Pancreatic tumors are classified as epithelial or non-epithelial based on their histogenesis, where pancreatic ductal adenocarcinoma (PDAC) is a most common malignant epithelial tumor of the pancreas accounting for more than 85% of all pancreatic malignancies (1). The cancerous exocrine duct cells lining the pancreas leads to PDAC and is the deadliest of all adult abdominal tumors (2). Moreover, PDAC has many subtypes, including classical and basal, squamoid-basaloid, treatment-enriched, and quasi-mesenchymal (3,4).

Furthermore, PDAC is the third-leading cause of cancer death in the age groups 50–64 and 65–79; while there is only 10.8% 5-year overall survival rate for PDAC for both metastatic and resectable cases (5,6). Strikingly, 62,210 cases were reported in the United States in 2023 and 49,380 deaths in 2022, while PDAC accounts for 2% of all cancer cases and results in 5% of all cancer deaths in the United States, therefore underlining the crucial need for earlier detection (5,7). Notably, PDAC is estimated to become the second-leading cause of cancer deaths by 2030, surpassing breast cancer, as mortality rates are on the rise, increasing 1% annually (8).

Major risk factors of PDAC include smoking, obesity, diabetes, family history of pancreatic cancer, and inherited genetic mutations such as those in *BRCA1*, *BRCA2,* and *PALB2* (9,10). Subsequently, the common symptoms of PDAC patients include jaundice due to bile duct obstruction, unexplained weight loss, abdominal or back pain, and new-onset diabetes. These symptoms typically arise at advanced stages, contributing to the poor prognosis associated with PDAC (11).

Currently, different treatment regimens, including surgery, chemotherapy and radiation therapy are used to treat PDAC; however, these treatment options remain ineffective due to highly resistant PDAC (12). Moreover, the gold standard to treat PDAC are polychemotherapy regimens, such as gemcitabine and abraxane or FOLFIRINOX (13). Recently, advances in immunotherapy have led to a shift in treatment regimens by gaining insights into the intricate interactions between the immune system and cancer cells, while PDAC has been resistant to immunotherapy and has demonstrated marginal efficacy in terms of survival (14,15).

Immunotherapy faces hurdles in PDAC treatment due to the immunosuppressive microenvironment that hinders the T-cell infiltration and activation, limiting the effectiveness of the combination therapy (16). The complex tumor immune microenvironment (TIME) of PDAC modulates the infiltration of the immunosuppressive cells and the activity of immune

regulatory molecules, leading to the dysfunctionality of the anti-tumor immune responses, including the exhaustion of the T-cells (17).

T-cell exhaustion is a dysfunctional state of the T-cells in chronic infections and cancer which results in cancer immune evasion, eventually leading to the poor prognosis in many cancers (18). In PDAC, the T-cells get activated which later differentiate into memory-like cells, eventually leading to differentiated exhausted T-cells, which result from the persistent exposure of antigens on cells with low proliferative capacity, loss of their cytotoxic function, increased apoptosis, and elevated expression of multiple inhibitory receptors such as immune checkpoints, including PD-1, CTLA-4, TIM-3, LAG-3, or TIGIT (19,20). The mechanism of T-cell exhaustion induced within the PDAC TIME is illustrated in **Figure 1**.

Moreover, PDAC is characterised by tumor-infiltrating CD4$^+$ and CD8$^+$ T-cells; however as PDAC progresses, percentage of Tregs elevates within the CD4$^+$ T-cell subset, while CD8$^+$ T-cells shifts in a decreased composition (21,22). Although T-cells are observed within the TIME of PDAC, it is considered as a poorly immune responsive cancer, as T-cells exhibit lack of activation or an exhausted phenotype (23).

Furthermore, a recent scRNA-seq study in combination with multiplex immunohistochemistry imaging, evaluated the T-cell landscape in PDAC, demonstrating that infiltrated CD8+ T-cells exhibited senescent or an exhausted phenotype with high expression levels of TIGIT$^+$ and CD39$^+$ alongside PD-1$^{low/intermediate}$ expression (24).

Another, scRNA-seq study revealed a reduction of ligand-receptor interactions after chemotherapy, especially between TIGIT on CD8$^+$ T-cells and its receptors on cancer cells, identifying TIGIT as a key inhibitory checkpoint molecule, suggesting that chemotherapy may contribute to immunotherapy resistance by impacting the immune interactions within the PDAC TIME (25).

Therefore, this study aimed to unravel the immune landscape, specifically T-cell landscape to identify the novel exhaustive T-cell markers, driven by the markers of cancer cells. This study utilised computational methods such as scRNA-seq analysis, differential gene expression, protein-protein interaction analysis, hub-genes identification, and expression validations of the identified hub-genes to gain insights into the molecular and functional aspects of T-cell exhaustion and to identify novel markers of cancer and T-cells associated with T-cell exhaustion. However, further clinical research is needed to validate the findings in this study.

Figure 1. The mechanism of T-cell exhaustion induced within the PDAC TIME shows that in PDAC, cancer cells release tumor-associated antigens (TAAs) into the tumor immune microenvironment (TIME), which are presented by antigen-presenting cells (APCs) to T-cells, causing chronic TCR stimulation. This triggers T-cell hyperactivation, amplified by PDAC-secreted cytokines like IL-6 and TGF-β, leading to metabolic stress. Prolonged activation induces upregulation of inhibitory receptors (PD-1, CTLA-4) on T-cells, while PDAC cells overexpress PD-L1. The PD-L1/PD-1 interaction suppresses T-cell effector functions, reducing cytokine production (e.g., IFN-γ) and cytotoxicity, enabling tumor immune evasion.

Methodology

ScRNA-seq dataset retrieval

The preprocessed (mapped) scRNA dataset of PDAC with an accession id GSE212966 was retrieved using the database, Gene Expression Omnibus (GEO) datasets (https://www.ncbi.nlm.nih.gov/gds) of National Center for Biotechnology Information (NCBI), a public repository that archives and distributes functional genomic data (26). The dataset consisted of two conditions (PDAC and control), with six PDAC samples and six control samples, comprising a total of 12 samples taken from untreated PDAC patients and sequenced using Illumina NovaSeq 6000 platform.

Data preprocessing and clustering

The "Seurat" package within R was utilised for the scRNA-seq analysis of the retrieved dataset, comprising 36601 genes and 57167 cells which were subjected to the quality control measures, including "nFeature_RNA > 200 & < 6000", "nCount_RNA > 1000", and mitochondrial reads < 10, filtering out the undesired and dead cells.

Moreover, the "LogNormalize" method was used for the normalization of the data, followed by the identification of 2000 highly variable features using the "variance stabilizing transformation (vst)" method. Furthermore, the data was scaled and linear dimensionality reduction was performed through principal component analysis (PCA). Subsequently, elbow plot was observed and first 15 PCs were utilised to identify the cell clusters using "FindNeighbors" and "FindClusters" functions at a resolution of 0.6 and utilising the Louvain algorithm.

Cell annotation and differentially expressed genes (DEGs)

The identified cell clusters were annotated using the "ScType" database (https://sctype.app/), a computational platform which provides automated cell-type identifications (27). After the identification of cell types, DEGs of cancer cells, CD8+ NKT-like cells, memory CD4+ T cells, and naive CD4+ T cells were identified using the "FindMarkers" function and the "wilcox" method with a threshold of ±0.25.

The differential gene expression analysis of cancer cells was performed in two groups, including cancer cells compared to all other cell types (cancer cells_vs_all-PDAC) within the PDAC condition and cancer cells within PDAC condition compared to all other cell types in

normal condition (cancer-PDAC_vs_all-normal). Moreover, the CD8+ NKT-like cells, memory CD4+ T cells, and naive CD4+ T cells within PDAC condition were compared to the same cells in normal condition (CD8+ NKT-like cells-PDAC_vs_CD8+ NKT-like cells-normal, memory CD4+ T cells-PDAC_vs_memory CD4+ T cells-normal, and naive CD4+ T cells-PDAC_vs_naive CD4+ T cells-normal).

The differential gene expression of the aforementioned combinations of cancer cells was performed to gain insights into the upregulated markers implicated in the heterogeneity and progression of cancer cells by exhausting the T-cells within the TIME. Furthermore, the differentially expressed genes of T-cells (CD8+ NKT-like cells, memory CD4+ T cells and naive CD4+ T cells) indicated the implication of upregulated markers in the loss of function of T-cells, which might be due to T-cell exhaustion, enabling cancer cells to grow and progress rapidly in TIME of PDAC.

Subsequently, common and unique upregulated markers of cancer cells from both groups were distinguished by utilising the Venn diagrams using the web-based tool, Venny 2.0 (https://bioinfogp.cnb.csic.es/tools/venny/index2.0.2.html).

Pathways enrichment analysis, protein-protein interaction (PPI) analysis and hub-genes identification

The identified upregulated genes of cancer cells, CD8+ NKT-like cells, memory CD4+ T cells, and naive CD4+ T cells were used for the pathways enrichment analysis using the GeneCodis4 (https://genecodis.genyo.es/), a web-based tool for the functional enrichment analysis allowing researchers to integrate different sources of annotations (28). The Reactome pathways were selected and the genes implicated in the top-10 upregulated pathways were used for PPI analysis using STRING (https://string-db.org), a database which collects, scores and integrates all the publicly available sources of PPI information, and to complement these with the computational predictions (29).

Lastly, the top-10 enriched hub-genes were identified from the proteins interaction network through the "Degree" method by using the Cytoscape, an open-source software which integrates the biomolecular interaction networks with high-throughput expression data (30).

Aberrant genes expression validation and survival rates in PDAC patients

The gene expression validations and survival analysis of the identified hub genes of cancer cells was performed using GEPIA2 (http://gepia2.cancer-pku.cn/#index) web server,

which is used for large-scale expression profiling and interactive analysis (31). Moreover, the genes expression validation of CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells was performed using Tumor Immune Single Cell Hub 2 (TISCH2, http://tisch.comp-genomics.org/), a scRNA-seq data resource from human and mouse tumors, enabling extensive characterization of gene expression in TIME (32). Furthermore, this was performed to shortlist the crucial hub genes implicated in the progression of PDAC.

Results

Data preprocessing and cell annotation

The retrieved dataset consisting of six PDAC samples and six control samples were merged using the "merge" function in the "Seurat" package, resulting in a combined matrix. This matrix was then converted into a "Seurat Object", which was further used for quality control measures retaining 38589 cells comprising 36601 genes. Subsequently, normalization, scaling, and highly variable features were identified. Among the 2000 highly variable features, the top-10 were *JCHAIN, IGKC, SPP1, EREG, APOC1, MZB1, IGHG1*, *TPSB2, CPA3* and *TPSAB1*. Moreover, the linear dimensionality reduction and cell clustering was performed based on gene expression profiles, resulting in a total of 26 cell clusters at a resolution of 0.6. The quality control, highly variable features, PCA plot, elbow plot, and t-SNE of cell clusters are shown in **Supplementary Figure S1(a-e)**.

Furthermore, the cell annotation resulted in 15 cell types, including Basophils, Cancer cells, CD8+ NKT-like cells, Endothelial, Macrophages, Memory CD4+ T cells, Myeloid Dendritic cells, Naive CD4+ T cells, Natural killer cells, Neutrophils, Plasma B cells, Plasmacytoid Dendritic cells, Platelets, Pre-B cells, and Progenitor cells. Interestingly, it was observed that the control condition exhibited trace amounts of cancer cells, indicating the spread of PDAC to control (healthy) regions of the pancreas, leading to metastasis.

Subsequently, it was observed that CD8+ NKT-like cells and memory CD4+ T cells exhibited a large and dense cluster in PDAC condition as compared to the control condition, suggesting the infiltration of T-cells at the tumor site. However, naive CD4+ T cells were observed to have a large and dense cluster of cells in control condition. The cell types of control and PDAC conditions are shown by using t-SNE method in **Figure 2a**.

Aberrant markers of cancer cells, CD8+ NKT-like cells, memory CD4+ T cells and naive CD4+ T cells

The differential gene expression of the "cancer cells_vs_all-PDAC" group resulted in 8,509 upregulated and 4,642 downregulated genes, totalling to 13,151 dysregulated genes. Among these genes the top 10 upregulated included *AC007529.2, LINC02747, FGA, LGALS7, RASSF10, FER1L6, AC036176.3, NPSR1, SERPINB3,* and *AADAC* with p-values of 3.31E-47, 8.77E-49, 4.02E-75, 3.06E-49, 6.12E-60, 0, 6.53E-121, 0, 0, and 0; while logarithm of fold change (log2FC) of 8.60, 8.12, 7.89, 7.88, 7.82, 7.76, 7.75, 7.72, 7.71, and 7.63, respectively.

Moreover, the "cancer-PDAC_vs_all-normal" group exhibited 9,268 upregulated and 4,221 downregulated genes, totalling to 13,489 dysregulated genes. Among these genes the top 10 upregulated included *CEACAM5, MUC5B, SERPINB3, KRT6B, TFF2, AL121761.1, TFF1, CEACAM6, EPS8L3,* and *SPRR3* with p-values of 0 except for *SPRR3* showing a p-value of 1.58E-108; while these genes showed log2FC of 14.42, 14.30, 13.77, 13.47, 13.38, 13.11, 13.10, 13.07, 13.01, and 12.60, respectively. The expression levels of both groups of cancer cell genes are shown in **Supplementary Figure S2 (a-b)**.

Furthermore, CD8+ NKT-like cells exhibited 2,598 upregulated and 3,771 downregulated genes, totalling to 6,369 dysregulated genes. The top 10 upregulated genes included *COL11A1, MUC5AC, CEACAM5, FXYD3, MUC5B, MUC4, KLK6, TSPAN1, GPX2,* and *CTSE* with p-values of 1.78E-26, 5.76E-30, 9.61E-21, 3.25E-283, 1.35E-16, 2.17E-15, 1.03E-09, 1.65E-08, 1.90E-06, and 2.55E-28, while log2FC of 8.81, 8.80, 8.40, 8.23, 8.01, 7.94, 7.35, 7.34, 7.34, and 7.24 respectively.

The memory CD4+ T cells showed 2,820 upregulated and 3,500 downregulated genes, totalling to 6,320 dysregulated genes. The top 10 upregulated genes included *FXYD3, TFF1, MUC1, OLFM4, AGR2, LCN2, CTSE, CEACAM6, KRT19,* and *CLDN4* with p-values of 1.53E-18, 1.22E-14, 1.33E-11, 7.02E-08, 1.90E-05, 6.27E-05, 0.005094647, 0.0036388, 7.40E-13, and 0.00020176; while log2FC of 8.42, 8.17, 7.51, 7.45, 6.97, 6.75, 6.51, 6.47, 6.37, and 5.39, respectively.

The naive CD4+ T cells resulted in 4,331 upregulated and 3,012 downregulated genes, totalling to 7,343 dysregulated genes. The top 10 upregulated genes included *KRT19, TFF1, LCN2, AGR2, TFF2, OLFM4, CLDN18, SPP1, C19orf33,* and *CEACAM6* with p-values of 1.14E-19, 4.28E-10, 7.87E-11, 4.89E-13, 1.45E-11, 3.76E-07, 1.45E-11, 2.12E-33, 7.87E-11, and 4.28E-10; while log2FC of 8.30, 7.95, 7.81, 7.75, 7.72, 7.61, 7.49, 7.49, 7.46, and 7.38, respectively. The expression levels of T-cells top 10 upregulated markers are shown in **Supplementary Figure S2 (c-e)**.

It was observed that the "cancer-PDAC_vs_all-normal" group exhibited the most and high intensity of dysregulation compared to the "cancer cells_vs_all-PDAC" group, indicating that markers in this group might be implicated in the rapid progression of cancer cells in TIME and may be highly implicated in signalling, leading to T-cell exhaustion. Nevertheless, it was observed that among T-cells, naive CD4+ T cells showed the highest number of dysregulated genes, indicating high influence of cancer cells on naive CD4+ T cells.

Subsequently, the upregulated markers of cancer cells showed 7,882 common genes among both groups, while 1,386 unique to "cancer-PDAC_vs_all-normal"; and 627 unique to

"cancer cells_vs_all-PDAC" group. Henceforth, all these genes were used separately for further analysis to identify unique and common markers in cancer cells while comparing different conditions. The venn diagram of "cancer-PDAC_vs_all-normal" and "cancer cells_vs_all-PDAC" groups is shown in **Supplementary Figure S2-f**.

Pathways dysregulation leading to T-cell exhaustion and PDAC progression

The upregulated markers were used to identify the enriched Reactome pathways. The common markers of cancer cells included Metabolism, Metabolism of lipids, Asparagine N-linked glycosylation, Post-translational protein modification, Membrane Trafficking, RHO GTPase cycle, Vesicle-mediated transport, Regulation of cholesterol biosynthesis by SREBP (SREBF), ER to Golgi Anterograde Transport, and RHOB GTPase cycle to be enriched top-10 pathways.

While the unique markers to "cancer cells_vs_all-PDAC" group included Gene expression (Transcription), Generic Transcription Pathway, RNA Polymerase II Transcription, Metabolism of RNA, DNA Repair, Antiviral mechanism by IFN-stimulated genes, SMAC (DIABLO) binds to IAPs, SMAC(DIABLO)-mediated dissociation of IAP complexes, Abasic sugar-phosphate removal via the single-nucleotide replacement pathway, and SMAC, XIAP-regulated apoptotic response to be top-10 enriched pathways.

Lastly, the unique markers to "cancer-PDAC_vs_all-normal" group included Extracellular matrix organization, Collagen formation, Assembly of collagen fibrils and other multimeric structures, Collagen degradation, Collagen biosynthesis and modifying enzymes, Degradation of the extracellular matrix, ECM proteoglycans, Collagen chain trimerization, MET activates PTK2 signaling, and Integrin cell surface interactions to be enriched top-10 pathways. The Reactome pathways of cancer cells markers are shown in **Figure 2 (b-d)**.

Moreover, the CD8+ NKT-like cells exhibited Extracellular matrix organization, Non-integrin membrane-ECM interactions, Assembly of collagen fibrils and other multimeric structures, Collagen formation, Signaling by Receptor Tyrosine Kinases, ECM proteoglycans, MET activates PTK2 signaling, Immune System, MET promotes cell motility, and Syndecan interactions top-10 upregulated pathways.

Furthermore, the top-10 upregulated pathways of memory CD4+ T cells included Signaling by Receptor Tyrosine Kinases, Neutrophil degranulation, Immune System, Syndecan interactions, Assembly of collagen fibrils and other multimeric structures, Potential

therapeutics for SARS, Collagen degradation, Extracellular matrix organization, Disease, and TP53 Regulates Transcription of Genes Involved in G2 Cell Cycle Arrest.

Lastly, the top-10 enriched pathways of naive CD4+ T cells exhibited Gene expression (Transcription), Metabolism of RNA, Post-translational protein modification, Immune System, Processing of Capped Intron-Containing Pre-mRNA, RNA Polymerase II Transcription, Cytokine Signaling in Immune system, Vesicle-mediated transport, Membrane Trafficking, and mRNA Splicing. The upregulated Reactome pathways of T-cells markers are shown in **Figure 2 (e-g)**.

Notably, it indicated that all the cancer cells pathways were uniquely implicated in cancer cells to evade the immune system and to promote its growth and progression in TIME by exhausting the T-cells. Moreover, it was observed that "Immune system" pathway was common among all T-cells, while "Extracellular matrix organization", "Assembly of collagen fibrils and other multimeric structures", "Signaling by Receptor Tyrosine Kinases", and "Syndecan interactions" pathways were found to be common among CD8+ NKT-like cells and memory CD4+ T cells.

This suggested that cancer cells were sending similar signals to upregulate similar pathways in CD8+ NKT-like cells and memory CD4+ T cells. However, naive CD4+ T-cells did not exhibit common pathways with other T-cells except for the "Immune system", suggesting the exhaustion of all three T-cells in this study, eventually leading to suppression of the immune system. This also indicates the heterogeneity that cancer cells dysregulated the immune cells (T-cells) through specific signals to specific T-cells, which creates complex interactions within the TIME of PDAC.

Figure 2. The analyses show annotated clusters and enriched pathways implicated in cancer and T-cells. **(a)** The t-SNE map showing annotated cells in Normal and PDAC conditions, **(b-d)** Upregulated Reactome pathways of cancer cells genes, **(e-g)** Upregulated Reactome pathways of CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells

Key candidate proteins associated with T-cell exhaustion within PDAC TIME

The genes from top-10 significant pathways of each cell type were subjected to PPI analysis resulting in different networks of proteins interacting with each other. The upregulated 1,960 common pathway genes of cancer cells resulted in a network of 1,949 proteins interacting with each other. Moreover, 124 upregulated genes of the "cancer cells_vs_all-PDAC" group exhibited a network of 91 proteins, while 69 upregulated genes of the "cancer-PDAC_vs_all-normal" group resulted in a network of 69 proteins interacting with each other. The PPI networks of cancer cell genes are shown in **Figure 3 (a-c)**.

Furthermore, 520 upregulated genes of CD8+ NKT-like cells exhibited a network of 513 proteins, while 657 enriched genes of memory CD4+ T cells showed a network of 639 proteins, and 1,454 upregulated genes of naive CD4+ T cells showed a network of 1,445

proteins. The protein networks of CD8+ NKT-like cells, memory CD4+ T cells, and naive CD4+ T cells are shown in **Figure 3 (d-f)**.

Subsequently, these protein networks of each cell type were used to identify the most interacting genes (hub-genes) using the "Degree" method. The protein network of common cancer cell genes showed top-10 hub genes, including *GAPDH, AKT1, EGFR, CS, RHOA, TPI1, SDHA, TFRC, FASN,* and *HIF1A*; showing 364, 284, 242, 212, 207, 200, 189, 189, 186, and 179 interactions in the network, respectively.

The protein network of unique cancer cell genes of "cancer cells_vs_all-PDAC" group exhibited top 10 hub genes, including *H4C6* (*HIST1H4A* or *HIST1H4C), MYC, H3C12* (*HIST1H3A* or *HIST1H3D), DDX21, USP7, RFC4, APEX1, CDK9, H2BC9* (*HIST1H2BH),* and *NOP2*. Among these proteins, *RFC4, APEX1, CDK9, HIST1H2BH, and NOP2* showed 12 interactions, while *DDX21* and *USP7* showed 16 interactions; lastly, *H4C6, MYC* and *H3C12* exhibited 29, 28, and 18 interactions, respectively.

The protein network of unique cancer cell genes of "cancer-PDAC_vs_all-normal" group showed top 10 hub genes, including *FN1, COL1A1, COL1A2, COL3A1, COL5A2, COL6A1, COL5A1, BGN, COL6A2,* and *FBN1*, showing 62, 59, 56, 55, 51, 50, 50, 48, 46, and 45 interactions within the protein network, respectively. The hub-genes of cancer cells are shown in **Figure 3 (g-i)**. The common and unique hub genes of cancer cells from both groups are mentioned in **Table 1**.

Moreover, the protein network of CD8+ NKT-like cells exhibited *TP53, FN1, MMP9, CD4, IFNG, NFKB1, HIF1A, HSP90AA1, ITGB1,* and *HSP90AB1* as the top 10 hub genes, showing 176, 170, 149, 147, 147, 142, 129, 126, 119, and 116 interactions, respectively. Furthermore, *AKT1, TP53, ACTB, CD4, JUN, FN1, MMP9, HSP90AA1, MAPK3,* and *HSP90AB1* were the top 10 hub genes in memory CD4+ T cells, exhibiting 200, 192, 180, 148, 146, 142, 126, 123, 121, and 113 interactions. Lastly, the protein network of naive CD4+ T cells showed *TP53, UBC, UBB, HSP90AA1, JUN, CTNNB1, NFKB1, HSP90AB1, HSPA8,* and *H3-3B* (*H3F3B*) top 10 hub genes, showing 452, 298, 279, 270, 260, 260, 251, 238, 229, and 212 interactions, respectively. The hub-genes of T-cells are shown in **Figure 3 (j-l)**. The hub genes of all three T-cells are mentioned in **Table 2**.

Notably, the sub-groups of cancer cells (common genes, cancer cells_vs_all-PDAC, and cancer-PDAC_vs_all-normal) indicated the unique functional implication of respective hub-genes within their specific group. This suggests the complexity and heterogeneity of cancers within the TIME, as multiple different genes are implicated in the progression of PDAC cancer by several factors, including T-cell exhaustion.

Subsequently, it was observed that *TP53, HSP90AA1,* and *HSP90AB1* were common among all three T-cells; *FN1, MMP9,* and *CD4* were common in CD8+ NKT-like cells and memory CD4+ T cells; *NFKB1* was common in CD8+ NKT-like cells and naive CD4+ T cells, while *JUN* was common in memory CD4+ T cells and naive CD4+ T cells. This indicated that multiple hub-genes of T-cells were common among each other, suggesting that markers of cancer cells might be implicated in upregulating common markers of different T-cells through various pathways which are essential for these T-cells.

Additionally, the unique markers of T-cells indicated that cancer cells also influence those markers of T-cells which are unique to respective T-cell types, to evade the immune response through T-cell exhaustion. The expression of cancer cells and T-cells hub-genes across different cell types exhibited in dot plots, violin plots, heatmaps, and box plots are shown in **Figures 4-6**.

Table 1. The top 10 hub genes of common cancer cell hub genes among "cancer cells_vs_all-PDAC" and "cancer-PDAC_vs_all-normal" groups and unique cancer cell hub genes in both groups along with their number of interactions

Common cancer cell hub genes among both groups	
Hub Genes	**Number of Interactions**
GAPDH	364
AKT1	284
EGFR	242
CS	212
RHOA	207
TPI1	200
SDHA	189
TFRC	189

FASN	186
HIF1A	179
Unique cancer cell genes of "cancer cells_vs_all-PDAC" group	
Hub Genes	**Number of Interactions**
H4C6	29
MYC	28
H3C12	18
DDX21	16
USP7	16
RFC4	12
APEX1	12
CDK9	12
H2BC9	12
NOP2	12
Unique cancer cell genes of "cancer-PDAC_vs_all-normal" group	
Hub Gene	**Number of Interactions**
FN1	62
COL1A1	59
COL1A2	56

COL3A1	55
COL5A2	51
COL6A1	50
COL5A1	50
BGN	48
COL6A2	46
FBN1	45

Table 2. The top 10 hub genes of CD8+ NKT-like cells, memory CD4+ T cells, and naive CD4+ T cells along with their number of interactions

CD8+ NKT-like Cells Hub Genes	
Hub Gene	**Number of Interactions**
TP53	176
FN1	170
MMP9	149
CD4	147
IFNG	147
NFKB1	142
HIF1A	129
HSP90AA1	126

ITGB1	119
HSP90AB1	116

Memory CD4+ T Cells Hub Genes	
Hub Gene	**Number of Interactions**
AKT1	200
TP53	192
ACTB	180
CD4	148
JUN	146
FN1	142
MMP9	126
HSP90AA1	123
MAPK3	121
HSP90AB1	113

Naive CD4+ T Cells Hub Genes	
Hub Gene	**Number of Interactions**
TP53	452
UBC	298
UBB	279

HSP90AA1	270
JUN	260
CTNNB1	260
NFKB1	251
HSP90AB1	238
HSPA8	229
H3-3B	212

Figure 3. The analyses showing PPI and hub genes identification of cancer and T-cells **(a-f)** The PPI networks of selected pathway genes from cancer and T-cells, **(g-l)** Hub-genes identification of the cancer and T-cells from respective PPI networks

Figure 4. Expression levels of all cancer cells hub-genes shown in dot plots, violin plots, box plots across conditions and cancer cells; and heatmap, **(a-b)** dot plots of cancer cell hub genes showing expression levels in different cell types, **(c-d)** violin plots of cancer cell hub genes showing expression levels in different cell types, **(e-f)** box plots of cancer cell hub genes showing expression levels in cancer cells, **(g-h)** box plots of cancer cell hub genes showing expression levels in conditions, **(i)** heatmap of cancer cell hub genes showing expression levels in different cell types

Figure 5. Expression levels of T-cells hub genes **(a-d)** Expression levels of CD8+ NKT-like cells hub genes shown in dot plot, violin plot, and box plots across conditions, **(e-h)** expression levels of memory CD4+ T-cells hub genes shown in dot plot, violin plot, and box plots across conditions

Figure 6. Expression levels of T-cells hub genes **(a-d)** Expression levels of naive CD4+ T-cells shown in dot plot, violin plot, and box plots across conditions, **(e)** Heatmap of all T-cells hub-genes showing expression in respective T-cell and other cell types

Expression profiling and survival correlation of key hub-genes

The GEPIA2 was used for the expression validation and survival analysis of cancer cell hub genes as it contains TCGA bulk-RNA datasets, while TISCH2 was utilised for the expression validation of T-cells and cancer cells hub genes as TISCH2 contains the scRNA-seq datasets. Moreover, the GEPIA2 showed all cancer cells hub-genes to be highly upregulated in the tumor compared to the control samples. The expression values of all the hub genes of cancer cells in tumor and control samples are mentioned in **Table 3**.

Furthermore, the survival analysis of the cancer cells hub-genes indicated that *GAPDH, AKT1, CS, RHOA, TPI1, SDHA, FASN, HIF1A, FN1, COL1A1, COL1A2, COL3A1, COL5A2, COL5A1, BGN, COL6A2, FBN1, USP7, CDK9, H2BC9,* and *NOP2* exhibited insignificant p-values, while *EGFR, TFRC, COL6A1, MYC, DDX21, RFC4,* and *APEX1* exhibited significant p-values including, 0.03, 0.04, 0.04, 0.01, 0.01, 0.04, and 0.04, respectively. Lastly, GEPIA2 showed no results for the overall survival analysis of *H4C6* and *H3C12*.

Subsequently, the significant hub-genes, including *EGFR* showed that high expression exhibited ~18% of patients survival for more than 60 months while low expression showed ~20% of patients survived for approximately more than 90 months. Moreover, the high expression of *TFRC* showed survival of ~18% patients for more than 70 months, while low expression showed the survival of ~18% patients for more than 90 months. Furthermore, the high expression of *COL6A1* indicated that none of the patients survived more than 70 months, while ~37% of the low expression patients survived for more than 90 months.

The high expression of the MYC showed ~11% of patients survived for more than 70 months, while ~29% of low expression patients survived for more than 90 months. The high expression of the *DDX21* showed that none of the patients survived more than 75 months, while ~38% of the low expression patients survived more than 90 months. The high expression of *RFC4* showed ~13% of patients survived for ~80 months, while ~32% of patients survived for more than 90 months. Lastly, the high expression of *APEX1* showed that none of the patients survived for more than 75 months, while ~35% of patients survived for more than 90 months. The overall survival plots of all the cancer cells hub-genes are shown in **Figure 7**.

Additionally, the TISCH2 indicated that all the hub-genes of cancer cells were upregulated in malignant cells, especially *GAPDH, RHOA, TPI1, H4C6, DDX21,* and *APEX1* showed considerable high expression levels in malignant cells. Moreover, the hub-genes of T-cells indicated the upregulation of all hub-genes, especially *ACTB, H3F3B, HSP90AA1, HSP90AB1, HSPA8, UBB,* and *UBC* exhibited high levels of expressions in CD8Tex cells. The feature plots of cancer cells and T-cells hub genes from TISCH2 are shown in **Supplementary Figure S3-4**. The violin plots of cancer cells and T-cells hub-genes are shown in **Figure 8**.

This indicated that the hub-genes of cancer cells and T-cells showed high expressions in datasets from GEPIA2 and TISCH2. However, the insignificant p-values in overall survival rate of the cancer cell hub-genes further need corroborations as these genes are well-reported to be expressed in PDAC patients leading to poor prognosis.

Table 3. Expression levels of cancer cell hub-genes in tumor and control samples mentioned in transcripts per million (TPM) using GEPIA2

Common Cancer Cell Hub-Genes	Tumor (TPM)	Normal (TPM)
GAPDH	2426.56	232.4
AKT1	80.5	34.08
EGFR	9.41	5.02
CS	58.68	34.33
RHOA	358.32	73.54
TPI1	461.79	92.66
SDHA	72.19	25.05
TFRC	24.72	7.09
FASN	43.01	13.89
HIF1A	65.46	9
cancer cells_vs_all-PDAC Hub-Genes	**Tumor (TPM)**	**Normal (TPM)**
H4C6	0.17	0
MYC	39.26	25.79
H3C12	0.07	0
DDX21	24.08	10.96
USP7	34.27	18.08
RFC4	13.18	5.83
APEX1	105.45	43.57
CDK9	38.76	21.68
H2BC9	0.35	0
NOP2	21.52	11.05
cancer-PDAC_vs_all-normal Hub-Genes	**Tumor (TPM)**	**Normal (TPM)**
FN1	978.95	17.23
COL1A1	1201.65	12.61
COL1A2	1166.36	18.25
COL3A1	1103.06	13.04
COL5A2	71.61	1.71
COL6A1	179.95	35.78
COL5A1	115.94	2.59

BGN	613.07	22.19
COL6A2	277.8	30.05
FBN1	56.58	2.7

Overall Survival Rate of Cancer cells Hub-Genes

Figure 7. The overall survival rate of all significant and insignificant cancer cells hub-genes, indicating the high and low expression of respective hub-gene, leading to overall survival rate in months

Expression of Cancer and T-cells Hub-Genes in TISCH2

Figure 8. Violin plots of cancer and T-cells hub-genes showing expression levels in malignant and CD8Tex cells, along with other cell types within PDAC TIME

Discussion

There is a high rate of recurrence and metastasis in PDAC patients after surgery, lacking efficient chemotherapy, radiotherapy, and immunotherapy (33). T-cell exhaustion, which is induced by the complex immunosuppressive TIME, may contribute to diminished responses of immunotherapy in PDAC patients (34). Therefore, this study aimed to identify the upregulated markers of cancer cells and T-cells implicated in the PDAC progression through T-cell exhaustion by utilising scRNA-seq analysis.

Subsequently, this study revealed enriched pathways of cancer cells implicated in crucial biological processes for maintaining cellular homeostasis, including lipid and protein metabolism, post-translational modifications, and membrane trafficking (35–37). Moreover, pathways such as Regulation of gene expression, RNA metabolism, and DNA repair are essential for efficient cell function and survival (38–40). This suggests that the cancer cells pathways collectively play a vital role in maintaining cellular integrity, communication, and responses to external signals, leading to cancer cells progression by evading the immune system.

Furthermore, the enriched pathways of CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells were implicated in gene regulation, cellular communication, and immune responses. The pathways, including Receptor tyrosine kinases and MET signaling pathways impact immune responses and cell movement by regulating cell motility, growth and survival (41,42). Additionally, the immune related pathways such as neutrophil degranulation and cytokine signaling were highly expressed in T-cells (43,44). This indicated that the high expression of immune related pathways might be leading to T-cell exhaustion, rendering them non-functional against cancer cells.

Notably, all three sub-groups (common, cancer cells_vs_all-PDAC, and cancer-PDAC_vs_all-normal) of cancer cells exhibiting hub-genes, indicated the heterogeneity and complexity of cancer cells within the TIME. Moreover, all the hub-genes, including *FN1, COL1A1, COL1A2, COL3A1, COL5A2, COL6A1, COL5A1, BGN, COL6A2,* and *FBN1* of "cancer-PDAC_vs_all-normal" group are reported to be implicated in PDAC patients leading to poor prognosis (45–51).

Similarly, the common hub-genes such as *GAPDH, AKT1, EGFR, RHOA, TPI1, SDHA, TFRC, FASN,* and *HIF1A* are reported to be implicated in oncogenic pathways leading to cancer cells proliferation and progression, playing crucial role in PDAC (52–60), while *CS* is not specifically reported in PDAC, resulting in a novel PDAC cancer cell marker.

Furthermore, the *MYC, DDX21, USP7, RFC4, APEX1, CDK9,* and *NOP2* of "cancer cells_vs_all-PDAC" group play critical role in the metabolic reprogramming of the PDAC, contributing to poor prognosis of PDAC (61–67). However, *H4C6, H3C12*, and *H2BC9* are not reported in cancer cells of PDAC patients and these novel markers might be implicated in cell proliferation and survival of cancer cells by sending signals, leading to exhaustion of T-cells.

Additionally, among the common hub-genes of all three or any two T-cell subtypes (CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells), *CD4* is reported to be activated in T-effs and strongly promote EMT-associated alterations in H6c7 cells and increased invasive behaviour (68). NFKB1 within the NFKB signaling is influenced in cytotoxic T-cells in the blood of patients with PDAC (69). *HSP90AA1* has been reported to be expressed in a new type of T-cell (HSP T or Thsp) within the PDAC environment (70), while *JUN* is reported to be implicated in the T-cell exhaustion mechanism within the PDAC environment (71).

However, *TP53, MMP9*, *FN1,* and *HSP90AB1* were observed to be novel markers as these markers are not specifically reported in CD8+ NKT-like cells, memory CD4+ T-cells, naive CD4+ T-cells or any related T-cell within PDAC patients, indicating the involvement of these novel markers in T-cell exhaustion and aberrant functional activity within PDAC TIME, eventually leading to PDAC cells proliferation and survival.

Moreover, the unique hub-genes of CD8+ NKT-like cells showed that *IFNG* and *HIF1A* are reported to be implicated in T-cells (72,73), while *ITGB1* turned out to be a novel marker of CD8+ NKT-like cells. Furthermore, the unique hub-genes of memory CD4+ T-cells such as *AKT1, ACTB*, and *MAPK3* were observed to be novel markers as these are not reported specifically in T-cells of PDAC patients. Lastly, the unique hub-genes of naive CD4+ T-cells only *HSPA8* is reported to be expressed in CD8+ T-cells in prostate cancer (74), while *UBC, UBB, CTNNB1*, and *H3-3B* are not reported, rendering them novel markers for naive CD4+ T-cells. This suggests that the aforementioned novel markers are potentially implicated in exhaustion of T-cells within TIME, eventually leading to cancer cells growth, proliferation, and survival by evading the immune responses in PDAC patients.

Subsequently, the expression and survival analysis of the cancer cells and T-cells hub-genes exhibited majority of the cancer cells overall survival to be insignificant (p-value > 0.05); however, aforementioned studies support the aberrant expression of these markers to be implicated in poor prognosis in PDAC patients. Additionally, further corroboration is needed,

especially for novel markers of cancer cells and T-cells within PDAC patients for diagnostic, prognostic, and therapeutic interventions.

Recent scRNA-seq studies on T-cells within PDAC patients have reported T-cell marker genes in CD8+ Tcm, CD4+ Tem, and γδT cells; and reported that the *CCL5/SDC1* receptor-ligand interactions in tumor infiltrating T-cells could promote tumor cells migration (75,76). However, this study reveals novel biomarkers of cancer cells, CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells implicated in T-cell exhaustion, rendering the growth, proliferation and progression of cancer cells.

In summary, the aforementioned novel biomarkers of cancer cells and T-cells might be involved in the complex TIME, leading to the poor prognosis of PDAC. These findings underscore the complexity and heterogeneity of the TIME and its influence on immune evasion and cancer cell survival. While existing studies support the role of some identified markers in poor prognosis, further clinical validation is required for PDAC management.

Conclusion

This study revealed novel biomarkers of cancer cells (*CS, H4C6, H3C12,* and *H2BC9*) and T-cells (*TP53, FN1, MMP9, ITGB1, HSP90AB1, AKT1, ACTB, MAPK3, UBC, UBB, CTNNB1,* and *H3-3B*), including CD8+ NKT-like cells, memory CD4+ T-cells, and naive CD4+ T-cells, which might be the key candidate biomarkers implicated in cancer cells progression through T-cell exhaustion. Further clinical investigations are needed to evaluate these novel biomarkers as potential therapeutic options in PDAC patients.

References

1. Schawkat K, Manning MA, Glickman JN, Mortele KJ. Pancreatic Ductal Adenocarcinoma and Its Variants: Pearls and Perils. RadioGraphics. 2020 Sep;40(5):1219–39.
2. Mukund A, Afridi MA, Karolak A, Park MA, Permuth JB, Rasool G. Pancreatic Ductal Adenocarcinoma (PDAC): A Review of Recent Advancements Enabled by Artificial Intelligence. Cancers. 2024 Jun 17;16(12):2240.
3. Taherian M, Wang H, Wang H. Pancreatic Ductal Adenocarcinoma: Molecular Pathology and Predictive Biomarkers. Cells. 2022 Sep 29;11(19):3068.
4. Espiau-Romera P, Courtois S, Parejo-Alonso B, Sancho P. Molecular and Metabolic Subtypes Correspondence for Pancreatic Ductal Adenocarcinoma Classification. J Clin Med. 2020 Dec 21;9(12):4128.
5. Nakaoka K, Ohno E, Kawabe N, Kuzuya T, Funasaka K, Nakagawa Y, et al. Current Status of the Diagnosis of Early-Stage Pancreatic Ductal Adenocarcinoma. Diagnostics [Internet]. 2023 Jan [cited 2024 Oct 16];13(2). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9857526/
6. Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. CA Cancer J Clin. 2024;74(1):12–49.
7. Zhao Z, Liu W. Pancreatic Cancer: A Review of Risk Factors, Diagnosis, and Treatment. Technol Cancer Res Treat. 2020 Dec 24;19:1533033820962117.
8. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. Cancer Res. 2014 Jun 1;74(11):2913–21.
9. Seeber A, Zimmer K, Kocher F, Puccini A, Xiu J, Nabhan C, et al. Molecular characteristics of *BRCA1/2* and *PALB2* mutations in pancreatic ductal adenocarcinoma. ESMO Open. 2020 Jan 1;5(6):e000942.
10. Singhi AD, Koay EJ, Chari ST, Maitra A. Early Detection of Pancreatic Cancer: Opportunities and Challenges. Gastroenterology. 2019 Feb 2;156(7):2024.
11. Park W, Chawla A, O'Reilly EM. Pancreatic Cancer: A Review. JAMA. 2021 Sep 7;326(9):851.
12. Anderson EM, Thomassian S, Gong J, Hendifar A, Osipov A. Advances in Pancreatic Ductal Adenocarcinoma Treatment. Cancers. 2021 Jan;13(21):5510.
13. Digiacomo G, Volta F, Garajova I, Balsano R, Cavazzoni A. Biological Hallmarks and New Therapeutic Approaches for the Treatment of PDAC. Life. 2021 Aug;11(8):843.
14. Royal RE, Levy C, Turner K, Mathur A, Hughes M, Kammula US, et al. Phase 2 Trial of Single Agent Ipilimumab (Anti-CTLA-4) for Locally Advanced or Metastatic Pancreatic Adenocarcinoma. J Immunother Hagerstown Md 1997. 2010 Oct;33(8):828–33.
15. Kunk PR, Bauer TW, Slingluff CL, Rahma OE. From bench to bedside a comprehensive review of pancreatic cancer immunotherapy. J Immunother Cancer. 2016 Mar 15;4:14.
16. Chouari T, La Costa FS, Merali N, Jessel MD, Sivakumar S, Annels N, et al. Advances in Immunotherapeutics in Pancreatic Ductal Adenocarcinoma. Cancers. 2023 Aug 25;15(17):4265.
17. Saka D, Gökalp M, Piyade B, Cevik NC, Arik Sever E, Unutmaz D, et al. Mechanisms of T-Cell Exhaustion in Pancreatic Cancer. Cancers. 2020 Aug 14;12(8):2274.
18. Jiang Y, Li Y, Zhu B. T-cell exhaustion in the tumor microenvironment. Cell Death Dis. 2015 Jun;6(6):e1792–e1792.
19. Schietinger A, Greenberg PD. Tolerance and Exhaustion: Defining Mechanisms of T cell Dysfunction. Trends Immunol. 2013 Nov 6;35(2):51.
20. Papalampros A, Vailas M, Ntostoglou K, Chiloeches ML, Sakellariou S, Chouliari NV, et al. Unique Spatial Immune Profiling in Pancreatic Ductal Adenocarcinoma with Enrichment of Exhausted and Senescent T Cells and Diffused CD47-SIRPα Expression. Cancers. 2020 Jul 7;12(7):1825.
21. Burles A Johnson III, Yarchoan M, Lee V, Laheru DA, Jaffee EM. Strategies for Increasing Pancreatic Tumor Immunogenicity. Clin Cancer Res Off J Am Assoc Cancer

Res. 2017 Apr 1;23(7):1656.

22. Balachandran VP, Łuksza M, Zhao JN, Makarov V, Moral JA, Remark R, et al. Identification of unique neoantigen qualities in long term pancreatic cancer survivors. Nature. 2017 Nov 8;551(7681):512.

23. Goulart MR, Stasinos K, Fincham REA, Delvecchio FR, Kocher HM. T cells in pancreatic cancer stroma. World J Gastroenterol. 2021 Dec 14;27(46):7956.

24. Sivakumar S, Abu-Shah E, Ahern DJ, Arbe-Barnes EH, Jainarayanan AK, Mangal N, et al. Activated Regulatory T-Cells, Dysfunctional and Senescent T-Cells Hinder the Immunity in Pancreatic Cancer. Cancers. 2021 Apr 8;13(8):1776.

25. Werba G, Weissinger D, Kawaler EA, Zhao E, Kalfakakou D, Dhara S, et al. Single-cell RNA sequencing reveals the effects of chemotherapy on human pancreatic adenocarcinoma and its tumor microenvironment. Nat Commun. 2023 Feb 13;14(1):797.

26. Clough E, Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, et al. NCBI GEO: archive for gene expression and epigenomics data sets: 23-year update. Nucleic Acids Res. 2024 Jan 5;52(D1):D138–44.

27. Nader K, Tasci M, Ianevski A, Erickson A, Verschuren EW, Aittokallio T, et al. ScType enables fast and accurate cell type identification from spatial transcriptomics data. Bioinformatics. 2024 Jul 1;40(7):btae426.

28. García-Moreno A, López-Domínguez R, Ramirez-Mena A, Pascual-Montano A, Aparicio-Puerta E, Hackenberg M, et al. GeneCodis 4: Expanding the modular enrichment analysis to regulatory elements [Internet]. bioRxiv; 2021 [cited 2024 Oct 6]. p. 2021.04.15.439962. Available from: https://www.biorxiv.org/content/10.1101/2021.04.15.439962v1

29. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2022 Nov 12;51(D1):D638–46.

30. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. Genome Res. 2003 Nov;13(11):2498–504.

31. Tang Z, Kang B, Li C, Chen T, Zhang Z. GEPIA2: an enhanced web server for large-scale expression profiling and interactive analysis. Nucleic Acids Res. 2019 Jul 2;47(W1):W556–60.

32. Han Y, Wang Y, Dong X, Sun D, Liu Z, Yue J, et al. TISCH2: expanded datasets and new tools for single-cell transcriptome analyses of the tumor microenvironment. Nucleic Acids Res. 2023 Jan 6;51(D1):D1425–31.

33. Hm K, S C, Gp W, M C. Pancreatic Cancer: A Review of Current Treatment and Novel Therapies. J Investig Surg Off J Acad Surg Res [Internet]. 2023 Dec 31 [cited 2024 Oct 11];36(1). Available from: https://pubmed.ncbi.nlm.nih.gov/36191926/?dopt=Abstract

34. Tasiheng Y, Lin X, Wang X, Zou X, Chen Y, Yan Y, et al. DNA hypo-methylation and expression of GBP4 induces T cell exhaustion in pancreatic cancer. Cancer Immunol Immunother. 2024 Aug 7;73(10):208.

35. Encarnación-Rosado J, Kimmelman AC. Harnessing metabolic dependencies in pancreatic cancers. Nat Rev Gastroenterol Hepatol. 2021 Jul;18(7):482–92.

36. Schneeweis C, Hassan Z, Schick M, Keller U, Schneider G. The SUMO pathway in pancreatic cancer: insights and inhibition. Br J Cancer. 2021 Feb 2;124(3):531–8.

37. Guo H, Zhou C, Zheng M, Zhang J, Wu H, He Q, et al. Insights into the role of derailed endocytic trafficking pathway in cancer: From the perspective of cancer hallmarks. Pharmacol Res. 2024 Mar 1;201:107084.

38. Stoof J, Harrold E, Mariottino S, Lowery MA, Walsh N. DNA Damage Repair Deficiency in Pancreatic Ductal Adenocarcinoma: Preclinical Models and Clinical Perspectives. Front Cell Dev Biol [Internet]. 2021 Oct 12 [cited 2024 Oct 11];9. Available from: https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2021.749490/full

39. Venkat S, Alahmari AA, Feigin ME. Drivers of gene expression dysregulation in

pancreatic cancer. Trends Cancer. 2021 Jul;7(7):594–605.

40.   Ali A, Chianese U, Papulino C, Toraldo A, Abakar MEA, Passaro E, et al. Metabolic Pathways as a Novel Landscape in Pancreatic Ductal Adenocarcinoma. Cancers. 2022 Aug 4;14(15):3799.

41.   Im K, Choi YJ, Kim DH, Kim DS, Ban K, Ji W, et al. AXL receptor tyrosine kinase inhibition improves the anti-tumor effects of CD8+ T cells by inducing CD103+ dendritic cell-mediated T cell priming. Biochem Biophys Res Commun. 2023 Nov 5;680:7–14.

42.   Shen Z, Xue W, Zheng Y, Geng Q, Wang L, Fan Z, et al. Molecular mechanism study of HGF/c-MET pathway activation and immune regulation for a tumor diagnosis model. Cancer Cell Int. 2021 Jul 14;21(1):374.

43.   Hoekstra ME, Vijver SV, Schumacher TN. Modulation of the tumor micro-environment by CD8+ T cell-derived cytokines. Curr Opin Immunol. 2021 Apr 13;69:65–71.

44.   Emmons TR, Giridharan T, Singel KL, Khan ANH, Ricciuti J, Howard K, et al. Mechanisms driving neutrophil-induced T-cell immunoparalysis in ovarian cancer. Cancer Immunol Res. 2021 Jul;9(7):790–810.

45.   Hu D, Ansari D, Zhou Q, Sasor A, Said Hilmersson K, Andersson R. Stromal fibronectin expression in patients with resected pancreatic ductal adenocarcinoma. World J Surg Oncol. 2019 Feb 8;17(1):29.

46.   The collagen landscape in cancer: profiling collagens in tumors and in circulation reveals novel markers of cancer- associated fibroblast subtypes - Thorlacius- Ussing - 2024 - The Journal of Pathology - Wiley Online Library [Internet]. [cited 2024 Oct 10]. Available from: https://pathsocjournals.onlinelibrary.wiley.com/doi/full/10.1002/path.6207

47.   Mansouri V, Arjmand B, Hamzeloo-Moghadam M, Razzaghi Z, Ahmadzadeh A, Ehsani Ardakani MJ, et al. Extracellular matrix is the main targeted environment in early stage of pancreatic ductal adenocarcinoma. Gastroenterol Hepatol Bed Bench. 2023;16(4):401–7.

48.   Maneshi P, Mason J, Dongre M, Öhlund D. Targeting Tumor-Stromal Interactions in Pancreatic Cancer: Impact of Collagens and Mechanical Traits. Front Cell Dev Biol [Internet]. 2021 Nov 25 [cited 2024 Oct 10];9. Available from: https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2021.787485/full

49.   Pan S, Chen R, Tamura Y, Crispin DA, Lai LA, May DH, et al. Quantitative Glycoproteomics Analysis Reveals Changes in N-Glycosylation Level Associated with Pancreatic Ductal Adenocarcinoma. J Proteome Res. 2014 Mar 7;13(3):1293–306.

50.   Collagen VI expression is negatively mechanosensitive in pancreatic cancer cells and supports the metastatic niche | Journal of Cell Science | The Company of Biologists [Internet]. [cited 2024 Oct 10]. Available from: https://journals.biologists.com/jcs/article/135/24/jcs259978/286155/Collagen-VI-expression-is-negatively

51.   Identification of Key Prognostic Biomarker and Its Correlation with Immune Infiltrates in Pancreatic Ductal Adenocarcinoma - Luan - 2020 - Disease Markers - Wiley Online Library [Internet]. [cited 2024 Oct 10]. Available from: https://onlinelibrary.wiley.com/doi/full/10.1155/2020/8825997

52.   Pacchiana R, Mullappilly N, Pinto A, Bova S, Forciniti S, Cullia G, et al. 3-Bromo-Isoxazoline Derivatives Inhibit GAPDH Enzyme in PDAC Cells Triggering Autophagy and Apoptotic Cell Death. Cancers. 2022 Jan;14(13):3153.

53.   Qi D, Song X, Xue C, Yao W, Shen P, Yu H, et al. AKT1/FOXP3 axis-mediated expression of CerS6 promotes p53 mutant pancreatic tumorigenesis. Cancer Lett. 2021 Dec 1;522:105–18.

54.   Handra-Luca A, Hammel P, Sauvanet A, Lesty C, Ruszniewski P, Couvelard A. EGFR expression in pancreatic adenocarcinoma. Relationship to tumour morphology and cell adhesion proteins. J Clin Pathol. 2014 Apr 1;67(4):295–300.

55.   Della Peruta M, Giagulli C, Laudanna C, Scarpa A, Sorio C. RHOA and PRKCZ control different aspects of cell motility in pancreatic cancer metastatic clones. Mol

Cancer. 2010 Mar 17;9(1):61.

56. Toyoda T, Miura N, Kato S, Masuda T, Ohashi R, Matsushita A, et al. Identification of TPI1 As a potential therapeutic target in pancreatic cancer with dependency of mutation using multi-omics analysis. Cancer Sci [Internet]. [cited 2024 Oct 11];n/a(n/a). Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/cas.16302

57. Frances A, Lumeau A, Bery N, Gayral M, Stuani L, Sorbara M, et al. Cytidine deaminase-dependent mitochondrial biogenesis as a potential vulnerability in pancreatic cancer cells. Commun Biol. 2024 Aug 30;7(1):1–11.

58. García-Montero C, Fraile-Martinez O, Cobo-Prieto D, De Leon-Oliva D, Boaru DL, De Castro-Martinez P, et al. Abnormal Histopathological Expression of Klotho, Ferroptosis, and Circadian Clock Regulators in Pancreatic Ductal Adenocarcinoma: Prognostic Implications and Correlation Analyses. Biomolecules. 2024 Aug;14(8):947.

59. Bian Y, Yu Y, Wang S, Li L. Up-regulation of fatty acid synthase induced by EGFR/ERK activation promotes tumor growth in pancreatic cancer. Biochem Biophys Res Commun. 2015 Aug 7;463(4):612–7.

60. Ma C, Guo Y, Zhang Y, Duo A, Jia Y, Liu C, et al. PAFAH1B2 is a HIF1a target gene and promotes metastasis in pancreatic cancer. Biochem Biophys Res Commun. 2018 Jun 27;501(3):654–60.

61. Zheng L, Li M, Wei J, Chen S, Xue C, Duan Y, et al. NOP2/Sun RNA methyltransferase 2 is a potential pan-cancer prognostic biomarker and is related to immunity. PLOS ONE. 2023 Sep 28;18(9):e0292212.

62. Kretz AL, Schaum M, Richter J, Kitzig EF, Engler CC, Leithäuser F, et al. CDK9 is a prognostic marker and therapeutic target in pancreatic cancer. Tumor Biol. 2017 Feb 1;39(2):1010428317694304.

63. Trilla-Fuertes L, Gámez-Pozo A, Lumbreras-Herrera MI, López-Vacas R, Heredia-Soto V, Ghanem I, et al. Identification of Carcinogenesis and Tumor Progression Processes in Pancreatic Ductal Adenocarcinoma Using High-Throughput Proteomics. Cancers. 2022 Jan;14(10):2414.

64. Yu L, Li J, Zhang M, Li Y, Bai J, Liu P, et al. Identification of RFC4 as a potential biomarker for pan- cancer involving prognosis, tumour immune microenvironment and drugs. J Cell Mol Med. 2024 Jun 21;28(12):e18478.

65. Chen H, Zhu X, Sun R, Ma P, Zhang E, Wang Z, et al. Ubiquitin-specific protease 7 is a druggable target that is essential for pancreatic cancer growth and chemoresistance. Invest New Drugs. 2020 Dec 1;38(6):1707–16.

66. Wu S, Sun X, Hua R, Hu C, Qin L. DDX21 functions as a potential novel oncopromoter in pancreatic ductal adenocarcinoma: a comprehensive analysis of the DExD box family. Discov Oncol. 2024 Aug 3;15(1):333.

67. Sodir NM, Kortlever RM, Barthet VJA, Campos T, Pellegrinet L, Kupczak S, et al. MYC Instructs and Maintains Pancreatic Adenocarcinoma Phenotype. Cancer Discov. 2020 Apr 1;10(4):588–607.

68. Full article: CD4+ T cells potently induce epithelial-mesenchymal-transition in premalignant and malignant pancreatic ductal epithelial cells–novel implications of CD4+ T cells in pancreatic cancer development [Internet]. [cited 2024 Oct 11]. Available from: https://www.tandfonline.com/doi/full/10.1080/2162402X.2014.1000083#d1e959

69. Ali LR, Lenehan PJ, Cardot-Ruffino V, Dias Costa A, Katz MHG, Bauer TW, et al. PD-1 Blockade Induces Reactivation of Nonproductive T-Cell Responses Characterized by NF-κB Signaling in Patients with Pancreatic Cancer. Clin Cancer Res. 2024 Feb 1;30(3):542–53.

70. Immune profiling and prognostic model of pancreatic cancer using quantitative pathology and single-cell RNA sequencing | Journal of Translational Medicine [Internet]. [cited 2024 Oct 11]. Available from: https://link.springer.com/article/10.1186/s12967-023-04051-4

71. Saka D, Gökalp M, Piyade B, Cevik NC, Arik Sever E, Unutmaz D, et al. Mechanisms of T-Cell Exhaustion in Pancreatic Cancer. Cancers. 2020 Aug;12(8):2274.

72. Palazon A, Tyrakis PA, Macias D, Veliça P, Rundqvist H, Fitzpatrick S, et al. An HIF-

1α/VEGF-A Axis in Cytotoxic T Cells Regulates Tumor Progression. Cancer Cell. 2017 Nov 13;32(5):669-683.e5.

73. Pearce H, Croft W, Nicol SM, Margielewska-Davies S, Powell R, Cornall R, et al. Tissue-Resident Memory T Cells in Pancreatic Ductal Adenocarcinoma Coexpress PD-1 and TIGIT and Functional Inhibition Is Reversible by Dual Antibody Blockade. Cancer Immunol Res. 2023 Apr 3;11(4):435–49.

74. Extracellular vesicles related gene HSPH1 exerts anti-tumor effects in prostate cancer via promoting the stress response of CD8 + T cells | Cellular Oncology [Internet]. [cited 2024 Oct 11]. Available from: https://link.springer.com/article/10.1007/s13402-023-00905-7

75. Zheng H, Li Y, Zhao Y, Jiang A. Single-cell and bulk RNA sequencing identifies T cell marker genes score to predict the prognosis of pancreatic ductal adenocarcinoma. Sci Rep. 2023 Mar 6;13(1):3684.

76. Chen K, Wang Y, Hou Y, Wang Q, Long D, Liu X, et al. Single cell RNA-seq reveals the CCL5/SDC1 receptor-ligand interaction between T cells and tumor cells in pancreatic cancer. Cancer Lett. 2022 Oct 1;545:215834.

Supplementary Document

Figure S1. ScRNA-seq dataset preprocessing steps showing **(a)** filtered cells based on number of features, RNA counts, and mitochondrial reads; **(b)** 2000 highly variable features with labelled top-10 features; **(c)** PCA plot; **(d)** Elbow plot showing the variable principle components; and **(e)** t-SNE map of the clusters identified at resolution of 0.6

Figure S2. Expression plots of top-10 cancer and T-cells genes showing **(a-b)** dot plot and heatmap of the top-10 cancer cell genes, respectively; **(c)** dot plot of top-10 T-cells genes; **(d)** dot plot of top-10 T-cells genes in individual PDAC and normal samples, indicating slight expression of genes within Naive CD4+ T-cells, leading to overall low expression, **(e)** as also seen in heatmap; **(f)** Venn diagram of cancer cells genes of "cancer-PDAC_vs_all-normal" and "cancer cells_vs_all-PDAC" groups, indicating the common and unique genes among both groups

Figure S3. Feature plots of each group of cancer cells hub-genes, showing expression levels of each gene in different cell types, including malignant cells

Figure S4. Feature plots of T-cells hub-genes showing the expression levels of each gene in different cell types, including CD8Tex cells