1

Predator-Induced Tail Coloration Toward Diverse Dragonfly Nymphs in Tadpoles of

the Japanese Tree Frog (*Dryophytes japonicus*)

Akihiro Noda[1,*] and Katsutoshi Watanabe[1]

1. Department of Zoology, Graduate School of Science, Kyoto University,

Kitashirakawa-oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan.

* Corresponding author. E-mail: noda@terra.zool.kyoto-u.ac.jp

Abstract

Anuran larvae can change their coloration and morphology to avoid predation.

Particularly, some tadpoles in the genus *Dryophytes* in the Americas develop deep tail fins and

bright orange tail coloration in response to predators such as dragonfly nymphs. These

conspicuous tails are hypothesized to attract predator attacks, protecting vital body parts from

fatal injuries. However, it remains unclear whether the extent of the induced phenotypic

changes differs between various predators with distinct feeding patterns and habitats. To

explore this, we reared tadpoles of the Japanese tree frog, *Dryophytes japonicus*, with four

dragonfly species, water bug, water scorpion, or newt, and measured phenotypic changes,

especially focusing on tail coloration. We found that the aeshnid *Anax nigrofasciatus*, a sit-

and-wait predator that climbs on aquatic vegetation, induced bright orange and broad tails of

tadpoles. Other odonate species that forage in and on the substrate also elicited tail coloration

or deeper larval tails, but it took more time to induce the tadpoles' response compared to *A.*

nigrofasciatus. In contrast, non-odonate predators did not induce color and morphological

changes. This study provides the first evidence of predator-induced tail coloration in an Asian

species of *Dryophytes*, which appears to be a defensive response specifically to dragonfly

nymphs, as visually adept predators. Moreover, the differences in the extent and timing of the

phenotypic changes among odonate species suggest that anti-predator phenotypes are

modulated based on relative predation risks, considering the predators' foraging patterns and

visual capabilities, thereby allowing tadpoles to minimize physiological costs and unintended

risks associated with the induced traits.

Introduction

The body color of organisms has diverse implications for predator–prey interactions. Many species exhibit cryptic body color, which suggests significant selective pressure from predators (Norris and Lowe, 1964). Cryptic coloration contributes to predation avoidance by representing background matching, countershading, and the disruption of body shape or outlines (Thibaudeau and Altig, 2012). In contrast, some organisms use bright colors and expendable body parts to divert predator attention away from more vulnerable areas (Hawlena et al., 2006). Consequently, body color contributes to predation avoidance by fulfilling either of two primary functions: enhancing prey crypticity or creating a defensive lure phenotype (Touchon and Warkentin, 2008).

Changes in body color are often induced by predators through phenotypic plasticity. The utilization of a prey's body coloration is highly dependent on its context, including the sensory capabilities of predators and the background color (e.g., Espanha et al., 2016). For example, although the poison dart frog (*Dendrobates tinctorius*) is well recognized for its typical aposematic signaling, its conspicuous coloration also functions as camouflage when viewed from a distance, blending with the background (Barnett et al., 2018). Especially in aquatic habitats, substantial variation exists in background conditions and the composition and abundance of predators across different localities, making it difficult for prey to achieve a uniform predation avoidance ability across their range (Van Buskirk, 2002; McIntyre et al., 2004). Therefore, such environmental uncertainty has generally been believed to favor plastic rather than constitutive anti-predator defenses (Agrawal and Karban, 1999; McIntyre et al., 2004).

Anuran larvae, or tadpoles, have served for decades as a model system for investigating induced defense mechanisms, as well as predator avoidance strategies and trade-offs mediated by plastic traits (Kasumovic, 2013; Kruger and Morin, 2020). Phenotypes such as body color and shape are generally subject to predator selection for survival and mate choice, but the latter can be ruled out in the case of tadpoles. They often rely on chemical cues

to evaluate predation risk through continually released predator-borne cues and digestion-

released, prey-borne cues, and then exhibit predator-induced defense mechanisms (Brönmark

and Hansson, 2000; Hettyey et al., 2010, 2015).

Predator-induced plastic responses in tadpoles can be categorized into four primary

domains: behavioral, morphological, physiological, and chemical defenses. Behavioral

responses encompass a reduction in overall activity (e.g., Lawler, 1989; Takahara et al.,

2012a), increased burst swimming speed (Takahara et al., 2008), and spatial avoidance of

predators (e.g., Laurila et al., 1998; Nunes et al., 2013). Morphological changes for predator

avoidance have been well documented in tadpoles of various species (Relyea, 2001; Van

Buskirk, 2002; Richter-Boix et al. 2007). For instance, tadpoles undergo changes in body size

(i.e., correlated with growth rate and metamorphosis timing) and shape, such as the

development of deeper tail fins (Lardner, 1998, 2000). Behavioral and morphological

responses are sometimes co-expressed (Hossie et al., 2017). Physiological responses include

changes in respiration rate and gut evacuation rate (Steiner, 2007; Steiner and Van Buskirk,

2009; Costello and Michel, 2013). These may arise due to trade-offs in predator avoidance

(Relyea and Auld, 2004) or as responses to partially offset the costs of lower energy intake

resulting from evasion (Barry and Syal, 2013). Additionally, tadpoles of certain groups,

particularly toads, evade predation through toxin production (Bókony et al., 2016; Hettyey et

al., 2019).

The types and intensities of induced responses in tadpoles often differ according to

predator types (Relyea, 2001; Teplitsky et al., 2004, 2005; Hettyey et al., 2011). For instance,

Kishida and Nishimura (2005) revealed that tadpoles of *Rana pirica* developed deeper tail

fins in response to dragonfly nymphs, whereas in the presence of salamander larvae, they

exhibited not only deeper tails but also bulging heads and bodies. Tadpoles expressing these

predator-specific traits should have a survival advantage over those with mismatched or non-

induced phenotypes when faced with predation. Conversely, expressing such defense traits

incurs energetic costs and can increase vulnerability to non-focal predators (Touchon and

Warkentin, 2008; Innes-Gold et al., 2019). Due to these costs, defense traits are expected to

remain plastic rather than fixed, and their types and intensities can vary depending on

predator species, sometimes exhibiting predator-specific responses.

Dragonfly nymphs, as one of the major predators of tadpoles, are known to induce

morphological responses characterized by deep tail fins across many anuran species (Relyea

and Werner, 2000; Van Buskirk, 2002). This induced broad tail has been repeatedly

hypothesized to improve the swimming performance of tadpoles and facilitate their escape

from predators (Smith and Van Buskirk, 1995; McCollum and Leimberger, 1997; Van Buskirk

and McCollum, 2000a). However, experimental evidence of this benefit is lacking (Van

Buskirk et al., 2003; Johnson et al., 2008). Moreover, a few tree frog species in the genus

Dryophytes respond to odonate predators by not only exhibiting deeper tails but also turning

their entire tails bright orange (McCollum and Van Buskirk, 1996; Van Buskirk and

McCollum, 1999; LaFiandra and Babbitt, 2004). Thus, as another function, the induced deep

and brightly colored fin is thought to serve as a lure, diverting predator attacks away from the

more vulnerable head and body regions (Van Buskirk et al., 2003, 2004; Johnson et al., 2008).

Given that tail fins tear easily (Doherty et al., 1998) and that tadpoles can maintain their

swimming performance even after partial tail loss (Hoff and Wassersug, 2000; Van Buskirk

and McCollim, 2000b), the tail is considered expendable relative to other body parts. Because

dragonfly nymphs are predominantly visual predators, exhibiting the ability to discern both

the shape and color of immobile as well as moving prey (Bybee et al., 2012; Espanha et al.,

2016), color and shape changes in tadpoles are likely to be effective in avoiding predation by

odonates.

Among dragonfly nymphs, several ecological types have been identified based on

microhabitat usage and foraging behavior (Pritchard, 1965; Corbet, 1999), including climbers

(e.g., Green darner, *Anax junius*), moving among aquatic vegetation and ambushing prey on

plants; sprawlers (e.g., Globe skimmer, *Pantala flavescens*; Sherratt and Harvey, 1989),

inhabiting pond bottoms and swimming through the water column to attack; and burrowers

(e.g., Golden-ringed dragonfly, *Cordulegaster boltonii*), burrowing in mud or bottom debris

and often lying in wait for prey with only the eyes protruding. The reliance on vision for

foraging varies among these types. For example, climbers that often have large compound

eyes rely extensively on vision for prey detection, while other types with smaller eyes may

rely more on tactile stimuli (Pritchard, 1966; Corbet, 1999). For tadpoles, the frequency of

encounters with and relative predation risk of these predators differ among habitats and

seasons. These differential predation risks likely determine the cost−benefit balance of

induced traits, thereby suggesting that tadpoles may exhibit different defense responses.

However, prior experimental studies have primarily focused on a single dragonfly species and

have not examined how tadpole responses may vary among odonate (and other) species.

The Japanese tree frog, *Dryophytes japonicus*, is a prevalent species with a wide

distribution encompassing entire regions of Japan (Honshu, Hokkaido, Shikoku, and Kyushu

islands), as well as regions outside Japan including Sakhalin Island, Jeju Island, and the

Korean Peninsula (Matsui and Maeda, 2018). Its breeding typically occurs from May to July

(in Honshu), with breeding pairs depositing small clutches of eggs at multiple sites. Tadpoles

of this species inhabit various environments such as rice paddies, marshes, ponds, puddles,

and other ephemeral water bodies (Matsui and Maeda, 2018). They are classified as

"nektonic" raspers, moving around the mid-water layer, characterized by distinctive body

shapes with tall tail fins that originate relatively anterior to the body (Orton, 1953). Tadpoles

of Japanese tree frog serve as important prey for numerous aquatic predators, including

dragonfly nymphs, water scorpions, and red-bellied newts (Ohba and Nakatsuji, 2006; Hirai

et al., 2020; Kita et al. 2020; Ozaki and Nishikawa, 2023). Morphological responses to

predators have also been documented in this species, with dragonfly nymphs inducing deep

tail fins, while chemical cues from goldfish prompt the development of shallow fins

(Takahara et al., 2012b). The occurrence of orange tail coloration has also been briefly

reported, although it has been attributed to ingested algae rather than predator-induced

plasticity (Doi et al., 2005; Yuasa, 2006, 2007). Despite being rarely documented, tadpoles

exhibiting brightly colored tails were frequently observed in our study site in Kyoto, central

Japan (Noda, personal observation).

144   To clarify the cause and function of the orange tail coloration in tadpoles of *D.*

japonicus, we conducted laboratory experiments with the tadpoles and several predator

species. These experiments aimed to achieve two primary objectives. First, we investigated

whether the tail coloration of tadpoles is induced by dragonfly nymphs, and if so, whether the

intensity and speed of this response vary among different odonate species with different

ecological characteristics. Simultaneously, we examined changes in body shape that are

known to be plastically induced by predators. The second objective was to assess whether

tadpoles exhibit color/morphological responses to predators characterized by a lower reliance

on vision for foraging compared to odonates. We used water bugs (*Appasus major*), water

scorpions (*Laccotrephes japonensis*), and newts (*Cynops pyrrhogaster*) as non-odonate

predators. For such predators, predation avoidance strategies involving tail coloration are

expected to be less effective than those against dragonfly nymphs. Thus, tadpoles may exhibit

weakened or no color/morphological responses to minimize the costs associated with

developing plastic traits. In both experiments, tadpoles were reared with a single predator

species, and their morphological traits of interest were compared with those of tadpoles raised

in predator-free tanks. By analyzing the extent of tail coloration in tadpoles exposed to diverse

predators, we revealed that tadpoles modulate their induced responses according to the risks

posed by predators with different foraging types and sensory capabilities.

Materials and Methods

165   Two series of laboratory experiments were carried out: Experiment I involved the

166   comparison of induction of color/morphological changes in tadpoles by four different

167   dragonfly species as predators, and Experiment II employed three non-odonate predators.

Experiment I

Adults of *Dryophytes japonicus* were captured in the paddy fields at Kyoto University (35°01'N, 135°47'E) on June 9 and 10, 2022. Following capture, mating pairs were housed overnight in plastic cages (19 × 19 × 14 cm; 6 cm water depth), and eggs were obtained the following day. After hatching, tadpoles were separately kept in plastic containers (43 × 31 × 16 cm; 6 cm water depth), as those from the same clutch were kept together in a single container. A room temperature of 20°C was maintained until the experiments began, and tadpoles were fed powdered food (Kyorin Co., Ltd.) every two days. In this experiment, four species of dragonfly nymphs were used as predators taking into account their foraging styles: *Anax nigrofasciatus* (Black-striped lesser emperor; climber), *Gynacantha japonica* (Blue-spotted dusk-hawker; climber), *Orthetrum albistylum speciosum* (White-tailed skimmer; burrower), and *Pantala flavescens* (Wandering glider; sprawler). Nymphs of *A. nigrofasciatus* collected in Sakura City, Chiba Prefecture, and those of *G. japonica* collected in Higashihiroshima City, Hiroshima Prefecture, were obtained from an online vendor, and *O. a. speciosum* and *P. flavescens* were collected from the same paddy field as the mating frog pairs. The dragonfly nymphs were maintained individually in plastic trays (23 × 17 × 8 cm) and fed tadpoles of *D. japonicus* every two days until the beginning of the induction experiment.

The induction experiment commenced on July 12, 2022. Each experimental tank housed 21 tadpoles, with three individuals randomly selected from each of the seven egg clutches to randomize genetic effects in each tank. Tadpoles in a tank were assigned to one of five treatments (Ax group with *A. nigrofasciatus*; Gy group with *G. japonica*; Or group with *O. a. speciosum*; Pf group with *P. flavescens*; and the control Ctl group) with eight replicates per treatment. Each experimental tank (31 × 18 × 24 cm; 10 cm water depth) were equipped with a transparent isolation case (17 × 9 × 10 cm) with approximately 5 mm diameter holes spaced about 5 mm apart, where one dragonfly nymph was introduced. This setup allowed visual and chemical-mediated interaction between the predator and tadpoles without physical

contact. The tanks were aerated and maintained under a 14-h light/10-h dark photoperiod at a

room temperature of 22°C, with half of the water in all tanks changed every three days. Air

was continuously supplied to the isolation case within each tank through an aquarium air

stone. Tanks designated for the Ct1 group lacked predators and utilized an empty isolation

case. Tadpoles in the experimental tanks were fed every two days with commercial food for

Axolotl, *Ambystoma mexicanum* (Kyorin Co., Ltd.), while dragonfly nymphs received two

tadpoles as food every two days.

Experiment II

The design of Experiment II mirrored that of Experiment I, with adjustments made

for sampling dates, the interval between hatching and the experiment's commencement, and

the selection of predators. For this experiment, adults of *D. japonicus* were captured from the

paddy fields at Kyoto University on May 18 and 20, 2022. Eggs were collected and tadpoles

from clutches were reared in the same way as in Experiment I until the experiments began.

Three predator species were chosen based on their foraging style and the primary sensory

organs used for prey detection: Ferocious water bug, *Appasus major* (sit-and-wait predator

using tactile and olfactory senses), water scorpion, *Laccotrephes japonensis* (sit-and-wait

predator using tactile and olfactory senses), and Japanese fire belly newt, *Cynops*

*pyrrhogaste*r (active tracker using olfactory senses). Aquatic insects, collected in

Higashihiroshima City, were purchased from the online vendor, while red-bellied newts were

collected from irrigation ditches in Kyoto City (35°05'N 135°47'E). These predators were

individually housed in plastic trays and fed with tadpoles every two days.

The induction experiment began on June 18, 2022. Each experimental tank housed

21 tadpoles, with seven individuals randomly selected from each of the three egg clutches.

Tadpoles in each tank were allocated to one of four treatments (Be group with *A. major*; Ws

group with *L. japonensis*; Fn group with *C. pyrrhogaster*; and the control Ct2 group) with

eight replicates per treatment. One predator was introduced into an isolation case in each tank.

Other experimental setups, i.e., tank design, room temperature, light condition, and feeding

frequency, were kept the same as in Experiment I.

Measuring Coloration and Morphological Responses

All tadpoles were photographed two and three weeks after the start of the experiment

using a Canon® EOS RP mirrorless camera. Individual tadpoles were placed in a narrow

acrylic aquarium with a ruler and color chart in the background. The distance between the

aquarium and the camera was kept constant for all individuals. All photos were taken with a

F/16 aperture, a 1/20th second exposure time, and ISO 3200. Consistency in artificial lighting

conditions was maintained for individuals photographed on the same day. Some individuals

were excluded from the analysis due to poor nutritional condition or significant tail loss

during rearing. The exclusion criteria included: (1) head and body length less than 6 mm, (2)

severe tail damage or loss, and (3) inability to maintain posture in water.

Tail parts excluding the tail muscle were trimmed using the image measuring

software, Natsumushi version 1.12.2 (Tanahashi and Fukatsu, 2018;

https://sites.google.com/site/mtahashilucanid/program/natsumushi). The color of the tail was

quantified using the average of CIELAB values (CIE 1976) calculated in Adobe Photoshop

25.5.1 (Adobe Inc.). In the CIELAB color space, color was represented using three

components: the lightness component L* and the color components a* and b*. The L*

component ranged from black to white between 0–100, while a* and b* components ranged

from green to red and blue to yellow, respectively, on a scale of −128 to 127. As Photoshop

displays images in 8-bit format, these values were rescaled to range from 0 to 255. Therefore,

L* was multiplied by 100/255 and a* and b* were adjusted by subtracting 128, respectively,

to facilitate analysis (Fig. 1). Due to an incorrect white balance, photographs taken at the

second week of Experiment II were excluded from the analysis of tail coloration. During the

third week of Experiment II, photographs of 11 individuals from the Ct2 group were also

taken with an incorrect white balance, so the color tone was later corrected using Photoshop,

referencing one of the other photographs from the Ct2 group. Additionally, morphological

measurements were conducted based on the photographs using Natsumushi. Morphological

traits included tadpole body length (BL), tail length (TAL), and tail depth (TD) were

measured using the scale within each image.

Statistical Analyses

256   All statistical analyses were conducted in R version 4.3.2 (R Core Team, 2023).

Linear mixed models (LMMs) were employed based on the measured traits, using the

lmerTest package (Kuznetsova et al., 2017) to estimate differences between the control group

and the other treatments. The experimental tank was treated as a random factor in the LMMs.

When color components (e.g., a^* and b^*) were utilized as the response variables, the

treatment (predator) was used as the explanatory variable in the models to compare the

treatment groups to the control group. In the same way, when body length (BL) was utilized

as the response variables, the treatment was used as the explanatory variable in the models.

On the other hand, in analyses for TAL and TD as the response variable, four models were

constructed for each trait. These models included: the model with only BL as the explanatory

variable, the model with only the treatment as the explanatory variable, the model with both

BL and treatment as explanatory variables, and the model incorporating BL, treatment, and

their interactions. The final model was determined via backward elimination, by applying

likelihood ratio (LR) tests. Specifically, all explanatory variables (with their interactions) were

entered into the full model, then removed one by one until a final model was selected. At each

step, the reduced model was compared with the former model by LR test. The significance

level was set at 5%. If a significant difference was found between the former and reduced

models, the former model was adopted as the final model for further analysis.

274   After constructing final models, comparisons between each treatment group and the

control were performed using Dunnett's test with the glht function in the multcomp package

(Hothorn et al., 2008). The significance level was set at 5%, with treatment groups considered

significantly different from the control group if the P values were below this threshold.

Orange tail coloration and increased tail depth were reported as predator-induced traits in

previous studies, so we decided to focus primarily on b* values and tail depth.

We also conducted a principal component analysis (PCA) for color and

morphological traits at the third week after the start of the experiment to concisely visualize

the multivariate responses of the tadpoles to the predator treatments. The analysis included

total length (TL; the sum of BL and TAL), standardized body length (sBL), standardized tail

length (sTAL), standardized tail depth (sTD), and coloration values (a* and b*), The sBL,

sTAL, and sTD were calculated by dividing the BL, TAL, and TD measurements by TL, then

subtracting their respective means and dividing by their standard deviations, in order to use

variables with different units together. Pearson's correlation matrices were used to compare

the principal component scores with the original variables to interpret the principal

components. We also used Pearson's correlation matrices to assess the relationships among

tadpole traits.

Results

Coloration

In Experiment I, the values of a* were lower in the Ax group compared to the Ct1

group (Dunnett's test, $P < 0.001$; Fig. 2). Additionally, the Ax group had a higher value of b*

($P < 0.001$). The latter indicates that tadpoles in the Ax group showed more orange tails than

in the control group. Also at the third week, the values of a* were lower ($P = 0.0283$) and b*

were higher ($P < 0.001$) in the Ax group than in the control group. In addition, the Or group

showed higher b* values than the Ct1 group ($P = 0.0443$), indicating that orange tails were

induced not only by *Anax nigrofasciatus*, but also by *Orthetrum albistylum speciosum*. Some

individuals of the Pf group also exhibited tail coloration, but this tendency was not

statistically significant ($P = 0.1426$ in b* value). However, in Experiment II, no group showed

significant differences from the Ct2 group in the a* or b* values at the third week (Fig. 3).

Morphological Responses

In Experiment I, whether tail length or height was used as the response variable,

models with BL and treatment as explanatory variables were selected as the final models at

both the second and third weeks after the start of the experiment. At the second week, the Pf

group had a larger body compared to the Ct1 group (Dunnett's test, $P = 0.0471$; Fig. 4), and

tails were shorter in the Ax and Pf groups than in the Ct1 group ($P < 0.001$). Moreover, the Ax

group had a higher tail than the Ct1 group ($P = 0.0092$). At the third week, only the Gy group

had a shorter tail than the Ct1 group ($P = 0.047$), while the Pf group, as well as the Ax group,

showed a higher tail than the Ct1 group ($P = 0.0215$ for Pf, $P < 0.001$ for Ax). The Or group

also tended to have higher tails compared to the Ct1 group, but the difference was not

statistically significant ($P = 0.1132$).

In Experiment II, the models with BL and treatment as explanatory variables were

chosen in the model selection for tail length at the second week. On the other hand, at the

third week, the models with only BL were selected. As for tail depth, the models with only BL

were also selected. Although the models with only BL as explanatory variable indicated that

the effect of the treatment on tail length or tail depth was not large, post-hoc tests were

conducted to assess differences between treatments. At the second week, the Be group

exhibited a shorter tail ($P < 0.001$), but no significant differences in tail depth were observed

between any group and the Ct2 group ($P > 0.05$ for all Be, Ws, and Fn; Fig. 5). At the third

week, there was no significant difference in tail depth compared to the Ct2 ($P > 0.05$ for all

Be, Ws, and Fn). Consequently, tadpoles in all non-odonate predator treatments were not

induced to develop significantly deeper tail fins compared to the control group.

Principal Component Analysis

In Experiment I, the first two principal components (PC1 and PC2) for larval coloration and morphology accounted for 67.3% of the variation, then combined with PC3, they explained 83% (Table 1). PC1 showed strong correlations with all morphological traits except sTD, indicating that it is likely influenced by the larval body proportion between body length and tail length. On the other hand, PC2 was strongly correlated with sTD (Pearson's correlation coefficients, -0.84) and b* (-0.65), suggesting that it represented the direction of predator-induced responses of *Dryophytes japonicus*. The 95% probability ellipse of the Ax group was conspicuously elongated in the negative direction along the PC2 axis, followed by the Pf, Or and Gy groups (Fig. 6). An increase in tail height and coloration change occurred together; the correlation coefficient between sTD and b* value was 0.34, ranking as the fourth highest after the coefficients between TL and other morphological traits (sBL, sTAL), and between TL and b* value (Table 2).

In Experiment II, the first two principal components explained 77.5% of the variation, and combined with PC3, they accounted for 90.4% (Table 1). PC1 largely explained color trait variation, as well as morphological traits except sTD. PC2 exhibited strong correlations with sBL, sTAL, and sTD, while PC3 showed relatively strong correlations with sTD and b*. Although PC3 may represent the direction of response in this species, its overall contribution to the total variation was limited (12.9%). Furthermore, the correlation between sTD and b* value was 0.16, which was not particularly large compared to other coefficients (Table 2), indicating that changes in tail color and shape did not necessarily occur together.

Discussion

Tail Coloration in Asian Species of *Dryophytes*

Our experiments revealed that tadpoles of *Dryophytes japonicus* developed induced changes in tail coloration as well as tail shape in response to dragonfly nymphs. Although tail coloration has rarely been documented in this species, its occurrence has been reported from

several locations in Japan, including Hokkaido, Shiga, Hyogo, Hiroshima, Shimane, Kouchi

and Ehime prefectures (Doi et al., 2005), as well as from our collection site in Kyoto (Noda,

personal observation). The phenotypes induced by *Anax nigrofasciatus*, i.e., deep tail fins and

bright orange tails, closely resemble those previously observed in other species in *Dryophytes*

in the New World, Cope's Gray Treefrog (*Dryophytes chrysoscelis*; McCollum and Van

Buskirk, 1996), Pine Woods Treefrog (*Dryophytes femoralis*; LaFiandra and Babbitt, 2004),

and Gray Treefrog (*Dryophytes versicolor*; Van Buskirk and McCollum, 1999). Thus, our

findings represent the first documented case of a predator-induced tail coloration phenotype in

Asian species of *Dryophytes*.

Our discoveries of the plasticity in the tail coloration in the Asian species of

Dryophytes suggests that the trait is symplesiomorphically shared between the North

American and ancestral Asian lineages, and has been inherited by present-day Asian species,

including *D. japonicus*. An ancestral stock of *Dryophytes* is inferred to have migrated from

North America to East Asia across the Bering Land Bridge during the Miocene (Duellman et

al. 2016; Dufresnes et al., 2016). Present-day Asian species of *Dryophytes* are monophyletic,

having diversified from this ancestral stock. It is plausible that tail coloration has persisted as

a phenotypic plasticity in both North American and East Asian species of *Dryophytes* due to

the consistent importance of dragonfly nymphs as predators.

It remains unknown whether tail coloration can be induced in the three other Asian

species, *Dryophytes flaviventris*, *Dryophytes immaculates*, and *Dryophytes suweonensis* (see

Borzée et al., 2020). In North America, Pine Barrens Treefrog, *Dryophytes andersonii* is

induced to have a darker tail, not an orange tail, by *Anax junius*, which induces orange tails in

other species of *Dryophytes* (Kruger and Morin, 2020). Furthermore, Squirrel Treefrog,

Dryophytes squirellus exhibits no morphological changes in response to odonate species

(McCoy and Bolker 2008; Baker, 2021). These cases indicate that the plastic responses in

Dryophytes have diversified, and some may have lost the ability to develop orange tails.

Future studies are needed to elucidate the interspecific variation in plastic responses to

predation risk and the evolutionary causes underlying this variation.

Differences in the Extent of Coloration by Predators

Our experiments demonstrated that the extent of induced responses varied depending

on the dragonfly species reared with tadpoles. Two weeks after the start of the experiment,

nymphs of *A. nigrofasciatus* induced both deep fin and tail coloration in tadpoles, whereas, at

the third week, *Pantala flavescens* also induced deep tail and *Orthetrum albistylum speciosum*

induced tail coloration. Furthermore, although weak, some tadpole individuals exposed to *P.*

flavescens showed tail coloration, and those exposed to *O. albistylum speciosum* also tended

to have higher tails. These results indicate that the defense traits (coloration and deep tail)

were induced by the nymphs of *A. nigrofasciatus* more rapidly and strongly, but similar

responses were also elicited by the other two predator species. The probability ellipses in PCA

further suggest that the direction of larval responses to these benthic predators (*P. flavescens*

and *O. albistylum speciosum*) were similar to that to *A. nigrofasciatus* (Fig. 6).

The differences in the degree and rate of responses induced by different dragonfly

species may reflect varying levels of predation risk for tadpoles posed by each species. Plastic

phenotypes with significant tail coloration should result in trade-offs because of energetic

costs associated with synthesizing pigmentation (Grether et al., 2001) or costs due to the

increased vulnerability to mismatched predators. Consequently, tadpoles are expected to

minimize these costs by developing effective defense traits for each predator as much as

possible based on the perceived predation risk (Touchon and Warkentin, 2008; Innes-Gold et

al., 2019). The predation risk for tadpoles would be different among dragonfly species

according to (1) the foraging style of dragonfly nymphs, (2) their color vision, and (3) the

total amount of intake related to their body size and occurrence period.

Firstly, the four dragonfly species used in our experiment exhibit different foraging

styles. According to the classification by Corbet (1999), *A. nigrofasciatus* and *G. japonica* are

climbers, sit-and-wait predators on aquatic vegetation, while *P. flavescens* is a sprawler,

foraging while moving along the bottom, and *O. albistylum speciosum* is a burrower, residing

within the substrate and ambushing prey. For sit-and-wait predators, there is often a time lag

between identifying prey and launching an attack, during which they can choose which part of

tadpole's body to target. Therefore, the lure effect of tadpole tails is likely to attract ambush

predators more effectively than those that actively chase their prey. In addition, as *A.*

nigrofasciatus prefers habitats abundant in floating-leaved plants (Ozono et al., 2021), it can

move freely along the plant structures from roots anchored at the bottom to leaves extending

above the water surface. As a result, the risk of predation on tadpoles could be present

throughout the entire water column, from the bottom to the surface, making behavioral

responses like moving to a specific depth insufficient as a means of avoiding predation.

Conversely, *P. flavescens* and *O. albistylum speciosum* primarily forage at the bottom, so

tadpoles may evade these predators by relocating to higher water layers. However, such

avoidance behavior often leads to a decrease in growth rate due to reduced foraging efficiency

(Teplitsky et al., 2003); i.e., for tadpoles that primarily feed on detritus, avoiding the bottom

layer may result in malnutrition. The combination of moderate morphological responses with

behavioral responses (the "double defense" strategy; Dewitt and Hucko, 1999) may help

mitigate the impact of malnutrition from behavioral changes while minimizing the costs of

defensive traits.

Secondly, differences in eyesight, particularly color vision among dragonfly species,

may influence the extent of the tail coloration changes in the tadpoles. Pritchard (1966) noted

a correlation between foraging habit and compound eye size in dragonfly nymphs, with

climbers typically possessing larger compound eyes compared to sprawlers and burrowers. He

also stated that the climbers, such as species in *Anax*, rely heavily on vision for prey

detection, whereas other foraging types depend more on tactile stimuli. Hence, it is plausible

that the lure effect of the brightly colored tails is more effective for climbers, which are more

visually oriented. Furthermore, Futahashi et al. (2015) investigated the genetic basis of color

vision in dragonflies and redocumented the number of opsin genes and the characteristic

spatiotemporal expression patterns in the nymphs of various species, including those of the

burrower *O. albistylim* and the climber *A. parthenope*, a species closely related to *A.*

nigrofasciatus used in our experiment. They found that dragonflies possess a greater number

of opsin genes than other insects, and *A. parthenope* possesses and expresses a greater number

of opsin genes, particularly those sensitive to long-wavelength light. Although the functions

of these diverse genes are not fully understood, this suggest that species of *Anax* may have

enhanced abilities to perceive reddish colors, potentially enabling tadpoles with bright orange

tails to more effectively avoid predation by using their tails as lures. Thus, the larval

antipredator defense of *D. japonicus*, which involves tail coloration, may have been driven by

the sensory systems of predatory dragonfly nymphs, and subsequently evolved specifically

against color-sensitive predators, represented by dragonfly nymphs of *Anax*.

Thirdly, the variation in the degree of induced responses may reflect the relative

predation risks associated with body sizes or occurrence periods of respective odonate

predators. In nature, it is generally difficult to determine which predators exert higher

predation pressure on tadpoles, and few studies have investigated such differences. However,

during the rearing process, we observed that the nymphs of *A. nigrofasciatus* exhibited higher

food demands and displayed more frequent attacks on tadpoles compared to other species.

Additionally, they are approximately twice the size of the nymphs of *P. flavescens* and *O.*

albistylum speciosum (see Ishida et al., 1988), suggesting that size-related predation pressures

could influence the necessity and effectiveness of tadpole morphological responses. However,

tadpoles did not exhibit significant responses to *G. japonica*, which is similar in size to *A.*

nigrofasciatus (see Ishida et al., 1988). This indicates that tadpoles likely respond not only to

size-related predation pressure but also to other characteristics of dragonfly species, such as

foraging style or habitat overlap, thereby modulating the extent of induced responses. Since

the habitat of nymphal *G. japonica* overlaps with that of tadpoles of *D. japonicus* (Nakagawa

et al., 2004), the lack of an apparent induced response may be due to the limited period in

which they co-occur. This species grows more rapidly compared to other species in Aeshnidae

in Japan, occurring in aquatic environments only from May to August (Ozono et al., 2021).

Considering that the nearly-metamorphosing nymphs cease foraging, the period during which

this species poses a predation threat to *D. japonicus* is even shorter. This brief coexistence

period suggests that this dragonfly species may not be an important predator for the tadpoles.

In addition to the tail coloration and depth we focused on, a tendency toward larger

body lengths and shorter tail lengths compared to those in the control group was observed,

although not statistically significant in most treatment groups. These changes may represent

general responses to predator treatments, as noted in previous studies. The increase in body

length could be attributed to enhanced availability of food resources resulting from leftovers

and feces of caged predators (e.g., McIntyre et al., 2004; Richardson, 2006). Similarly, shorter

tail lengths in the presence of predators have been reported in several frog species (Laurila et

al., 2002; LaFiandra and Babbitt, 2004).

Responses Toward Non-Odonate Predators

Tadpoles of *D. japonicus* showed neither conspicuous tail coloration nor deep fin

when exposed to newts or aquatic insects other than dragonfly nymphs in our experiment,

likely because the costs of expressing these defenses outweighed their benefits for avoiding

predation. Although Experiment II did not include a positive control (i.e., treatment with

dragonfly nymphs, such as *A. nigrofasciatus*), we believe that the absence of color induction

was unlikely caused by experimental failure, because preliminary experiments demonstrated

the high reproducibility of the induced responses when using dragonfly nymphs. Therefore,

our results suggest that changes in body color and tail depth are not strongly induced by these

predators and may not serve as effective predation avoidance strategies against them. An

important point in understanding why tadpoles did not show tail coloration is that defensive

traits come with physiological costs and may increase predation pressure from non-focal

predators. If changes in tail color and shape do not significantly contribute to non-odonate

predator avoidance, tadpole should minimize these disadvantages by refraining from

developing these antipredator defenses.

Given that the non-odonate predators are presumably less dependent on vision than

odonate species when foraging, the strategy of using colored deep tails as lures may be

ineffective and provides only limited benefits against them. For instance, diving beetles, water

bugs, and water scorpions capture prey by grasping the entire body with their arms, so the tail

of tadpoles should not serve as a lure for them. Additionally, they use tactile, visual, and

olfactory senses for locating prey (Babbitt and Jordan, 1996; Kehr and Gómez, 2009; Mogali

et al., 2019; Nowińska and Brożek, 2019), and possess well-developed organs for detecting

auditory and mechanical signals (Yager, 1999; Cocroft and Rodríguez, 2005), suggesting they

are relatively less dependent on vision. Previous studies have reported predator-induced

changes in larval morphology in response to these predators (Benard, 2006; McCoy, 2007;

Michel, 2012), but few have reported effects on larval tail coloration (Touchon and

Warkentin, 2011). Similarly, newts and salamanders, both as larvae and adults, are active

foragers that swallow their prey whole like fish (Wilson et al., 2005; Kishida and Nishimura,

2007). Consequently, the tails of the tadpoles are unlikely to function as a lure for these

predators. Newts tend to rely on olfaction cues as well as vision for food recognition

(Copeland, 1913; Lindquist and Bachmann, 1982; Schwarz et al., 2020). Few studies have

reported that tadpoles exhibit induced defensive traits in response to these predators, and none

have documented induced changes in body coloration (Kishida and Nishimura, 2004; Hettyey

et al., 2011; Urban et al., 2017). These facts support the hypothesis that the brightly colored

tails of D. japonicus do not function as lures against the non-odonate predators, and that, as a

result, tadpoles do not express costly defensive traits in response to them.

The lack of induction of defensive phenotypes may also be attributed to

physiological costs associated with developing tail coloration, although the magnitude of such

costs in tadpoles remains poorly understood. Since most animals are unable to biosynthesize

carotenoids, the pigments thought to be causing the bright orange coloration, they metabolize

and accumulate these pigments by absorbing them from their diet (Bertrand et al., 2006;

Thibaudeau and Altig, 2012). Thus, the allocation costs of expressing coloration by

transporting these pigments may not be as high as commonly assumed. On the other hand, the

potential for increased vulnerability to non-focal predators may also interfere with the

expression of defensive traits (Innes-Gold et al., 2019). This disadvantage may sometimes

outweigh the energetic costs of expressing a defensive phenotype. Carfagno et al. (2011)

found that tadpoles of *Acris* exhibit darker a tail spot, which is considered a defensive trait

against dragonfly nymphs, even in the absence of predators. However, the presence of fish

predators induces the loss of this spot. This suggests that the cost of expressing defensive

traits may be lower than typically expected, and the disadvantage of increased vulnerability to

mismatched predators may prevent these phenotypes from being constitutively expressed.

Similarly, in tadpoles of *D. japonicus*, the elevated risk of predation from non-odonate

predators, considered non-focal predators, due to expressing bright tail coloration, rather than

its energetic cost, could explain why they did not exhibit deep and brightly-colored tail in

response to these predators. These defense traits against dragonfly nymphs may not only be

ineffective in preventing predation by non-odonate predators but could also increase the

potential vulnerability to them. For example, the deepened tail may increase the vibrations

produced during swimming, thereby heightening predation pressure on tadpoles by tactile or

olfactory predators (i.e., water bugs, water scorpions, and newts). Additionally, for other

predators such as fish, tail coloration may make the tadpoles more vulnerable by increasing

their visibility. However, further experiments are warranted to assess and compare the

effectiveness of these defense traits toward each predator, including evaluating how tadpole

coloration influence their survival rate in the presence of various aquatic predators.

Acknowledgements

We thank the Experimental Farm, Graduate School of Agriculture, Kyoto University, for providing the research environment, and Mr. Kagata, Mr. Yasui, and Mr. Sakakibara, technical staff, for their assistance with observations. This work was supported by JST SPRING, Grant Number JPMJSP2110.

Literature Cited

Agrawal, A. A., and R. Karban. 1999. Why induced defenses may be favored over constitutive strategies in plants, p. 45−61. *In*: Ecology and Evolution of Inducible Defenses. R. Tollrian and C. D. Harvell (eds.). Princeton University Press, Princeton, New Jersey.

Babbitt, K. J., and F. Jordan. 1996. Predation on *Bufo terrestris* tadpoles: effects of cover and predator identity. Copeia 1996:485−488.

Baker, E. A. 2021. Evaluating a Predator-Induced Phenotype in a Mixed Species Context. Unpubl. Master's Thesis, East Carolina University, Greenville, North Carolina.

Barnett, J. B., C. Michalis, N. E. Scott-Samuel, and I. C. Cuthill. 2018. Distance dependent defensive coloration in the poison frog *Dendrobates tinctorius*, Dendrobatidae. Proceedings of the National Academy of Sciences 115:6416−6421.

Barry, M. J., and S. Syal. 2013. Metabolic responses of tadpoles to chemical predation cues. Hydrobiologia 700:267−276.

Benard, M. F. 2006. Survival trade-offs between two predator-induced phenotypes in Pacific treefrogs (*Pseudacris regilla*). Ecology 87:340−346.

Bertrand, S., B. Faivre, and G. Sorci. 2006. Do carotenoid-based sexual traits signal the availability of non-pigmentary antioxidants? Journal of Experimental Biology, 209:4414−4419.

Bókony, V., Á. M. Móricz, Z. Tóth, Z. Gál, A. Kurali, Z. Mikó, K. Pásztor, M. Szederkényi, Z. Tóth, J. Ujszegi, B. Üveges, D. Krüzselyi, R. J. Capon, H. Hoi, and

A. Hettyey. 2016. Variation in chemical defense among natural populations of common toad, *Bufo bufo*, tadpoles: the role of environmental factors. Journal of Chemical Ecology 42:329−338.

Borzée, A., K. R. Messenger, S. Chae, D. Andersen, J. Groffen, Y. I. Kim, J. An, S. N. Othman, K. Ri, T. Y. Nam, Y. Bae, J. L. Ren, J. T. Li, M. F. Chuang . . . M. S. Min. 2020. Yellow sea mediated segregation between North East Asian *Dryophytes* species. PLoS One 15:e0234299.

Brönmark, C., and L. A. Hansson. 2000. Chemical communication in aquatic systems: an introduction. Oikos 88:103−109.

Bybee, S. M., K. K. Johnson, E. Gering, M. F. Whiting, and K. A. Crandall. 2012. All the better to see you with: a review of odonate color vision with transcriptomic insight into the odonate eye. Organisms Diversity & Evolution 12:241−250.

Carfagno, G. L. F., J. M. Carithers, L. J. Mycoff, and R. M. Lehtinen. 2011. How the cricket frog lost its spot: the inducible defense hypothesis. Herpetologica 67:386−396.

Cocroft, R. B., and R. L. Rodríguez. 2005. The behavioral ecology of insect vibrational communication. Bioscience 55:323−334.

Copeland, M. 1913. The olfactory reactions of the spotted newt *Diemyctylus viridescens* (Rafinesque). Journal of Animal Behavior 3:260−273.

Corbet, P. S. 1999. Dragonflies: behaviour and ecology of Odonata. Cornell University Press, Ithaca, New York.

Costello, D. M., and M. J. Michel. 2013. Predator-induced defenses in tadpoles confound body stoichiometry predictions of the general stress paradigm. Ecology 94:2229–2236.

Dewitt, T. J., A. Sih, and J. A. Hucko. 1999. Trait compensation and cospecialization in a freshwater snail: size, shape and antipredator behaviour. Animal Behaviour 58: 397–407.

Doherty, P. A., R. J. Wassersug, and J. M. Lee. 1998. Mechanical properties of the tadpole tail fin. Journal of Experimental Biology 201:2691−2699.

Doi, T., H. Okuyama, and S. Murakami. 2005. Red tail fin patches in Japanese tree frog

tadpoles (*Hyla japonica*). Bulletin of the Herpetological Society of Japan 2005:4−7.

Duellman, W. E., A. B. Marion, and S. B. Hedges. 2016. Phylogenetics, classification, and

biogeography of the treefrogs (Amphibia: Anura: Arboranae). Zootaxa 4104:1−109.

Dufresnes, C., S. N. Litvinchuk, A. Borzée, Y. Jang, J. T. Li, I. Miura, N. Perrin, and M.

Stöck. 2016. Phylogeography reveals an ancient cryptic radiation in East-Asian tree frogs

(*Hyla japonica* group) and complex relationships between continental and island

lineages. BMC Evolutionary Biology 16:253.

Espanha, J., M. F. de Vasconcelos, and P. C. Eterovick. 2016. The role of tadpole

coloration against visually oriented predators. Behavioral Ecology and Sociobiology

70:255−267.

Futahashi, R., R. Kawahara-Miki, M. Kinoshita, K. Yoshitake, S. Yajima, K. Arikawa,

and T. Fukatsu. 2015. Extraordinary diversity of visual opsin genes in dragonflies.

Proceedings of the National Academy of Sciences 112:E1247−E1256.

Grether, G. F., J. Hudon, and J. A. Endler. 2001. Carotenoid scarcity, synthetic pteridine

pigments and the evolution of sexual coloration in guppies (*Poecilia reticulata*).

Proceedings of the Royal Society of London Series B 268:1245−1253.

Hawlena, D., R. Boochnik, Z. Abramsky, and A. Bouskila. 2006. Blue tail and striped

body: why do lizards change their infant costume when growing up? Behavioral Ecology

17:889−896.

Hettyey, A., Z. Tóth, K. E. Thonhauser, J. G. Frommen, D. J. Penn, and J. Van Buskirk.

2015. The relative importance of prey-borne and predator-borne chemical cues for

inducible antipredator responses in tadpoles. Oecologia 179:699−710.

Hettyey, A., B. Üveges, Á. M. Móricz, L. Drahos, R. J. Capon, J. Van Buskirk, Z. Tóth,

and V. Bókony. 2019. Predator-induced changes in the chemical defence of a vertebrate.

Journal of Animal Ecology 88:1925−1935.

Hettyey, A., K. Vincze, S. Zsarnóczai, H. Hoi, and A. Laurila. 2011. Costs and benefits of

defences induced by predators differing in dangerousness. Journal of Evolutionary

Biology 24:1007−1019.

Hettyey, A., S. Zsarnóczai, K. Vincze, H. Hoi, and A. Laurila. 2010. Interactions between

the information content of different chemical cues affect induced defences in tadpoles.

Oikos 119:1814−1822.

Hirai, N., T. Morioka, and M. Ishii. 2020. Species diversity of Odonata in Nakaikemi

Marsh, Fukui Prefecture, Japan. Japanese Journal of Environmental Entomology and

Zoology 31:1−12.

Hoff, K. V., and R. J. Wassersug. 2000. Tadpole locomotion: axial movement and tail

functions in a largely vertebraeless vertebrate. American Zoologist 40:62−76.

Hossie, T., K. Landolt, and D. L. Murray. 2017. Determinants and co-expression of anti-

predator responses in amphibian tadpoles: a meta-analysis. Oikos 126:173–184.

Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous inference in general parametric

models. Biometrical Journal 50:346–363.

Innes-Gold, A. A., N. Y. Zuczek, and J. C. Touchon. 2019. Right phenotype, wrong place.

Proceedings of the Royal Society B 286: 20192347.

Ishida, S., K. Ishida, K. Kojima, and M. Sugimura. 1988. Illustrated guide for

identification of the Japanese Odonata. Tokai University Press, Shinjuku, Tokyo.

Johnson, J. B., D. B. Burt, and T. J. Dewitt. 2008. Form, function, and fitness: pathways to

survival. Evolution 62:1243−1251.

Kasumovic, M. M. 2013. The multidimensional consequences of the juvenile environment:

towards an integrative view of the adult phenotype. Animal Behaviour 85:1049–1059.

Kehr, A. I., and V. I. Gómez. 2009. Intestinal, body and tail plasticity in *Rhinella schneideri*

(Bufonidae) tadpoles induced by a predator insect (Belostoma elegans). Advanced

Studies in Biology 1:85–94.

Kishida, O., and K. Nishimura. 2004. Bulgy tadpoles: inducible defense morph. Oecologia
140:414–421.

Kishida, O., and K. Nishimura. 2005. Multiple inducible defences against multiple
predators in the anuran tadpole, *Rana pirica*. Evolutionary Ecology Research 7:619–631.

Kishida, O., and K. Nishimura. 2007. Flexible inducible morphological defense in anuran
larvae. Japanese Journal of Ecology 57:40–47.

Kita, A., M. Nakahara, and M. Tokuda. 2020. Changes in Odonata abundance. between
2000 and 2015–2016 in Saga Plain, northern Kyushu, Japan. Journal of Insect
Conservation 24:575–583.

Kruger, A., and P. J. Morin. 2020. Predators induce morphological changes in tadpoles of
Hyla andersonii. Copeia 108: 316–325.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests
in Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

LaFiandra, E. M., and K. J. Babbitt. 2004. Predator induced phenotypic plasticity in the
pinewoods tree frog, *Hyla femoralis*: necessary cues and the cost of development.
Oecologia 138:350–359.

Lardner, B. 1998. Plasticity or fixed adaptive traits? Strategies for predation avoidance in
Rana arvalis tadpoles. Oecologia 117:119–126.

Lardner, B. 2000. Morphological and life history responses to predators in larvae of seven
anurans. Oikos 88:169–180.

Laurila, A., J. Kujasalo, and E. Ranta. 1998. Predator-induced changes in life history in
two anuran tadpoles: effects of predator diet. Oikos 83:307–317.

Laurila, A., S. Pakkasmaa, P. A. Crochet, and J. Merilä. 2002. Predator-induced plasticity
in early life history and morphology in two anuran amphibians. Oecologia 132:524–530.

Lawler, S. P. 1989. Behavioural responses to predators and predation risk in four species of
larval anurans. Animal Behaviour 38:1039–1047.

Lindquist, S. B., and M. D. Bachmann. 1982. The role of visual and olfactory cues in the

prey catching behavior of the tiger salamander, *Ambystoma tigrinum*. Copeia 1982:81–

90.

Matsui, M., and N. Maeda. 2018. Encyclopedia of Japanese frogs. Bun-ichi Co., Ltd.,

Shinjuku, Tokyo.

McCollum, S. A., and J. D. Leimberger. 1997. Predator-induced morphological changes in

an amphibian: predation by dragonflies affects tadpole shape and color. Oecologia

109:615–621.

McCollum, S. A., and J. Van Buskirk. 1996. Costs and benefits of a predator-induced

polyphenism in the gray treefrog *Hyla chrysoscelis*. Evolution 50:583–593.

McCoy, M. W. 2007. Conspecific density determines the magnitude and character of

predator-induced phenotype. Oecologia 153:871–878.

McCoy, M. W., and B. M. Bolker. 2008. Trait-mediated interactions: influence of prey size,

density and experience. Journal of Animal Ecology 77:478–486.

McIntyre, P. B., S. Baldwin, and A. S. Flecker. 2004. Effects of behavioral and

morphological plasticity on risk of predation in a Neotropical tadpole. Oecologia

141:130–138.

Michel, M. J. 2012. Phenotypic plasticity in complex environments: effects of structural

complexity on predator-and competitor-induced phenotypes of tadpoles of the wood frog,

Rana sylvatica. Biological Journal of the Linnean Society 105:853–863.

Mogali, S., B. Shanbhag, and S. Saidapur. 2019. Experience of predacious cues and

accessibility to refuge minimize mortality of *Hylarana temporalis* tadpoles. Acta

Herpetologica 14:15–19.

Nakagawa, M., T. Motobayashi, Y. Arai, and T. Nishimura. 2004. Ecology of Eggs of

Gynacantha japaonica Bartenef in Paddy Fields. Transactions of the Japanese Society of

Irrigation, Drainage and Reclamation Engineering 229:71–77.

Norris, K. S., and C. H. Lowe. 1964. An analysis of background color-matching in

amphibians and reptiles. Ecology 45:565–580.

Nowińska, A., and J. Brożek. 2019. Antennal sensory structures in water bugs of Nepoidea

(Insecta: Hemiptera: Nepomorpha), their morphology and function. Zoomorphology

138:307–319.

Nunes, A. L., A. Richter-Boix, A. Laurila, and R. Rebelo. 2013. Do anuran larvae respond

behaviourally to chemical cues from an invasive crayfish predator? A community-wide

study. Oecologia 171:115–127.

Ohba, S., and F. Nakasuji. 2006. Dietary items of predacious aquatic bugs (Nepoidea:

Heteroptera) in Japanese wetlands. Limnology 7:41–43.

Orton, G. L. 1953. The systematics of vertebrate larvae. Systematic Zoology 2:63–75.

Ozaki, K., and K. Nishikawa. 2023. Feeding Habits of the Japanese Fire-belly Newt.

(Ampbihia: Urodela: Salamandridae) in Central Honshu, Japan. Current Herpetology

42:162–170.

Ozono, A., I. Kawashima, and R. Futahashi. 2021. Dragonflies of Japan. Bun-ichi Co. Ltd.,

Shinjuku, Tokyo.

Pritchard, G. 1965. Prey capture by dragonfly larvae (Odonata; Anisoptera). Canadian

Journal of Zoology 43:271–289.

Pritchard, G. 1966. On the morphology of the compound eyes of dragonflies (Odonata:

Anisoptera), with special reference to their role in prey capture. Proceedings of the Royal

Entomological Society of London. Series A, General Entomology 41:1–8.

R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R- project.org/.

Relyea, R. A. 2001. Morphological and behavioral plasticity of larval anurans in response to

different predators. Ecology 82:523–540.

Relyea, R. A., and J. R. Auld. 2004. Having the guts to compete: how intestinal plasticity

explains costs of inducible defences. Ecology Letters 7:869–875.

Relyea, R. A., and E. E. Werner. 2000. Morphological plasticity in four larval anurans

distributed along an environmental gradient. Copeia 2000:178–190.

Richardson, J. L. 2006. Novel features of an inducible defense system in larval tree frogs

(*Hyla chrysoscelis*). Ecology 87:780–787.

Richter-Boix, A., G. A. Llorente, and A. Montori. 2007. A comparative study of predator-

induced phenotype in tadpoles across a pond permanency gradient. Hydrobiologia

583:43–56.

Schwarz, D., S. N. Gorb, A. Kovalev, N. Konow, and E. Heiss. 2020. Flexibility of intraoral

food processing in the salamandrid newt *Triturus carnifex*: effects of environment and

prey type. Journal of Experimental Biology 223:jeb232868.

Sherratt, T. N., and I. F. Harvey. 1989. Predation by larvae of *Pantala flavescens* (Odonata)

on tadpoles of *Phyllomedusa trinitatis* and *Physalaemus pustulosus*: the influence of

absolute and relative density of prey on predator choice. Oikos 56:170–176.

Smith, D. C., and J. Van Buskirk. 1995. Phenotypic design, plasticity, and ecological

performance in two tadpole species. The American Naturalist 145:211–233.

Steiner, U. K. 2007. Linking antipredator behaviour, ingestion, gut evacuation and costs of

predator-induced responses in tadpoles. Animal Behaviour 74:1473–1479.

Steiner, U. K., and J. Van Buskirk. 2009. Predator-induced changes in metabolism cannot

explain the growth/predation risk tradeoff. PLoS One 4:e6160.

Takahara, T., Y. Kohmatsu, A. Maruyama, H. Doi, H. Yamanaka, and R. Yamaoka.

2012a. Inducible defense behavior of an anuran tadpole: cue-detection range and cue

types used against predator. Behavioral Ecology 23:863–868.

Takahara, T., Y. Kohmatsu, A. Maruyama, and R. Yamaoka. 2008. Benefit of suites of

defensive behavior induced by predator chemical cues on anuran tadpoles, *Hyla*

japonica. Behavioral Ecology and Sociobiology 63:235–240.

Takahara, T., Y. Kohmatsu, A. Maruyama, and R. Yamaoka. 2012b. Predator-specific

inducible morphological changes in two Japanese anuran tadpoles. Herpetology Notes

5:43–47.

Tanahashi, M., and T. Fukatsu. 2018: *Natsumushi*: An image measuring software for

entomological studies. Entomological Science 21:347–360.

Teplitsky, C., S. Plenet, and P. Joly. 2003. Tadpoles' responses to risk of fish introduction.

Oecologia 134:270–277.

Teplitsky, C., S. Plenet, and P. Joly. 2004. Hierarchical responses of tadpoles to multiple

predators. Ecology 85:2888–2894.

Teplitsky, C., S. Plenet, J. P. Léna, N. Mermet, E. Malet, and P. Joly. 2005. Escape

behaviour and ultimate causes of specific induced defences in an anuran tadpole. Journal

of Evolutionary Biology 18:180–190.

Thibaudeau, G., and R. Altig. 2012. Coloration of anuran tadpoles (Amphibia):

development, dynamics, function, and hypotheses. International Scholarly Research

Notices 2012:1–12.

Touchon, J. C., and K. M. Warkentin. 2008. Fish and dragonfly nymph predators induce

opposite shifts in color and morphology of tadpoles. Oikos 117:634–640.

Touchon, J. C., and K. M. Warkentin. 2011. Thermally contingent plasticity: temperature

alters expression of predator-induced colour and morphology in a Neotropical treefrog

tadpole. Journal of Animal Ecology 80:79–88.

Urban, M. C., J. L. Richardson, N. A. Freidenfelds, D. L. Drake, J. F. Fischer, and P. P.

Saunders. 2017. Microgeographic adaptation of Wood Frog tadpoles to an apex predator.

Copeia 105:451–461.

Van Buskirk, J. 2002. A comparative test of the adaptive plasticity hypothesis: relationships

between habitat and phenotype in anuran larvae. The American Naturalist 160:87–102.

Van Buskirk, J., P. Anderwald, S. Lüpold, L. Reinhardt, and H. Schuler. 2003. The lure

effect, tadpole tail shape, and the target of dragonfly strikes. Journal of Herpetology

37:420–424.

Van Buskirk, J., J. Aschwanden, I. Buckelmüller, S. Reolon, and S. Rüttiman. 2004. Bold

tail coloration protects tadpoles from dragonfly strikes. Copeia 2004:599–602.

Van Buskirk, J., and S. A. McCollum. 1999. Plasticity and selection explain variation in

tadpole phenotype between ponds with different predator composition. Oikos 85:31–39.

Van Buskirk, J., and S. A. McCollum. 2000a. Functional mechanisms of an inducible

defence in tadpoles: morphology and behaviour influence mortality risk from predation.

Journal of Evolution Biology 13:336–347.

Van Buskirk, J., and S. A. McCollum. 2000b. Influence of tail shape on tadpole swimming

performance. Journal of Experimental Biology 203:2149–2158.

Wilson, R. S., P. G. Kraft, and R. Van Damme. 2005. Predator-specific changes in the

morphology and swimming performance of larval *Rana lessonae*. Functional Ecology

19:238–244.

Yager, D. D. 1999. Structure, development, and evolution of insect auditory systems.

Microscopy research and Technique 47:380–400.

Yuasa, Y. 2006. On the red tails of tadpoles of the frog, *Hyla japonica*. Hyogo Freshwater

Biology 58:87–90.

Yuasa, Y. 2007. Food of red-tailed larvae of the Japanese tree frog *(Hyla japonica)*. Hyogo

Freshwater Biology 59:143–144.

Fig. 1. Photographs of tadpoles of *Dryophytes japonicus* reared without predator (top) and

with *Anax nigrofasciatus* (bottom) and their corresponding Photoshop Lab color histograms.

The mean values of tail coloration are L = 61.2, a* = -2.5, and b* = 7.96 (without predator),

and L = 57.96, a* = 1.75, and b* = 35.51 (with *A. nigrofasciatus*), respectively.

Fig. 2. (A) Scatterplots of tail coloration parameters (a* and b* values) of tadpoles two weeks (top) and three weeks (bottom) after the start of the Experiment I, using different dragonfly species. Tadpoles were reared in the presence of no predators (the control group; Ct1), *Anax nigrofasciatus* (Ax), *Gynacantha japonica* (Gy), *Orthetrum albistylum speciosum* (Or), or *Pantala flavescens* (Pf). (B) Box-and-whisker plots displaying the range of the tail color traits (a* and b* values) two weeks (top) and three weeks (bottom) after the start of the Experiment I. The asterisks in the figure indicate that there were significant differences compared to the control group, which were led by the LMM analysis (see Materials and Methods). *$P < 0.05$, ***$P < 0.001$.

Fig. 3. (A) Scatterplots of tail coloration parameters (a* and b* values) of tadpoles two

weeks (top) and three weeks (bottom) after the start of the Experiment II, using non-odonate

species. Tadpoles were reared in the presence of no predators (the control group; Ct2), water

bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt

(*Cynops pyrrhogaster*; Fn). (B) Box-and-whisker plots displaying the range of the tail color

traits (a* and b* values) three weeks after the start of the Experiment II.

Fig. 4. Box-and-whisker plots showing the range of tadpole morphological traits, BL (body

length), TAL (tail length), and TD (tail depth) in Experiment I, using different dragonfly

species. Tadpoles were reared in the presence of no predators (the control group; Ct1), *Anax*

nigrofasciatus (Ax), *Gynacantha japonica* (Gy), *Orthetrum albistylum speciosum* (Or), or

Pantala flavescens (Pf). TAL and TD values were divided by body length to account for size

differences among individual tadpoles. The asterisks in the figure indicate that there were

significant differences compared to the control group, which were led by the LMM analysis

(see Materials and Methods). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Fig. 5. Box-and-whisker plots showing the range of tadpole morphological traits, BL (body length), TAL (tail length), and TD (tail depth) in Experiment II, using non-odonate predators. Tadpoles were reared in the presence of no predators (the control group; Ct2), water bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt (*Cynops pyrrhogaster*; Fn). TAL and TD values were divided by body length to account for size differences among individual tadpoles. The asterisks in the figure indicate that there were significant differences compared to the control group, which were led by the LMM analysis (see Materials and Methods). ***$P < 0.001$.

Fig. 6. Plots of principal components 1 and 2, with their contribution shown in parentheses, based on four morphological traits, total length [TL; the sum of body length (BL) and tail length (TAL)], standardized body length (sBL; BL/TL), standardized tail length (sTAL; TAL/TL), and standardized tail depth (sTD; TD/TL), and the coloration values (a* and b*), sampled three weeks after the start of Experiment I (left) and Experiment II (right). In Experiment I, tadpoles were reared in the presence of no predators (the control group; Ct1), *Anax nigrofasciatus* (Ax), *Gynacantha japonica* (Gy), *Orthetrum albistylum speciosum* (Or), or *Pantala flavescens* (Pf). In Experiment II, they were reared in the presence of no predators (the control group; Ct2), water bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt (*Cynops pyrrhogaster*; Fn). Ellipses represent the 95% probability ellipses of each treatment group. The red arrows indicate the direction in which the variables of each trait contribute to the principal components. Data for one individual in the Ax group, which appeared as an outlier in the bottom left-hand side, was omitted in this figure.

864   **Table 1.** Results of principal component analysis of coloration and morphological traits of

865   tadpoles in Experiment I (top) and Experiment II (bottom). PC1, PC2, and PC3 indicate the

866   first three principal components. The values in the table are the correlation coefficients of

867   each morphological trait with the principal component scores.

Experiment I

Response variable	PC1	PC2	PC3
Total Length	−0.72	−0.34	0.19
Standardized body length	0.93	−0.28	0.10
Standardized tail length	−0.93	0.28	−0.10
Standardized tail depth	0.05	−0.84	0.07
a*	0.09	0.43	0.88
b*	−0.44	−0.65	0.31
% of variance explained	41.0%	26.3%	15.7%

Experiment II

Response variable	PC1	PC2	PC3
Total Length	0.85	0.18	0.06
Standardized body length	−0.78	0.57	0.22
Standardized tail length	0.78	−0.57	−0.22
Standardized tail depth	0.11	0.68	−0.72
a*	−0.77	−0.43	−0.15
b*	0.74	0.45	0.36
% of variance explained	51.8%	25.7%	12.9%

Table 2. The Pearson's correlation coefficients between morphological trait pairs, total

length [TL; the sum of body length (BL) and tail length (TAL)], standardized body length

(sBL; BL/TL), standardized tail length (sTAL; TAL/TL), and standardized tail depth (sTD;

TD/TL), and the coloration values (a* and b*). Diagonal cells, representing self-correlations,

are denoted by dashes. Results from Experiment I are shown above the diagonal, while results

from Experiment II are presented below the diagonal.

Experiment I

	TL	sBL	sTAL	sTD	a*	b*
TL	–	−0.47	0.47	0.18	−0.07	0.43
sBL	−0.51	–	−1.00	0.20	0.03	−0.18
sTAL	0.51	−1.00	–	−0.20	−0.03	0.18
sTD	0.16	0.14	−0.14	–	−0.21	0.34
a*	−0.62	0.33	−0.33	−0.26	–	−0.10
b*	0.67	−0.26	0.26	0.16	−0.72	–

Experiment II