Predator-Induced Tail Coloration Toward Diverse Dragonfly Nymphs in Tadpoles of the

East Japan Tree Frog (*Dryophytes leopardus*)

Akihiro Noda[1,] and Katsutoshi Watanabe[1]

Running head: **Predator-induced tail coloration in tadpoles of *Dryophytes leopardus***

1.   Department of Zoology, Graduate School of Science, Kyoto University, Kitashirakawa-

oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan.

(AN)Email: kapurira@outlook.jp. Send reprint requests to this address

Abstract

Anuran larvae can change their coloration and morphology to avoid predation.

Particularly, some tadpoles in the genus *Dryophytes* in the Americas develop deep tail fins and

bright orange tail coloration in response to predators such as dragonfly nymphs. These

conspicuous tails are hypothesized to attract predator attacks, protecting vital body parts from

fatal injuries. However, it remains unclear whether the extent of the induced phenotypic

changes differs between various predators with distinct feeding patterns and habitats. To

explore this, we reared tadpoles of the East Japan tree frog, *Dryophytes leopardus*, with four

dragonfly species, water bug, water scorpion, or newt, and measured phenotypic changes,

especially focusing on tail coloration. We found that the aeshnid *Anax nigrofasciatus*, a sit-

and-wait predator that climbs on aquatic vegetation, induced bright orange and broad tails of

tadpoles. Other odonate species that forage in and on the substrate also elicited tail coloration

or deeper larval tails, but it took more time to induce the tadpoles' response compared to *A.*

nigrofasciatus. In contrast, non-odonate predators did not induce color and morphological

changes. This study provides the first evidence of predator-induced tail coloration in an Asian

species of *Dryophytes*, which appears to be a defensive response specifically to dragonfly

nymphs, as visually adept predators. Moreover, the differences in the extent and timing of the

phenotypic changes among odonate species suggest that anti-predator phenotypes are

modulated based on relative predation risks, considering the predators' foraging patterns and

visual capabilities, thereby allowing tadpoles to minimize physiological costs and unintended

risks associated with the induced traits.

The utilization of prey's body color and shape is highly dependent on its context, including the sensory capabilities of predators and the background condition (e.g., Espanha et al., 2016; Barnett et al., 2018). Especially in aquatic habitats, substantial variation exists in background conditions and the composition and abundance of predators across different localities, making it difficult for prey to achieve a uniform predation avoidance ability across their range (Van Buskirk, 2002; McIntyre et al., 2004). Therefore, such environmental uncertainty has generally been believed to favor plastic rather than constitutive anti-predator defenses (Agrawal and Karban, 1999; McIntyre et al., 2004).

Anuran larvae, or tadpoles, have served for decades as a model system for investigating induced defense mechanisms, as well as predator avoidance strategies and trade-offs mediated by plastic traits (Kasumovic, 2013; Kruger and Morin, 2020). They often rely on chemical cues to evaluate predation risk through continually released predator-borne cues and digestion-released, prey-borne cues, and then exhibit predator-induced defense mechanisms (Brönmark and Hansson, 2000; Hettyey et al., 2010, 2015). Predator-induced plastic responses in tadpoles can be categorized into four primary domains: behavioral, morphological, physiological, and chemical defenses. Behavioral responses encompass changes in overall activity levels (e.g., Lawler, 1989; Takahara et al., 2012a; Reuben and Touchon, 2021), increased burst swimming speed (Takahara et al., 2008), and spatial avoidance of predators (e.g., Laurila et al., 1998; Nunes et al., 2013). Morphological changes for predator avoidance have been well documented in tadpoles of various species (Relyea, 2001; Van Buskirk, 2002; Richter-Boix et al. 2007). For instance, tadpoles undergo changes in body size (i.e., correlated with growth rate and metamorphosis timing), body shape such as the development of deeper tail fins (Lardner, 1998, 2000), and coloration (McCollum and Van Buskirk, 1996; Touchon and Warkentin, 2008). Behavioral and morphological responses are sometimes co-expressed (Hossie et al., 2017). Physiological responses include changes in

respiration rate and gut evacuation rate (Steiner, 2007; Steiner and Van Buskirk, 2009;

Costello and Michel, 2013). These may arise due to trade-offs in predator avoidance (Relyea

and Auld, 2004) or as responses to partially offset the costs of lower energy intake resulting

from evasion (Barry and Syal, 2013). Additionally, tadpoles of certain groups, particularly

toads, evade predation through toxin production (Bókony et al., 2016; Hettyey et al., 2019).

The types and intensities of induced responses in tadpoles often differ according to

predator types (Relyea, 2001; Teplitsky et al., 2004, 2005; Hettyey et al., 2011). For instance,

Kishida and Nishimura (2005) revealed that tadpoles of *Rana pirica* developed deeper tail

fins in response to dragonfly nymphs, whereas in the presence of salamander larvae, they

exhibited not only deeper tails but also bulging heads and bodies. Tadpoles expressing these

predator-specific traits should have a survival advantage over those with mismatched or non-

induced phenotypes when faced with predation. Conversely, expressing such defense traits

incurs energetic costs and can increase vulnerability to non-focal predators (Touchon and

Warkentin, 2008; Innes-Gold et al., 2019). Due to these costs, defense traits are expected to

remain plastic rather than fixed, and their types and intensities can vary depending on

predator species, sometimes exhibiting predator-specific responses.

Dragonfly nymphs, as one of the major predators of tadpoles, are known to induce

morphological responses characterized by deep tail fins across many anuran species (Relyea

and Werner, 2000; Van Buskirk, 2002). This induced broad tail has been repeatedly

hypothesized to improve the swimming performance of tadpoles and facilitate their escape

from predators (Smith and Van Buskirk, 1995; McCollum and Leimberger, 1997; Van Buskirk

and McCollum, 2000a). However, experimental evidence of this benefit is lacking (Van

Buskirk et al., 2003; Johnson et al., 2008). Moreover, a few tree frog species in the genus

Dryophytes respond to odonate predators by not only exhibiting deeper tails but also turning

their entire tails bright orange (McCollum and Van Buskirk, 1996; Van Buskirk and

McCollum, 1999; LaFiandra and Babbitt, 2004). Thus, as another function, the induced deep

and brightly colored fin is thought to serve as a lure, diverting predator attacks away from the

more vulnerable head and body regions (Van Buskirk et al., 2003, 2004; Hawlena et al., 2006;

Johnson et al., 2008). Given that tail fins tear easily (Doherty et al., 1998) and that tadpoles

can maintain their swimming performance even after partial tail loss (Hoff and Wassersug,

2000; Van Buskirk and McCollim, 2000b), the tail is considered expendable relative to other

body parts. Because dragonfly nymphs are predominantly visual predators, exhibiting the

ability to discern both the shape and color of immobile as well as moving prey (Bybee et al.,

2012; Espanha et al., 2016), color and shape changes in tadpoles are likely to be effective in

avoiding predation by odonates.

Among dragonfly nymphs, several ecological types have been identified based on

microhabitat usage and foraging behavior (Pritchard, 1965; Corbet, 1999), including climbers

(e.g., Green darner, *Anax junius*), moving among aquatic vegetation and ambushing prey on

plants; sprawlers (e.g., Globe skimmer, *Pantala flavescens*; Sherratt and Harvey, 1989),

inhabiting pond bottoms and swimming through the water column to attack; and burrowers

(e.g., Golden-ringed dragonfly, *Cordulegaster boltonii*), burrowing in mud or bottom debris

and often lying in wait for prey with only the eyes protruding. The reliance on vision for

foraging varies among these types (Pritchard, 1966; Corbet, 1999). For tadpoles, the

frequency of encounters with and relative predation risk of these predators differ among

habitats and seasons. These differential predation risks likely determine the cost−benefit

balance of induced traits, thereby suggesting that tadpoles may exhibit different defense

responses. However, prior experimental studies have primarily focused on a single dragonfly

species and have not examined how tadpole responses may vary among odonate (and other)

species.

The East Japan tree frog, *Dryophytes leopardus*, which was recently recognized as

distinct from *Dryophytes japonicus*, is a prevalent species with a wide distribution

encompassing northeastern Japan (central to eastern Honshu and Hokkaido), as well as

regions outside Japan including Sakhalin Island (Shimada et al., 2025). Tadpoles of this

species inhabit various environments such as rice paddies, marshes, ponds, puddles, and other

ephemeral water bodies (Matsui and Maeda, 2018). Tadpoles, including this species, serve as

important prey for numerous aquatic predators, including dragonfly nymphs, water scorpions,

and red-bellied newts (Ohba and Nakatsuji, 2006; Hirai et al., 2020; Kita et al., 2020; Ozaki

and Nishikawa, 2023). Morphological responses to predators have also been documented in

this species, with dragonfly nymphs inducing deep tail fins, while chemical cues from

goldfish prompt the development of shallow fins (Takahara et al., 2012b). The occurrence of

orange tail coloration has also been briefly reported, although it has been attributed to

ingested algae rather than predator-induced plasticity (Doi et al., 2005; Yuasa, 2006, 2007).

Despite being rarely documented, tadpoles exhibiting brightly colored tails were frequently

observed at our study site in Kyoto, central Japan (Noda, personal observation).

To clarify the cause and function of the orange tail coloration in tadpoles of *D.*

leopardus, we conducted laboratory experiments with the tadpoles and several predator

species. These experiments aimed to achieve two primary objectives. First, we investigated

whether the tail coloration of tadpoles is induced by dragonfly nymphs, and if so, whether the

intensity and speed of this response vary among different odonate species with different

ecological characteristics. Simultaneously, we examined changes in body shape that are

known to be plastically induced by predators. The second objective was to assess whether

tadpoles exhibit color/morphological responses to predators characterized by different

foraging styles compared to odonates. We used water bugs (*Appasus major*), water scorpions

(*Laccotrephes japonensis*), and newts (*Cynops pyrrhogaster*) as non-odonate predators. Water

bugs and scorpions capture prey by enclosing them with their frontal legs, and newts swallow

prey whole. Accordingly, predation avoidance strategies involving tail coloration, which are

based on diverting attacks toward a specific body part, are expected to be less effective

against these predators than against dragonfly nymphs. Therefore, tadpoles may exhibit

weakened or no color/morphological responses to minimize the costs associated with

developing plastic traits. In both experiments, tadpoles were reared with a single predator

species, and their morphological traits of interest were compared with those of tadpoles raised

in predator-free tanks. By analyzing the extent of tail coloration in tadpoles exposed to diverse

predators, we revealed that tadpoles modulate their induced responses according to the risks

posed by predators with different foraging types and sensory capabilities.

MATERIALS AND METHODS

Two series of laboratory experiments were carried out: Experiment I involved the

comparison of induction of color/morphological changes in tadpoles by four different

dragonfly species as predators, and Experiment II employed three non-odonate predators.

Experiment I.—Seven mating pairs of *Dryophytes leopardus* were captured in the paddy

fields at Kyoto University (35°01'N, 135°47'E) on June 9 and 10, 2022. Following capture,

mating pairs were housed overnight in plastic cages (19 × 19 × 14 cm; 6 cm water depth), and

eggs were obtained the following day. The eggs hatched in about 2−3 days, and after hatching,

tadpoles were separately kept in plastic containers (43 × 31 × 16 cm; 6 cm water depth), as

those from the same clutch were kept together in a single container. A room temperature of

20°C was maintained until the experiments began, and tadpoles were fed powdered food

(Kyorin Co., Ltd.) every two days. In this experiment, four species of dragonfly nymphs were

used as predators taking into account their foraging styles: *Anax nigrofasciatus* (Black-striped

lesser emperor; climber), *Gynacantha japonica* (Blue-spotted dusk-hawker; climber),

Orthetrum albistylum speciosum (White-tailed skimmer; burrower), and *Pantala flavescens*

(Wandering glider; sprawler). Nymphs of *A. nigrofasciatus* collected in Sakura City, Chiba

Prefecture, and those of *G. japonica* collected in Higashihiroshima City, Hiroshima

Prefecture, were obtained from an online vendor, and *O. a. speciosum* and *P. flavescens* were

collected from the same paddy field as the mating frog pairs. The dragonfly nymphs were

maintained individually in plastic trays (23 × 17 × 8 cm) and fed tadpoles of *D. leopardus*

every two days until the beginning of the induction experiment.

The induction experiment commenced on July 12, 2022. Each experimental tank

housed 21 tadpoles, with three individuals randomly selected from each of the seven egg

clutches to randomize genetic effects in each tank. Tadpoles in a tank were assigned to one of

five treatments (Ax group with *A. nigrofasciatus*; Gy group with *G. japonica*; Or group with

O. a. speciosum; Pf group with *P. flavescens*; and the control Ct1 group) with eight replicates

per treatment. Each experimental tank (31 × 18 × 24 cm; 10 cm water depth) was equipped

with a transparent isolation case (17 × 9 × 10 cm) with approximately 5 mm diameter holes

spaced about 5 mm apart, where one dragonfly nymph was introduced. This setup allowed

visual and chemical-mediated interaction between the predator and tadpoles without physical

contact. The tanks were aerated and maintained under a 14-h light/10-h dark photoperiod at a

room temperature of 22°C, with half of the water in all tanks changed every three days. Air

was continuously supplied to the isolation case within each tank through an aquarium air

stone. Tanks designated for the Ct1 group lacked predators and utilized an empty isolation

case. Tadpoles in the experimental tanks were fed every two days with commercial food for

Axolotl, *Ambystoma mexicanum* (Kyorin Co., Ltd.), while dragonfly nymphs received two

tadpoles as food every two days.

Experiment II.—The design of Experiment II mirrored that of Experiment I, with

adjustments made for sampling dates, the interval between hatching and the experiment's

commencement, and the selection of predators. For this experiment, three mating pairs of *D.*

leopardus were captured from the paddy fields at Kyoto University on May 18 and 20, 2022.

Eggs were collected and tadpoles from clutches were reared in the same way as in Experiment

I until the experiments began. Three predator species were chosen based on their foraging

style and prey capture behavior: Ferocious water bug, *Appasus major* (sit-and-wait predator

that seizes prey with its frontal legs), water scorpion, *Laccotrephes japonensis* (sit-and-wait

predator that grasps prey with its frontal legs), and Japanese fire belly newt, *Cynops*

*pyrrhogaste*r (active tracker that swallows prey whole). Aquatic insects, collected in

Higashihiroshima City, were purchased from the online vendor, while red-bellied newts were

collected from irrigation ditches in Kyoto City (35°05'N 135°47'E). These predators were

individually housed in plastic trays and fed with tadpoles every two days.

The induction experiment began on June 18, 2022. Each experimental tank housed

21 tadpoles, with seven individuals randomly selected from each of the three egg clutches.

Tadpoles in each tank were allocated to one of four treatments (Be group with *A. major*; Ws

group with *L. japonensis*; Fn group with *C. pyrrhogaster*; and the control Ct2 group) with

eight replicates per treatment. One predator was introduced into an isolation case in each tank.

Other experimental setups, i.e., tank design, room temperature, light condition, and feeding

frequency, were kept the same as in Experiment I.

Measuring Coloration and Morphological Responses.—All tadpoles were photographed two

and three weeks after the start of the experiment using a Canon® EOS RP mirrorless camera.

Individual tadpoles were placed in a narrow acrylic aquarium (20.5 × 4.5 × 13.5 cm) with a

ruler and color chart in the background. To minimize the effect of posture variation among

individuals, multiple photographs were taken per tadpole, and the image most suitable for

measurement was selected. Each trait was measured once per individual. The distance

between the aquarium and the camera was kept constant for all individuals. All photos were

taken with a F/16 aperture, a 1/20th second exposure time, and ISO 3200. Consistency in

artificial lighting conditions was maintained for individuals photographed on the same day,

with the white balance manually set and standardized at 4600 K. Some individuals were

excluded from the analysis due to poor nutritional condition or significant tail loss during

rearing. The exclusion criteria included: (1) head and body length less than 6 mm, (2) severe

tail damage or loss, and (3) inability to maintain posture in water.

Tail fin parts, defined as all tail areas excluding the central tail muscle, were trimmed

manually using the image measuring software, Natsumushi version 1.12.2 (Tanahashi and

Fukatsu, 2018; https://sites.google.com/site/mtahashilucanid/program/natsumushi) and

selected for color measurement. The color of the tail was quantified using the average of

CIELAB values (CIE 1976) calculated in Adobe Photoshop 25.5.1 (Adobe Inc.). In the

CIELAB color space, color was represented using three components: the lightness component

L* and the color components a* and b*. The L* component ranged from black to white

between 0–100, while a* and b* components ranged from green to red and blue to yellow,

respectively, on a scale of −128 to 127. As Photoshop displays images in 8-bit format, these

values were rescaled to range from 0 to 255. Therefore, L* was multiplied by 100/255 and a*

and b* were adjusted by subtracting 128, respectively, to facilitate analysis (Fig. 1). Due to

incorrect white balance, photographs taken at the second week of Experiment II were

excluded from the analysis of tail coloration. During the third week of Experiment II,

photographs of 11 individuals from the Ct2 group were also taken with an incorrect white

balance, so the color tone was later corrected using Photoshop. We used the "Match Color"

function in the "Image Adjustments" menu, referencing one of the other photographs from the

Ct2 group that was taken under the same conditions, with the white balance properly set to

4600 K. Additionally, morphological measurements were conducted based on the photographs

using Natsumushi. Morphological traits included tadpole body length (BL), tail length (TAL),

and tail depth (TD) were measured using the scale within each image.

Statistical Analyses.—All statistical analyses were conducted in R version 4.3.2 (R Core

Team, 2023). Linear mixed models (LMMs) were employed based on the measured traits,

using the lmerTest package (Kuznetsova et al., 2017) to estimate differences between the

control group and the other treatments. The experimental tank was treated as a random factor

in the LMMs. When color components (e.g., a* and b*) were utilized as the response

variables, the treatment (predator) was used as the explanatory variable in the models to

compare the treatment groups to the control group. In the same way, when body length (BL)

was utilized as the response variables, the treatment was used as the explanatory variable in

the models. On the other hand, in analyses for TAL and TD as the response variable, four

models were constructed for each trait. These models included: the model with only BL as the

explanatory variable, the model with only the treatment as the explanatory variable, the model

with both BL and treatment as explanatory variables, and the model incorporating BL,

treatment, and their interactions. The final model was determined via backward elimination,

by applying likelihood ratio (LR) tests. The significance level was set at 5%. If a significant

difference was found between the former and reduced models, the former model was adopted

as the final model for further analysis.

After constructing final models, we evaluated the overall effect of predator treatment

by performing ANOVA via the anova function in the lmerTest package. Subsequently,

comparisons between each treatment group and the control were performed using Dunnett's

test with estimated marginal means calculated by the emmeans package (Lenth, 2025).

Marginal means were derived under the asymptotic framework. The significance level was set

at 5%, with treatment groups considered significantly different from the control group if the P

values were below this threshold. Orange tail coloration and increased tail depth were

reported as predator-induced traits in previous studies (McCollum and Van Buskirk, 1996;

Touchon and Warkentin, 2008), so we decided to focus primarily on b* values and tail depth.

To evaluate the covariation between tail coloration and morphology at the third week of the

experiment, we also conducted Pearson's correlation tests between residuals of b* values and

tail depth (TD), after removing the effects of body length (BL) via linear regression. This

analysis was performed separately for each predator treatment that showed a significant

difference from the control group in either trait.

RESULTS

Coloration.—Experiment I showed that predator presence induced changes in tadpole tail

coloration. At the second week, the overall effects of predator treatment were significant for

both a* and b* values (a*: $F = 11.41$, $P < 0.001$; b*: $F = 12.05$, $P < 0.001$). The response was

strongest with *Anax nigrofasciatus* (Ax group), where tadpoles developed significantly more

orange tails than controls, as indicated by a consistent decrease in a* and increase in b* values

(Dunnett's test, $P < 0.05$; Table 1; Fig. 2). At the third week, a significant overall effect of

predator treatment remained only in b* values ($F = 11.31$, $P < 0.001$), not in a* values ($F =$

2.18, $P = 0.096$). Nevertheless, tadpoles in the Ax group still showed significantly lower a*

values ($P = 0.028$) and higher b* values ($P < 0.001$) than controls. A few individuals in the Ax

group exhibited particularly high values in both a* and b*, corresponding to intensely vivid

orange tails. These data points likely represent the upper range of predator-induced color

expression, rather than artifacts. A weaker, yet still significant, effect was observed with

tadpoles exposed to *Orthetrum albistylum speciosum* (Or group), which showed elevated b*

values compared to the control ($P = 0.045$). In the tadpoles with *Pantala flavescens* (Pf

group), a similar trend was noted, but the differences from the control group were not

statistically significant ($P = 0.143$ in b* value). In contrast to the results of Experiment I,

overall treatment effects in Experiment II were not significant (a*: $F = 0.49$, $P = 0.689$; b*: F

$= 1.84$, $P = 0.166$), and no group showed significant differences from the Ct2 group in the a*

or b* values at the third week (Table 2; Fig. 3).

Morphological Responses.—In Experiment I, predators had minimal effects on body length

(BL). The overall effect of predator treatment on BL was not statistically significant at either

week (Week 2: $F = 2.35$, $P = 0.075$; Week 3: $F = 2.35$, $P = 0.074$), although a weak trend was

observed. Tadpoles exposed to *P. flavescens* (Pf) developed slightly larger bodies than

controls at the second week (Dunnett's test, $P = 0.048$), but this difference was no longer

significant at the third week (Table 3; Fig. 4). Similarly, although predator treatment had a

significant overall effect on tail length, its effects were limited to certain groups or time

points. When tail length was used as the response variable, models with body length and

treatment as explanatory variables were selected as the final models at both the second and

third weeks after the start of the experiment. In both models, tail length showed a strong

positive association with body length ($P < 0.001$), indicating that larger tadpoles tended to

have longer tails. In these models, predator treatment had a significant overall effect at both

the second week ($F = 6.76$, $P < 0.001$) and the third week ($F = 3.28$, $P = 0.023$). Independent

of BL, tadpoles in the Pf and *A. nigrofasciatus* (Ax) groups had shorter tails than controls at

the second week ($P < 0.001$), but not at the third week. In contrast, tadpoles exposed to

Gynacantha japonica (Gy group) showed reduced tail length only at the third week ($P =$

0.005).

The impact of predator treatment on tail depth was more pronounced. When tail

depth was used as the response variable, BL and treatment were chosen as explanatory

variables at both the second and third weeks after the start of the experiment. In both models,

tail depth also showed a strong positive association with BL ($P < 0.001$), and the effects of

predator treatments were interpreted independently of body size. Predator treatment had a

strong overall effect on tail depth at both the second week ($F = 4.86$, $P = 0.003$) and the third

week ($F = 9.60$, $P < 0.001$). Post-hoc tests revealed that tadpoles in the Ax group developed

deeper tails than controls at both time points ($P < 0.01$). Moreover, tadpoles in the Pf group

also developed deeper tail fins than controls at the third week ($P = 0.022$). A similar trend was

observed in the *O. albistylum* (Or) group, although this difference did not reach statistical

significance ($P = 0.113$).

In Experiment II, predators had minimal effects not only on BL, but also on tail

morphology. Predator treatment had no significant effect on BL (Week 2: $F = 1.82$, $P = 0.168$;

Week 3: $F = 2.18$, $P = 0.118$). For tail length, models including both BL and treatment as

explanatory variables were chosen at the second week. In this model, tail length showed a

strong positive association with BL ($P < 0.001$), and predator treatment had a significant

overall effect ($F = 7.43$, $P < 0.001$). Post-hoc tests revealed that a significant reduction in tail

length was observed in the Ferocious water bug (Be) group ($P < 0.001$; Fig. 5). At the third

week, however, models with only BL were selected, indicating that the effect of the treatment

on tail length appeared limited. Similarly, for tail depth, models with only BL were selected in

both weeks, and tail depth was positively associated with body length ($P < 0.001$). Post-hoc

tests also confirmed that tail depth was largely unaffected, with no significant differences

observed. Thus, tadpoles in all non-odonate predator treatments were not induced to develop

significantly deeper tail fins compared to the control group.

The orange coloration and deeper tail morphology tended to be co-expressed within

individuals. Significant, though weak, positive correlations between b* values and tail depth

were detected in all three odonate predator treatments that induced changes in either trait by

the third week (Pf: $r = 0.393$, $P < 0.001$; Ax: $r = 0.294$, $P = 0.003$; Or: $r = 0.277$, $P = 0.011$).

DISCUSSION

Tail Coloration in Asian Species of Dryophytes.—Our experiments revealed that tadpoles of

Dryophytes leopardus developed induced changes in tail coloration as well as tail shape in

response to dragonfly nymphs. Although tail coloration has rarely been documented in this

species or in its close relative, *Dryophytes japonicus*, its occurrence has been reported from

several locations in Japan, including Hokkaido, Shiga, Hyogo, Hiroshima, Shimane, Kouchi

and Ehime prefectures (Doi et al., 2005), as well as from our collection site in Kyoto (Noda,

personal observation). The phenotypes induced by *Anax nigrofasciatus*, i.e., deep tail fins and

bright orange tails, closely resemble those previously observed in other species in *Dryophytes*

in the New World, Cope's Gray Treefrog (*Dryophytes chrysoscelis*; McCollum and Van

Buskirk, 1996), Pine Woods Treefrog (*Dryophytes femoralis*; LaFiandra and Babbitt, 2004),

and Gray Treefrog (*Dryophytes versicolor*; Van Buskirk and McCollum, 1999). Thus, our

findings represent the first documented case of a predator-induced tail coloration phenotype in

Asian species of *Dryophytes*.

Our discoveries of the plasticity in the tail coloration in the Asian species of

Dryophytes suggests that the trait is symplesiomorphically shared between the North

American and ancestral Asian lineages, and has been inherited by present-day Asian species,

including *D. leopardus*. An ancestral stock of *Dryophytes* is inferred to have migrated from

North America to East Asia across the Bering Land Bridge during the Miocene (Duellman et

al. 2016; Dufresnes et al., 2016). Present-day Asian species of *Dryophytes* are monophyletic,

having diversified from this ancestral stock. It is plausible that tail coloration has persisted as

a phenotypic plasticity in both North American and East Asian species of *Dryophytes* due to

the consistent importance of dragonfly nymphs as predators.

It remains unknown whether tail coloration can be induced in the three other Asian

species, *Dryophytes flaviventris*, *Dryophytes immaculates*, and *Dryophytes suweonensis* (see

Borzée et al., 2020). In North America, Pine Barrens Treefrog, *Dryophytes andersonii* are

induced to have a darker tail, not an orange tail, by *Anax junius*, which induces orange tails in

other species of *Dryophytes* (Kruger and Morin, 2020). Furthermore, Squirrel Treefrog,

Dryophytes squirellus exhibits no morphological changes in response to odonate species

(McCoy and Bolker 2008; Baker, 2021). These cases indicate that the plastic responses in

Dryophytes have diversified, and some may have lost the ability to develop orange tails.

Future studies are needed to elucidate the interspecific variation in plastic responses to

predation risk and the evolutionary causes underlying this variation.

Differences in the Extent of Coloration by Predators.—Our experiments demonstrated that

the extent and timing of induced responses varied depending on the dragonfly species reared

with tadpoles. Two weeks after the start of the experiment, nymphs of *A. nigrofasciatus*

induced both deep fin and tail coloration in tadpoles. Additionally, by the third week, similar

but weaker responses were observed in the other predator treatments: *Pantala flavescens* also

induced deep tail and *Orthetrum albistylum speciosum* induced tail coloration. Furthermore,

although weak, some tadpole individuals exposed to *P. flavescens* showed tail coloration, and

those exposed to *O. albistylum speciosum* also tended to have deeper tails. These results

indicate that the defense traits (coloration and deep tail) were induced by the nymphs of *A.*

nigrofasciatus more rapidly and strongly, but similar responses were also elicited by the other

two predator species. Furthermore, the significant positive correlation observed between the

b* value and tail depth suggests that these traits tended to be co-expressed in a coordinated

manner. These findings support the idea that the expression of these anti-predator defenses is

governed by a shared, integrated strategy, whose intensity varies depending on predator

identity.

The differences in the degree and rate of responses induced by different dragonfly

species may reflect varying levels of predation risk for tadpoles posed by each species. Plastic

phenotypes with significant tail coloration should result in trade-offs because of energetic

costs associated with synthesizing pigmentation (Grether et al., 2001) or costs due to the

increased vulnerability to mismatched predators. Consequently, tadpoles are expected to

minimize these costs by developing effective defense traits for each predator as much as

possible based on the perceived predation risk (Touchon and Warkentin, 2008; Innes-Gold et

al., 2019). The predation risk for tadpoles would be different among dragonfly species

according to (1) the foraging style of dragonfly nymphs, (2) their color vision, and (3) the

total amount of intake related to their body size and occurrence period.

Firstly, the four dragonfly species used in our experiment exhibit different foraging

styles. According to the classification by Corbet (1999), *A. nigrofasciatus* and *G. japonica* are

climbers, sit-and-wait predators on aquatic vegetation, while *P. flavescens* is a sprawler,

foraging while moving along the bottom, and *O. albistylum speciosum* is a burrower, residing

within the substrate and ambushing prey. For sit-and-wait predators, there is often a time lag

between identifying prey and launching an attack, during which they can choose which part of

tadpole's body to target. Therefore, the lure effect of tadpole tails is likely to attract ambush

predators more effectively than those that actively chase their prey. In addition, as *A.*

nigrofasciatus prefers habitats abundant in floating-leaved plants (Ozono et al., 2021), it can

move freely along the plant structures from roots anchored at the bottom to leaves extending

above the water surface. As a result, the risk of predation on tadpoles could be present

throughout the entire water column, from the bottom to the surface, making behavioral

responses like moving to a specific depth insufficient as a means of avoiding predation.

Conversely, *P. flavescens* and *O. albistylum speciosum* primarily forage at the bottom, so

tadpoles may evade these predators by relocating to higher water layers. However, such

avoidance behavior often leads to a decrease in growth rate due to reduced foraging efficiency

(Teplitsky et al., 2003); i.e., for tadpoles that primarily feed on detritus, avoiding the bottom

layer may result in malnutrition. The combination of moderate morphological responses with

behavioral responses (the "double defense" strategy; Dewitt and Hucko, 1999) may help

mitigate the impact of malnutrition from behavioral changes while minimizing the costs of

defensive traits.

Secondly, differences in eyesight, particularly color vision among dragonfly species,

may influence the extent of the tail coloration changes in the tadpoles. Pritchard (1966) noted

a correlation between foraging habit and compound eye size in dragonfly nymphs, with

climbers typically possessing larger compound eyes compared to sprawlers and burrowers. He

also stated that the climbers, such as species in *Anax*, rely heavily on vision for prey

detection, whereas other foraging types depend more on tactile stimuli. Hence, it is plausible

that the lure effect of the brightly colored tails is more effective for climbers, which are more

visually oriented. Furthermore, Futahashi et al. (2015) investigated the genetic basis of color

vision in dragonflies and redocumented the number of opsin genes and the characteristic

spatiotemporal expression patterns in the nymphs of various species, including those of the

burrower *O. albistylim* and the climber *A. parthenope*, a species closely related to *A.*

nigrofasciatus used in our experiment. They found that dragonflies possess a greater number

of opsin genes than other insects, and *A. parthenope* possesses and expresses a greater number

of opsin genes, particularly those sensitive to long-wavelength light. Although the functions

of these diverse genes are not fully understood, this suggest that species of *Anax* may have

enhanced abilities to perceive reddish colors, potentially enabling tadpoles with bright orange

tails to more effectively avoid predation by using their tails as lures. Thus, the larval

antipredator defense of *D. leopardus*, which involves tail coloration, may have been driven by

the sensory systems of predatory dragonfly nymphs, and subsequently evolved specifically

against color-sensitive predators, represented by dragonfly nymphs of *Anax*.

Thirdly, the variation in the degree of induced responses may reflect the relative

predation risks associated with body sizes or occurrence periods of respective odonate

predators. In nature, it is generally difficult to determine which predators exert higher

predation pressure on tadpoles, and few studies have investigated such differences. However,

during the rearing process, we observed that the nymphs of *A. nigrofasciatus* exhibited higher

food demands and displayed more frequent attacks on tadpoles compared to other species.

Additionally, they are approximately twice the size of the nymphs of *P. flavescens* and *O.*

albistylum speciosum (see Ishida et al., 1988), suggesting that size-related predation pressures

could influence the necessity and effectiveness of tadpole morphological responses. However,

tadpoles did not exhibit significant responses to *G. japonica*, which is similar in size to *A.*

nigrofasciatus (see Ishida et al., 1988). This indicates that tadpoles likely respond not only to

size-related predation pressure but also to other characteristics of dragonfly species, such as

foraging style or habitat overlap, thereby modulating the extent of induced responses. Since

the habitat of nymphal *G. japonica* overlaps with that of tadpoles of *D. leopardus* (Nakagawa

et al., 2004), the lack of an apparent induced response may be due to the limited period in

which they co-occur. This species grows more rapidly compared to other species in Aeshnidae

in Japan, occurring in aquatic environments only from May to August (Ozono et al., 2021).

Considering that the nearly-metamorphosing nymphs cease foraging, the period during which

this species poses a predation threat to *D. leopardus* is even shorter. This brief coexistence

period suggests that this dragonfly species may not be an important predator for the tadpoles.

In addition to the tail coloration and depth we focused on, a tendency toward larger

body lengths and shorter tail lengths compared to those in the control group was observed,

although not statistically significant in most treatment groups. These changes may represent

general responses to predator treatments, as noted in previous studies. The increase in body

length could be attributed to enhanced availability of food resources resulting from leftovers

and feces of caged predators (e.g., McIntyre et al., 2004; Richardson, 2006). Similarly, shorter

tail lengths in the presence of predators have been reported in several frog species (Laurila et

al., 2002; LaFiandra and Babbitt, 2004).

Responses Toward Non-Odonate Predators.—Tadpoles of *D. leopardus* showed neither

conspicuous tail coloration nor deep fin when exposed to newts or aquatic insects other than

dragonfly nymphs in our experiment, likely because the costs of expressing these defenses

outweighed their benefits for avoiding predation. Although Experiment II did not include a

positive control (i.e., treatment with dragonfly nymphs, such as *A. nigrofasciatus*), we believe

that the absence of color induction was unlikely caused by experimental failure, because

preliminary experiments demonstrated the high reproducibility of the induced responses when

using dragonfly nymphs. Therefore, our results suggest that changes in body color and tail

depth are not strongly induced by these predators and may not serve as effective predation

avoidance strategies against them. An important point in understanding why tadpoles did not

show tail coloration is that defensive traits come with physiological costs and may increase

predation pressure from non-focal predators. If changes in tail color and shape do not

significantly contribute to non-odonate predator avoidance, tadpole should minimize these

disadvantages by refraining from developing these antipredator defenses.

485   Given that the non-odonate predators do not target specific body parts of their prey

during attacks, the strategy of using colored deep tails as lures may be ineffective and provide

only limited benefits against them. For instance, diving beetles, water bugs, and water

scorpions capture prey by grasping the entire body with their arms, so the tails of tadpoles

should not serve as lures for them. Additionally, they use tactile, visual, and olfactory senses

for locating prey (Babbitt and Jordan, 1996; Kehr and Gómez, 2009; Mogali et al., 2019;

Nowińska and Brożek, 2019), and possess well-developed organs for detecting auditory and

mechanical signals (Yager, 1999; Cocroft and Rodríguez, 2005). Previous studies have

reported predator-induced changes in larval morphology in response to water bugs (Benard,

2006; McCoy, 2007; Michel, 2012), but few have reported effects on larval tail coloration

(McIntyre et al., 2004; Touchon and Warkentin, 2011). Similarly, newts and salamanders, both

as larvae and adults, are active foragers that swallow their prey whole like fish (Wilson et al.,

2005; Kishida and Nishimura, 2007). Consequently, the tails of the tadpoles are unlikely to

function as a lure for these predators. Newts tend to rely on olfactory cues as well as vision

for food recognition (Copeland, 1913; Lindquist and Bachmann, 1982; Schwarz et al., 2020).

Few studies have reported that tadpoles exhibit induced defensive traits in response to these

predators, and none have documented induced changes in body coloration (Kishida and

Nishimura, 2004; Hettyey et al., 2011; Urban et al., 2017). These facts support the hypothesis

that the brightly colored tails of *D. leopardus* do not function as lures against the non-odonate

predators, and that, as a result, tadpoles do not express costly defensive traits in response to

them.

506   The lack of induction of defensive phenotypes may also be attributed to

physiological costs associated with developing tail coloration, although the magnitude of such

costs in tadpoles remains poorly understood. Since most animals are unable to biosynthesize

carotenoids, the pigments thought to be causing the bright orange coloration, they metabolize

and accumulate these pigments by absorbing them from their diet (Bertrand et al., 2006;

Thibaudeau and Altig, 2012). Thus, the allocation costs of expressing coloration by

transporting these pigments may not be as high as commonly assumed. On the other hand, the

potential for increased vulnerability to non-focal predators may also interfere with the

expression of defensive traits (Innes-Gold et al., 2019). This disadvantage may sometimes

outweigh the energetic costs of expressing a defensive phenotype. Carfagno et al. (2011)

found that tadpoles of *Acris* exhibit darker a tail spot, which is considered a defensive trait

against dragonfly nymphs, even in the absence of predators. However, the presence of fish

predators induces the loss of this spot. This suggests that the cost of expressing defensive

traits may be lower than typically expected, and the disadvantage of increased vulnerability to

mismatched predators may prevent these phenotypes from being constitutively expressed.

Similarly, in tadpoles of *D. leopardus*, the elevated risk of predation from non-odonate

predators, considered non-focal predators, due to expressing bright tail coloration, rather than

its energetic cost, could explain why they did not exhibit deep and brightly-colored tail in

response to these predators. These defense traits against dragonfly nymphs may not only be

ineffective in preventing predation by non-odonate predators but could also increase the

potential vulnerability to them. For example, the deepened tail may increase the vibrations

produced during swimming, thereby heightening predation pressure on tadpoles by tactile or

olfactory predators (i.e., water bugs, water scorpions, and newts). Additionally, for other

predators such as fish, tail coloration may make the tadpoles more vulnerable by increasing

their visibility. However, further experiments are warranted to assess and compare the

effectiveness of these defense traits toward each predator, including evaluating how tadpole

coloration influence their survival rate in the presence of various aquatic predators.

ACKNOWLEDGMENTS

We followed the ASIH "Guidelines for Live Amphibians and Reptiles in Field and Laboratory Research" (https://bit.ly/ASIH_Herps). The described methods were conducted in compliance with the Regulation on Animal Experimentation at Kyoto University. Because the aim of the experiments was to induce the anti-predator phenotypes of tadpoles, we needed to use live tadpoles and some of them were eaten by their predators in the experiments. However, the process was the same as a predation event occurring in nature, which did not cause any unnatural pain to the tadpoles. We thank the Experimental Farm, Graduate School of Agriculture, Kyoto University, for providing the research environment, and Mr. Kagata, Mr. Yasui, and Mr. Sakakibara for their assistance with observations. This work was supported by JST SPRING, Grant Number JPMJSP2110.

LITERATURE CITED

Agrawal, A. A., and R. Karban. 1999. Why induced defenses may be favored over constitutive strategies in plants, p. 45−61. *In*: Ecology and Evolution of Inducible Defenses. R. Tollrian and C. D. Harvell (eds.). Princeton University Press, Princeton, New Jersey.

Babbitt, K. J., and F. Jordan. 1996. Predation on *Bufo terrestris* tadpoles: effects of cover and predator identity. Copeia 1996:485−488.

Baker, E. A. 2021. Evaluating a Predator-Induced Phenotype in a Mixed Species Context. Unpubl. Master's Thesis, East Carolina University, Greenville, North Carolina.

Barnett, J. B., C. Michalis, N. E. Scott-Samuel, and I. C. Cuthill. 2018. Distance

dependent defensive coloration in the poison frog *Dendrobates tinctorius*,

Dendrobatidae. Proceedings of the National Academy of Sciences 115:6416–6421.

Barry, M. J., and S. Syal. 2013. Metabolic responses of tadpoles to chemical predation cues.

Hydrobiologia 700:267–276.

Benard, M. F. 2006. Survival trade-offs between two predator-induced phenotypes in Pacific

treefrogs (*Pseudacris regilla*). Ecology 87:340–346.

Bertrand, S., B. Faivre, and G. Sorci. 2006. Do carotenoid-based sexual traits signal the

availability of non-pigmentary antioxidants? Journal of Experimental Biology, 209:4414–

4419.

Bókony, V., Á. M. Móricz, Z. Tóth, Z. Gál, A. Kurali, Z. Mikó, K. Pásztor, M.

Szederkényi, Z. Tóth, J. Ujszegi, B. Üveges, D. Krüzselyi, R. J. Capon, H. Hoi, and

A. Hettyey. 2016. Variation in chemical defense among natural populations of common

toad, *Bufo bufo*, tadpoles: the role of environmental factors. Journal of Chemical

Ecology 42:329–338.

Borzée, A., K. R. Messenger, S. Chae, D. Andersen, J. Groffen, Y. I. Kim, J. An, S. N.

Othman, K. Ri, T. Y. Nam, Y. Bae, J. L. Ren, J. T. Li, M. F. Chuang . . . M. S. Min.

2020. Yellow sea mediated segregation between North East Asian *Dryophytes* species.

PLoS One 15:e0234299.

Brönmark, C., and L. A. Hansson. 2000. Chemical communication in aquatic systems: an

introduction. Oikos 88:103–109.

Bybee, S. M., K. K. Johnson, E. Gering, M. F. Whiting, and K. A. Crandall. 2012. All the

better to see you with: a review of odonate color vision with transcriptomic insight into

the odonate eye. Organisms Diversity & Evolution 12:241–250.

Carfagno, G. L. F., J. M. Carithers, L. J. Mycoff, and R. M. Lehtinen. 2011. How the

cricket frog lost its spot: the inducible defense hypothesis. Herpetologica 67:386–396.

Cocroft, R. B., and R. L. Rodríguez. 2005. The behavioral ecology of insect vibrational

communication. Bioscience 55:323–334.

Copeland, M. 1913. The olfactory reactions of the spotted newt *Diemyctylus viridescens*

(Rafinesque). Journal of Animal Behavior 3:260–273.

Corbet, P. S. 1999. Dragonflies: behaviour and ecology of Odonata. Cornell University Press,

Ithaca, New York.

Costello, D. M., and M. J. Michel. 2013. Predator-induced defenses in tadpoles confound

body stoichiometry predictions of the general stress paradigm. Ecology 94:2229–2236.

Dewitt, T. J., A. Sih, and J. A. Hucko. 1999. Trait compensation and cospecialization in a

freshwater snail: size, shape and antipredator behaviour. Animal Behaviour 58: 397–407.

Doherty, P. A., R. J. Wassersug, and J. M. Lee. 1998. Mechanical properties of the tadpole

tail fin. Journal of Experimental Biology 201:2691–2699.

Doi, T., H. Okuyama, and S. Murakami. 2005. Red tail fin patches in Japanese tree frog

tadpoles (*Hyla japonica*). Bulletin of the Herpetological Society of Japan 2005:4–7.

Duellman, W. E., A. B. Marion, and S. B. Hedges. 2016. Phylogenetics, classification, and

biogeography of the treefrogs (Amphibia: Anura: Arboranae). Zootaxa 4104:1–109.

Dufresnes, C., S. N. Litvinchuk, A. Borzée, Y. Jang, J. T. Li, I. Miura, N. Perrin, and M.

Stöck. 2016. Phylogeography reveals an ancient cryptic radiation in East-Asian tree frogs

(*Hyla japonica* group) and complex relationships between continental and island

lineages. BMC Evolutionary Biology 16:253.

Espanha, J., M. F. de Vasconcelos, and P. C. Eterovick. 2016. The role of tadpole

coloration against visually oriented predators. Behavioral Ecology and Sociobiology

70:255−267.

Futahashi, R., R. Kawahara-Miki, M. Kinoshita, K. Yoshitake, S. Yajima, K. Arikawa,

and T. Fukatsu. 2015. Extraordinary diversity of visual opsin genes in dragonflies.

Proceedings of the National Academy of Sciences 112:E1247−E1256.

Grether, G. F., J. Hudon, and J. A. Endler. 2001. Carotenoid scarcity, synthetic pteridine

pigments and the evolution of sexual coloration in guppies (*Poecilia reticulata*).

Proceedings of the Royal Society of London Series B 268:1245−1253.

Hawlena, D., R. Boochnik, Z. Abramsky, and A. Bouskila. 2006. Blue tail and striped

body: why do lizards change their infant costume when growing up? Behavioral Ecology

17:889−896.

Hettyey, A., Z. Tóth, K. E. Thonhauser, J. G. Frommen, D. J. Penn, and J. Van Buskirk.

2015. The relative importance of prey-borne and predator-borne chemical cues for

inducible antipredator responses in tadpoles. Oecologia 179:699−710.

Hettyey, A., B. Üveges, Á. M. Móricz, L. Drahos, R. J. Capon, J. Van Buskirk, Z. Tóth,

and V. Bókony. 2019. Predator-induced changes in the chemical defence of a vertebrate.

Journal of Animal Ecology 88:1925−1935.

Hettyey, A., K. Vincze, S. Zsarnóczai, H. Hoi, and A. Laurila. 2011. Costs and benefits of

defences induced by predators differing in dangerousness. Journal of Evolutionary

Biology 24:1007–1019.

Hettyey, A., S. Zsarnóczai, K. Vincze, H. Hoi, and A. Laurila. 2010. Interactions between

the information content of different chemical cues affect induced defences in tadpoles.

Oikos 119:1814–1822.

Hirai, N., T. Morioka, and M. Ishii. 2020. Species diversity of Odonata in Nakaikemi

Marsh, Fukui Prefecture, Japan. Japanese Journal of Environmental Entomology and

Zoology 31:1–12.

Hoff, K. V., and R. J. Wassersug. 2000. Tadpole locomotion: axial movement and tail

functions in a largely vertebraeless vertebrate. American Zoologist 40:62–76.

Hossie, T., K. Landolt, and D. L. Murray. 2017. Determinants and co-expression of anti-

predator responses in amphibian tadpoles: a meta-analysis. Oikos 126:173–184.

Innes-Gold, A. A., N. Y. Zuczek, and J. C. Touchon. 2019. Right phenotype, wrong place:

predator-induced plasticity is costly in a mismatched environment. Proceedings of the

Royal Society B 286:20192347.

Ishida, S., K. Ishida, K. Kojima, and M. Sugimura. 1988. Illustrated guide for

identification of the Japanese Odonata. Tokai University Press, Shinjuku, Tokyo.

Johnson, J. B., D. B. Burt, and T. J. Dewitt. 2008. Form, function, and fitness: pathways to

survival. Evolution 62:1243–1251.

Kasumovic, M. M. 2013. The multidimensional consequences of the juvenile environment:

towards an integrative view of the adult phenotype. Animal Behaviour 85:1049–1059.

Kehr, A. I., and V. I. Gómez. 2009. Intestinal, body and tail plasticity in *Rhinella schneideri*

(Bufonidae) tadpoles induced by a predator insect (Belostoma elegans). Advanced

Studies in Biology 1:85–94.

Kishida, O., and K. Nishimura. 2004. Bulgy tadpoles: inducible defense morph. Oecologia

140:414–421.

Kishida, O., and K. Nishimura. 2005. Multiple inducible defences against multiple

predators in the anuran tadpole, *Rana pirica*. Evolutionary Ecology Research 7:619–631.

Kishida, O., and K. Nishimura. 2007. Flexible inducible morphological defense in anuran

larvae. Japanese Journal of Ecology 57:40–47.

Kita, A., M. Nakahara, and M. Tokuda. 2020. Changes in Odonata abundance. between

2000 and 2015–2016 in Saga Plain, northern Kyushu, Japan. Journal of Insect

Conservation 24:575–583.

Kruger, A., and P. J. Morin. 2020. Predators induce morphological changes in tadpoles of

Hyla andersonii. Copeia 108: 316–325.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests

in Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

LaFiandra, E. M., and K. J. Babbitt. 2004. Predator induced phenotypic plasticity in the

pinewoods tree frog, *Hyla femoralis*: necessary cues and the cost of development.

Oecologia 138:350–359.

Lardner, B. 1998. Plasticity or fixed adaptive traits? Strategies for predation avoidance in

Rana arvalis tadpoles. Oecologia 117:119–126.

Lardner, B. 2000. Morphological and life history responses to predators in larvae of seven

anurans. Oikos 88:169–180.

Laurila, A., J. Kujasalo, and E. Ranta. 1998. Predator-induced changes in life history in

two anuran tadpoles: effects of predator diet. Oikos 83:307–317.

Laurila, A., S. Pakkasmaa, P. A. Crochet, and J. Merilä. 2002. Predator-induced plasticity

in early life history and morphology in two anuran amphibians. Oecologia 132:524–530.

Lawler, S. P. 1989. Behavioural responses to predators and predation risk in four species of

larval anurans. Animal Behaviour 38:1039–1047.

Lenth, R. 2025. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.10.7. https://CRAN.R-project.org/package=emmeans

Lindquist, S. B., and M. D. Bachmann. 1982. The role of visual and olfactory cues in the

prey catching behavior of the tiger salamander, *Ambystoma tigrinum*. Copeia 1982:81–

90.

Matsui, M., and N. Maeda. 2018. Encyclopedia of Japanese frogs. Bun-ichi Co., Ltd.,

Shinjuku, Tokyo.

McCollum, S. A., and J. D. Leimberger. 1997. Predator-induced morphological changes in

an amphibian: predation by dragonflies affects tadpole shape and color. Oecologia

109:615–621.

McCollum, S. A., and J. Van Buskirk. 1996. Costs and benefits of a predator-induced

polyphenism in the gray treefrog *Hyla chrysoscelis*. Evolution 50:583–593.

McCoy, M. W. 2007. Conspecific density determines the magnitude and character of

predator-induced phenotype. Oecologia 153:871–878.

McCoy, M. W., and B. M. Bolker. 2008. Trait-mediated interactions: influence of prey size,

density and experience. Journal of Animal Ecology 77:478–486.

McIntyre, P. B., S. Baldwin, and A. S. Flecker. 2004. Effects of behavioral and

morphological plasticity on risk of predation in a Neotropical tadpole. Oecologia

141:130–138.

Michel, M. J. 2012. Phenotypic plasticity in complex environments: effects of structural

complexity on predator-and competitor-induced phenotypes of tadpoles of the wood frog,

Rana sylvatica. Biological Journal of the Linnean Society 105:853–863.

Mogali, S., B. Shanbhag, and S. Saidapur. 2019. Experience of predacious cues and

accessibility to refuge minimize mortality of *Hylarana temporalis* tadpoles. Acta

Herpetologica 14:15–19.

Nakagawa, M., T. Motobayashi, Y. Arai, and T. Nishimura. 2004. Ecology of Eggs of

Gynacantha japaonica Bartenef in Paddy Fields. Transactions of the Japanese Society of

Irrigation, Drainage and Reclamation Engineering 229:71–77.

Nowińska, A., and J. Brożek. 2019. Antennal sensory structures in water bugs of Nepoidea

(Insecta: Hemiptera: Nepomorpha), their morphology and function. Zoomorphology

138:307–319.

Nunes, A. L., A. Richter-Boix, A. Laurila, and R. Rebelo. 2013. Do anuran larvae respond

behaviourally to chemical cues from an invasive crayfish predator? A community-wide

study. Oecologia 171:115–127.

Ohba, S., and F. Nakasuji. 2006. Dietary items of predacious aquatic bugs (Nepoidea:

Heteroptera) in Japanese wetlands. Limnology 7:41–43.

Ozaki, K., and K. Nishikawa. 2023. Feeding Habits of the Japanese Fire-belly Newt.

(Ampbihia: Urodela: Salamandridae) in Central Honshu, Japan. Current Herpetology

42:162–170.

Ozono, A., I. Kawashima, and R. Futahashi. 2021. Dragonflies of Japan. Bun-ichi Co. Ltd.,

Shinjuku, Tokyo.

Pritchard, G. 1965. Prey capture by dragonfly larvae (Odonata; Anisoptera). Canadian

Journal of Zoology 43:271–289.

Pritchard, G. 1966. On the morphology of the compound eyes of dragonflies (Odonata:

Anisoptera), with special reference to their role in prey capture. Proceedings of the Royal

Entomological Society of London. Series A, General Entomology 41:1–8.

R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Relyea, R. A. 2001. Morphological and behavioral plasticity of larval anurans in response to

different predators. Ecology 82:523–540.

Relyea, R. A., and J. R. Auld. 2004. Having the guts to compete: how intestinal plasticity

explains costs of inducible defences. Ecology Letters 7:869–875.

Relyea, R. A., and E. E. Werner. 2000. Morphological plasticity in four larval anurans

distributed along an environmental gradient. Copeia 2000:178–190.

Reuben, P. L., and J. C. Touchon. 2021. Nothing as it seems: behavioural plasticity appears

correlated with morphology and colour, but is not in a Neotropical tadpole. Proceedings

of the Royal Society B 288:20210246.

Richardson, J. L. 2006. Novel features of an inducible defense system in larval tree frogs

(*Hyla chrysoscelis*). Ecology 87:780–787.

Richter-Boix, A., G. A. Llorente, and A. Montori. 2007. A comparative study of predator-

induced phenotype in tadpoles across a pond permanency gradient. Hydrobiologia

583:43–56.

Schwarz, D., S. N. Gorb, A. Kovalev, N. Konow, and E. Heiss. 2020. Flexibility of intraoral

food processing in the salamandrid newt *Triturus carnifex*: effects of environment and

prey type. Journal of Experimental Biology 223:jeb232868.

Sherratt, T. N., and I. F. Harvey. 1989. Predation by larvae of *Pantala flavescens* (Odonata)

on tadpoles of *Phyllomedusa trinitatis* and *Physalaemus pustulosus*: the influence of

absolute and relative density of prey on predator choice. Oikos 56:170–176.

Shimada, T., M. Matsui, and K. Tanaka. 2025. Genetic and morphological variation

analyses of *Dryophytes japonicus* (Anura, Hylidae) with description of a new species

from northeastern Japan. Zootaxa 5590:61–84.

Smith, D. C., and J. Van Buskirk. 1995. Phenotypic design, plasticity, and ecological

performance in two tadpole species. The American Naturalist 145:211–233.

Steiner, U. K. 2007. Linking antipredator behaviour, ingestion, gut evacuation and costs of

predator-induced responses in tadpoles. Animal Behaviour 74:1473–1479.

Steiner, U. K., and J. Van Buskirk. 2009. Predator-induced changes in metabolism cannot

explain the growth/predation risk tradeoff. PLoS One 4:e6160.

Takahara, T., Y. Kohmatsu, A. Maruyama, H. Doi, H. Yamanaka, and R. Yamaoka.

2012a. Inducible defense behavior of an anuran tadpole: cue-detection range and cue

types used against predator. Behavioral Ecology 23:863–868.

Takahara, T., Y. Kohmatsu, A. Maruyama, and R. Yamaoka. 2008. Benefit of suites of

defensive behavior induced by predator chemical cues on anuran tadpoles, *Hyla japonica*.

Behavioral Ecology and Sociobiology 63:235–240.

Takahara, T., Y. Kohmatsu, A. Maruyama, and R. Yamaoka. 2012b. Predator-specific

inducible morphological changes in two Japanese anuran tadpoles. Herpetology Notes

5:43–47.

Tanahashi, M., and T. Fukatsu. 2018: *Natsumushi*: An image measuring software for

entomological studies. Entomological Science 21:347–360.

Teplitsky, C., S. Plenet, and P. Joly. 2003. Tadpoles' responses to risk of fish introduction.

Oecologia 134:270–277.

Teplitsky, C., S. Plenet, and P. Joly. 2004. Hierarchical responses of tadpoles to multiple

predators. Ecology 85:2888–2894.

Teplitsky, C., S. Plenet, J. P. Léna, N. Mermet, E. Malet, and P. Joly. 2005. Escape

behaviour and ultimate causes of specific induced defences in an anuran tadpole. Journal

of Evolutionary Biology 18:180–190.

Thibaudeau, G., and R. Altig. 2012. Coloration of anuran tadpoles (Amphibia):

development, dynamics, function, and hypotheses. International Scholarly Research

Notices 2012:1–12.

Touchon, J. C., and K. M. Warkentin. 2008. Fish and dragonfly nymph predators induce

opposite shifts in color and morphology of tadpoles. Oikos 117:634–640.

Touchon, J. C., and K. M. Warkentin. 2011. Thermally contingent plasticity: temperature

alters expression of predator-induced colour and morphology in a Neotropical treefrog

tadpole. Journal of Animal Ecology 80:79–88.

Urban, M. C., J. L. Richardson, N. A. Freidenfelds, D. L. Drake, J. F. Fischer, and P. P.

Saunders. 2017. Microgeographic adaptation of Wood Frog tadpoles to an apex predator.

Copeia 105:451–461.

Van Buskirk, J. 2002. A comparative test of the adaptive plasticity hypothesis: relationships

between habitat and phenotype in anuran larvae. The American Naturalist 160:87–102.

Van Buskirk, J., P. Anderwald, S. Lüpold, L. Reinhardt, and H. Schuler. 2003. The lure

effect, tadpole tail shape, and the target of dragonfly strikes. Journal of Herpetology

37:420–424.

Van Buskirk, J., J. Aschwanden, I. Buckelmüller, S. Reolon, and S. Rüttiman. 2004. Bold

tail coloration protects tadpoles from dragonfly strikes. Copeia 2004:599–602.

Van Buskirk, J., and S. A. McCollum. 1999. Plasticity and selection explain variation in

tadpole phenotype between ponds with different predator composition. Oikos 85:31–39.

Van Buskirk, J., and S. A. McCollum. 2000a. Functional mechanisms of an inducible

defence in tadpoles: morphology and behaviour influence mortality risk from predation.

Journal of Evolution Biology 13:336–347.

Van Buskirk, J., and S. A. McCollum. 2000b. Influence of tail shape on tadpole swimming

performance. Journal of Experimental Biology 203:2149–2158.

Wilson, R. S., P. G. Kraft, and R. Van Damme. 2005. Predator-specific changes in the

morphology and swimming performance of larval *Rana lessonae*. Functional Ecology

19:238–244.

Yager, D. D. 1999. Structure, development, and evolution of insect auditory systems.

Microscopy research and Technique 47:380–400.

Yuasa, Y. 2006. On the red tails of tadpoles of the frog, *Hyla japonica*. Hyogo Freshwater

Biology 58:87–90.

Yuasa, Y. 2007. Food of red-tailed larvae of the Japanese tree frog *(Hyla japonica)*. Hyogo

Freshwater Biology 59:143–144.

Fig. 1. Photographs of tadpoles of *Dryophytes leopardus* reared without predator (top) and

with *Anax nigrofasciatus* (bottom) and their corresponding Photoshop Lab color histograms.

The mean values of tail coloration are L = 61.2, a* = -2.5, and b* = 7.96 (without predator),

and L = 57.96, a* = 1.75, and b* = 35.51 (with *A. nigrofasciatus*), respectively.

Fig. 2. (A) Scatterplots of tail coloration parameters (a* and b* values) of tadpoles two weeks

(top) and three weeks (bottom) after the start of Experiment I, using different dragonfly

species. Tadpoles were reared in the presence of no predators (the control group; Ct1), *Anax*

nigrofasciatus (Ax), *Gynacantha japonica* (Gy), *Orthetrum albistylum speciosum* (Or), or

Pantala flavescens (Pf). (B) Box-and-whisker plots displaying the range of the tail color traits

(a* and b* values) two weeks (top) and three weeks (bottom) after the start of Experiment I.

The asterisks in the figure indicate that there were significant differences compared to the

control group, which were led by Dunnett's test (see Materials and Methods). *$P < 0.05$, ***P

<0.001.

Fig. 3. (A) Scatterplots of tail coloration parameters (a* and b* values) of tadpoles three weeks (bottom) after the start of Experiment II, using non-odonate species. Tadpoles were reared in the presence of no predators (the control group; Ct2), water bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt (*Cynops pyrrhogaster*; Fn). (B) Box-and-whisker plots displaying the range of the tail color traits (a* and b* values) three weeks after the start of Experiment II.

Fig. 4. Box-and-whisker plots showing the distribution of tadpole morphological traits, BL

(body length), TAL (tail length), and TD (tail depth) in Experiment I, using different

dragonfly species. Tadpoles were reared in the presence of no predators (the control group;

Ct1), *Anax nigrofasciatus* (Ax), *Gynacantha japonica* (Gy), *Orthetrum albistylum speciosum*

(Or), or *Pantala flavescens* (Pf). Boxplots represent the raw data distributions, while black

points and error bars indicate estimated marginal means and 95% confidence intervals derived

from linear mixed-effects models (see Materials and Methods). The asterisks in the figure

indicate that there were significant differences compared to the control group, which were led

by Dunnett's test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Fig. 5. Box-and-whisker plots showing the distribution of tadpole morphological traits, BL

(body length), TAL (tail length), and TD (tail depth) in Experiment II, using non-odonate

predators. Tadpoles were reared in the presence of no predators (the control group; Ct2), water

bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt

(*Cynops pyrrhogaster*; Fn). Boxplots represent the raw data distributions, while black points

and error bars indicate estimated marginal means and 95% confidence intervals derived from

linear mixed-effects models (see Materials and Methods). The asterisk in the figure indicates a

significant difference compared to the control group based on Dunnett's test. ***$P < 0.001$.

Table 1. Summary of model results for tail coloration (a* and b* values) measured at Week 2

and Week 3 after the start of Experiment I. Tadpoles were reared in the presence of no

predators (the control group), *Anax nigrofasciatus* (Ax), *Gynacantha japonica* (Gy),

Orthetrum albistylum speciosum (Or), or *Pantala flavescens* (Pf). Fixed effects include the

corresponding estimates (Est.), standard errors (SE), degrees of freedom (df), t-values (*t*), and

model P-values (P_m). Post hoc comparisons were performed using Dunnett's test to compare

each predator treatment to the control group (P_D). Variance (Var.) and standard deviation

(SD) for the random effect (Tank) and residuals are also shown. The control group served as

the reference level in all models.

a* value

Week 2

Effect	Est.	SE	df	*t*	P_m	P_D
(Intercept)	−2.158	0.052	32.153	−41.753	<0.001	
Ax	−0.462	0.074	33.245	−6.256	<0.001	<0.001
Gy	−0.157	0.076	37.975	−2.062	0.046	0.129
Or	−0.112	0.076	37.689	−1.462	0.152	0.392
Pf	−0.074	0.075	34.351	−0.991	0.329	0.689

Random Effect	Var.	SD
Tank	<0.001	0.009
Residual	0.378	0.615

Week 3

Effect	Est.	SE	df	*t*	P_m	P_D
(Intercept)	−1.998	0.063	25.632	−31.725	<0.001	
Ax	−0.247	0.093	26.952	−2.659	0.013	0.028
Gy	−0.141	0.091	27.399	−1.558	0.131	0.338
Or	−0.145	0.097	32.740	−1.495	0.145	0.373
Pf	−0.035	0.094	26.841	−0.369	0.715	0.966

Random Effect	Var.	SD
Tank	0.002	0.043
Residual	0.429	0.655

b* value

Week 2

Effect	Est.	SE	df	*t*	P_m	P_D
(Intercept)	7.048	0.296	31.896	23.786	<0.001	
Ax	2.568	0.422	32.707	6.083	<0.001	<0.001
Gy	0.620	0.431	35.535	1.439	0.159	0.405
Or	0.219	0.431	35.478	0.507	0.615	0.929
Pf	0.299	0.425	33.461	0.705	0.486	0.849

Random Effect	Var.	SD
Tank	0.269	0.519
Residual	7.679	2.771

Week 3

Effect	Est.	SE	df	*t*	P_m	P_D
(Intercept)	6.798	0.363	28.806	18.712	<0.001	
Ax	3.398	0.532	31.019	6.391	<0.001	<0.001
Gy	0.658	0.520	30.095	1.266	0.215	0.513
Or	1.355	0.543	34.275	2.494	0.018	0.045
Pf	1.083	0.538	29.914	2.012	0.053	0.143

Random Effect	Var.	SD
Tank	0.500	0.707
Residual	7.967	2.823

Table 2. Summary of model results for tail coloration (a* and b* values) measured at Week 3 after the start of Experiment II. Tadpoles were reared in the presence of no predators (the control group), water bug (*Appasus major*; Be), water scorpion (*Laccotrephes japonensis*; Ws), and fire belly newt (*Cynops pyrrhogaster*; Fn). Fixed effects include the corresponding estimates (Est.), standard errors (SE), degrees of freedom (df), t-values (*t*), and model P-values (P_m). Post hoc comparisons were performed using Dunnett's test to compare each predator treatment to the control group (P_D). Variance (Var.) and standard deviation (SD) for the random effect (Tank) and residuals are also shown. The control group served as the reference level in all models.

a* value
Week 3

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	−1.739	0.067	32.760	−25.970	<0.001	
Be	−0.031	0.094	31.019	−0.336	0.739	0.954
Ws	−0.097	0.092	29.311	−1.053	0.301	0.578
Fn	−0.086	0.092	29.619	−0.933	0.358	0.655

Random Effect	Var.	SD
Tank	0.015	0.123
Residual	0.258	0.508

b* value
Week 3

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	5.871	0.216	28.429	27.198	<0.001	
Be	0.117	0.301	26.657	0.387	0.702	0.938
Ws	0.452	0.295	25.105	1.532	0.138	0.294
Fn	0.605	0.296	25.383	2.043	0.052	0.108

Random Effect	Var.	SD
Tank	0.131	0.363
Residual	3.026	1.740

Table 3. Summary of model results for morphological traits (body length, tail length, and tail

depth) measured at Week 2 and Week 3 after the start of Experiment I. Tadpoles were reared

in the presence of no predators (the control group), *Anax nigrofasciatus* (Ax), *Gynacantha*

japonica (Gy), *Orthetrum albistylum speciosum* (Or), or *Pantala flavescens* (Pf). Fixed effects

include the corresponding estimates (Est.), standard errors (SE), degrees of freedom (df), t-

values (t), and model P-values (P_m). Post hoc comparisons were performed using Dunnett's

test to compare each predator treatment to the control group (P_D). Variance (Var.) and standard

deviation (SD) for the random effect (Tank) and residuals are also shown. The control group

served as the reference level in all models.

Body length

Week 2

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	9.365	0.176	29.472	53.098	<0.001	
Ax	0.221	0.251	30.283	0.879	0.386	0.757
Gy	0.128	0.257	33.237	0.499	0.621	0.932
Or	–0.098	0.257	33.152	–0.381	0.706	0.964
Pf	0.626	0.253	31.049	2.472	0.019	0.048

Random Effect	Var.	SD
Tank	0.077	0.277
Residual	3.047	1.746

Week 3

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	10.283	0.214	29.421	47.990	<0.001	
Ax	–0.116	0.316	31.356	–0.367	0.716	0.967
Gy	0.233	0.308	31.190	0.757	0.455	0.824
Or	0.627	0.326	36.584	1.921	0.063	0.174
Pf	0.673	0.319	30.756	2.111	0.043	0.116

Random Effect	Var.	SD
Tank	0.082	0.288
Residual	4.089	2.022

Tail length

Week 2

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	–0.410	0.352	441.321	–1.165	0.245	
Body length	2.230	0.035	639.859	64.157	<0.001	
Ax	–0.824	0.191	34.547	–4.315	<0.001	<0.001
Gy	–0.436	0.197	39.064	–2.214	0.033	0.091
Or	–0.376	0.197	38.784	–1.910	0.063	0.178
Pf	–0.860	0.194	36.388	–4.439	<0.001	<0.001

Random Effect	Var.	SD
Tank	0.006	0.079
Residual	2.416	1.554

Week 3

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	–2.121	0.409	316.445	–5.182	<0.001	
Body length	2.287	0.036	488.929	63.557	<0.001	
Ax	–0.307	0.257	29.473	–1.192	0.243	0.560
Gy	–0.810	0.251	29.271	–3.224	0.003	0.005
Or	–0.224	0.266	34.751	–0.842	0.405	0.778
Pf	–0.020	0.261	29.368	–0.078	0.938	0.999

Random Effect	Var.	SD
Tank	0.064	0.253
Residual	2.586	1.608

Tail depth

Week 2

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	–1.229	0.141	373.721	–8.733	<0.001	
Body length	0.827	0.014	642.972	61.130	<0.001	
Ax	0.266	0.088	31.452	3.029	0.005	0.009
Gy	0.061	0.090	34.327	0.683	0.499	0.860
Or	0.025	0.090	34.238	0.280	0.781	0.983
Pf	–0.114	0.089	32.718	–1.288	0.207	0.499

Random Effect	Var.	SD
Tank	0.010	0.100
Residual	0.358	0.598

Week 3

Effect	Est.	SE	df	t	P_m	P_D
(Intercept)	–1.545	0.154	328.542	–10.038	<0.001	
Body length	0.801	0.014	488.464	58.376	<0.001	
Ax	0.519	0.091	27.792	5.723	<0.001	<0.001
Gy	0.068	0.089	28.102	0.763	0.452	0.821
Or	0.200	0.095	34.000	2.119	0.042	0.113
Pf	0.253	0.092	28.144	2.751	0.010	0.022

Random Effect	Var.	SD
Tank	0.004	0.061
Residual	0.380	0.617