SMG6-dependent nonsense-mediated decay maintains dsRNA homeostasis and
suppresses immunogenic transcripts in a hepatocellular carcinoma model

Enes S. Arpa [1], Virginie Ricci [1], Ilja E. Shapiro [2], Marija Jokic [3], Yahya Mohammadzadeh [4],
Lisa Bertrand [1], Irene Gonzalez-Menendez [5,6], Leticia Quintanilla-Martinez [5,6], Alexandre P.
Bénéchet [7], Jan Rehwinkel [8], Tatiana V. Petrova [4], Sanjiv A. Luther [3], Michal Bassani-
Sternberg [2], David Gatfield [1,*]

[1] Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland.
[2] Ludwig Institute for Cancer Research, University of Lausanne, Lausanne, Switzerland; Department
of Oncology, Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland; Agora Cancer
Research Centre, 1011 Lausanne, Switzerland.
[3] Department of Immunobiology, University of Lausanne, Epalinges, Switzerland.
[4] Department of Oncology, Ludwig Institute for Cancer Research Lausanne and University of
Lausanne, Lausanne, Switzerland.
[5] Department of Pathology and Neuropathology, University Hospital Tübingen, University of Tübingen,
Tübingen, Germany.
[6] Cluster of Excellence iFIT (EXC 2180) "Image-Guided and Functionally Instructed Tumor Therapies",
Eberhard-Karls University of Tübingen, Tübingen, Germany.
[7] In Vivo Imaging Facility (IVIF), Department of Research and Training, Lausanne University Hospital
and University of Lausanne, Lausanne, Switzerland.
[8] Medical Research Council Translational Immune Discovery Unit, Medical Research Council
Weatherall Institute of Molecular Medicine, Radcliffe Department of Medicine, University of Oxford,
Oxford, UK.
[*] Correspondence: david.gatfield@unil.ch

Abstract

The nonsense-mediated mRNA decay (NMD) pathway degrades transcripts bearing
premature termination codons (PTCs), typically arising from mutations and RNA processing
errors, to prevent the accumulation of aberrant proteins. Its role in cancer is complex, with both
tumor-promoting and tumor-suppressive functions described. Here, we selectively inhibit NMD
in the liver by conditionally inactivating its endonuclease SMG6 in a genetic mouse model of
hepatocellular carcinoma. SMG6 inactivation completely abrogates tumor formation and elicits
both innate and adaptive immune responses. Mechanistically, SMG6 deficiency activates the
type I interferon response via the double-stranded RNA (dsRNA) sensor MDA5, uncovering a
physiological role for NMD in regulating cytoplasmic dsRNA levels. In parallel, stabilization and
translation of NMD-target transcripts generate non-canonical immunopeptides presented by
MHC-I, promoting potent CD8[+] T-cell responses. These findings establish SMG6-dependent
NMD as a core orchestrator of immune tolerance in cancer and reveal that its selective

inhibition in cancer cells unleashes immunogenic transcript expression with therapeutic potential.

Introduction

Eukaryotic cells employ nonsense-mediated mRNA decay (NMD) as a critical post-transcriptional quality control mechanism that degrades transcripts harboring premature termination codons (PTCs), thereby preventing the accumulation of truncated, potentially toxic proteins [1, 2]. PTCs typically arise from mutations, transcriptional errors, or splicing defects. In addition to its canonical surveillance function, NMD is increasingly recognized as a broad regulator of gene expression, shaping the transcriptome through selective degradation of physiological mRNAs [3]. NMD is executed by a multiprotein machinery centered on the helicase UPF1, and elicits mRNA cleavage through the endonuclease SMG6, followed by further exonucleolytic degradation [2].

In cancer, the role of NMD is multifaceted and context-dependent [4]. On the one hand, it may suppress tumorigenesis by limiting the expression of mutated, potentially oncogenic transcripts. On the other, it can promote immune evasion by eliminating mRNAs that encode neoantigens, particularly those arising from frameshift-inducing insertion/deletion (indel) mutations that are common in cancer genomes. By suppressing such immunogenic RNAs, NMD may dampen anti-tumor immune surveillance and reduce responsiveness to immunotherapies [5-9].

Previous studies investigating NMD in cancer have primarily relied on global depletion of core NMD components such as UPF1 or SMG proteins in cell lines or transplantable tumor models [10-14]. These approaches are limited in their ability to capture early tumorigenic events and the dynamics of immune surveillance. Moreover, these models often conflate NMD-specific effects with pleiotropic roles of UPF/SMG proteins in genome stability, telomere maintenance, and other essential processes [15].

To address these limitations, we developed a liver-specific, inducible mouse model that selectively disables the endonuclease activity of SMG6 without ablating the protein entirely [16]. We combined this model with a genetic background of hepatocellular carcinoma (HCC) driven by hepatic deletion of *Pten* and *Tsc1*, which closely mirrors human disease progression [17]. Using this physiologically relevant system, we demonstrate that selective inactivation of SMG6 triggers robust innate and adaptive immune responses and potently suppresses liver tumorigenesis. Mechanistically, for the innate immune response, we uncover that SMG6 loss induces a type I interferon (IFN) response via activation of the cytosolic double-stranded RNA (dsRNA) sensor MDA5. This reveals an unexpected physiological role for NMD in maintaining

cytoplasmic dsRNA homeostasis and preventing aberrant activation of antiviral immunity. In

validation of the long-standing neoantigen hypothesis, we find that the adaptive immune

response is driven by the stabilization and translation of NMD-targeted transcripts, which give

rise to non-canonical peptides presented on MHC-I and elicit infiltration by CD8[+] T-cells.

Together, our findings establish SMG6-dependent NMD as a central regulator of immune

tolerance in the liver and highlight its selective inhibition as a promising strategy to promote

immunogenicity in cancer.

Results

HCC suppression through mutant SMG6

We recently developed a conditional NMD loss-of-function mouse model that specifically

targets the endonucleolytic activity of SMG6 (**Extended Data Fig. 1a)** [16]. This is achieved

through Cre/loxP-mediated conversion of a wild-type protein-expressing *Smg6^flox* allele to

Smg6^mut, in which point mutations inactivate the catalytic residues of the PIN (PilT N-terminus)

nuclease domain. This approach overcomes the limitations of full NMD factor knockouts, which

often impair general cellular fitness due to both NMD-dependent and -independent functions

of these proteins [15, 18]. Liver-specific *Smg6^mut* mice (i.e., *Smg6^flox/flox*; *Alb^CreERT2* animals treated

with tamoxifen to activate CreERT2 in hepatocytes) display overall mild phenotypes [16],

underscoring the model's suitability for probing NMD function *in vivo*, including in disease-

relevant contexts*.

To investigate the consequences of SMG6 inactivation on tumorigenesis, we combined this

NMD allele with a genetic model of hepatocellular carcinoma (HCC). This model (**Fig. 1a**)

employs conditional knockout alleles of the tumor suppressors *Pten* and *Tsc1*, which encode

inhibitors of the mTOR pathway. Liver-specific knockout of these genes (hereafter referred to

as L-dKO) promotes lipogenesis and drives the progression from non-alcoholic fatty liver

disease (NAFLD) at early stages, to non-alcoholic steatohepatitis (NASH) and, ultimately, HCC

at late stages [17], closely recapitulating disease evolution in humans [19].

We first assessed the impact of *Smg6^mut* on HCC formation and progression at late stages, i.e.

beginning from week 12 post-tamoxifen (**Fig. 1a**). Weekly magnetic resonance imaging (MRI)

revealed liver tumors in L-dKO *Smg6^wt* mice (**Fig. 1b**), whose rapid progression necessitated

euthanasia by week 17 due to advanced-stage HCC (**Fig. 1c**). In contrast, L-dKO *Smg6^mut*

mice developed no tumors and none required sacrifice, indicating complete suppression of the

HCC phenotype (**Fig. 1c**).

Macroscopic and histological analyses confirmed HCC in L-dKO *Smg6^wt* (**Fig. 1d**, top). In L-

dKO *Smg6^mut* mice, persisting until the endpoint at week 21 post-tamoxifen, we observed

prominent duct proliferation and moderate perisinusoidal and periportal fibrosis (**Fig. 1d**,

bottom). These changes disrupted the hepatic parenchyma and resulted in a nodular tissue architecture. Notably, no histological evidence of HCC was detected, and L-dKO $Smg6^{mut}$ livers showed elevated CD8$^+$ T-cell infiltration compared to L-dKO $Smg6^{wt}$ (**Fig. 1e, f**), consistent with sustained adaptive immune activation. Serum markers of liver injury (ALT, AST, LDH; **Extended Data Fig. 1b-d**) were similarly elevated in L-dKO $Smg6^{mut}$ and L-dKO $Smg6^{wt}$ animals at the late stage, and both groups exhibited comparable hepatomegaly (**Extended Data Fig. 1e**). As essential controls, L-dKO $Smg6^{mut}$ livers retained expression of the recombined, mutant $Smg6$ mRNA (**Extended Data Fig. 1f**) and lacked PTEN and TSC1 proteins from hepatocytes (**Figures 1g-i**), with remaining protein localized to non-parenchymal cells such as Kupffer cells (**Figure 1j**). These findings ruled out trivial explanations for the absence of HCC – such as synthetic lethality of the $Smg6^{mut}$; $Pten^{ko}$; $Tsc1^{ko}$ genotype and regeneration by non-mutant tissue – and demonstrate that tumor suppression is specifically due to the interaction between mutant SMG6 and the tumor-prone L-dKO background. In summary, we concluded that inhibition of NMD via $Smg6$ mutation prevents cancer formation in this genetic model of HCC.

SMG6 inactivation promotes early immune infiltration

Given the striking phenotypic divergence at late stages – tumors in $Smg6$ wild-type mice but none in $Smg6$ mutants – we next investigated the early stages in our genetic model, at 3 weeks post-tamoxifen (**Fig. 1a**), with the aim of understanding the molecular and cellular changes occurring in NMD-deficient livers prior to tumor development. As expected, Western blot analysis at this early time point confirmed strong depletion of TSC1 and PTEN proteins in L-dKO livers, regardless of $Smg6$ genotype (**Fig. 2a, b**).

Macroscopic and histological analyses revealed comparable NASH features in both $Smg6^{wt}$ or $Smg6^{mut}$ L-dKO livers, including focal microsteatosis, hepatocyte ballooning, inflammatory infiltration, and necrosis (**Fig. 2c; Extended Data Fig. 2a**). However, inflammation was already markedly more pronounced in L-dKO $Smg6^{mut}$ livers at this early stage, as shown by increased staining for the neutrophil/monocyte marker myeloperoxidase (MPO; **Fig. 2d**, left panels) and the cytotoxic T-cell marker CD8 (**Fig. 2d**, right panels: **Fig. 2e**). Flow cytometry further confirmed elevated hepatic leukocyte counts in L-dKO $Smg6^{mut}$ livers (**Fig. 2f-h; Extended Data Fig. 2b, c**). Consistent with this heightened inflammatory state, serum levels of liver injury markers also peaked specifically in L-dKO $Smg6^{mut}$ animals at this early time point (**Extended Data Fig. 1b-d**).

Interestingly, increased immune cell infiltration was also observed in $Smg6^{mut}$ livers lacking the L-dKO alleles, indicating that NMD inhibition alone promotes a pro-inflammatory hepatic environment (**Fig. 2i, j**). Notably, when $Smg6$ mutation was combined with L-dKO alleles, infiltration was further enhanced – particularly for innate immune populations such as

neutrophils and Ly6C^{high} and Ly6C^{low} monocytes (**Fig. 2g-j**). In addition, L-dKO *Smg6^{mut}* livers

displayed elevated apoptotic markers (**Extended Data Fig. 2d-h**) and increased expression of

pro-fibrotic genes (**Extended Data Fig. 2i**), consistent with heightened cell death and liver

injury. Together, these findings demonstrate that SMG6 inactivation potentiates early liver

inflammation, immune cell infiltration, and apoptosis in a tumor-prone background – at a time

point well before overt HCC develops in this model.

Smg6 mutation induces CD8⁺ T-cell responses through non-canonical immunopeptide

presentation

To investigate the mechanism underlying the immune-altered state in *Smg6^{mut}* livers, we first

focused on adaptive immune activation – particularly the infiltration of CD8⁺ T-cells (**Fig. 2j**),

key effectors of anti-tumor immunity. We hypothesized that NMD inhibition enables

hepatocytes to present novel MHC-I-bound immunopeptides derived from stabilized NMD-

target transcripts.

To test this, we employed a multi-omics approach (**Fig. 3a**), combining transcriptomics (RNA-

seq, to uncover NMD-regulated transcripts), ribosome profiling (Ribo-seq, to detect translated

open reading frames), and immunopeptidomics (mass spectroscopy-based identification of

MHC-I-bound peptides). This strategy enables the identification of peptides uniquely presented

in *Smg6^{mut}* livers and to trace their origin to NMD-regulated transcripts.

Immunopeptidomics analyses identified over 2,000 MHC-I-bound peptides (**Fig. 3b**), most of

which were shared between *Smg6^{mut}* and *Smg6^{wt}* livers and mapped to canonical coding

sequences (**Fig. 3c; Supplementary Table 1**). However, 18 peptides were detected

exclusively in *Smg6^{mut}* livers across all replicates, compared to only 4 peptides unique to

Smg6^{wt} – a highly significant enrichment of *Smg6^{mut}*-specific peptides (P = 5.4e-05; Fisher's

exact test). Among these, four peptides originated from non-canonical coding regions: three

from exonic sequences and one from an intronic region (**Fig. 3c**, middle and right panels).

For example, the peptide SAPDGIKTPM mapped to the *D17H6S56E-5* gene, previously

annotated as a long non-coding RNA with sequence similarity to a mouse leukemia virus (MLV)

envelope gene [20]. RNA-seq analysis revealed elevated expression of this transcript in *Smg6^{mut}*

livers (**Fig. 3d**), especially for exon-mapping reads, whereas intronic read levels were only

modestly increased (**Fig. 3e**). Since intron-mapping reads can serve as a proxy for

transcriptional output [21], these data suggest that *D17H6S56E-5* upregulation is primarily post-

transcriptional, consistent with its identity as an NMD target. Ribo-seq further confirmed

translation of a 596-amino acid (aa) open reading frame (ORF) containing the SAPDGIKTPM

epitope near its N-terminus (**Fig. 3d**, right panel). Similarly, the VVITFSKM peptide originated

from the *Gm6545/A330040F15Rik* locus, a pseudogene copy of *Fam111a*, which showed

increased RNA expression and ribosome occupancy of a 554-aa ORF (**Extended Data Fig.**

3a). The remaining two peptides – VSTTNFTHA and AGPSNSEIHL – mapped to an out-of-
frame product from the *Hemojuvelin (Hjv)* locus and an intronic sequence of *Transferrin (Trf)*,
respectively (**Supplementary Table 1**). Together, these findings indicate that NMD inhibition
enables the expression and translation of non-canonical transcripts, which give rise to novel
MHC-I-presented peptides – potentially accounting for the observed CD8$^+$ T-cell activation.
To test their immunogenicity, we assessed whether the four *Smg6^{mut}*-specific non-canonical
peptides could activate CD8$^+$ T-cells, using leukocytes isolated from the spleen and liver of
Smg6^{wt} and *Smg6^{mut}* mice three weeks post-tamoxifen (**Fig. 3f**). First, we noted that the
NetMHCpan rank predictions indicated strong binding affinity for all four peptides (rank < 2%;
Fig. 3g). Next, we chemically synthesized the four non-canonical peptides and used them as
a pool to determine their immunogenic potential. To assess specificity, we compared with two
control pools in our assay: one with commonly detected peptides and one with *Smg6^{mut}*-
specific canonical peptides (**Fig. 3c, g**). Upon stimulation with the *Smg6^{mut}*-specific non-
canonical peptide pool, CD8$^+$ T-cells from both spleen and liver of *Smg6^{mut}* mice showed a
robust increase in IFNγ expression (**Fig. 3h; Extended Data Fig. 3b, c**). In contrast, leukocytes
from *Smg6^{wt}* mice showed no such response, and neither of the control peptide pools elicited
IFNγ$^+$CD8$^+$ T-cell activation in either genotype (**Fig. 3h**). Together, these results establish that
Smg6^{mut}-specific non-canonical peptides are processed and presented via MHC-I and can
elicit robust CD8$^+$ T-cell activation *in vivo*.

SMG6 inactivation triggers a cell-autonomous type I IFN response
We next investigated whether the observed innate immune infiltration might also stem from a
cell-intrinsic, SMG6-dependent mechanism. Supporting this hypothesis, RNA-seq analysis
revealed broad upregulation of transcripts indicative of an activated type I interferon response
(**Fig. 4a**) [22]. Quantitative real-time PCR (qPCR) confirmed strong *Smg6^{mut}*-dependent induction
of canonical IFN-stimulated genes (ISGs), including *Ifih1*, *Oas1a*, *Ifi44*, *Zbp1*, and *Ifit1* (**Fig.**
4b), consistent with robust activation of the type I IFN pathway.
Given that whole liver tissue contains both *Smg6^{mut}* hepatocytes and *Smg6^{wt}* immune cells,
ISG induction could result either from hepatocyte-leukocyte interactions or from a cell-
autonomous effect of SMG6 loss-of-function. To distinguish between these possibilities, we
used *Smg6$^{flox/flox}$* primary tail fibroblasts transduced with tamoxifen-activatable *CreERT2* (**Fig.**
4c) [16]. Tamoxifen treatment efficiently abrogated NMD, as evidenced by >10-fold increase in
accumulation of an NMD-sensitive *Hnrnpl* splice variant (**Fig. 4d**). Crucially, NMD inactivation
triggered massive ISG upregulation, with *Ifih1*, *Oas1a*, *Ifi44*, *Zbp1*, and *Ifit1* increasing by up
to several hundredfold (**Fig. 4e**), establishing that loss of SMG6 activity is sufficient to induce
a type I IFN response in a cell-autonomous manner. To assess IFN pathway activation at the
protein level, we measured CXCL10, an IFN-inducible chemokine and well-established

surrogate for type I IFN signaling [23]. CRE-treated *Smg6^flox* fibroblasts displayed markedly elevated CXCL10 levels compared to all relevant controls (**Fig. 4f**), further supporting autonomous IFN pathway activation under *Smg6^mut* conditions. Importantly, all genotypes responded comparably to poly I:C stimulation, reaching similarly high CXCL10 levels (**Fig. 4f**), indicating that the downstream IFN signaling machinery remains intact and inducible across conditions. Together, these findings demonstrate that SMG6 endonuclease inactivation drives a cell-intrinsic type I IFN response, independent of immune cell interactions, in both liver tissue and fibroblasts.

Increased abundance of intron-retaining transcripts in *Smg6^mut* livers

The type I IFN response is a key antiviral pathway triggered when cytoplasmic sensors detect atypical nucleic acids, such as dsRNA. Although typically associated with viral infection, endogenous sources of dsRNA – arising from dysregulated RNA processing – can similarly enhance anti-tumor immunity or induce apoptosis. One such mechanism involves splicing inhibition, which promotes leakage of intron-containing mRNAs into the cytoplasm. These transcripts can form dsRNA structures, often due to repetitive element sequences embedded within introns [24-26].

Transcripts with retained introns are well-established NMD substrates in mammals [27, 28], as also shown in our previous work using *Smg6^mut* fibroblasts [16]. We therefore asked whether a similar accumulation of intron-retaining transcripts occurs in *Smg6^mut* livers. Using hepatic RNA-seq data, we performed differential expression analysis focusing on annotated intron-retaining isoforms. This revealed a marked increase in over 500 such isoforms in *Smg6^mut* livers, compared to only 81 showing significantly reduced abundance (**Fig. 4g**). Inspection of representative gene loci confirmed that exonic read coverage – shared by canonical and intron-retaining isoforms – remained largely unchanged, while intronic read coverage was markedly elevated (**Fig. 4h, i**), supporting specific enrichment of intron-containing transcripts. Notably, we identified introns predicted to form dsRNA structures due to the presence of oppositely oriented repetitive elements. For example, the retained intron in *Phkg2-208* (**Fig. 4i**) harbors two SINE/B1 elements in sense and antisense orientation, resulting in extensive complementarity predicted to support intramolecular dsRNA formation (**Fig. 4j, k**). Additional transcript category analyses from the RNA-seq dataset, revealing a global increase in lncRNAs and NMD-targets, alongside a reduction in protein-coding isoforms, are shown in (**Extended Data Fig. 4a-c**).

Together, these findings demonstrate that SMG6 inactivation leads to widespread accumulation of intron-retaining transcripts in the liver, which may serve as endogenous sources of immunostimulatory dsRNA and contribute to type I IFN activation.

MDA5 is required for the SMG6-dependent type I IFN response

The widespread accumulation of intron-retaining transcripts in *Smg6^mut* livers and fibroblasts suggests a source of endogenous dsRNA capable of triggering innate immune signaling. To identify how this immunogenic RNA is sensed, we focused on the cytoplasmic dsRNA receptors MDA5 and RIG-I, both of which activate type I IFN responses via MAVS [29]. Since RIG-I preferentially detects short dsRNAs with 5′-triphosphate ends – a hallmark of viral RNAs [29] – we hypothesized that MDA5, which may recognize longer, structured dsRNA species [30], mediates the response to *Smg6* inactivation.

To test this, we generated *Smg6^flox/flox* fibroblast lines lacking *Ifih1*, the gene encoding MDA5 (**Fig. 5a**, **Extended Data Fig. 5a, b**), and induced *Smg6* mutation using recombinant, cell-permeable TAT-Cre protein (**Fig. 5b**). NMD inhibition remained intact in MDA5-deficient cells, as evidenced by increased accumulation of the NMD-sensitive *Hnrnpl* splice variant (**Fig. 5c**). Strikingly, however, ISG induction – robust in MDA5-proficient *Smg6^mut* cells – was nearly abolished in the absence of MDA5 (**Fig. 5d**), demonstrating that this sensor is essential for the SMG6-dependent IFN response. Consistent with the accumulation of immunogenic RNA, cytoplasmic dsRNA levels were significantly elevated in *Smg6^mut* fibroblasts, as detected by increased J2 antibody staining (**Fig. 5e, f**) [31].

These results identify MDA5 as the critical sensor linking SMG6 inactivation to type I IFN activation, via detection of endogenous dsRNA. Moreover, the specificity of this response – and its absence in Cre-treated MDA5-deficient cells – rules out confounding activation of alternative pathways such as cGAS-STING, previously associated with Cre-mediated DNA damage [32]. Together, our findings establish SMG6-dependent NMD as a key RNA quality control mechanism that prevents the accumulation of endogenous dsRNA and inappropriate activation of MDA5 and type I IFN signaling — thereby preserving innate immune quiescence in non-infected cells.

Discussion

Using a unique combination of tissue-specific NMD inactivation with a fully immunocompetent genetic cancer model for HCC, we provide *in vivo* evidence that loss of the endonuclease activity of SMG6 triggers both innate and adaptive immune mechanisms and potently acts to prevent tumor development. Specifically, our study validates the long-standing "neoantigen hypothesis", showing that aberrant transcripts normally degraded by NMD give rise to non-canonical peptides presented on MHC-I and elicit CD8+ T-cell responses. Unexpectedly, we also uncover a second, previously unrecognized regulatory function of NMD, which may arguably represent the more fundamental function of the pathway: SMG6 curtails aberrant innate immune activation by preventing the cytoplasmic accumulation of immunostimulatory dsRNAs. These dsRNAs, which can endogenously derive from intron-retaining transcripts

enriched in repetitive elements, are sensed by MDA5 and trigger a type I interferon response.

Together, these findings establish SMG6 as a molecular gatekeeper that maintains immune

quiescence by eliminating immunogenic RNA species before they engage dsRNA sensing

pathways or are translated into novel immunopeptide-containing proteins.

This dual role of SMG6 significantly reframes our understanding of NMD. Traditionally viewed

as a translation-coupled protein quality control system, NMD has been understood primarily in

terms of its ability to eliminate aberrant transcripts containing premature termination codons,

thereby preventing the accumulation of truncated, potentially toxic proteins. Our results extend

this view to include an RNA-centered function: controlling the abundance of specific

cytoplasmic RNAs (that is, dsRNAs) to prevent background activation of immune sensors. In

this light, SMG6 joins other RNA-modifying or -processing factors such as ADAR1 that act to

silence endogenous dsRNA signals and uphold self-tolerance [33].

Viewed from an evolutionary perspective, this RNA-centric function may represent a key

selective pressure driving the conservation of the pathway. While the suppression of toxic

proteins remains a valid functional rationale, the more immediate threat posed by the activation

of innate immune sensors – such as MDA5 and PKR – by endogenous RNAs may have

provided strong evolutionary pressures as well. These sensors are exquisitely tuned to

recognize features common to viral nucleic acids but can be inadvertently activated by

endogenous transcripts when RNA processing or degradation fails. In this context, SMG6-

mediated degradation acts as a buffer, maintaining the cytoplasmic RNA environment below

an immunostimulatory threshold. The parallels to mechanisms implicated in autoimmune and

inflammatory pathologies suggest that dysregulation of NMD could contribute to a wider

spectrum of diseases than previously appreciated [34].

Finally, our study positions SMG6 as a potential therapeutic target in oncology. In our

immunocompetent HCC model, liver-specific inactivation of SMG6 prevents tumor

development through robust immune activation. This supports the concept that NMD inhibition

can convert immunologically "cold" tumors into "hot" ones by exposing them to both innate and

adaptive immune surveillance. The challenge ahead lies in achieving selective modulation:

while transient, tissue-restricted inhibition of SMG6 could unleash anticancer immunity, it also

risks triggering inflammation, as evidenced by the development of steatohepatitis and fibrosis

in our model. Nonetheless, our results point to the exciting possibility that pharmacological

inhibition of SMG6's endonuclease activity may be harnessed to therapeutically manipulate

the immunogenicity of cancer, provided its immune-stimulatory effects can be balanced

against potential toxicity.

Methods

Animals

All animal experiments were conducted in accordance with the cantonal guidelines of the Canton of Vaud, Switzerland, license VD3739. Healthy adult male mice of age 8 to 10 weeks were used. Animals with $Alb^{CreERT2}$; $Smg6^{wt}$ and $Alb^{CreERT2}$; $Smg6^{flox}$ alleles were maintained on a C57BL/6J background. Mice with $Alb^{CreERT2}$; $Pten^{flox}$; $Smg6^{wt}$; $Tsc1^{flox}$ and $Alb^{CreERT2}$; $Pten^{flox}$; $Smg6^{flox}$; $Tsc1^{flox}$ alleles were maintained on a hybrid background (C57BL/6J, 129SvJ, and BALB/2c). The alleles $Alb^{CreERT2}$, $Pten^{flox}$, $Smg6^{flox}$, and $Tsc1^{flox}$ have been previously described [16, 35-37].

Induction of hepatocyte-specific *Pten* & *Tsc1* double knockout and *Smg6* mutation

8- to 10-week-old male L-dKO $Smg6^{wt}$ and $Smg6^{flox}$ mice, carrying the liver-specific *Albumin*-driven *CreERT2*, and their control littermates received four intraperitoneal injections of tamoxifen, dissolved in corn oil at a dosage of 75 mg/kg of body weight, in the consecutive days. The early-stage mice were euthanized 3 weeks after tamoxifen injections. In the late-stage experiment, L-dKO $Smg6^{wt}$ mice were euthanized based on the progression of HCC (latest at week 17). Animals in the other groups (Ctrl $Smg6^{wt}$, Ctrl $Smg6^{mut}$ and L-dKO $Smg6^{mut}$) were euthanized 21 weeks after tamoxifen injections.

Magnetic resonance imaging (MRI)

Mice were scanned weekly (starting from week 12), to monitor HCC evolution and progression, using a 3 Tesla small animal MRI scanner (Bruker BioSpin MRI, Ettlingen, Germany) with a 40-mm volume coil as a transmitter. Animals were anaesthetized using isoflurane (Attane) at 3-4% for 2 min. When fully sedated, animals were transferred into the MRI machine. The imaging procedure was performed under 1.5% isoflurane in oxygen. Data was acquired via Paravision 360 v2.0 software.

Histopathology and immunohistochemistry analyses

Livers were fixed in 4% buffered formalin for 24 h and embedded in paraffin. For histology 3-5 µm-thick sections were cut and stained with Hematoxylin and Eosin (H&E) and Masson's Trichrome (MTC). Immunohistochemistry was performed on an automated immunostainer (Ventana Medical Systems, Inc.) according to the company's protocols for open procedures with slight modifications. All slides were stained with the antibodies CD4 (SP35; Zytomed, Berlin, Germany), CD8 (C8/144B; Dako, Glostrup, Denmark), PTEN (Cell Signaling Technology, 9188), Cleaved-Caspase3 (Cell Signaling Technology, 9661), and MPO (anti-MPO Ab-1, Lab Vision UK, Ltd., Newmarket, Suffolk, Great Britain). Appropriate positive and negative controls were used to confirm the adequacy of the staining. The histologic samples

were analyzed by an experienced pathologist (L. Quintanilla-Martinez). All samples were scanned with the Ventana DP200 (Roche, Basel, Switzerland) and processed with the Image Viewer MFC Application. Final image preparation was performed with Adobe Photoshop 2024.

Intrahepatic leukocyte isolation and flow cytometry staining

Leukocytes in liver were isolated as described previously [38]. Mice were euthanized and subsequently perfused with 10 ml 1x PBS via the inferior vena cava. 1 g of liver tissue was excised from each mouse and placed into a tube containing 10 ml RPMI. The tissues were then crushed using 70 µm cell strainers and a plunger, ensuring all pieces were passed through the strainer. The resulting cell suspensions were centrifuged at 1600 x g for 5 min. The cell pellets were resuspended in a solution containing 2 mg collagenase, 25 µl DNase and 10 ml RPMI. The mixture was incubated at 37°C for 40 min, constantly stirring. After incubation, 25 ml RPMI was added to dilute the mixture. The samples were then centrifuged at 300 x g for 3 min to precipitate the connective tissue. The supernatants were transferred to new tubes and centrifuged at 1600 x g for 5 min. The cell pellets were then resuspended in a Percoll density gradient solution (36% Percoll [Cytiva], 4% 10x PBS, 60% RPMI) and centrifuged at 2000 x g for 20 min without brake. The pellets were resuspended with 2 ml of Ammonium-Chloride-Potassium (ACK) lysis buffer for 2 min to lyse remaining red blood cells. Following that, 8 ml RPMI was added to the tubes, and the samples were centrifuged at 1600 x g for 5 min. The purified leukocyte pellets were resuspended in 1 ml HBSS. The cells were blocked with anti-CD16/32 and then washed with flow cytometry buffer (1x PBS, 2% FCS, 2mM EDTA). Next, the cells were incubated with following primary antibodies for 25 min at 4°C in the dark: CD19-BV605 (BD biosciences, 563148), TCRb-PE (BD biosciences, 12-5961-81), CD4-biotin (homemade), CD8a (Biolegend, 100750), CD44-FITC (homemade), CD11b-FITC (Biolegend, 101206), CD45-PE-Cy7 (Biolegend, 10114), Ly6C-BV785 (Biolegend, 128041), Ly6G-APC-Cy7 (Biolegend, 127624), CD19-PE (Immunotools, 22190194), TCRb-PE (BD biosciences, 12-5961-81), MHCII-BV711 (Biolegend, 107643), CD11c-biotin (homemade), NK1.1 (BV605, 108753), Streptavidin (APC, 405207) and BV510-Zombie aqua (Biolegend, 423102). Following washes with flow cytometry buffer, the cells were incubated with secondary antibodies for 20 min at 4°C in the dark. The cells were fixed with BD fixation kit (554714).

Serum parameters

Following the euthanasia of the animals, blood was quickly collected from the decapitated body, and then transferred into a lithium-heparin (LiHep) added tube (BD microtainer) to obtain serum. After 30 min to 1 h incubation at room temperature (RT), samples were centrifuged (2,000 g x 10 min, RT) to separate serum. Following that, 100 µl of serum was loaded to a Cobas c111 analyzer to measure the levels of following enzymes: ALT, AST and LDH.

Smg6 cDNA genotyping

Total RNA was prepared as described previously [39]. For each RNA sample, the following reaction was assembled: 4.5 µg RNA, 1.5 µl random hexamer (100 µg/ml), 1.5 µl 10 mM dNTPs, RNase-free water to 18 µl. Samples were heated for 5 min at 65°C, and then immediately put on ice. A master mix containing the following was added to each tube: 6 µl 5x First-strand buffer, 3 µl 0.1 M DTT, 0.375 µl RNase inhibitor, 1.5 µl RNase-free water. After a short incubation (2 min, at RT), 1.125 µl Superscript II reverse transcriptase enzyme was added to each reaction. Samples were incubated 10 min at 25°C, 50 min at 42°C, 15 min at 70°C, respectively. *Smg6* genotyping polymerase chain reaction (PCR) was performed as described previously [16]. The primer sequences – encompassing both mutant sites – are as follows: forward: CCA GAT ACC AAC GGC TTC AT; reverse: TCT GGG TGG ATT GGT AGC TC. Reactions were run on a 0.8% agarose gel, at 100 V for 100 minutes. Bands corresponding to ~620 nt were excised, and then purified using Qiaquick Gel Extraction Kit (QIAGEN, 28704). DNAs were Sanger-sequenced by Microsynth.

Quantitative Reverse Transcriptase PCR

Total RNA was prepared, and cDNA was synthesized as mentioned above. 10-times diluted cDNA templates and a master mix containing SYBR green master mix (2x), forward and reverse primers (3.0 µM) were mixed, and then a 384-well plate was prepared using the Tecan Freedom EVO software. RT-PCR was performed using QuantStudio6 real-time systems. Relative mRNA expression levels were obtained by normalizing each CT value (mean of three technical replicates) to *Rpl13* and *Rpl27* expressions. The qPCR primer sequences can be found in Supplementary Table 2.

Transcriptomics analysis

Total RNA was prepared as mentioned above. RNA-seq libraries were prepared from individual mouse livers according to the standard protocols. Single-end sequencing (150 bases) was performed on NovaSeq6000 (Illumina) with a sequencing depth of 400 million cDNA-mapping reads in total. Sequencing reads were trimmed for the first two low quality (nucleotides) nt. Trimmed reads were then mapped using STAR [40] (v2.7.11b; --twopassMode Basic --seedSearchStartLmax 50 --outSAMmultNmax 1) in a sequential manner: mouse rRNA (Ensembl version 39.111), human rRNA (Ensembl version 38.111), mouse tRNA, and mouse genome. Read counts per genes were obtained using summarizeOverlaps() of the R (v4.4.0) library GenomicAlignments (mode = 'Union') and low read counts were excluded (≤5 counts in less than three samples). Differential gene expression analysis was performed using DEseq2 [41].

Ribosome profiling

The libraries for ribosome profiling were generated as [42] with slight changes. Frozen liver tissues (200 mg) from individual mice were homogenized with 5 strokes using a Teflon homogenizer in 3 volumes of polysome buffer (150 mM NaCl, 20 mM Tris-HCl [pH 7.4], 5 mM DTT, complete EDTA-free protease inhibitors [Roche] and 40 U/ml RNasin plus [Promega]) supplemented with 1% Triton X-100 and 0.5% Na deoxycholate. Liver lysates were incubated on ice for 10 min and cleared by centrifugation at 10,000 g, 4°C for 10 min. Supernatants were flash-frozen and stored in liquid nitrogen. For each sample, 15 OD_{260} of liver lysate was digested with RNase I (Ambion) for 45 min at RT under constant shaking. RNase I-digested RNA was separated on 15% urea-polyacrylamide gels to excise monosome (~30 nt) footprints. From the gel slices, RNA was extracted overnight at 4°C on a rotating wheel before precipitation in isopropanol for 1 h, at -20°C. RNA 3'-end repair was performed with 2 U/µl of T4 PNK (Lucigen) before 2 h of adapter ligation at 25°C using T4 RNA Ligase 1 (NEB) and T4 RNA Ligase 2 Deletion Mutant (Lucigen) and 1 µl of a 20 µM 5'-adenylated DNA adapter. Adapter removal was performed by treating individual libraries with 5'deadenylase (NEB) and RecJf exonuclease (NEB) for 1 h at 30°C and 1 h at 37°C. Samples were purified and pooled using Zymo Clean & Concentrator columns. Ribosomal RNA was depleted according to siTools Biotech rRNA depletion kit specifications with a custom-made riboPOOL. The clean-up step was performed using Zymo Clean & Concentrator columns. Further library preparation steps were performed as described. The amplification of libraries was performed using i5 and i7 [NexteraD502 or NexteraD503] primers. Libraries were sequenced on a NovaSeq6000 (Illumina).

Ribosome profiling data analysis

Sequencing reads were cleaned and trimmed. Briefly, adapter sequences were removed using cutadapt (v3.5) [43] with the following parameters --match-read-wildcards --overlap 8 --discard-untrimmed --minimum-length 30. Reads were filtered by quality using FASTX Toolkit (v0.0.14; -Q33 -q 30 -p 90) and by size using an in-house Python script (length 26-35 nt). Cleaned reads were then mapped using STAR (--twopassMode Basic –seedSearchStartLmax 28 –outSAMmultNmax 1) [40] in a sequential manner: mouse rRNA (Ensembl version 39.111), human rRNA (Ensembl version 38.111), mouse tRNA, and mouse genome. STAR was used for the mapping to splice junctions, as neoantigens can be produced from both exons and introns. The A-site position of mapped reads to the mouse genome was retrieved using the R (v4.4.0) package ribosomeProfilingQC (v1.16.0; estimatePsite() and getPsiteCoordinate(), and addition of 3 nt to recover the A-site position). The A-sites were assigned to the counting bins and the A-site count per bin was retrieved. Counting bins were defined using an in-house R script (main functions: makeTxDbFromGFF() from txdbmaker [v1.1.1]; exonicParts() from

GenomicFeatures [v1.57.0]; and reduce(), gaps(), and findOverlaps() from IRanges [v2.39.2])

and A-site counts were obtained using summarizeOverlaps() of GenomicAlignments (v1.41.0;

mode = 'Union') [44]. The low A-site counts (≤5 counts in less than three samples) were excluded

and the differential analysis was performed using DESeq2 (v.1.45.3). To identify putative

neoantigens, the median A-site counts were calculated for each condition and the bins with at

least one A-site count in L-dKO $Smg6^{mut}$ were retained. The nucleotide sequences of each bin

were extracted, translated in the three reading frames, and the resulting amino acid sequences

were then split by stop codon, and retained if the peptide sequence lengths were ≥ 8 amino

acids.

MHC-I antibody production

MHC-I monoclonal antibodies were purified from the supernatant of HB-51 and HIB 34.1.2

hybridoma cells (ATCC), respectively, with Protein A-Sepharose 4B beads (Invitrogen). HB-51

has specificity for H2-Db and H2-Kb, HIB 34.1.2 has specificity for H2-Dd, H2-Kd, and H2-Ld.

Each antibody type was cross-linked separately to Protein A-Sepharose 4B beads at

concentrations of 5 mg of antibodies per 1 ml of bead volume. To this end, the antibodies were

incubated with Protein A-Sepharose 4B beads for 1 hour, at RT. Antibodies and beads were

cross-linked with dimethyl pimelimidate dihydrochloride (Sigma-Aldrich) in 0.2 M sodium

borate buffer pH 9 (Sigma-Aldrich) to achieve a final concentration of 20 mM for 30 min. The

reaction was quenched via incubation with 0.2 M ethanolamine pH 8 (Sigma-Aldrich) for 2

hours. Cross-linked antibodies were stored at 4°C.

Purification of MHC-I-eluted peptides

200 mg of liver tissue per biological replicate was lysed in 2 ml phosphate buffered saline with

0.5% sodium deoxycholate (Sigma-Aldrich), 0.2 mM iodoacetamide (Sigma-Aldrich), 1 mM

EDTA, 1:200 Protease Inhibitor Cocktail (Sigma-Aldrich), 1 mM phenylmethylsulfonylfluoride

(Roche), and 1% octyl-beta-D glucopyranoside (Sigma-Aldrich) at 4°C for 1 hour, after

homogenization with an IKA T10 standard ULTRA TURRAX. Lysates were cleared by

centrifugation with a table-top centrifuge (Eppendorf Centrifuge) at 4°C at 20,000 g for 50 min.

MHC-I molecules purification was done with HB51 antibodies for C57BL/6 and a 1:1 mix of

HB51, HiB antibodies for C57BL/6; BALB/2c; 129_SvJae. Immunoprecipitations in 96-well

plates followed the protocol of [45]. Shortly, after plate conditioning, cross-linked beads and then

lysates were added to single wells, followed by washes at different sodium chloride

concentrations and an acidic elution step. Peptides were then desalted and purified on C18-

columns. Recovered peptides were dried using vacuum centrifugation (Thermo Fisher

Scientific) and stored at -80°C.

Mass-spectrometry acquisition

For Mass spectrometry (MS) analysis, MHC-I peptide samples were resuspended in 8 µl of 2% ACN and 0.1% formic acid (FA). 3 µl of peptide resuspension was injected into the LC-MS setup per measurement, counting two in DDA and one in DIA. The LC-MS setup consists of an Easy-nLC 1200 (Thermo Fisher Scientific) connected to a Q Exactive HF-X (Thermo Fisher Scientific). Chromatographic separation was done with a 450 mm homemade column of 75 µm inner diameter packed with ReproSil Pur C18-AQ 1.9µm resin (Dr. Maisch GmbH). For DDA measurements, the chromatographic separation was performed over 2 hours using a gradient of H_2O:FA, 99.9%:0.1% (solvent A) and CH_3CN:FA 80%:0.1% (solvent B). The gradient consisted of sequential, linear steps: 0 to 110 min (2-25% B), 110 to 114 min (25-35% B), 114 to 115 min (35-100% B), and 115 to 120 min (constant at 100% B) at a flow rate of 250 nl/min. The mass spectrometer had the following parameters set: full-scan MS spectra were acquired from m/z = 300-1,650 at a resolution of 60,000 (m/z = 200) with maximum injection times of 80 ms. Auto gain control (AGC) target value was tuned to 3×10^6 ions. We acquired MS/MS spectra at a resolution of 30,000 (m/z = 200), a top 20 method and isolation windows of 1.2 m/z, as well as a collision energy of 27 (HCD). All MS/MS scans accumulated ions to an AGC target value of 2×10^5 at a maximum injection time of 120 ms. Precursors of charge 4 and above were not selected for fragmentation. The dynamic exclusion was set to 20s. For DIA, the chromatographic separation was performed as in DDA measurements. The DIA acquisition cycles included one full-scan MS spectra, acquired from m/z = 300-1,650 (Resolution = 120,000, ion accumulation time = 60 ms), and 22 MS/MS scans. For all MS/MS scans the AGC target value was set to 3×10^6 ion, mass resolution at 30,000 and a stepped normalized collision energy (25.5, 27, 30) was applied. The maximum ion accumulation was set to auto and their first mass was fixed to 200 m/z. Consecutive MS/MS scans overlapped in their m/z windows by 1 m/z.

Immunopeptidomics data processing

All DDA files were used to generate a library with a database search using FragPipe (v22.0), including MSFragger (v4.1) [46]. The FASTA file was compiled as described in 'Ribosome profiling data analysis'. For search space generation, unspecific digestion with peptide lengths of 8 to 14 amino acids was used. Variable modifications included methionine oxidation, N-terminal acetylation, and cysteine carbamidomethylation. No fixed modifications were applied. Peptide, ion, and psm FDRs were set to 0.03 while applying a group-specific FDR as indicated in the FASTA header with PE = X, X being the respective protein group. No protein FDR was applied. All other settings were set to default. DIA files were searched with the library described above using DIA-NN (v1.8) with a precursor q-value cut-off of 0.01. Peptide binding affinities were predicted with MixMHC [47]. Peptides were considered as binders when they had a %-rank

below 2 for at least one HLA allele present in the sample haplotypes. Data analyses were done

with Julia and R, visualizations with R.

Protein extraction and Western Blot

Total proteins from mouse liver samples were isolated according to the NUN (NaCl, Urea,

Nonidet P-40) procedure, as [16]: Livers were homogenized in two tissue volumes of 10 mM

Hepes (pH 7.6), 100 mM KCl, 0.1 mM EDTA, 10% glycerol, 0.15 mM spermine, and 0.5 mM

spermidine for 15 s using a Teflon homogenizer. Four tissue volumes of 2x NUN buffer (2 M

urea, 2% NP-40, 0.6 M NaCl, 50 mM Hepes [pH 7.6], 2 mM dithiothreitol, and 0.1 mM

phenylmethylsulfonyl fluoride; supplemented with complete protease inhibitor tablets, Roche)

were added dropwise on a vortex with low speed. Lysates were incubated 1 h on ice and

cleared through centrifugation at 10,000 rpm, 4°C, for 20 min. Aliquots of the lysates (40 μg of

protein loaded per lane) were separated by SDS-polyacrylamide gel electrophoresis and

transferred to PVDF membrane using an iBlot 2 gel transfer device. The membrane was

blocked using 5% milk in Tris-buffered saline with 0,1% Tween 20 (TBST), 1 h at RT, followed

by an incubation overnight at 4°C with appropriate dilutions of primary antibodies, including

anti-Cleaved CASPASE3 (Cell Signaling Technology, 9661), anti-MDA5 (Cell Signaling

Technology, 5321), anti-PTEN (Cell Signaling Technology, 9188), anti-Hamartin/TSC1 (Cell

Signaling Technology, 6935), anti-GAPDH (Cell Signaling Technology, 2118), and anti-

VINCULIN (Abcam, ab129002). Following TBST washing (5 min, three times), the membrane

was incubated with the appropriate secondary antibody conjugated with horseradish

peroxidase for 60 min at RT, followed by TBST washing (15 min, three times).

Chemiluminescence signal was detected using Supersignal West Femto Maximum Sensitivity

Substrate (Thermofisher Scientific, 34095), as described by the manufacturer. The

quantification of signal was performed using ImageJ software.

Deletion of *Ifih1* in *Smg6^flox/flox*; *CreERT2* fibroblasts via CRISPR

First, lentiCRISPRv2-Blast-mU6 empty backbone (kind gift from J. Van Leeuwen, Addgene

plasmid #206806) [48] was digested with BsmBI (NEB, 0580) enzyme. The digestion products

were run on an agarose gel (0.8%), and the band corresponding to the open vector (~11 kb)

was excised, followed by DNA purification using QIAquick Gel Extraction Kit. Following

annealing reaction was assembled in a tube: 1 μl Forward primer (100 μM), 1 μl Reverse primer

(100 μM), 1 μl 10x T4 ligation buffer (NEB, B0202S), 0.5 μl T4 PNK (NEB, B0201S), and 6.5

μl H$_2$O. Primers were annealed with the following protocol: incubate 30 min at 37°C, and 5 min

at 95°C, respectively, then cool to 25°C at a rate of 5°C/min. Annealed primers were ligated

into the open vector using the following mixture: 50 ng open vector, 2 μl 10x T4 ligation buffer,

1 μl T4 DNA ligase (NEB, M0202S), 1 μl annealed primers (1:100 diluted), and nuclease-free

H$_2$O to 20 µl. Ligation mix was incubated at RT for 1 h. 10 µl ligation mix was added to 100 µl

Stbl3 competent cells, and incubated on ice for 30 min, followed by a heat-shock for 30 sec at

42°C. After a short incubation (5 min, on ice), 1 ml S.O.C. medium was added to the mixture,

and the tube was incubated at 37°C, for 1 h with shaking. 100 µl of the transformation mixture

was plated on an agar plate containing Ampicillin and incubated overnight at 37°C. Positive

colonies were selected and cultured overnight with shaking at 37°C, in LB medium containing

Ampicillin, then purified using a Midiprep kit (QIAGEN, 12943). Cells were transfected with the

plasmid using Lipofectamine 3000 (Thermo Fisher Scientific, L3000001), according to the

manufacturer's protocol, for 24 h, then were selected with Blasticidin (5 µg/ml) for 96 h. *Ifih1$^{ko/ko}$*

cells were obtained through single cell cloning. *Ifih1* sgRNA primer sequences: forward, CAC

CGC GTA GAC GAC ATA TTA CCA G; reverse, AAA CCT GGT AAT ATG TCG TCT ACG C.

Induction of Cre-mediated recombination in fibroblasts

Two different methods were used to induce Cre-dependent recombination:

1-Direct delivery of recombinant CRE protein (TAT-CRE, kind gift from I. Lopez-Mejia)*:*

Smg6$^{w/wt}$ and *Smg6$^{flox/flox}$* (-/+ *Ifih1$^{ko/ko}$*) fibroblasts were seeded in 6-well plates. The standard

culture medium (Thermo Fisher Scientific, 41965062) was replaced with Opti-MeM (Thermo

Fisher Scientific, 31985062). Cells were subsequently treated with 2.5 µM TAT-CRE or vehicle

for 8 h. After treatment, OptiMeM was replaced with the standard culture medium.

2-Recombination via 4-hydroxytamoxifen (4-OHT): *Smg6$^{flox/flox}$*; *CreERT2* cells were treated

with 2 µM 4-OHT or DMSO for 3 days (4-OHT containing media was refreshed every 24 h).

In both methods, cells were maintained in culture for at least one-week post-treatment to

ensure efficient recombination. Cells were subsequently collected for RNA extraction (as

explained above).

J2 immunofluorescence

Immunofluorescence of dsRNA was performed as [49]. First, *Smg6$^{flox/flox}$* cells were seeded in a

6-well plate, then treated with 2.5 µM TAT-CRE or vehicle as above. One week after the

treatment, cells were transferred to a new 6-well plate containing coverslips, at a confluency

of ca. 20%. Next day, the culture medium was removed, and cells were fixed using 4%

paraformaldehyde (PFA) for 15 min at RT, then washed three times (5 min each time), with 1x

PBS. Fixed cells were permeabilized with 500 µl PBSTX, followed by a 10 min incubation at

RT. PBSTX was removed, 500 µl of IFA blocking buffer was added into each well, and

incubated for 1 h, at RT with rocking. After blocking, cells were incubated with 500 µl primary

antibody solution (anti-dsRNA monoclonal antibody J2: Jena Bioscience, RNT-SCI-10010200)

in IFA blocking buffer (1:1000), overnight, at 4°C with rocking. Primary antibody solution was

removed, and cells were washed with 500 µl PBSTX (three times, for 5 min each). Cells were

incubated with the secondary antibody solution in PBSTX (1:300, Goat anti-mouse IgG Alexa
Fluor 488 for detection of J2, followed by washes with 500 µl PBSTX 3 times, for 5 min each).
To prepare the slides for imaging, coverslips were removed from 6-well plate, and cleaned with
70% ethanol, then placed face down onto the glasses, sealed with a nail polish. Images were
captured using a fluorescence microscope. The quantification of J2 intensity was performed
on ImageJ software.

Enzyme-linked immunosorbent assay (ELISA)
CXCL10 ELISA (R&D, DY466) was performed using R&D DuoSet™ ELISA development
system, according to manufacturer's protocol. 96-well ELISA plate (R&D, DY990) was coated
with 100 µl of capture antibody solution and incubated overnight at room temperature.
Following washes using wash buffer (R&D, WA126) three times, plate was blocked with 300
µl block buffer (R&D, DY995) for 1 hour at room temperature. Washes were repeated as above.
100 µl of diluted samples and standards were added to the plate and then incubated for 2
hours at room temperature. Washes were repeated as above. 100 µl of detection antibody
solution was added to the plate and then incubated for 2 hours at room temperature. Washes
were repeated as above. 100 µl of Streptavidin-HRP B solution was added to the plate and
then incubated for 20 minutes at room temperature. Washes were repeated as above. 100 µl
of substrate solution (R&D, DY999B) was added to the plate and then incubated for 20 minutes
at room temperature. The reaction was stopped via stopping solution (R&D, DY994). Protein
levels were detected using a microplate reader at 450 nm optical density.

Ex vivo peptide stimulation of leukocytes
Intrahepatic leukocytes isolated from mice as explained above. Spleens were mechanically
disrupted and filtered through a 70 µm cell strain to obtain single-cell suspensions. Both
intrahepatic leukocytes and spleens were washed with PBS. 2 million cells per mouse per
condition were seeded in round-bottom 96-well plates in T-cell medium (RPMI supplemented
with 25 µM 2-mercaptoethanol, 1 mM sodium pyruvate, MEM non-essential amino acids
(Gibco™, 11140050), 10% FBS, 1% Penicillin/Streptomycin, and L-glutamine. A pool of non-
canonical, canonical, and common peptides was added to the appropriate wells at a final
concentration of 2 µg/ml per peptide. CD28 agonist antibody (1 µg/ml; BioLegend, 102115)
and Brefeldin A were added to all wells. Cells were incubated at 37°C for 8 hours. After
incubation, cells were collected and stained with Zombie UV™ Fixable Viability Kit (1:500;
BioLegend, 423107) to stain dead cells. Then, extracellular and intracellular staining was
performed using BD Cytofix/Cytoperm™ Fixation/Permeabilization Kit (Cat# 554714)
according to the manufacturer's instructions. Data were acquired on a BD LSR Fortessa flow
cytometer and analyzed using FlowJo software. Antibodies used were: CD3-PE (BioLegend,

100206), CD45-Pacific Blue (BioLegend, 103126), CD8a-Brilliant Violet 510 (BioLegend,

100714), and IFNγ-APC (BioLegend, 505810).

RNA structure prediction

636   RNA structure prediction was performed via RNAfold web server using default parameters

637   (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi).

Repetitive element analysis

639   Repetitive elements within individual transcripts were detected using RepeatMasker [50] web

640   server (https://www.repeatmasker.org/cgi-bin/WEBRepeatMasker).

Acknowledgements

We thank the Lausanne Genomic Technologies Facility for library preparation and sequencing,

University of Lausanne Peptide & Tetramer Core Facility for the synthesis of synthetic

peptides, F. Lammers for help with genotyping, C. Roger for histology analysis, G.

Niederhäuser for serum measurements, Anne-Claire Godet at www.threoninedesign.fr for the

graphical illustration in Figure 5g, I. Lopez-Mejia for providing TAT-CRE, J. Van Leeuwen for

providing lentiCRISPRv2-Blast-mU6 plasmid, M. Hall and D. Mossmann for providing L-dKO

mice and for scientific inputs, O. Mühlemann and E. Karousis for important discussions. D.G.

acknowledges funding by Swiss National Science Foundation (SNSF) NCCR RNA & Disease

Phase III funding (grant number 205601), SNSF project grant (212423) and by Novartis

Foundation for Medical-Biological Research grant (21B066).

Author Contributions

653   Conceptualization: E.S.A. and D.G.; methodology: E.S.A., V.R., I.E.S., M.J., Y.M., L.B., I.G.-

654   M., and A.P.B.; investigation: E.S.A., V.R., I.E.S., M.J., Y.M., I.G.-M., and L.Q.-M.;

655   visualization: E.S.A., V.R., I.E.S., M.J., Y.M., and I.G.-M.; supervision: D.G., L.Q.-M., J.R.,

656   T.V.P., S.A.L., and M.B.-S.; writing: E.S.A. and D.G.

Declaration of Interests

658   The authors declare no competing interests.

Data Availability

The transcriptomics and ribo-seq data discussed in this publication have been deposited in

NCBI's Gene Expression Omnibus [51] and are accessible through GEO Series accession

numbers GSE278996 and GSE278997. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium via the PRIDE [52] partner repository with the

dataset identifier PXD057903.

Code Availability

All computational scripts are available on GitHub (https://github.com/NMD_Smg6/).

References

1.   Kurosaki, T., Popp, M.W. & Maquat, L.E. Quality and quantity control of gene expression by nonsense-mediated mRNA decay. *Nature reviews. Molecular cell biology* **20**, 406-420 (2019).

2.   Karousis, E.D. & Muhlemann, O. The broader sense of nonsense. *Trends Biochem Sci* **47**, 921-935 (2022).

3.   Boehm, V. *et al.* SMG5-SMG7 authorize nonsense-mediated mRNA decay by enabling SMG6 endonucleolytic activity. *Nature communications* **12**, 3965 (2021).

4.   Supek, F., Lehner, B. & Lindeboom, R.G. To NMD or Not To NMD: nonsense-mediated mRNA decay in cancer and other genetic diseases. *Trends in Genetics* **37**, 657-668 (2021).

5.   Lindeboom, R.G.H., Vermeulen, M., Lehner, B. & Supek, F. The impact of nonsense-mediated mRNA decay on genetic disease, gene editing and cancer immunotherapy. *Nat Genet* **51**, 1645-1651 (2019).

6.   Wu, C.C. *et al.* Immuno-genomic landscape of osteosarcoma. *Nat Commun* **11**, 1008 (2020).

7.   Litchfield, K. *et al.* Escape from nonsense-mediated decay associates with anti-tumor immunogenicity. *Nat Commun* **11**, 3800 (2020).

8.   Oka, M. *et al.* Aberrant splicing isoforms detected by full-length transcriptome sequencing as transcripts of potential neoantigens in non-small cell lung cancer. *Genome Biol* **22**, 9 (2021).

9.   Becker, J.P. *et al.* NMD inhibition by 5-azacytidine augments presentation of immunogenic frameshift-derived neoepitopes. *iScience* **24**, 102389 (2021).

10. Zhang, Y. *et al.* Immunotherapy for breast cancer using EpCAM aptamer tumor-targeted gene knockdown. *Proc Natl Acad Sci U S A* **118** (2021).

11. Pastor, F., Kolonias, D., Giangrande, P.H. & Gilboa, E. Induction of tumour immunity by targeted inhibition of nonsense-mediated mRNA decay. *Nature* **465**, 227-230 (2010).

12. Meraviglia-Crivelli, D. *et al.* IL-6/STAT3 signaling in tumor cells restricts the expression of frameshift-derived neoantigens by SMG1 induction. *Mol Cancer* **21**, 211 (2022).

13. Meraviglia-Crivelli, D. *et al.* A pan-tumor-siRNA aptamer chimera to block nonsense-mediated mRNA decay inflames and suppresses tumor progression. *Mol Ther Nucleic Acids* **29**, 413-425 (2022).

14. Nasif, S. *et al.* Inhibition of nonsense-mediated mRNA decay reduces the tumorigenicity of human fibrosarcoma cells. *NAR Cancer* **5**, zcad048 (2023).

15. Hwang, J. & Maquat, L.E. Nonsense-mediated mRNA decay (NMD) in animal embryogenesis: to die or not to die, that is the question. *Curr Opin Genet Dev* **21**, 422-430 (2011).

16. Katsioudi, G. *et al.* A conditional Smg6 mutant mouse model reveals circadian clock regulation through the nonsense-mediated mRNA decay pathway. *Science advances* **9**, eade2828 (2023).

17. Guri, Y. *et al.* mTORC2 Promotes Tumorigenesis via Lipid Synthesis. *Cancer Cell* **32**, 807-823 e812 (2017).

18. Li, T. *et al.* Smg6/Est1 licenses embryonic stem cell differentiation via nonsense-mediated mRNA decay. *The EMBO journal* **34**, 1630-1647 (2015).

19. Postic, C. & Girard, J. The role of the lipogenic pathway in the development of hepatic steatosis. *Diabetes Metab* **34**, 643-648 (2008).

20. Snoek, M., van Dinten, L. & van Vugt, H. A novel gene, G7e, resembling a viral
envelope gene, is located at the recombinational hot spot in the class III region of the
mouse MHC. *Genomics* **38**, 5-12 (1996).
21. Du, N.H., Arpat, A.B., De Matos, M. & Gatfield, D. MicroRNAs shape circadian
hepatic gene expression on a transcriptome-wide scale. *eLife* **3**, e02510 (2014).
22. Schoggins, J.W. *et al.* A diverse range of gene products are effectors of the type I
interferon antiviral response. *Nature* **472**, 481-485 (2011).
23. Sung, J.H. *et al.* Chemokine guidance of central memory T cells is critical for antiviral
recall responses in lymph nodes. *Cell* **150**, 1249-1263 (2012).
24. Bowling, E.A. *et al.* Spliceosome-targeted therapies trigger an antiviral immune
response in triple-negative breast cancer. *Cell* **184**, 384-403 e321 (2021).
25. Zheng, R. *et al.* hnRNPM protects against the dsRNA-mediated interferon response by
repressing LINE-associated cryptic splicing. *Mol Cell* **84**, 2087-2103 e2088 (2024).
26. Dias Junior, A.G., Sampaio, N.G. & Rehwinkel, J. A Balancing Act: MDA5 in Antiviral
Immunity and Autoinflammation. *Trends Microbiol* **27**, 75-85 (2019).
27. Lewis, B.P., Green, R.E. & Brenner, S.E. Evidence for the widespread coupling of
alternative splicing and nonsense-mediated mRNA decay in humans. *Proceedings of*
the National Academy of Sciences **100**, 189-192 (2003).
28. Zhou, H. & Deng, X.W. Intron Retention, an Orchestrated Program of Gene Expression
Regulation. *Bioessays* **47**, e202400248 (2025).
29. Rehwinkel, J. & Gack, M.U. RIG-I-like receptors: their regulation and roles in RNA
sensing. *Nat Rev Immunol* **20**, 537-551 (2020).
30. Pichlmair, A. *et al.* Activation of MDA5 requires higher-order RNA structures
generated during virus infection. *J Virol* **83**, 10761-10769 (2009).
31. Schonborn, J. *et al.* Monoclonal antibodies to double-stranded RNA as probes of RNA
structure in crude nucleic acid extracts. *Nucleic Acids Res* **19**, 2993-3000 (1991).
32. Pepin, G. *et al.* Cre-dependent DNA recombination activates a STING-dependent innate
immune response. *Nucleic Acids Res* **44**, 5356-5364 (2016).
33. Rehwinkel, J. & Mehdipour, P. ADAR1: from basic mechanisms to inhibitors. *Trends*
Cell Biol **35**, 59-73 (2025).
34. Rigby, R.E. & Rehwinkel, J. RNA degradation in antiviral immunity and autoimmunity.
Trends Immunol **36**, 179-188 (2015).
35. Schuler, M., Dierich, A., Chambon, P. & Metzger, D. Efficient temporally controlled
targeted somatic mutagenesis in hepatocytes of the mouse. *Genesis* **39**, 167-172 (2004).
36. Horie, Y. *et al.* Hepatocyte-specific Pten deficiency results in steatohepatitis and
hepatocellular carcinomas. *J Clin Invest* **113**, 1774-1783 (2004).
37. Kwiatkowski, D.J. *et al.* A mouse model of TSC1 reveals sex-dependent lethality from
liver hemangiomas, and up-regulation of p70S6 kinase activity in Tsc1 null cells. *Hum*
Mol Genet **11**, 525-534 (2002).
38. Benechet, A.P. *et al.* Dynamics and genomic landscape of CD8(+) T cells undergoing
hepatic priming. *Nature* **574**, 200-205 (2019).
39. Gatfield, D. *et al.* Integration of microRNA miR-122 in hepatic circadian gene
expression. *Genes Dev* **23**, 1313-1326 (2009).
40. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
41. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
42. Snieckute, G. *et al.* ROS-induced ribosome impairment underlies ZAKalpha-mediated
metabolic decline in obesity and aging. *Science* **382**, eadf3208 (2023).

43.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
764   reads. *EMBnet.journal* **17**, 10-12 (2011).

765   44.   Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS*
766   *Comput Biol* **9**, e1003118 (2013).

767   45.   Chong, C. *et al.* High-throughput and Sensitive Immunopeptidomics Platform Reveals
768   Profound Interferongamma-Mediated Remodeling of the Human Leukocyte Antigen
769   (HLA) Ligandome. *Mol Cell Proteomics* **17**, 533-548 (2018).

770   46.   Kong, A.T., Leprevost, F.V., Avtonomov, D.M., Mellacheruvu, D. & Nesvizhskii, A.I.
771   MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry-
772   based proteomics. *Nat Methods* **14**, 513-520 (2017).

773   47.   Gfeller, D. *et al.* Improved predictions of antigen presentation and TCR recognition
774   with MixMHCpred2.2 and PRIME2.0 reveal potent SARS-CoV-2 CD8(+) T-cell
775   epitopes. *Cell Syst* **14**, 72-83 e75 (2023).

776   48.   Unlu, B. *et al.* Global analysis of suppressor mutations that rescue human genetic
777   defects. *Genome Med* **15**, 78 (2023).

778   49.   de Faria, I.J.S., Imler, J.L. & Marques, J.T. Protocol for the analysis of double-stranded
779   RNAs in virus-infected insect cells using anti-dsRNA antibodies. *STAR Protoc* **4**,
780   102033 (2023).

781   50.   Smit, A., Hubley, R. & Green, P. RepeatMasker Open-3.0.
782   **http://www.repeatmasker.org** (1996-2004).

783   51.   Edgar, R., Domrachev, M. & Lash, A.E. Gene Expression Omnibus: NCBI gene
784   expression and hybridization array data repository. *Nucleic Acids Res* **30**, 207-210
785   (2002).

786   52.   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass
787   spectrometry-based proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).

Figures and Figure Legends

Figure 1. *Smg6* mutation suppresses cancer in genetic HCC model.

(a) Schematic of the experimental design. Inactivation of tumor suppressor genes *Tsc1* and *Pten* via Cre-mediated recombination in liver (L-dKO) leads to tumors/HCC from week 12 [17]. Cre-mediated recombination of the *Smg6^flox^* allele to *Smg6^mut^* introduces point mutations in catalytically important residues of the PIN domain [16]. Both L-dKO *Smg6^wt/wt^* and *Smg6^flox/flox^* mice received 4 intraperitoneal tamoxifen injections to activate hepatocyte-specific CreERT2 (driven from *Albumin* promoter). Early-stage experiments were conducted 3 weeks post-tamoxifen. Late-stage animals were scanned via MRI from week 12 to 21 (endpoint). In L-dKO *Smg6^wt/wt^*, livers show NAFLD and NASH as indicated.

(b) Representative longitudinal MRI scans for a tumor-bearing L-dKO *Smg6^wt^* mouse (top) and a tumor-free L-dKO *Smg6^mut^* animal. Tumors in L-dKO *Smg6^wt^* liver are indicated with yellow arrows (upper). Fibrotic plaques marked by yellow line in L-dKO *Smg6^mut^* liver (lower).

(c) Kaplan-Meier plot, from two different cohorts, displays survival percentage of L-dKO *Smg6^wt^* (n = 11) and *Smg6^mut^* (n = 13) mice throughout the experiment. *P*-value by a log-rank (Mantel-Cox) test. Asterisk indicates 3 animals that deceased for other reasons than HCC in one of the cohorts (stress in animal handling).

(d) Representative whole liver images, Masson's Trichrome (MTC) staining for fibrosis (at 20x magnification) and H&E images (at 2x magnification) from late-stage L-dKO *Smg6^wt^* and *Smg6^mut^* mice. Dashed lines surround tumors in L-dKO *Smg6^wt^*. Arrowheads indicate fibrotic areas in L-dKO *Smg6^mut^*.

(e) Representative immunohistochemistry (IHC) images of livers from late-stage mice, showing CD8-positive cells, at 20x magnification. Arrowheads mark CD8+ T-cells (brown).

(f) Quantification of CD8 IHC images. For each sample, the number of CD8-positive T-cells was counted in 15 randomly selected images, at 20x magnification (L-dKO *Smg6^wt^*, n = 5 mice; L-dKO *Smg6^mut^*, n = 6 mice). The *P*-value was calculated with two-tailed unpaired *t*-test.

(g) Western blot analysis of total liver proteins from late-stage mice for PTEN and TSC1. VINCULIN and GAPDH served as loading controls (Control, n = 2 mice; L-dKO *Smg6^wt^*, n = 6 mice; L-dKO *Smg6^mut^*, n = 3 mice).

(h) Quantification of PTEN and VINCULIN Western blots from (g)

(i) Quantification of TSC1 and GAPDH Western blots from (g).

(j) Representative IHC images of livers from late-stage mice stained with anti-PTEN antibodies. Light brown staining shows PTEN expression in hepatocytes in wild-type

control (blue arrowhead). Dark brown staining indicates PTEN expression in non-hepatocyte cells in both wild-type control and L-dKO (black arrowhead). Images were captured at 20x (large images) and 80x (small images) magnifications.

For (f), (h), and (i) data are plotted as means, and the error bars indicate SEM. The *P*-values were calculated with two-tailed unpaired *t*-test.

Figure 2. SMG6-dependent early inflammation.

(a) Western blot analysis on total liver proteins, for PTEN and TSC1, at early-stage post-tamoxifen. GAPDH serves as loading control (Control, n = 2 mice; L-dKO $Smg6^{wt}$, n = 6 mice; L-dKO $Smg6^{mut}$, n = 5 mice).

(b) Quantification of Western blots from (a).

(c) Representative images of whole-livers from early-stage $Smg6^{wt}$ and $Smg6^{mut}$ mice, without and with L-dKO background, as well as Hematoxylin and Eosin (H&E)-stained sections (at 10x magnification) from early-stage L-dKO $Smg6^{wt}$ and $Smg6^{mut}$ mice.

(d) Representative IHC images of livers from early-stage L-dKO $Smg6^{wt}$ and $Smg6^{mut}$ mice, using neutrophil marker Myeloperoxidase (MPO) and T-cell marker CD8, at 20x magnification. Arrowheads mark MPO-positive or CD8-positive cells (brown).

(e) Quantification of CD8-stained IHC images. For each sample, the number of CD8+ T-cells was counted in 15 randomly selected images, at 20x magnification (mean values from 4 mice per group).

(f) Schematic of the experimental design for intrahepatic leukocyte isolation and flow cytometry-based counting. $Smg6^{wt/wt}$ and $Smg6^{flox/flox}$ mice (with or without L-dKO) received 5 intraperitoneal tamoxifen injections. Livers were dissected 3 weeks later, and leukocytes were isolated for flow cytometric quantification. FACS, fluorescence-associated cell sorting.

(g) Overall leukocyte numbers per gram liver tissue (mean values from 7 mice per group), as assessed by flow cytometry. Cells are identified according to indicated color code. DC, dendritic cells; NK-cells, natural killer cells.

(h) Relative leukocyte counts for individual cell types as assessed by flow cytometry. Graphs show normalized numbers (set to 1 in L-dKO $Smg6^{wt/wt}$) of indicated leukocyte type (n = 7 mice for each group).

(i) Overall leukocyte numbers per gram liver tissue (mean values from 5 mice per group), as assessed by flow cytometry. Cells are identified according to indicated color code.

(j) Relative leukocyte counts for individual cell types as assessed by flow cytometry. Graphs show normalized numbers (set to 1 in $Smg6^{wt/wt}$) of indicated leukocyte type (n = 5 mice for each group. Data are plotted as means, and the error bars indicate SEM.

For (b), (e), (h), and (j) data are plotted as means, and the error bars indicate SEM. The *P*-values were calculated with two-tailed unpaired *t*-test.

Figure 3. Multiomics-based identifcation and validation of SMG6-dependent immunopeptides

(a) Schematic of the multiomics approach used to identify SMG6-dependent immunopeptide candidates by combining RNA-seq, Ribosome profiling (Ribo-seq) and immunopeptidomics data.

(b) MHC-I binder peptide count and distribution across *Smg6^{wt/wt}* and *Smg6^{mut/mut}* liver replicates (n = 3 mice per genotype). Enrichment of unique peptides in *Smg6^{mut/mut}* samples (18 in all replicates, orange circle) as compared to *Smg6^{wt/wt}* (4 in all replicates, blue circle) is highly significant (*P* = 0.000053; Fisher's exact test). The heatmap scale is colored according to peptide count, for better visual representation.

(c) Scatter plots of peptide intensity values, showing peptide categories canonical (protein-coding), non-canonical (exonic) and non-canonical (intronic), in *Smg6^{wt/wt}* (x-axis) vs, *Smg6^{mut/mut}* (y-axis). Data are plotted as medians from the three replicates. Unique peptides for one of the genotypes that are found in all three replicates are colored orange (*Smg6^{mut/mut}* -specific peptides) and blue (*Smg6^{wt/wt}*-specific peptides), with specific peptides that are used in the downstream pools highlighted and annotated with their amino acid sequence.

(d) RNA-seq (top panels) and Ribo-seq (bottom panels) read pileup plots show the strongly increased transcript and footprint abundance of *D17H6S56E-5* in *Smg6^{mut/mut}*. Left panel shows the larger genomic region, including the CDS (highlighted in turquoise) from which the non-canonical polypeptide is translated, while the right panel zooms into the specific region where the immunopeptide is translated (highlighted in light magenta). Plots were generated using merged sequencing data from the three biological replicates (individual livers) per group. Please note gene orientation from right to left.

(e) Normalized exonic and intronic RNA-seq read counts of *D17H6S56E-5*. Each dot represents the merged sequencing data from three biological replicates/livers.

(f) Schematic of the immunogenicity quantification experiment. *Smg6^{wt/wt}* and *Smg6^{mut/mut}* mice were sacrificed 3 weeks post-tamoxifen. Spleens and livers were extracted from the animals and leukocytes were isolated. The cells were then incubated with peptide pools indicated in (g), for 6 hours. The percentages of IFNγ⁺CD8⁺ T-cells were used as a readout for the immunogenicity of each peptide pool. Common peptides were used as negative control (*Smg6^{wt/wt}*, n = 3 mice, *Smg6^{mut/mut}*, n = 4 mice).

(g) Table showing peptide sequences derived from the indicated genes, transcript types, rank percentages from NetMHCpan prediction and their intensity in each of three

replicates of *Smg6^{wt/wt}* and *Smg6^{mut/mut}* livers. Each group of four peptides was used as a pool in the experiment in (h).

(h) Overall IFNγ⁺CD8⁺ T-cell percentage per tissue (left panel, spleen; right panel, liver), as assessed by flow cytometry, upon treatment with no peptide pool (-), canonical common peptide pool (com), canonical *Smg6^{mut}*-specific peptide pool (can), and non-canonical *Smg6^{mut}*-specific peptide pool (non-can). Bars indicate means, and the error bars SEM; individual datapoints are measurements from independent animals (*Smg6^{wt/wt}*, n = 3 mice, *Smg6^{mut/mut}*, n = 4 mice). The *P*-values were calculated with two-tailed unpaired *t*-test.

Figure 4. *Smg6* mutation elicits type I IFN response and increased intron abundance

(a) Heatmap showing ISG expression in control and mutant livers (n = 3 livers per group). RNA-seq data was plotted after Z-score transformation of normalized expression levels across the 6 samples.

(b) Relative hepatic mRNA levels of indicated ISGs (n = 3 mice for each group), detected via qPCR. *Smg6*$^{wt/wt}$ (control) samples were set to 1.

(c) Schematic of the experiment to analyze SMG6-dependent ISG expression. To induce *Smg6* mutation in *Smg6*$^{flox/flox}$; *CreERT2* fibroblasts, cells were treated with 2 µM 4-hydroxytamoxifen (4-OHT) during days 1 to 4, and cells were collected 6 days later (day 10) for RNA isolation and qPCR. Control treatment was DMSO.

(d) Relative mRNA expression of the endogenous, diagnostic NMD-target, an *Hnrnpl* splice form, in samples collected as described in (c). Three biological replicates were used per group.

(e) Relative mRNA expression of indicated ISGs in same samples as in (d). Three biological replicates were used per group.

(f) Analysis of CXCL10. At day 1, *Smg6*$^{wt/wt}$ and *Smg6*$^{flox/flox}$ cells were treated with TAT-CRE (2.5 µM, 8 hours) or vehicle. At day 7, indicated groups were treated with poly I:C to serve as positive control (1 µM, 8 hours). At day 10, supernatants were collected, and CXCL10 levels were detected via ELISA. Data are plotted as medians (n = 3 biological replicates for each group). The *P*-values were calculated with two-tailed unpaired *t*-test.

(g) Differential expression analysis at the transcript level, specifically on retained intron containing mRNAs, comparing *Smg6*$^{wt/wt}$ and *Smg6*$^{mut/mut}$ livers (n = 3 mice per genotype). Significantly more or less abundant transcripts are indicated in olive, showing strong skew to upregulation. Two transcripts are individually labeled, as they are further shown in panels (h) and (i)

(h) RNA-seq pileup plots show the upregulated transcript isoform (containing intron) for *H2-Q10*. Top panels show the larger genomic region, including the retained intron, while bottom panels zoom in the retained intron area. Plots were generated using merged sequencing data from three biological replicates per group. Please note gene orientation from left to right.

(i) RNA-seq pileup plots as in (h) show the increased abundance of the intron-containing transcript isoform for *Phkg2.* Genomic area where SINE/B1 elements are located in intronic region is highlighted. Plots were generated using merged sequencing data from three biological replicates per group. Please note gene orientation from left to right.

(j) RNA structure prediction of *Phkg2-208* by RNAfold web server, showing extensive predicted dsRNA region.

(k) Predicted double-stranded structure from SINE/B1 elements located in the intronic region of *Phkg2-208* transcript.

For (b), (d) and (e) data are plotted as means, and the error bars indicate SEM. The *P*-values were calculated with two-tailed unpaired *t*-test.

Figure 5. *Smg6* mutation drives type I IFN response through dsRNA sensor MDA5

(a) Western blot validation of the loss of MDA5 expression in *Ifih1^ko/ko^* fibroblasts (in background of *Smg6^flox/flox^* fibroblasts). GAPDH serves as loading control.

(b) Schematic of the experiment. Cells with indicated genotypes were treated with either TAT-CRE (2.5 µM) or vehicle for 8 hours, and then harvested at day 10 for RNA extraction and qPCR analyses.

(c) Relative mRNA expression of the endogenous, diagnostic NMD-target, an *Hnrnpl* splice form, in samples collected as described in (b). Three biological replicates were used per group.

(d) Relative mRNA expression of indicated ISGs. Arrows indicate the fold-change in the expression of indicated genes between cells nontreated vs. treated with TAT-CRE. Three biological replicates were used per group.

(e) Representative immunofluorescence images using dsRNA-specific antibody J2 (red) on *Smg6^flox/flox^* fibroblasts that were not treated (top) or treated (bottom) with cell-permeable TAT-CRE to induce *Smg6* mutation. DAPI (blue) serves to visualize nuclei.

(f) Quantification of J2 signal intensity (n = 238 cells for TAT-CRE-negative, n = 185 cells for TAT-CRE-positive). Median and third quartiles of each group are marked by dotted lines. The *P*-value was calculated with two-tailed unpaired *t*-test.

(g) Model for how the suppression of NMD activity via endonuclease-inactive SMG6 activates both innate and adaptive branches of the immune system. For innate responses, aberrant mRNAs (shown here: intron-retained transcripts known to contain repetitive elements) can form double-stranded configurations, causing activation of cytoplasmic dsRNA sensor MDA5, which upregulates a type I IFN response, resulting in innate immune cell infiltration. With the accumulation of aberrant or otherwise NMD-suppressed mRNAs in the cytoplasm, their translation leads to the production of unusual/non-canonical peptides which are then subsequently processed for MHC-I presentation on the cell surface, which attracts CD8+ T-cells to infiltrate the tissue, leading to adaptive immune cell recognition. Please note that for simplicity, prior steps required for CD8+ T-cell activation (i.e., antigen presentation by dendritic cells) have been omitted from the schematic.

Extended Data Figure 1. Regression of liver damage marker levels in *Smg6* mutant mice from early- to late-stage L-dKO.

(a) Schematic of the genetic NMD-deficient mouse allele based on *Smg6*. *Smg6^flox^* expresses wild-type SMG6 protein encoded by the blue exons; after Cre-mediated recombination to *Smg6^mut^*, point-mutated exons 18 and 19 (orange) lead to the expression of mutant SMG6 (D1352A, 1391A). The right panel shows the structure of the human protein (PDB accession 2HWW) with the corresponding mutated aspartic acid residues (in orange) that are within the catalytic triade of the PIN nuclease domain. Please note that human Asp1392 corresponds to mouse Asp1391, and human Asp1353 to mouse Asp1352. Figure adapted from [16].

(b-d) Comparison of the levels of indicated liver damage enzymes, detected in serum, between early- and late-stage mice. Arrows display the fold-change alteration in the

level of liver damage markers from early- to late-stage, in the same group (n = 4 mice for control, n = 5 mice for early-stage groups; L-dKO $Smg6^{wt}$ late, n = 10 mice; L-dKO $Smg6^{mut}$ late, n = 9 mice).

(e) Comparison of the ratio of liver-over-body weight between early- and late-stage mice, within the same group (n = 5 mice for early-stage groups; L-dKO $Smg6^{wt}$ late, n = 10 mice; L-dKO $Smg6^{mut}$ late, n = 9 mice).

(f) Sanger sequencing chromatograms on cDNA, from representative late-stage livers, show the conversion of aspartic acid (Asp) encoding wild-type codon (GAT) to alanine (Ala) encoding mutant codon (GCC), at the position in exon 18 that codes for one of the Asp residues of the catalytic center of SMG6 that was targeted in the genetic model, through tamoxifen-induced CreERT2 activation in liver.

Extended Data Figure 2. Cellular changes following *Smg6* mutation in liver.

(a) Representative H&E images from early-stage L-dKO livers, at 20x magnification. Black arrowheads indicate hepatocyte ballooning in both livers, while blue arrowheads mark apoptotic hepatocytes in L-dKO *Smg6^{mut}* tissue.

(b) Flow cytometry gating strategy of intrahepatic myeloid cells.

(c) Flow cytometry gating strategy of intrahepatic lymphocytes.

(d) Representative IHC images of livers from early-stage mice, stained with antibodies directed against cleaved Caspase-3, at 20x magnification. Arrowheads show cleaved Caspase-3+ cells (brown).

(e) Quantification of cleaved Caspase-3 IHC images as that shown in (d). For each sample, the number of cleaved Caspase-3+ cells was counted in 10 randomly selected images, at 20x magnification (mean values from 7 mice per group).

(f) Western blot analysis of total proteins, for cleaved Caspase-3. VINCULIN served as loading control (L-dKO *Smg6^{wt}*, n = 6 mice; L-dKO *Smg6^{mut}*, n = 5 mice).

(g) Quantification of Western blot from (f).

(h) Relative hepatic mRNA levels of *Bax*, from early-stage mice (n = 5 mice per group).

(i) Relative hepatic mRNA levels of indicated pro-fibrotic genes, from early-stage mice (n = 4 mice per group).

For (e), (g), (h), and (i) data are plotted as means, and the error bars indicate SEM. The *P*-values were calculated with two-tailed unpaired *t*-test.

Extended Data Figure 3. Increased non-canonical antigen expression in *Smg6* mutant livers.

(a) RNA-seq (top panels) and Ribo-seq (bottom panels) read coverage plots show upregulated transcript and ribosome footprint abundance for *Gm6545* in *Smg6^{mut/mut}* compared to *Smg6^{wt/wt}*. Left panel shows the whole genomic region, including the CDS (highlighted in turquoise) where the non-canonical polypeptide is translated, while right panel zooms into the specific region where the immunopeptide is translated (highlighted in light magenta). Read coverage plots were generated by merging sequencing data from three biological replicates (individual livers) per group.

(b, c) Representative flow cytometry gating strategies for one liver and one spleen from each group for IFNγ$^+$CD8$^+$ T-cells, treated with non-canonical peptide pool.

Extended Data Figure 4. RNA-seq analysis from *Smg6^{mut}* vs. *Smg6^{wt}* livers for different transcript biotypes.

(a-c) Differential expression analysis of liver transcriptomics data for protein coding transcripts (a), lncRNAs (b), and annotated NMD isoform transcripts (c). Significantly changed transcripts are highlighted in olive.

a

b

Extended Data Figure 5. Deletion of *Ifih1* in *Smg6^flox/flox* fibroblasts via CRISPR.

    (a) Western blot confirmation of the loss of MDA5 activity in different fibroblast single-cell clones. GAPDH serves as loading control.

    (b) Quantification of Western blot from (a).