Action mechanism of a novel agrichemical quinofumelin against

Fusarium graminearum

Qian Xiu [a,1], Xiaoru Yin [a,1], Yuanyuan Chen [b,1], Ziyang Zhang [a], Yushuai Mao [a], Tianshi Wang [b], Jie Zhang [a], Mingguo Zhou [a,2], Yabing Duan [a,2]

[a] College of Plant Protection, State Key Laboratory of Agricultural and Forestry Biosecurity, Nanjing Agricultural University, Nanjing, 210095, China.

[b] College of Science, Nanjing Agricultural University, Nanjing, 210095, China.

[1] Qian Xiu, Xiaoru Yin and Yuanyuan Chen contributed equally to this work.

[2] Corresponding authors: E-mail address: mgzhou@njau.edu.cn (Mingguo Zhou); dyb@njau.edu.cn (Yabing Duan)

Abstract

Modern fungicides have made significant contributions to crop disease management, but the development of resistant fungal strains has caused their failure in disease control. Therefore, developing fungicides with novel action mechanisms is the most effective measure to manage resistance. Quinofumelin, a novel quinoline fungicide, exhibits exceptional antifungal activity against phytopathogens. However, there is currently no available information on its mechanism of action. Here, we used transcriptome and metabolome analysis to observe a co-enrichment pattern of differentially expressed genes (DEGs) and differentially accumulated metabolites (DAMs) within pyrimidine biosynthesis pathway (PBP), identifying down-regulation of dihydroorotate dehydrogenase (DHODH). Exogenous uridine monophosphate (UMP), uridine or uracil (metabolites in PBP) successfully restored quinofumelin-induced inhibition of mycelial growth in *Fusarium graminearum* and *Fusarium asiaticum*. Additionally, the deletion of *FgDHODHII* was determined to be lethal; however, mycelial growth of ΔFgDHODHII mutants could be restored by adding UMP, uridine or uracil. These findings indicate that the deficiencies in *FgDHODHII* are functionally equivalent to complete inhibition of its activity by quinofumelin. Finally, molecular docking, surface plasmon resonance (SPR) and microscale thermophoresis (MST) results strongly support the precise interaction between quinofumelin and FgDHODHII. Collectively, these findings provide compelling evidence for the involvement of *de novo* uracil biosynthesis as mechanism of action for quinofumelin while identifying FgDHODHII as its specific target.

Key words: Action mechanism; Fungicide target; Quinofumelin; *Fusarium graminearum*

Introduction

Agrochemical fungicides have a history spanning over 3000 years in the control of plant diseases, and have undergone a development process from traditional protective fungicides with multiple action sites to modern selective fungicides with single action site (Lamberth, 2022). The introduction of the first fully systemic seed treatment agent (carboxin, a succinate dehydrogenase inhibitor) in 1966 and the launch of the first broad-spectrum foliar systemic fungicide (benomyl, a tubulin polymerization inhibitor) in 1968 represent significant milestones in the advancement of modern selective fungicides with single action site (Lamberth et al., 2021; Morton et al., 2008). Owing to their exceptional selectivity, potent activity, specific target site, and low toxicity, modern selective fungicides have gained widespread acceptance and witnessed an increasing market share year after year. However, like every coin has two sides, these types of fungicides encounter substantial challenges related to serious resistance issues when deployed in agricultural fields (Lucas et al., 2015). Numerous reports have been published on plant pathogens developing

resistance to modern selective fungicides, which often result in failures or reductions in the
effectiveness of chemical control against plant diseases and lead to decreased crop yields and
quality (https://www.frac.info) (Duan et al., 2019; King et al., 2021; Mao et al., 2024). Therefore,
it is crucial to expedite the development of new-generation selective fungicides with innovative
modes of action. Based on a single fungicide target, numerous modern selective fungicides can be
developed (Umetsu et al., 2020). For instance, 24 commercialized available fungicides have been
created so far based on the respiratory chain complex II as the target. Similarly, another 20
commercialized fungicides have been developed targeting the respiratory chain complex III.
Additionally, there are already 37 commercialized fungicides that specifically target the fungicide
target C14-demethylase in ergosterol biosynthesis (https://www.frac.info). However, identifying
specific targets that efficiently control plant pathogens while ensuring safety for non-target
organisms remains a significant challenge. The rate of discovery for new targets lags far behind
current demands.
According to the latest information released by the Fungicide Resistance Action Committee
(FRAC), over 30 fungicide selective targets have been identified and acknowledged in the field of
fungicide biological research spanning more than 130 years. Currently, these targets have been
identified based on the successful development of fungicides and are all related proteins that are
essential for pathogen growth or pathogenicity. Extensive identification and functional analysis of
essential genes for growth or pathogenicity in plant pathogens have been achieved with rapid
advancements in molecular biology techniques; however, there is no precedent for successfully
developing commercialized fungicides targeting genes involved in growth or pathogenicity. The
creation of highly active modern selective fungicides currently relies on the structural
characteristics of existing targets and is accomplished through technologies such as AI drug
discovery and design (AIDD), and computer aided drug design (CADD) (Yang et al., 2019). The
biological functions of drug targets are highly conserved, although variations may occur at critical
amino acid sites during genetic evolution. Therefore, it is crucial to explore drug targets and
analyze their structural characteristics in order to develop novel fungicides with enhanced activity
and selectivity. Recently, the Fungicide Biology group from Nanjing Agricultural University
proposed a new concept for targeting pesticides based on the principles of targeted drugs in the
medicine field (Zhong et al., 2021). The objective is to create specialized targeted pesticides that
exhibit higher activity, greater selectivity, stronger specialization, lower toxicity, and
environmental friendliness by leveraging the structural characteristics of pesticide targets. This
approach aims to provide essential technical support for scientifically controlling crop diseases.
Quinofumelin (CAS 861647-84-9) is a novel quinoline fungicide developed by Mitsui Chemicals
Co., Ltd. Japan (Ito et al., 2023). Due to their significant bioactivity, quinoline compounds and
their derivatives have found extensive applications in medicine and pesticides (Tao et al., 2021;
Wheeler et al., 2003). In our previous study, quinofumelin not only exhibited excellent antifungal
activity against the mycelial growth and spore germination of *F. graminearum*, but also inhibited
the biosynthesis of deoxynivalenol (DON) (Xiu et al., 2021). In addition, quinofumelin exhibits no
cross-resistance to currently extensively used fungicides, indicating its unique action target against
phytopathogenic fungi (Tao et al., 2021). In previous studies, quinofumelin was identified as a
specific inhibitor of dihydroorotate dehydrogenase class II (DHODHII) in *Pyricularia oryzae*,
suppressing fungal growth by blocking the conversion of dihydroorotate to orotate (Higashimura
et al., 2022). Gene disruption of *DHODHII* (*PoPYR4*) further confirmed its critical role in fungal
pathogenicity and validated it as the action target site of quinofumelin. However, previous

investigations did not report any detailed methodologies for target identification. Additionally,
there was a lack of information regarding the precise interaction between quinofumelin and
DHODHII. In this study, we used transcriptomic and metabolomic analyses to investigate the co-
enrichment patterns of differentially expressed genes (DEGs) and differentially accumulated
metabolites (DAMs). This approach enabled the identification of potential metabolic pathways
were identified. Ultimately, the pyrimidine biosynthesis pathway was identified as the target.
Therefore, this study unveils that the mechanism of action of quinofumelin involves *de novo*
synthesis of uracil, thereby identifing DHODH as its specific target. As a selective fungicidal
target, DHODH offers a valuable resource for the design and development of novel fungicides.

Results

Transcriptomic analysis of *F. graminearum* as affected by quinofumelin

RNA-seq experiments have emerged as the standard approach for quantifying and comparing gene
expression levels across a wide range of species and conditions, providing critical insights into
cellular phenomena (Pertea et al., 2016). To investigate the impact of quinofumelin on gene
expression levels in *F. graminearum*, we performed an RNA-Seq experiment. For quality
assessment of the RNA-Seq data, we conducted a comprehensive quality control analysis. In the
principal component analysis, PC1 and PC2 accounted for 45.59% and 16.98% of the total
variance, respectively. The distinct separation observed between the groups indicates significant
differences in gene expression between the controls and experimental samples, which could be
used for subsequent investigations (Supplementary Information, Fig. S1). The clustering heat map
effectively categorized the controls and experimental samples into distinct clusters based on their
gene expression profiles, demonstrating robust biological replication with minimal intra-group
variability and ensuring the reliability of the data. Moreover, notable inter-group dissimilarities
highlighted variations among different treatment samples (Fig. 1c).

The differential analysis unveiled a total of 234 significantly differentially expressed genes (DEGs)
between the control and experimental samples. Among these findings, there were identified as up-
regulated for a count of 97 genes while down-regulation was observed in case of another set
comprising around 137 genes (Fig. 1a). A graphical representation was employed to depict the
statistical distribution pattern exhibited by these DEGs (Fig. 1b). To gain deeper insights into gene
functionalities, Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG)
enrichment analyses were performed on these DEGs. The results showed that they are notably
enriched in seventeen distinct GO terms spanning across two major categories; eleven molecular
functions along with six processes (Fig. 1d). Furthermore, the enrichment analysis highlighted
significant associations with various functional attributes including transition metal ion binding,
oxidoreductase activity, modified amino acid binding, amide binding, monooxygenase activity,
and heme binding. Moreover, the enriched biological processes primarily involved transcriptional
regulation driven by nucleic-enabled mechanisms as well as monocarboxylic acid metabolism and
RNA biosynthesis. Additionally, the KEGG enrichment analysis identified specific pathways
associated with thiamine metabolism, tryptophan metabolism, nitrogen metabolism, amino acid
sugar and nucleotide sugar metabolism, pantothenic acid and CoA biosynthesis, as well as
nucleotide sugar production compound synthesis (Fig. 1e). In the GO and KEGG enrichment
analyses, the detailed counts of down- and up-regulated DEGs for each metabolic pathway are
provided in Supplementary Table S1 and S2.

Metabolomic analysis of *F. graminearum* as affected by quinofumelin

According to the findings from the KEGG enrichment analysis of DEGs derived from

transcriptomic data, these DEGs are associated with metabolism pathways as above described. To
specifically investigate the metabolic pathways involved action mechanism of quinofumelin, we
performed further metabolomic experiments. In metabolomics principal component analysis, the
contribution rate of PC1 was determined to be 76.6% in positive ion mode and 51.2% in negative
ion mode, respectively. Moreover, the quality control (QC) samples exhibited highly concentrated
results, indicating excellent instrument stability and remarkable data repeatability. Furthermore, a
distinct separation between groups was observed, particularly prominent in the positive ion mode,
indicating significant variations in metabolites among different groups that warrant further
investigation (Supplementary Information, Fig. S2a and S2b). The findings from Pearson
correlation analysis were consistent with those obtained from principal component analysis
(Supplementary Information, Fig. S2c and S2d). Strong correlations were identified within
samples belonging to the same group, while weak correlations were observed between different
groups.
Metabolomics analysis revealed a total of 713 differential metabolites, with 14 up-regulated and
699 down-regulated in the positive ion mode, as well as 230 differential metabolites, including
103 up-regulated and 127 down-regulated in the negative ion mode (Supplementary Information,
Fig. 2a and Fig. S3a). Based on the classification of the identified 943 different metabolites, it was
observed that amino acids and their derivatives accounted for the highest proportion (17%),
followed by organic acids and their derivatives (11%), benzene and substituted derivatives (11%),
aldehydes, ketones, esters (11%), and heterocyclic compounds (10%) (Fig. 2c). The volcano plot
visually represents the distribution of statistical differences in metabolite abundance between the
two groups (Supplementary Information, Fig. 2b and Fig. S3b).
KEGG pathways enriched by differential metabolites primarily include arachidonic acid
metabolism, pyrimidine metabolism, alanine, aspartate and glutamate metabolism, purine
metabolism, citrate cycle, arginine and proline metabolism, fructose and mannose metabolism,
nitrogen metabolism, and metabolic pathways (Fig. 2d). The classification of enrichment
pathways for differential metabolites revealed that the majority of these metabolites were
associated with nucleotide metabolism (25%), amino acid metabolism (25%), carbohydrate
metabolism (25%), and energy metabolism (12.5%) (Supplementary Information, Fig. S3c).
Conjoint analysis of transcriptome and metabolome integration
Through transcriptome and metabolome data analysis, we observed a co-enrichment of
differentially expressed genes and differentially accumulated metabolites in the pyrimidine
metabolism pathway. In the expression calorigrams of metabolites and genes depicting the *de novo*
biosynthesis pathway of pyrimidine, we identified down-regulation of dihydroorotate
dehydrogenase (DHODH) in the experimental groups. Through in vitro RT-qPCR verification, it
was found that after treating with quinofumelin at EC_{50} (0.035 μg/mL) and EC_{90} (1 μg/mL)
concentrations, the expression of the DHODH gene was respectively decreased by 11.91% and

33.77% ($P < 0.05$) (Supplementary Information, Fig. S4). This down-regulation hinders the

conversion of dihydroorotate to orotate, thereby inhibiting the biosynthesis of downstream
metabolites orotidine-5P and uracil, leading to their decreased abundance (Fig. 2e). Thus, we
conclude that the action mechanism of quinofumelin may involve inhibition of the *de novo*
biosynthesis pathway of pyrimidine.
Identification of DHODHII in *F. graminearum*
The *FgDHODHII* gene (FGSG_09678) was identified through the integrated analysis of

transcriptome and metabolome data. The gene sequence for *FgDHODHII* was retrieved from the
National Center for Biotechnology Information (NCBI) GenBank database. *FgDHODHII* is a
gene consisting of 1419 bp without introns and is predicted to encode a protein consisting of 472
amino acids. Orthologs of *FgDHODHII* were obtained using the BLAST algorithm. The
phylogenetic tree was constructed based on the amino acid sequences of DHODHII homologous
protein with Mega X using the neighbor-joining method. Motif pattern information was generated
using the MEME suite (Bailey et al., 2009), while functional domain information and protein
coordinates were obtained through CD-search (Marchler-Bauer et al., 2009). TBtools software
was used for visualizing the tree representation (Chen et al., 2020). All DHODHII proteins possess
the dihydroorotate dehydrogenase domain (Fig. 3), indicating high conservation among fungi.
Additionally, the amino acid sequence of the FgDHODHII exhibits 55.41% similarity to that of
DHODHII from *Pyricularia oryzae*, as previously reported (Higashimura et al., 2022).

Recovery test of mycelial growth suppressed by quinofumelin in *Fusarium genus*
The mycelial growth of *F. graminearum* PH-1 was significantly inhibited on CZA plates
containing quinofumelin at concentrations of 1 μg/mL (Fig. 4). To investigate the potential
recovery of mycelial growth by exogenous dihydroorotate, uridine monophosphate (UMP),
uridine, or uracil, CZA plates supplemented with quinofumelin were treated with these compounds.
Despite the addition of exogenous dihydroorotate at 50 μg/mL failing to restore mycelial growth,
the addition of exogenous UMP, uridine or uracil at 50 μg/mL successfully recovered
quinofumelin-induced inhibition of mycelial growth. Similar results were observed in other strains
of *F. graminearum* or *F. asiaticum* (Fig. 4 and Fig. S5). Thus, it can be concluded that
quinofumelin effectively inhibits the pyrimidine biosynthesis pathways.

The absence of *FgDHODHII* is lethal for mycelial growth in *F. graminearum*
To further investigate the biological functions of *FgDHODHII* in *F. graminearum*, we employed a
homologous recombination strategy to generate target gene deletion mutants. Through PCR
analysis, two putative *FgDHODHII* deletion mutants were successfully identified. The
ΔFgDHODHII mutants exhibited distinct phenotypic differences compared to the wild-type
progenitor PH-1 and the respective complemented strain (Fig. 5a). These mutants displayed an
inability to grow on CZA plate; however, their mycelial growth was restored when supplemented
with 50 μg/mL of exogenous UMP, uridine or uracil, while no recovery was observed with 50
μg/mL of exogenous dihydroorotate (Fig. 5b).

Molecular docking
To facilitate the molecular docking process between FgDHODHII and quinofumelin, ten distinct
binding models were generated using AutoDockTools-1.5.7 software, each accompanied by its
respective binding energies calculation results. The most favorable conformational arrangement
was determined based on its relative ranking in terms of the computed binding energy values and
subsequently visualized through PyMOL 2.5.4 software (Fig. 6a). Notably, the obtained docking
results revealed a robust interaction between FgDHODHII protein and quinofumelin as evidenced
by a calculated negative free energy value of -6.8 kcal/mol.

The precise interaction between FgDHODHII and quinofumelin was verified by Surface
plasmon resonance analysis
Surface plasmon resonance (SPR) was utilized to evaluate the interaction affinity between
quinofumelin and FgDHODHII in this study. The experiment was conducted using a BIAcore
T200 (Cytiva) instrument for biomolecular interaction analysis. FgDHODHII was immobilized as
the ligand on the chip surface, while quinofumelin continuously flowed through as the analyte

solution. Dilute quinofumelin to different concentration gradients (62.5, 32.15, 15.63, 7.81, 3.91
μM), measure the binding energy, and then fit the binding equilibrium dissociation constant (K_d)
value (Fig. 6b). The SPR results revealed that quinofumelin exhibited fast-binding and fast-
dissociation modes when interacting with the target protein FgDHODHII. The calculated K_D value
was determined as 6.606×10^{-6} M, indicating specific binding of quinofumelin to FgDHODHII.

The interaction between FgDHODHII and quinofumelin was confirmed by microscale
thermophoresis

In this study, microscale thermophoresis (MST) was further used to evaluate the interaction
affinity between FgDHODHII and quinofumelin. The strength of the binding can be inferred from
the Kd value. The experiment was conducted using MO.Control v1.6.1 and data analysis was
performed using MO. Affinity Analysis v2.3. By introducing a fluorescent label GFP to the
FgDHODHII protein while maintaining its concentration constant, non-fluorescent labeled small
molecules were diluted into 16 concentration gradients. Subsequently, the protein solution and
small molecule diluent were mixed at a ratio of 19:1 before being loaded into the capillary for
further analysis in the machine. The MST assay demonstrated that FgDHODHII exhibited specific
binding to quinofumelin in vitro, with a dissociation constant (Kd) of 0.471 ± 0.268 μM, whereas
no binding was observed with phenamacril which lacks cross-resistance with quinofumelin (Fig.
6c).

Discussion

In over 130 years of research on fungicide biology, more than 30 novel targets for selective
fungicides have been identified. Based on these newly discovered targets, over 200 modern
selective fungicides have been developed. Consequently, the identification of novel targets for
fungicides poses a significantly greater challenge compared to the development of new fungicides.
Currently, modern selective fungicides with single action site are widely applied in agricultural
practices for crop disease management. However, these specific types of fungicides encounter
substantial obstacles associated with resistance issues when deployed in agricultural fields.
Numerous studies have extensively documented cases of plant pathogens developing resistance to
modern selective fungicides, resulting in failures or reductions in their efficacy against plant
diseases and subsequently leading to decreased crop yields and quality (Duan et al., 2019; King et
al., 2021; Mao et al., 2024). For instance, *Botrytis cinerea*, a significant plant pathogenic fungus,
has emerged as multi-resistant population that is resistant to all currently used modern selective
fungicides (Shao et al., 2020; Weber et al., 2023). This poses a formidable challenge for the
control of crop grey mold disease caused by *B. cinerea*. Therefore, there is an urgent need for the
development of new fungicides with novel mechanisms of action to effectively manage multiple
resistance.

The novel quinoline, quinofumelin, developed by Mitsui Chemicals Co., Ltd. Japan, has
demonstrated remarkable antifungal activity against plant pathogens such as *F. graminearum* and
Sclerotinia sclerotiorum (Tao et al., 2021; Xiu et al., 2021). Moreover, our research group found
that quinofumelin exhibits no cross-resistance to all types of currently used modern selective
fungicides (Tao et al., 2021). These findings indicate that quinofumelin possesses a distinctive
mode of action against plant pathogens. Quinofumelin has been identified as a agrochemical
fungicide targeting DHODHII in *Pyricularia oryzae* (Higashimura et al., 2022). Similarly,
ipflufenoquin, another agrochemical fungicide with an analogous mechanism of action to
quinofumelin, has been reported to target DHODH in *Aspergillus fumigatus* (Rhijn et al., 2023).
To elucidate the mode of action of quinofumelin against *F. graminearum*, we conducted a

comprehensive investigation using integrated transcriptomic and metabolomic analyses. The
transcriptome data revealed that DEGs were involved in 'tryptophan metabolism' and 'amino
sugar and nucleotide sugar metabolism'. Furthermore, the metabolome data identified DAMs
enriched in 'nucleotide metabolism' and 'amino acid metabolism'. Interestingly, we observed a
co-enrichment of DEGs and DAMs in the pyrimidine metabolism pathway. Specifically, we
identified a down-regulation of dihydroorotate dehydrogenase (DHODH) in the experimental
groups, which impeded the conversion of dihydroorotate to orotate. Consequently, this inhibition
led to decreased biosynthesis of downstream metabolites orotidine-5P and uracil. Therefore, we
conducted an experiment using *F. graminearum* standard strain PH-1 to quantify mycelial growth
by supplementing exogenous uracil, which is the end product of the uracil biosynthesis pathway.
This was performed in the presence of complete inhibition of mycelial growth caused by
quinofumelin. Surprisingly, the addition of exogenous uracil restored the inhibited mycelial
growth induced by quinofumelin (Fig. 4). Meanwhile, similar results were successfully verified
with four other *F. graminearum* strains and five *F. asiaticum* strains (Fig. 4 and Fig. S4). Based on
our depicted pyrimidine biosynthesis pathway (Fig. 2E), we also supplemented other intermediate
products such as dihydroorotate, orotate, UMP, and uridine externally. Except for dihydroorotate
and orotate, supplementation of exogenous UMP and uridine effectively restored mycelial growth
inhibited by quinofumelin as well. These findings strongly support that quinofumelin acts through
modulation of the uracil biosynthesis pathway. Based on these findings and in conjunction with
previous studies ((Higashimura et al., 2022; Rhijn et al., 2023), these agrochemical fungicides are
categorized under FRAC group 52. This classification targets dihydroorotate dehydrogenase
(DHODH) during *de novo* pyrimidine biosynthesis pathway, as outlined by the Fungicide
Resistance Action Committee (https://www.frac.info/).As a key enzyme in the *de novo* pyrimidine
biosynthesis pathway, DHODH has previously been investigated as a potential target for inhibiting
infectious organisms and as an antitumor agent in mammals (Chen et al., 1992; Mao et al., 2021).
Inhibiting DHODH can effectively impede the synthesis of DNA, RNA, and glycoproteins,
thereby regulating aberrant cell metabolism and proliferation to ultimately achieve therapeutic
objectives (Boukalova et al., 2020). Within mammalian cells, DHODH is closely associated with
the mitochondrial membrane and plays a pivotal role in the respiratory chain through its
interaction with ubiquinone (Chen et al., 1976). Here, using amino acid sequences from *Homo*
sapiens DHODHII as a reference query, we have identified FgDHODHII (FGSG_09678), a gene
comprising of 1419 bp without any introns, which encodes 472 amino acids in *F. graminearum*.
FgDHODHII possesses a typical DHOD-2 like domain (NCBI: cd04738) and belongs to class 2
DHODs. Class 2 DHODHs are membrane-bound enzymes, whereas class 1 enzymes are located
in the cytosol (Nørager et al., 2002). The DHODH enzyme catalyzes a unique redox reaction in
the *de novo* pyrimidine biosynthesis pathway(Dorasamy et al., 2017). The DHOD-2 like domain
can bind orotate and flavin mononucleotide (FMN), facilitating the oxidation of dihydroorotate to
orotate and the reduction of the FMN cofactor (Alves et al., 2015). Based on the aforementioned
findings, we have concluded that DHODHII functions as the target of action for quinofumelin
against *F. graminearum*. Consequently, we performed a protoplast transformation assay to delete
FgDHODHII in *F. graminearum* and further validate the precise target of action for quinofumelin.
Typically, most fungicide targets are either lethal genes or essential pathogenic genes. Therefore, it
is plausible that the inactivation of FgDHODHII may be lethal. In our genetic transformation
experiment, exogenous uridine was supplemented to successfully obtain deletion mutants of the
FgDHODHII gene. Fortunately, two ΔFgDHODHII mutants were successfully obtained. These

mutants exhibited an inability to grow on PDA or CZA medium; however, mycelial growth was
fully restored upon addition of uridine to the medium (Fig. 5A). Additionally, exogenous
supplementation of UMP or uracil effectively restored mycelial growth. However, the
supplementation of dihydroorotate to the medium did not alleviate the growth defect observed in
ΔFgDHODHII mutants (Fig. 5B). Furthermore, mycelial growth in two ΔFgDHODHII mutants
was restored by exogenously adding UMP, uracil or uridine under quinofumelin treatments. These
findings indicate that the defects of FgDHODHII are functionally equivalent to complete
inhibition of its activity by quinofumelin and further confirm that the mechanism of action for
quinofumelin involves *de novo* synthesis of uracil while identifying FgDHODHII as its specific
target.
Molecular docking is a computational procedure employed to predict the optimal binding
orientation of a ligand with its macromolecular target (receptor), thereby forming a stable complex.
In concise terms, docking is a molecular modeling technique employed to predict the interaction
between a protein (enzyme) and small molecules (ligands). In this study, we conducted molecular
docking analysis between quinofumelin and the target protein FgDHODHII. The docking results
unveiled a robust affinity between FgDHODHII and quinofumelin. Furthermore, these findings
provided additional evidence that the strong affinity of quinofumelin with FgDHODHII disrupts
its activity and function.
Surface Plasmon Resonance (SPR) technique is employed for real-time measurement of molecular
interactions (Olaru et al., 2015). The SPR sensor map characterizes intermolecular interaction
rates primarily by conveying the kinetic rate constant, while it characterizes the binding level
between molecules mainly by conveying the affinity constant (Guo, 2012; Piliarik et al., 2009). In
this study, we used SPR technology to measure the binding energy and kinetics between
FgDHODHII and quinofumelin. The binding kinetics of FgDHODHII and quinofumelin
demonstrated fast-binding and fast-dissociation modes. Soluble FgDHODHII exhibited an ideal
affinity (Kd) of 6.606×10^{-6} M with quinofumelin, falling within the suitable range of 10^{-3} M to 10^{-6} M for biomacromolecule-small molecule interactions. Furthermore, the binding affinity between
FgDHODHII and quinofumelin was confirmed by a microscale thermophoresis (MST) assay.
Therefore, these data strongly support the precise interaction between quinofumelin and its target
protein FgDHODHII. This newly discovered selective target for antimicrobial agents offers a
valuable resource for the design and development of targeted pesticides.
Unlike many other dehydrogenases that require NAD or NADP as an electron acceptor, DHODH
transfers electrons to ubiquinone, which then transfers electrons into respiratory complex III and
eventually to oxygen(Wang et al., 2016). Thus, the production of orotate in pyrimidine
biosynthesis is coupled with the electron transport system and energy production, exerting
additional control over nucleotide synthesis and ensuring its coupling to cellular energy
metabolism. In the pyrimidine biosynthesis pathway, DHODHII regulates the conversion from
dihydroorotate to orotate. Our findings demonstrated that DHODHII is target of action for
quinofumelin. A previous study reported that exogenous orotate but not exogenous dihydroorotate
recovered mycelial growth inhibited by quinofumelin, indicating that quinofumelin inhibits the
formation of orotate in the pyrimidine biosynthesis pathway (Higashimura et al., 2022). However,
we observed that the supplementation of orotate to the medium did not alleviate the growth defect
of ΔFgDHODHII mutants. Considering the disparity in mycelial growth inhibition between *F.*
graminearum and other fungi, our data suggests that a nonclassical two-site ping-pong mechanism
is the most suitable model for DHODH as previously reported (Chen et al., 1992). In the classical

ping-pong mechanism, DHODH interacts with dihydroorotate and ubiquinol in an identical
confirmation, while orotate and ubiquinone combine with an alternative form (Cleland, 1963).
However, similar to brequinar sodium inhibiting the activity of DHODH isolated from mammalian
sources, FgDHODHII may follow a nonclassical, two-site ping-pong mechanism. In this
nonclassical ping-pong mechanism, the two substrates may not share chemical resemblance and
therefore do not bind to a single binding site but likely bind to two adjacent sites that facilitate
reactant transfer. The precise nature of the interaction between quinofumelin and FgDHODHII
may require further elucidation studies.

Materials and methods

Strains, culture conditions and fungicides

F. graminearum sensitive strain PH-1 and 9 wild-type strains (*F. graminearum* and *F. asiaticum*),
and the PH-1 deletion mutants ΔFgDHODHII were maintained in Fungicide Biology Laboratory,
Nanjing Agricultural University.
The media used in this study included potato dextrose agar (PDA) medium, Yeast extract peptone
dextrose (YEPD) medium, Czapek Solution Agar (CZA) medium and LB medium.
Chemical-grade quinofumelin (Mitsui Agricultural Chemical Company of Japan) was prepared
with dimethyl sulfoxide at a concentration of 1×10^4 μg/mL and stored at 4°C.

Transcriptomic assay of mycelial growth in *F. graminearum* as affected by quinofumelin

A total of 20 mL of YEPD medium containing 1 mL of conidia suspension (1×10^5 conidia/mL)
was incubated with shaking (175 rpm/min) at 25°C. After 24 h, the medium was added with
quinofumelin (EC_{90}) at a concentration of 1 μg/mL, while an equal amount of dimethyl sulfoxide
was added as the control (CK). The incubation continued for another 48 h, followed by filtration
and collection of hyphae. Subsequently, Genedenovo Biotechnology Co., Ltd. (Guangzhou, China)
performed sequencing on the Illumina platform. Quality control measures were implemented to
remove linker sequences and low-quality bases from the original reads in order to obtain clean
data readings. The filtered data underwent analysis for base composition and mass distribution to
visualize data quality. Based on the gene expression data, we used R (http://www.r-project.org/)
for conducting principal component analysis, while the "pheatmap" package was used to generate
the clustering heatmap. DESeq2 was applied to analyze group differences based on quantitative
expression results. The screening threshold of P value <0.05 and FC≥1.5 or FC≤0.67 was used to
identify unigenes with significant differential expression between the two groups using a t-test.
Finally, functional enrichment analysis of GO and KEGG pathways was performed on these
differentially expressed unigenes.

Metabonomic assay of mycelial growth in *F. graminearum* as affected by quinofumelin

Transcriptome and metabolome samples were obtained from the same treatment group. The
widely targeted metabolome analysis was conducted by Metware Biotechnology Co., Ltd. (Wuhan,
China). The frozen sample stored at -80°C was thawed on ice. A 400 μL solution (Methanol: Water
= 7:3, v/v) containing an internal standard was added to a 20 mg sample and vortexed for 3 min.
Subsequently, the sample was sonicated in an ice bath for 10 min and vortexed for 1 min before
being placed at -20°C for 30 min. After centrifugation at 12000 rpm for 10 min (4°C). The
sediment was removed, followed by another centrifugation of the supernatant at 12000 rpm for 3
min (4°C). A volume of 200 μL aliquots of supernatant was transferred for LC-MS analysis. All
samples were acquired using the LC-MS system according to machine orders. The original data
file acquired by LC-MS was converted into mzML format using ProteoWizard software. Peak

extraction, peak alignment, and retention time correction were performed by XCMS program.

The identified metabolites were analyzed using MBEole2.0

(https://csbg.cnb.csic.es/mbrole2/index.php) and KEGG (http://www.kegg.jp). PLS-DA analysis

was used to identify the differentially accumulated metabolites (DAMs), with a screening

threshold of P value<0.05 and FC≥1.5 or FC≤0.67. Each sample was replicated six times for

biological consistency.

Recovery assay of quinofumelin-induced mycelial growth inhibition in *F. graminearum*

F. graminearum sensitive strain PH-1 and 9 wild-type strains (*F. graminearum* and *F. asiaticum*),

and the PH-1 deletion mutants ΔFgDHODHII, were preincubated on a PDA plate to form mycelial

colonies. Mycelial plugs (5 mm in diameter) from the edge of a 3-day-old colony were removed

and inoculated onto a CZA plate containing quinofumelin (1 μg/mL) and dihydroorotate, UMP,

uridine or uracil at 50 μg/mL, followed by measurement of colony diameters. Three replicates

were conducted for each concentration.

Construction of deletion mutant vectors

The deletion mutant of *FgDHODHII* was generated through protoplast transformation, as

previously described with some modifications in the transformation procedures(Zhang et al.,

2016). For the transformation process, washed protoplasts were suspended in 160 μL sorbitol-tris-

calcium chloride (STC) solution and 40 μL sorbitol-peg-tris-calcium chloride (SPTC) solution (40%

polyethylene glycol 6000 in STC), along with 2 μg of the aforementioned DNA fragments and 10

μL of heparin (5 mg/mL in STC) to form protoplast suspensions. Subsequently, these mixtures

were incubated on ice for 30 min, followed by mixing with 1 mL of SPTC and further incubation

at room temperature for 20 min. The resulting suspensions were gently mixed into regeneration

medium (RM) (100 mL) containing 50 μg/mL uridine and poured into plates. After incubating for

24 h at 25°C, the plates were overlaid with selective regeneration medium (SRM) containing 50

μg/mL uridine and continuously incubated for an additional 4 days. The putative transformants

could be observed growing on SRM. All primers used are listed in Table 1.

Molecular docking

The binding affinity of quinofumelin with FgDHODHII protein was assessed through molecular

docking using AutoDockTools-1.5.7 software. The three-dimensional structure of quinofumelin

(PubChem CID: 23160856) used for docking was obtained from PubChem database, while the

three-dimensional structural model of *F. graminearum* DHODHII protein predicted by AlphaFold

(Jumper et al., 2021) was obtained from the UniProt database (Bateman et al., 2019). In the

docking studies, binding energy was employed as a metric to evaluate the strength of protein-

ligand interaction, where lower values indicated higher affinity between proteins and ligands. The

resulting docking outcomes were visualized using PyMOL software version 2.5.4.

Surface plasmon resonance (SPR) verification of specific interactions between FgDHODHII

and quinofumelin

The cDNA sequence of FgDHODHII was analyzed and codon optimization was performed for the

E. coli expression system (Kyriakopoulos et al., 2013; Yang et al., 2021). Subsequently, it was

ligated into the pCold-9×his-MBP plasmid and the pET-28a plasmid at the NdeI and EcoRI sites

within the multiple cloning sites to construct recombinant plasmids: pCold-9×his-MBP-TEV-

DHODHII (Fang et al., 2018), pET-28a (+)-DHODHII and pET-28a (+)-DHODHII[124-467] (with

removal of insoluble fragments) (Supplementary Information, Fig. S6).

Plasmid DNA was extracted using the E.Z.N.A Plasmid Extraction DNA Mini Kit I protocol

(Omega, China). For protein overexpression, the exacted plasmids were separately transformed

into *E. coli* BL21 strain (Zhu et al., 2014). A single clone obtained from each transformation was
used for protein overexpression and purification. Upon reaching an OD600 of 1.0, 0.6 mM
isopropyl-β-D-thiogalactopyranoside (IPTG) was added to induce protein overexpression.
Following induction at 16°C for 9 h, cells were harvested, lysed, and purified using a Ni-NTA
column (QIAGEN).
The interaction between FgDHODHII and quinofumelin was evaluated using Biacore T200
instrument (GE Healthcare). The CM7 sensor chip was employed for immobilizing 50 µg/mL of
purified FgDHODHII protein via amino coupling method at a flow rate of 10 µL/min, utilizing
1×PBS-P+ containing 5% DMSO as running buffer. The desired coupling amount of FgDHODHII
protein on the chip surface was targeted to be within the range of 12000-24000 RU. To investigate
binding affinity, quinofumelin concentrations were diluted from 3.90625 to 62.5 µM in a constant
flow rate running buffer containing 5% DMSO. Following sample injection, a switching buffer
was introduced to allow spontaneous dissociation of quinofumelin from FgDHODHII on the chip
surface, and this dissociation process was continuously monitored in real time by measuring
response values. The affinity value (Kd) was determined through kinetic analysis or steady-state
affinity method.

Identification of specific interactions between FgDHODHII and quinofumelin using
microscale thermophoresis (MST)

To construct the FgDHODHII-GFP cassette, the FgDHODHII fragment containing the native
promoter and ORF (without the stop codon) was amplified. Subsequently, the resulting PCR
product was co-transformed with XhoI-digested pYF11 into *E. coli* DH5α as previously reported
(Tang et al., 2020). The fusion vector was then used to transform PH-1 to obtain the FgDHODHII-
GFP strain. The GFP fusion protein was extracted from mycelia. Two Monolith NT.115 standard
capillaries were employed for repeated absorption of a specific protein concentration, and placed
in positions 1 and 2 of the sample tray in the MonoTemper Monolith NT.115 instrument. Sixteen
PCR tubes were prepared, with 10 µL DMSO added to PCR tubes 2-16 and 20 µL of 5000 µM
quinofumelin (or phenamacril) dissolved in DMSO added to PCR tube 1. Then, 10 µL of
quinofumelin from PCR tube 1 was transferred and thoroughly mixed with PCR tube 2 using a
pipette. These steps were repeated for PCR tubes 3-16. Following mixing with PCR tube 16
(discarding the remaining 10 µL), each PCR tube was thoroughly mixed with 190 µL DHODHII-
GFP protein, and incubated at room temperature for 30 min. Monolith NT.115 standard capillaries
were inserted into each PCR tube to allow solution entry into the capillary. The capillaries 1-16
were subsequently placed in the sample tray of the device and the program was initiated. All
measurements were performed using 80% (Auto-detect) Nano-RED. MonoTemper Monolith
NT.115 software was used to evaluate the binding affinity between the target protein and
quinofumelin using the Kd values and the signal to noise ratio , and the results were combined and
analyzed using Mo. Affinity Analysis v2.3 software (Rotem et al., 2015).

Statistical analysis

The data in the study were analyzed using analysis of variance (ANOVA) with the SPSS 14.0
software (SPSS Inc. Chicago, IL, USA). When ANOVA yielded a significant result ($p = 0.05$),
means were compared using Fisher's protected least significant difference (PLSD).

Acknowledgements

This work was supported by the National Key Research and Development Program of China
(2022YFD1400100), and the National Natural Science Foundation of China (32072448).

Conflicts of interest

The authors declare no competing interest.

Data availability

Data will be made available on request.

References

Alves, C., Silva, J., & Roitberg, A. (2015). Insights into the Mechanism of Oxidation of Dihydroorotate
to Orotate Catalysed by Human Class 2 Dihydroorotate Dehydrogenase: A QM/MM Free
Energy Study. *Physical Chemistry Chemical Physics*, *17*. https://10.1039/C5CP02016F

Bailey, T. L., Boden, M., Buske, F. A., Frith, M., Grant, C. E., Clementi, L., Ren, J., Li, W. W., & Noble,
W. S. (2009). MEME Suite: tools for motif discovery and searching. *Nucleic Acids Research*,
37(suppl_2), W202-W208. https://10.1093/nar/gkp335

Bateman, A., Martin, M. J., Orchard, S., Magrane, M., Alpi, E., Bely, B., Bingley, M., Britto, R.,
Bursteinas, B., Busiello, G., Bye-A-Jee, H., Da Silva, A., De Giorgi, M., Dogan, T., Castro, L.
G., Garmiri, P., Georghiou, G., Gonzales, D., Gonzales, L., Hatton-Ellis, E., Ignatchenko, A.,
Ishtiaq, R., Jokinen, P., Joshi, V., Jyothi, D., Lopez, R., Luo, J., Lussi, Y., MacDougall, A.,
Madeira, F., Mahmoudy, M., Menchi, M., Nightingale, A., Onwubiko, J., Palka, B., Pichler, K.,
Pundir, S., Qi, G. Y., Raj, S., Renaux, A., Lopez, M. R., Saidi, R., Sawford, T., Shypitsyna, A.,
Speretta, E., Turner, E., Tyagi, N., Vasudev, P., Volynkin, V., Wardell, T., Warner, K., Watkins,
X., Zaru, R., Zellner, H., Bridge, A., Xenarios, I., Poux, S., Redaschi, N., Aimo, L., Argoud-
Puy, G., Auchincloss, A., Axelsen, K., Bansal, P., Baratin, D., Blatter, M. C., Bolleman, J.,
Boutet, E., Breuza, L., Casals-Casas, C., de Castro, E., Coudert, E., Cuche, B., Doche, M.,
Dornevil, D., Estreicher, A., Famiglietti, L., Feuermann, M., Gasteiger, E., Gehant, S.,
Gerritsen, V., Gos, A., Gruaz, N., Hinz, U., Hulo, C., Hyka-Nouspikel, N., Jungo, F., Keller,
G., Kerhornou, A., Lara, V., Lemercier, P., Lieberherr, D., Lombardot, T., Martin, X., Masson,
P., Morgat, A., Neto, T. B., Paesano, S., Pedruzzi, I., Pilbout, S., Pozzato, M., Pruess, M.,
Rivoire, C., Sigrist, C., Sonesson, K., Stutz, A., Sundaram, S., Tognolli, M., Verbregue, L., Wu,
C. H., Arighi, C. N., Arminski, L., Chen, C. M., Chen, Y. X., Cowart, J., Garavelli, J. S.,
Huang, H. Z., Laiho, K., McGarvey, P., Natale, D. A., Ross, K., Vinayaka, C. R., Wang, Q. H.,
Wang, Y. Q., Yeh, L. S., Zhang, J., & UniProt, C. (2019). UniProt: a worldwide hub of protein
knowledge. *Nucleic Acids Research*, *47*(D1), D506-D515. https://10.1093/nar/gky1049

Boukalova, S., Hubackova, S., Milosevic, M., Ezrova, Z., Neuzil, J., & Rohlena, J. (2020).
Dihydroorotate dehydrogenase in oxidative phosphorylation and cancer. *Biochimica et*
Biophysica Acta (BBA) - Molecular Basis of Disease, *1866*(6), 165759.
https://doi.org/10.1016/j.bbadis.2020.165759

Chen, C., Chen, H., Zhang, Y., Thomas, H. R., Frank, M. H., He, Y., & Xia, R. (2020). TBtools: An
Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. *Molecular*
Plant, *13*(8), 1194-1202. https://doi.org/10.1016/j.molp.2020.06.009

Chen, J., & Jones, M. E. (1976). The cellular location of dihydroorotate dehydrogenase: Relation to de
novo biosynthesis of pyrimidines. *Archives of Biochemistry and Biophysics*, *176*(1), 82-90.
https://doi.org/10.1016/0003-9861(76)90143-0

Chen, S., Perrella, F., Behrens, D., & Papp, L. (1992). Inhibition of Dihydroorotate Dehydrogenase
Activity by Brequinar Sodium. *Cancer Research*, *52*, 3521-3527.

Cleland, W. W. (1963). The kinetics of enzyme-catalyzed reactions with two or more substrates or
products: III. Prediction of initial velocity and inhibition patterns by inspection. *Biochimica et*

Biophysica Acta, *67*, 188-196. https://doi.org/10.1016/0926-6569(63)90227-X

Dorasamy, M. S., Choudhary, B., Nellore, K., Subramanya, H., & Wong, P.-F. (2017). Dihydroorotate dehydrogenase Inhibitors Target c-Myc and Arrest Melanoma, Myeloma and Lymphoma cells at S-phase. *Journal of Cancer*, *8*(15), 3086-3098. https://10.7150/jca.14835

Duan, Y., Xin, W., Lu, F., Li, T., Li, M., Wu, J., Wang, J., & Zhou, M. (2019). Benzimidazole- and QoI-resistance in *Corynespora cassiicola* populations from greenhouse-cultivated cucumber: An emerging problem in China. *Pesticide Biochemistry and Physiology*, *153*, 95-105. https://doi.org/10.1016/j.pestbp.2018.11.006

Fang, K., Li, Y., Yin, X., Samad, A., & Jin, T. (2018). High-Level Prokaryotic Expression and Purification of Death Domain Superfamily with MBP Tag. *Clinical Laboratory*, *64*(4), 467-475. https://10.7754/clin.lab.2017.170922

Guo, X. (2012). Surface plasmon resonance based biosensor technique: A review. *Journal of Biophotonics*, *5*(7), 483-501. https://doi.org/10.1002/jbio.201200015

Higashimura, N., Hamada, A., Ohara, T., Sakurai, S., Ito, H., & Banba, S. (2022). The target site of the novel fungicide quinofumelin, *Pyricularia oryzae* class II dihydroorotate dehydrogenase. *Journal of Pesticide Science*, *47*(4), 190-196. https://10.1584/jpestics.D22-027

Ito, H., Takada, T., Morimoto, M., Komai, H., Kajino, F., Ohara, T., Tamagawa, Y., Tsuda, M., & Banba, S. (2023). Design and biological activity of a novel fungicide, quinofumelin. *Journal of Pesticide Science*, *48*(1), 22-27. https://10.1584/jpestics.D22-042

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P., & Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583-589. https://10.1038/s41586-021-03819-2

King, K., Bucur, D., Ritchie, F., Hawkins, N., Kaczmarek, A., Duan, Y., Kildea, S., West, J., & Fraaije, B. (2021). Fungicide resistance status and chemical control options for the brassica pathogen *Pyrenopeziza brassicae*. *Plant Pathology*, *70*. https://10.1111/ppa.13441

Kyriakopoulos, S., & Kontoravdi, C. (2013). Analysis of the landscape of biologically-derived pharmaceuticals in Europe: Dominant production systems, molecule types on the rise and approval trends. *European Journal of Pharmaceutical Sciences*, *48*(3), 428-441. https://doi.org/10.1016/j.ejps.2012.11.016

Lamberth, C. (2022). Latest Research Trends in Agrochemical Fungicides: Any Learnings for Pharmaceutical Antifungals? *ACS Medicinal Chemistry Letters Journal*, *13*(6), 895-903. https://10.1021/acsmedchemlett.2c00113

Lamberth, C., Rendine, S., & Sulzer-Mosse, S. (2021). Chapter 4 - Agrochemical disease control: The story so far. In P. Maienfisch & S. Mangelinckx (Eds.), (pp. 65-85): Adv Vet Emd-Acad Pr.

Lucas, J. A., Hawkins, N. J., & Fraaije, B. A. (2015). The Evolution of Fungicide Resistance. In S. Sariaslani & G. M. Gadd (Eds.), *Advances in Applied Microbiology, Vol 90*, vol. 90 (pp. 29-92).

Mao, C., Liu, X., Zhang, Y., Lei, G., Yan, Y., Lee, H., Koppula, P., Wu, S., Zhuang, L., Fang, B., Poyurovsky, M. V., Olszewski, K., & Gan, B. (2021). DHODH-mediated ferroptosis defence is a targetable vulnerability in cancer. *Nature*, *593*(7860), 586-590. https://10.1038/s41586-021-03539-7

Mao, Y., Meng, X., Zhang, Z., Zhang, L., Feng, Y., Wang, T., Zhang, J., Cai, Y., Wang, J., Sheng, G., Shen, Y., Zhou, M., & Duan, Y. (2024). Genetic variation of drug target CYP51 conferring resistance to ergosterol biosynthesis inhibitors in *Botrytis cinerea*, causing lily gray mould. *Industrial Crops and Products*, *208*, 117797. https://doi.org/10.1016/j.indcrop.2023.117797

Marchler-Bauer, A., Anderson, J. B., Chitsaz, F., Derbyshire, M. K., DeWeese-Scott, C., Fong, J. H., Geer, L. Y., Geer, R. C., Gonzales, N. R., Gwadz, M., He, S., Hurwitz, D. I., Jackson, J. D., Ke, Z., Lanczycki, C. J., Liebert, C. A., Liu, C., Lu, F., Lu, S., Marchler, G. H., Mullokandov, M., Song, J. S., Tasneem, A., Thanki, N., Yamashita, R. A., Zhang, D., Zhang, N., & Bryant, S. H. (2009). CDD: specific functional annotation with the Conserved Domain Database. *Nucleic Acids Research*, *37*(suppl_1), D205-D210. https://10.1093/nar/gkn845

Morton, V., & Staub, T. (2008). A Short History of Fungicides. *APSnet Features*. https://10.1094/APSnetFeature-2008-0308

Nørager, S., Jensen, K. F., Björnberg, O., & Larsen, S. (2002). *E. coli* Dihydroorotate Dehydrogenase Reveals Structural and Functional Distinctions between Different Classes of Dihydroorotate Dehydrogenases. *Structure*, *10*(9), 1211-1223. https://doi.org/10.1016/S0969-2126(02)00831-6

Olaru, A., Bala, C., Jaffrezic-Renault, N., & Aboul-Enein, H. Y. (2015). Surface Plasmon Resonance (SPR) Biosensors in Pharmaceutical Analysis. *Critical Reviews in Analytical Chemistry*, *45*(2), 97-105. https://10.1080/10408347.2014.881250

Pertea, M., Kim, D., Pertea, G. M., Leek, J. T., & Salzberg, S. L. (2016). Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature Protocols*, *11*(9), 1650-1667. https://10.1038/nprot.2016.095

Piliarik, M., Vaisocherová, H., & Homola, J. (2009). Surface Plasmon Resonance Biosensing. In A. Rasooly & K. E. Herold (Eds.), *Biosensors and Biodetection*, (pp. 65-88). Totowa, NJ: Humana Press.

Rotem, O., Nesper, J., Borovok, I., Gorovits, R., Kolot, M., Pasternak, Z., Shin, I., Glatter, T., Pietrokovski, S., Jenal, U., & Jurkevitch, E. (2015). An Extended Cyclic Di-GMP Network in the Predatory Bacterium *Bdellovibrio bacteriovorus*. *Journal of Bacteriology*, *198*(1), 127-137. https://10.1128/jb.00422-15

Shao, W., Zhao, Y., & Ma, Z. (2020). Advances in Understanding Fungicide Resistance in Botrytis cinerea in China. *Phytopathology*, *111*(3), 455-463. https://10.1094/PHYTO-07-20-0313-IA

Tang, G., Chen, A., Dawood, D. H., Liang, J., Chen, Y., & Ma, Z. (2020). Capping proteins regulate fungal development, DON-toxisome formation and virulence in *Fusarium graminearum*. *Molecular Plant Pathology*, *21*(2), 173-187. https://doi.org/10.1111/mpp.12887

Tao, X., Zhao, H., Xu, H., Li, Z., Wang, J., Song, X., Zhou, M., & Duan, Y. (2021). Antifungal Activity and Biological Characteristics of the Novel Fungicide Quinofumelin Against *Sclerotinia sclerotiorum*. *Plant Disease*, *105*(9), 2567-2574. https://10.1094/PDIS-08-20-1821-RE

Umetsu, N., & Shirai, Y. (2020). Development of novel pesticides in the 21st century. *Journal of Pesticide Science*, *45*(1-2), 54-74. https://10.1584/jpestics.D20-201

Wang, Y., & Hekimi, S. (2016). Understanding Ubiquinone. *Trends in Cell Biology*, *26*(5), 367-378. https://doi.org/10.1016/j.tcb.2015.12.007

Weber, R. W. S., & Petridis, A. (2023). Fungicide Resistance in *Botrytis spp.* and Regional Strategies for Its Management in Northern European Strawberry Production. In *BioTech*, vol. 12 (pp. 64).

Wheeler, I. E., Hollomon, D. W., Gustafson, G., Mitchell, J. C., Longhurst, C., Zhang, Z., & Gurr, S. J. (2003). Quinoxyfen perturbs signal transduction in barley powdery mildew (*Blumeria*

graminis f.sp. *hordei*). *Molecular Plant Pathology*, *4*(3), 177-186. https://doi.org/10.1046/j.1364-3703.2003.00165.x

Xiu, Q., Bi, L., Xu, H., Li, T., Zhou, Z., Li, Z., Wang, J., Duan, Y., & Zhou, M. (2021). Antifungal Activity of Quinofumelin against *Fusarium graminearum* and Its Inhibitory Effect on DON Biosynthesis. *Toxins*, *13*(5), 348. https:// https://doi.org/10.3390/toxins13050348

Yang, D., Prabowo, C. P. S., Eun, H., Park, S. Y., Cho, I. J., Jiao, S., & Lee, S. Y. (2021). *Escherichia coli* as a platform microbial host for systems metabolic engineering. *Essays in Biochemistry*, *65*(2), 225-246. https://10.1042/EBC20200172

Yang, X., Wang, Y. F., Byrne, R., Schneider, G., & Yang, S. Y. (2019). Concepts of Artificial Intelligence for Computer-Assisted Drug Discovery. *Chemical Reviews*, *119*(18), 10520-10594. https://10.1021/acs.chemrev.8b00728

Zhang, L., Li, B., Zhang, Y., Jia, X., & Zhou, M. (2016). Hexokinase plays a critical role in deoxynivalenol (DON) production and fungal development in *Fusarium graminearum*. *Molecular Plant Pathology*, *17*(1), 16-28. https://doi.org/10.1111/mpp.12258

Zhong, L., Li, Y. S., Xiong, L., Wang, W. J., Wu, M., Yuan, T., Yang, W., Tian, C. Y., Miao, Z., Wang, T. Q., & Yang, S. Y. (2021). Small molecules in targeted cancer therapy: advances, challenges, and future perspectives. *Signal Transduction and Targeted Therapy*, *6*(1). https://10.1038/s41392-021-00572-w

Zhu, L., Zhang, X., Chang, L., Wang, A., Feng, P., & Han, L. (2014). Molecular cloning, prokaryotic expression and promoter analysis of squalene synthase gene from *Schizochytrium Limacinum*. *Applied Biochemistry and Microbiology*, *50*(4), 411-419. https://10.1134/S0003683814040140

Figures:

Fig. 1 Analysis of transcriptome sequencing data in *F. graminearum* PH-1 treated with
quinofumelin.
a Bar chart illustrating the comparison of differentially expressed genes between EC$_{90}$ treatments
and the control.
b Volcano plot depicting up-regulated DEGs as red scattered dots and down-regulated DEGs as
blue scattered dots.
c Cluster heat maps displaying gene relative expression values using RNA-seq data scale-
standardized values.
d Bubble map presenting the results of GO enrichment analysis for DEGs.
e Bubble map showcasing the findings of KEGG enrichment analysis for DEGs.

Fig.2 Difference analysis and enrichment analysis of metabolome data in *F. graminearum* PH-1 treated with quinofumelin.

a Bar chart illustrating the number of differential accumulation metabolites (DAMs) quantity in experimental groups (EC_{90}) and control groups (CK) in positive ion mode.

b Volcanic map of all metabolites in positive ion mode. The red scatter points represent up-regulated DAMs, while the blue scatter points represent down-regulated DAMs.

c Classification and proportion of DAMs.

d Bubble diagram of differential accumulation metabolite KEGG enrichment pathway.

e Pyrimidine *de novo* biosynthesis pathway in *Fusarium graminearum*. The left box plots represent the relative content of 8 metabolite samples, while the right box represents the relative content of 6 gene samples. The color gradient from navy blue to firebrick red indicates a progression from low to high content.

Fig. 3 Phylogenic tree of DHODHII proteins. The phylogenetic tree was constructed based on the amino acid sequences of DHODHII homologous protein with Mega X using the neighbor-joining method. The bootstrap values from 1000 replications are indicated on the branches. Motif pattern information was generated using the MEME suite12, while functional domain information and protein coordinates were obtained through CD-search. The final phylogenetic tree was visualized using TBtools. FgDHODHII was highlighted in red. The amino acid sequence of *DHODHII* from *Alternaria alternata* (AA0117_g542), *Aspergillus nidulans* (ANIA_05909), *Aspergillus oryzae* (AO1008_05965), *Blumeria graminis* (BLGH_00861), *Botrytis cinerea* (Bcin15g04150), *Botrytis elliptica* (BELL_0162g00150), *Colletotrichum asianum* (GQ607_007053), *Colletotrichum fioriniae* (CFIO01_03831), *Colletotrichum higginsianum* (CH063_10143), *Corynespora cassiicola* (BS50DRAFT_576390), *Erysiphe necator* (EV44_g2092), *Erysiphe pulchra* (EPUL_004094), *Fusarium euwallaceae* (BHE90_002825), *Fusarium floridanum* (CEP51_012005), *Fusarium fujikuroi* (FFMR_05040), *Fusarium graminearum* (FGSG_09678), *Fusarium nygamai* (FNYG_10990), *Fusarium odoratissimum* (FOIG_08270), *Fusarium oxysporum* (FOXG_08390), *Fusarium pseudograminearum* (FPSE_04283), *Fusarium solani* (NechaG15121), *Fusarium verticillioides* (FVEG_06288), *Magnaporthe oryzae* (MGG_08814), *Neurospora crassa* (NCU06532), *Oidium neolycopersici* (OnM2_089032), *Pestalotiopsis fici* (PFICI_04984), *Podospora anserina* (PODANS_6_7920), *Puccinia graminis* (PGTG_16739), *Puccinia triticina* (PTTG_02070), *Rhizoctonia solani* (RSOLAG22IIIB_08327), *Schizosaccharomyces pombe* (SPAC57A10.12c), *Ustilaginoidea virens* (UVI_02052880), *Verticillium alfalfae* (VDBG_01322), *Verticillium dahliae* (VDAG_07935), and *Zymoseptoria tritici* (Mycgr3G72548) were accessed in EnsemblFungi database.

Fig. 4 Recovery test of mycelial growth suppressed by quinofumelin.
All strains were incubated on CZA plates at 25°C for 3 days. *F. graminearum* strains PH-1, 512E-8, SX2117, XY-4, and R10. The left image shows the colony morphology, while the right image is a bar chart of colony diameters.

703
Fig. 5 Mycelial growth of FgDHODHII deletion mutants on CZA plates.
a Mycelial growth of the FgDHODHII deletion mutants and the parental strain was observed in
the presence or absence of 50 μg/mL uridine.
b Recovery of mycelial growth of the FgDHODHII deletion mutants was observed in the presence
of 50 μg/mL dihydroorotate, UMP, uridine or uracil. All strains were incubated at 25°C for 3
days.

Fig. 6 Validation and analysis of quinofumelin's binding affinity to FgDHODHII.

a Three-dimensional model of the wild-type FgDHODHII are depicted, with an enlarged view highlighting the binding site for quinofumelin on the corresponding FgDHODHII models.

b The SPR response (RU) analysis and fitting results demonstrate the interaction between quinofumelin and FgDHODHII.

c Microscale thermophoresis (MST) analysis to detect the binding of quinofumelin and FgDHODHII.

720 **Fig. 7 A model of the pyrimidine pathway that quinofumelin attacks.**

722   **Tables:**

723   **Table 1 Primers used in the study**

Primer	Sequence (5'-3')	Use
FgDHODHII-UF	GGCAGTGAAAGTCTTGTTCAAG	PCR primers to
FgDHODHII-UR	GCTCCTTCAATATCATCTTCTGTGGC	amplify FgDHODHII
	TGGGAAGTAATGCTGGG	upstream fragment
FgDHODHII-DF	GAGACAATACCGGAAGGAACAC	PCR primers to
	ATGCTATAGCAAGCAACA	amplify FgDHODHII
FgDHODHII-DR	AAACTTAACAAATCATCTTTGCC	downstream fragment
SS-F	ACAGAAGATGATATTGAAGGAGC	PCR primers to
SS-R	GAGACAATACCGGAAGGAA	amplify Hph-Hsv-tk
		fragment
FgDHODHII-YZ-F	TTTGCCCGATCTCTTGGTCT	PCR primers to
FgDHODHII-YZ-R	AACCACCAGTCTCCATGAGG	verify FgDHODHII
FgDHODHII-GFP-F	CTCATCACCATCACCATCACAT	
	GTCTGCCGCTCTCCTACG	Constructed
		FgDHODHII-GFP
FgDHODHII-GFP-R	AGCTCCTCGCCCTTGCTCACCC	fusion vector
	TCGAATTCTTCACACCAC	
T7	TAATACGACTCACTATAGG	
T7t	CCGCTAGTTATTGCTCAGC	
pET-28a-F	CCTGGTGCCGCGCGGCAGCCATATGCTGG	
	CTGTCAATGTTTTTGGAACAG	PCR primers to
pET-28a-R	AAGCTTGTCGACGGAGCTCGAATTCCTAT	construct FgDHODHII
	TTTAACTCCTGCTTGATCCTG	expression vector
pCold-F	GTAAAGCACGCCATATCGC	
pCold-R	CCAAATGGCAGGGATCTTAG	

724