Developmentally cascading structures do not lose evolutionary potential, but compound

developmental instability in rat molars

Natasha S. Vitek[1], Ella Saks[1], Amy Dong[1], Robert W. Burroughs [1], Devin L Ward[2], Emma

Pomeroy[3], Malgorzata Martin-Gronert[4], Susan E. Ozanne[4]

[1] Department of Ecology & Evolution, Stony Brook University, Stony Brook, NY.

[2] Department of Anthropology, University of Toronto, Toronto, Canada

[3] Department of Archaeology and Newnham College, University of Cambridge, Cambridge, UK

[4] Metabolic Research Laboratories and MRC Metabolic Diseases Unit, Institute of Metabolic

Science, University of Cambridge, Cambridge, UK

ORCID:

NSV: 0000-0003-0587-8907

RWB: 0000-0003-4384-3430

DLW: 0000-0002-2343-8238

EP: 0000-0001-6251-2165

MM-G: 0000-0002-6763-8215

SEO: 0000-0001-8753-5144

Keywords: evolvability, phenotypic plasticity, nutrition, growth, molar, Mammalia

Abstract

Increasing variability down serially segmented structures, such as mammalian molar teeth and

vertebrate limb segments, is a much-replicated pattern. The same phenotypic pattern has

conflicting interpretations at different evolutionary scales. Macroevolutionary patterns are

thought to reflect greater evolutionary potential in later-forming segments, but microevolutionary

patterns are thought to reflect less evolutionary potential and greater phenotypic plasticity. We

address this conflict by recalculating evolutionary potential (evolvability) from a systematic

review of published mammalian molar sizes, then directly measure phenotypic plasticity from a

controlled feeding experiment. Effects on lengths and widths are discordant in a way that

suggests general growth pathways have a role in phenotypically plastic dental responses to

nutrition. Effects on successive trait means do not necessarily increase downstream, contrary to

long-standing hypotheses. We confirm prior findings of increasing non-inherited variance

downstream, showing decoupling between effects on trait mean and variance. These patterns can

be explained by a cascading model of tooth development compounding the effect of

developmental instability as an influence separate from general environmental effects on the

developing embryo. When evaluated in terms of evolvability, later-developing molars are

equally or more evolvable than earlier-developing molars, aligning their microevolutionary

potential with macroevolutionary patterns in other serially segmented structures.

Introduction

Animal bodies contain numerous serially segmented structures, such as vertebrate teeth, limb

bones, and vertebrae (Young *et al.*, 2015). The semi-independent nature of serial structures

makes them important systems for understanding how diverse phenotypes evolve in structures

that share a significant proportion of their genetic and developmental basis (Kavanagh *et al.*,

2007; Böhmer *et al.*, 2015).

In mammalian teeth as an example, divergent perspectives emerge on the relative

evolutionary potential of different segments in a module. In macroevolution, where phenotype is

largely understood to have a genetic basis, later-developing segments such as the third molars

(M_3) are considered to have greater evolutionary potential than early-developing, upstream

segments such as the first or second molars (M_1, M_2). Support comes from observations of faster

evolutionary rates downstream ($M_3 > M_2 > M_1$) sometimes documented directly (Sofaer *et al.*,

1971; Mongle *et al.*, 2022; Amaral *et al.*, 2024), sometimes implied by greater taxonomic utility

of the M_3 as a more sensitive indicator of species differences, especially in rodents and suoids

(Hershkovitz, 1967; Barnosky, 1987; Van der Made, 1996; Polly, 2003). These observations

translate into higher macroevolutionary diversity, or variance, in M_3s vs. earlier-developing

segments (Billet & Bardin, 2021).

In microevolutionary contexts, where phenotype likely has both inherited genetic basis and

non-inherited plastic basis, molars usually, but not always, share the same predicted pattern of

higher phenotypic variation in later-forming segments (ex., (Schuman & Brace, 1954; Bader &

Lehmann, 1965; Gingerich, 1974; Cuozzo, 2008) but see (Emery-Wetherell & Davis, 2018) for

a counter-example). Along the same sequences of segments, two other patterns emerge. First,

successive molars generally (though not exclusively) have lower levels of heritability (h^2), or the

proportion of phenotypic variance that can be attributed to additive genetic variation and can

respond to selection over generations (Alvesalo & Tigerstedt, 1974; Hlusko *et al.*, 2011). Second,

successive molars have higher levels of fluctuating asymmetry, or non-inherited phenotypic

variation usually attributed to developmental instability (Sofaer *et al.*, 1971; Alibert *et al.*, 1994).

In microevolutionary contexts, the higher variance has been attributed to longer exposure to or

greater phenotypically plastic sensitivity to environmental influences (Butler, 1983; Townsend *et*

al., 2009), or "diminished genetic control" (Harris, 2003). The emerging interpretation is that

greater phenotypic variation between individuals at the microevolutionary level either does not

indicate anything about relative evolutionary potential, because it is a signal of relative amounts

of non-inherited variation, or indicates less evolutionary potential, directly conflicting with the

macroevolutionary perspective of greater evolutionary potential.

Put another way, the similar patterns of phenotypic variation at both micro- and

macroevolutionary levels are attributed to two very different causes at each level, one inherited,

producing evolvable phenotypes, and one not. Similar findings and discrepancies in

interpretation have been found in vertebrate limbs, another segmented structure (Hallgrímsson *et*

al., 2002; Stepanova & Womack, 2020; Rothier *et al.*, 2023). The largely separated nature of

macroevolutionary and microevolutionary inquiry (Rolland *et al.*, 2023) may explain why this

discrepancy has not been noticed before, to our knowledge. It highlights the need for studies that

bridge the two scales.

To address this apparent conflict in interpretation, in this study we examine variation in

serially segmented structures at the within-species, microevolutionary level. Specifically, we use

two complementary approaches to test two complementary explanations for the

microevolutionary interpretation of reduced evolutionary potential in successive segments

(hereafter: Reduced Potential model) using the mammalian molar tooth module as a study

system. Both explanations focus on how decreasing heritability is partitioned among causes. The

first could be that reduced heritability could be caused by reduced levels of additive genetic

variation in successive molars, indicating a stronger discordance between microevolution and

macroevolution. The second explanation is that reduced heritability could be caused by increased

levels of environmentally induced variation, which may mask concordance between the two

levels of evolution.

Predictions for Evolvability

One interpretation of a Reduced Potential model is that later-forming structures carry less

additive genetic variation, the kind of variation that can respond to selection and evolve over

generations (Futuyma & Kirkpatrick, 2017). To address this interpretation, we conducted a

systematic review of patterns of heritability down the molar row, then used these results to

calculate evolvability. Heritability has long served as a comparative metric for such purposes, but

the evolvability (I_A) metric has emerged as a more appropriate, direct evaluation of evolutionary

potential (Houle, 1992; Hansen & Wagner, 2023). Both heritability and evolvability are

calculated from quantitative genetic studies partitioning phenotypic variation into additive

genetic (V_A) and other (V_E) components (note that V_E is sometimes called the environmental

component but does not refer solely to an ecological environment with which a species

interacts). Evolvability makes the additive genetic component comparable between traits and

samples by mean-scaling evolvable variance (V_A), using the same mean-scaling logic that

underlies coefficients of variation (CV) (Houle, 1992). In contrast, heritability conflates potential

for both an evolved response and a non-evolved, phenotypically plastic response by scaling

evolvable variance by total phenotypic variance, which includes both sources of variation

(Hansen *et al.*, 2011). Given the Reduced Potential model, our approach tests:

Prediction 1: We expect to observe decreased evolvability along the molar row, similar to

published patterns for heritability.

Predictions for Phenotypically Plastic Patterns

A complementary interpretation of the Reduced Potential model is that later-forming structures are more phenotypically variable because their formation is more strongly affected by their environment. To address this interpretation, we isolate phenotypically plastic variation and characterize its pattern along the molar row using a controlled feeding study of inbred lab rats (*Rattus norvegicus*). Phenotypically plastic variation occurs when different environments induce different phenotypes from an identical genotype (West-Eberhard, 1989). Environments may refer to climatic conditions, such temperature inducing trait change (Gupta & Lewontin, 1982), as well as other external conditions, such as nutrition availability (Sciulli *et al.*, 1979), as well as highly local conditions, such as slight differences in the developmental microenvironment surrounding symmetric structures, known as developmental instability (Leamy & Klingenberg, 2005). Resulting structures share genotypes and should be phenotypically perfectly symmetrical, but often carry a small amount of fluctuating asymmetry (Van Valen, 1962).

Increasing downstream phenotypic plasticity could refer to at least two different patterns. It is important to differentiate between them to better link patterns to processes. In this case, differentiation allows us to study types of developmental processes that form molar teeth. We assembled predictions using knowledge of molar development and its relationship to growth as well as the relationship between growth and nutrition, given our study system of nutrition's phenotypically plastic effect on molar sizes (Holloway *et al.*, 1961; Sciulli *et al.*, 1979; Blankenship-Sefczek *et al.*, 2024).

Specifically, trait means may not respond to the environment in the same way as trait variances (Fig. 1). An environmental condition that induces consistent changes in developmental pathways can cause a change in trait means in a sample. For example, reduced growth rates

could affect trait means, but would not necessarily affect variance (Miller & German, 1999;

Schwab & Moczek, 2016). The same mechanism could also affect variance if, for example, later-

forming segments are smaller or if a compounding effect reduces means, resulting in an increase

in mean-scaled measures of variance such as CV (Polly, 1998). We refer to a situation where a

single environmental change has an increasingly strong effect on downstream segments as one of

decreasing environmental canalization (Hallgrímsson *et al.*, 2002).

Developmental instability is considered as a separate phenomenon. If an environmental

condition increases developmental instability without inducing other changes, then the random

changes will increase the variance but should not affect the mean (Palmer & Strobeck, 1986;

Hallgrímsson *et al.*, 2002). In the case of poor nutrition, increased developmental instability

could be caused by decoherence, or a loss of co-regulation of traits driven by disruptions to gene

regulatory networks when homeostasis is lost (Lea *et al.*, 2019). Therefore, in this study we

assess our expectations through both means and variances.

Second, more specific to our study system, molar tooth width and length have potentially

different relationships to nutrition and its effect on development. Width, but not length, is

significantly genetically correlated with body size (Hlusko *et al.*, 2006). Molars also reach their

maximum length earlier in development, before they reach maximum width (Christensen *et al.*,

2023), similarly supporting the hypothesis that length and width are determined by different sets

of developmental pathways. If genetic correlations between body size and molar width, but not

molar length, occur through sharing pathways that affect growth in general, and if those

pathways are affected by nutrition (Miller & German, 1999; Lui & Baron, 2011), then we expect

that molar traits more strongly correlated with body size to also be more strongly affected by

nutrition than other molar traits. Therefore:

Prediction 2: We expect poor nutrition will have a stronger effect on molar width than molar length.

Third, developmental relationships between teeth may result in increasing phenotypic plasticity in downstream segments (Fig. 1). Segmented structures form from the iterative expression of similar or identical sets of developmental pathways (Thesleff & Sharpe, 1997; Young *et al.*, 2015). Importantly, early-forming structures have the capacity to interact with later-forming structures, producing downstream phenotypes that may differ between individuals only because of variation in phenotype of the initial segment, not because of any change in genetic basis between segments or because of different relationships of different segments to the environment (Kavanagh *et al.*, 2007). This cascading process could compound a phenotypically plastic effect in later-forming segments, producing a stronger effect and less environmental canalization in the phenotypes of later-forming segments (Hallgrímsson *et al.*, 2002).

Alternatively, effects may not compound. Although there is evidence of compounding in processes producing activation/inhibition ratios that permit enamel knot formation (Kavanagh *et al.*, 2007), that system does not require compounding in other processes like a general growth mechanism which might also affect molar sizes in conjunction with enamel knot dynamics. This hypothesis of multiple influences on size, some compounding and some not, is consistent with prior findings. Molar size phenotypes are incompletely described by a single model of compounding developmental process (Roseman & Delezene, 2019; Vitek *et al.*, 2020). In addition, molar positions have some level of genetic independence from one another, albeit often small, implying that the sets of developmental pathways that control their growth may be slightly different (Hlusko *et al.*, 2011; Hardin, 2020). These deviations from a cascading model fit an alternative proposal that a different effect of phenotypic plasticity on different molars is instead

related to different levels of exposure, or the amount of time each tooth spends developing,

implying that the molars are equally canalized against environmental effects per unit of exposure

time (Butler, 1983; Townsend *et al.*, 2009). Either model could explain previously observed

patterns of decreasing heritability and increasing phenotypic variance down the tooth row

(Kavanagh *et al.*, 2007; Riga *et al.*, 2014; Denes *et al.*, 2018), leading to:

Prediction 3: We expect the phenotypically plastic effect of any environmentally induced

change will increase along the tooth row from M_1 to M_3.

Overall, this approach of separating expectations for evolvable phenotypic variation and

plastic phenotypic variation may clarify potential connections between microevolutionary and

macroevolutionary patterns, resolving apparent disconnects in interpretation.

Methods

Evolvability

To test Prediction 1, we conducted a systematic literature review (Supplementary Methods

(Page *et al.*, 2021)). The objective of this systematic review was (A) to assess the consistency of

the evidence for the common perception that heritability decreases down the molar row, and (B)

to use reported parameter values to calculate evolvability and evaluate whether it has any

consistent pattern down the molar row. Trait heritability values and their standard errors were

collated for analysis. To meet minimum criteria for calculating evolvability, publications needed

to report the following for buccolingual width or mesiodistal length for at least two lower molar

positions: (1) trait means (m), (2) trait variances (V_P), (3) trait heritability ($h^2 = V_A/V_P$). From

these values, we could calculate evolvability, $I_A = V_A/m^2$, from heritability using the equation: I_A

$= h^2 * V_P /m^2$. Error in evolvability was estimated by resampling: a simulated sample of 1,000

additive variance estimates was calculated by sampling a normal distribution with the mean and

standard error of a given sample from the literature. The resulting simulated estimates were then

summarized as a standard deviation, which was plotted around each estimate. Given the range of

study designs, we did not conduct a meta-analysis of patterns across successive teeth. Instead, we

tabulated the number of datasets that reported higher, lower, or ambiguous levels of heritability

or evolvability in downstream molars. Some datasets are right and left sides from the same

individuals, which we visualized separately for clarity but for which we included only one side,

the left, in the tabulation.

Trait Plasticity

To test Predictions 2 and 3, we leveraged data collected from a previously conducted

controlled experiment originally intended to study the effect of maternal malnutrition on insulin

action and adipose tissue in male offspring of Wistar Han rats (*Rattus norvegicus*; Crl:WI[Han]

strain). This type of laboratory study used inbred lines to control for genetic variation between

individuals, allowing us to assume that all phenotypic variation was due to some component of

environment. Thus, by comparing phenotypic patterns between traits and between experimental

groups we could study the impact of environment on traits. In this case, the main environmental

variable that differed between control and experimental groups was the quality of the maternal

diet throughout gestation and suckling (8% low-protein experimental diet group vs. 20% protein

control diet group), and we measured its effect on offspring phenotype (Martin-Gronert *et al.*,

2016; Ward *et al.*, 2021). Nutritional variation is known to influence dental phenotypes,

providing a high likelihood of observing environmental impacts in our study (Holloway *et al.*,

1961; Miller & German, 1999; Calsa *et al.*, 2022). The time window of environmental

perturbation (Day 1 of gestation, E1, through weaning on postnatal day P21) is appropriate for

studying the effect of environment on rat molars because molar final size is determined during a

finite window of growth and then remains unchanged for the remainder of an individual's life

(Larson & Bader, 1976). In rats, which develop 1-2 days slower than mice, molars initiate

formation as the first molar (M_1) placode on embryonic day E13-14, then a tooth bud at E16-17

(Larson & Bader, 1976; Christensen *et al.*, 2023). The M_2 tooth bud appears at E17-19, and the

M_3 tooth bud is visible approximately 10 days later, at postnatal day P5-7 (Larson & Bader,

1976; Kavanagh *et al.*, 2007). Each molar reaches an inflection point and slows growth about 5

days after the bud stage, and achieves its final size about 8 days after bud stage (Christensen *et*

al., 2023), meaning that in the rat, final M_1 size is achieved by P1-2, M_2 size by P3-4, and M_3

size by P13-15. Therefore, we consider each molar tooth to be equally exposed to a consistent

environmental perturbation.

To collect phenotypic trait data, we used µCT scans of offspring sacrificed at age 3 months

(N = 25 control, 16 low-protein) to generate 3D models from which linear measures could be

collected. Further details of experimental design and µCT scan collection are reported in

(Martin-Gronert *et al.*, 2016; Ward *et al.*, 2021). From µCT image stacks, volumes of the left

molar row were segmented into 3D surfaces using automated thresholding in Avizo® version

2019.3 (FEI, Hilsboro, USA). Smoothing of the region of interest was conducted in each of the

three standard planes, but after a surface was extracted from that region we conducted no further

smoothing. From these surfaces, mesiodistal length and buccolingual width were measured in

triplicate by a single observer using MeshLab (Cignoni *et al.*, 2008). To evaluate whether

measurement error might contribute an undue amount of spurious noise to our estimates of

variation, as has been proposed to explain some patterns in variation across molars (Polly, 1998),

we calculated percent measurement error for each trait (Yezerinac *et al.*, 1992).

We calculated CV for each experimental group separately to ensure that potentially different

amounts of change in mean trait size did not spuriously influence estimates of variation. To

evaluate whether CVs differed between samples, we used a bootstrap resampling approach (N

replicates = 10,000) to test the hypothesis that one trait was more variable (larger CV) than

another trait compared to what would be expected by chance. We specifically tested hypotheses

that CV should increase between successive widths (one-tailed), between successive lengths

(one-tailed), that it should differ between length and width (two-tailed), and whether it differed

between control and experimental groups (two-tailed). Bootstrap resamples of low-protein CVs

were used to generate the null hypothesis distribution for between-trait comparisons.

To determine if it was necessary to test hypotheses of mean effect on any specific trait, we

first established whether the nutritional environment had a significant effect on trait means by

comparing experimental groups. For mean trait size, we used a t-test. To compare effect between

teeth, it was necessary to take the different sizes of each molar position into account ($M_1 > M_2 >$

M_3), because the same magnitude of difference has a different meaning for effect on the M_1 vs.

the M_3. We calculated a percent reduction statistic [mean control value – mean experimental

value] / [mean experimental value] to mean-scale the effect, similar to the mean scaling

performed for variation (CV) and evolvability (I_A) (Houle, 1992). To evaluate whether effect size

differences were significant, we used a bootstrapping resampling approach. To test Prediction 2,

we compared width to length values within each tooth. To test Prediction 3, we compared

successive length values to each other and successive width values to each other. When each

hypothesis was evaluated using multiple tests, we report p-values corrected using the Bonferroni

approach (Holm, 1979). All analyses and visualizations were conducted in R version 4.4.1 (R

Core Team, 2015) using packages `dplyr`, `reshape2`, `ggplot2`, `ggthemes`, and `patchwork`

(Wickham, 2007, 2009; Wickham *et al.*, 2023; Arnold, 2024; Pedersen, 2024).

Results

Evolvability

Eight reports of 536 items recovered in literature searches reported length or width

heritability for multiple molars, including two records found by searching the citations of six

studies recovered from database searches (Dempsey & Townsend, 2001; Hlusko *et al.*, 2011) (SI

Fig. 1). Seven samples from five of those eight reports met all three criteria for calculating

evolvability (Bader, 1965; Leamy & Bader, 1968; Leamy & Touchberry, 1974; Hlusko *et al.*,

2011; Hardin, 2019). Evolvability (I_A) patterns were generally ambiguous or increasing along the

molar row for both lengths and widths (Fig. 2). Heritability (h^2) patterns were generally

ambiguous or decreasing along the molar row (SI Table 1-2).

Trait Plasticity

Measurement error was low across traits (<2%, Table 1). All six traits were significantly

smaller in the low-protein offspring group compared to the control offspring group (Table 1).

Effect size, as measured by percent reduction from control to low-protein, was not consistent

down the toothrow (Table 2, Fig. 3A). Length of M_2 was significantly more strongly affected

than the length of the M_1. Widths of M_2 and M_3 were more strongly affected than width of the

M_1. No other comparison between upstream and downstream effects was significant. Widths

were generally more strongly affected than lengths, but only significantly so in the M_1 and M_3.

Variance, as measured by CV, did not differ significantly between the control and low-protein

group, with the single exception of M_3 length. CV significantly increased down the molar row

for lengths, but not for widths (Table 2, Fig. 3B). Widths were only more variable than lengths

for the M_1.

Discussion

Our results largely confirm prior patterns for statistics related to phenotypic variances

(Townsend *et al.*, 2009). Phenotypic variances are higher in the lengths of later-forming teeth.

However, a more systematic review and comparison of published heritability values does not

support a consistent link between higher downstream variances and lower downstream

heritability or evolvability. The largely ambiguous patterns for widths in both variance-based

measures appears to be more common than has been previously recognized. Our additional

results show that molar size heritability and phenotypic variance do not reliably predict molar

size evolvability, nor do they predict patterns of effect on trait means. In short, Prediction 1 was

not supported, and Prediction 3 was supported for measures of variance but not trait means. The

notably different patterns between lengths and widths generally supports Prediction 2, though

support is not universal. Our results are not attributable to measurement error, as has potentially

been an issue in other studies of variation (Polly, 1998).

Phenotypic Plasticity in Serially Segmented Structures

Multiple processes are potentially mixed together in the characterization of later-forming

molars as more sensitive to environmental influence. The discordant patterns of phenotypic

plasticity in trait means vs. trait variances allows us to distinguish between two different types of

environmentally related processes and their potentially different relationship to serial segments.

This finding is robust to the limitations of our study. We acknowledge that because the

experimental design did not include observations of how long each molar took to form, we

cannot test the alternative hypothesis of environmental canalization scaled by exposure time.

With a single sample of one species included in our study of plasticity, it is also possible that we

did not capture a stronger effect in later-forming molar traits that may be present in other species

or genotypes (Holloway *et al.*, 1961; Tonge & McCance, 1973). However, none of these limits to

generalizability contradict the point that the phenotypically plastic responses of trait means and

variances are decoupled in later-forming molars. Different developmental mechanisms either

respond differently to the environment or respond to different components of the environment.

Instead of a single mechanism underlying a Reduced Potential model and impacting both

trait means and variances, we hypothesize that at least two separate types of environmentally

sensitive processes impact molar size traits. The first process we hypothesize is an effect of

nutritional environment on general organismal growth, which affects molars to the degree that

they are environmentally canalized, or developmentally independent of overall organismal

growth. The developmental non-independence from general growth across molars need not

compound down the molar row, allowing for nonlinear patterns like those we observed in our

study, especially for width measurements (Fig. 3A, Table 1), with no requirement for a shared

pattern across structures like teeth, limbs, and vertebral segments with very different

developmental underpinnings.

The second process we hypothesize is a model of developmental instability compounding

down the development of the molar row, explaining patterns of phenotypic variance, heritability,

and fluctuating asymmetry documented in this and other studies (Sofaer *et al.*, 1971; Gingerich,

1974; Hlusko *et al.*, 2011). Our data collection protocol did not permit us to evaluate fluctuating

asymmetry, a more standard statistic for inferring developmental instability (Palmer & Strobeck,

1986; Hallgrímsson *et al.*, 2002), but prior studies have found a consistent pattern of downstream

increase, consistent with our explanation (Sofaer *et al.*, 1971; Alibert *et al.*, 1994). The other

phenomenon that would produce such an effect, antisymmetry, is not a feature of mammalian

tooth sizes (Van Valen, 1962; Leamy *et al.*, 2005).

Unlike environmental canalization, developmental instability might compound more

generally across serial structures if it were caused by iterative expression of a gene regulatory

network. Notably, molar length fits expected patterns of this model (Table 2), given that length is

more conceptually closely tied to a model of compounding activation/inhibition dynamics than

molar width (Kavanagh *et al.*, 2007). The lack of significant difference in CV between the two

experimental groups (Table 1) suggests that this compounding instability is not related to

nutrition, but instead may be a more general feature of certain developmental mechanisms

(Hallgrímsson *et al.*, 2002). When successive segments interact with each other through

expression of one gene regulatory network, any stochastic developmental instability in the

network affecting the M_1 would add on to the stochasticity affecting the M_2 itself, and so on

down the tooth row. No tooth would have developmental instability less than zero, and therefore

developmental instability would only ever remain stable or increase down a set of segments,

resulting in higher downstream phenotypic variance if all else were equal among molars. The

nature of developmental instability would mean that this random process averages out with no

accumulating downstream effect on trait means in a sample, even though the effect accumulates

downstream in any given individual.

In sum, our results contradict the commonly held hypothesis that increasing developmental

instability is mechanistically related to increasing sensitivity to environment and decreasing

environmental canalization (Hallgrímsson *et al.*, 2002). Although compounding developmental

instability can potentially explain patterns of variance-related statistics, it cannot explain either

the patterns of environmental canalization or of evolvability. If this model holds, then any set of

serial segments that share a single gene regulatory network may show this pattern of increasing

variance, not necessarily tethered to increasing sensitivity to the environment.

Plasticity & Organismal Growth Reflected in Teeth

Prior work found stronger links between molar width and overall body size than molar length

in terms of genetic correlations (Hlusko *et al.*, 2006). Further work built on these links to focus

on studying more tooth-specific evolutionary patterns, rather than patterns that might reflect

more mixed evolutionary signals of body size and tooth size evolution (Hlusko *et al.*, 2016). In

our study, all molar size traits were significantly, phenotypically plastically reduced by poor

nutrition, highlighting that no trait is fully buffered from non-evolved, size-related responses to

environment, consistent with heritability estimates (Hlusko *et al.*, 2011). However, the stronger

impact on widths than lengths supports the interpretation that focusing on lengths better isolates

the evolution of teeth separate from evolution of other organismal properties (Hlusko *et al.*,

2016).

The stronger effect of poor nutrition on molar width than length may reflect how overall

organismal growth pathways are specifically expressed during tooth growth. This hypothesis

would make our phenotypic patterns a potential phenocopy (sensu (Hallgrímsson *et al.*, 2002)) of

evolved changes whose underlying mutations affected the same developmental pathways. In the

case of teeth and growth, insulin-like growth factor (IGF) signaling and expression is a

prominent candidate for future study. Insulin-like growth factors are ubiquitous during

embryonic development and regulation of IGF is critical for cellular proliferation of developing

tissue and organs, including dentition (Brown *et al.*, 2017; Oyanagi *et al.*, 2019; Christensen *et*

al., 2023). Poor nutrition can impact IGF regulation by reducing the rate of protein

phosphorylation, resulting in a reduction of available IGF binding proteins and potentially

resulting in a negative feedback loop (Tagliabracci *et al.*, 2015; Chrudinová *et al.*, 2024).

Reduced IGF results in systematically smaller sizes for developing embryos without impacting

the fundamental timing or process of development due to the role of IGF in proper organogenesis

(Estrella *et al.*, 2024). By reducing phosphorylation, poor nutrition could directly influence

developing tooth size without otherwise inducing significant alteration of gene expression in

other developmental pathways, similar to its role in genetically-based patterning of teeth

(Christensen *et al.*, 2023).

Reconciling Macro- and Microevolutionary Interpretations

Overall, we found equal or increasing evolutionary potential down the molar row in

microevolutionary contexts. This interpretation reconciles microevolutionary and

macroevolutionary interpretations of molars and segment patterns, with implications for other

serially segmented structures (Mongle *et al.*, 2022). Later-forming segments may be less

heritable and statistically noisier because of their inflated variance, but their mean values are not

necessarily poorer reflections of evolutionary history than the mean values of earlier-forming

segments. To the contrary, none of the samples had lower evolvabilities in the M_3 than in

upstream molars. Instead, the M_3 either had equal evolvability, suggesting a role for relaxed

selection on the M_3 compared to upstream molars (Gingerich & Winkler, 1979; Mongle *et al.*,

2022), or had increased evolvability, suggesting that there is also comparatively more additive

genetic variation for selection to act on in these teeth (Fig. 2). This perspective may explain why

the M_3 has been considered so taxonomically useful in certain rodents (Barnosky, 1993; Polly,

2003), despite their general characterization as a noisy, non-ideal choice for taxonomic study

(Gingerich, 1974; Vezzosi & Kerber, 2018). Mammalian molars appear to be another case where

heritability is not as informative of evolutionary potential as evolvability (Hansen *et al.*, 2011).

In limbs, the macroevolutionary pattern of higher evolutionary rates in later-forming

segments is attributed to function, where more distal, downstream segments have more direct

interaction with substrates and other sources of selective pressure (Rothier *et al.*, 2023). This

selection pressure may make their potential more easily observed, in contrast to the molar

module where function is often conserved down the tooth row and treated as a single functional

complex along with other cheek teeth (Ungar, 2010). Investigating the macroevolution of taxa

that differ in their degree of functional variation down the molar row may provide an important

comparison to patterns in the limb, filling in a remaining gap in our understanding of how

microevolutionary and macroevolutionary diversification align.

Acknowledgments

Thanks to M.T. Silcox for connecting the co-authors of this study, C.J. Percival for conversations

that helped structure the discussion. Thanks to funding from the Henry Sidgwick Research

Fellowship from Newnham College, Cambridge, University of Toronto PhD Pilot Research

Grant, and startup funding from Stony Brook University. Author RWB was funded via an

Institutional Research and Academic Career Development Award (IRACDA) made to Stony

Brook University from the National Institute of General Medical Sciences of the National

Institutes of Health [K12GM102778]. The content is solely the responsibility of the authors and

does not necessarily represent official views of the National Institutes of Health.

Funding Sources:

Henry Sidgwick Research Fellowship from Newnham College, Cambridge

University of Toronto PhD Pilot Research Grant

Startup funding from Stony Brook University

Data accessibility statement: Data and code sufficient to reproduce results in this study are

contained in a Dryad publicly accessible digital repository [*private reviewer link: XXXXXX*]

Author contributions: NSV conducted conceptualization, scripting, validation, formal analysis,

and visualization components of the study. NSV, EP, MM-G, and SEO contributed to

methodology. NSV, ES, AD contributed to phenotypic data collection. DW, EP, MM-G, and SEO

contributed resources. NSV and DW contributed to data curation. NSV and SEO contributed

supervision and project administration. EP and SEO contributed funding acquisition for the

nutrition study. NSV and RWB wrote the original draft, and all authors contributed to review and

editing.

Conflicts of interest statement: The authors declare no conflicts of interest.

Citations

Alibert, P., Renaud, S., Dod, B., Bonhomme, F. & Auffray, J.-C. (1994) Fluctuating asymmetry in the *Mus*

musculus hybrid zone: a heterotic effect in disrupted co-adapted genomes. *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, **258**, 53–59.

Alvesalo, L. & Tigerstedt, P.M.A. (1974) Heritabilities of human tooth dimensions. *Hereditas*, **77**, 311–

318.

Amaral, E.D.O.S., Jara-Espejo, M. & Line, S.R.P. (2024) The minimum free energy of G-quadruplex-forming

sequences of the MSX1 intron is associated with relative molar and premolar size in eutherians. *Journal*

of Mammalian Evolution, **31**, 22.

Arnold, J.B. (2024) *ggthemes: Extra Themes, Scales and Geoms for "ggplot2."*

Bader, R.S. (1965) Heritability of dental characters in the house mouse. *Evolution*, **19**, 378–384.

Bader, R.S. & Lehmann, W.H. (1965) Phenotypic and genotypic variation in odontometric traits of the

house mouse. *American Midland Naturalist*, **74**, 28–38.

Barnosky, A.D. (1987) Punctuated equilibrium and phyletic gradualism: some facts from the Quaternary

mammalian record. In *Current Mammalogy* (ed. by Genoways, H.H.). Springer US, Boston, MA, pp. 109–

147.

Barnosky, A.D. (1993) Mosaic evolution at the population level in *Microtus pennsylvanicus*. In

Morphological Change in Quaternay Mammals of North America (ed. by Martin, R.A. & Barnosky, A.D.).

Cambridge University Press, Cambridge, U.K., pp. 24–59.

Billet, G. & Bardin, J. (2021) Segmental Series and Size: Clade-Wide Investigation of Molar Proportions

Reveals a Major Evolutionary Allometry in the Dentition of Placental Mammals. *Systematic Biology*, **70**,

1101–1109.

Blankenship-Sefczek, E.C., Goodman, A.H., Hubbe, M., Hunter, J.P. & Guatelli-Steinberg, D. (2024)

Nutritional supplementation, tooth crown size, and trait expression in individuals from Tezonteopan,

Mexico. *PLOS ONE*, **19**, e0305123.

Böhmer, C., Rauhut, O.W.M. & Wörheide, G. (2015) Correlation between *Hox* code and vertebral

morphology in archosaurs. *Proceedings of the Royal Society B: Biological Sciences*, **282**, 20150077.

Brown, S.M., Peters, R. & Lawrence, A.B. (2017) Up-regulation of IGF-1 in the frontal cortex of piglets

exposed to an environmentally enriched arena. *Physiology & Behavior*, **173**, 285–292.

Butler, P.M. (1983) Evolution and mammalian dental morphology. *Journal De Biologie Buccale*, **11**, 285–

302.

Calsa, B., Bortolança, T.J., Masiero, B.C., Esquisatto, M.A.M., De Oliveira, C.A., Catisti, R., *et al.* (2022)

Maxillary and dental development in the offspring of protein-restricted female rats. *European Journal of*

Oral Sciences, **130**, e12895.

Christensen, M.M., Hallikas, O., Das Roy, R., Väänänen, V., Stenberg, O.E., Häkkinen, T.J., *et al.* (2023) The

developmental basis for scaling of mammalian tooth size. *Proceedings of the National Academy of*

Sciences, **120**, e2300374120.

Chrudinová, M., Kirk, N.S., Chuard, A., Venugopal, H., Zhang, F., Lubos, M., *et al.* (2024) A viral insulin-like

peptide inhibits IGF-1 receptor phosphorylation and regulates IGF1R gene expression. *Molecular*

Metabolism, **80**, 101863.

Cignoni, P., Callieri, M., Corsini, M., Dellepiane, M., Ganovelli, F. & Ranzuglia, G. (2008) MeshLab: an

Open-Source Mesh Processing Tool. In *Eurographics Italian Chapter Conference* (ed. by Scarano, V.,

Chiara, R.D. & Erra, U.). The Eurographics Association, pp. 129–136.

Cuozzo, F.P. (2008) Using extant patterns of dental variation to identify species in the primate fossil

record: a case study of middle Eocene *Omomys* from the Bridger Basin, southwestern Wyoming.

Primates, **49**, 101–115.

Dempsey, P.J. & Townsend, G.C. (2001) Genetic and environmental contributions to variation in human

tooth size. *Heredity*, **86**, 685–693.

Denes, B.J., Lagou, A., Dorotheou, D. & Kiliaridis, S. (2018) A longitudinal study on timing and velocity of

rat molar eruption: Timing of rat molar eruption. *Laboratory Animals*, **52**, 394–401.

Emery-Wetherell, M. & Davis, E. (2018) Dental measurements do not diagnose modern artiodactyl

species: Implications for the systematics of Merycoidodontoidea. *Palaeontologia Electronica*.

Estrella, C.A.S., Gatford, K.L., Xiang, R., Javadmanesh, A., Ghanipoor-Samami, M., Nattrass, G.S., *et al.*

(2024) Asymmetric growth-limiting development of the female conceptus. *Frontiers in Endocrinology*, **14**,

1306513.

Futuyma, D.J. & Kirkpatrick, M. (2017) *Evolution*. Fourth edition. Sinauer Associates, Inc., Publishers,

Sunderland, Massachusetts.

Gingerich, P.D. (1974) Size variability of the teeth in living mammals and the diagnosis of closely related

fossil species. *Journal of Paleontology*, **48**, 895–903.

Gingerich, P.D. & Winkler, D.A. (1979) Patterns of variation and correlation in the dentition of the red fox,

Vulpes vulpes. *Journal of Mammalogy*, **60**, 691–704.

Gupta, A.P. & Lewontin, R.C. (1982) A study of reaction norms in natural populations of *Drosophila*

pseudoobscura. *Evolution*, **36**, 934–948.

Hallgrímsson, B., Willmore, K. & Hall, B.K. (2002) Canalization, developmental stability, and

morphological integration in primate limbs. *American Journal of Physical Anthropology*, **119**, 131–158.

Hansen, T.F., Pélabon, C. & Houle, D. (2011) Heritability is not evolvability. *Evolutionary Biology*, **38**, 258–

277.

Hansen, T.F. & Wagner, G.P. (2023) The Evolution of Evolvability. In *Evolvability: A Unifying Concept in*

Evolutionary Biology? (ed. by Hansen, T.F., Houle, D., Pavličev, M. & Pélabon, C.). The MIT Press.

Hardin, A.M. (2019) Genetic contributions to dental dimensions in brown-mantled tamarins (*SAGUINUS*

FUSCICOLLIS) and rhesus macaques (*MACACA MULATTA*). *American Journal of Physical Anthropology*, **168**,

292–302.

Hardin, A.M. (2020) Genetic correlations in the rhesus macaque dentition. *Journal of Human Evolution*,

148, 102873.

Harris, E.F. (2003) Where's the Variation? Variance Components in Tooth Sizes of the Permanent

Dentition. *Dental Anthropology Journal*, **16**, 84–94.

Hershkovitz, P. (1967) Dynamics of Rodent Molar Evolution: A Study Based on New World Cricetinae,

Family Muridae. *Journal of Dental Research*, **46**, 829–842.

Hlusko, L.J., Lease, L.R. & Mahaney, M.C. (2006) Evolution of genetically correlated traits: Tooth size and

body size in baboons. *American Journal of Physical Anthropology*, **131**, 420–427.

Hlusko, L.J., Sage, R.D. & Mahaney, M.C. (2011) Modularity in the mammalian dentition: mice and

monkeys share a common dental genetic architecture. *Journal of Experimental Zoology Part B: Molecular*

and Developmental Evolution, **316B**, 21–49.

Hlusko, L.J., Schmitt, C.A., Monson, T.A., Brasil, M.F. & Mahaney, M.C. (2016) The integration of

quantitative genetics, paleontology, and neontology reveals genetic underpinnings of primate dental

evolution. *Proceedings of the National Academy of Sciences*, 201605901.

Holloway, P.J., Shaw, J.H. & Sweeney, E.A. (1961) Effects of various sucrose:casein ratios in purified diets

on the teeth and supporting structures of rats. *Archives of Oral Biology*, **3**, 185–200.

Holm, S. (1979) A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics*,

6, 65–70.

Houle, D. (1992) Comparing evolvability and variability of quantitative traits. *Genetics*, **130**, 195–204.

Kavanagh, K.D., Evans, A.R. & Jernvall, J. (2007) Predicting evolutionary patterns of mammalian teeth

from development. *Nature*, **449**, 427–432.

Larson, M.J. & Bader, R.S. (1976) Maternal Effects in Odontometric Traits of House Mice. *American*

Midland Naturalist, **96**, 21.

Lea, A., Subramaniam, M., Ko, A., Lehtimäki, T., Raitoharju, E., Kähönen, M., *et al.* (2019) Genetic and

environmental perturbations lead to regulatory decoherence. *eLife*, **8**, e40538.

Leamy, L.J. & Bader, R.S. (1968) Components of Variance of Odontometric Traits in a Wild-Derived

Population of Peromyscus leucopus. *Evolution*, **22**, 826.

Leamy, L.J. & Klingenberg, C.P. (2005) The genetics and evolution of fluctuating asymmetry. *Annual*

Review of Ecology, Evolution, and Systematics, **36**, 1–21.

Leamy, L.J. & Touchberry, R.W. (1974) Additive and non-additive genetic variance in odontometric traits

in crosses of seven inbred lines of house mice. *Genetical Research*, **23**, 207–217.

Leamy, L.J., Workman, M.S., Routman, E.J. & Cheverud, J.M. (2005) An epistatic genetic basis for

fluctuating asymmetry of tooth size and shape in mice. *Heredity*, **94**, 316–325.

Lui, J.C. & Baron, J. (2011) Mechanisms Limiting Body Growth in Mammals. *Endocrine Reviews*, **32**, 422–

440.

Martin-Gronert, M.S., Fernandez-Twinn, D.S., Bushell, M., Siddle, K. & Ozanne, S.E. (2016) Cell-

autonomous programming of rat adipose tissue insulin signalling proteins by maternal nutrition.

Diabetologia, **59**, 1266–1275.

Miller, J.P. & German, R.Z. (1999) Protein Malnutrition Affects the Growth Trajectories of the Craniofacial

Skeleton in Rats. *The Journal of Nutrition*, **129**, 2061–2069.

Mongle, C.S., Nesbitt, A., Machado, F.A., Smaers, J.B., Turner, A.H., Grine, F.E., *et al.* (2022) A common

mechanism drives the alignment between the micro- and macroevolution of primate molars. *Evolution*,

evo.14600.

Oyanagi, T., Takeshita, N., Hara, M., Ikeda, E., Chida, T., Seki, D., *et al.* (2019) Insulin-like growth factor 1

modulates bioengineered tooth morphogenesis. *Scientific Reports*, **9**, 368.

Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., *et al.* (2021) The

PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Systematic Reviews*, **10**,

89.

Palmer, A.R. & Strobeck, C. (1986) Fluctuating asymmetry: Measurement, Analysis, Patterns. *Annual*

Review of Ecology and Systematics, **17**, 391–421.

Pedersen, T.L. (2024) *patchwork: The Composer of Plots*.

Polly, P.D. (1998) Variability in mammalian dentitions: size-related bias in the coefficient of variation.

Biological Journal of the Linnean Society, **64**, 83–99.

Polly, P.D. (2003) Paleophylogeography: the tempo of geographic differentiation in marmots (*Marmota*).

Journal of Mammalogy, **84**, 369–284.

R Core Team. (2015) R: A Language and Environment for Statistical Computing.

Riga, A., Belcastro, M.G. & Moggi-Cecchi, J. (2014) Environmental stress increases variability in the

expression of dental cusps: Environmental Influence on Dental Morphology. *American Journal of Physical*

Anthropology, **153**, 397–407.

Rolland, J., Henao-Diaz, L.F., Doebeli, M., Germain, R., Harmon, L.J., Knowles, L.L., *et al.* (2023)

Conceptual and empirical bridges between micro- and macroevolution. *Nature Ecology & Evolution*, **7**,

1181–1193.

Roseman, C.C. & Delezene, L.K. (2019) The inhibitory cascade model is not a good predictor of molar size

covariation. *Evolutionary Biology*, **46**, 229–238.

Rothier, P.S., Fabre, A.-C., Clavel, J., Benson, R.B. & Herrel, A. (2023) Mammalian forelimb evolution is

driven by uneven proximal-to-distal morphological diversity. *eLife*, **12**, e81492.

Schuman, E.L. & Brace, C.L. (1954) Metric and morphologic variations in the dentition of the Liberian

chimpanzee; comparisons with anthropoid and human dentition. *Huam Biology*, **26**, 239–268.

Schwab, D.B. & Moczek, A.P. (2016) Nutrient Stress During Ontogeny Alters Patterns of Resource

Allocation in two Species of Horned Beetles. *Journal of Experimental Zoology Part A: Ecological Genetics*

and Physiology, **325**, 481–490.

Sciulli, P.W., Doyle, W.J., Kelley, C., Siegel, P. & Siegel, M.I. (1979) The interaction of stressors in the

induction of increased levels of fluctuating asymmetry in the laboratory rat. *American Journal of Physical*

Anthropology, **50**, 279–284.

Sofaer, J.A., Bailit, H.L. & MacLean, C.J. (1971) A developmental basis for differential tooth reduction

during hominid evolution. *Evolution*, **25**, 509–517.

Stepanova, N. & Womack, M.C. (2020) Anuran limbs reflect microhabitat and distal, later-developing

bones are more evolutionarily labile. *Evolution*, **74**, 2005–2019.

Tagliabracci, V.S., Wiley, S.E., Guo, X., Kinch, L.N., Durrant, E., Wen, J., *et al.* (2015) A Single Kinase

Generates the Majority of the Secreted Phosphoproteome. *Cell*, **161**, 1619–1632.

Thesleff, I. & Sharpe, P. (1997) Signalling networks regulating dental development. *Mechanisms of*

development, **67**, 111–123.

Tonge, C.H. & McCance, R.A. (1973) Normal development of the jaws and teeth in pigs, and the dela and

malocclusion produced by calorie deficiencies. *Journal of Anatomy*, **115**, 1–22.

Townsend, G., Harris, E.F., Lesot, H., Clauss, F. & Brook, A. (2009) Morphogenetic fields within the human

dentition: A new, clinically relevant synthesis of an old concept. *Archives of Oral Biology*, **54**, S34–S44.

Ungar, P.S. (2010) *Mammal teeth: origin, evolution, and diversity*. Johns Hopkins University Press,

Baltimore.

Van der Made, J. (1996) Listriodontinae (Suidae, Mammalia), their evolution, systematics and

distribution in time and space. *Contributions to Tertiary and Quaternary Geology*, **33**, 3–254.

Van Valen, L. (1962) A study of fluctuating asymmetry. *Evolution*, **16**, 125–142.

Vezzosi, R.I. & Kerber, L. (2018) The southernmost record of a large erethizontid rodent (Hystricomorpha:

Erethizontoidea) in the Pleistocene of South America: Biogeographic and paleoenvironmental

implications. *Journal of South American Earth Sciences*, **82**, 76–90.

Vitek, N.S., McDaniel, S.F. & Bloch, J.I. (2020) Is variation in molar tooth crown morphology of the

Grasshopper Mouse (*Onychomys leucogaster*) a reflection of selection or drift? In *Society of Integrative*

and Comparative Biology Annual Meeting. Presented at the Society of Integrative and Comparative

Biology Annual Meeting, Austin, TX.

Ward, D.L., Schroeder, L., Pomeroy, E., Roy, J.E., Buck, L.T., Stock, J.T., *et al.* (2021) Early life malnutrition

and fluctuating asymmetry in the rat bony labyrinth. *The Anatomical Record*, ar.24601.

West-Eberhard, M.J. (1989) Phenotypic plasticity and the origins of diversity. *Annual review of Ecology*

and Systematics, 249–278.

Wickham, H. (2007) Reshaping Data with the reshape Package. *Journal of Statistical Software*, **21**, 1–20.

Wickham, H. (2009) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. (2023) *dplyr: A Grammar of Data*

Manipulation.

Yezerinac, S.M., Lougheed, S.C. & Handford, P. (1992) Measurement error and morphometric studies:

statistical power and observer experience. *Systematic Biology*, **41**, 471–482.

Young, N.M., Winslow, B., Takkellapati, S. & Kavanagh, K. (2015) Shared rules of development predict

patterns of evolution in vertebrate segmentation. *Nature Communications*, **6**, 6690.

Figures and Tables with Captions

Table 1. Summary statistics for molar size traits.

Trait	Tooth	PME	C vs. LP x̄ p-value	LP vs. C % Reduced	CV C	CV LP	C vs. LP CV p-value
Length	M_1	0.58	**$< 1 * 10^{-5}$**	5.53	2.76	3.04	1
	M_2	0.34	**$< 1 * 10^{-5}$**	8.81	3.88	3.39	1
	M_3	0.36	**$< 1 * 10^{-5}$**	6.67	4.75	5.88	**0.0216**
Width	M_1	1.29	**$< 1 * 10^{-5}$**	9.56	3.3	3.84	0.4008
	M_2	0.89	**$< 1 * 10^{-5}$**	7.4	3.58	4.32	0.6102
	M_3	0.38	**$< 1 * 10^{-5}$**	11.74	4.83	5.25	1

Significance values reported post-Bonferroni correction. Significant p-values are in bold.

Abbreviations: C, control group; CV, coefficient of variation; LP, low protein group; p, p-value;

PME, percent measurement error; x̄, mean trait value

Table 2. P-values assessing significance of differences between pairs of traits. Low-protein CVs were used for between-trait comparisons.

	Length vs. Width			Length		Width	
	% Reduced	CV		% Reduced	CV	% Reduced	CV
M_1	< 0.0003	0.0051	M_1 vs. M_2	0.0003	0.036	1	1
M_2	1	1	M_2 vs. M_3	1	0.0021	0.0003	0.0003
M_3	< 0.0003	1	M_1 vs. M_3	1	0.0003	0.0003	0.1437

Significance values reported post-Bonferroni correction. Abbreviations: CV, coefficient of variation.

Figure 1. Conceptual diagram of two different ways in which a low-protein diet, a kind of

environmental perturbation, could generate a pattern of increasing coefficients of variation (CV)

in downstream tooth sizes, matching previously published patterns of variation in tooth size. A.

Control group, in which each tooth sample has a mean size (black solid outline) and variance

(gray dashed ellipse). B., Low Protein group, in which all tooth sizes are reduced, but either (left)

downstream teeth are more strongly reduced in size than upstream teeth, resulting in an increase

in CV because of a more strongly reduced denominator in the CV equation, or (right)

downstream teeth have more strongly increased variance than upstream teeth, resulting in an

increase in CV because of a more strongly increased numerator in the same equation.

Abbreviations: CV, coefficient of variation; s, standard deviation; $\bar{x}$, sample mean.

Figure 2. Evolvability (I_A) and heritability (h^2) calculated from a systematic review of published samples. Lines connect traits measured from the same sample. Top panel shows values from studies that did not report enough information to calculate evolvability. Middle panel shows samples with paired evolvability estimates (bottom panel). Bars indicate standard error, jittered to better show overlap. Note that the y-axis is on a logarithmic scale to better display the range of evolvability estimates.

Figure 3. Effect of low-protein nutrition on the mean and variance of molar size traits,

standardized by mean trait size. Box and whisker plots show median (thick bar) and mean

(triangle). Corresponding density plots are to the right. Note that density plots height may not

match box plot median because of the different way each approach summarizes the distribution.

A. Percent that low-protein group is reduced compared to control group. B. Coefficient of

variation. Boxes with upward-facing triangles show the control group. Boxes with downward-

facing triangles show the low-protein group. The pair of density plots show control (left) and

low-protein (right) groups. Abbreviations: CV, coefficient of variation; L, length; W, width.

Supplementary Figure 1. PRISMA 2020 flow diagram of report retrieval.

Supplementary Table 1. Characteristics of studies recovered from the systematic literature

search.

Supplementary Table 2. Results of each sign test for consistent downstream decreases in

heritability (h^2) and evolvability (I_A) in the molar tooth row of previously published samples.