Developmentally cascading structures do not lose evolutionary potential, but compound

developmental instability in rat molars

Natasha S. Vitek[1], Ella Saks[1], Amy Dong[1], Robert W. Burroughs [1], Devin L Ward[2], Emma

Pomeroy[3], Malgorzata Martin-Gronert[4], Susan E. Ozanne[4]

[1] Department of Ecology & Evolution, Stony Brook University, Stony Brook, NY.

[2] Department of Anthropology, University of Toronto, Toronto, Canada

[3] Department of Archaeology and Newnham College, University of Cambridge, Cambridge, UK

[4] Metabolic Research Laboratories and MRC Metabolic Diseases Unit, Institute of Metabolic

Science, University of Cambridge, Cambridge, UK

ORCID:

NSV: 0000-0003-0587-8907

RWB: 0000-0003-4384-3430

DLW: 0000-0002-2343-8238

EP: 0000-0001-6251-2165

MM-G: 0000-0002-6763-8215

SEO: 0000-0001-8753-5144

Keywords: evolvability, phenotypic plasticity, nutrition, growth, molar, Mammalia

Abstract

Increasing variability down serially segmented structures, such as mammalian molar teeth and

vertebrate limb segments, is a much-replicated pattern. The same phenotypic pattern has

conflicting interpretations at different evolutionary scales. Macroevolutionary patterns are

thought to reflect greater evolutionary potential in later-forming segments, but microevolutionary

patterns are thought to reflect less evolutionary potential and greater phenotypic plasticity. We

address this conflict by recalculating evolutionary potential (evolvability) from a systematic

review of published mammalian molar sizes, then directly measure phenotypic plasticity from a

controlled feeding experiment. Effects on lengths and widths are discordant in a way that

suggests general growth pathways have a role in phenotypically plastic dental responses to

nutrition. Effects on successive trait means do not necessarily increase downstream, contrary to

long-standing hypotheses. We confirm prior findings of increasing non-inherited variance

downstream, showing decoupling between effects on trait mean and variance. These patterns can

be explained by a cascading model of tooth development compounding the effect of

developmental instability as an influence separate from general environmental effects on the

developing embryo. When evaluated in terms of evolvability, later-developing molars are equally

or more evolvable than earlier-developing molars, aligning their microevolutionary potential

with macroevolutionary patterns in other serially segmented structures.

Introduction

Animal bodies contain numerous serially segmented structures, such as vertebrate teeth, limb

bones, and vertebrae (Young et al. 2015). The semi-independent nature of serial structures makes

them important systems for understanding how diverse phenotypes evolve in structures that share a significant proportion of their genetic and developmental basis (Kavanagh et al. 2007; Böhmer et al. 2015).

In mammalian teeth, some of the most studied vertebrate serial structures, conflicting perspectives from different evolutionary scales emerge on the relative evolutionary potential of different segments in a module (Fig. 1). In macroevolution, where phenotype is largely understood to have a genetic basis, later-developing segments such as the third molars (M_3) are considered to have greater evolutionary potential than early-developing, upstream segments such as the first or second molars (M_1, M_2; Fig. 1B). Support comes from observations of faster evolutionary rates downstream ($M_3 > M_2 > M_1$) sometimes documented directly (Sofaer et al. 1971; Mongle et al. 2022; Amaral et al. 2024), sometimes implied by greater taxonomic utility of the M_3 as a more sensitive indicator of species differences, especially in rodents and suoids (Hershkovitz 1967; Barnosky 1987; Van der Made 1996; Polly 2003; Willis and Swindler 2004). These observations translate into higher macroevolutionary diversity, or variance, in M_3s vs. earlier-developing segments (Billet and Bardin 2021).

Figure 1. Conceptual diagram of the conflicting interpretations of patterns of variation down

mammalian molars, an example of a segmented structure. The conflict is highlighted in the

dashed box. Increases or decreases in gray indicate observed patterns. Increases or decreases in

black indicate interpretations of those patterns. A, line drawing of the three lower molars of

Rattus norvegicus, which develop in sequential order from top to bottom. Both evolutionary

scales share a general pattern of increasing phenotypic variation in downstream teeth. B,

macroevolutionary perspective on increasing phenotypic variation. C, microevolutionary

perspective. Abbreviations: FA, fluctuating asymmetry; h^2, heritability.

In microevolutionary contexts, where phenotype likely has both inherited genetic basis and

non-inherited plastic basis, molars usually share the same predicted pattern of higher phenotypic

variation in later-forming segments (Fig. 1A ex., Schuman and Brace 1954; Bader and Lehmann

1965; Gingerich 1974; Cuozzo 2008 but see Emery-Wetherell and Davis 2018 for a counter-

example). Along the same sequence of segments, two other patterns emerge. First, successive

molars generally (though not exclusively) have lower levels of heritability (h^2), or the proportion

of phenotypic variance that can be attributed to additive genetic variation and can respond to

selection over generations (Fig. 1C; Alvesalo and Tigerstedt 1974; Hlusko et al. 2011). Second,

successive molars have higher levels of fluctuating asymmetry, or non-inherited phenotypic

variation usually attributed to developmental instability (Fig. 1C; Sofaer et al. 1971; Alibert et al.

1994). In microevolutionary contexts, the higher variance has been attributed to longer exposure

to or greater phenotypically plastic sensitivity to environmental influences (Butler 1983;

Townsend et al. 2009), or "diminished genetic control" (Harris 2003, p. 91). The emerging

interpretation is that greater phenotypic variation between individuals at the microevolutionary

level does not indicate anything about relative evolutionary potential, because it is a signal of

relative amounts of non-inherited variation, or indicates less evolutionary potential, directly

conflicting with the macroevolutionary perspective of greater evolutionary potential (Soulé 1982;

Kieser and Groeneveld 1987).

Put another way, the similar patterns of phenotypic variation at both micro- and

macroevolutionary levels are attributed to two very different causes at each level, one inherited,

producing evolvable phenotypes, and one not (Fig. 1). This particular discrepancy between

evolutionary scales has not been noticed before, to our knowledge. Similar patterns have been

found in vertebrate limbs, another segmented structure where phenotypic variation increases

downstream both within and between species, as does fluctuating asymmetry within species, but

heritability decreases (Hallgrímsson et al. 2002; Stepanova and Womack 2020; Rothier et al.

2023). However, the two perspectives have not been compared to each other to our knowledge in

limbs, either. Other inquiries into connections between evolutionary scales have focused on lines

of least resistance or primary axes of intraspecific variation as predictors of macroevolutionary

divergence (Schluter 1996; Renaud and Auffray 2013; Houle et al. 2017; Hayden et al. 2020;

Tsuboi et al. 2024), as well as explaining unanticipated stasis (Estes and Arnold 2007; Hunt and

Rabosky 2014; Hansen 2024; Voje et al. 2024) and slowdowns of evolutionary rates (Gingerich

2001; Hansen and Houle 2004; Rolland et al. 2023). Patterns of intraspecific variation in

segments have a long history of documentation and a search for causes but not, as far as we can

tell, a comparison of patterns across evolutionary scales that would reveal this discrepancy

(Gingerich 1974; Yablokov 1974; Polly 1998).

To address this apparent conflict in interpretation, in this study we examine variation in

serially segmented structures at the within-species, microevolutionary level. Specifically, we use

two complementary approaches to test two complementary explanations for the

microevolutionary interpretation of reduced evolutionary potential in successive segments

(hereafter: Reduced Potential model) using the mammalian molar tooth module as a study

system. Both explanations focus on how decreasing heritability is partitioned among causes. The

first explanation is that reduced heritability could be caused by reduced levels of additive genetic

variation in successive molars, indicating a stronger discordance between microevolution and

macroevolution. The second explanation is that reduced heritability could be caused by increased

levels of environmentally induced variation, which may mask concordance between the two

levels of evolution.

Predictions for Evolvability

One interpretation of a Reduced Potential model is that later-forming structures carry less

additive genetic variation, the kind of variation that can respond to selection and evolve over

generations (Futuyma and Kirkpatrick 2017). Heritability has long served as a comparative

metric for such purposes, but the evolvability (I_A) metric has emerged as a more appropriate,

direct evaluation of evolutionary potential (Houle 1992; Hansen and Wagner 2023). Both

heritability and evolvability are calculated from quantitative genetic studies partitioning

phenotypic variation into additive genetic (V_A) and other (V_E) components (note that V_E is

sometimes called the environmental component but does not refer solely to an ecological

environment with which a species interacts). Evolvability makes the additive genetic component

comparable between traits and samples by mean-scaling evolvable variance (V_A), using the same

mean-scaling logic that underlies coefficients of variation (CV; Houle 1992). In contrast,

heritability conflates potential for both an evolved response and a non-evolved, phenotypically

plastic response by scaling evolvable variance by total phenotypic variance, which includes both

sources of variation (Hansen et al. 2011). Therefore, the observation that heritability decreases

downstream in both teeth and limbs may not actually indicate a pattern of reduced evolvability or

evolutionary potential (Alvesalo and Tigerstedt 1974; Hallgrímsson et al. 2002). To better

address this interpretation we conducted a systematic review of patterns of heritability down the

molar row, then used these results to calculate evolvability. If the Reduced Potential model

operates through reduced downstream levels of additive genetic variation, indicating reduced

evolutionary potential, then:

Prediction 1: We expect to observe decreased evolvability along the molar row, similar to

published patterns for heritability (Fig. 1).

 Predictions for Phenotypically Plastic Patterns

A complementary interpretation of the Reduced Potential model is that later-forming

structures are more phenotypically variable because their formation is more strongly affected by

their environment. This expectation has historically been based on patterns of variance or

standardized CV (Soulé 1982). However, if the underlying mechanism is correct, then the model

also predicts that the environment should also more strongly affect mean trait values, which

might also contribute to patterns of variance (Fig. 2). To address this interpretation, we isolate

phenotypically plastic variation and characterize its pattern along the molar row using a

controlled feeding study of inbred lab rats (*Rattus norvegicus*). Phenotypically plastic variation

occurs when different environments induce different phenotypes from an identical genotype

(West-Eberhard 1989). Environments may refer to climatic conditions, such as temperature

inducing trait change (Gupta and Lewontin 1982), as well as other external conditions, such as

nutrition availability (Sciulli et al. 1979), as well as highly local conditions, such as slight

differences in cell locations and other features of the developmental microenvironment

surrounding symmetric structures, known as developmental instability (Leamy and Klingenberg

2005; Cano-Fernández et al. 2023). Resulting structures share genotypes and should be

phenotypically perfectly symmetrical, but often carry a small amount of fluctuating asymmetry

(Van Valen 1962).

Figure 2. Conceptual diagram of two different ways in which a low-protein diet, a kind of

environmental perturbation, could generate a pattern of increasing coefficients of variation (CV)

in downstream tooth sizes, matching previously published patterns of variation in tooth size. A.

Control group, in which each tooth sample has a mean size (black solid outline) and variance

(gray dashed ellipse). B., Low Protein group, in which all tooth sizes are reduced, but either (left)

downstream teeth are more strongly reduced in mean size than upstream teeth, resulting in an

increase in CV because of a more strongly reduced denominator in the CV equation, or (right)

downstream teeth have more strongly increased variance than upstream teeth, resulting in an

increase in CV because of a more strongly increased numerator in the same equation.

Abbreviations: CV, coefficient of variation; s, standard deviation; x̄, sample mean.

Use of a controlled feeding study perturbing protein amounts in the diet provided a plausible

case in which mean trait values as well as variance might be differentially affected by the

environment through poor nutrition. It allowed us to develop predictions based on prior work

relating molar growth and nutrition (Holloway et al. 1961; Sciulli et al. 1979; Blankenship-

Sefczek et al. 2024). For example, reduced growth rates could affect trait means, but would not

necessarily affect variance (Miller and German 1999; Schwab and Moczek 2016). The same

mechanism could also affect variance if, for example, later-forming segments are smaller or if a

compounding effect reduces means, resulting in an increase in mean-scaled measures of variance

such as CV (Fig. 2; Polly 1998). We refer to a situation where a single environmental change has

an increasingly strong mean effect on downstream segments as one of decreasing environmental

canalization (Hallgrímsson et al. 2002).

Poor nutrition might also affect variances without affecting mean effect if it changed levels of

developmental instability. Developmental instability is considered as a separate phenomenon

from environmental canalization, with separate underlying developmental mechanisms

(Hallgrímsson et al. 2002; Cano-Fernández et al. 2023). If an environmental condition increases

developmental instability without inducing other changes, then the random changes will increase

the variance but should not affect the mean (Palmer and Strobeck 1986; Hallgrímsson et al.

2002). In the case of poor nutrition, increased developmental instability could be caused by

decoherence, or a loss of co-regulation of traits driven by disruptions to gene regulatory

networks when homeostasis is lost (Lea et al. 2019). Therefore, in this study we assess our

expectations through both means and variances, using the two separate patterns to help interpret

potential developmental causes of increasing downstream variation addressed by the Reduced

Potential model.

Increased downstream effect size could also be inherent to serially segmented structures

which share iterative expression of similar or identical sets of developmental pathways (Thesleff

and Sharpe 1997; Young et al. 2015). Early-forming structures have the capacity to interact with

later-forming structures, producing downstream phenotypes that may differ between individuals

only because of variation in phenotype of the initial segment, not because of any change in

genetic basis between segments or because of different relationships of different segments to the

environment (Kavanagh et al. 2007). This cascading process could compound a phenotypically

plastic effect in later-forming segments, producing a stronger effect and less environmental

canalization in the phenotypes of later-forming segments (Hallgrímsson et al. 2002). This

compounding would predict a pattern of increasing effects of environmental impact downstream,

following the Reduced Potential model, regardless of whether environmental canalization or

developmental instability is part of the underlying mechanism.

Alternatively, effects may not compound, but instead the environment could have different

direct effect on each segment. Although there is evidence of compounding in processes

producing activation/inhibition ratios that permit enamel knot formation (Kavanagh et al. 2007),

that system does not require compounding in other processes like a general growth mechanism

which might also affect molar sizes in conjunction with enamel knot dynamics. This hypothesis

of multiple influences on size, some compounding and some not, is consistent with prior

findings. Molar size phenotypes are incompletely described by a single model of compounding

developmental process (Roseman and Delezene 2019; Vitek et al. 2020). In addition, molar

positions have some level of genetic independence from one another, albeit often small, implying

that the sets of developmental pathways that control their growth may be slightly different

(Hlusko et al. 2011; Hardin 2020). These deviations from a cascading model fit an alternative

proposal that a potentially different effect of phenotypic plasticity on different molars is instead

related to different levels of exposure, or the amount of time each tooth spends developing,

implying that the molars are equally canalized against environmental effects per unit of exposure

time (Butler 1983; Townsend et al. 2009). Either model could explain previously observed

patterns of decreasing heritability and increasing phenotypic variance down the tooth row

(Kavanagh et al. 2007; Riga et al. 2014; Denes et al. 2018). Following either model and

consistent with previously documented patterns for variance:

Prediction 2: We expect the phenotypically plastic effect of any environmentally induced

change in mean and variance will increase along the tooth row from M_1 to M_3.

Finally, if the differential environmental sensitivity prediction of the Reduced Potential

model is true, then we expect to observe different impacts of a poor-nutrition environment on

molar length and width based on prior studies relating molars to growth. Width, but not length, is

significantly genetically correlated with body size, and therefore more likely to be related to

general growth factors affected by nutrition (Hlusko et al. 2006). Molars also reach their

maximum length earlier in development, before they reach maximum width, with a strong

constraints imposed by the width of the surrounding bone (Renvoisé et al. 2017; Christensen et

al. 2023), similarly supporting the hypothesis that length and width are determined by different

sets of developmental pathways. If genetic correlations between body size and molar width, but

not molar length, occur through sharing pathways that affect growth in general, and if those

pathways are affected by nutrition (Miller and German 1999; Lui and Baron 2011), then we

expect that molar traits more strongly correlated with body size to also be more strongly affected

by nutrition than other molar traits. Therefore, if a nutrition-specific application of the Reduced

Potential model predicts a stronger effect on environment in some traits but not others, then:

Prediction 3: We expect poor nutrition will have a stronger effect on molar width than molar

length.

Overall, this approach of separating expectations for evolvable phenotypic variation and

plastic phenotypic variation may clarify potential connections between microevolutionary and

macroevolutionary patterns, resolving apparent disconnects in interpretation.

Methods

Evolvability

To test Prediction 1, we conducted a systematic literature review (Supplementary Methods;

Page et al. 2021). The objective of this systematic review was (A) to assess the consistency of the

evidence for the common characterization that heritability decreases down the molar row (Paul et

al. 2023), and (B) to use reported parameter values to calculate evolvability and evaluate whether

it has any consistent pattern down the molar row. Trait heritability values and their standard

errors were collated for analysis. To meet minimum criteria for calculating evolvability,

publications needed to report the following for buccolingual width or mesiodistal length for at

least two lower molar positions: (1) trait means (m), (2) trait variances (V_P), (3) trait heritability

($h^2 = V_A/V_P$). From these values, we could calculate evolvability, $I_A = V_A/m^2$, from heritability

using the equation: $I_A = h^2 * V_P /m^2$. Error in evolvability was estimated by resampling: a

simulated sample of 1,000 additive variance estimates was calculated by sampling a normal

distribution with the mean and standard error of a given sample from the literature. The resulting

simulated estimates were then summarized as a standard deviation, which was plotted around

each estimate. Given the range of study designs, we did not conduct a meta-analysis of patterns

across successive teeth. Instead, we tabulated the number of datasets that reported higher, lower,

or ambiguous levels of heritability or evolvability in downstream molars. Limitations of sample

size also prevented us from assessing between-taxon differences, although phylogenetic

distinctions in tooth anatomy and development exist (Ungar 2010). Some datasets are right and

left sides from the same individuals, which we visualized separately for clarity but for which we

included only one side, the left, in the tabulation.

Trait Plasticity

To test Predictions 2 and 3, we leveraged data collected from a previously conducted

controlled experiment originally intended to study the effect of maternal malnutrition on insulin

action and adipose tissue in male offspring of Wistar Han rats (*Rattus norvegicus*; Crl:WI[Han]

strain). This type of laboratory study used inbred lines to control for genetic variation between

individuals, allowing us to assume that all phenotypic variation was due to some component of

environment. Thus, by comparing phenotypic patterns between traits and between experimental

groups we could study the impact of environment on traits. In this case, the environmental

variable that differed between control and experimental groups was the quality of the maternal

diet throughout gestation and suckling (8% low-protein experimental diet group vs. 20% protein

control diet group), and we measured its effect on offspring phenotype (Martin-Gronert et al.

2016; Ward et al. 2021). Nutritional variation is known to influence dental phenotypes, providing

a high likelihood of observing environmental impacts in our study (Holloway et al. 1961; Miller

and German 1999; Calsa et al. 2022). The time window of environmental perturbation (Day 1 of

gestation, E1, through weaning on postnatal day P21) is appropriate for studying the effect of

environment on rat molars because molar final size is determined during a finite window of

growth and then remains unchanged for the remainder of an individual's life (Larson and Bader

1976). In rats, which develop 1-2 days slower than mice, molars initiate formation as the first

molar (M_1) placode on embryonic day E13-14, then a tooth bud at E16-17 (Larson and Bader

1976; Christensen et al. 2023). The M_2 tooth bud appears at E17-19, and the M_3 tooth bud is

visible approximately 10 days later, at postnatal day P5-7 (Larson and Bader 1976; Kavanagh et

al. 2007). Each molar reaches an inflection point and slows growth about 5 days after the bud

stage, and achieves its final size about 8 days after bud stage (Christensen et al. 2023), meaning

that in the rat, final M_1 size is achieved by P1-2, M_2 size by P3-4, and M_3 size by P13-15.

Therefore, we consider each molar tooth to be equally exposed to a consistent environmental

perturbation.

To collect phenotypic trait data, we used μCT scans of offspring sacrificed at age 3 months

(N = 41 total, 25 control, 16 low-protein) to generate 3D models from which linear measures

could be collected. Further details of experimental design and μCT scan collection are reported

in (Martin-Gronert et al. 2016; Ward et al. 2021). From μCT image stacks, volumes of the left

molar row were segmented into 3D surfaces using automated thresholding in Avizo® version

2019.3 (FEI, Hilsboro, USA). Smoothing of the region of interest was conducted in each of the

three standard planes, but after a surface was extracted from that region we conducted no further

smoothing. From these surfaces, mesiodistal length and buccolingual width were measured in

triplicate by a single observer using MeshLab (Cignoni et al. 2008). To evaluate whether

measurement error might contribute an undue amount of spurious noise to our estimates of

variation, as has been proposed to explain some patterns in variation across molars (Polly 1998),

we calculated percent measurement error for each trait (Yezerinac et al. 1992).

We calculated CV for each experimental group separately to ensure that potentially different

amounts of change in mean trait size did not spuriously influence estimates of variation. To

evaluate whether CVs differed between samples, we used a bootstrap resampling approach (N

replicates = 10,000) to test the hypothesis that one trait was more variable (larger CV) than

another trait compared to what would be expected by chance. We specifically tested hypotheses

that CV should increase between successive widths (one-tailed), between successive lengths

(one-tailed), that it should differ between length and width (two-tailed), and whether it differed

between control and experimental groups (two-tailed). Bootstrap resamples of low-protein CVs

were used to generate the null hypothesis distribution for between-trait comparisons.

To determine if it was necessary to test hypotheses of mean effect on any specific trait, we

first established whether the nutritional environment had a significant effect on trait means by

comparing experimental groups. For mean trait size, we used a t-test. To compare effect between

teeth, it was necessary to take the different sizes of each molar position into account ($M_1 > M_2 >$

M_3), because the same magnitude of difference has a different meaning for effect on the M_1 vs.

the M_3. We calculated a percent reduction statistic [mean control value – mean experimental

value] / [mean experimental value] to mean-scale the effect, similar to the mean scaling

performed for variation (CV) and evolvability (I_A) (Houle 1992). To evaluate whether effect size

differences were significant, we used a bootstrapping resampling approach. To test Prediction 2,

we compared successive length values to each other and successive width values to each other.

To test Prediction 3, we compared width to length values within each tooth. When each

hypothesis was evaluated using multiple tests, we report p-values corrected using the Bonferroni

approach (Holm 1979). All analyses and visualizations were conducted in R version 4.4.1 (R

Core Team 2015) using packages `dplyr`, `reshape2`, `ggplot2`, `ggthemes`, and `patchwork`

(Wickham 2007, 2009; Wickham et al. 2023; Arnold 2024; Pedersen 2024).

Results

Evolvability

Eight reports out of 536 items recovered in literature searches reported length or width

heritability for multiple molars, including two records found by searching the citations of six

studies recovered from database searches (Dempsey and Townsend 2001; Hlusko et al. 2011) (SI

Fig. 2). Seven samples from five of those eight reports met all three criteria for calculating

evolvability (Bader 1965; Leamy and Bader 1968; Leamy and Touchberry 1974; Hlusko et al.

2011; Hardin 2019). Evolvability (I_A) patterns were generally ambiguous or increasing along the

molar row for both lengths and widths (Fig. 3). Heritability (h^2) patterns were generally

ambiguous or decreasing along the molar row, given the large standard errors around mean

estimates (SI Table 1-2).

Figure 3. Evolvability (I_A) and heritability (h^2) calculated from a systematic review of published samples. Lines connect traits measured from the same sample. Top panel shows values from studies that did not report enough information to calculate evolvability. Middle panel shows samples with paired evolvability estimates (bottom panel). Bars indicate standard error, jittered to better show overlap. Note that the y-axis is on a logarithmic scale to better display the range of evolvability estimates. Circles indicate length measurements. Triangles indicate width measurements.

Trait Plasticity

Measurement error was low across traits (<2%, Table 1). All six traits were significantly

smaller in the low-protein offspring group compared to the control offspring group (Table 1).

Effect size, as measured by percent reduction from control to low-protein, was not consistent

down the toothrow (Table 2, Fig. 4A). Length of M_2 was significantly more strongly affected

than the length of the M_1. Widths of M_2 and M_3 were more strongly affected than width of the

M_1. No other comparison between upstream and downstream effects was significant. Widths

were generally more strongly affected than lengths, but only significantly so in the M_1 and M_3.

Variance, as measured by CV, did not differ significantly between the control and low-protein

group, with the single exception of M_3 length. CV significantly increased down the molar row

for lengths, but not for widths (Table 2, Fig. 4B). Widths were only more variable than lengths

for the M_1.

Table 1. Summary statistics for molar size traits.

Trait	Tooth	PME	C vs. LP x^2 p-value	LP vs. C % Reduced	CV C	CV LP	C vs. LP CV p-value
Length	M_1	0.58	**< 1 * 10⁻⁵**	5.53	2.76	3.04	1
	M_2	0.34	**< 1 * 10⁻⁵**	8.81	3.88	3.39	1
	M_3	0.36	**< 1 * 10⁻⁵**	6.67	4.75	5.88	**0.0216**
Width	M_1	1.29	**< 1 * 10⁻⁵**	9.56	3.3	3.84	0.4008

M_2	0.89	**$< 1 * 10^{-5}$**	7.4	3.58	4.32	0.6102
M_3	0.38	**$< 1 * 10^{-5}$**	11.74	4.83	5.25	1

365 Significance values reported post-Bonferroni correction. Significant p-values are in bold.

366 Abbreviations: C, control group; CV, coefficient of variation; LP, low protein group; p, p-value;

367 PME, percent measurement error; x̄, mean trait value

369 Table 2. P-values assessing significance of differences between pairs of traits. Low-protein CVs

370 were used for between-trait comparisons.

	Length vs. Width			Length		Width	
	% Reduced	CV		% Reduced	CV	% Reduced	CV
M_1	< 0.0003	0.0051	M_1 vs. M_2	0.0003	0.036	1	1
M_2	1	1	M_2 vs. M_3	1	0.0021	0.0003	0.0003
M_3	< 0.0003	1	M_1 vs. M_3	1	0.0003	0.0003	0.1437

371 Significance values reported post-Bonferroni correction. Abbreviations: CV, coefficient of

372 variation.

Figure 4. Effect of low-protein nutrition on the mean and variance of molar size traits,

standardized by mean trait size. Box and whisker plots show median (thick bar) and mean

(triangle). Corresponding density plots are to the right. Note that density plots height may not

match box plot median because of the different way each approach summarizes the distribution.

A. Percent that low-protein group is reduced compared to control group. B. Coefficient of

variation. Boxes with upward-facing triangles show the control group. Boxes with downward-

facing triangles show the low-protein group. The pair of density plots show control (left) and

low-protein (right) groups. Abbreviations: CV, coefficient of variation; L, length; W, width.

Discussion

Our results largely confirm prior patterns for statistics related to phenotypic variances

(Townsend et al. 2009). Phenotypic variances are higher in the lengths of later-forming teeth.

However, a more systematic review and comparison of published heritability values does not

support a consistent link between higher downstream variances and lower downstream

heritability or evolvability. The largely ambiguous patterns for widths in both variance-based

measures may be more common than has been previously recognized. Our additional results

suggest that molar size heritability and phenotypic variance do not reliably predict molar size

evolvability, nor do they predict patterns of effect on trait means. In short, Prediction 1 was not

supported, although the small sample size of heritability and evolvability values limits the

strength of our interpretations beyond that point. Prediction 2 was supported for measures of

variance but not trait means. The notably different patterns between lengths and widths generally

supports Prediction 3, though support is not universal. Our results are not attributable to

measurement error, as has potentially been an issue in other studies of variation (Polly 1998).

Phenotypic Plasticity in Serially Segmented Structures

Multiple processes are potentially mixed together in the characterization of later-forming

molars as more sensitive to environmental influence. The discordant patterns of phenotypic

plasticity in trait means vs. trait variances allows us to distinguish between two different types of

environmentally related processes and their potentially different relationship to serial segments.

This finding is robust to the limitations of our study. We acknowledge that because the

experimental design did not include observations of how long each molar took to form, we

cannot test the alternative hypothesis of environmental canalization scaled by exposure time.

With a single sample of one sex of one species included in our study of plasticity, it is also

possible that we did not capture a stronger effect in later-forming molar traits that may be present

in other sexes, species or genotypes (Holloway et al. 1961; Tonge and McCance 1973). It is also

likely that testing of only a single environmental cue (percent protein in the diet) does not

capture all possible patterns of covariation between traits.

However, none of these limits to generalizability contradict the point that the phenotypically

plastic responses of trait means and variances are decoupled in later-forming molars in our

sample, and therefore could be decoupled in other samples. Different developmental mechanisms

have the capacity to either respond differently to the environment or respond to different

components of the environment. Although it is possible that these responses could be coupled

under other conditions, our results demonstrate that coupling is not a necessary result of any

shared developmental cause. Importantly, this result means that one pattern should not be

assumed to represent the other, an assumption that may have contributed to the apparent conflict

between macroevolutionary and microevolutionary interpretation of segmented variation.

Instead of a single mechanism underlying a Reduced Potential model and impacting both

trait means and variances, we hypothesize that at least two separate types of environmentally

sensitive processes impact molar size traits. The first process we hypothesize is an effect of

nutritional environment on general organismal growth, which affects molars to the degree that

they are environmentally decanalized, or developmentally dependent on overall organismal

growth. The developmental dependence on general growth across molars need not compound

down the molar row, allowing for nonlinear patterns like those we observed in our study,

especially for width measurements (Fig. 4A, Table 1), with no requirement for a shared pattern

across structures like teeth, limbs, and vertebral segments with very different developmental

underpinnings.

The second process we hypothesize is a model of developmental instability compounding

down the development of the molar row, explaining patterns of phenotypic variance, heritability,

and fluctuating asymmetry documented in this and other studies (Sofaer et al. 1971; Gingerich

1974; Hlusko et al. 2011). Our data collection protocol did not permit us to evaluate fluctuating

asymmetry, a more standard statistic for inferring developmental instability (Palmer and

Strobeck 1986; Hallgrímsson et al. 2002), but prior studies have found a consistent pattern of

downstream increase, consistent with our explanation (Sofaer et al. 1971; Alibert et al. 1994).

The other phenomenon that would produce such an effect, antisymmetry, is not a feature of

mammalian tooth sizes (Van Valen 1962; Leamy et al. 2005).

Unlike environmental canalization, developmental instability might compound more

generally across serial structures if it were caused by iterative expression of a gene regulatory

network. Notably, molar length fits expected patterns of this model (Table 2), given that length is

more conceptually closely tied to a model of compounding activation/inhibition dynamics than

molar width (Kavanagh et al. 2007). The lack of significant difference in CV between the two

experimental groups (Table 1) suggests that this compounding instability is not related to

nutrition, but instead may be a more general feature of certain developmental mechanisms

(Hallgrímsson et al. 2002). When successive segments interact with each other through

expression of one gene regulatory network, any stochastic developmental instability in the

network affecting the M_1 would add on to the stochasticity affecting the M_2 itself, and so on

down the tooth row. No tooth would have developmental instability less than zero, and therefore

developmental instability would only ever remain stable or increase down a set of segments,

resulting in higher downstream phenotypic variance if all else were equal among molars. The

nature of developmental instability would mean that this random process averages out with no

accumulating downstream effect on trait means in a sample, even though the effect accumulates

downstream in any given individual.

In sum, our results contradict the commonly held hypothesis that increasing developmental

instability is mechanistically related to increasing sensitivity to environment and decreasing

environmental canalization (Hallgrímsson et al. 2002). Although compounding developmental

instability can potentially explain patterns of variance-related statistics, it cannot explain either

the patterns of environmental canalization or of evolvability. If this model holds, then any set of

serial segments that share a single gene regulatory network may show this pattern of increasing

variance, not necessarily tethered to increasing sensitivity to the environment.

Plasticity & Organismal Growth Reflected in Teeth

Prior work found stronger genetic correlations between molar width and overall body size

than between molar length and body size (Hlusko et al. 2006). Further work built on these links

to focus on more tooth-specific evolutionary patterns, rather than patterns that might reflect more

mixed evolutionary signals of body size and tooth size evolution (Hlusko et al. 2016). In our

study, all molar size traits were significantly, phenotypically plastically reduced by poor

nutrition, highlighting that no trait is fully buffered from non-evolved, size-related responses to

environment, consistent with heritability estimates (Hlusko et al. 2011). However, the stronger

impact on widths than lengths supports the interpretation that focusing on lengths better isolates

the evolution of teeth separate from evolution of other organismal properties (Hlusko et al.

2016).

The stronger effect of poor nutrition on molar width than length may reflect how overall

organismal growth pathways are specifically expressed during tooth growth. This hypothesis

would make our phenotypic patterns a potential phenocopy (sensu Hallgrímsson et al. 2002) of

evolved changes whose underlying mutations affected the same developmental pathways. Such

patterns may be a molar-specific instance of a broader pattern of allometric variation serving as a

case of plasticity-led evolution, which has also been hypothesized for rodent mandibular

variation (Renaud and Auffray 2013; Levis and Pfennig 2021).

In the case of teeth and growth, insulin-like growth factor (IGF) signaling and expression is a

prominent candidate for future study. Insulin-like growth factors are ubiquitous during

embryonic development and regulation of IGF is critical for cellular proliferation of developing

tissue and organs, including dentition (Brown et al. 2017; Oyanagi et al. 2019; Christensen et al.

2023). Poor nutrition can impact IGF regulation by reducing the rate of protein phosphorylation,

resulting in a reduction of available IGF binding proteins and potentially resulting in a negative

feedback loop (Tagliabracci et al. 2015; Chrudinová et al. 2024). Reduced IGF results in

systematically smaller sizes for developing embryos without impacting the fundamental timing

or process of development due to the role of IGF in proper organogenesis (Estrella et al. 2024).

By reducing phosphorylation, poor nutrition could directly influence developing tooth size

without otherwise inducing significant alteration of gene expression in other developmental

pathways, similar to its role in genetically-based patterning of teeth (Christensen et al. 2023). In

addition, poor nutrition could indirectly influence developing tooth size by affecting growth of

the surrounding bone, which may limit space for growth of the tooth (Renvoisé et al. 2017).

Further work is needed to understand mechanisms that link nutrition to tooth size.

Reconciling Macro- and Microevolutionary Interpretations

Overall, we found equal or increasing evolutionary potential down the molar row in

microevolutionary contexts. This interpretation reconciles microevolutionary and

macroevolutionary interpretations of molars and segment patterns, with implications for other

serially segmented structures (Mongle et al. 2022). The macroevolutionary perspective of

increasing evolutionary potential is better supported across evolutionary scales by a fuller panel

of available evolvability data, sparse though it may be. Our separation of plastic effects on

downstream mean and variance explain how patterns of heritability and fluctuating asymmetry

could be real patterns, but still consistent with a model of increasing evolutionary potential.

Later-forming segments may be less heritable and statistically noisier because of their inflated

variance, but their mean values are not necessarily poorer reflections of evolutionary history than

the mean values of earlier-forming segments. To the contrary, none of the samples had lower

evolvabilities in the M_3 than in upstream molars. Instead, the M_3 either had equal evolvability,

suggesting a role for relaxed selection on the M_3 compared to upstream molars (Gingerich and

Winkler 1979; Mongle et al. 2022), or had increased evolvability, suggesting that there is also

comparatively more additive genetic variation for selection to act on in these teeth (Fig. 3). This

perspective may explain why the M_3 has been considered so taxonomically useful in certain

rodents (Barnosky 1993; Polly 2003), despite their general characterization as a noisy, non-ideal

choice for taxonomic study (Gingerich 1974; Vezzosi and Kerber 2018). Mammalian molars

appear to be another case where heritability is not as informative of evolutionary potential as

evolvability (Hansen et al. 2011).

It is not clear if the same explanation applies to vertebrae limbs, given the many differences

between limb and molar development. In limbs, the macroevolutionary pattern of higher

evolutionary rates in later-forming segments is attributed to function, where more distal,

downstream segments have more direct interaction with substrates and other sources of selective

pressure (Rothier et al. 2023). This selection pressure may make their potential more easily

observed, in contrast to the molar module where function is often conserved down the tooth row

and treated as a single functional complex along with other cheek teeth (Ungar 2010).

Investigating the macroevolution of taxa that differ in their degree of functional variation down

the molar row may provide an important comparison to patterns in the limb, filling in a

remaining gap in our understanding of how microevolutionary and macroevolutionary

diversification align.

Data accessibility statement: Data and code sufficient to reproduce results in this study are

contained in a Dryad publicly accessible digital repository [*private reviewer link: in submission*]

Conflicts of interest statement: The authors declare no conflicts of interest.

Citations

Alibert, P., S. Renaud, B. Dod, F. Bonhomme, and J.-C. Auffray. 1994. Fluctuating asymmetry in
the *Mus musculus* hybrid zone: a heterotic effect in disrupted co-adapted genomes.
Proceedings of the Royal Society of London. Series B: Biological Sciences 258:53–59.

Alvesalo, L., and P. M. A. Tigerstedt. 1974. Heritabilities of human tooth dimensions. Hereditas
77:311–318.

Amaral, E. D. O. S., M. Jara-Espejo, and S. R. P. Line. 2024. The minimum free energy of G-
quadruplex-forming sequences of the MSX1 intron is associated with relative molar and
premolar size in eutherians. Journal of Mammalian Evolution 31:22.

Arnold, J. B. 2024. ggthemes: Extra Themes, Scales and Geoms for "ggplot2."

Bader, R. S. 1965. Heritability of dental characters in the house mouse. Evolution 19:378–384.

Bader, R. S., and W. H. Lehmann. 1965. Phenotypic and genotypic variation in odontometric
traits of the house mouse. American Midland Naturalist 74:28–38.

Barnosky, A. D. 1987. Punctuated equilibrium and phyletic gradualism: some facts from the
Quaternary mammalian record. Pages 109–147 *in* H. H. Genoways, ed. Current
Mammalogy. Springer US, Boston, MA.

Barnosky, A. D. 1993. Mosaic evolution at the population level in *Microtus pennsylvanicus*.

Pages 24–59 *in* R. A. Martin and A. D. Barnosky, eds. Morphological Change in

Quaternay Mammals of North America. Cambridge University Press, Cambridge, U.K.

Billet, G., and J. Bardin. 2021. Segmental Series and Size: Clade-Wide Investigation of Molar

Proportions Reveals a Major Evolutionary Allometry in the Dentition of Placental

Mammals. (L. Davalos, ed.)Systematic Biology 70:1101–1109.

Blankenship-Sefczek, E. C., A. H. Goodman, M. Hubbe, J. P. Hunter, and D. Guatelli-Steinberg.

2024. Nutritional supplementation, tooth crown size, and trait expression in individuals

from Tezonteopan, Mexico. (J. Cray, ed.)PLOS ONE 19:e0305123.

Böhmer, C., O. W. M. Rauhut, and G. Wörheide. 2015. Correlation between *Hox* code and

vertebral morphology in archosaurs. Proceedings of the Royal Society B: Biological

Sciences 282:20150077.

Brown, S. M., R. Peters, and A. B. Lawrence. 2017. Up-regulation of IGF-1 in the frontal cortex

of piglets exposed to an environmentally enriched arena. Physiology & Behavior

173:285–292.

Butler, P. M. 1983. Evolution and mammalian dental morphology. Journal De Biologie Buccale

11:285–302.

Calsa, B., T. J. Bortolança, B. C. Masiero, M. A. M. Esquisatto, C. A. De Oliveira, R. Catisti, and

M. Santamaria Jr. 2022. Maxillary and dental development in the offspring of

protein restricted female rats. European Journal of Oral Sciences 130:e12895.

Cano-Fernández, H., T. Tissot, M. Brun-Usan, and I. Salazar-Ciudad. 2023. On the origins of

developmental robustness: modeling buffering mechanisms against cell-level noise.

Development 150:dev201911.

Christensen, M. M., O. Hallikas, R. Das Roy, V. Väänänen, O. E. Stenberg, T. J. Häkkinen, J.-C.

François, et al. 2023. The developmental basis for scaling of mammalian tooth size.

Proceedings of the National Academy of Sciences 120:e2300374120.

Chrudinová, M., N. S. Kirk, A. Chuard, H. Venugopal, F. Zhang, M. Lubos, V. Gelfanov, et al.

2024. A viral insulin-like peptide inhibits IGF-1 receptor phosphorylation and regulates

IGF1R gene expression. Molecular Metabolism 80:101863.

Cignoni, P., M. Callieri, M. Corsini, M. Dellepiane, F. Ganovelli, and G. Ranzuglia. 2008.

MeshLab: an Open-Source Mesh Processing Tool. Pages 129–136 *in* V. Scarano, R. D.

Chiara, and U. Erra, eds. Eurographics Italian Chapter Conference. The Eurographics

Association.

Cuozzo, F. P. 2008. Using extant patterns of dental variation to identify species in the primate

fossil record: a case study of middle Eocene *Omomys* from the Bridger Basin,

southwestern Wyoming. Primates 49:101–115.

Dempsey, P. J., and G. C. Townsend. 2001. Genetic and environmental contributions to variation

in human tooth size. Heredity 86:685–693.

Denes, B. J., A. Lagou, D. Dorotheou, and S. Kiliaridis. 2018. A longitudinal study on timing and

velocity of rat molar eruption: Timing of rat molar eruption. Laboratory Animals 52:394–

401.

Emery-Wetherell, M., and E. Davis. 2018. Dental measurements do not diagnose modern

artiodactyl species: Implications for the systematics of Merycoidodontoidea.

Palaeontologia Electronica.

Estes, S., and S. J. Arnold. 2007. Resolving the paradox of stasis: models with stabilizing

selection explain evolutionary divergence on all timescales. The American Naturalist

169:227–244.

Estrella, C. A. S., K. L. Gatford, R. Xiang, A. Javadmanesh, M. Ghanipoor-Samami, G. S.

Nattrass, E. Shuaib, et al. 2024. Asymmetric growth-limiting development of the female

conceptus. Frontiers in Endocrinology 14:1306513.

Futuyma, D. J., and M. Kirkpatrick. 2017. Evolution (Fourth edition.). Sinauer Associates, Inc.,

Publishers, Sunderland, Massachusetts.

Gingerich, P. D. 1974. Size variability of the teeth in living mammals and the diagnosis of

closely related fossil species. Journal of Paleontology 48:895–903.

Gingerich, P. D. 2001. Rates of evolution on time scales of the evolutionary process. Genetics

112–113:127–144.

Gingerich, P. D., and D. A. Winkler. 1979. Patterns of variation and correlation in the dentition of

the red fox, *Vulpes vulpes*. Journal of Mammalogy 60:691–704.

Gupta, A. P., and R. C. Lewontin. 1982. A study of reaction norms in natural populations of

Drosophila pseudoobscura. Evolution 36:934–948.

Hallgrímsson, B., K. Willmore, and B. K. Hall. 2002. Canalization, developmental stability, and

morphological integration in primate limbs. American Journal of Physical Anthropology

119:131–158.

Hansen, T. F. 2024. Three modes of evolution? Remarks on rates of evolution and time scaling.

(L. Holman & T. Latrille, eds.)Journal of Evolutionary Biology voae071.

Hansen, T. F., and D. Houle. 2004. Evolvability, stabilizing selection, and the problem of stasis.

Pages 130–154 *in* M. Pigliucci and K. Preston, eds. Phenotypic Integration: Studying the

Ecology and Evolution of Complex Phenotypes. Oxford University Press, New York.

Hansen, T. F., C. Pélabon, and D. Houle. 2011. Heritability is not evolvability. Evolutionary

Biology 38:258–277.

Hansen, T. F., and G. P. Wagner. 2023. The Evolution of Evolvability. *in* T. F. Hansen, D. Houle,

M. Pavličev, and C. Pélabon, eds. Evolvability: A Unifying Concept in Evolutionary

Biology? The MIT Press.

Hardin, A. M. 2019. Genetic contributions to dental dimensions in brown☐mantled tamarins (

SAGUINUS FUSCICOLLIS) and rhesus macaques (*MACACA MULATTA*). American Journal of

Physical Anthropology 168:292–302.

———. 2020. Genetic correlations in the rhesus macaque dentition. Journal of Human Evolution

148:102873.

Harris, E. F. 2003. Where's the Variation? Variance Components in Tooth Sizes of the Permanent

Dentition. Dental Anthropology Journal 16:84–94.

Hayden, L., K. Lochovska, M. Sémon, S. Renaud, M.-L. Delignette-Muller, M. Vilcot, R.

Peterkova, et al. 2020. Developmental variability channels mouse molar evolution. eLife

9:e50103.

Hershkovitz, P. 1967. Dynamics of Rodent Molar Evolution: A Study Based on New World

Cricetinae, Family Muridae. Journal of Dental Research 46:829–842.

Hlusko, L. J., L. R. Lease, and M. C. Mahaney. 2006. Evolution of genetically correlated traits:

Tooth size and body size in baboons. American Journal of Physical Anthropology

131:420–427.

Hlusko, L. J., R. D. Sage, and M. C. Mahaney. 2011. Modularity in the mammalian dentition:

mice and monkeys share a common dental genetic architecture. Journal of Experimental

Zoology Part B: Molecular and Developmental Evolution 316B:21–49.

Hlusko, L. J., C. A. Schmitt, T. A. Monson, M. F. Brasil, and M. C. Mahaney. 2016. The

integration of quantitative genetics, paleontology, and neontology reveals genetic

underpinnings of primate dental evolution. Proceedings of the National Academy of

Sciences 201605901.

Holloway, P. J., J. H. Shaw, and E. A. Sweeney. 1961. Effects of various sucrose:casein ratios in

purified diets on the teeth and supporting structures of rats. Archives of Oral Biology

3:185–200.

Holm, S. 1979. A simple sequentially rejective multiple test procedure. Scandinavian journal of

statistics 6:65–70.

Houle, D. 1992. Comparing evolvability and variability of quantitative traits. Genetics 130:195–

204.

Houle, D., G. H. Bolstad, K. Van Der Linde, and T. F. Hansen. 2017. Mutation predicts 40

million years of fly wing evolution. Nature 548:447–450.

Hunt, G., and D. L. Rabosky. 2014. Phenotypic evolution in fossil species: pattern and process.

Annual Review of Earth and Planetary Sciences 42:421–441.

Kavanagh, K. D., A. R. Evans, and J. Jernvall. 2007. Predicting evolutionary patterns of

mammalian teeth from development. Nature 449:427–432.

Kieser, J. A., and H. T. Groeneveld. 1987. A metric analysis of *Cercopithecus aethiops* dentition.

International journal of primatology 8:261–280.

Larson, M. J., and R. S. Bader. 1976. Maternal Effects in Odontometric Traits of House Mice.

American Midland Naturalist 96:21.

Lea, A., M. Subramaniam, A. Ko, T. Lehtimäki, E. Raitoharju, M. Kähönen, I. Seppälä, et al.

2019. Genetic and environmental perturbations lead to regulatory decoherence. eLife

8:e40538.

Leamy, L. J., and R. S. Bader. 1968. Components of Variance of Odontometric Traits in a Wild-

Derived Population of Peromyscus leucopus. Evolution 22:826.

Leamy, L. J., and C. P. Klingenberg. 2005. The genetics and evolution of fluctuating asymmetry.

Annual Review of Ecology, Evolution, and Systematics 36:1–21.

Leamy, L. J., and R. W. Touchberry. 1974. Additive and non-additive genetic variance in

odontometric traits in crosses of seven inbred lines of house mice. Genetical Research

23:207–217.

Leamy, L. J., M. S. Workman, E. J. Routman, and J. M. Cheverud. 2005. An epistatic genetic

basis for fluctuating asymmetry of tooth size and shape in mice. Heredity 94:316–325.

Levis, N. A., and D. W. Pfennig. 2021. Innovation and Diversification Via Plasticity-Led

Evolution. Pages 211–240 *in* D. W. Pfennig, ed. Phenotypic Plasticity & Evolution:

Causes, Consequences, Controversies (1st ed.). CRC Press.

Lui, J. C., and J. Baron. 2011. Mechanisms Limiting Body Growth in Mammals. Endocrine

Reviews 32:422–440.

Martin-Gronert, M. S., D. S. Fernandez-Twinn, M. Bushell, K. Siddle, and S. E. Ozanne. 2016.

Cell-autonomous programming of rat adipose tissue insulin signalling proteins by

maternal nutrition. Diabetologia 59:1266–1275.

Miller, J. P., and R. Z. German. 1999. Protein Malnutrition Affects the Growth Trajectories of the

Craniofacial Skeleton in Rats. The Journal of Nutrition 129:2061–2069.

Mongle, C. S., A. Nesbitt, F. A. Machado, J. B. Smaers, A. H. Turner, F. E. Grine, and J. C.

Uyeda. 2022. A common mechanism drives the alignment between the micro☐ and

macroevolution of primate molars. Evolution evo.14600.

Oyanagi, T., N. Takeshita, M. Hara, E. Ikeda, T. Chida, D. Seki, M. Yoshida, et al. 2019. Insulin-

like growth factor 1 modulates bioengineered tooth morphogenesis. Scientific Reports

9:368.

Page, M. J., J. E. McKenzie, P. M. Bossuyt, I. Boutron, T. C. Hoffmann, C. D. Mulrow, L.

Shamseer, et al. 2021. The PRISMA 2020 statement: an updated guideline for reporting

systematic reviews. Systematic Reviews 10:89.

Palmer, A. R., and C. Strobeck. 1986. Fluctuating asymmetry: Measurement, Analysis, Patterns.

Annual Review of Ecology and Systematics 17:391–421.

Paul, K. S., R. Feezell, T. Hughes, and A. H. Brook. 2023. Integrating genealogy and dental

variation: contributions to biological anthropology. American Journal of Biological

Anthropology 181:145–179.

Pedersen, T. L. 2024. patchwork: The Composer of Plots.

Polly, P. D. 1998. Variability in mammalian dentitions: size-related bias in the coefficient of

variation. Biological Journal of the Linnean Society 64:83–99.

Polly, P. D. 2003. Paleophylogeography: the tempo of geographic differentiation in marmots

(*Marmota*). Journal of Mammalogy 84:369–284.

R Core Team. 2015. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Renaud, S., and J.-C. Auffray. 2013. The direction of main phenotypic variance as a channel to

morphological evolution: case studies in murine rodents. Hystrix, the Italian Journal of

Mammalogy 24:85–93.

Renvoisé, E., K. D. Kavanagh, V. Lazzari, T. J. Häkkinen, R. Rice, S. Pantalacci, I. Salazar-

Ciudad, et al. 2017. Mechanical constraint from growing jaw facilitates mammalian

dental diversity. Proceedings of the National Academy of Sciences 201707410.

Riga, A., M. G. Belcastro, and J. Moggi-Cecchi. 2014. Environmental stress increases variability

in the expression of dental cusps: Environmental Influence on Dental Morphology.

American Journal of Physical Anthropology 153:397–407.

Rolland, J., L. F. Henao-Diaz, M. Doebeli, R. Germain, L. J. Harmon, L. L. Knowles, L. H.

Liow, et al. 2023. Conceptual and empirical bridges between micro- and macroevolution.

Nature Ecology & Evolution 7:1181–1193.

Roseman, C. C., and L. K. Delezene. 2019. The inhibitory cascade model is not a good predictor

of molar size covariation. Evolutionary Biology 46:229–238.

Rothier, P. S., A.-C. Fabre, J. Clavel, R. B. Benson, and A. Herrel. 2023. Mammalian forelimb

evolution is driven by uneven proximal-to-distal morphological diversity. eLife

12:e81492.

Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. Evolution 50:1766.

Schuman, E. L., and C. L. Brace. 1954. Metric and morphologic variations in the dentition of the

Liberian chimpanzee; comparisons with anthropoid and human dentition. Huam Biology

26:239–268.

Schwab, D. B., and A. P. Moczek. 2016. Nutrient Stress During Ontogeny Alters Patterns of

Resource Allocation in two Species of Horned Beetles. Journal of Experimental Zoology

Part A: Ecological Genetics and Physiology 325:481–490.

Sciulli, P. W., W. J. Doyle, C. Kelley, P. Siegel, and M. I. Siegel. 1979. The interaction of

stressors in the induction of increased levels of fluctuating asymmetry in the laboratory

rat. American Journal of Physical Anthropology 50:279–284.

Sofaer, J. A., H. L. Bailit, and C. J. MacLean. 1971. A developmental basis for differential tooth

reduction during hominid evolution. Evolution 25:509–517.

Soulé, M. E. 1982. Allomeric Variation. 1. The Theory and Some Consequences. The American

Naturalist 120:751–764.

Stepanova, N., and M. C. Womack. 2020. Anuran limbs reflect microhabitat and distal,

later☐developing bones are more evolutionarily labile. Evolution 74:2005–2019.

Tagliabracci, V. S., S. E. Wiley, X. Guo, L. N. Kinch, E. Durrant, J. Wen, J. Xiao, et al. 2015. A

Single Kinase Generates the Majority of the Secreted Phosphoproteome. Cell 161:1619–

1632.

Thesleff, I., and P. Sharpe. 1997. Signalling networks regulating dental development.

Mechanisms of development 67:111–123.

Tonge, C. H., and R. A. McCance. 1973. Normal development of the jaws and teeth in pigs, and

the dela and malocclusion produced by calorie deficiencies. Journal of Anatomy 115:1–

22.

Townsend, G., E. F. Harris, H. Lesot, F. Clauss, and A. Brook. 2009. Morphogenetic fields within

the human dentition: A new, clinically relevant synthesis of an old concept. Archives of

Oral Biology 54:S34–S44.

Tsuboi, M., J. Sztepanacz, S. De Lisle, K. L. Voje, M. Grabowski, M. J. Hopkins, A. Porto, et al.

2024. The Paradox of Predictability Provides a Bridge Between Micro- and

Macroevolution. Journal of Evolutionary Biology voae103.

Ungar, P. S. 2010. Mammal teeth: origin, evolution, and diversity. Johns Hopkins University

Press, Baltimore.

Van der Made, J. 1996. Listriodontinae (Suidae, Mammalia), their evolution, systematics and

distribution in time and space. Contributions to Tertiary and Quaternary Geology 33:3–

254.

Van Valen, L. 1962. A study of fluctuating asymmetry. Evolution 16:125–142.

Vezzosi, R. I., and L. Kerber. 2018. The southernmost record of a large erethizontid rodent

(Hystricomorpha: Erethizontoidea) in the Pleistocene of South America: Biogeographic

and paleoenvironmental implications. Journal of South American Earth Sciences 82:76–

90.

Vitek, N. S., S. F. McDaniel, and J. I. Bloch. 2020. Is variation in molar tooth crown morphology

of the Grasshopper Mouse (*Onychomys leucogaster*) a reflection of selection or drift?

Society of Integrative and Comparative Biology Annual Meeting. Presented at the

Society of Integrative and Comparative Biology Annual Meeting, Austin, TX.

Voje, K. L., M. Saito-Kato, and T. L. Spanbauer. 2024. Evolution in fossil time series reconciles

observations in micro- and macroevolution. (L. Holman & T. Gaboriau, eds.)Journal of

Evolutionary Biology voae087.

Ward, D. L., L. Schroeder, E. Pomeroy, J. E. Roy, L. T. Buck, J. T. Stock, M. Martin□Gronert, et

al. 2021. Early life malnutrition and fluctuating asymmetry in the rat bony labyrinth. The

Anatomical Record ar.24601.

West-Eberhard, M. J. 1989. Phenotypic plasticity and the origins of diversity. Annual review of

Ecology and Systematics 249–278.

Wickham, H. 2007. Reshaping Data with the reshape Package. Journal of Statistical Software

21:1–20.

———. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wickham, H., R. François, L. Henry, K. Müller, and D. Vaughan. 2023. dplyr: A Grammar of

Data Manipulation.

Willis, M. S., and D. R. Swindler. 2004. Molar size and shape variations among Asian colobines.

American Journal of Physical Anthropology 125:51–60.

Yablokov, A. V. 1974. Variability of Mammals. (L. Van Valen, ed.). Amerind Publishing

Company, New Delhi.

Yezerinac, S. M., S. C. Lougheed, and P. Handford. 1992. Measurement error and morphometric

studies: statistical power and observer experience. Systematic Biology 41:471–482.

Young, N. M., B. Winslow, S. Takkellapati, and K. Kavanagh. 2015. Shared rules of

development predict patterns of evolution in vertebrate segmentation. Nature

Communications 6:6690.

Supplementary Figures and Tables with Captions

Supplementary Figure 1. PRISMA 2020 flow diagram of report retrieval.

Supplementary Table 1. Characteristics of studies recovered from the systematic literature

search.

Supplementary Table 2. Results of each sign test for consistent downstream decreases in

heritability (h^2) and evolvability (I_A) in the molar tooth row of previously published samples.

Acknowledgments

Thanks to M.T. Silcox for connecting the co-authors of this study, C.J. Percival for conversations

that helped structure the discussion. Thanks to funding from the Henry Sidgwick Research

Fellowship from Newnham College, Cambridge, University of Toronto PhD Pilot Research

Grant, and startup funding from Stony Brook University. Author RWB was funded via an

Institutional Research and Academic Career Development Award (IRACDA) made to Stony

Brook University from the National Institute of General Medical Sciences of the National

Institutes of Health [K12GM102778]. The content is solely the responsibility of the authors and

does not necessarily represent official views of the National Institutes of Health.

Funding Sources:

Henry Sidgwick Research Fellowship from Newnham College, Cambridge

University of Toronto PhD Pilot Research Grant

Startup funding from Stony Brook University

Author contributions: NSV conducted conceptualization, scripting, validation, formal analysis,

and visualization components of the study. NSV, EP, MM-G, and SEO contributed to

methodology. NSV, ES, AD contributed to phenotypic data collection. DW, EP, MM-G, and SEO

contributed resources. NSV and DW contributed to data curation. NSV and SEO contributed

supervision and project administration. EP and SEO contributed funding acquisition for the

nutrition study. NSV and RWB wrote the original draft, and all authors contributed to review and

editing.