Global molecular landscape of early MASLD progression in obesity

Qing Zhao[1,2,3*], William De Nardo[4*], Ruoyu Wang[5,10*], Yi Zhong[8], Umur Keles[10], Gabriele Sakalauskaite[10], Li Na Zhao[10], Huiyi Tay[1,2], Sonia Youhanna[8], Mengchao Yan[9], Ye Xie[9], Youngrae Kim[1,2], Sungdong Lee[1,2], Rachel Liyu Lim[1,2], Guoshou Teo[1,2], Pradeep Narayanaswamy[7], Paul R. Burton[6], Volker M. Lauschke[8,9], Hyungwon Choi[1,2,3#], Matthew J. Watt[4,#], and Philipp Kaldis[10,#,&]

[1]Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[2]Singapore Lipidomics Incubator, Life Sciences Institute, National University of Singapore, Singapore
[3]Cardiovascular & Metabolic Disease Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[4]Department of Anatomy and Physiology, School of Biomedical Sciences, Faculty of Medicine Dentistry and Health Sciences, University of Melbourne, Melbourne, Victoria, Australia
[5]Department of Hepatology, The First Hospital of Hunan University of Chinese Medicine, Changsha, Hunan, China
[6]Department of Surgery, School of Translational Medicine, Monash University, Melbourne, Victoria, 3004, Australia and Bariatric Unit, Department of General Surgery, The Alfred Hospital, Melbourne, Victoria, 3004, Australia
[7]SCIEX, R&D, Singapore
[8]Department of Physiology and Pharmacology and Centre of Molecular Medicine, Karolinska Institutet, 171 77 Stockholm, Sweden
[9]Dr. Margarete Fischer-Bosch Institute of Clinical Pharmacology, Stuttgart, Germany.
[10]Department of Clinical Sciences Malmö and Lund University Diabetes Centre (LUDC), Lund University, Clinical Research Centre (CRC), 202 13 Malmö, Sweden

#Correspondence should be addressed to PK (phone +46737768101; philipp.kaldis@med.lu.se), MJW (matt.watt@unimelb.edu.au) or HC (hyung_won_choi@nus.edu.sg) – &Lead Contact

Running title: Dual-omics in obese individuals with MASLD

Keywords: Metabolic dysfunction-associated steatotic liver disease; Lipid metabolism; Fibrosis; GTPase signaling; Obesity

1 **Summary**
2
3

Metabolic dysfunction-associated steatotic liver disease (MASLD) is often
asymptomatic early on but can progress to irreversible conditions like cirrhosis. Due
to limited access to human liver biopsies, systematic and integrative molecular
resources remain scarce. In this study, we performed transcriptomic analyses on liver
and metabolomic analyses on liver and plasma samples from morbidly obese
individuals without liver pathology or at early-stage MASLD. While, the plasma
metabolomic profile did not fully mirror liver histological features, dual-omics
integration of liver samples revealed significantly remodeled lipid and amino acid
metabolism pathways. Integrative network analysis uncoupled metabolic remodeling
and gene expression as independent features of hepatic steatosis and fibrosis
progression, respectively. Notably, GTPases and their regulators emerged as a novel
class of genes linked to early liver fibrosis. This study offers a detailed molecular
landscape of early MASLD in obesity and highlights potential targets of obesity-linked
liver fibrosis.

Introduction
Metabolic dysfunction-associated steatotic liver disease (MASLD) is the most common
chronic liver disease, affecting over 30% of adults worldwide (Miao et al., 2024). The
prevalence of MASLD is closely associated with the epidemic of obesity, with an
estimated 75% prevalence among individuals with obesity compared to 32% in the
general population (Riazi et al., 2022). As liver manifestation of the metabolic
syndrome, MASLD can cause, and is also caused by concurrent metabolic
abnormalities commonly seen in obese individuals, such as insulin resistance,
diabetes, dyslipidemia, and hypertension (Targher et al., 2024; Yki-Jarvinen, 2014).
Early stages of MASLD are characterized by hepatic steatosis, which is the
excessive deposition of triacylglycerol (TAG) rich lipid droplets (LD) within the liver
parenchyma and is usually reversible. However, in a subset of patients, the disease
can develop into inflammatory and ballooning stages termed metabolic dysfunction-
associated steatohepatitis (MASH), a more severe form of the disease with an
increased incidence and severity of fibrosis. Fibrosis can be present at any stage of
the MASLD spectrum of disease and at each stage is increasingly associated with
treatment complications, liver-related and overall mortality (Angulo et al., 2015a;
Ekstedt et al., 2015; Hagstrom et al., 2017). Unresolved fibrosis can progress to end-
stage diseases including hepatic cirrhosis, and hepatocellular carcinoma, lethal
malignancies with limited treatment options (Feng et al., 2024; Rodriguez et al., 2024).
Metabolic dysfunction is a primary feature and contributor of MASLD
pathogenesis (Eslam et al., 2020). Hepatic steatosis is often accompanied by
increased glucose production (Scoditti et al., 2024), elevated de novo lipogenesis
(Lambert et al., 2014), and disrupted cholesterol homeostasis (Sakuma et al., 2025).
Mitochondrial respiration is adaptively upregulated to prevent lipid accumulation in
obesity (Koliaki et al., 2015), but with MASLD progression, mitochondria can become
dysfunctional and exacerbate metabolic dysregulation (Fromenty & Roden, 2023).
These interconnected metabolic phenotypes form the basis for the development of
MASLD (Bril et al., 2017; Younossi et al., 2018). Recent advancements in
metabolomics have greatly enhanced the understanding of the molecular systems
biology of MASLD (McGlinchey et al., 2022; Vvedenskaya et al., 2021), however, few
studies have combined such analysis with transcriptomics to provide integrative
insights into disease pathogenesis in humans.
Fibrosis is the only histological feature of MASLD associated with liver-related
mortality and morbidity (Angulo et al., 2015b). In the injured liver, damaged and
apoptotic hepatocytes modulate the crosstalk between hepatocytes, liver
macrophages (Kupffer cells), and hepatic stellate cells (HSCs) by releasing fibrogenic
cytokines (e.g., TGF-β) and activating multiple signaling pathways (Bataller & Brenner,
2005; Subramanian et al., 2022). As a result of the multicellular response, HSCs
transdifferentiate into active, collagen-producing myofibroblasts, driving fibrogenesis
and the excessive deposition of extracellular matrix (ECM) (Subramanian et al., 2022).
Since fibrosis is a key prognostic marker of MASLD progression, the shift from
steatosis to fibrosis marks a critical point in the disease, offering a key opportunity for
intervention to prevent further progression (Powell et al., 2021). Despite the availability
of certain guidelines for clinical management of MASLD (European Association for the
Study of the Liver . Electronic address et al., 2024; Rinella et al., 2023), the molecular
and metabolic changes underpinning the key transitions driving disease progression
in the context of obesity remain poorly understood. This poses a challenge to early
disease management and prevention.

In this study, we generated a comprehensive map of gene expression and
metabolomic profiles to delineate the molecular events associated with early-stage
MASLD progression in obesity. Using this multi-omic resource, we focused our
investigation on key molecular changes associated with (a) the transition from obesity
with normal hepatic histology to MASLD and (b) the onset of liver fibrosis. Our data
reveal distinct molecular signatures underlying steatosis and fibrosis progression,
offering a detailed molecular portrait of the liver in early MASLD and highlighting
potential therapeutic targets for reversing fibrosis at initial stages.

Results
Overview of the study
We analyzed samples from 109 obese individuals recruited before bariatric surgery at
The Avenue, Cabrini, or Alfred Hospitals in Melbourne, Australia (**Table 1** and
Supplementary file 1). Following the exclusion criteria described in Materials and
Methods, 33 individuals lacked histological abnormalities (no MASLD) and 76 had
MASLD (**Figure 1A**). Notably, 83 individuals (76%) were females in this cohort. Most
individuals with obesity were in the early disease stages, with 74 individuals (68%)
displaying grade 0-1 steatosis and 83 (76%) grade 0-1 fibrosis. Hepatic inflammation
and ballooning were mild, with 9 cases exhibiting grade 2 or higher inflammation and
only one case showing grade 2 ballooning (**Figure 1B** and **Table 1**). In clinical tests,
MASLD patients displayed worse liver function (higher alanine transaminase
(ALT)/aspartate aminotransferase (AST) ratio and gamma-glutamyl transpeptidase
(GGT) levels), higher non-high-density (HDL) lipoprotein cholesterol and blood
triglyceride levels, higher C-peptide, and insulin resistance compared to those with 'No
MASLD' (**Table 1**, $p < 0.05$), confirming correct classification. Liver fibrosis was
strongly correlated with insulin resistance, while steatosis grades were most correlated
with the levels of plasma lipids (**Figure 1C**).
We performed parallel transcriptomic profiling of liver slices and metabolomic
analysis of liver and plasma samples (see Materials and Methods). Untargeted
metabolomics (covering polar & non-polar metabolites including lipids) enabled semi-
quantification of 1104 liver and 962 plasma metabolites, respectively (**Figure 1D**).
Plasma urea and creatinine levels from untargeted metabolomics closely correlated
with clinical assays (**Figure 2–figure supplement 1A**), supporting data reliability.
Principal component analysis (PCA) revealed distinct clustering of the liver
metabolome by MASLD status, but not in plasma (**Figure 1E**). The liver transcriptome
was modestly separated between individuals with and without MASLD along the main
principal components (**Figure 1F**), indicating that transcriptional programs may be
shaped by both MASLD histological progression and confounding metabolic and
biological processes in obese individuals (**Supplementary file 2**).
As steatosis and fibrosis are the key histological features of MASLD, our
statistical analyses focused on identifying molecular determinants linearly associated
with their progression (**Supplementary file 3-5**). Models were adjusted for patient
characteristics (age, sex, and BMI) and type 2 diabetes, the latter due to its differing
prevalence across disease groups (**Table 1**). Moreover, we compared our
transcriptomic data with those from two additional published cohorts: the Virginia
cohort (VA cohort, GSE130970) (Hoang et al., 2019) with a disease spectrum
comparable to that of our study, and the European cohort (EU cohort, GSE135251)
(Govaere et al., 2020; Govaere et al., 2023) with a broader spectrum of the disease
covering advanced MASLD pathologies (**Figure 1G**).

Global investigation of the metabolome in obese MASLD patients

As expected, the liver metabolome was extensively remodeled in individuals with MASLD. Steatosis is the primary histological feature linked to liver metabolome, with 206 metabolites showing significant positive associations and 242 showing negative associations (q < 0.05, **Figure 2A** and **Figure 2–figure supplement 1B**). For example, we observed higher levels of glycerolipids (GLs, e.g. TAGs, diacylglycerols [DAGs]) and lower levels of membrane lipid classes, especially glycerophospholipids (GPLs) (**Figure 2A-B** and **Figure 2–figure supplement 1B**). Specifically, TAGs with 0 to 5 double bonds in the fatty acyl chains were positively associated with the histological outcomes, whereas TAGs containing at least one polyunsaturated fatty acid (PUFA) chain (e.g. number of double bonds > 5) were negatively associated with disease progression, particularly with steatosis (**Figure 2–figure supplement 1C**). Our findings are likely the results of increased de novo lipogenesis and elevated hepatotoxicity associated with saturated fatty acids (Roumans et al., 2020; Wang et al., 2006).

The plasma metabolome displayed limited associations with the histological features (**Figure 2A**), with statistically significant associations primarily with steatosis, such as TAGs (q value < 0.05, n = 22) and ether-linked TAG (TAG-O, q value < 0.05, n = 8) species. To better interpret the plasma metabolome data, we performed partial correlation network analysis to assess the associations among circulating metabolites, clinical variables, and hepatic histology in individuals with obesity (Lee et al., 2025). The network demonstrates the highest connectivity of plasma metabolites with clinical variables and fewer connections with hepatic histology features (**Figure 2C**). This implies that plasma metabolome largely reflects other metabolic conditions such as kidney function, dyslipidemia, and insulin resistance rather than hepatic features. Most plasma lipid classes were correlated with blood lipoprotein cholesterol levels (**Figure 2D**). However, in the context of obesity, blood lipoprotein levels did not show strong correlation with MASLD progression (**Figure 2–figure supplement 2**), indicating that their broad impact on plasma metabolome may mask the signals from hepatic abnormalities.

We next compared steatosis- and fibrosis-associated metabolite changes in the liver and plasma (**Figure 2E**). Plasma GLs exhibited similar but weaker associations with liver steatosis compared to hepatic GLs, whereas the depletion of PUFA-containing TAGs in the liver was not mirrored in circulation. Although plasma polar metabolites showed changes in the context of steatosis, potential circulating markers emerged for liver fibrosis with mild statistical significance. With fibrosis progression, there was a trend toward increased levels of tyrosine, quinolinic acid, lactic acid, and kynurenic acid in plasma (p < 0.05, q > 0.05) (**Supplementary file 3**), consistent with their reported roles as key hepatic metabolites implicated in MASLD progression and as proposed circulating biomarkers for MASLD severity or biopsy-proven MASH (Gou et al., 2025; Huang et al., 2025; Lahdou et al., 2013; Tan et al., 2025). Overall, despite elevated TAG levels and candidate fibrosis indicators, the circulating metabolome is less indicative of early-stage MASLD-related alterations in individuals with obesity and obesity-related co-morbidities.

Integrative view of liver metabolism remodeling

To characterize metabolic remodeling in obese individuals during early disease development, we integrated the hepatic metabolome and transcriptome.

Lipid metabolism

In the liver, accumulation of GLs and mild reduction of GPLs were the main features
of the metabolome changes (see **Figure 2**). Consistent with this observation, we
identified genes implicated in the homeostasis of GLs and GPLs (**Figure 3A** and
Figure 3–figure supplement 1). Genes such as *DGAT2*, *PNPLA3*, and *PLIN3* play a
role in LD formation and TAG and DAG metabolism (**Figure 3–figure supplement 1B**)
(Mashek, 2021). GPL metabolism was also markedly altered, including key genes
such as *LPCAT1*, *PLD2*, *PCYT2*, *ETNK*2 and that are implicated in a compensatory
response to the shift in phospholipid metabolism and the increased turnover of GPLs
(Holdaway et al., 2025; van der Veen et al., 2012). In individuals with advanced hepatic
fibrosis, expression levels of the genes involved in primary bile acid biosynthesis, such
as *SLC27A5, AMACR*, *ACOX2*, *AKR1C4*, and *BAAT*, were lower. In addition, a group
of cytochrome P450 (CYP) genes were downregulated during the progression of
fibrosis, particularly those involved in CYP-dependent PUFA metabolism (i.e., linoleic
acids and arachidonic acids into bioactive molecules) and steroid hormone
biosynthesis (Dos Santos & Fleming, 2020; Hajeyah et al., 2020) (**Figure 3A**). Overall,
our data highlights a number of genes encoding the enzyme subunits of anabolic and
catabolic lipid metabolism in early MASLD, and some of these gene signals were
consistently observed in both the EU and VA cohorts (Govaere et al., 2020; Hoang et
al., 2019).

Dysregulated metabolic pathways during steatosis progression
To explore the variation in metabolic activities using dual-omic descriptors, we
prioritized key pathways with both dysregulated metabolites and gene expressions
(**Figure 3B**). In addition to significant GL and GPL remodeling, we also identified
altered pathways of amino acid metabolism at both omics levels. Elevated levels of
amino acids including glycine, glutamic acid, arginine, serine, and alanine in steatotic
livers may reflect increased collagen synthesis and ECM remodeling during MASLD
progression (Albaugh et al., 2017; de Paz-Lugo et al., 2018) (**Figure 3C-D**). Notably,
aromatic amino acids including phenylalanine, tyrosine, and tryptophan were lower in
subjects with advanced steatosis (**Figure 3D**). Dual-omics data revealed consistent
downregulation of tryptophan metabolism, encompassing both the indole and
melatonin pathways (**Figure 3C**). However, quinolinic acid, a key metabolite in the
kynurenine pathway and a product of tryptophan catabolism, was significantly
elevated in association with both hepatic steatosis and fibrosis. Collectively, early
MASLD development is associated with reduced aromatic amino acid levels and the
altered tryptophan catabolic flux in the liver, potentially reflecting the aberrant gut–liver
axis communication in obese MASLD patients (Arto et al., 2024; Schnabl et al., 2025;
Yanko et al., 2023).
Moreover, homocysteine and its upstream metabolite S-adenosylhomocysteine
(SAH) levels were significantly higher in steatotic livers. The ratio of S-
adenosylmethionine (SAM) to SAH and that of methionine to homocysteine indicated
progressively decreasing trends with advancing steatosis (**Figure 3–figure**
supplement 2), indicating impaired methylation potential under steatotic conditions
(Walker, 2017). Furthermore, key regulators of SAM/SAH homeostasis, including
GNMT, *MAT1A*, and *AHCY*, were downregulated in association with fibrosis (**Figure**
3C), which have been linked to liver pathogenesis in *in vivo* models (Alarcon-Vila et
al., 2023; Matthews et al., 2009; Robinson et al., 2023; Varela-Rey et al., 2010).
Collectively, these findings suggest that insufficient methylation modifications of
DNAs, RNAs and proteins may be a mediator of MASLD pathogenesis, particularly in
obesity.

In addition, we observed increasing levels of the antioxidants including
coenzyme Q10 (CoQ10; ubiquinone), taurine, and alpha-tocopherol (vitamin E) with
steatosis progression (**Figure 3D**). Yet another indicator of oxidative stress,
hydroxybutyric acid, was increased in advanced steatosis and is also recognized as
an early marker of insulin resistance (Sousa et al., 2021). Overall, the progression of
liver steatosis is accompanied by extensive changes in amino acid metabolism and
oxidative stress regulation.

Dysregulated genes involved in mitochondrial function and autophagy
In the liver metabolome, lower levels of hepatic long-chain acyl-carnitine (CAR)
species were observed with advanced steatosis (CAR 18:2 and 20:4, **Supplementary**
file 4), suggesting dysregulated fatty acid transmembrane transport and β-oxidation
(Houten et al., 2016; Knottnerus et al., 2018). To systematically assess mitochondrial
functions beyond β-oxidation, we mapped steatosis and fibrosis-associated genes to
MitoCarta3.0 (Rath et al., 2021). Distinct mitochondrial dysfunction patterns emerged:
steatosis involved 26 downregulated and 55 upregulated mitochondrial-function
genes, including 16 related to mtDNA maintenance, while fibrosis was linked to 151
downregulated (dark green) and only 15 upregulated (orange) genes (**Figure 4A** and
Supplementary file 6). These transcriptional shifts suggest that hepatic fibrosis
involves broad mitochondrial impairment, contributing to oxidative stress and reduced
ATP production, which could favor the exacerbation of steatohepatitis (Fromenty &
Roden, 2023).
Autophagy serves as a critical cellular response mechanism to the overload of
intrahepatic TAG and cholesterol (Martinez-Lopez & Singh, 2015; Trivedi et al., 2021).
Metabolomic analysis showed increased free cholesterol and reduced levels of CE
18:1 and 18:2, the predominant hepatic CE species (Xie et al., 2002) (**Figure 4–figure**
supplement 1A), supporting prior findings that CE de-esterification contributes to
elevated free cholesterol in MASLD (Min et al., 2012). Although autophagy has been
linked to HSC activation (Kang & Chen, 2009; Tuohetahuntila et al., 2017), its
involvement in MASLD progression remains unclear. Using a list of predictive genes
for autophagy activation (Bordi et al., 2021), we observed upregulation of MTOR, a
central member gene of mTORC1 complex that tightly regulates autophagy, with pro-
autophagic markers such as *ULK1* and *PINK1* during steatosis progression (**Figure**
4B). Further investigation of over 600 autophagy genes (Bordi et al., 2021)
(**Supplementary file 7** and **Figure 4–figure supplement 1B**) revealed active
autophagy initiation in early MASLD, where 119 genes across pathways including
mTORC and upstream effectors, lysosome, and autophagy core assembly genes
displayed altered expression levels. This suggests that autophagy serves as an
adaptive response to hepatic lipotoxicity during steatosis, but its activation may decline
as the disease progresses, as evidenced by the downregulation of key regulators such
as *NR1H4*, *SIRT5*, *FOS*, and *EGR1* (**Figure 4–figure supplement 1B**), thereby
impairing liver metabolism (Singh et al., 2009).

Molecular signatures of hepatic steatosis and fibrosis are mutually independent
To gain insights into the dual-omic data in the liver further, we performed a partial-
correlation network analysis to integrate liver transcriptomic, metabolomic, and clinical
data (Lee et al., 2025). The resulting subnetwork highlighted distinct molecular
signatures correlated with steatosis and fibrosis (**Figure 5A**). Steatosis was primarily
associated with hepatic neutral lipid accumulation and related metabolomic
alterations, whereas fibrosis was predominantly linked to transcriptional changes,

including the upregulation of genes involved in ECM remodeling (e.g., *TGFB3*,
FSTL1), signal transduction (e.g., *RHOU*, *DOK3*), and metabolism (e.g., *CYP2C19*,
PNPLA4, *SGPL1*). These findings underscore the presence of distinct molecular
pathways driving the progression of steatosis and fibrosis.
In our transcriptomics analysis, 992 and 1,251 gene markers were identified as
significantly associated with the histological features of steatosis and fibrosis,
respectively, with limited overlap between the two sets (**Figure 5B** and
Supplementary file 5, q value < 0.05), suggesting unique transcriptomic regulations
underlie each feature. Functional enrichment analysis further supported this
distinction, revealing largely non-overlapping biological processes associated with
each gene set (**Figure 5–figure supplement 1** and **Supplementary file 8**).
To distinguish gene signatures uniquely linked to steatosis or fibrosis, we
included the other histological feature as a covariate in the regression model. This
allowed us to define steatosis-specific genes (independent of fibrosis) and fibrosis-
specific genes (independent of steatosis) (**Supplementary file 9**). Since steatosis
precedes fibrosis in the pathophysiology of MASLD, enrichment analysis of steatosis-
and fibrosis-specific genes revealed a 'pseudo-temporal' progression of biological
processes (**Figure 5C**). More pathways were found to be associated with fibrosis than
with steatosis. Steatosis-specific upregulated genes were enriched in protein
glycosylation, inflammatory response, lipid catabolism, and lysosome organization. In
contrast, fibrosis-specific genes showed downregulation of various metabolic
pathways and upregulation of processes related to ECM remodeling, hypoxia,
signaling, and cell morphogenesis. Additionally, apoptosis-related pathways were
suppressed in steatosis but activated in fibrosis, while nucleotide metabolism showed
the opposite trend, suggesting dynamic regulation of cell death and proliferation during
MASLD development. These findings implicate distinct and lesion-specific gene
regulatory programs in early MASLD progression.
Moreover, consistent with previous observations showing that hepatic fibrosis
correlates with insulin resistance parameters in clinical assays (**Figure 1C**), we found
that individuals with diabetic MASLD exhibited a greater number of downregulated
genes as fibrosis progressed than non-diabetic MASLD individuals (**Supplementary
file 10 and Figure 5–figure supplement 2**). Since most of the suppressed genes in
the diabetic subgroup are involved in metabolism (e.g., *BAAT*, *G6PC1*, *SULT2A1*,
MAT1A), we hypothesize that diabetes may exacerbate the metabolic dysfunction
associated with hepatic fibrosis progression.

Gene signatures of fibrosis initiation
To further explore representative gene signatures in the development of fibrosis, we
identified 213 genes as progressive markers by comparisons of gene expression
levels against two different baselines (no MASLD and steatosis; details provided in
the methods; **Supplementary file 11**). Among them, 75 of these markers overlapped
with fibrosis-associated genes in the VA cohort and 35 overlapped with the EU cohort
(**Figure 6A**) (Hoang et al., 2019), resulting in 130 novel fibrosis markers. Pathway
analysis of these fibrosis markers highlighted prominent roles for signal transduction
and ECM organization/ disassembly, as expected (**Figure 6–figure supplement 1A**).
To infer the cell type-of-origin for signals from bulk sequencing, we mapped
progressive fibrosis markers onto a human liver single-cell atlas (MacParland et al.,
2018). Upregulated genes appeared to reflect changes from different cell types,
whereas downregulated markers were predominantly hepatocyte-specific (**Figure 6A**,
left panel). This suggests that in the early stages of liver fibrosis development, the

transcriptional programs are activated in different cell types, while hepatocyte
functions potentially become muted during the process. Meanwhile, we observed
fibrosis-associated upregulation of genes involved in TGF-β and SMAD signaling
cascades (*TGFB1*, *TGFB3*, and *INHBA*), ECM activators (*ITGAV*, *LOXL2*, *THBS1*,
MMP9, and *MMP14*) (Li et al., 2022; Munger et al., 1999; Wen et al., 2018), regulators
(*CDK8*, *CTDSP2*, *HDAC1*, and *HDAC7*), and downstream targets (*TIMP1*, *MMP9*, and
COL5A2) (**Figure 6–figure supplement 1B**). This aligns well with previous reports
that TGF-β signaling and HSC activation drive fibrosis during MASLD progression
(Kisseleva & Brenner, 2021; Massague & Sheppard, 2023).
Given that the transition from simple steatosis to the onset of fibrosis marks a
critical window for disease management, we compared gene expression profiles
between individuals with fibrosis accompanied by steatosis and those with steatosis
but without fibrosis (**Supplementary file 12**). Notably, the top-enriched pathways in
this comparison emerged as GTPase signaling and its regulation, signal transduction,
and innate immune response (**Figure 6B**) among other pathways. Moreover, we
identified 37 GTPase-related genes displaying significant associations with fibrosis
progression (**Figure 6–figure supplement 2A** and **Supplementary file 13**). To
explore the potential role of GTPase signaling in fibrosis, we examined an inventory
of 251 genes encoding GTPases, GTPase-activating proteins (GAPs), and guanine
nucleotide exchange factors (GEFs), analyzing their interactome using the Human
Cell Map (Go et al., 2021) and a protein-protein interaction network (Huttlin et al.,
2017; Razick et al., 2008).
Among the 6,971 proteins that have been tested for physical interaction or
subcellular proximity with GTPase-related genes, 508 of these associated significantly
with fibrosis progression (**Figure 6C**). The network indicates that
GTPases and their regulators interact with a wide range of fibrosis genes, with *ARF3*
and *RAC1* being central hubs of the network. Functional enrichment of the core nodes
revealed that the GTPase network regulates intracellular transport, actin cytoskeleton
organization, exocytosis, and other biological processes during fibrosis progression
(**Figure 6–figure supplement 2B**). Therefore, GTPases and their regulators are co-
regulated with fibrosis-related genes encoding their protein interaction partners,
supporting the likelihood of a functional link between GTPases and hepatic fibrosis.
This is supported by TGF-β genes positively correlated with co-expressed GTPase-
related genes (**Figure 6–figure supplement 2C**), suggesting a potential link between
TGF-β signaling and GTPase pathways.
To experimentally validate our findings, the altered expression of GTPase-
related genes was explored in an independent system using an established model of
3D liver spheroids in which hepatic cells remain viable and functional for multiple
weeks (Bell et al., 2016; Kemas et al., 2021; Vorrink et al., 2017). Specifically, we co-
cultured primary fully-differentiated human hepatocytes with Kupffer and stellate cells
isolated from adult patients with histologically confirmed MASH (Youhanna et al.,
2025) and examined the overall expression profile of GTPase-regulated genes
between our human cohort and the ex vivo system. The expression of GTPases were
largely concordant when assessing human patients and the 3D spheroid system
(**Figure 6D**). In comparison to the control group, spheroids from MASH patients
demonstrated an upregulation of 24 genes encoding GTPases and their regulators
(**Figure 6E**). Remarkably, the treatment of liver spheroids with elafibranor, a dual
PPARα/δ agonist (Ratziu et al., 2016), restored the expression levels of 17 (70%)
GTPase-related genes back to the baseline (**Figure 6E**). This independent

experimental system further suggests that GTPases may play a role in MASLD
pathogenesis and the fibrotic response.
Since HSCs are the main liver cell type responsible for activating fibrosis, we
next assessed whether inhibition of GTPase activity in an immortalized HSC cell line,
the LX-2 cells, could attenuate markers of fibrogenesis (**Figure 6–figure supplement
3A**). Selective GTPase inhibitors targeting Rac1 (NSC23766) and Cdc42 (ML141)
reduced the mRNA expression of *COL1A1* and *COL1A2*, as well as pro-collagen
secretion (**Figure 6–figure supplement 3**) under basal conditions. TGF-β1 mediated
activation induced the expected increase in collagen secretion and gene expression
which was attenuated by Rac1 inhibitor (NSC23766) and to some extent by ML141.
In addition, examination of previously published transcriptomic data of HSCs isolated
from CCL$_4$-mediated liver fibrosis in mice (De Smet et al., 2021), revealed the
upregulation of GTPases comparable to the steatosis-to-fibrosis transition in our
human cohort, with a temporal pattern aligning with hepatic collagen deposition
(**Figure 6–figure supplement 4A**). Further, in a human liver organoid model, TGF-β
induced increased expression of GTPase-related genes in hepatocytes and HSCs,
but not in fibroblasts (Hess et al., 2023) (**Figure 6–figure supplement 4B**), suggesting
a potential feedforward loop involving the TGF-β/GTPase axis between hepatocytes
and HSCs. To investigate intercellular crosstalk in GTPase regulation, we examined
key GTPase-related genes in LX-2 cells, hepatocyte monocultures, and spheroid co-
cultures (including hepatocytes, HSCs, Kupffer cells). As shown in **Figure 6–figure
supplement 5**, TGF-β1 induced a potential increase in *VAV1* and *DOCK2* consistent
in co-cultures, hepatocytes, and LX-2 cells, while *RAC1*, *RAB32*, and *RHOU* showed
cell type-specific responses. These findings indicate that multiple hepatic cell types
mediate GTPase regulation, underscoring intercellular crosstalk, which requires
further detailed investigation.
Overall, these additional experiments and analyses of datasets identified
upregulated GTPase-related genes during fibrosis initiation (**Figure 6–figure
supplement 4C**). However, further in-depth mechanistic studies are needed to
validate this association to determine how TGF-β is regulating GTPases and how
GTPase control the secretion of collagen, leading to fibrosis.

Discussion

In this study, we performed a comprehensive omics-based analysis of a cohort of 109
obese individuals with early MASLD. Through integrative dual-omics approaches, we
mapped the liver and plasma metabolomes and identified distinct hepatic molecular
features associated with steatosis and fibrosis progression. Notably, fibrosis was
closely connected to global reprogramming of hepatic gene expression, with GTPase-
related genes emerging as possible mediators of fibrosis initiation.
The strength of our study is the characterization of hepatic pathophysiology in
relation to steatosis and fibrosis via transcriptome and metabolome-wide variations.
Consistent with general expectation, our work highlighted altered hepatic lipid
metabolism in human liver tissues of obese patients at both metabolite and gene levels
(Liu et al., 2023; Radosavljevic et al., 2024; Saliba-Gustafsson et al., 2024). Consistent
with the variations in GLs, higher expression levels of *DGAT2*, *PNPLA3*, and *PLIN3*
were associated with steatosis progression. The higher expression of *DGAT2*, which
encodes a key enzyme catalyzing the last step of de novo TAG synthesis, implies
enhanced integration of TAGs into LDs in the endoplasmic reticulum, which
presumably alleviates lipid-induced ER stress and evades accumulation of lipotoxic
lipids (Scorletti & Carr, 2022). Upregulation of *PLIN3* was correlated with higher grades

of both fibrosis and steatosis. The protein encoded by *PLIN3*, along with other PLIN
proteins (Van Woerkom et al., 2024), plays an important role in LD stabilization and
the prevention of TAG hydrolysis, thus potentially contributing to hepatic steatosis
(Carr et al., 2012). In addition, although previous animal studies have reported
increased levels of sphingolipids in MASLD models (Babiy et al., 2023; McGlinchey et
al., 2022), we did not observe this in patients with obesity at early stages of the disease.
Sphingolipid alterations varied across cohorts with differing patient compositions
(McGlinchey et al., 2022; Ooi et al., 2021; Vvedenskaya et al., 2021), suggesting that
sphingolipid metabolism may be dependent on disease stage, obesity status, and
exhibit discrepancies between humans and mice.
Beyond lipid metabolism, other metabolic pathways and processes also
exhibited dysregulation at the dual-omics level. Specifically, with excessive lipid
deposition, we observed higher levels of taurine and vitamin E, both of which have
antioxidant effects and likely reflect the hepatic response to oxidative stress (Al-Baiaty
et al., 2021; Arroyave-Ospina et al., 2021; Wei et al., 2024). Importantly, amino acid
metabolism was markedly dysregulated as steatosis progressed, characterized by
elevated hepatic levels of serine, arginine, glutamic acid, glycine, and alanine, along
with decreased levels of the aromatic amino acids phenylalanine, tryptophan, and
tyrosine. A previous *in vitro* study also demonstrated that palmitic acid
supplementation disrupted the metabolism of the aromatic amino acids in PH5CH8
and HepG2 cells (Aggarwal et al., 2023). As microbiota-derived metabolites,
dysregulation in these amino acid may reflect disrupted gut function and may affect
the de novo synthesis of nicotinamide adenine dinucleotide in MASLD patients (Xue
et al., 2023). Moreover, we also observed evidence of autophagy activation concurrent
with intrahepatic lipid and cholesterol accumulation, wherein lysosome-associated
genes were uniquely associated with steatosis, as shown in **Figure 4B** and highlighted
in **Figure 5C** by the enrichment of lysosome organization among steatosis-specific
processes. This autophagic response most likely represents a protective mechanism
to ameliorate steatosis (Gual et al., 2017); however, the downregulation of autophagy
regulators during fibrosis development may exacerbate liver metabolic dysfunction.
As liver fibrosis progressed, the expression levels of genes involved in primary
bile acid biosynthesis, PUFA metabolism, and steroid hormone biosynthesis were
lower, especially *ACADS*, *ACADSB*, and *ACADM* which encode key enzymes in the
initial stage of catalyzing fatty acid β-oxidation, and *HADH* and *ACAA1*, which encode
enzymes that act in the late stage of β-oxidation (Adeva-Andany et al., 2019).
Moreover, mitochondrial-related genes were downregulated with fibrosis progression,
particularly those related to the electron transport chain of the mitochondrial inner
membrane. This may indicate that pathological oxidative stress and impaired ATP
production caused by early mitochondrial dysfunction may aggravate fatty liver
inflammation (Koliaki et al., 2015), and the augmentation of the inflammatory response
may become an important cause of liver fibrosis.
Through univariate hypothesis testing and partial correlation analysis, we
revealed distinct molecular signatures correlated with fibrosis and steatosis. Liver
fibrosis was associated with gene expression alteration, from which we identified over
200 progressive markers tracking its gradual advancement. Furthermore, GTPase-
related genes were dysregulated at the mRNA level with the emergence and
progression of fibrosis. Increased expression of GTPase regulators was independently
confirmed in 3D primary human liver spheroids, with elafibranor restoring their
expression to baseline levels (**Figure 6E**). The downregulated expression of GTPase

regulators following elafibranor treatment suggests a potential anti-fibrotic mechanism
involving GTPase signaling.
GTPase proteins are categorized into several subfamilies such as Rho-, Ras-,
and Arf-GTPase based on their sequence and structure (Gray et al., 2020). GAPs and
GEFs are the key regulators of GTPase activity that modulate intrinsic GTPase
functions and the GTP-bound state (Bos et al., 2007). Only a few studies have
elucidated the role of GTPases in liver pathology (Agarwal et al., 2023; Huang et al.,
2018; Peng et al., 2021; Schwerbel et al., 2020). For instance, *Rap1a* was identified
as a signaling molecule that suppresses both gluconeogenesis and hepatic steatosis
(Agarwal et al., 2023). The Rab23-specific GAP, *GP73*, has been implicated in
triggering non-obese MASLD (Peng et al., 2021). Additionally, Rho-GTPase signaling
through the ROCK1/AMPK axis regulates de novo lipogenesis during overnutrition
(Huang et al., 2018), while an immune-related GTPase triggers lipophagy and
prevents hepatic lipid storage (Schwerbel et al., 2020). However, evidence regarding
the role of GTPase regulation in human liver is limited, particularly in the context of
liver fibrosis initiation and progression, and a thorough mechanistic investigation is
warranted to establish causality.
In recent years, dual- and multi-omics strategies have gradually emerged in
MASLD research. A proteo-transcriptomic map constructed from the EU cohort's liver
RNA sequencing data and circulating proteome identified four promising protein
biomarkers (Govaere et al., 2023). Among these, *AKR1B10* was also identified in our
study as a key gene related to lipid metabolism, showing strong associations with both
steatosis and fibrosis (**Figure 3A**), which highlights its potential as a biomarker even
in the early stages of MASLD. Additionally, we observed mitochondrial dysregulation
linked to the progression of liver fibrosis (**Figure 4A**). Consistent with our findings, a
multi-omics study in obese individuals with MASLD also revealed mitochondrial
dysfunction at the hepatic protein level. The study further emphasized similar
mitochondrial disruptions in adipose tissue, suggesting the liver-adipose tissue
interaction in obesity-related MASLD (Castane et al., 2025). Beyond inter-tissue
interactions, Raverdy *et al.* reported the heterogeneity of MASLD molecular signatures
in relation to varying degrees of cardiovascular risk and type 2 diabetes (Raverdy et
al., 2024). This further highlights the impact of systemic metabolic conditions on
MASLD progression. In our analysis of diabetic individuals, we identified more
downregulated metabolic genes associated with fibrosis, supporting the notion that
insulin resistance may exacerbate metabolic dysregulation and interferes with the
disease trajectory.
Overall, our integrative investigation offers one of the most detailed molecular
landscapes of early stage MASLD in individuals with obesity, with comprehensive
descriptors of the early remodeling of hepatic metabolism associated with initiation of
fibrosis in the steatotic liver.

Limitations of the study

This study focused on 109 patients with early-stage MASLD, potentially overlooking
molecular changes associated with later disease stages. To address this, we cross-
referenced our findings with two external cohorts (EU and VA). Moreover, as the
results were based predominantly on participants without diabetes, their validity in
diabetic populations warrants additional evaluation since a potential association
between liver fibrosis and insulin resistance was observed. However, because of the
limited number of diabetic cases and the imbalanced distribution of fibrosis grades

between diabetic and non-diabetic groups, the interaction between fibrosis and
diabetes could not be evaluated in this study.
We identified GTPase-related genes as potential future targets for reversing
liver fibrosis. While preliminary validation has been conducted in different in vitro
systems, further investigation is required to confirm the causal role of specific
GTPases in fibrosis initiation especially in HSCs and to elucidate how GTPase
signaling contributes to collagen production throughout the progression of MASLD.
We used patient derived liver spheroids as an independent experimental
system to validate some of our data (see Figure 6D-E). This system has been
extensively validated (Bell et al., 2018; Bell et al., 2016; Bell et al., 2017; Messner et
al., 2018; Vorrink et al., 2017) and consists of adult patient-derived hepatocytes,
hepatic stellate cells, and Kupffer cells that are fully differentiated. These spheroids
are stable for several weeks in culture but as any in vitro culture system also has
disadvantages. Given that the spheroids consists of 3 cell types, we cannot determine
which cells are mainly driving the expression of the genes we are measuring. This is
demonstrated in **Figure 6–figure supplement 5,** where some genes display higher
expression in the Co-culture, in hepatocytes (Mono-culture), or LX-2 cells,
respectively. To understand better how these cell types signal to each other, additional
experiments will need to be performed.

Materials and Methods
Ethics statement
Participants provided written and verbal informed consent. The study protocol
conforms to the ethical guidelines of the 1975 Declaration of Helsinki and was
approved by the University of Melbourne Human Ethics Committee (ethics ID
1851533), The Avenue Hospital Human Research Ethics Committee (Ramsay Health;
ethics ID WD00006, HREC reference number 249), the Alfred Hospital Human
Research Ethics Committee (ethics ID GO00005), and Cabrini Hospital Human
Research Ethics Committees (ethics ID 09-31-08-15). All work on the data from human
samples in Sweden was approved by Etikprövningsmyndigheten (Dnr 2024-07917-
01).

Cohort recruitment
A detailed medical history was taken, and metabolic comorbidities were noted
including the presence of previously diagnosed hypertension and diabetes assessed
by oral glucose tolerance testing. Exclusion criteria included: age <18 years, previous
gender reassignment, other causes of chronic liver disease and/or hepatic steatosis
including Wilson's disease, α-1-antitrypsin deficiency, viral hepatitis, human
immunodeficiency virus, primary biliary cholangitis, autoimmune hepatitis, genetic iron
overload, hypo- or hyperthyroidism, coeliac disease, as well as recent (within three
months of screening visit) or concomitant use of agents known to cause hepatic
steatosis including corticosteroids, amiodarone, methotrexate, tamoxifen, valproic
acid and/or high dose oestrogens (Chalasani et al., 2017). Further exclusion criteria
included potential for alcohol-induced liver disease, which was assessed through a
modified version of the alcohol use disorders identification test (AUDIT) (Chalasani et
al., 2017; Saunders et al., 1993). Following histological assessment, patients with liver
slice weighing less than 2mg were excluded from the omics analysis.
Eligible patients with obesity scheduled for primary or secondary sleeve
gastrectomy, gastric bypass, or the insertion of a laparoscopic-adjustable band were
prospectively enrolled. All patients were fasted for 8-12 h overnight and venous blood

was taken before the induction of anaesthesia. Blood was transferred to 2 x K2E
Ethylenediaminetetraacetic acid (EDTA), 2 x SST™ II Advance and 1 x FX 5 mg bio-
containers for subsequent storage or clinical/biochemical assessments. All blood
samples were sent to Melbourne PathologyTM (Victoria, Australia) for standardised
measurement of biochemical and metabolic variables, except for one bio-container of
EDTA. Standard blood analyses were performed for electrolytes, full blood
examination, glucose, glycosylated haemoglobin, insulin, C-peptide, cholesterol,
triacylglycerols, and liver function assessed by ALT, AST, GGT, and alkaline
phosphatase (ALP), and screening blood tests for liver disease. The remaining blood
within the EDTA tube was spun at 8000 x g for 10 min and the plasma was collected
and stored at -80°C for mass spectrometry analyses.

Liver biopsy collection and histological feature assessment

An ~ 1 cm^3 wedge liver biopsy was collected from the left lobe of the liver during
surgery. All liver samples were collected between 8am and 1pm. The liver was cut into
two portions. One portion was placed in formalin and transported to TissuPath™
(Mount Waverley, Victoria), paraffin embedded and processed for histological
analysis. Samples were graded according to the Clinical Research Network NAFLD
activity score (NAS) (Brunt et al., 2011) and Kleiner classification of liver fibrosis
(Kleiner et al., 2005) by a liver pathologist at TissuPath. Patients with no MASLD were
defined by steatosis score of 0, regardless of inflammation or fibrosis grade 1. MASL
patients were defined by a steatosis score ≥1 with or without lobular inflammation. The
remaining portion of liver slices was used for bulk RNA sequencing and untargeted
metabolomics analysis.

Sample preparation for untargeted metabolomics

For liver tissue biopsies, each sample containing 50 ± 5 mg of tissue was extracted in
0.8 mL 50:50 (v/v) methanol:chloroform (Wu et al., 2019; Xu et al., 2016). Samples
were homogenized for 10 min at 25 Hz with a single 3-mm tungsten carbide bead per
tube. Separation of phases was achieved by the addition of 0.4 mL of water followed
by vortex-mixing and centrifugation (2400 g, 15 min, 4 °C). After separation, the upper
phase (the metabolite-containing fraction) and the lower phase (the lipid-containing
fraction) were transferred into separate tubes and dried using SpeedVac.
For plasma samples, 225 µL of methanol was added to 50 µl plasma, and the
mixture was vortexed for 10 sec (Lee et al., 2014). Subsequently, 450 µL chloroform
was added and the mixture was incubated for 1h in a shaker. To induce phase
separation, 187.5 µL water was added and the mixture was centrifuged at 12,000 rcf
for 15 min at 4 °C. The upper phase (the metabolite-containing fraction) phase and
lower (the lipid-containing fraction) were collected into separate fresh tubes and dried
using SpeedVac.
Dry extracts from the organic fraction were resuspended in the mixed solvent
of 90:10 isopropanol: acetonitrile. For the aqueous fraction, 90% acetonitrile was used
to reconstitute dry extracts. Plasma samples were reconstituted in 1:4 dilution
(plasma-to-solvent: 50 µL/200 µL), and the reconstitution volumes for liver biopsy
samples were corrected by the weights of liver slices (tissue-to-solvent: 50 mg/1000
µL). After reconstitution, samples were sonicated for 15 min and centrifuged at 12,000
g for 10 min at 4 °C.

Metabolomics data acquisition and processing

A 1 µL aliquot of the extract was subjected to LC-MS analysis using the Sciex
TripleTOF 6600 system coupled with the Agilent 1290 HPLC. For polar metabolite

analysis, mobile phase A was water with 10 mM ammonium formate and mobile phase
B was acetonitrile with 0.1% formic acid. Flow rate was 0.250 ml/min. The LC gradient
started at 3% A, increased to 30% A from 1 to 12 min, then to 90% A from 12 to 15
min, remained at 90% until 18.5 min, and returned to 3%, holding until the end of the
24-min run. For lipid analysis, mobile phase A was prepared by 60:40 water:
acetonitrile with 10 mM ammonium formate, and mobile phase B was made by 90:10
isopropanol: acetonitrile with 10 mM ammonium formate. Flow rate was 0.250 ml/min
and the total run time is 15.80 min. The LC gradient started at 80% A, decreased to
40% A at 2 min, further decreased to 0% from 2 to 12 min, remained at 0% A until 14
min, then returned to 80% A at 14.10 min, and held at 80% A until the end of run. Mass
spectrometry settings were as follows: gas1 50 V, gas2 60V, curtain gas 25V, source
temperature 500 °C, IonSpray voltage 5500 V. The collision energy was set to 30 V
and 45 V for polar metabolites and lipids, respectively. Fifteen precursors were
selected for MSMS fragmentation in the DDA acquisition. The mass range for the RP
method was 300 – 1000 m/z and for the HILIC method was 50 – 1000 m/z. In the
SWATH acquisition, fixed windows were applied with window widths of 21 Da for the
HILIC method and 10 Da for the RP method.
Each tissue type (i.e., plasma and liver) was analyzed as a separate batch.
Following sample reconstitution, the supernatants were collected into MS vials and
pooled as quality control (QC) samples for each tissue type. For each analytical batch,
patient samples were injected in a randomized sequence, along with one QC injection
approximately every 10 samples for monitoring instrument stability and calculating the
coefficient of variation (CoV). Both reverse phase (RP) and hydrophilic interaction
liquid chromatography (HILIC) columns were used for chromatographic separations of
organic and aqueous fractions, respectively. For the analysis of aqueous fraction,
metabolites were separated on SeQuant ZIC-cHILIC (3 µm, 100Å, 100x2.1 PEEK)
column with a flow rate of 0.25 mL/min in a 24-minute run. For the organic fraction,
metabolites were separated on RRHD Eclipse Plus C18 (2.1 x 50 mm, 1.8 µm).
Data acquisition was performed in both positive and negative ionization modes.
Data-dependent acquisition (DDA) was performed on pooled QC samples for
compound identification and SWATH-MS analysis was applied to individual samples
for relative quantification. The mass range for the RP method was 300 – 1000 m/z and
for the HILIC method was 50 – 1000 m/z. Data was converted into mzML format and
was then processed by MetaboKit (Narayanaswamy et al., 2020)
(https://github.com/MetaboKit). To gain broad metabolite coverage, we curated a data
processing pipeline for metabolites identified via spectral matches (MSMS-level, with
915 identifications postprocessing) and those identified solely through mass matches
(MS1-level, with 548 identifications postprocessing). For metabolite quantification, the
project-specific spectral library generated from DDA analysis was used to extract MS1
and MS2 features from the SWATH data using the MetaboKit DIA module. Compound
transitions with a CoV > 30% were excluded from the analysis. For each compound,
the precursor or fragment ion with the lowest CoV was selected as the quantifier. For
a compound identified in both positive and negative modes using the same column,
the peak feature with the lower CoV were selected. Overlapping metabolites in the RP
and HILIC datasets are mainly semi-polar lipids, as RP detects compounds with m/z
> 300. For lipids identified by both methods, the ones measured on the RP column
was selected.

RNA sequencing

Transcriptome sequencing service was provided by Biomarker Technologies (BMK),
GmbH. Briefly, RNA were extracted from liver biopsies using Trizol reagent. The
quality and quantity of extracted RNA was assessed by Nanodrop (quantity and purity)
and Labchip GX (Quality). The cDNA libraries were prepared using Hieff NGS Ultima
Dual-mode mRNA Library Prep Kit from Illumina (Yeasen) as per manufacturer's
instructions. RNA sequencing was performed on NoveSeq6000 platform (PE 150
mode). For raw data processing, FastQC (v0.11.9) was used as a quality control check
for raw sequencing. Alignment to the reference genome (GRCh38, Ensembl release
77) was performed using STAR (v2.7.9a), and gene expression levels (transcripts-
per-million, TPM) were estimated using RSEM (v1.3.1). Gene expression data
underwent a logarithmic transformation to $log2(TPM + 1)$ and mean-centering
normalization for downstream analyses.

Statistical analysis
All analyses in this study were conducted using R version 4.3.1. Linear regression
model was used to evaluate the linear association between histological grades and
molecular abundance levels with adjustment for age, sex, BMI, and diabetes.
Statistically significantly changed metabolites or genes were defined by controlling the
overall type I error at 5% or 10% (q value < 0.05 or 0.1). Spearman rank correlation
was used to assess the relationship between the same metabolite in the liver and
plasma. This accounts for matrix effects across different tissue types in untargeted
analysis by comparing the relative intensity ranking of a metabolite across all samples
within each tissue type. Gene overrepresentation analyses were performed using
gene ontology recourse by in-house software (Gene Ontology et al., 2023). ClueGO,
a plugin app in Cytoscape, was used to visualize enrichment maps (Bindea et al.,
2009). Partial correlation network analysis was employed to integrate the two omics
layers using an in-house partial correlation network algorithm ACCORD (Lee et al.,
2025). KEGG pathway database was used for the analysis of lipid metabolism and
pathway mapping (Kanehisa et al., 2016). All network visualizations in this study were
done using Cytoscape (Otasek et al., 2019).
To investigate MASLD subgroup signatures related to diabetic status, we
analyzed linear associations between gene signatures and histological features
separately in non-diabetic (n = 71) and diabetic individuals (n = 23). Statistical power
was estimated by comparing variance explained in full (y ~ x + a + b + c) versus
reduced (y ~ a + b + c) regression models, converting the incremental R^2 into Cohen's
f^2, and applying pwr.f2.test at α = 0.05. We further compared gene expression profiles
between diabetic (n = 21) and non-diabetic (n = 43) MASLD patients, identifying 166
differentially expressed genes ($p < 0.05$, $|log_2FC| > 0.32$). Of these, 54 genes
(**Supplementary file 10**) were both differentially expressed and significantly
associated with fibrosis progression, and thus marked as signatures in diabetic
MASLD.
To identify progressive markers of liver fibrosis, we performed pairwise
analyses for fibrosis stages using two distinct baselines: patients with no MASLD and
those with steatosis but no fibrosis. 213 markers were characterized as progressive
based on the following criteria: a) significant linear association with fibrosis grades,
and b) significant differential expression (p value < 0.05 in t-test) in at least two fibrosis
stages (e.g., F1 and F2, F2 and F3) compared to both baselines.
To identify independent gene signatures associated specifically with steatosis
and fibrosis progression, we applied linear regression models for each outcome
adjusting for the other. Steatosis-specific signatures were defined as genes

significantly associated with steatosis but not with fibrosis, and vice versa. Metascape
was used for meta-analysis of enrichment pathways (Zhou et al., 2019). Pathways
with at least 10 hits and p value < 0.05 were considered steatosis- or fibrosis-specific.
For meta-analysis of reference human cohorts, the two RNA-seq datasets of
the liver transcriptome, the VA cohort (GSE130970) and the EU cohort (GSE135251),
were downloaded from Gene Expression Omnibus, and associations between gene
expression (log2 TPM) and fibrosis grades or NAS were analyzed using linear
regression. In the EU cohort, the patients without MASH diagnosis were removed from
the meta-analysis for the purpose of identifying fibrosis signatures in MASH patients.
Human liver single cell dataset was obtained from GSE115469 (MacParland et al.,
2018). The mean log2 TPM was calculated for each cell type, and cell specificity was
determined by subtracting the mean expression levels across all cell types.

Cross-referencing datasets of mouse models and human liver organoids
The RNA sequencing data (raw counts) for HSCs isolated from liver fibrosis mouse
models (De Smet et al., 2021) was downloaded from GSE176042 and analyzed using
the DESeq2 R package. This previously published study investigated HSC initiation
and perpetuation using acute and chronic models, respectively. For the acute model,
mice were injected with CCl_4 once and samples were taken at 24 h, 72 h, and 1 w. For
the chronic model, mice received a regimen of semi-weekly injection of CCl_4 for 4
weeks. After this regimen, mice were sacrificed and samples were taken at 24 h, 72
h, and 2 w. The statistical results from single cell dataset of stem cell-derived human
liver organoids were obtained from GSE207889 (Hess et al., 2023).

3D liver spheroid MASH model
3D liver spheroids were generated based on co-cultures of cryopreserved primary
human hepatocytes (PHH) and liver non-parenchymal cells (NPC) as previously
described (Youhanna et al., 2025). Briefly, cells were seeded at a PHH:NPC ratio of
6:1 in 96-well ultra-low attachment plates (CLS7007-24EA Corning) with a total of
1500 cells/well in culture medium (William's E medium containing 11 mM glucose, 100
nM dexamethasone, 10 µg/mL insulin, 5.5 mg/L transferrin, 6.7 µg/L selenite, 2 mM
L-glutamine, 100 U/mL penicillin, 0.1 mg/mL streptomycin) supplemented with 10%
FBS. After spheroid formation FBS was phased out and cells were maintained in
medium supplemented with albumin-conjugated free-fatty acids (FFA) including 80µM
palmitic acid and 80 µM oleic acid. At the end of the experiment, spheroids were
collected for RNA extraction (Zymo R105). The elafibranor treatment protocol was
described in the previous study (Youhanna et al., 2025). Each experimental condition
(control, MASH, and elafibranor treatment) was performed in quadruplicate. The
functional characterization of hepatocytes in the liver spheroid culture model has been
published previously (Bell et al., 2018; Bell et al., 2016; Bell et al., 2017; Messner et
al., 2018; Vorrink et al., 2017).
40

1 **Table 1. Patient characteristics**

	Patients (N)	No MASLD (N = 33)	MASLD (N = 76)	P value
Patient information				
Age (years), median ± MAD	109	36 (14.8)	41.5 (9.6)	0.942
Sex (male), n (%)	108	4 (12.1)	22 (29.3)	0.092
Diabetes, n (%)	109	2 (6.1)	27 (35.5)	0.003
Hypertension, n (%)	93	6 (18.2)	16 (26.7)	0.505
BMI (kg/m2), median ± MAD	108	42.6 (6.6)	44.5 (9.1)	0.045
Clinical chemistry parameters				
ALT (U/L), median ± MAD	104	24 (11.9)	42 (25.2)	0.507
AST (U/L), median ± MAD	104	24 (7.4)	30 (13.3)	0.39
ALT/AST, median ± MAD	104	1 (0.3)	1.33 (0.4)	0.005
GGT (U/L), median ± MAD	104	23 (10.4)	28 (13.3)	0.041
Total cholesterol (mmol/L), median ± MAD	104	4 (0.9)	4.15 (1.0)	0.072
HDL (mmol/L), median ± MAD	102	1.04 (0.2)	0.94 (0.2)	0.053
LDL (mmol/L), median ± MAD	101	2.2 (0.7)	2.6 (0.7)	0.081
Non-HDL cholesterol (mmol/L), median ± MAD	108	2.76 (0.9)	3.27 (1.0)	0.003
Blood triglyceride (mmol/L), median ± MAD	103	1.2 (0.6)	1.5 (0.6)	0.013
Insulin (mU/L), median ± MAD	100	7.95 (5.6)	10.35 (7.1)	0.043
FBG (mmol/L), median ± MAD	102	4.9 (0.7)	5.2 (1.0)	0.114
C-peptide (nmol/L), median ± MAD	101	0.8 (0.4)	1.12 (0.4)	0.002
HOMA2–IR, median ± MAD	92	1.13 (0.7)	1.43 (0.9)	0.034
Liver histology				
Steatosis (S0/S1/S2/S3)	109	33/0/0/0	0/41/27/8	<0.001
Inflammation (I0/I1/I2/I3)	109	30/3/0/0	21/46/7/2	<0.001
Ballooning (B0/B1/B2)	109	33/0/0	65/10/1	0.07
NAS (N0/N1/N2/N3/ N4/N5/N6/N7)	109	30/3/0/0/ 0/0/0/0	0/15/29/13/ 14/2/2/1	<0.001
Fibrosis (F0/F1/F2/F3)	109	27/6/0/0	23/27/20/6	<0.001

Additional patient data is available in Supplementary file 1.
MASLD: Metabolic dysfunction-associated steatotic liver disease; MAD: Median
absolute deviation; BMI: Body mass index; ALT: Alanine transaminase; AST:
Aspartate aminotransferase; GGT: Gamma-glutamyl transpeptidase; HOMA2 – IR:
Homeostasis model assessment 2 of insulin resistance; HDL: High-density lipoprotein;
LDL: Low-density lipoprotein; FBG: Fasting blood glucose; NAS: Nonalcoholic fatty
liver disease activity score.
9

Figure legends

Figure 1: Study overview. (A) The overall study design. (B) Alluvial diagram of patient
composition of groupings across liver histological features. (C) Relationship between
clinical test results and stages of liver histological features. Node size was determined
by –log10(p value) in ANOVA tests. Color indicates the significance degree [–log10(p
value)] and the direction of change from early to late stages. (D) Number of
metabolites analyzed in the liver and plasma. (E-F) Principal component analysis of
liver metabolome, plasma metabolome, and liver transcriptome. (G) Disease spectra
covered in this cohort and comparison to two published datasets. All source data are
available in Supplementary file 2.

Figure 2: Hepatic and circulating metabolome in obese individuals with MASLD.
(A) Number of metabolites in each class that were significantly associated with
histological features in the liver and plasma (q value < 0.05). Metabolite classes with
at least 5 metabolites with significance are shown in the plot. (B) Associations between
steatosis grades and lipid species in each lipid class. Dots were colored by double
bond numbers (left) and carbon numbers (right) of lipid species, respectively. (C)
Partial correlation network of the plasma metabolome and clinical covariates. Node
size reflects the degree of connectivity, with larger nodes indicating connections to a
greater number of metabolite nodes. (D) Heatmaps for pairwise analysis between
plasma lipid classes and clinical variables. Linear regression analysis was performed
for numerical variables, while logistic regression was conducted for binary variables.
Significance values refer to –log10(p value)*sign(coefficients) from regression models.
(E) Comparisons of liver metabolome and plasma metabolome regarding their
associations with liver steatosis and fibrosis. All source data are available in
Supplementary file 3 and 4.

Figure 2–figure supplement 1. Metabolomic analysis in this MASLD cohort. (A)
Plasma urea and creatinine from untargeted metabolomics correlated with clinical
assays. (B) Liver and plasma metabolites associated with the disease progression in
each matrix (q < 0.05 or q < 0.1). Compound classes with at least 5 hits associated
with any histological feature were shown in the plot. (C) Associations between fatty
acid composition of triglycerides (TAGs) and histological outcomes. Significance
values refer to –log10(p value)*sign(coefficient) from linear regression models.

Figure 2–figure supplement 2. Blood lipoprotein cholesterol levels in patients
with different steatosis and fibrosis grades. HDL, high-density lipoprotein
cholesterol. LDL, low-density lipoprotein cholesterol. NonHDL-Chol, non-high-density
lipoprotein cholesterol. TotalChol, total blood lipoprotein cholesterol.

Figure 3: Integrative view of key metabolic pathways implicated in liver
metabolism in obese individuals with MASLD. (A) Heatmap of log2 fold changes
from pair-wise analysis of lipid metabolism-related genes. Association with steatosis
(S) and fibrosis (F) is indicated by black dots (q value < 0.05, top). Results were cross-
referenced with two published cohorts (VA cohort, GSE130970; EU cohort,
GSE135251, bottom). Asterisks indicate genes with a q-value < 0.05 and a consistent
change in direction within our cohort. (B) KEGG metabolic pathways with at least 4
genes or metabolites significantly associated with steatosis [S] or fibrosis [F]. (C)
Integrative map of gene and metabolite alterations associated with steatosis (left half

of each box/circle) and fibrosis (right half of each box/circle). (D) Metabolite alterations
corresponding to the advancement of steatosis grades (x-axis). All source data are
available in Supplementary file 4 and 5.
4
Figure 3–figure supplement 1. Integrative map of liver metabolism in obese MASLD
patients. (A) An integrative map of transcriptomics and metabolomics of glycerolipid
and glycerophospholipid metabolism in early MASLD. (B) Gene expression changes
in fatty acid β-oxidation, mitochondrial respiratory chain, and lipid droplet (LD)
metabolism associated with steatosis and fibrosis. There was a notable
downregulation of genes involved in the electron transport chain of the inner
mitochondrial membrane, specifically in complex I (NDUF family), complex II (SDH
family), complex III (UQCR), complex IV (COX), and complex V (ATP5 family) as
fibrosis progressed. Genes involved in respiratory electron transport were colored by
the significance [–log10(p value)*sign(coefficient)] in relation to fibrosis grades. Inset:
gene expression changes in the electron transport chain are depicted.
16
17 **Figure 3–figure supplement 2**. Ratios of metabolites in one-carbon metabolism in
18 individuals with different steatosis grades. SAM, S-adenosylmethionine. SAH, S-
19 adenosylhomocysteine.
20
Figure 4: Dysregulated mitochondrial function and autophagy during the
progression of steatosis and fibrosis. Gene expression patterns related to
mitochondrial metabolism (A), mitochondria-related biological processes (A),
autophagy activation (B), and lysosomal biogenesis (B) with statistical association to
steatosis and fibrosis. S, steatosis. F, fibrosis. All source data are available in
Supplementary file 6 and 7.
27
Figure 4–figure supplement 1. Autophagy regulation in the liver. (A) Hepatic free
cholesterol, CE 18:1, and CE 18:2 levels in this cohort in relation to steatosis grade
(x-axis). (B) Heatmap of autophagy genes in relation to histological features, indicating
active lipophagy in the liver of obese individuals. Asterisks in the main heatmap denote
the significance (p < 0.05) from comparisons of each subgroup. Association with
steatosis (S), NAS (N) and fibrosis (F) was indicated by black dots (q < 0.05, left).
Results were cross-referenced with two published cohorts (VA cohort, GSE130970;
EU cohort, GSE135251) (*q < 0.05 and with consistent direction of changes as in our
cohort, bottom).
37
Figure 5: Steatosis and fibrosis as independent processes. (A) Network of genes,
metabolites, and histological features (fibrosis and steatosis) using partial correlation
analysis as described in the methods. (B) Venn diagrams depicting the number of
genes significantly associated with steatosis and fibrosis (q value < 0.05) in the liver
transcriptome. (C) Gene ontology (GO) enrichment of gene sets specific to steatosis
or fibrosis. All source data are available in Supplementary file 8, 9, and 10.
44
45 **Figure 5–figure supplement 1. Functional enrichment of gene sets associated**
46 **with steatosis and fibrosis in the liver transcriptome.** Node size is depicted by
47 circles and P value by color.
48
Figure 5–figure supplement 2. Statistical power and subgroup analysis of
associations between gene expressions and steatosis or fibrosis grades.

Statistical power in the original linear regression analysis of steatosis and fibrosis was
plotted against p-values (A). Differential features associated with steatosis ($q < 0.05$)
showed power > 0.8, whereas those associated with fibrosis showed power > 0.7.
Subgroup analyses of participants with and without diabetes were conducted to
evaluate associations between gene expression and steatosis (B) and fibrosis (C).
The x-axis shows coefficients from the original analysis including all participants.
Corresponding coefficients from the analyses of non-diabetic (left panel) and diabetic
(middle-left panel) individuals are displayed alongside the statistical power for each
subgroup analysis (middle-right and right panels). Red dots represent genes that were
significant in the original analysis ($q < 0.05$). Subgroup analyses suggest that the
results in the non-diabetic subgroup (n = 71) were highly consistent with findings from
the original analysis (n = 94, adjusted for diabetes), indicating that the originally
reported gene signatures, after correction for diabetic status, remain valid in non-
diabetic individuals.
15
Figure 6: Liver fibrosis signatures and potential therapeutic targets of fibrosis
initiation. (A) Top 60 fibrosis markers in the liver. A total of 213 genes were identified,
from which the top 30 upregulated and top 30 downregulated genes are visualized.
The middle plot (blue bar on top) shows the comparison of individuals with fibrosis and
without pathology (baseline - "no MASLD"). For the right plot (red bar), individuals with
fibrosis were compared to those with steatosis but no fibrosis (baseline – "steatosis
but no fibrosis"). Results were cross-referenced with two published cohorts (VA cohort,
GSE130970; EU cohort, GSE135251, right). Asterisks indicate genes with a q-value
< 0.05 and a consistent change in direction within our cohort. The average gene
expressions of liver cell types were obtained from GSE115469 at the log2CPM level
(left). Dots in the single cell map indicate zero expression in the corresponding cell
types. (B) Enrichment of pathways in the transition from simple steatosis to the onset
of fibrosis. (C) Protein-protein interaction network of GTPases and their regulators.
Nodes are colored based on gene type, with borders indicating the direction of gene
regulation. Node size corresponds to the significance of the genes in relation to fibrosis
grades within this cohort. GEFs: Guanine nucleotide exchange factors. GAPs:
GTPase-activating proteins. (D) Comparison of expression level changes in GTPase-
related genes between this human cohort and an independent 3D spheroid MASH
system. Log2 fold change for the human cohort was calculated by comparing patients
with grade 2 or 3 fibrosis to those without fibrosis or steatosis. Genes with the same
direction of change are colored in red, while others are colored in purple. (E)
Expression of GTPase-related genes in patient-derived 3D liver spheroids: control
spheroids, spheroids from patients with MASH, and Elafibranor-treated MASH
spheroids (n = 4). Sixty-eight genes with a p-value < 0.05 from the ANOVA test were
plotted in the diagram, with blue lines highlighting 24 genes that exhibited increased
expression in the MASH group compared to the control group. All source data are
available in Supplementary file 11, 12, and 13.
43
44   **Figure 6–figure supplement 1. Liver fibrosis pathways and gene signatures.** (A)
45   Pathway analysis of the enrichment of pathways for the 213 fibrosis gene signatures.
46   (B) TGF-β and SMAD related gene expression associated with fibrosis grades (x-axis)
47   is depicted. Significance is indicated by black dots; • $p < 0.05$, •• $p < 0.01$, ••• $p <$
48   0.001.
49

Figure 6–figure supplement 2. GTPases and their regulation emerges as a
potential target for liver fibrosis. (A) Heatmap of GTPase-related genes display a
better correlation with fibrosis grades. Asterisks in the main heatmap denote the
significance ($p < 0.05$) from comparisons of each subgroup. (B) Enrichment map of
selected genes constituting the core of the protein-protein interaction network of
GTPase-related genes. Nodes are colored according to BH (Benjamini-Hochberg)
adjusted p-value, and node sizes are proportional to the number of genes in the term.
(C) Correlation matrix of gene expression levels for TGF-β and GTPase-related genes.
The correlation plot was created of Pearson's r values for GTPase-related genes and
TGF-β genes (TGFB1 and TGFB3). The left annotation bar indicates the significance
of associations between gene expression and fibrosis grades.
12
Figure 6–figure supplement 3. GTPase inhibition and Elafibranor treatment in
HSC-derived LX2 cells. (A) Workflow of the LX2 experiment in panel A-F (n = 8-10).
(B) GTPase inhibitors NSC23766 (Rac1 inhibitor) and ML141 (Cdc42 inhibitor)
significantly reduced pro-collagen secretion from HSC-like LX-2 cells and (C)
decreased gene expression of COL1A1 and COL1A2 under basal conditions. (D)
TGF-β administration increased pro-collagen secretion by 32% and (E) doubled
collagen gene expression in LX2 cells. (F) NSC23766-mediated GTPase inhibition
impairs pro-collagen secretion from LX-2 cells after TGF-β treatment. Pro-collagen
secretion was determined by ELISA and gene expression levels were assessed by
qPCR. Asterisks (*) denote p value < 0.05 for statistical significance from one-way
ANOVA with Holms-Sidak multiple comparisons (B&C), unpaired t-test (D&E) Kruskal
Wallis Test with Dunns multiple comparisons (F). (G) Gene expression of selected
GTPases in LX-2 cells with and without Elafibranor (n = 3).
26
Figure 6–figure supplement 4. GTPase-related genes in external systems. (A)
Human fibrosis signatures from the present study were partially reflected in a mouse
study (GSE176042), where RNA sequencing was performed on hepatic stellate cells
(HSCs) isolated from both acute and chronic mouse fibrosis models. GTPase-related
genes, especially those involved in the initiation of fibrosis, were also upregulated in
HSCs isolated from both acute and chronic liver fibrosis mouse models at different
time points. Row annotations 'Linear' and 'Transition' indicate whether gene
expression is linearly associated with fibrosis grades (Linear = 1) or shows a significant
difference between patients with fibrosis and steatosis versus simple steatosis
(Transition = 1). Asterisk denotes genes with adjusted p value < 0.05. (B) GTPase-
related genes induced by TGF-β were elevated in hepatocytes and HSCs at different
stages of a human liver organoid system (GSE207889). Significance was calculated
by –log10(p value)*sign(fold change). AH, adult hepatocyte-like; cAH, cycling adult
hepatocyte-like; CHOL, cholangiocyte-like; DC, ductal cell-like; FH1, fetal hepatocyte
1-like; FIB, fibroblast-like; HB1, hepatoblast 1-like; HB2, hepatoblast 2-like; HSC,
hepatic stellate cell-like; SMC, smooth muscle cell-like. (C) Schematic illustration of
the hypothesized role of GTPase signaling in regulating liver fibrogenesis. TGF-β
production is initiated in non-parenchymal liver cells, leading to the overexpression of
GTPase regulators in both hepatic stellate cells and hepatocytes. This activation of
GTPase signaling amplifies downstream fibrogenic effects, creating a feed-forward
loop that intensifies fibrosis progression. Further investigation is needed to prove by
which mechanisms GTPases regulate collagen expression and therefore fibrosis.
49

**Figure 6–figure supplement 5. Expression of GTPase-related genes in spheroid
co-culture, hepatocyte monoculture, and LX-2 cells.** RNA-sequencing-based
expression of GTPase-related genes in spheroid co-culture, hepatocyte monoculture
upon control (Ctrl), TGF-β1 or TGFβ – inhibitor treatment (A, B). Expression of GTPase-
related genes in control (Ctrl) and TGF-β1- treated (24h) LX2 cells measured by qPCR
(C). Asterisks (*) denote p value < 0.05, (**) $p < 0.005$, (***) $p < 0.0001$ for statistical
significance from unpaired t-test, numerical values above the bars indicate p-value. Bars
were removed where data did not show strong directionality and statistical reliability, as
indicated by (^) symbol. Symbol (@) indicates one or more expression values were not
detected in the original dataset. Please note that ACTA2 expression in LX-2 cells is time
dependent (unpublished results) and therefore we also included fibronectin (FN1) as a
control for TGF-β1 treatment.

Acknowledgments

14 This manuscript is the result of a wonderful international collaboration and we thank
15 all present and past members of the Watt, Choi, Lauschke, and Kaldis laboratories for
16 discussions, input, and support. P.K. thanks Åke Nilsson for discussions and
17 suggestions.

Abbreviations

20	ALT	Alanine aminotransferase
21	AST	Aspartate aminotransferase
22	ALP	Alkaline phosphatase
23	CAR	Acylcarnitines
24	CCl_4	Carbon tetrachloride
25	Cer	Ceramide
26	CM	Conditioned medium
27	CYP	Cytochrome P
28	DAG/DG	Diacylglycerol
29	DDA	Data-dependent acquisition
30	ECM	Extracellular matrix
31	FBS	Fetal bovine serum
32	FFA	Free-fatty acids
33	FXR	Farnesoid X receptor
34	GAP	GTPase-activating protein
35	GEF	Guanine nucleotide exchange factor
36	GGT	Gamma-glutamyltransferase
37	GL	Glycerolipid
38	GPL	Glycerophospholipid
39	HSC	Hepatic stellate cell
40	HILIC	Hydrophilic interaction liquid chromatography
41	LD	Lipid droplet
42	LNAPE	N-acyl-lysophosphatidylethanolamine
43	LPC-O	Ether-linked lysophosphatidylcholine
44	LPI	Ether-linked lysophosphatidylinositol
45	LPS	Ether-linked lysophosphatidylserine
46	LPC	Lysophosphatidylcholine
47	LPE	Lysophosphatidylethanolamine
48	MG	Monoacylglycerol
49	MASH	Metabolic dysfunction-associated steatohepatitis
50	MASLD	Metabolic dysfunction-associated steatotic liver disease
51	NAS	NAFLD activity score

1	NPC	Non-parenchymal cells
2	NAE	N-acyl ethanolamines
3	PCA	Principal component analysis
4	PHH	Primary human hepatocytes
5	PUFA	Polyunsaturated fatty acid
6	PC	Phosphatidylcholine
7	PC-O	Ether-linked phosphatidylcholine
8	PE	Phosphatidylethanolamine
9	PE-O	Ether-linked phosphatidylethanolamine
10	PG	Phosphatidylglycerol
11	PI	Phosphatidylinositol
12	PS	Phosphatidylserine
13	QC	Quality control
14	RP	Reverse phase
15	SM	Sphingomyelin
16	SL	Sphingolipid
17	SPB	Sphingoid base
18	SAM	S-adenosylmethionine
19	SAH	S-adenosylhomocysteine
20	TAG/TG	Triacylglycerol/Triglyceride
21	TAG-O	Ether-linked triacylglycerol
22	Vitamin E	Alpha-tocopherol

Conflict of interest statement:

VML is co-founder, CEO and shareholder of HepaPredict AB, as well as co-founder and shareholder of Shanghai Hepo Biotechnology Ltd. All other authors declare no conflict of interest.

Additional files

Supplementary files

Supplementary file 1. Additional patient characteristics.

Supplementary file 2. Top-enriched pathways of genes with nonzero loading scores on PC1 and PC2 in the sparse PCA of the liver transcriptome.

Supplementary file 3. Statistical analysis of metabolomics data for plasma samples.

Supplementary file 4. Statistical analysis of metabolomics data for liver samples.

Supplementary file 5. Statistical analysis of transcriptomics data for liver samples

Supplementary file 6. Associations between mitochondrial function-related genes and liver histological grades.

Supplementary file 7. Associations between autophagy-related genes and liver histological grades.

Supplementary file 8. Over-representation analysis enrichment analysis of genes linearly associated with steatosis or fibrosis grades.

Supplementary file 9. Steatosis-specific and fibrosis-specific gene signatures

Supplementary file 10. Subgroup statistical analysis of liver transcriptome in diabetic and non-diabetic individuals.

Supplementary file 11. Progressive gene markers associated with liver fibrosis in MASLD patients.

Supplementary file 12. Genes involved in the transition from fibrosis-free steatosis to fibrosis.

Supplementary file 13. Associations between GTPase-related genes and liver histological grades.

Appendix 1 – supplementary methods

Appendix 2 – discussion of plasma metabolome

Data availability

All data are available within the manuscript, supplementary figures, and

supplementary tables. RNA sequencing data to this article have been submitted to

SRA (PRJNA1185558) and is deposited with GEO (GSE281797). Untargeted LC-

HRMS data is deposited with Zenodo, including IDA data (DOI

10.5281/zenodo.14091962), SWATH data for the organic fraction (DOI

10.5281/zenodo.14096635), and SWATH data for the aqueous fraction (DOI

10.5281/zenodo.14096753, DOI 10.5281/zenodo.14136832). All clinical data,

processed omics datasets, and code are available at

https://github.com/SLINGhub/MASLD_dual_omics.

The following datasets were generated:

Author(s)	Year	Dataset title	Dataset URL	Dataset and identifier
Zhao Q, De Nardo W, Choi H, Watt MJ, Kaldis P	2025	Transcriptomic profiles of liver biopsies in obese patients with metabolic dysfunction-associated steatotic liver disease	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE281797	NCBI Gene Expression Omnibus, GSE281797
Zhao Q, De Nardo W, Choi H, Watt MJ, Kaldis P	2024	Untargeted LC-HRMS-based metabolomic profiling reveals distinct metabolomic profiles in patients with metabolic dysfunction-associated steatotic liver disease (MASLD)	https://zenodo.org/records/14091963	Zenodo DOI 10.5281/zenodo.14091962, DOI 10.5281/zenodo.14096635, DOI 10.5281/zenodo.14096753, DOI 10.5281/zenodo.14136832

Financial support statement:

PK is supported in part by the Novo Nordisk Foundation (NNF24OC0092365),

Swedish Research Council (2021-01331), the Swedish Cancer Society

(Cancerfonden; 21-1566Pj and 24-3605Pj), the Crafoord Foundation (Ref. No.

20220628), the Swedish Foundation for Strategic Research Dnr IRC15-0067, and

Swedish Research Council, Strategic Research Area EXODIAB, Dnr 2009-1039. LNZ

is supported in part by the IngaBritt och Arne Lundbergs Forskningsstiftelse LU2020-

0013, the Crafoord Foundation (Ref. No. 20210516, Ref. No. 20220582), Stiftelsen

Längmanska kulturfonden (BA21-0148), and the Åke Wibergs Stiftelse. VML

acknowledges support from the Swedish Research Council (2021-02801 and 2023-

03015), the Ruth och Richard Julins Foundation for Gastroenterology (2021-00158),

Knut and Alice Wallenberg Foundation (VC-2021-0026), by the SciLifeLab and

Wallenberg National Program for Data-Driven Life Science (WASPDDLS22:006), a

Novo Nordisk Pioneer Innovator Grant (NNF23OC0085944), the Robert Bosch

Foundation, as well as the Innovative Medicines Initiative 2 Joint Undertaking (JU)

under grant agreement No. 875510. The JU receives support from the European

Union's Horizon 2020 research and innovation programme and EFPIA, Ontario
Institute for Cancer Research, Royal Institution for the Advancement of Learning
McGill University, Kungliga Tekniska Hoegskolan, and Diamond Light Source Limited.
HC is supported in part by Singapore Ministry of Education (T2EP202121-0016,
T2EP20223-0010), National Research Foundation and Agency for Science,
Technology and Research (I1901E0040), and National Medical Research Council
(CG21APR1008). MJW is supported by the National Health and Medical Research
Council of Australia (APP1162511). The funders had no role in study design, data
collection and interpretation, or the decision to submit the work for publication.
10
Author contributions:
QZ*: designing research studies, conducting experiments, acquiring data, analyzing
data, writing the manuscript
WD*: designing research studies, conducting experiments, acquiring data, analyzing
data, writing the manuscript
RW*: analyzing data, writing the manuscript
YZ: conducting experiments, acquiring data, analyzing data
UK: analyzing data, writing the manuscript
GS: conducting experiments, acquiring data, analyzing data
LNZ: analyzing data, writing the manuscript
HT: conducting experiments, acquiring data, analyzing data
SY: conducting experiments, acquiring data, analyzing data
MY: conducting experiments, acquiring data, analyzing data
YX: conducting experiments, acquiring data, analyzing data
YK: conducting experiments, acquiring data, analyzing data
SL: conducting experiments, acquiring data, analyzing data
RL: conducting experiments, acquiring data, analyzing data
GT: conducting experiments, acquiring data, analyzing data
PN: conducting experiments, acquiring data, analyzing data
PRB: providing reagents, cohort management
VMK: designing research studies, analyzing data, writing the manuscript, supervision,
funding acquisition
HC: designing research studies, analyzing data, writing the manuscript, supervision,
funding acquisition
MJW: designing research studies, analyzing data, writing the manuscript, supervision,
funding acquisition
PK: designing research studies, analyzing data, writing the manuscript, supervision,
funding acquisition, leading of collaboration
39
QZ, WD, and RW are co-first authors based on their contributions, which was writing
original draft, LC-MS data acquisition and bioinformatic analysis of transcriptomic,
metabolomic, and clinical data (QZ), organizing cohort samples, clinical parameters,
managing, and LX-2 experiments (WD), and writing the original draft and analyzing
data (RW).

ORCID
2
Qing Zhao[1,2,3*]   https://orcid.org/0000-0003-1038-1209
William de Nardo[4*]   https://orcid.org/0000-0002-1441-1295
Ruoyu Wang[5,10]   https://orcid.org/0009-0003-5579-0492
Yi Zhong[7]   https://orcid.org/0000-0002-0929-8224
Umur Keles[10]   https://orcid.org/0000-0002-6771-3721
Gabrielé Sakalauskaite[10] https://orcid.org/0009-0002-5066-6663
Li Na Zhao[10]   https://orcid.org/0000-0001-6552-0929
Huiyi Tay[1,2]   https://orcid.org/0009-0009-9256-2118
Sonia Youhanna[7]   https://orcid.org/0000-0002-9376-8478
Mengchao Yan[8]   nd
Ye Xie[8]   nd
Youngrae Kim[1,2]   https://orcid.org/0000-0002-7816-9602
Sungdong Lee[1,2]   https://orcid.org/0000-0003-0655-5050
Rachel Liyu Lim[1,2]   https://orcid.org/0009-0009-0241-9079
Guoshou Teo[1,2]   https://orcid.org/0000-0003-3891-1494
Pradeep Narayanaswamy[6] https://orcid.org/0000-0002-3300-2278
Paul R. Burton[9]   nd
Volker M. Lauschke[7,8] https://orcid.org/0000-0002-1140-6204
Hyungwon Choi[1,2,3#] https://orcid.org/0000-0002-6687-3088
Matthew J. Watt[4,#]   https://orcid.org/0000-0003-4064-4188
Philipp Kaldis[10,#,&]   http://orcid.org/0000-0002-7247-7591
24
25
26

References

Adeva-Andany, M. M., Carneiro-Freire, N., Seco-Filgueira, M., Fernandez-Fernandez,
C., & Mourino-Bayolo, D. (2019). Mitochondrial beta-oxidation of saturated fatty
acids   in   humans.   *Mitochondrion*,   *46*,   73-90.
https://doi.org/10.1016/j.mito.2018.02.009
Agarwal, H., Wang, Y., & Ozcan, L. (2023). Rap1 Activation Protects Against Fatty
Liver   and   Non-Alcoholic   Steatohepatitis   Development.   *bioRxiv.*,   DOI
10.1101/2023.1110.1124.563728. https://doi.org/10.1101/2023.10.24.563728
Aggarwal, S., Yadav, V., Maiwall, R., Rastogi, A., Pamecha, V., Bedi, O., Maras, J. S.,
Trehanpati, N., & Ramakrishna, G. (2023). Metabolomic analysis shows
dysregulation in amino acid and NAD+ metabolism in palmitate treated
hepatocytes and plasma of non-alcoholic fatty liver disease spectrum. *Biochem*
Biophys   Res   Commun,   *643*,   129-138.
https://doi.org/10.1016/j.bbrc.2022.12.078
Al-Baiaty, F. D. R., Ismail, A., Abdul Latiff, Z., Muhammad Nawawi, K. N., Raja Ali, R.
A., & Mokhtar, N. M. (2021). Possible Hepatoprotective Effect of Tocotrienol-
Rich Fraction Vitamin E in Non-alcoholic Fatty Liver Disease in Obese Children
and   Adolescents.   *Front   Pediatr*,   *9*,   667247.
https://doi.org/10.3389/fped.2021.667247
Alarcon-Vila, C., Insausti-Urkia, N., Torres, S., Segales-Rovira, P., Conde de la Rosa,
L., Nunez, S., Fucho, R., Fernandez-Checa, J. C., & Garcia-Ruiz, C. (2023).
Dietary and genetic disruption of hepatic methionine metabolism induce acid

sphingomyelinase to promote steatohepatitis. *Redox Biol, 59*, 102596. https://doi.org/10.1016/j.redox.2022.102596

Albaugh, V. L., Mukherjee, K., & Barbul, A. (2017). Proline Precursors and Collagen Synthesis: Biochemical Challenges of Nutrient Supplementation and Wound Healing. *J Nutr, 147*(11), 2011-2017. https://doi.org/10.3945/jn.117.256404

Angulo, P., Kleiner, D. E., Dam-Larsen, S., Adams, L. A., Bjornsson, E. S., Charatcharoenwitthaya, P., Mills, P. R., Keach, J. C., Lafferty, H. D., Stahler, A., Haflidadottir, S., & Bendtsen, F. (2015a). Liver Fibrosis, but No Other Histologic Features, Is Associated With Long-term Outcomes of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology, 149*(2), 389-397.e310. https://doi.org/10.1053/j.gastro.2015.04.043

Angulo, P., Kleiner, D. E., Dam-Larsen, S., Adams, L. A., Bjornsson, E. S., Charatcharoenwitthaya, P., Mills, P. R., Keach, J. C., Lafferty, H. D., Stahler, A., Haflidadottir, S., & Bendtsen, F. (2015b). Liver Fibrosis, but No Other Histologic Features, Is Associated With Long-term Outcomes of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology, 149*(2), 389-397 e310. https://doi.org/10.1053/j.gastro.2015.04.043

Arroyave-Ospina, J. C., Wu, Z., Geng, Y., & Moshage, H. (2021). Role of Oxidative Stress in the Pathogenesis of Non-Alcoholic Fatty Liver Disease: Implications for Prevention and Therapy. *Antioxidants, 10*(2), 174. https://doi.org/10.3390/antiox10020174

Arto, C., Rusu, E. C., Clavero-Mestres, H., Barrientos-Riosalido, A., Bertran, L., Mahmoudian, R., Aguilar, C., Riesco, D., Chicote, J. U., Parada, D., Martinez, S., Sabench, F., Richart, C., & Auguet, T. (2024). Metabolic profiling of tryptophan pathways: Implications for obesity and metabolic dysfunction-associated steatotic liver disease. *Eur J Clin Invest, 54*(11), e14279. https://doi.org/10.1111/eci.14279

Babiy, B., Ramos-Molina, B., Ocana, L., Sacristan, S., Burgos-Santamaria, D., Martinez-Botas, J., Busto, R., Perna, C., Frutos, M. D., Albillos, A., & Pastor, O. (2023). Dihydrosphingolipids are associated with steatosis and increased fibrosis damage in non-alcoholic fatty liver disease. *Biochim Biophys Acta Mol Cell Biol Lipids, 1868*(7), 159318. https://doi.org/10.1016/j.bbalip.2023.159318

Bataller, R., & Brenner, D. A. (2005). Liver fibrosis. *J Clin Invest, 115*(2), 209-218. https://doi.org/10.1172/JCI24282

Bell, C. C., Dankers, A. C. A., Lauschke, V. M., Sison-Young, R., Jenkins, R., Rowe, C., Goldring, C. E., Park, K., Regan, S. L., Walker, T., Schofield, C., Baze, A., Foster, A. J., Williams, D. P., van de Ven, A. W. M., Jacobs, F., Houdt, J. V., Lahteenmaki, T., Snoeys, J., . . . Ingelman-Sundberg, M. (2018). Comparison of Hepatic 2D Sandwich Cultures and 3D Spheroids for Long-term Toxicity Applications: A Multicenter Study. *Toxicol Sci, 162*(2), 655-666. https://doi.org/10.1093/toxsci/kfx289

Bell, C. C., Hendriks, D. F., Moro, S. M., Ellis, E., Walsh, J., Renblom, A., Fredriksson Puigvert, L., Dankers, A. C., Jacobs, F., Snoeys, J., Sison-Young, R. L., Jenkins, R. E., Nordling, A., Mkrtchian, S., Park, B. K., Kitteringham, N. R., Goldring, C. E., Lauschke, V. M., & Ingelman-Sundberg, M. (2016). Characterization of primary human hepatocyte spheroids as a model system for drug-induced liver injury, liver function and disease. *Sci Rep, 6*, 25187. https://doi.org/10.1038/srep25187

Bell, C. C., Lauschke, V. M., Vorrink, S. U., Palmgren, H., Duffin, R., Andersson, T. B., & Ingelman-Sundberg, M. (2017). Transcriptional, Functional, and

Mechanistic Comparisons of Stem Cell-Derived Hepatocytes, HepaRG Cells, and Three-Dimensional Human Hepatocyte Spheroids as Predictive In Vitro Systems for Drug-Induced Liver Injury. *Drug Metab Dispos*, *45*(4), 419-429. https://doi.org/10.1124/dmd.116.074369

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W. H., Pages, F., Trajanoski, Z., & Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics*, *25*(8), 1091-1093. https://doi.org/10.1093/bioinformatics/btp101

Bordi, M., De Cegli, R., Testa, B., Nixon, R. A., Ballabio, A., & Cecconi, F. (2021). A gene toolbox for monitoring autophagy transcription. *Cell Death Dis*, *12*(11), 1044. https://doi.org/10.1038/s41419-021-04121-9

Bos, J. L., Rehmann, H., & Wittinghofer, A. (2007). GEFs and GAPs: critical elements in the control of small G proteins. *Cell*, *129*(5), 865-877. https://doi.org/10.1016/j.cell.2007.05.018

Bril, F., Barb, D., Portillo-Sanchez, P., Biernacki, D., Lomonaco, R., Suman, A., Weber, M. H., Budd, J. T., Lupi, M. E., & Cusi, K. (2017). Metabolic and histological implications of intrahepatic triglyceride content in nonalcoholic fatty liver disease. *Hepatology*, *65*(4), 1132-1144. https://doi.org/10.1002/hep.28985

Brunt, E. M., Kleiner, D. E., Wilson, L. A., Belt, P., & Neuschwander-Tetri, B. A. (2011). Nonalcoholic fatty liver disease (NAFLD) activity score and the histopathologic diagnosis in NAFLD: distinct clinicopathologic meanings. *Hepatology*, *53*(3), 810-820. https://doi.org/10.1002/hep.24127

Carr, R. M., Patel, R. T., Rao, V., Dhir, R., Graham, M. J., Crooke, R. M., & Ahima, R. S. (2012). Reduction of TIP47 improves hepatic steatosis and glucose homeostasis in mice. *Am J Physiol Regul Integr Comp Physiol*, *302*(8), R996-1003. https://doi.org/10.1152/ajpregu.00177.2011

Castane, H., Jimenez-Franco, A., Hernandez-Aguilera, A., Martinez-Navidad, C., Cambra-Cortes, V., Onoiu, A. I., Jimenez-Aguilar, J. M., Paris, M., Hernandez, M., Parada, D., Guilarte, C., Zorzano, A., Hernandez-Alvarez, M. I., Camps, J., & Joven, J. (2025). Multi-omics profiling reveals altered mitochondrial metabolism in adipose tissue from patients with metabolic dysfunction-associated steatohepatitis. *EBioMedicine*, *111*, 105532. https://doi.org/10.1016/j.ebiom.2024.105532

Chalasani, N., Younossi, Z., Lavine, J. E., Charlton, M., Cusi, K., Rinella, M., Harrison, S. A., Brunt, E. M., & Sanyal, A. J. (2017). The diagnosis and management of nonalcoholic fatty liver disease: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology*, *67*(1), 328-357. https://doi.org/10.1002/hep.29367

de Paz-Lugo, P., Lupianez, J. A., & Melendez-Hevia, E. (2018). High glycine concentration increases collagen synthesis by articular chondrocytes in vitro: acute glycine deficiency could be an important cause of osteoarthritis. *Amino Acids*, *50*(10), 1357-1365. https://doi.org/10.1007/s00726-018-2611-x

De Smet, V., Eysackers, N., Merens, V., Kazemzadeh Dastjerd, M., Halder, G., Verhulst, S., Mannaerts, I., & van Grunsven, L. A. (2021). Initiation of hepatic stellate cell activation extends into chronic liver disease. *Cell Death Dis*, *12*(12), 1110. https://doi.org/10.1038/s41419-021-04377-1

Dos Santos, L. R. B., & Fleming, I. (2020). Role of cytochrome P450-derived, polyunsaturated fatty acid mediators in diabetes and the metabolic syndrome.

Prostaglandins Other Lipid Mediat, *148*, 106407. https://doi.org/10.1016/j.prostaglandins.2019.106407

Ekstedt, M., Hagström, H., Nasr, P., Fredrikson, M., Stål, P., Kechagias, S., & Hultcrantz, R. (2015). Fibrosis stage is the strongest predictor for disease-specific mortality in NAFLD after up to 33 years of follow-up. *Hepatology*, *61*(5), 1547-1554.

Eslam, M., Newsome, P. N., Sarin, S. K., Anstee, Q. M., Targher, G., Romero-Gomez, M., Zelber-Sagi, S., Wai-Sun Wong, V., Dufour, J. F., Schattenberg, J. M., Kawaguchi, T., Arrese, M., Valenti, L., Shiha, G., Tiribelli, C., Yki-Jarvinen, H., Fan, J. G., Gronbaek, H., Yilmaz, Y., . . . George, J. (2020). A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement. *J Hepatol*, *73*(1), 202-209. https://doi.org/10.1016/j.jhep.2020.03.039

European Association for the Study of the Liver . Electronic address, e. e. e., European Association for the Study of, D., European Association for the Study of, O., & European Association for the Study of the, L. (2024). EASL-EASD-EASO Clinical Practice Guidelines on the management of metabolic dysfunction-associated steatotic liver disease (MASLD). *J Hepatol*, *81*(3), 492-542. https://doi.org/10.1016/j.jhep.2024.04.031

Feng, G., Valenti, L., Wong, V. W., Fouad, Y. M., Yilmaz, Y., Kim, W., Sebastiani, G., Younossi, Z. M., Hernandez-Gea, V., & Zheng, M. H. (2024). Recompensation in cirrhosis: unravelling the evolving natural history of nonalcoholic fatty liver disease. *Nat Rev Gastroenterol Hepatol*, *21*(1), 46-56. https://doi.org/10.1038/s41575-023-00846-4

Fromenty, B., & Roden, M. (2023). Mitochondrial alterations in fatty liver diseases. *J Hepatol*, *78*(2), 415-429. https://doi.org/10.1016/j.jhep.2022.09.020

Gene Ontology, C., Aleksander, S. A., Balhoff, J., Carbon, S., Cherry, J. M., Drabkin, H. J., Ebert, D., Feuermann, M., Gaudet, P., Harris, N. L., Hill, D. P., Lee, R., Mi, H., Moxon, S., Mungall, C. J., Muruganugan, A., Mushayahama, T., Sternberg, P. W., Thomas, P. D., . . . Westerfield, M. (2023). The Gene Ontology knowledgebase in 2023. *Genetics*, *224*(1), iyad031. https://doi.org/10.1093/genetics/iyad031

Go, C. D., Knight, J. D. R., Rajasekharan, A., Rathod, B., Hesketh, G. G., Abe, K. T., Youn, J. Y., Samavarchi-Tehrani, P., Zhang, H., Zhu, L. Y., Popiel, E., Lambert, J. P., Coyaud, E., Cheung, S. W. T., Rajendran, D., Wong, C. J., Antonicka, H., Pelletier, L., Palazzo, A. F., . . . Gingras, A. C. (2021). A proximity-dependent biotinylation map of a human cell. *Nature*, *595*(7865), 120-124. https://doi.org/10.1038/s41586-021-03592-2

Gou, Y., Li, A., Dong, X., Hao, A., Li, J., Xiang, H., Rahaman, S., He, T. C., & Fan, J. (2025). Lactate transporter MCT4 regulates the hub genes for lipid metabolism and inflammation to attenuate intracellular lipid accumulation in non-alcoholic fatty liver disease. *Genes Dis*, *12*(4), 101554. https://doi.org/10.1016/j.gendis.2025.101554

Govaere, O., Cockell, S., Tiniakos, D., Queen, R., Younes, R., Vacca, M., Alexander, L., Ravaioli, F., Palmer, J., Petta, S., Boursier, J., Rosso, C., Johnson, K., Wonders, K., Day, C. P., Ekstedt, M., Oresic, M., Darlay, R., Cordell, H. J., . . . Anstee, Q. M. (2020). Transcriptomic profiling across the nonalcoholic fatty liver disease spectrum reveals gene signatures for steatohepatitis and fibrosis. *Sci Transl Med*, *12*(572), eaba4448. https://doi.org/10.1126/scitranslmed.aba4448

Govaere, O., Hasoon, M., Alexander, L., Cockell, S., Tiniakos, D., Ekstedt, M., Schattenberg, J. M., Boursier, J., Bugianesi, E., Ratziu, V., Investigators, L., Daly, A. K., & Anstee, Q. M. (2023). A proteo-transcriptomic map of non-alcoholic fatty liver disease signatures. *Nat Metab*, *5*(4), 572-578. https://doi.org/10.1038/s42255-023-00775-1

Gray, J. L., von Delft, F., & Brennan, P. E. (2020). Targeting the Small GTPase Superfamily through Their Regulatory Proteins. *Angew Chem Int Ed Engl*, *59*(16), 6342-6366. https://doi.org/10.1002/anie.201900585

Gual, P., Gilgenkrantz, H., & Lotersztajn, S. (2017). Autophagy in chronic liver diseases: the two faces of Janus. *Am J Physiol Cell Physiol*, *312*(3), C263-C273. https://doi.org/10.1152/ajpcell.00295.2016

Hagstrom, H., Nasr, P., Ekstedt, M., Hammar, U., Stal, P., Hultcrantz, R., & Kechagias, S. (2017). Fibrosis stage but not NASH predicts mortality and time to development of severe liver disease in biopsy-proven NAFLD. *J Hepatol*, *67*(6), 1265-1273. https://doi.org/10.1016/j.jhep.2017.07.027

Hajeyah, A. A., Griffiths, W. J., Wang, Y., Finch, A. J., & O'Donnell, V. B. (2020). The Biosynthesis of Enzymatically Oxidized Lipids. *Front Endocrinol (Lausanne)*, *11*, 591819. https://doi.org/10.3389/fendo.2020.591819

Hess, A., Gentile, S. D., Ben Saad, A., Rahman, R. U., Habboub, T., Pratt, D. S., & Mullen, A. C. (2023). Single-cell transcriptomics stratifies organoid models of metabolic dysfunction-associated steatotic liver disease. *EMBO J*, *42*(24), e113898. https://doi.org/10.15252/embj.2023113898

Hoang, S. A., Oseini, A., Feaver, R. E., Cole, B. K., Asgharpour, A., Vincent, R., Siddiqui, M., Lawson, M. J., Day, N. C., Taylor, J. M., Wamhoff, B. R., Mirshahi, F., Contos, M. J., Idowu, M., & Sanyal, A. J. (2019). Gene Expression Predicts Histological Severity and Reveals Distinct Molecular Profiles of Nonalcoholic Fatty Liver Disease. *Sci Rep*, *9*(1), 12541. https://doi.org/10.1038/s41598-019-48746-5

Holdaway, C. M., Leonard, K. A., Nelson, R., van der Veen, J., Das, C., Watts, R., Clugston, R. D., Lehner, R., & Jacobs, R. L. (2025). Alterations in phosphatidylethanolamine metabolism impacts hepatocellular lipid storage, energy homeostasis, and proliferation. *Biochim Biophys Acta Mol Cell Biol Lipids*, *1870*(4), 159608. https://doi.org/10.1016/j.bbalip.2025.159608

Houten, S. M., Violante, S., Ventura, F. V., & Wanders, R. J. (2016). The Biochemistry and Physiology of Mitochondrial Fatty Acid beta-Oxidation and Its Genetic Disorders. *Annu Rev Physiol*, *78*, 23-44. https://doi.org/10.1146/annurev-physiol-021115-105045

Huang, H., Lee, S. H., Sousa-Lima, I., Kim, S. S., Hwang, W. M., Dagon, Y., Yang, W. M., Cho, S., Kang, M. C., Seo, J. A., Shibata, M., Cho, H., Belew, G. D., Bhin, J., Desai, B. N., Ryu, M. J., Shong, M., Li, P., Meng, H., . . . Kim, Y. B. (2018). Rho-kinase/AMPK axis regulates hepatic lipogenesis during overnutrition. *J Clin Invest*, *128*(12), 5335-5350. https://doi.org/10.1172/JCI63562

Huang, Q., Qadri, S. F., Bian, H., Yi, X., Lin, C., Yang, X., Zhu, X., Lin, H., Yan, H., Chang, X., Sun, X., Ma, S., Wu, Q., Zeng, H., Hu, X., Zheng, Y., Yki-Jarvinen, H., Gao, X., Tang, H., & Xia, M. (2025). A metabolome-derived score predicts metabolic dysfunction-associated steatohepatitis and mortality from liver disease. *J Hepatol*, *82*(5), 781-793. https://doi.org/10.1016/j.jhep.2024.10.015

Huttlin, E. L., Bruckner, R. J., Paulo, J. A., Cannon, J. R., Ting, L., Baltier, K., Colby, G., Gebreab, F., Gygi, M. P., Parzen, H., Szpyt, J., Tam, S., Zarraga, G., Pontano-Vaites, L., Swarup, S., White, A. E., Schweppe, D. K., Rad, R.,

Erickson, B. K., . . . Harper, J. W. (2017). Architecture of the human interactome defines protein communities and disease networks. *Nature*, *545*(7655), 505-509. https://doi.org/10.1038/nature22366

Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., & Tanabe, M. (2016). KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res*, *44*(D1), D457-462. https://doi.org/10.1093/nar/gkv1070

Kang, Q., & Chen, A. (2009). Curcumin eliminates oxidized LDL roles in activating hepatic stellate cells by suppressing gene expression of lectin-like oxidized LDL receptor-1. *Lab Invest*, *89*(11), 1275-1290. https://doi.org/10.1038/labinvest.2009.93

Kemas, A. M., Youhanna, S., Zandi Shafagh, R., & Lauschke, V. M. (2021). Insulin-dependent glucose consumption dynamics in 3D primary human liver cultures measured by a sensitive and specific glucose sensor with nanoliter input volume. *FASEB J*, *35*(3), e21305. https://doi.org/10.1096/fj.202001989RR

Kisseleva, T., & Brenner, D. (2021). Molecular and cellular mechanisms of liver fibrosis and its regression. *Nat Rev Gastroenterol Hepatol*, *18*(3), 151-166. https://doi.org/10.1038/s41575-020-00372-7

Kleiner, D. E., Brunt, E. M., Van Natta, M., Behling, C., Contos, M. J., Cummings, O. W., Ferrell, L. D., Liu, Y. C., Torbenson, M. S., Unalp-Arida, A., Yeh, M., McCullough, A. J., Sanyal, A. J., & Nonalcoholic Steatohepatitis Clinical Research, N. (2005). Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology*, *41*(6), 1313-1321. https://doi.org/10.1002/hep.20701

Knottnerus, S. J. G., Bleeker, J. C., Wust, R. C. I., Ferdinandusse, S., L, I. J., Wijburg, F. A., Wanders, R. J. A., Visser, G., & Houtkooper, R. H. (2018). Disorders of mitochondrial long-chain fatty acid oxidation and the carnitine shuttle. *Rev Endocr Metab Disord*, *19*(1), 93-106. https://doi.org/10.1007/s11154-018-9448-1

Koliaki, C., Szendroedi, J., Kaul, K., Jelenik, T., Nowotny, P., Jankowiak, F., Herder, C., Carstensen, M., Krausch, M., Knoefel, W. T., Schlensak, M., & Roden, M. (2015). Adaptation of hepatic mitochondrial function in humans with non-alcoholic fatty liver is lost in steatohepatitis. *Cell Metab*, *21*(5), 739-746. https://doi.org/10.1016/j.cmet.2015.04.004

Lahdou, I., Sadeghi, M., Oweira, H., Fusch, G., Daniel, V., Mehrabi, A., Jung, G., Elhadedy, H., Schmidt, J., Sandra-Petrescu, F., Iancu, M., Opelz, G., Terness, P., & Schefold, J. C. (2013). Increased serum levels of quinolinic acid indicate enhanced severity of hepatic dysfunction in patients with liver cirrhosis. *Hum Immunol*, *74*(1), 60-66. https://doi.org/10.1016/j.humimm.2012.09.009

Lambert, J. E., Ramos-Roman, M. A., Browning, J. D., & Parks, E. J. (2014). Increased de novo lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver disease. *Gastroenterology*, *146*(3), 726-735. https://doi.org/10.1053/j.gastro.2013.11.049

Lee, D. Y., Kind, T., Yoon, Y. R., Fiehn, O., & Liu, K. H. (2014). Comparative evaluation of extraction methods for simultaneous mass-spectrometric analysis of complex lipids and primary metabolites from human blood plasma. *Anal Bioanal Chem*, *406*(28), 7275-7286. https://doi.org/10.1007/s00216-014-8124-x

Lee, S., Bang, J., Kim, Y., Choi, H., Oh, S.-Y., & Won, J.-H. (2025). Learning massive-scale partial correlation networks in clinical multiomics studies with HP-ACCORD. *The Annals of Applied Statistics*, *19*(4), 2759-2781, 2723. https://doi.org/10.1214/25-AOAS2089

Li, Y., Fan, W., Link, F., Wang, S., & Dooley, S. (2022). Transforming growth factor beta latency: A mechanism of cytokine storage and signalling regulation in liver homeostasis and disease. *JHEP Rep*, *4*(2), 100397. https://doi.org/10.1016/j.jhepr.2021.100397

Liu, J., Hu, S., Chen, L., Daly, C., Prada Medina, C. A., Richardson, T. G., Traylor, M., Dempster, N. J., Mbasu, R., Monfeuga, T., Vujkovic, M., Tsao, P. S., Lynch, J. A., Voight, B. F., Chang, K. M., Million, V. A., Cobbold, J. F., Tomlinson, J. W., van Duijn, C. M., & Howson, J. M. M. (2023). Profiling the genome and proteome of metabolic dysfunction-associated steatotic liver disease identifies potential therapeutic targets. *medRxiv.*, DOI: 10.1101/2023.1111.1130.23299247. https://doi.org/10.1101/2023.11.30.23299247

MacParland, S. A., Liu, J. C., Ma, X. Z., Innes, B. T., Bartczak, A. M., Gage, B. K., Manuel, J., Khuu, N., Echeverri, J., Linares, I., Gupta, R., Cheng, M. L., Liu, L. Y., Camat, D., Chung, S. W., Seliga, R. K., Shao, Z., Lee, E., Ogawa, S., . . . McGilvray, I. D. (2018). Single cell RNA sequencing of human liver reveals distinct intrahepatic macrophage populations. *Nat Commun*, *9*(1), 4383. https://doi.org/10.1038/s41467-018-06318-7

Martinez-Lopez, N., & Singh, R. (2015). Autophagy and Lipid Droplets in the Liver. *Annu Rev Nutr*, *35*, 215-237. https://doi.org/10.1146/annurev-nutr-071813-105336

Mashek, D. G. (2021). Hepatic lipid droplets: A balancing act between energy storage and metabolic dysfunction in NAFLD. *Mol Metab*, *50*, 101115. https://doi.org/10.1016/j.molmet.2020.101115

Massague, J., & Sheppard, D. (2023). TGF-beta signaling in health and disease. *Cell*, *186*(19), 4007-4037. https://doi.org/10.1016/j.cell.2023.07.036

Matthews, R. P., Lorent, K., Manoral-Mobias, R., Huang, Y., Gong, W., Murray, I. V., Blair, I. A., & Pack, M. (2009). TNFalpha-dependent hepatic steatosis and liver degeneration caused by mutation of zebrafish S-adenosylhomocysteine hydrolase. *Development*, *136*(5), 865-875. https://doi.org/10.1242/dev.027565

McGlinchey, A. J., Govaere, O., Geng, D., Ratziu, V., Allison, M., Bousier, J., Petta, S., de Oliviera, C., Bugianesi, E., Schattenberg, J. M., Daly, A. K., Hyotylainen, T., Anstee, Q. M., & Oresic, M. (2022). Metabolic signatures across the full spectrum of non-alcoholic fatty liver disease. *JHEP Rep*, *4*(5), 100477. https://doi.org/10.1016/j.jhepr.2022.100477

Messner, S., Fredriksson, L., Lauschke, V. M., Roessger, K., Escher, C., Bober, M., Kelm, J. M., Ingelman-Sundberg, M., & Moritz, W. (2018). Transcriptomic, Proteomic, and Functional Long-Term Characterization of Multicellular Three-Dimensional Human Liver Microtissues. *Appl In Vitro Toxicol*, *4*(1), 1-12. https://doi.org/10.1089/aivt.2017.0022

Miao, L., Targher, G., Byrne, C. D., Cao, Y. Y., & Zheng, M. H. (2024). Current status and future trends of the global burden of MASLD. *Trends Endocrinol Metab*, *35*(8), 697-707. https://doi.org/10.1016/j.tem.2024.02.007

Min, H. K., Kapoor, A., Fuchs, M., Mirshahi, F., Zhou, H., Maher, J., Kellum, J., Warnick, R., Contos, M. J., & Sanyal, A. J. (2012). Increased hepatic synthesis and dysregulation of cholesterol metabolism is associated with the severity of nonalcoholic fatty liver disease. *Cell Metab*, *15*(5), 665-674. https://doi.org/10.1016/j.cmet.2012.04.004

Munger, J. S., Huang, X., Kawakatsu, H., Griffiths, M. J., Dalton, S. L., Wu, J., Pittet, J. F., Kaminski, N., Garat, C., Matthay, M. A., Rifkin, D. B., & Sheppard, D.

(1999). The integrin alpha v beta 6 binds and activates latent TGF beta 1: a mechanism for regulating pulmonary inflammation and fibrosis. *Cell*, *96*(3), 319-328. https://doi.org/10.1016/s0092-8674(00)80545-0

Narayanaswamy, P., Teo, G., Ow, J. R., Lau, A., Kaldis, P., Tate, S., & Choi, H. (2020). MetaboKit: a comprehensive data extraction tool for untargeted metabolomics. *Mol Omics*, *16*(5), 436-447. https://doi.org/10.1039/d0mo00030b

Ooi, G. J., Meikle, P. J., Huynh, K., Earnest, A., Roberts, S. K., Kemp, W., Parker, B. L., Brown, W., Burton, P., & Watt, M. J. (2021). Hepatic lipidomic remodeling in severe obesity manifests with steatosis and does not evolve with non-alcoholic steatohepatitis. *J Hepatol*, *75*(3), 524-535. https://doi.org/10.1016/j.jhep.2021.04.013

Otasek, D., Morris, J. H., Boucas, J., Pico, A. R., & Demchak, B. (2019). Cytoscape Automation: empowering workflow-based network analysis. *Genome Biol*, *20*(1), 185. https://doi.org/10.1186/s13059-019-1758-4

Peng, Y., Zeng, Q., Wan, L., Ma, E., Li, H., Yang, X., Zhang, Y., Huang, L., Lin, H., Feng, J., Xu, Y., Li, J., Liu, M., Liu, J., Lin, C., Sun, Z., Cheng, G., Zhang, X., Liu, J., . . . Wei, C. (2021). GP73 is a TBC-domain Rab GTPase-activating protein contributing to the pathogenesis of non-alcoholic fatty liver disease without obesity. *Nat Commun*, *12*(1), 7004. https://doi.org/10.1038/s41467-021-27309-1

Powell, E. E., Wong, V. W., & Rinella, M. (2021). Non-alcoholic fatty liver disease. *Lancet*, *397*(10290), 2212-2224. https://doi.org/10.1016/S0140-6736(20)32511-3

Radosavljevic, T., Brankovic, M., Samardzic, J., Djuretic, J., Vukicevic, D., Vucevic, D., & Jakovljevic, V. (2024). Altered Mitochondrial Function in MASLD: Key Features and Promising Therapeutic Approaches. *Antioxidants*, *13*(8), 906. https://doi.org/10.3390/antiox13080906

Rath, S., Sharma, R., Gupta, R., Ast, T., Chan, C., Durham, T. J., Goodman, R. P., Grabarek, Z., Haas, M. E., Hung, W. H. W., Joshi, P. R., Jourdain, A. A., Kim, S. H., Kotrys, A. V., Lam, S. S., McCoy, J. G., Meisel, J. D., Miranda, M., Panda, A., . . . Mootha, V. K. (2021). MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic Acids Res*, *49*(D1), D1541-D1547. https://doi.org/10.1093/nar/gkaa1011

Ratziu, V., Harrison, S. A., Francque, S., Bedossa, P., Lehert, P., Serfaty, L., Romero-Gomez, M., Boursier, J., Abdelmalek, M., Caldwell, S., Drenth, J., Anstee, Q. M., Hum, D., Hanf, R., Roudot, A., Megnien, S., Staels, B., Sanyal, A., & Group, G.-I. S. (2016). Elafibranor, an Agonist of the Peroxisome Proliferator-Activated Receptor-alpha and -delta, Induces Resolution of Nonalcoholic Steatohepatitis Without Fibrosis Worsening. *Gastroenterology*, *150*(5), 1147-1159 e1145. https://doi.org/10.1053/j.gastro.2016.01.038

Raverdy, V., Tavaglione, F., Chatelain, E., Lassailly, G., De Vincentis, A., Vespasiani-Gentilucci, U., Qadri, S. F., Caiazzo, R., Verkindt, H., Saponaro, C., Kerr-Conte, J., Baud, G., Marciniak, C., Chetboun, M., Oukhouya-Daoud, N., Blanck, S., Vandel, J., Olsson, L., Chakaroun, R., . . . Pattou, F. (2024). Data-driven cluster analysis identifies distinct types of metabolic dysfunction-associated steatotic liver disease. *Nat Med*, *30*(12), 3624-3633. https://doi.org/10.1038/s41591-024-03283-1

Razick, S., Magklaras, G., & Donaldson, I. M. (2008). iRefIndex: a consolidated protein interaction database with provenance. *BMC Bioinformatics*, *9*, 405. https://doi.org/10.1186/1471-2105-9-405

Riazi, K., Azhari, H., Charette, J. H., Underwood, F. E., King, J. A., Afshar, E. E., Swain, M. G., Congly, S. E., Kaplan, G. G., & Shaheen, A. A. (2022). The prevalence and incidence of NAFLD worldwide: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol*, *7*(9), 851-861. https://doi.org/10.1016/S2468-1253(22)00165-0

Rinella, M. E., Neuschwander-Tetri, B. A., Siddiqui, M. S., Abdelmalek, M. F., Caldwell, S., Barb, D., Kleiner, D. E., & Loomba, R. (2023). AASLD Practice Guidance on the clinical assessment and management of nonalcoholic fatty liver disease. *Hepatology*, *77*(5), 1797-1835. https://doi.org/10.1097/HEP.0000000000000323

Robinson, A. E., Binek, A., Ramani, K., Sundararaman, N., Barbier-Torres, L., Murray, B., Venkatraman, V., Kreimer, S., Ardle, A. M., Noureddin, M., Fernandez-Ramos, D., Lopitz-Otsoa, F., Gutierrez de Juan, V., Millet, O., Mato, J. M., Lu, S. C., & Van Eyk, J. E. (2023). Hyperphosphorylation of hepatic proteome characterizes nonalcoholic fatty liver disease in S-adenosylmethionine deficiency. *iScience*, *26*(2), 105987. https://doi.org/10.1016/j.isci.2023.105987

Rodriguez, L. A., Schmittdiel, J. A., Liu, L., Macdonald, B. A., Balasubramanian, S., Chai, K. P., Seo, S. I., Mukhtar, N., Levin, T. R., & Saxena, V. (2024). Hepatocellular Carcinoma in Metabolic Dysfunction-Associated Steatotic Liver Disease. *JAMA Netw Open*, *7*(7), e2421019. https://doi.org/10.1001/jamanetworkopen.2024.21019

Roumans, K. H. M., Lindeboom, L., Veeraiah, P., Remie, C. M. E., Phielix, E., Havekes, B., Bruls, Y. M. H., Brouwers, M., Stahlman, M., Alssema, M., Peters, H. P. F., de Mutsert, R., Staels, B., Taskinen, M. R., Boren, J., Schrauwen, P., & Schrauwen-Hinderling, V. B. (2020). Hepatic saturated fatty acid fraction is associated with de novo lipogenesis and hepatic insulin resistance. *Nat Commun*, *11*(1), 1891. https://doi.org/10.1038/s41467-020-15684-0

Sakuma, I., Gaspar, R. C., Nasiri, A. R., Dufour, S., Kahn, M., Zheng, J., LaMoia, T. E., Guerra, M. T., Taki, Y., Kawashima, Y., Yimlamai, D., Perelis, M., Vatner, D. F., Petersen, K. F., Huttasch, M., Knebel, B., Kahl, S., Roden, M., Samuel, V. T., . . . Shulman, G. I. (2025). Liver lipid droplet cholesterol content is a key determinant of metabolic dysfunction-associated steatohepatitis. *Proc Natl Acad Sci U S A*, *122*(18), e2502978122. https://doi.org/10.1073/pnas.2502978122

Saliba-Gustafsson, P., Justesen, J. M., Ranta, A., Sharma, D., Bielczyk-Maczynska, E., Li, J., Najmi, L. A., Apodaka, M., Aspichueta, P., Bjorck, H. M., Eriksson, P., Schurr, T. M., Franco-Cereceda, A., Gloudemans, M., Mujica, E., den Hoed, M., Assimes, T. L., Quertermous, T., Carcamo-Orive, I., . . . Knowles, J. W. (2024). A functional genomic framework to elucidate novel causal metabolic dysfunction-associated fatty liver disease genes. *Hepatology*, : in press. https://doi.org/10.1097/HEP.0000000000001066

Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction*, *88*(6), 791-804. https://doi.org/10.1111/j.1360-0443.1993.tb02093.x

Schnabl, B., Damman, C. J., & Carr, R. M. (2025). Metabolic dysfunction-associated steatotic liver disease and the gut microbiome: pathogenic insights and therapeutic innovations. *J Clin Invest*, *135*(7). https://doi.org/10.1172/JCI186423

Schwerbel, K., Kamitz, A., Krahmer, N., Hallahan, N., Jahnert, M., Gottmann, P., Lebek, S., Schallschmidt, T., Arends, D., Schumacher, F., Kleuser, B., Haltenhof, T., Heyd, F., Gancheva, S., Broman, K. W., Roden, M., Joost, H. G., Chadt, A., Al-Hasani, H., . . . Schurmann, A. (2020). Immunity-related GTPase induces lipophagy to prevent excess hepatic lipid accumulation. *J Hepatol*, *73*(4), 771-782. https://doi.org/10.1016/j.jhep.2020.04.031

Scoditti, E., Sabatini, S., Carli, F., & Gastaldelli, A. (2024). Hepatic glucose metabolism in the steatotic liver. *Nat Rev Gastroenterol Hepatol*, *21*(5), 319-334. https://doi.org/10.1038/s41575-023-00888-8

Scorletti, E., & Carr, R. M. (2022). A new perspective on NAFLD: Focusing on lipid droplets. *J Hepatol*, *76*(4), 934-945. https://doi.org/10.1016/j.jhep.2021.11.009

Singh, R., Kaushik, S., Wang, Y., Xiang, Y., Novak, I., Komatsu, M., Tanaka, K., Cuervo, A. M., & Czaja, M. J. (2009). Autophagy regulates lipid metabolism. *Nature*, *458*(7242), 1131-1135. https://doi.org/10.1038/nature07976

Sousa, A. P., Cunha, D. M., Franco, C., Teixeira, C., Gojon, F., Baylina, P., & Fernandes, R. (2021). Which Role Plays 2-Hydroxybutyric Acid on Insulin Resistance? *Metabolites*, *11*(12). https://doi.org/10.3390/metabo11120835

Subramanian, P., Hampe, J., Tacke, F., & Chavakis, T. (2022). Fibrogenic Pathways in Metabolic Dysfunction Associated Fatty Liver Disease (MAFLD). *Int J Mol Sci*, *23*(13), 6996. https://doi.org/10.3390/ijms23136996

Tan, Q., Deng, S., & Xiong, L. (2025). Role of Kynurenine and Its Derivatives in Liver Diseases: Recent Advances and Future Clinical Perspectives. *Int J Mol Sci*, *26*(3). https://doi.org/10.3390/ijms26030968

Targher, G., Byrne, C. D., & Tilg, H. (2024). MASLD: a systemic metabolic disorder with cardiovascular and malignant complications. *Gut*, *73*(4), 691-702. https://doi.org/10.1136/gutjnl-2023-330595

Trivedi, P., Wang, S., & Friedman, S. L. (2021). The Power of Plasticity-Metabolic Regulation of Hepatic Stellate Cells. *Cell Metab*, *33*(2), 242-257. https://doi.org/10.1016/j.cmet.2020.10.026

Tuohetahuntila, M., Molenaar, M. R., Spee, B., Brouwers, J. F., Wubbolts, R., Houweling, M., Yan, C., Du, H., VanderVen, B. C., Vaandrager, A. B., & Helms, J. B. (2017). Lysosome-mediated degradation of a distinct pool of lipid droplets during hepatic stellate cell activation. *J Biol Chem*, *292*(30), 12436-12448. https://doi.org/10.1074/jbc.M117.778472

van der Veen, J. N., Lingrell, S., & Vance, D. E. (2012). The membrane lipid phosphatidylcholine is an unexpected source of triacylglycerol in the liver. *J Biol Chem*, *287*(28), 23418-23426. https://doi.org/10.1074/jbc.M112.381723

Van Woerkom, A., Harney, D. J., Nagarajan, S. R., Hakeem-Sanni, M. F., Lin, J., Hooke, M., Pulpitel, T., Cooney, G. J., Larance, M., Saunders, D. N., Brandon, A. E., & Hoy, A. J. (2024). Hepatic lipid droplet-associated proteome changes distinguish dietary-induced fatty liver from glucose tolerance in male mice. *Am J Physiol Endocrinol Metab*, *326*(6), E842-E855. https://doi.org/10.1152/ajpendo.00013.2024

Varela-Rey, M., Martinez-Lopez, N., Fernandez-Ramos, D., Embade, N., Calvisi, D. F., Woodhoo, A., Rodriguez, J., Fraga, M. F., Julve, J., Rodriguez-Millan, E., Frades, I., Torres, L., Luka, Z., Wagner, C., Esteller, M., Lu, S. C., Martinez-Chantar, M. L., & Mato, J. M. (2010). Fatty liver and fibrosis in glycine N-methyltransferase knockout mice is prevented by nicotinamide. *Hepatology*, *52*(1), 105-114. https://doi.org/10.1002/hep.23639

Vorrink, S. U., Ullah, S., Schmidt, S., Nandania, J., Velagapudi, V., Beck, O., Ingelman-Sundberg, M., & Lauschke, V. M. (2017). Endogenous and xenobiotic metabolic stability of primary human hepatocytes in long-term 3D spheroid cultures revealed by a combination of targeted and untargeted metabolomics. *FASEB J, 31*(6), 2696-2708. https://doi.org/10.1096/fj.201601375R

Vvedenskaya, O., Rose, T. D., Knittelfelder, O., Palladini, A., Wodke, J. A. H., Schuhmann, K., Ackerman, J. M., Wang, Y., Has, C., Brosch, M., Thangapandi, V. R., Buch, S., Zullig, T., Hartler, J., Kofeler, H. C., Rocken, C., Coskun, U., Klipp, E., von Schoenfels, W., . . . Shevchenko, A. (2021). Nonalcoholic fatty liver disease stratification by liver lipidomics. *J Lipid Res, 62*, 100104. https://doi.org/10.1016/j.jlr.2021.100104

Walker, A. K. (2017). 1-Carbon Cycle Metabolites Methylate Their Way to Fatty Liver. *Trends Endocrinol Metab, 28*(1), 63-72. https://doi.org/10.1016/j.tem.2016.10.004

Wang, D., Wei, Y., & Pagliassotti, M. J. (2006). Saturated fatty acids promote endoplasmic reticulum stress and liver injury in rats with hepatic steatosis. *Endocrinology, 147*(2), 943-951. https://doi.org/10.1210/en.2005-0570

Wei, W., Lyu, X., Markhard, A. L., Fu, S., Mardjuki, R. E., Cavanagh, P. E., Zeng, X., Rajniak, J., Lu, N., Xiao, S., Zhao, M., Moya-Garzon, M. D., Truong, S. D., Chou, J. C., Wat, L. W., Chidambaranathan-Reghupaty, S., Coassolo, L., Xu, D., Shen, F., . . . Long, J. Z. (2024). PTER is a N-acetyltaurine hydrolase that regulates feeding and obesity. *Nature, 633*(8028), 182-188. https://doi.org/10.1038/s41586-024-07801-6

Wen, X., Liu, Y., Bai, Y., Li, M., Fu, Q., & Zheng, Y. (2018). LOXL2, a copper-dependent monoamine oxidase, activates lung fibroblasts through the TGF-beta/Smad pathway. *Int J Mol Med, 42*(6), 3530-3541. https://doi.org/10.3892/ijmm.2018.3927

Wu, Z. E., Kruger, M. C., Cooper, G. J. S., Poppitt, S. D., & Fraser, K. (2019). Tissue-Specific Sample Dilution: An Important Parameter to Optimise Prior to Untargeted LC-MS Metabolomics. *Metabolites, 9*(7), 124. https://doi.org/10.3390/metabo9070124

Xie, C., Woollett, L. A., Turley, S. D., & Dietschy, J. M. (2002). Fatty acids differentially regulate hepatic cholesteryl ester formation and incorporation into lipoproteins in the liver of the mouse. *J Lipid Res, 43*(9), 1508-1519. https://doi.org/10.1194/jlr.m200146-jlr200

Xu, J., Begley, P., Church, S. J., Patassini, S., Hollywood, K. A., Jullig, M., Curtis, M. A., Waldvogel, H. J., Faull, R. L., Unwin, R. D., & Cooper, G. J. (2016). Graded perturbations of metabolism in multiple regions of human brain in Alzheimer's disease: Snapshot of a pervasive metabolic disorder. *Biochim Biophys Acta, 1862*(6), 1084-1092. https://doi.org/10.1016/j.bbadis.2016.03.001

Xue, C., Li, G., Zheng, Q., Gu, X., Shi, Q., Su, Y., Chu, Q., Yuan, X., Bao, Z., Lu, J., & Li, L. (2023). Tryptophan metabolism in health and disease. *Cell Metab, 35*(8), 1304-1326. https://doi.org/10.1016/j.cmet.2023.06.004

Yanko, R., Levashov, M., Chaka, O. G., Nosar, V., Khasabov, S. G., & Khasabova, I. (2023). Tryptophan Prevents the Development of Non-Alcoholic Fatty Liver Disease. *Diabetes Metab Syndr Obes, 16*, 4195-4204. https://doi.org/10.2147/DMSO.S444278

Yki-Jarvinen, H. (2014). Non-alcoholic fatty liver disease as a cause and a consequence of metabolic syndrome. *Lancet Diabetes Endocrinol, 2*(11), 901-910. https://doi.org/10.1016/S2213-8587(14)70032-4

Youhanna, S., Kemas, A. M., Wright, S. C., Zhong, Y., Klumpp, B., Klein, K., Motso,
A., Michel, M., Ziegler, N., Shang, M., Sabatier, P., Kannt, A., Sheng, H., Oliva-
Vilarnau, N., Buttner, F. A., Seashore-Ludlow, B., Schreiner, J., Windbergs, M.,
Cornillet, M., . . . Lauschke, V. M. (2025). Chemogenomic Screening in a
Patient-Derived 3D Fatty Liver Disease Model Reveals the CHRM1-TRPM8
Axis as a Novel Module for Targeted Intervention. *Adv Sci (Weinh)*, *12*(3),
e2407572. https://doi.org/10.1002/advs.202407572
Younossi, Z., Anstee, Q. M., Marietti, M., Hardy, T., Henry, L., Eslam, M., George, J.,
& Bugianesi, E. (2018). Global burden of NAFLD and NASH: trends,
predictions, risk factors and prevention. *Nat Rev Gastroenterol Hepatol*, *15*(1),
11-20. https://doi.org/10.1038/nrgastro.2017.109
Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O.,
Benner, C., & Chanda, S. K. (2019). Metascape provides a biologist-oriented
resource for the analysis of systems-level datasets. *Nat Commun*, *10*(1), 1523.
https://doi.org/10.1038/s41467-019-09234-6
16

Graphical abstract

Dual-omics profiling

Metabolomics + *Transcriptomics*

Steatosis

Fibrosis

TAG ↑

TAG and DAG ↑ GPL ↓
Amino acid metabolism ↑ ↓
Tryptophan metabolism ↑ ↓
Autophagy activation ↑

Mitochondrial function ↓
Small molecule metabolism ↓
Signal transduction ↑
Extracellular matrix remodeling ↑

Regulation of GTPase-related genes

Figure 1. Study overview

(A) The overall study design. (B) Alluvial diagram of patient composition of groupings across liver histological features. (C) Relationship between clinical test results and stages of liver histological features. Node size was determined by –log10(p value) in ANOVA tests. Color indicates the significance degree [–log10(p value)] and the direction of change from early to late stages. (D) Number of metabolites analyzed in the liver and plasma. (E-F) Principal component analysis of liver metabolome, plasma metabolome, and liver transcriptome. (G) Disease spectra covered in this cohort and comparision to two published datasets.

Figure 2. Hepatic and circulating metabolome in obese MASLD patients

(A) Number of metabolites in each class that were significantly associated with histological features in the liver (q value < 0.05). Metabolite classes with at least 5 metabolites with significance are shown in the plot. (B) Associations between steatosis grades and lipid species in each lipid class. Dots were colored by double bond numbers (left) and carbon numbers (right) of lipid species, respectively. (C) Partial correlation network of the plasma metabolome and clinical covariates. Node size reflects the degree of connectivity, with larger nodes indicating connections to a greater number of metabolite nodes. (D) Heatmaps for pairwise analysis between plasma lipid classes and clinical variables. Linear regression analysis was performed for numerical variables, while logistic regression was conducted for binary variables. Significance values refer to −log10(p value)*sign(coefficients) from regression models. (E) Comparisons of liver metabolome and plasma metabolome regarding their associations with liver steatosis and fibrosis.

Figure 3. Integrative view of key metabolic pathways implicated in liver metabolism in obese individuals with MASLD
(A) Heatmap of log2 fold changes from pair-wise analysis of lipid metabolism-related genes. Association with steatosis (S) and fibrosis (F) is indicated by black dots (q value < 0.05, top). Results were cross-referenced with two published cohorts (VA cohort, GSE130970; EU cohort, GSE135251, bottom) Asterisks indicate genes with a q-value < 0.05 and a consistent change in direction within our cohort. (B) KEGG metabolic pathways with at least 4 genes or metabolites significantly associated with steatosis (S) or fibrosis (F). (C) Integrative map of gene and metabolite alterations associated with steatosis (left half of each box/circle) and fibrosis (right half of each box/circle). (D) Metabolite alterations corresponding to the advancement of steatosis grades (x-axis).

Figure 4. **Dysregulated mitochondrial function and autophagy during the progression of steatosis and fibrosis.**
Gene expression patterns related to mitochondrial metabolism (A), mitochondria-related biological processes (A), autophagy activation (B), and lysosomal biogenesis (B) with statistical association to steatosis and fibrosis. S, steatosis. F, fibrosis.

Figure 5. **Steatosis and fibrosis as independent processes**
(A) Network of genes, metabolites, and histological features (fibrosis and steatosis) using partial correlation analysis as described in the methods. (B) Venn diagrams depicting the number of genes significantly associated with steatosis and fibrosis (q value < 0.05). (C) Gene ontology (GO) enrichment of gene sets specific to steatosis or fibrosis. H, healthy obese controls. S, steatosis. F, fibrosis.

Figure 6. Liver fibrosis signatures and a potential therapeutic target of fibrosis initiation.
(A) Top 60 fibrosis markers in the liver. A total of 213 genes were identified, from which the top 30 upregulated and top 30 downregulated genes are visualized. The left side of the plot (blue bar on top) shows the comparison of individuals with fibrosis and individuals without pathology (baseline - "no MASLD"). For the right side (red bar), individuals with fibrosis were compared to those with steatosis but no fibrosis (baseline – "steatosis but no fibrosis"). Results were cross-referenced with two published cohorts (VA cohort, GSE130970; EU cohort, GSE135251, right). Asterisks indicate genes with a q-value < 0.05 and a consistent change in direction within our cohort. The average gene expressions of liver cell types were obtained from GSE115469 at the log2CPM level. Dots in the single cell map indicate zero expression in the corresponding cell types. (B) Enrichment of pathways in the transition from simple steatosis to the onset of fibrosis.

(C) Protein interaction network of GTPases and their regulators. Nodes are colored based on gene type, with borders indicating the direction of gene regulation. Node size corresponds to the significance of the genes in relation to fibrosis grades within this cohort. GEFs: Guanine nucleotide exchange factors. GAPs: GTPase-activating proteins. (D) Comparison of expression level changes in GTPase-related genes between this human cohort and an independent 3D spheroid MASH system. Log2 fold change for the human cohort was calculated by comparing patients with grade 2 or 3 fibrosis to those without fibrosis or steatosis. Genes with the same direction of change are colored in red, while others are colored in purple. (E) Expression of GTPase-related genes in patient-derived 3D liver spheroids (controls; MASH; MASH treated with Elafibranor; n = 4). Sixty-eight genes with a p-value < 0.05 from the ANOVA test were plotted in the diagram, with blue lines highlighting 24 genes that exhibited increased expression in the MASH group compared to the control group.

Figure 2–figure supplement 1. Metabolomic analysis in this obese MASLD cohort
(A) Plasma urea and creatinine from untargeted metabolomics correlated with clinical assays. (B) Metabolites associated with the disease progression in each matrix (q < 0.05 or q < 0.1). Compound classes with at least 5 hits associated with any histological feature were shown in the plot. (C) Associations between fatty acid composition of triacylglycerides (TAGs) and histological outcomes. Significance values refer to –log10(p value)*sign(coefficient) from linear regression models.

Figure 2–figure supplement 2. Blood lipoprotein cholesterol levels in patients with different steatosis and fibrosis grades. HDL, high-density lipoprotein cholesterol. LDL, low-density lipoprotein cholesterol. NonHDL-Chol, non-high-density lipoprotein cholesterol. TotalChol, total blood lipoprotein cholesterol.

A

B

Figure 3–figure supplement 1. Integrative map of liver metabolism in obese individuals with MASLD.
(A) An integrative map of transcriptomics and metabolomics of glycerolipid and glycerophospholipid metabolism in early MASLD. (B) Gene expression changes in fatty acid β-oxidation, mitochondrial respiratory chain, and lipid droplet (LD) metabolism associated with steatosis and fibrosis. There was a notable downregulation of genes involved in the electron transport chain of the inner mitochondrial membrane, specifically in complex I (NDUF family), complex II (SDH family), complex III (UQCR), complex IV (COX), and complex V (ATP5 family) as fibrosis progressed. Genes involved in respiratory electron transport were colored by the significance ($-\log10$(p value)*sign(coefficient)) in relation to fibrosis grades. Inset: gene expression changes in the electron transport chain are depicted.

Figure 3–figure supplement 2. Ratios of metabolites in one-carbon metabolism in individuals with different steatosis grades. SAM, S-adenosylmethionine. SAH, S-adenosylhomocysteine.

Figure 4–figure supplement 1. Autophagy regulation in the liver.
(A) Hepatic free cholesterol, CE 18:1, and CE 18:2 levels in this cohort in relation to steatosis grade (x-axis). (B) Heatmap of autophagy genes in relation to histological features, indicating active lipophagy in the liver of obese individuals. Asterisks in the main heatmap denote the significance (p < 0.05) from comparisons of each subgroup. Association with steatosis (S), NAS (N) and fibrosis (F) was indicated by black dots (q < 0.05, left). Results were cross-referenced with two published cohorts (VA cohort, GSE130970; EU cohort, GSE135251) (*q < 0.05 and with consistent direction of changes as in our cohort, bottom).

Figure 5–figure supplement 1. Functional enrichment of gene sets associated with steatosis and fibrosis in the liver transcriptome. Node size is depicted by circles and P value by color.

Figure 5–figure supplement 2. Statistical power and subgroup analysis of associations between gene expressions and steatosis or fibrosis grades.
Statistical power in the original linear regression analysis of steatosis and fibrosis was plotted against p-values (A). Differential features associated with steatosis (q < 0.05) showed power > 0.8, whereas those associated with fibrosis showed power > 0.7. Subgroup analyses of participants with and without diabetes were conducted to evaluate associations between gene expression and steatosis (B) and fibrosis (C). The x-axis shows coefficients from the original analysis including all participants. Corresponding coefficients from the analyses of non-diabetic (left panel) and diabetic (middle-left panel) individuals are displayed alongside the statistical power for each subgroup analysis (middle-right and right panels). Red dots represent genes that were significant in the original analysis (q < 0.05). Subgroup analyses suggest that the results in the non-diabetic subgroup (n = 71) were highly consistent with findings from the original analysis (n = 94, adjusted for diabetes), indicating that the originally reported gene signatures, after correction for diabetic status, remain valid in non-diabetic individuals.

Figure 6–figure supplement 1. Liver fibrosis pathways and gene signatures
(A) Pathway analysis of the enrichment of pathways for the 213 fibrosis signatures. (B) TGF-β and SMAD related gene expression associated with fibrosis grades (x-axis) is depicted. Significance is indicated by black dots; * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 6–figure supplement 2. GTPases and their regulation emerges as a potential target for liver fibrosis.

(A) Heatmap of GTPase-related genes display a better correlation with fibrosis grades. Asterisks in the main heatmap denote the significance (p < 0.05) from comparisons of each subgroup. (B) Enrichment map of selected genes constituting the core of the protein-protein interaction network of GTPase-related genes. Nodes are colored according to BH (Benjamini-Hochberg) adjusted p-value, and node sizes are proportional to the number of genes in the term. (C) Correlation matrix of gene expression levels for TGF-β and GTPase-related genes. The correlation plot was created of Pearson's r values for GTPase-related genes and TGF-β genes (TGFB1 and TGFB3). The left annotation bar indicates the significance of associations between gene expression and fibrosis grades.

Figure 6–figure supplement 3. GTPase inhibition and Elafibranor treatment in LX2 cells.
(A) Workflow of the LX2 experiment in panel A-F (n = 8-10). (B) GTPase inhibitors NSC23766 (Rac1 inhibitor) and ML141 (Cdc42 inhibitor) significantly reduced pro-collagen secretion from HSC-like LX-2 cells and (C) inhibited gene expression of COL1A1 and COL1A2 under basal conditions. (D) TGFβ administration increased pro-collagen secretion by 32% and (E) collagen gene expression in LX2 cells. (F) NSC23766-mediated GTPase inhibition impairs pro-collagen secretion from LX-2 cells after TGF-β treatment. Pro-collagen secretion was determined by ELISA and gene expression levels were assessed by qPCR. Asterisks (*) denote p value < 0.05 for statistical significance from one-way ANOVA with Holms-Sidak multiple comparisons (B&C), unpaired t-test (D&E) Kruskal Wallis Test with Dunns multiple comparisons (F). (G) Gene signatures in LX-2 cells with and without Elafibranor (n = 3).

Figure 6–figure supplement 4. **GTPase-related genes in external systems**

(A) Human fibrosis signatures from the present study were partially reflected in a mouse study (GSE176042), where RNA sequencing was performed on hepatic stellate cells (HSCs) isolated from both acute and chronic mouse fibrosis models. GTPase-related genes, especially those involved in the initiation of fibrosis, were also upregulated in HSCs isolated from both acute and chronic liver fibrosis mouse models at different time points. Row annotations 'Linear' and 'Transition' indicate whether gene expression is linearly associated with fibrosis grades (Linear = 1) or shows a significant difference between patients with fibrosis and steatosis versus simple steatosis (Transition = 1). Asterisk denotes genes with adjusted p value < 0.05. (B) GTPase-related genes induced by TGF-β were elevated in hepatocytes and HSCs at different stages of a human liver organoid system (GSE207889). Significance was calculated by –log10(p value)*sign(fold change). Asterisk denotes genes with adjusted p value < 0.05. AH, adult hepatocyte-like; CHOL, cholangiocyte-like; DC, ductal cell-like; FH1, fetal hepatocyte 1-like; FIB, fibroblast-like; HB1, hepatoblast 1-like; HB2, hepatoblast 2-like; HSC, hepatic stellate cell-like; SMC, smooth muscle cell-like. (C) Schematic illustration of the hypothesized role of GTPase signaling in regulating liver fibrogenesis. TGF-β production is initiated in non-parenchymal liver cells, leading to the overexpression of GTPase regulators in both hepatic stellate cells and hepatocytes. This activation of GTPase signaling amplifies downstream fibrogenic effects, creating a feed-forward loop that intensifies fibrosis progression. Further investigation is needed to prove by which mechanisms GTPases regulate collagen expression and therefore fibrosis.

Figure 6–figure supplement 5. **Expression of GTPase-related genes in spheroid co-culture, hepatocyte monoculture, and LX-2 cells.** RNA-sequencing-based expression of GTPase-related genes in spheroid co-culture, hepatocyte monoculture upon control (Ctrl), TGF-β1 or TGFβ – inhibitor treatment (A, B). Expression of GTPase-related genes in control (Ctrl) and TGF-β1- treated (24h) LX2 cells measured by qPCR (C). Asterisks (*) denote p value < 0.05, (**) p < 0.005, (***) p < 0.0001 for statistical significance from unpaired t-test, numerical values above the bars indicate p-value. Bars were removed where data did not show strong directionality and statistical reliability, as indicated by (^) symbol. Symbol (@) indicates one or more expression values were not detected in the original dataset. Please note that ACTA2 expression in LX-2 cells is time dependent (unpublished results) and therefore we also included fibronectin (FN1) as a control for TGF-β1 treatment.