Title: From macro to micro: De novo genomes of Aedes mosquitoes enable comparative
genomics among close and distant relatives

Authors: Gen Morinaga[1], Darío Balcazar[2], Athanase Badolo[3], Diana Iyaloo[4], Theo Mouillaud[5],
Maria Sharakhova[6], Scott M. Geib[7], Christophe Paupy[5], Diego Ayala[5,8], Jeffrey R. Powell[2],
Andrea Gloria-Soria[2,9], John Soghigian[1]

Author Affiliations: [1]Faculty of Veterinary Medicine, University of Calgary, Calgary, AB,
Canada; [2]Department of Ecology & Evolution, Yale University, New Haven, CT, USA;
[3]Laboratoire d'Entomologie Fondamentale et Appliquée, Université Joseph Ki-Zerbo,
Ouagadougou, Burkina Faso; [4]Vector Biology & Control Division, Ministry of Health & Quality of
Life, Curepipe, Mauritius; [5]L'Institut de recherche pour le développment, UMR MIVEGEC,
Montpellier, France; [6]Department of Entomology, Virginia Polytechnic and State University,
Blacksburg, VA, USA; [7]USDA-ARS Tropical Pest Genetics and Molecular Biology Research Unit,
Hilo, HI, USA; [8]Medical Entomology Unit, Institut Pasteur de Madagascar, Antananarivo,
Madagascar; [9]The Connecticut Agricultural Experiment Station, New Haven, CT, USA.

Key words: genomes, yellow fever mosquito, comparative genomics

Word count (excludes literature cited): 8322/10000 (Main text + table + figure captions)
words. Abstract: 248/250 words. Significance statement 137/150 words.

2 Tables, 6 Figures

129/100 references

Abstract

The yellow fever mosquito (*Aedes aegypti*) is an organism of high medical importance because it is the primary vector for diseases such as yellow fever, Zika, dengue, and chikungunya. Its medical importance has made it a subject of numerous efforts to understand their biology. One such effort, was the development of a high-quality reference genome (AaegL5). However, this reference genome was sourced from a highly inbred laboratory strain with unknown geographic origin. Thus, the reference is not representative of a wild mosquito, let alone one from its native range in sub-Saharan Africa. To better understand the genetic architecture of *Ae. aegypti* and their sister species, we developed two *de novo* chromosome-scale genomes with sequences sourced from single individuals: one of *Ae. aegypti formosus* (Aaf) from Burkina Faso and one of *Ae. mascarensis* (Am) from Mauritius. Both genomes exhibit high contiguity and gene completeness, comparable to AaegL5. While Aaf exhibits high degree of synteny to AaegL5, it also exhibits several large inversions. We further conducted comparative genomic analyses using our genomes and other publicly available culicid reference genomes to find extensive chromosomal rearrangements between major lineages. Overrepresentation analysis of expanded genes in Aaf, AaegL5, and Am revealed that while the overarching category of genes that have expanded are similar, the specific genes that have expanded differ. Our findings elucidate novel insights into chromosome evolution at both microevolutionary and macroevolutionary scales. The genomic resources we present are additions to the arsenal of biologists in understanding mosquito biology and genome evolution.

Significance

Aedes aegypti is a major arboviral disease vector found throughout the tropics and sub-tropics. Its subspecies differ ecologically, as native sub-Saharan African form feeds on mammals generally and inhabit both sylvatic and domestic areas and the global invasive form preferentially feeds on humans and lives primarily domestic areas. Their medical importance has prompted the development of a high-quality reference genome, but it was sourced from an inbred laboratory strain of unknown origin. Here, we leveraged PacBio HiFi sequencing and HiC sequencing to develop the first de novo genome of *Ae. aegypti* sampled its native range in Burkina Faso. We also present a de novo genome of *Ae. mascarensis*, its sister species. Our genomes are comparably contiguous and complete to the reference genome. Comparative genomic analysis using our genomes and other culicid reference genomes reveal extensive chromosomal rearrangements.

Introduction

Mosquitoes (family Culicidae) are major vectors of arboviruses and parasites that cause minor to severe illnesses that affect millions of people globally (Yee et al. 2022). To mitigate the impact of mosquitoes on global public health, tremendous efforts have been made to elucidate mosquito biology. One such effort is the development of numerous genomic resources of mosquito species, especially those considered medically important such as *Anopheles* and *Aedes* mosquitoes. An example of these resources is the reference genome for *Aedes aegypti* (henceforth, 'AaegL5'; Matthews et al. 2018), the primary vector of arboviruses, including yellow fever, dengue, chikungunya, and Zika viruses (Pierson & Diamond 2020).

At the time of its development, AaegL5 utilized available technologies to achieve an accurate, complete, and contiguous chromosome-level genome assembly. Beyond improving upon the contiguity of previous versions of the reference (Dudchenko et al. 2017), AaegL5 also provided an improved set of gene annotations (Matthews et al. 2018), which allows for finer-scale mapping of genes and gene families. Indeed, the development of this reference has been invaluable—allowing researchers to study in detail *Ae. aegypti* transcriptomics (Herre et al. 2022), developmental biology (Herre et al. 2022), population genetics (Schmidt et al. 2020; Soghigian et al. 2020; Gloria-Soria et al. 2022), species distribution modeling (Rose et al. 2020), and phylogenomics (Soghigian et al. 2023). This resource is also valuable for elucidating biological differences between the subspecies *Ae. aegypti formosus* and *Ae. aegypti aegypti* (Aaf and Aaa, respectively) which differ in their bionomics—Aaf is a native to sub-Saharan Africa, inhabits both forested and domestic areas, and take blood meals from a variety of mammals; Aaa is a found globally outside of Africa, inhabits urban areas, and females have a strong affinity to feed on humans (Powell et al. 2018).

Despite its value as a genomic resource, AaegL5 is not without its shortcomings. The sequencing technology available at the time precluded the authors from using wild mosquitoes for two, co-related reasons. First, library construction requires high DNA input—a major hurdle for small insects such as flies and mosquitoes where a single individual may not be sufficient to extract enough source DNA, leading to assemblies where multiple individuals are pooled. Second, the long-read sequencing platforms available at the time were error-prone and required polishing using other, more accurate sequencing technologies (i.e., Illumina short-reads), which again would require more input DNA from additional individuals. Matthews et al., (2018) mitigated both challenges by sequencing 80 male siblings from a highly inbred laboratory strain (Liverpool), thereby reducing mis-assemblies resulting from high levels of heterozygosity (Whibley et al. 2021) whilst also extracting enough input DNA to meet library construction requirements.

A corollary of using an inbred laboratory colony for a reference genome, of course, is the reduction of heterozygosity and genetic diversity that it represents. This is not problematic *per se*—a single individual is a mere snapshot of the complex ebbs and flows of population dynamics within a single population. However, when laboratory colonies have measurably diverged from wild or source populations, they may no longer be representative of wild populations, leading to erroneous conclusions and problems with reproducibility (Brekke et al. 2018). Recent work (Gloria-Soria et al. 2019) has shown that laboratory strains of *Ae. aegypti* exhibit significantly lower genetic diversity than their wild counterparts. Furthermore, some strains do not cluster with any wild mosquito populations (CDC, Liverpool, and Orlando) or have diverged and/or become contaminated as strains were passed among laboratories (Liverpool

and Rockefeller; Gloria-Soria et al. 2019). Lastly, Gloria-Soria et al., (2019) showed that the
Liverpool strain, originally thought to have been sourced from West Africa (Macdonald 1962),
does not cluster with any Aaf populations.

Within Culicidae, chromosome-scale genome assemblies are taxonomically
concentrated in the Anophelinae, and these assemblies have allowed biologists to uncover
some of the intricate evolutionary dynamics that bridge the divide between microevolution and
macroevolution (Neafsey et al. 2015; Lukyanchikova et al. 2022). Recent efforts to expand the
phylogenetic sample of mosquitos have produced numerous chromosome-scale assemblies in
the Culicinae (e.g., Peng et al. 2021; Ryazansky et al. 2024). This effort to expand the
phylogenetic scope beyond Anophelinae allows biologists to further investigate the structural
variation between members of major clades (e.g., subgenera, tribes).

Here, we present the first *de novo* genome assembly of a wild *Ae. aegypti formosus*
(Aaf) specimen from West Africa and a *de novo* assembly of *Ae. mascarensis* (Am)—a partially
reproductively isolated sister species of *Ae. aegypti* found on the island of Mauritius in the south
west Indian Ocean (Hartberg & Craig Jr 1970). *Aedes mascarensis* diverged from Aaf roughly
8–10 MY (Soghigian et al. 2020). These two species are members of what is now known as the
Aegypti group (Le Goff et al. 2013; Soghigian et al. 2020). Each assembly was built from a
single wild-caught mosquito from their respective locales (Aaf: Burkina Faso; *Ae. mascarensis*:
Mauritius) using the recently developed Pacific Biosciences high fidelity (PacBio HiFi)
sequencing platform (Wenger et al. 2019). The high accuracy (99.999%) of PacBio HiFi reads
allows us to sequence a relatively highly heterozygous individual while the length of the reads
(>13 kbp) helps to span and resolve highly repetitive regions ubiquitous in the *Ae. aegypti*
genome (Matthews et al. 2018). We also compare new genomes with those of other mosquitoes
and place them in an evolutionary context to understand how genes and genomic structure
have changed across major culicid lineages. Lastly, we present a method for comparing draft
assemblies and implement it as a new, publicly available, *asmidx* package for R.

Methods

Sample collection

We sampled for wild mosquitoes in two locations: Ouagadougou, Burkina Faso, for *Aedes*
aegypti formosus (Oct. 2021); and Chamarel, Mauritius for *Ae. mascarensis* (May 2022). In both
locations, we collected mosquito eggs by placing ovitraps lined-up with seed-germination papr.
These eggs were shipped to The Connecticut Agricultural Experiment Station (CAES) for
rearing. We reared mosquitoes from eggs at CAES by submerging the paper containing the
eggs in deionized water and provided TetraMin® Tropical Flakes as *ad libidum* as food source
for larvae. Once mosquitoes pupated, we transferred the pupae into medicine cups filled with
larval water and placed them into insect rearing cages (12 x 12 x 12 inches/28 liters) where the
adults emerged. Larval trays and cages were maintained in an incubator at 27^{o}C with a 12:8
light/dark cycle throughout the rearing process. We provided adults with *ad libitum* sugar water
for 3–5 days until they were collected in dry ice for the DNA extraction protocol.

DNA sequencing for genome assembly and Hi-C genome-wide DNA cross-linking

To generate DNA sequences for PacBio HiFi for both Aaf and Am, we collected and froze adult
female mosquitoes on dry ice, then homogenized individuals with a sterile DNAse/RNAse free
plastic pestle. We eluted the homogenate using 180 µl of PBS and processed the solution using

the MagAttract HMW DNA Kit (Qiagen, Germantown, MD, USA) following the frozen tissue
protocol from the manufacturer. We then sent the extracted DNA to the Maryland Institute of
Genome Sciences of the University of Maryland for low input library preparation, where two
Sequel II 8M SMRT Cell runs (CCS/HiFi mode - 30 hour movie) were used to obtain sequences.

For HiC sequencing, we pooled multiple individuals (four females for *Ae. aegypti*
formosus and eight females for *Ae. mascarensis*) together and pulverized them in dry ice. We
then cross-linked samples and prepared *Ae. aegypti formosus* library with the Arima High
Coverage HiC kit and Arima HiC+ kit (Arima Genomics, San Diego, CA, USA), following
manufacturer protocols. For *Ae. mascarensis*, we prepared HiC libraries using the xGen ssDNA
& Low Input DNA Library Prep Kit (IDT, San Diego, CA, USA). We then sequenced both HiC
libraries at the Yale Center for Genome Analysis to achieve 100 million 150 bp paired-end reads.

Genome assembly

The PacBio HiFi sequencing platform and the programs (or the specific modes) built to handle
these data are still nascent, so we used four different assemblers and compared the outputted
draft assembly from each program. We used *HiCanu* (Nurk et al. 2020), *flye* (Kolmogorov et al.
2019), *hifiasm* (Cheng et al. 2021), and *IPA* (available at:
https://github.com/PacificBiosciences/pbipa). For all programs, we specified an estimated
haploid genome size of 1.3Gbp and used default settings and set flags necessary to assemble
PacBio HiFi reads (Supplementary Table S1).

We also compared the performance of two purging programs designed to identify and
remove duplicate haplotypes from the draft assemblies—*purge_haplotigs* (Roach et al. 2018)
and *purge_dups* (Guan et al. 2020) (henceforth, *ph* and *pd* respectively). Note that *hifiasm* and
IPA employ a purging step as part of their respective assembly pipelines by default—*hifiasm*
employs a variant of *pd* with a different algorithm for haplotype identification, and *IPA* simply
uses *pd*. Thus, in our workflow, assemblies output by these programs were purged twice, which
allowed us to assess "out-of-the-box" performance of all programs.

To assemble the mitochondrial genomes, we used the program *mitohifi* (Uliano-Silva et
al. 2023) in 'reads' mode and input the PacBio HiFi reads. This program was reference guided,
so for both species we used the *Ae. aegypti* complete mitochondrion found on GenBank
(OR544945.1).

***asmidx*: A holistic approach to assessing genome quality based on user input**

To quantify assembly metrics, we fed each intermediate assembly to the program *Inspector* ver.
1.0.1 (Chen et al. 2021). This program quantified basic assembly metrics, detects assembly
errors at the structural (expansion, collapses, inversions, and haplotype switches ≥50Bp) and
small scales (base substitutions, expansions, collapses <50Bp), and attempts to correct them.
We also assessed gene content completeness for each assembly generated using *BUSCO* ver.
5.2.2 and the Diptera OrthoDB data set ver. 10 (n = 3,285 single-copy orthologs).

Common genomic metrics (e.g., N50) and gene content of an assembly can be effective
indicators of assembly quality, but there is no consensus on which of these characteristics (or
what set of them) best characterizes a "good" genome assembly. Furthermore, it is unknown
whether each assembly program used in tandem with a purging program outputs similar quality
genome assemblies. To facilitate identifying the "best" assembly derived from the same set of

reads, we wrote an application using the *shiny* package ver. 1.7.4 for the R statistical
programming language ver. 4.2.1, which we call *asmidx* (available at
https://github.com/genmor/asmidx). The application takes as input a user-generated data set in
tabular format, with column headers where each row contains an assembly, and each column
contains a metric describing that assembly. The user can then select assembly metrics that
should be maximized and minimized to assess quality. Additionally, the user can identify a
genome size column and input a known genome size (e.g., from a reference genome) which will
be used to compute relative genome size differences, where smaller differences in relative
genome size will be considered better. Each chosen metric is then feature-normalized to be
between 0 (the worst) and 1 (the best). A row sum is then computed and multiplied by 100
resulting in normalized scores which are used to rank each assembly by quality. We also allow
users to differentially weight each of the selected characters. To make this process intuitive, we
allow for any positive value for weighting. The weights are multiplied to their respective feature-
normalized columns, row sums re-computed, and multiplied by 100 to output a weighted score.
For both weighted and unweighted scores, higher values will be associated with better
assemblies based on the quality metrics supplied (and weighted) by the user. The application
outputs these rankings in tabular format and visually represents them using a lollipop plot, both
of which the user can download. In addition to this shiny application, we wrote helper functions
to convert output from Inspector into tabular format.

To rank our assemblies, we considered four sets of metrics: gene content (duplicated,
fragmented, and missing BUSCOs), structural errors in the assembly (total number of expanded
bases, collapsed bases, and inverted bases), contiguity (N50), and genome size difference
relative to the reference. Our rankings maximized N50 and minimized all other metrics. We
weighted all structural errors and N50 by 0.1, duplicated and missing BUSCOs by 0.15,
fragmented BUSCOs by 0.125, and relative genome size difference by 0.175. For each
specimen, we used the assembly with the highest weighted score for scaffolding using Hi-C.

Scaffolding contigs to chromosomes via Hi-C and post processing

To prepare cross-linked, paired-end Illumina short-reads for use in scaffolding, we trimmed the
first five bases from the reads using the program *fastp* (ver. 0.23.4; Chen et al. 2018). We then
followed the Arima Genomics mapping pipeline (available at
https://github.com/ArimaGenomics/mapping_pipeline). This pipeline relies on *BWA* (ver. 0.7.17-
r1198-dirty; Li and Durbin 2009), *samtools* (ver. 1.15; Danecek et al. 2021), *picard* (ver. 2.2.4;
Broad Institute 2019), and custom Perl scripts for aligning the short-reads to a draft contig-level
genome assembly and preparing it for scaffolding (Table Program details). To scaffold, we used
the program *Yahs* (ver. 1.2a.2; Zhou et al. 2023). We manually curated the outputted scaffold-
level assembly using the programs *Juicer tools* and *Juicebox* (ver. 1.11.08; Durand et al. 2016)
to further remove duplicate contigs and correct mis-assemblies (i.e., inverted and mis-joined
contigs and scaffolds) and generated finalized contact maps for visualization purposes using
HapHiC (ver. 1.0.5; Zeng et al. 2024). After manually curating the scaffolds, we used *TGS-
gapcloser* (ver. 1.2.1; Xu et al. 2020) to close gaps in the assembly. We inspected gene content
of this assembly using *BUSCO* and then finalized the gap-filled assemblies by using a custom
script to remove "debris" sequences (i.e., contigs and scaffolds with duplicate HiC signal) and
scaffolds/contigs containing only duplicate BUSCOs that were not located on the chromosomal
scaffolds. We concatenated these "debris" sequences together with the duplicates detected by
the purging program. We then fed these assemblies to the *BlobToolKit* (ver. 4.2.1; Challis et al.

2020) suite to determine whether our assemblies contained sequences from contaminants or
endosymbionts and to output final summary statistics.

Validating structural variation and departures from AaegL5

To ascertain the validity of the structural variations we observed in Aaf and Am (relative to
AaegL5), we used NCBI BLAST (ver. 2.12.0+; Camacho et al. 2009) to create databases from
our scaffolded *Aaf* and *Am* assemblies to find positional hits of 88 *Ae. aegypti* bacterial artificial
chromosome (BAC) clone sequences (Matthews et al. 2018; Supplementary table S2). We
retained only the best hits (i.e., highest bit score) for each BAC clone and visualized their
positional order in both assemblies using the R package *ChromoMap* (ver. 4.1.1; Anand and
Rodriguez Lopez 2022; Supplementary figure S1). After this validation step, we aligned the
scaffolded *Aaf* assembly to AaegL5 using *minimap2* (Li 2018), and the resultant alignment file
fed into *SyRI* (ver. 1.6.3 ; Goel et al. 2019) which identified nucleotide synteny and structural
variation (i.e., duplications, translocations, and inversions). We performed the same analysis for
Am but chose not to interpret nucleotide synteny because it is too divergent from *Ae. aegypti*,
even after using less stringent *minimap2* settings (i.e., -asm20) and found the resulting output
uninterpretable (see Supplementary figure S2).

Genome structural annotation

We used the *RepeatModeler* pipeline (ver. 2.0.4; Flynn et al. 2020) to model and identify
repetitive elements in the genomes. After generating a *de novo* repeat database from our draft
assemblies, we soft-masked the assemblies using *RepeatMasker* (Smit et al. 2013) in four
iterations, passing each outputted soft-masked fasta files to the subsequent step: 1) mask only
simple repeats; 2) mask repeats using the -species flag with 'diptera' which queries the Dfam
database (ver. 3.7; Storer et al. 2021) for dipteran repeat sequences; 3) mask repeats identified
in *Ae. aegypti* (Nene et al. 2007) from TEfam repeat database; 4) mask repeats based on the *de*
novo repeat database created from *RepeatModeler*. We then quantified the diversity and
divergence relative to the consensus sequences of the repetitive content in Aaf and Am using
the '*calcDivergenceFromAlign.pl*' script included with *RepeatMasker.* This script estimated
divergence using the Kimura (K81) model of sequence evolution modified to account for the
high mutability of "CG" sites (Tsunoyama et al. 2001).

We input the final, soft-masked assemblies into the *BRAKER2* genome annotation
pipeline (ver. 3.0.3; Stanke et al. 2006b, 2008; Hoff et al. 2016, 2019; Brůna et al. 2021) with the
Arthropoda protein data set obtained from *OrthoDB* (ver. 11; Kuznetsov et al. 2023). *Braker*
uses *GeneMark-ES* (Lomsadze et al. 2005) and *ProtHint* (Brůna et al. 2020) to predict protein
coding genes in the assembly, then aligns these predicted proteins and regions using
DIAMOND (ver. 0.9.24.125; Buchfink et al. 2015) and *SPALN* (Iwata & Gotoh 2012). High-
confidence hits output by these programs are then fed into *GeneMark-EP+* (Brůna et al. 2020)
and *Augustus* (ver. 3.4.0; Stanke et al. 2006a) to output gene predictions. Following
recommendations from the program authors, we used the final output from *Augustus*. We then
used a Python script '*selectSupportedSubsets.py*' included with *Braker* to exclude genes
predicted without any external support (i.e., no support from OrthoDB). We used this output for
all analysis that involved the proteome. We assessed the quality of these annotations using
BUSCO in protein mode and *AGAT* (ver. 1.2.0; Dainat et al. 2023) to quantify annotation metrics
after we subset the output from *BRAKER2* to exclude genes supported only through
computational predictions and keeping only the longest isoforms.

Comparative genomic analysis

We compared the Aaf and Am assemblies with other Culicidae, including AaegL5, and nine
(eight culicid; one outgroup) other annotated, high-quality, chromosome-level reference
assemblies available on NCBI that represent the breadth of phylogenetic diversity of the family.
These assemblies were: *Anopheles cruzii* (subgenus *Kerteszia*)*, An. darlingi* (subgenus
Nyssorhynchus)*, An. gambiae* (subgenus *Cellia*)*, An. ziemanni* (subgenus *Anopheles*)*,
Armigeres subalbatus (tribe Aedini)*, Sabethes cyaneus* (tribe Sabethini)*, Culex pipiens pallens*
(tribe Culicini)*, Cx. quinquefasciatus* (tribe Culicini)*, and the sandfly, *Phlebotomous papatasi*
(Supplementary table S3). We assessed gene order synteny using the R package *GENESPACE*
(ver. 1.2.3; Lovell et al. 2022). *GENESPACE* uses *Orthofinder* (Emms & Kelly 2019) to identify
orthologous genes across a set of species and assesses synteny and collinearity of the
orthologs between all pairwise combinations of species using *MCScanX* (Wang et al. 2012). We
set AaegL5 as the reference for the riparian plot output by *GENESPACE.*

We also assessed the evolution of gene families across these genomes using part of the
compare_genomes pipeline (Paril et al. 2023). This pipeline chained several programs to: 1)
identify orthologous genes (*Orthofinder*); 2) infer a dated phylogeny using the orthologs (*IQ-
Tree2*; Minh et al. 2020); 3) assess gene family expansion and contraction (*Cafe5*; Mendes et al.
2021); and 4) use the PANTHER classification system (Mi et al. 2013) to assign gene family and
function. For phylogenetic inference, the pipeline set IQ-Tree2 to use a multi-partition model
(Chernomor et al. 2016) and performed model selection using ModelFinder (Kalyaanamoorthy
et al. 2017). It additionally used ultrafast bootstrap approximation (Hoang et al. 2018) to
estimate branch support and a least squares algorithm to date the inferred phylogeny (To et al.
2016). To analyze evolutionary gene expansion and contraction, we modeled genes families to
evolve at different rate categories through a γ-parameter with K = 4 categories in *Cafe5*. For
gene function and ontology, we used the biological process set (GO:0008150) from *An.
gambiae* (taxon ID: 7165), and the pipeline classified gene function using PantherHMM 17 (Mi
et al. 2013, 2019). Although *compare_genomes* performs GO enrichment and
overrepresentation as part of the pipeline, we elected to use the intermediate output to perform
our own. We did this by querying the PANTHER DB web tool (www.pantherdb.org; ver. 18.0;
accessed: Apr. 22, 2024) with a list of significantly expanded orthogroups (from *Cafe5*) for
AaegL5, Aaf, and Am (separately and together) to perform a statistical overrepresentation test
using Fisher's exact test using the 'GO biological process complete' annotation set of *An.
gambiae* and corrected for multiple testing by accounting for false discovery rate (Benjamini &
Hochberg 1995). We downloaded the full data table of results for each taxon and filtered each
list in R to include only those where $P_{FDR} < 0.01$ for a given taxon. We then examined the
semantic similarity of these overrepresented GO terms based on the method of Wang et al.
(2007), used K-means clustering to determine similar sets of terms, then corrected the P-values
a final time for multiple comparisons (again accounting for false discovery rate) using the R
package *simplifyEnrichment* (ver. 1.12; Gu and Hübschmann 2023). For a more holistic analysis
of the Aegypti group, we repeated the above overrepresentation test on PANTHER DB, this time
included all three expanded sets and outputting only significantly overrepresented GO terms
($P_{FDR} < 0.01$). Lastly, we took advantage of the fact that we identified gene families and their
positions in the Aaf genome, so we mapped the ones specifically located in putatively inverted
regions (relative to AaegL5). We matched gene identity and names by creating a local blast
database from AaegL5 and queried the orthologs in the inverted regions, limiting output to a
single alignment with e-values less than 1e-60.

Results

Identifying the best combination of HiFi assembler and purging program

We generated twelve draft assemblies each for Aaf and Am—four outputs directly from the assemblers and four of each output from purging programs *ph* and *pd* after taking the outputs from the assemblers as input and removing duplicated contigs and haplotigs. We provide detailed comparisons in the supplement (Supplemental results; Supplementary table S4). In brief, *asmidx* allowed us to identify *HiCanu* paired with *ph* output the best Aaf assembly, and *hifiasm* paired with *ph* output the best Am assembly (Fig. 1). While not free of arbitrary decisions, *asmidx* allows users to compare draft assemblies transparently and flexibly.

Final assembly characteristics

Scaffolding the assemblies using Hi-C, and manually curating the scaffolds using *Juicebox* (Durand et al. 2016) substantially improved the contiguity of the assemblies. For Aaf, N50 saw an 82-fold improvement and reduced L50 from 82 to 2, yielding an assembly with 706 scaffolds, 1,124 contigs, and total assembly size of 1.24 Gbp (Supplementary figures S3 and S4; Table 1). In terms of gene content, we detected 3,127 (95.2%) complete single-copy, 55 (1.7%) duplicated, 45 (1.4%) fragmented, and 58 (1.7%) missing BUSCOs from Diptera orthodb10 ortholog set (n = 3,285). For Am, scaffolding improved N50 by 30-fold and reduced L50 from 22 to 2, outputting an assembly with 74 scaffolds, 269 contigs, and a total assembly size of 1.29 Gbp (Supplementary figures S3 and S4; Table 1). We detected 2,990 (91%) single-copy, 179 (5.4%) duplicated, 48 (1.5%) fragmented, and 68 (2.1%) missing BUSCOs. We attempted to fill gaps in both assemblies using the HiFi reads, but neither resulted in dramatic reduction in scaffold count, although we were able to fill some gaps between contigs in both assemblies (Supplementary figure S3; Table 1). We additionally assembled the mitochondrial genome of these individuals, which yielded mitogenomes at roughly 16 kbp total size for both species.

Structural annotation of both assemblies showed their compositions were proportionally similar to one another (Fig. 2). Repetitive elements comprise the majority of genomic content (up to nearly 80%)—approximately 20% of the genomic content are classified as LTR retrotransposons and roughly 15% attributed to non-LTR retrotransposons (Fig. 2). In both assemblies, DNA transposons accounted for 6% of the genomic content, and approximately 1% of the assemblies were classified as helitrons (Fig. 2). Additionally, approximately 30% of both assemblies were considered repetitive, but unable to be classified (Fig. 2). For both assemblies, genomic repeat contents appear to have accumulated recently relative to the consensus repeat sequences with the peak occurring at 5% and 3% (Aaf and Am, respectively; Fig. 2). Unmasked, genomic content for both assemblies accounted for 22% (274 Mbp) and 21% (276 Mbp) of their genomes (Aaf and Am, respectively; Fig. 2). Exonic content accounted for 28 and 27 Mbp for Aaf and Am respectively (roughly 2% for both genomes) and intronic content accounting for 15% (211 Mbp) and 16% (230 Mbp) for Aaf and Am respectively. Lastly, monoexonic genes constituted 17% (3,120/17,009) and 16% (2,764/17,672) of the gene contents of Aaf and Am respectively.

Identifying putative endosymbiont and contaminant sequences

We assessed both the gap-filled assemblies and the alternate assemblies (e.g., assemblies consisting of haplotigs, duplicates, and 'debris') for any potential endosymbionts using the Blobtoolkit pipeline (Challis et al. 2020). Of the sequences in the primary Aaf assembly,

we traced 1.18 Gbp to Arthropoda, while the rest (54.4 Mbp) yielded no hits. In the alternate
assembly, we detected two sequences, a contig and a scaffold, totaling 1.59 Mbp originating
from α-proteobacteria (Supplementary figure S5). We isolated and removed this sequence from
the alternate Aaf assembly and queried the first 4400 bp of the scaffold sequence on BLAST.
The highest hit (99.89% identity) was for a *Rickettsia* endosymbiont found in *Cimex lectularius*
(GenBank Accession #: CP084572.1), while the next two highest hits (88.73%, 88.32%), were
similarly for *Rickettsia* endosymbionts isolated from *Oedothorax gibbosus* and *Culicoides*
impuctatus respectively (GenBank Accession #:OW370493.1; CP084573.1). In the primary Am
assembly, we traced 1.28 Gbp of the sequences to Arthropoda and the rest (7.74 Mbp) yielded
no hits. Unlike the Aaf alternate assembly, the Am alternate assembly did not contain any
sequences from that did not originate from either Arthropoda or yielded no hits.

Assessing structural rearrangement between AaegL5 and Aaf

We found a substantial degree of synteny between AaegL5 and Aaf, amounting to 1.08 Gbp or
87% of the total Aaf assembly exhibiting synteny with AaegL5 (Fig. 3). The Aaf assembly also
exhibited 175 inversions (most of which are small), totaling 44 Mbp or 3.5% of the total
assembly length. We found 870 translocations totaling 4.2 Mbp in length (0.34% of total
assembly length). The Aaf assembly also exhibited 490 instances of duplicated sequences
which totaled 1.9 Mbp in length (0.15% of total assembly length). Approximately 5% (61 Mbp
total length) of the Aaf assembly was not syntenic with AaegL5. We note two relatively large
inversions on chromosome 1—one located at 1*p*34 and one at 1*q*42 (1.63 and 1.84 Mbp long,
respectively; Fig. 3; Supplementary Table S5). We also detect a series of smaller inversions on
the telomeric end of the q-arm of chromosome 1. Additionally, we detected smaller inversions on
chromosome 3 located near 3*p*44 and two located on the telomeric end of the q-arm. As noted
in the methods, we make no attempt to interpret nucleotide synteny or structural variation at the
nucleotide level between Am and AaegL5 because they are too diverged. Nevertheless, we
show the synteny map between these assemblies in supplementary figure S2.

Gene order synteny across the Culicidae

Our analysis of gene order synteny revealed largely conserved patterns of chromosome
evolution within mosquito clades, but patterns between clades showed substantial chromosomal
rearrangement (Fig.5; Supplementary figure S6). Within the Aedini (*Aedes* and *Armigeres*) and
Culicini (*Culex*), our analysis showed largely syntenic patterns (i.e., similar gene order) between
the assemblies. However, *Anopheles* mosquitoes showed substantial chromosomal
rearrangements between the subgenera that we included (Fig. 5). We found whole-arm
translocations between chromosomes 2 and 3. In fact, translocation whole-arm translocations
between chromosomes 2 and 3 appear to be pervasive when viewed at greater evolutionary
scales. Without account for the translocation of chromosome 1 genes in Aedini onto
chromosome 3 in *Anopheles,* we find five distinct arm associations in chromosomes 2 and 3
(Fig. 5; Supplementary figure S6)—one arm association among the Aedini, one for *Sabethes*,
one for *Culex*, one for *An. cruzii* and *An. darlingi*, and one for *An. gambiae.* Interestingly, our
analysis suggests *An. ziemanni* have the same autosomal arm associations as the Aedini while
maintaining the same chromosome composition as other *Anopheles* (Fig. 5; Supplementary
figure S6). Our analysis showed largely syntenic gene order arrangement between AaegL5, Aaf,
and Am. This analysis found a similar set of inversions between AaegL5 and Aaf on
chromosome 1 (Fig. 5; Supplementary figure S6). The same regions appeared to be inverted
between Aaf and Am (Fig. 5; Supplementary figure S6). We detected an additional inversion

between Aaf and Am at the distal end of the q-arm of chromosome 1. Lastly, we note that *P.*
papatasi has N=5 chromosomes—three macrochromosomes (> 40 Mbp) and two
microchromosomes (< 20 Mbp), but only three macrochromosomes are shown in figure 5. This
was likely because too few orthologs were detected on the microchromosomes to adequately
assess synteny. Nevertheless, we detected considerable gene order rearrangement in *P.*
papatasi (Fig. 5)—genes originating from AaegL5 chromosome 2 comprised much of
chromosomes 2 and 3 in *P. papatasi*, while chromosome 1 of *P. papatasi* was composed of
genes originating from chromosomes 1 and 3 of AaegL5.

Gene family evolution in the Culicidae

We found 6,559 common orthologs between the twelve species included in our analysis
(Supplementary figure 7), 3,537 of which were single copy (Fig. 5). We found that both Aaf and
Am had more orthologs in total (17,672 and 17,009 respectively) when compared to AaegL5
(14,626) (Fig. 5). We also found that Aaf and Am had more orthologs exclusively in common
with one another than with AaegL5 (Fig. 5). Our analysis found 12,924 multi-copy orthologs in
Aaf, 12,639 multi-copy orthologs in Am, and 10,538 multi-copy orthologs in AaegL5.
Furthermore, we detected 639 paralogs unique to Aaf, 447 paralogs unique in Am, and 285
unique paralogs in AaegL5 (Supplementary figure S7. Our analysis failed to assign orthology to
639 genes in Aaf, 447 genes in Am, and 285 genes in AaegL5 (Supplementary figure S7).

Across the culicid assemblies we included, *Ar. subalbatus* had the highest ortholog count
(19,040), followed by Aaf and Am (Supplementary figure S7). These three assemblies also
comprised the top three in terms of unique paralogs and unassigned genes. Anophelines had
138 orthologs exclusively common among them, while species in the Culicini had 320 orthologs
exclusively common among them. At more granular evolutionary scales, *Culex* species
exhibited 953 exclusive orthologs, while *Aedes* species exhibited 218 exclusive orthologs.

We used the maximum likelihood phylogeny output from IQ-Tree2 with 8,742,672 sites
aligned across 3,534 partitions for the taxa in our analyses. We re-rooted the outputted
phylogeny using *P. papatasi* as the outgroup, which showed strong monophyly of the Culicidae
(Fig. 5). The topology of our phylogeny was largely congruent to that of Soghigian et al., (2023)
with the exception of the placement of *Sabethes* (Fig. 5). Our phylogeny placed *Sa. cyaneus*
(thus, Sabethini) as sharing a more recent common ancestor with the Aedini than the Culicini,
contrary to the results of Soghigian et al. (2023), and more similar to those of Reidenbach et al.
(2009).

Our analysis of gene family expansion and contraction showed substantial variability in
gene family gains and losses, the majority of which occurred at the species level (Fig. 5). In
addition to raw numbers of gains and losses of gene families, rapid gene family evolution (i.e.,
gene families with statistically significant changes in count AND categorized to have higher than
average rates – reflected by blue numbers to the right of nodes in Fig. 5) appears to have
happened at or near the tips rather than toward the root (Fig. 5). Indeed, our analysis showed
that deeper internal nodes tended to have very few quickly evolving gene families than more
relatively shallow nodes, such as the nodes leading to *Aedes* and the ancestor of *Culex* (Fig. 5).
Despite the lack of quickly evolving gene families, our analysis did suggest an overall gene
family expansion in the subfamily Culicinae, and contraction in the subfamily Anophelinae (Fig.
5). Between tribes in the Culicinae, our results indicated that the Aedini saw much greater gene
family expansion than contractions (Fig. 5). Of the three *Aedes* genomes, both Aaf and AaegL5

saw roughly similar number of gene family contractions and expansions, Am saw a higher
number of expansions than contractions (Fig. 5). Among tips, Aaf and AaegL5 have the highest
number of rapidly evolving genes—most other taxa had roughly half or fewer rapidly evolving
genes (Fig. 5). The distribution of these appear to differ between assemblies, though in general,
rapid changes in gene count appear to be gains (Supplementary figure S8). Notably, the
assemblies that exhibit rapid losses tend to be those derived from laboratory strains (i.e.,
AaegL5, *Cx. pipiens pallens*, *Cx. quinquefasciatus*, *Sa. cyaneus;* Supplementary figure S8).

We assessed the biological process gene ontology (GO) terms associated with each
gene family and highlighted the most reoccurring, significantly expanded or contracted gene
families with GO annotations by total number of copies represented among the species included.
In general, while many gene families have expanded and contracted since diverging from each
taxon's recent common ancestor, we did not detect any changes that would be indicative of a
pattern particular of any group of taxa (Supplementary figure S9), reflecting that most
differences appeared to be between tips, rather than between genera or higher taxonomic
rankings. In our analysis, we detected two sets of orthologs encoding odorant receptors, totaling
five instances of significant count changes (Supplementary figure S6). We detected two different
orthologs of rho-guanine exchange factor-related protein that had three total instances of
significant copy number changes (Supplementary figure S9). E3 ubiquitin-protein ligase trip12
and glucose-methanol-choline oxidoreductase were each assigned to two different orthologs,
each significantly changing in copy number once (Supplementary figure S9). We found seven
other protein families that were each assigned orthologs whose copy number significantly
changed twice: cyclic nucleotide-gated cation channel subunit A, fatty acid acyl transferase-
related, malic enzyme-related, nipped-b-like protein delangin SCC2-related, oxidoreductase
Glyr-1-related, scaffold attachment factor B-related, and voltage gated potassium channel
(Supplementary figure S9).

Comparison of gene ontology between Aaf, Am, and AaegL5

We assessed gene function at a finer scale across the three *Aedes* genomes by assessing
significantly overrepresented GO terms among the set of significantly expanded genes common
to these genomes. We used K-means clustering to group the 194 GO-terms common across the
three genomes into 11 clusters, three of which appeared to describe metabolic processes (Fig.
6). The remaining eight clusters were loosely described as processes vital to behavior—sensory
perception and detection of chemicals, ion transport, male mating and reproductive behavior,
synaptic signaling and signal transduction, and cellular organization and biogenesis (Fig. 6). We
did not detect any notable commonalities between the taxa and the sets of overrepresented GO
terms (Fig. 6). When we repeated the overrepresentation test with all three taxa in a single
analysis, we found similar results, albeit with substantially fewer GO terms overrepresented
(Table 2; Supplementary table S6).

Gene families located in putative inversions in Aaf

By taking advantage of the gene families identified, we mapped 413 genes in the putatively
inverted regions of Aaf chromosomes (relative to AaegL5), of which 354 were also identified in
AaegL5 (Supplementary table S7) Notable genes among those found (Supplementary figure
S10; supplementary table S8) were different odorant receptors (*Or4*) odorant binding proteins
(Obp 56a and d), and ion channels (Shaker, *NaCh*) [important for signal transduction (Bohbot et
al. 2007; McBride et al. 2014; Matthews et al. 2016)]; E3 ubiquitin ligases (*RNF 19B* and *RNF*

126) [similar E3 ubiquitin ligases are implicated in susceptibility to flavivirus infection (Giraldo et
al. 2020; Dubey et al. 2022)]; heat shock proteins [important for heat and dehydration tolerance
(Zhao et al. 2009; Benoit et al. 2010, 2011)]; and cytochrome P450 and adult and larval cuticle
proteins [important for insecticide resistance (Poupardin et al. 2010)].

Discussion

De novo assembly of wild, individual *Aedes* mosquitoes

Small body size and a high input DNA requirement have been major hurdles to producing high-
quality, chromosome-scale genome assemblies from many wild insects. Recent advances in
sequencing technologies that generate highly accurate, long-reads, like PacBio HiFi (Wenger et
al. 2019), allowed us to obtain enough high quality reads from a single mosquito to use for the
first *de novo* genome assembly of *Aedes aegypti formosus* and *Ae. mascarensis*, avoiding the
need of rearing colonies in the laboratory to obtain sufficient material. One additional roadblock,
albeit minor compared to issues such as DNA input requirements, is determining which
combinations of varied software tools produces the best assembly, particularly when
considering numerous genome assembly metrics. *asmidx* allowed us to overcome this hurdle,
choosing the best assembly from a range of excellent assemblies produced by a variety of
genomic tools. Combined with Hi-C aided scaffolding (Burton et al. 2013; Dudchenko et al.
2017), the resulting genome assemblies from our pipelines are both highly contiguous and
highly complete (Table 2; Supplementary figures S3 and S4). Using these chromosome-level
assemblies together with other high quality Culicid reference genomes, we conducted a series
of phylogenomic and comparative genomic analyses. The phylogenomic analysis (Fig. 5)
showed minor differences to those recently published (Soghigian et al. 2023), likely due to
substantial differences in taxonomic (i.e., number of taxa and lineages) and genetic sampling
(i.e., whole genomes vs. sequence capture). The comparative genomic analysis revealed
notable structural differences across large phylogenetic distances (Fig. 4) and numerous
insights on the evolution gene families in the Culicidae (Figs. 5 and 6, Supplementary figures S8
and S9). We detail the implications of these findings below.

Comparing Aaf, Am, and AaegL5

The assemblies of *Aedes aegypti formosus* (Aaf) and *Ae. mascarensis* (Am) had genome sizes
comparable to AaegL5 assembly size (Supplementary figure S3; Table 1) and within range of
genome size estimates from flow cytometry (Matthews et al. 2018). Both assemblies exhibited
comparable gene content and accuracy to the *Ae. aegypti* reference genome, AaegL5
(Supplementary figure S3; Table 1) with a high degree of synteny between homologous
chromosomes (Fig. 3). However, we also found evidence of inversions on Aaf chromosomes
relative to AaegL5, though they require further testing via PCR for veracity. Recent studies
(Redmond et al. 2020; Liang et al. 2024) have described numerous inversions in each of the
chromosomes African and global populations of *Ae. aegypti*. In our *Ae. aegypti formosus*
assembly (Aaf), we detected several of the same relatively large inversions that they discovered
on 1*p*34, 1*q*42, and 3*q*43 (Fig. 3; Liang et al. 2024). The inversions on 1*p*34 and 3*q*43 are
common among African populations of *Ae. aegypti* (referred to as 1pA and 3qG respectively by
Liang et al. 2024). Furthermore, the inversion we detected on 1*q*42 is positioned similar to 1qF
or 1qG detected in Burkina Faso populations Ouahigouya (OHI) and Ouaga-dougou (OGD).
Structural variations, such as inversions (Supplementary table S5), rearrange gene order, which
in turn can lead to adaptive phenotypes that are shielded from recombination (reviewed in:

Wellenreuther & Bernatchez 2018; Wellenreuther et al. 2019). This phenomenon is well-
documented among *Anopheles* mosquitoes, wherein inversions are associated with numerous
local adaptations (Powell et al. 1999; Cheng et al. 2018; Ayala et al. 2014) and genomic
diversity (The Anopheles gambiae 1000 Genomes Consortium 2020). Inversions in *Ae. aegypti*
have a long history of interest (Macdonald & Sheppard 1965) with recent research focusing on
identifying inversions in different populations in both subspecies (Dickson et al. 2016; Redmond
et al. 2020; Liang et al. 2024). The adaptive effect of these inversions remains unclear, however
some genes that have been identified occur in inverted regions and have identified phenotypes.
For example, odorant receptor 4 (*Or4*; McBride et al. 2014) located near the 1*q*42 position have
been linked to preference for human odor. Other genes may have implications for vector
management and adaptation. For example, over-expression and diversification of cuticle
proteins are implicated in insecticide resistance in many insects (reviewed in: Balabanidou et al.
2018). Similarly, upregulation or increased copy number of heat shock proteins may contribute
to more readily adaptable populations under increasing global temperatures (but see: Ware-
Gilmore et al. 2023). A targeted, transcriptomic approach is necessary to further interrogate how
genes in these inverted regions are expressed and their phenotypic consequences.

Aedes aegypti and *Ae. mascarensis* diverged approximately 8–10MY (Soghigian et al.
2020), thus we expected overall similarity between genome structure, especially in genic
regions between the three *Aedes* assemblies. Indeed, our results showed a high degree of gene
order synteny between these assemblies (Fig. 5) and the holistic set of GO term clusters show
key clusters of genes with similar functions appear to be overrepresented (Fig. 6; Table 2;
Supplementary table S6). For example, GO term clusters that describe mating and reproductive
behavior as well as sensory perception appeared to be commonly overrepresented in all three
taxa, perhaps because they are highly consequential to fitness (Cabrera & Jaffe 2007), and
could represent differences in mating behaviors unique either to *Aedes* mosquitoes or to the
Aegypti Group. However, an equally intriguing observation is that while many of the overarching
clusters (i.e., mating/reproductive behavior, sensory perception, metabolism) are similar, the
specific set of genes and GO terms assigned to them appear to vary in the Aegypti group, even
between Aaf and AaegL5 (Fig. 6). These differences may be the manifestations of local
adaptations—to environmental conditions where the mosquitoes were sampled (for Aaf and Am)
or to the laboratory (for AaegL5). Indeed, numerous studies have documented local adaptations
in *Aedes aegypti aegypti* to climatic/environmental (Soudi et al. 2023), altitude (Kramer et al.
2023), and vector competency for dengue serotypes (Lambrechts et al. 2009). Another
possibility is that the source population for AaegL5 is not necessarily representative of wild *Ae.
aegypti* (Gloria-Soria et al. 2019). Expanding the taxonomic sampling to other members of the
Aegypti group may shed light on what functional genetic differences exist within the group.

An examination of the repetitive content in Aaf and Am found a notable departure from
what was originally reported in AaegL5 (Fig. 2; Matthews et al. 2018). Indeed, whereas
Matthews et al. (2018) found that 65% of the AaegL5 genomic sequence was considered
repetitive, our assemblies showed nearly 80% of the genomic content to be repetitive. Other *Ae.
aegypti* assemblies whose repeat content is characterized (Aag2 cell line: Whitfield et al. 2017;
ROCK chromosome-scale assembly: Fisher et al. 2022) reported levels similar to Matthews et
al. (2018). However, a recent re-examination found 78% of the AaegL5 genome to be repetitive
DNA (Ryazansky et al. 2024) and thus similar to what we detected in Aaf and Am. The repeat
landscapes for both Aaf and Am assemblies (Fig. 2 A, B) are similar to those reported by
Whitfield et al. (2017), with sequence divergence peaking close to zero relative to the

consensus sequences. These landscapes, particularly that of long terminal repeat (LTR)
retrotransposons, represent recent activation and thus may represent recent infection from an
RNA virus (Whitfield et al. 2017). In similar vein, the

A *Rickettsia* endosymbiont found in *Aedes aegypti formosu*s

We discovered an endosymbiont in our *Aaf* assembly (Supplementary figure S5). The genome
size of this endosymbiont (1.59 Mbp) was similar to the Torix-group *Rickettsia* endosymbiont
first detected in *Culicoides impunctatus* (Davison et al. 2022). While *Rickettsia* are better known
for causing typhus fever and spotted fever (Raoult & Roux 1997), work conducted in the past
20+ years has revealed their extensive association with invertebrates and their tendency to
manipulate host reproduction (see Perlman et al. 2006 and sources cited therein). This body of
work has revealed their taxonomic diversity (Perlman et al. 2006; Weinert et al. 2009; Davison
et al. 2022) and the diversity of their host range (e.g., Kikuchi et al. 2002; Thongprem et al.
2021), but their effects on hosts are still understudied. The most documented effect of many
invertebrate-affecting *Rickettsia* is host reproductive manipulation, similar to those of *Wolbachia*
(reviewed in: Werren 1997; Perlman et al. 2006). These endosymbionts are vertically
transmitted from infected mother to her offspring, thus hijacking host reproduction to benefit
themselves. These effects generally lead to a female-biased sex-ratio either by killing males
(Werren et al. 1994) or inducing parthenogenesis (Hagimori et al. 2006; Aguin-Pombo et al.
2021). Torix-group *Rickettsia* have a direct effect on body size of host leech species, wherein
infected individuals exhibiting larger body sizes (Kikuchi et al. 2002), and low dispersal among
infected spiders (Goodacre et al. 2006). The prevalence or the effect of this *Rickettsia* in *Ae.*
aegypti formosus from Burkina Faso is unknown.

Repeated genomic rearrangement in Culicidae

Recent advances in sequencing technologies, physical mapping, and three dimensional
chromosomal structure inference have made evident the extensive evolutionary rearrangement
of chromosomes in Culicidae (Sharakhov et al. 2002; Neafsey et al. 2015; Palatini et al. 2020;
Yurchenko et al. 2023; Ryazansky et al. 2024; Lukyanchikova et al. 2022). Microscale structural
variants that confer local adaptations (Ayala et al. 2014; Powell et al. 1999; Cheng et al. 2018)
and macroscale whole-arm translocations detected between species are well-studied in
Anopheles (Sharakhov et al. 2002, 2016; Neafsey et al. 2015; Wei et al. 2017; Artemov et al.
2018). In the Culicinae, Arensburger et al. (2010) detected whole-arm translocations between
Ae. aegypti and *Cx. quinquefasciatus,* which Ryazansky et al. (2024) recently confirmed. Our
analysis expand on the scope of these studies by including more taxa from across the Culicidae
and show several intriguing trends. First, associations of chromosome arms have repeatedly
changed throughout the evolutionary history of the Culicidae (Fig. 4; Supplementary figure S6).
Indeed, relative to AaegL5, all non-Aedini genomes we investigated showed whole-arm
translocations between chromosomes 2 and 3 (Fig. 4, Supplementary figure S6; also see:
Arensburger et al. 2010; Neafsey et al. 2015; Ryazansky et al. 2024). Second, in stark contrast
to our first point, chromosomes 2 and 3 showed stability in the Aedini, as none of the
assemblies in the tribe showed whole-arm translocations (Fig. 4; Supplementary figure S6). Our
taxonomic sample cover roughly 50MY of evolutionary history in the Aedini, and in a similar
timeframe, each of the anophelines in our dataset evolved to exhibit unique chromosomes 2
and 3 arm associations, again in line with the findings of Neafsey et al. (2015), wherein syntenic
blocks rapidly decayed in that timeframe. Lastly, despite multiple major rearrangements, most
culicids exhibit a karyotype 2N = 6, with the sole exception being *Chagasia bathana* (subfamily

Anophelinae), where 2N = 8 (Rai & Black 1999). This level of conservation is remarkable in the
context of other arthropods such as Coleoptera (Blackmon et al. 2024), Lepidoptera (Wright et
al. 2024), and within Diptera (Morelli et al. 2022). To better-interrogate chromosome evolution
within Culicidae, more high quality, chromosome-scale genome assemblies are required,
especially within the Culicinae.

Evolutionary changes in distribution and copy number of gene families

Changes in copy number of key gene families may play key adaptive roles in mosquitoes
leading to differences in vector effectiveness between species (Arcà et al. 2017; Palatini et al.
2017; Catapano et al. 2023) and populations (Lambrechts et al. 2009; Bennett et al. 2021). We
compared our *Ae. aegypti formosus* (Aaf) and *Ae. mascarensis* (Am) genomes to high quality
mosquito genome assemblies that were publicly available and found striking differences both in
the distribution of rapidly evolving orthologs (Supplementary figure S8) and ortholog copy
number (Supplementary figure S9).

Rapid changes in copy number may also be an indication of adaptation (Simon et al.
2015; Xie et al. 2018). Rapid gains, in particular have been attributed to adaptive evolutionary
changes, as duplicated gene copies are "released" from stabilizing selective pressures may
respond adaptively as the environment or the context in which they are expressed changes
(Guo & Kim 2007; Vieira et al. 2007), however other works have also shown rapid losses to also
lead to adaptation (McBride & Arguello 2007; Goldman-Huertas et al. 2015). Rapid gains in
orthologs appear in all mosquito assemblies in our data set (Supplementary figure S8), however
rapid losses appear to have happened more often among laboratory strains. It is unclear
whether this is a pattern of adaptation to laboratory conditions (Gloria-Soria et al. 2019; Ross et
al. 2019) or an artifact (e.g., sampling bias) of the available genomic resources of mosquitoes. A
more detailed examination with more diverse sampling of laboratory strains would be necessary
if there is a tendency for rapid gene loss among laboratory strains compared to wild populations.

At more granular levels, we detected eleven different reoccurring rapidly evolving gene
families in our data set (Supplementary figure S9). Among them, we detected those involved in
sensory processes (odorant receptors) and signaling cascades (CNG cation channels)
reoccurred most often (Fig. 5). These proteins are crucial for detecting and transducing olfactory
signal (Zwiebel & Takken 2004; Sato et al. 2008) and thus key for host detection. Our analysis
showed two sets of orthologs that encode odorant receptors—one that has contracted in *Cx.*
quinquefasciatus and *Ar. subalbatus* but expanded in *Cx. pipiens pallens*, and another that has
expanded in *Cx. quinquefasciatus* and *Ae. aegypti* (AaegL5) (Supplementary figure S9). No
clear pattern of blood host affinity (based on: Soghigian et al. 2023) arises from these
combination of taxa and the orthologs we detected.

Conclusion

Here, we presented a *de novo* genome assembly of a wild caught *Ae. aegypti formosus* (*Aaf*)
and *Ae. mascarensis* (*Am*) each derived from a single, wild-caught individual. Our assemblies
are comparable to the reference *Ae. aegypti* assembly (AaegL5; Matthews et al. 2018) in terms
of contiguity and gene content but differ in that the Aaf assembly exhibits genomic structural
variation particular to West Africa (Liang et al. 2024), and notable differences in ortholog counts
and their functions. At 8–10 MY diverged, we view the nucleotide synteny between Am and
AaegL5 as unreliable, but gene order in *Ae. mascarensis* is highly conserved and show a high
degree of synteny with *Ae. aegypti*. With the three *Aedes* genome assemblies we also showed

variation in gene family expansion, and the function of those expanded gene families, showing
population- (between AaegL5 and Aaf) and species- (between *Ae. aegpyti* and *Ae.*
mascarensis) wide differences. These assemblies will be valuable assets for future studies to
understand the biology and evolution of *Ae. aegypti*—the *Aaf* assembly more closely reflects
natural populations of *Ae. aegypti* in the ancestral range, and the Am assembly provides the
closest genome output to date for this species. We used our newly assembled genomes along
with ten other reference genomes evolutionary changes within the Culicidae and find repeated
bouts of major chromosomal rearrangements, particularly between chromosomes 2 and 3. The
genomes we present here represent initial steps toward the development genomic resources for
all of the currently described taxa in the Aegypti group (Soghigian et al. 2020). Further
development within this group can elucidate genomic architecture that differentiates the ecology
and behavior of the African and the global invasive sub-species.

Data Availability

The PacBio HiFi reads generated for this project will be deposited in GenBank within
BioProjects (Am_MascCH02 principal: PRJNA1199517, Am_MascCH02 alternate:
PRJNA1199516, Aaf_Bf05 principal: PRJNA1199519, Aaf_Bf05 alternate PRJNA1199518)
under SRA accession XXXXX and XXXXX. The GenBank accession number for the assemblies
we generated are: Am_MascCH02 principal: XXXX, Am_MascCH02 alternate: XXXXXX,
Aaf_Bf05 principal: XXXXXX, Aaf_Bf05 alternate: XXXXXXX, and all available at NCBI.

Acknowledgements

Financial support for this project was provided by NIAID R01 AI101112 awarded to JRP. We also
acknowledge the support of the Natural Sciences and Engineering Research Council of Canada
(NSERC), funding reference number RMS21-73779779 [Cette recherche a été financée par le
Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), numéro de
référence RMS21-73779779]. GM was supported by NIAID R01 AI155562 awarded to JRP, AG-
S, and JS. We thank T. Petruff, J. Brophy, S. Arent, and R. Pellegrini for support with sample
processing. We also thank L. Jackson for providing comments on an earlier version of this
manuscript. Lastly, we thank the Research Computing Services group at the University of
Calgary.

Literature Cited

Aguin-Pombo D, Rodrigues MCPA, Voetdijk B, Breeuwer JAJ. 2021. Parthenogenesis and Sex-Ratio Distorting Bacteria in *Empoasca* (Hemiptera: Cicadellidae) Leafhoppers. Annals of the Entomological Society of America. 114:738–749. doi: 10.1093/aesa/saab025.

Anand L, Rodriguez Lopez CM. 2022. ChromoMap: an R package for interactive visualization of multi-omics data and annotation of chromosomes. BMC Bioinformatics. 23:1–9. doi: 10.1186/s12859-021-04556-z.

Arcà B, Lombardo F, Struchiner CJ, Ribeiro JMC. 2017. Anopheline salivary protein genes and gene families: an evolutionary overview after the whole genome sequence of sixteen Anopheles species. BMC Genomics. 18:153. doi: 10.1186/s12864-017-3579-8.

Arensburger P et al. 2010. Sequencing of *Culex quinquefasciatus* Establishes a Platform for Mosquito Comparative Genomics. Science. 330:86–88. doi: 10.1126/science.1191864.

Artemov GN et al. 2018. Partial-arm translocations in evolution of malaria mosquitoes revealed by high-coverage physical mapping of the Anopheles atroparvus genome. BMC Genomics. 19:278. doi: 10.1186/s12864-018-4663-4.

Ayala D, Ullastres A, González J. 2014. Adaptation through chromosomal inversions in *Anopheles*. Frontiers in Genetics. 5. https://www.frontiersin.org/articles/10.3389/fgene.2014.00129 (Accessed October 4, 2022).

Balabanidou V, Grigoraki L, Vontas J. 2018. Insect cuticle: a critical determinant of insecticide resistance. Current Opinion in Insect Science. 27:68–74. doi: 10.1016/j.cois.2018.03.001.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological). 57:289–300. doi: 10.1111/j.2517-6161.1995.tb02031.x.

Bennett KL, McMillan WO, Loaiza JR. 2021. The genomic signal of local environmental adaptation in *Aedes aegypti* mosquitoes. Evolutionary Applications. 14:1301–1313. doi: 10.1111/eva.13199.

Benoit JB et al. 2011. Drinking a hot blood meal elicits a protective heat shock response in mosquitoes. Proceedings of the National Academy of Sciences. 108:8026–8029. doi: 10.1073/pnas.1105195108.

Benoit JB, Lopez-Martinez G, Phillips ZP, Patrick KR, Denlinger DL. 2010. Heat shock proteins contribute to mosquito dehydration tolerance. Journal of Insect Physiology. 56:151–156. doi: 10.1016/j.jinsphys.2009.09.012.

Blackmon H, Jonika MM, Alfieri JM, Fardoun L, Demuth JP. 2024. Drift drives the evolution of chromosome number I: The impact of trait transitions on genome evolution in Coleoptera. Journal of Heredity. 115:173–182. doi: 10.1093/jhered/esae001.

Bohbot J et al. 2007. Molecular characterization of the *Aedes aegypti* odorant receptor gene family. Insect Molecular Biology. 16:525–537. doi: 10.1111/j.1365-2583.2007.00748.x.

Brekke TD, Steele KA, Mulley JF. 2018. Inbred or Outbred? Genetic Diversity in Laboratory Rodent Colonies. G3 Genes|Genomes|Genetics. 8:679–686. doi: 10.1534/g3.117.300495.

Broad Institute. 2019. Picard Toolkit. https://broadinstitute.github.io/picard/ (Accessed February 20, 2024).

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR Genomics and Bioinformatics. 3:lqaa108. doi: 10.1093/nargab/lqaa108.

Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP+: eukaryotic gene prediction with self-training in the space of genes and proteins. NAR Genomics and Bioinformatics. 2:lqaa026. doi: 10.1093/nargab/lqaa026.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. Nat Methods. 12:59–60. doi: 10.1038/nmeth.3176.

Burton JN et al. 2013. Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. Nat Biotechnol. 31:1119–1125. doi: 10.1038/nbt.2727.

Cabrera M, Jaffe K. 2007. An aggregation pheromone modulates lekking behavior in the vector mosquito *Aedes aegypti* (Diptera: Culicidae). moco. 23:1–10. doi: 10.2987/8756-971X(2007)23[1:AAPMLB]2.0.CO;2.

Camacho C et al. 2009. BLAST+: architecture and applications. BMC Bioinformatics. 10:421. doi: 10.1186/1471-2105-10-421.

Catapano PL et al. 2023. De novo genome assembly of the invasive mosquito species *Aedes japonicus* and *Aedes koreicus*. Parasites Vectors. 16:427. doi: 10.1186/s13071-023-06048-w.

Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. 2020. BlobToolKit – Interactive Quality Assessment of Genome Assemblies. G3 Genes|Genomes|Genetics. 10:1361–1374. doi: 10.1534/g3.119.400908.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 34:i884–i890. doi: 10.1093/bioinformatics/bty560.

Chen Y, Zhang Y, Wang AY, Gao M, Chong Z. 2021. Accurate long-read de novo assembly evaluation with Inspector. Genome Biology. 22:312. doi: 10.1186/s13059-021-02527-4.

Cheng C, Tan JC, Hahn MW, Besansky NJ. 2018. Systems genetic analysis of inversion polymorphisms in the malaria mosquito *Anopheles gambiae*. Proceedings of the National Academy of Sciences. 115:E7005–E7014. doi: 10.1073/pnas.1806760115.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nat Methods. 18:170–175. doi: 10.1038/s41592-020-01056-5.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. Systematic Biology. 65:997–1008. doi: 10.1093/sysbio/syw037.

Dainat J et al. 2023. NBISweden/AGAT: AGAT-v1.2.0. doi: 10.5281/zenodo.8178877.

Danecek P et al. 2021. Twelve years of SAMtools and BCFtools. GigaScience. 10:giab008. doi: 10.1093/gigascience/giab008.

Davison HR et al. 2022. Genomic diversity across the *Rickettsia* and '*Candidatus* Megaira' genera and proposal of genus status for the Torix group. Nat Commun. 13:2630. doi: 10.1038/s41467-022-30385-6.

Dickson LB et al. 2016. Reproductive Incompatibility Involving Senegalese *Aedes aegypti* (L) Is Associated with Chromosome Rearrangements. PLOS Neglected Tropical Diseases. 10:e0004626. doi: 10.1371/journal.pntd.0004626.

Dubey SK, Mehta D, Chaudhary S, Hasan A, Sunil S. 2022. An E3 Ubiquitin Ligase Scaffolding Protein Is Proviral during Chikungunya Virus Infection in *Aedes aegypti*. Microbiology Spectrum. 10:e00595-22. doi: 10.1128/spectrum.00595-22.

Dudchenko O et al. 2017. De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. Science. 356:92–95. doi: 10.1126/science.aal3327.

Durand NC et al. 2016. Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. Cell Syst. 3:99–101. doi: 10.1016/j.cels.2015.07.012.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol. 20:1–14. doi: 10.1186/s13059-019-1832-y.

Fisher CR, Wilson M, Scott JG. 2022. A chromosome-level assembly of the widely used Rockefeller strain of *Aedes aegypti*, the yellow fever mosquito. G3 Genes|Genomes|Genetics. 12:jkac242. doi: 10.1093/g3journal/jkac242.

Flynn JM et al. 2020. RepeatModeler2 for automated genomic discovery of transposable element families. Proceedings of the National Academy of Sciences. 117:9451–9457. doi: 10.1073/pnas.1921046117.

Giraldo MI et al. 2020. Envelope protein ubiquitination drives entry and pathogenesis of Zika virus. Nature. 585:414–419. doi: 10.1038/s41586-020-2457-8.

Gloria-Soria A et al. 2022. Origins of high latitude introductions of *Aedes aegypti* to Nebraska and Utah during 2019. Infection, Genetics and Evolution. 103:105333. doi: 10.1016/j.meegid.2022.105333.

Gloria-Soria A, Soghigian J, Kellner D, Powell JR. 2019. Genetic diversity of laboratory strains and implications for research: The case of *Aedes aegypti*. PLOS Neglected Tropical Diseases. 13:e0007930. doi: 10.1371/journal.pntd.0007930.

Goel M, Sun H, Jiao W-B, Schneeberger K. 2019. SyRI: finding genomic rearrangements and local sequence differences from whole-genome assemblies. Genome Biology. 20:277. doi: 10.1186/s13059-019-1911-0.

Goldman-Huertas B et al. 2015. Evolution of herbivory in Drosophilidae linked to loss of behaviors, antennal responses, odorant receptors, and ancestral diet. Proceedings of the National Academy of Sciences. 112:3026–3031. doi: 10.1073/pnas.1424656112.

Goodacre SL, Martin OY, Thomas CFG, Hewitt GM. 2006. *Wolbachia* and other endosymbiont infections in spiders. Molecular Ecology. 15:517–527. doi: 10.1111/j.1365-294X.2005.02802.x.

Gu Z, Hübschmann D. 2023. *simplifyEnrichment*: A Bioconductor Package for Clustering and Visualizing Functional Enrichment Results. Genomics, Proteomics & Bioinformatics. 21:190–202. doi: 10.1016/j.gpb.2022.04.008.

Guan D et al. 2020. Identifying and removing haplotypic duplication in primary genome assemblies. Bioinformatics. 36:2896–2898. doi: 10.1093/bioinformatics/btaa025.

Guo S, Kim J. 2007. Molecular Evolution of Drosophila Odorant Receptor Genes. Molecular Biology and Evolution. 24:1198–1207. doi: 10.1093/molbev/msm038.

Hagimori T, Abe Y, Date S, Miura K. 2006. The First Finding of a *Rickettsia* Bacterium Associated with Parthenogenesis Induction Among Insects. Curr Microbiol. 52:97–101. doi: 10.1007/s00284-005-0092-0.

Hartberg WK, Craig Jr GeorgeB. 1970. Reproductive Isolation in *Stegomyia* Mosquitoes. II. Hybrid Breakdown Between *Aedes aegypti* and *A. mascarensis*. Evolution. 24:692–703.

Herre M et al. 2022. Non-canonical odor coding in the mosquito. Cell. 185:3104-3123.e28. doi: 10.1016/j.cell.2022.07.024.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol. 35:518–522. doi: 10.1093/molbev/msx281.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. Bioinformatics. 32:767–769. doi: 10.1093/bioinformatics/btv661.

Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-Genome Annotation with BRAKER. In: Gene Prediction: Methods and Protocols. Kollmar, M, editor. Methods in Molecular Biology Springer: New York, NY pp. 65–95. doi: 10.1007/978-1-4939-9173-0_5.

Iwata H, Gotoh O. 2012. Benchmarking spliced alignment programs including Spaln2, an extended version of Spaln that incorporates additional species-specific features. Nucleic Acids Research. 40:e161. doi: 10.1093/nar/gks708.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods. 14:587–589. doi: 10.1038/nmeth.4285.

Kikuchi Y, Sameshima S, Kitade O, Kojima J, Fukatsu T. 2002. Novel Clade of *Rickettsia* spp. from Leeches. Appl Environ Microbiol. 68:999–1004. doi: 10.1128/AEM.68.2.999-1004.2002.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol. 37:540–546. doi: 10.1038/s41587-019-0072-8.

Kramer IM et al. 2023. Genomic profiling of climate adaptation in Aedes aegypti along an altitudinal gradient in Nepal indicates nongradual expansion of the disease vector. Molecular Ecology. 32:350–368. doi: 10.1111/mec.16752.

Kuznetsov D et al. 2023. OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. Nucleic Acids Research. 51:D445–D451. doi: 10.1093/nar/gkac998.

Lambrechts L et al. 2009. Genetic specificity and potential for local adaptation between dengue viruses and mosquito vectors. BMC Evol Biol. 9:160. doi: 10.1186/1471-2148-9-160.

Le Goff G, Brengues C, Robert V. 2013. Stegomyia mosquitoes in Mayotte, taxonomic study and description of Stegomyia pia n. sp. Parasite. 20:31. doi: 10.1051/parasite/2013030.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34:3094–3100. doi: 10.1093/bioinformatics/bty191.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics. 25:1754–1760. doi: 10.1093/bioinformatics/btp324.

Liang J et al. 2024. Discovery and characterization of chromosomal inversions in the arboviral vector mosquito *Aedes aegypti*. 2024.02.16.580682. doi: 10.1101/2024.02.16.580682.

Lomsadze A, Ter-Hovhannisyan V, Chernoff YO, Borodovsky M. 2005. Gene identification in novel eukaryotic genomes by self-training algorithm. Nucleic Acids Research. 33:6494–6506. doi: 10.1093/nar/gki937.

Lovell JT et al. 2022. GENESPACE tracks regions of interest and gene copy number variation across multiple genomes Weigel, D, editor. eLife. 11:e78526. doi: 10.7554/eLife.78526.

Lukyanchikova V et al. 2022. *Anopheles* mosquitoes reveal new principles of 3D genome organization in insects. Nat Commun. 13:1960. doi: 10.1038/s41467-022-29599-5.

Macdonald WW. 1962. The Selection of a Strain of *Aëdes aegypti* Susceptible to Infection with Semi-Periodic Brugia Malayi. Annals of Tropical Medicine & Parasitology. 56:368–372. doi: 10.1080/00034983.1962.11686134.

Macdonald WW, Sheppard PM. 1965. Cross-over values in the sex chromosomes of the mosquito *Aedes aegypti* and evidence of the presence of inversions. Annals of Tropical Medicine & Parasitology. 59:74–87. doi: 10.1080/00034983.1965.11686285.

Matthews BJ et al. 2018. Improved reference genome of *Aedes aegypti* informs arbovirus vector control. Nature. 563:501–507. doi: 10.1038/s41586-018-0692-z.

Matthews BJ, McBride CS, DeGennaro M, Despo O, Vosshall LB. 2016. The neurotranscriptome of the *Aedes aegypti* mosquito. BMC Genomics. 17:1–20. doi: 10.1186/s12864-015-2239-0.

McBride CS et al. 2014. Evolution of mosquito preference for humans linked to an odorant receptor. Nature. 515:222–227. doi: 10.1038/nature13964.

McBride CS, Arguello JR. 2007. Five *Drosophila* Genomes Reveal Nonneutral Evolution and the Signature of Host Specialization in the Chemoreceptor Superfamily. Genetics. 177:1395–1416. doi: 10.1534/genetics.107.078683.

Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2021. CAFE 5 models variation in evolutionary rates among gene families. Bioinformatics. 36:5516–5518. doi: 10.1093/bioinformatics/btaa1022.

Mi H et al. 2019. Protocol Update for large-scale genome and gene function analysis with the PANTHER classification system (v.14.0). Nat Protoc. 14:703–721. doi: 10.1038/s41596-019-0128-8.

Mi H, Muruganujan A, Casagrande JT, Thomas PD. 2013. Large-scale gene function analysis with the PANTHER classification system. Nat Protoc. 8:1551–1566. doi: 10.1038/nprot.2013.092.

Minh BQ et al. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution. 37:1530–1534. doi: 10.1093/molbev/msaa015.

Morelli MW, Blackmon H, Hjelmen CE. 2022. Diptera and Drosophila Karyotype Databases: A Useful Dataset to Guide Evolutionary and Genomic Studies. Front. Ecol. Evol. 10. doi: 10.3389/fevo.2022.832378.

Neafsey DE et al. 2015. Highly evolvable malaria vectors: The genomes of 16 *Anopheles* mosquitoes. Science. 347:1258522. doi: 10.1126/science.1258522.

Nene V et al. 2007. Genome Sequence of *Aedes aegypti*, a Major Arbovirus Vector. Science. 316:1718–1723. doi: 10.1126/science.1138878.

Nurk S et al. 2020. HiCanu: accurate assembly of segmental duplications, satellites, and allelic variants from high-fidelity long reads. Genome Res. 30:1291–1305. doi: 10.1101/gr.263566.120.

Palatini U et al. 2017. Comparative genomics shows that viral integrations are abundant and express piRNAs in the arboviral vectors *Aedes aegypti* and *Aedes albopictus*. BMC Genomics. 18:512. doi: 10.1186/s12864-017-3903-3.

Palatini U et al. 2020. Improved reference genome of the arboviral vector *Aedes albopictus*. Genome Biol. 21:215. doi: 10.1186/s13059-020-02141-w.

Paril J, Zare T, Fournier-Level A. 2023. Compare_Genomes: A Comparative Genomics Workflow to Streamline the Analysis of Evolutionary Divergence Across Eukaryotic Genomes. Current Protocols. 3:e876. doi: 10.1002/cpz1.876.

Peng C et al. 2021. A Draft Genome Assembly of *Culex pipiens pallens* (Diptera: Culicidae) Using PacBio Sequencing Lavrov, D, editor. Genome Biology and Evolution. 13:evab005. doi: 10.1093/gbe/evab005.

Perlman SJ, Hunter MS, Zchori-Fein E. 2006. The emerging diversity of *Rickettsia*. Proceedings of the Royal Society B: Biological Sciences. 273:2097–2106. doi: 10.1098/rspb.2006.3541.

Pierson TC, Diamond MS. 2020. The continued threat of emerging flaviviruses. Nat Microbiol. 5:796–812. doi: 10.1038/s41564-020-0714-0.

Poupardin R, Riaz MA, Vontas J, David JP, Reynaud S. 2010. Transcription profiling of eleven cytochrome P450s potentially involved in xenobiotic metabolism in the mosquito Aedes aegypti. Insect Molecular Biology. 19:185–193. doi: 10.1111/j.1365-2583.2009.00967.x.

Powell JR, Gloria-Soria A, Kotsakiozi P. 2018. Recent History of *Aedes aegypti*: Vector Genomics and Epidemiology Records. BioScience. 68:854–860. doi: 10.1093/biosci/biy119.

Powell JR, Petrarca V, della Torre A, Caccone A, Coluzzi M. 1999. Population structure, speciation, and introgression in the *Anopheles gambiae* complex. Parassitologia. 41:101–113.

Rai KS, Black WC. 1999. 1 - Mosquito Genomes: Structure, Organization, and Evolution. In: Advances in Genetics. Hall, JC, Dunlap, JC, Friedmann, T, & Giannelli, F, editors. Vol. 41 Academic Press pp. 1–33. doi: 10.1016/S0065-2660(08)60149-2.

Raoult D, Roux V. 1997. Rickettsioses as paradigms of new or emerging infectious diseases. Clinical Microbiology Reviews. 10:694–719. doi: 10.1128/cmr.10.4.694.

Redmond SN et al. 2020. Linked-read sequencing identifies abundant microinversions and introgression in the arboviral vector *Aedes aegypti*. BMC Biol. 18:1–14. doi: 10.1186/s12915-020-0757-y.

Reidenbach KR et al. 2009. Phylogenetic analysis and temporal diversification of mosquitoes (Diptera: Culicidae) based on nuclear genes and morphology. BMC Evol Biol. 9:298. doi: 10.1186/1471-2148-9-298.

Roach MJ, Schmidt SA, Borneman AR. 2018. Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies. BMC Bioinformatics. 19:460. doi: 10.1186/s12859-018-2485-7.

Rose NH et al. 2020. Climate and Urbanization Drive Mosquito Preference for Humans. Current Biology. 30:3570-3579.e6. doi: 10.1016/j.cub.2020.06.092.

Ross PA, Endersby-Harshman NM, Hoffmann AA. 2019. A comprehensive assessment of inbreeding and laboratory adaptation in *Aedes aegypti* mosquitoes. Evolutionary Applications. 12:572–586. doi: 10.1111/eva.12740.

Ryazansky SS et al. 2024. The chromosome-scale genome assembly for the West Nile vector *Culex quinquefasciatus* uncovers patterns of genome evolution in mosquitoes. BMC Biol. 22:16. doi: 10.1186/s12915-024-01825-0.

Sato K et al. 2008. Insect olfactory receptors are heteromeric ligand-gated ion channels. Nature. 452:1002–1006. doi: 10.1038/nature06850.

Schmidt TL, Chung J, Honnen A-C, Weeks AR, Hoffmann AA. 2020. Population genomics of two invasive mosquitoes (*Aedes aegypti* and *Aedes albopictus*) from the Indo-Pacific. PLOS Neglected Tropical Diseases. 14:e0008463. doi: 10.1371/journal.pntd.0008463.

Sharakhov IV et al. 2002. Inversions and Gene Order Shuffling in *Anopheles gambiae* and *A. funestus.* Science. 298:182–185. doi: 10.1126/science.1076803.

Sharakhov IV, Artemov GN, Sharakhova MV. 2016. Chromosome evolution in malaria mosquitoes inferred from physically mapped genome assemblies. J. Bioinform. Comput. Biol. 14:1630003. doi: 10.1142/S0219720016300033.

Simon J-C et al. 2015. Genomics of adaptation to host-plants in herbivorous insects. Briefings in Functional Genomics. 14:413–423. doi: 10.1093/bfgp/elv015.

Smit A, Hubley R, Green P. 2013. RepeatMasker. http://www.repeatmasker.org.

Soghigian J et al. 2020. Genetic evidence for the origin of *Aedes aegypti*, the yellow fever mosquito, in the southwestern Indian Ocean. Molecular Ecology. 29:3593–3606. doi: 10.1111/mec.15590.

Soghigian J et al. 2023. Phylogenomics reveals the history of host use in mosquitoes. Nat Commun. 14:6252. doi: 10.1038/s41467-023-41764-y.

Soudi S et al. 2023. Genomic signatures of local adaptation in recent invasive *Aedes aegypti* populations in California. BMC Genomics. 24:311. doi: 10.1186/s12864-023-09402-5.

Stanke M, Keller O, et al. 2006. AUGUSTUS: ab initio prediction of alternative transcripts. Nucleic Acids Research. 34:W435–W439. doi: 10.1093/nar/gkl200.

Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically mapped cDNA alignments to improve de novo gene finding. Bioinformatics. 24:637–644. doi: 10.1093/bioinformatics/btn013.

Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. BMC Bioinformatics. 7:1–11. doi: 10.1186/1471-2105-7-62.

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of transposable element families, sequence models, and genome annotations. Mobile DNA. 12:1–14. doi: 10.1186/s13100-020-00230-y.

The Anopheles gambiae 1000 Genomes Consortium. 2020. Genome variation and population structure among 1142 mosquitoes of the African malaria vector species Anopheles gambiae and Anopheles coluzzii. Genome Res. 30:1533–1546. doi: 10.1101/gr.262790.120.

Thongprem P, Davison HR, Thompson DJ, Lorenzo-Carballa MO, Hurst GDD. 2021. Incidence and Diversity of Torix *Rickettsia*–Odonata Symbioses. Microb Ecol. 81:203–212. doi: 10.1007/s00248-020-01568-9.

To T-H, Jung M, Lycett S, Gascuel O. 2016. Fast Dating Using Least-Squares Criteria and Algorithms. Systematic Biology. 65:82–97. doi: 10.1093/sysbio/syv068.

Tsunoyama K, Bellgard MI, Gojobori T. 2001. Intragenic Variation of Synonymous Substitution Rates Is Caused by Nonrandom Mutations at Methylated CpG. J Mol Evol. 53:456–464. doi: 10.1007/s002390010235.

Uliano-Silva M et al. 2023. MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads. BMC Bioinformatics. 24:1–13. doi: 10.1186/s12859-023-05385-y.

Vieira FG, Sánchez-Gracia A, Rozas J. 2007. Comparative genomic analysis of the odorant-binding protein family in 12 *Drosophila* genomes: purifying selection and birth-and-death evolution. Genome Biol. 8:R235. doi: 10.1186/gb-2007-8-11-r235.

Wang JZ, Du Z, Payattakool R, Yu PS, Chen C-F. 2007. A new method to measure the semantic similarity of GO terms. Bioinformatics. 23:1274–1281. doi: 10.1093/bioinformatics/btm087.

Wang Y et al. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Research. 40:e49. doi: 10.1093/nar/gkr1293.

Ware-Gilmore F, Novelo M, Sgrò CM, Hall MD, McGraw EA. 2023. Assessing the role of family level variation and heat shock gene expression in the thermal stress response of the mosquito *Aedes aegypti*. Philosophical Transactions of the Royal Society B: Biological Sciences. 378:20220011. doi: 10.1098/rstb.2022.0011.

Wei Y et al. 2017. Comparative physical genome mapping of malaria vectors *Anopheles sinensis* and *Anopheles gambiae*. Malar J. 16:235. doi: 10.1186/s12936-017-1888-7.

Weinert LA, Werren JH, Aebi A, Stone GN, Jiggins FM. 2009. Evolution and diversity of *Rickettsia* bacteria. BMC Biol. 7:6. doi: 10.1186/1741-7007-7-6.

Wellenreuther M, Bernatchez L. 2018. Eco-Evolutionary Genomics of Chromosomal Inversions. Trends in Ecology & Evolution. 33:427–440. doi: 10.1016/j.tree.2018.04.002.

Wellenreuther M, Mérot C, Berdan E, Bernatchez L. 2019. Going beyond SNPs: The role of structural genomic variants in adaptive evolution and species diversification. Molecular Ecology. 28:1203–1209. doi: 10.1111/mec.15066.

Wenger AM et al. 2019. Accurate circular consensus long-read sequencing improves variant detection and assembly of a human genome. Nat Biotechnol. 37:1155–1162. doi: 10.1038/s41587-019-0217-9.

Werren JH. 1997. Biology of *Wolbachia*. Annual Review of Entomology. 42:587–609. doi: 10.1146/annurev.ento.42.1.587.

Werren JH et al. 1994. Rickettsial relative associated with male killing in the ladybird beetle (*Adalia bipunctata*). Journal of Bacteriology. 176:388–394. doi: 10.1128/jb.176.2.388-394.1994.

Whibley A, Kelley JL, Narum SR. 2021. The changing face of genome assemblies: Guidance on achieving high-quality reference genomes. Molecular Ecology Resources. 21:641–652. doi: 10.1111/1755-0998.13312.

Whitfield ZJ et al. 2017. The Diversity, Structure, and Function of Heritable Adaptive Immunity Sequences in the *Aedes aegypti* Genome. Current Biology. 27:3511-3519.e7. doi: 10.1016/j.cub.2017.09.067.

Wright CJ, Stevens L, Mackintosh A, Lawniczak M, Blaxter M. 2024. Comparative genomics reveals the dynamics of chromosome evolution in Lepidoptera. Nat Ecol Evol. 8:777–790. doi: 10.1038/s41559-024-02329-4.

Xie W et al. 2018. The invasive MED/Q *Bemisia tabaci* genome: a tale of gene loss and gene gain. BMC Genomics. 19:68. doi: 10.1186/s12864-018-4448-9.

Xu M et al. 2020. TGS-GapCloser: A fast and accurate gap closer for large genomes with low coverage of error-prone long reads. GigaScience. 9:giaa094. doi: 10.1093/gigascience/giaa094.

Yee DA et al. 2022. Robust network stability of mosquitoes and human pathogens of medical importance. Parasites & Vectors. 15:216. doi: 10.1186/s13071-022-05333-4.

Yurchenko AA et al. 2023. Phylogenomics revealed migration routes and adaptive radiation timing of Holarctic malaria mosquito species of the Maculipennis Group. BMC Biol. 21:63. doi: 10.1186/s12915-023-01538-w.

Zeng X et al. 2024. Chromosome-level scaffolding of haplotype-resolved assemblies using Hi-C data without reference genomes. Nat. Plants. 10:1184–1200. doi: 10.1038/s41477-024-01755-3.

Zhao L, Pridgeon JW, Becnel JJ, Clark GG, Linthicum KJ. 2009. Identification of Genes Differentially Expressed During Heat Shock Treatment in *Aedes aegypti*. Journal of Medical Entomology. 46:490–495. doi: 10.1603/033.046.0312.

Zhou C, McCarthy SA, Durbin R. 2023. YaHS: yet another Hi-C scaffolding tool. Bioinformatics. 39:btac808. doi: 10.1093/bioinformatics/btac808.

Zwiebel LJ, Takken W. 2004. Olfactory regulation of mosquito–host interactions. Insect Biochemistry and Molecular Biology. 34:645–652. doi: 10.1016/j.ibmb.2004.03.017.

Table 1. Final assembly metrics for Aaf (*Aedes aegypti formosus*) and Am (*Ae. mascarensis*) compared to those of AaegL5 (*Ae. aegypti*).

Metric	AaegL5	Aaf	Am
# Scaffolds	2,310	706	74
# Contigs	2,539	1,124	269
Total length (Gbp)	1.279	1.239	1.289
Gap %	0.002	0.007	0.003
L50	2	2	2
N50 (Mbp)	40.978	39.859	42.609
GC ± SD	0.382 ± 0.029	0.381 ± 0.074	0.381 ± 0.04
Mitogenome length (Mbp)	16,790	16,617	16,428
# Exons	94,104	63,767	62,375
# Genes	19,203	17,672	17,009

Table 2. Overrepresentation test of terms common to all three Aegypti group assemblies (*Aedes aegypti formosus:* Aaf; *Ae. aegypti* AaegL5; *Ae. mascarensis* Am) done on the PANTHER DB web interface (release 20240807). The reference annotation set is from *Anopheles gambiae*. Significance testing done using Fisher's Exact Test and the resulting P-values were corrected using false discovery rate (FDR). Here, 'Fold' refers to fold enrichment calculated as the actual number in the sample divided by the expected number from the same relative to the reference set. Here, the em-dashes ("—") reflect the nested nature of GO terms and alternating shading of the rows separate basal-most GO terms.

GO	GO ID	Aaf		AaegL5		Am	
		Fold	P_{FDR}	Fold	P_{FDR}	Fold	P_{FDR}
male courtship behavior	GO:0008049	10.92	<0.0001	9.70	<0.0001	8.95	<0.0001
—male mating behavior	GO:0060179	10.92	<0.0001	9.70	<0.0001	8.95	<0.0001
——mating behavior	GO:0007617	10.68	<0.0001	9.49	<0.0001	8.75	<0.0001
———reproductive behavior	GO:0019098	10.68	<0.0001	9.49	<0.0001	8.75	<0.0001
————multicellular organismal reproductive process	GO:0048609	7.67	<0.0001	5.32	<0.0001	4.22	<0.0001
—————reproductive process	GO:0022414	5.62	<0.0001	3.99	<0.0001	3.18	<0.0001
————behavior	GO:0007610	7.74	<0.0001	7.21	<0.0001	6.95	<0.0001
————multicellular organismal process	GO:0032501	2.99	<0.0001	2.54	<0.0001	1.57	0.0001
—courtship behavior	GO:0007619	10.92	<0.0001	9.70	<0.0001	8.95	<0.0001
chemosensory behavior	GO:0007635	9.27	<0.0001	8.64	<0.0001	8.34	<0.0001
—response to chemical	GO:0042221	2.25	<0.0001	4.04	<0.0001	2.16	<0.0001
sensory perception of taste	GO:0050909	8.05	<0.0001	7.31	<0.0001	7.62	<0.0001
—sensory perception of chemical stimulus	GO:0007606	4.77	<0.0001	6.05	<0.0001	2.67	<0.0001
——sensory perception	GO:0007600	4.69	<0.0001	5.13	<0.0001	2.28	<0.0001
———nervous system process	GO:0050877	4.72	<0.0001	5.12	<0.0001	2.38	<0.0001
————system process	GO:0003008	4.63	<0.0001	4.97	<0.0001	2.29	<0.0001
excitatory postsynaptic potential	GO:0060079	6.99	0.0008	6.21	0.0027	5.73	0.0048
—chemical synaptic transmission, postsynaptic	GO:0099565	6.99	0.0008	6.21	0.0027	5.73	0.0047
——synaptic signaling	GO:0099536	3.86	<0.0001	2.37	0.0183	2.18	0.0378
———signaling	GO:0023052	1.90	<0.0001	1.51	0.0004	1.46	0.0001
————cell communication	GO:0007154	1.88	<0.0001	1.51	0.0003	1.45	0.0013
——signal transduction	GO:0007165	1.89	<0.0001	1.49	0.0020	1.44	0.0047
—regulation of postsynaptic membrane potential	GO:0060078	5.73	<0.0001	5.09	0.0003	4.7	0.0006
——regulation of membrane potential	GO:0042391	3.80	<0.0001	3.04	0.0012	2.65	0.0101
cellular component organization	GO:0016043	0.61	0.0001	0.74	0.0409	0.66	0.0006
—cellular component organization or biogenesis	GO:0071840	0.59	<0.0001	0.72	0.0112	0.71	0.0043

A. *Aedes aegypti formosus* (Aaf)

B. *Aedes mascarensis* (Am)

Fig.1. Lollipop plot ranking each draft assemblies for (A) *Aedes aegypti formosus* (Aaf) and (B) *Ae. mascarensis* (Am). The two best assemblies for each taxon is indicated by larger, blue circles. The scores based on duplicated (0.15), fragmented (0.125), missing (0.15), collapses (0.10), expansions (0.1), inversions (0.1), N50 (0.1), and relative genome size (0.175) [metric (weight)].

Fig. 2 Genomic content for *Aedes aegypti formosus* (Aaf; A), *Ae. mascarensis* (Am; B) *Aedes aegypti* reference genome found by Matthews et al., (2018) [AaegL5; C] and repeat landscape plots (A and B only). In the landscape plots, sequence divergence is shown in 1% intervals. Sequence divergence of the landscape plot was estimated using the Kimura model of sequence evolution modified to account for the high mutability of CpG sites. Landscape plots do not account for "Unmasked DNA". Categories with trivial number of bases are shown as "~0%" (A and B), while "0%" for C are actual 0s.

Fig. 3 Synteny between chromosomes of *Aedes aegypti* (AaegL5) and *Ae. aegypti formosus* (Aaf). Bacterial artificial chromosomes (BAC; Matthews et al. 2018) positions represented as black horizontal bars.

Fig. 4 Riparian plot showing gene order synteny between eleven culicid genomes and the outgroup *Phlebotomus papatasi*. Synteny is assessed relative to the *Aedes aegypti* reference genome (AaegL5)—all genes originating from chromosome 1 on AaegL5 are shown in purple, all genes originating from chromosome 2 on AaegL5 are shown in green, and all genes originating from chromosome 3 are shown in yellow. Chromosomes with "*" in their names have been inverted to facilitate visibility. Similarly, some chromosomes appear out of order to facilitate visibility. Note that chromosomes 4 and 5 of *P. papatasi* are microchromosomes and not shown because too few orthologs were detected on them to adequately assess synteny with any of the chromosomes of AaegL5.

Fig. 5 Maximum likelihood phylogeny output from IQTree2, re-rooted (to *Phlebotomus papatasi*) and ultrametricized using the *ape* package for R. Values shown at all nodes and tips represent rapidly evolving orthologs (blue) and expansions or contractions. All circles represent the ratio of expansions and contractions of ortholog copy number at each node and tip. Size of each circle represents the magnitude of the ratio (calculated as the greater of the two numbers divided by the lesser), while color represents direction (contraction-biased ratios are negative and darker, expansion-biased ratios are lighter and positive). Bold letters denote ancestral nodes for taxonomic groups: Aedini (A), Culicini (B), Culicinae (C), Anophelinae (D).

Fig. 6 Heatmaps of GO term similarity and overrepresentation for the *Aedes aegypti* reference genome (AaegL5), *Ae. aegypti formosus* (Aaf), and *Ae. mascarensis* (Am). Similarity of 194 GO terms were assessed using semantic similarity outlined by Wang et al. (2007) and used K-means clustering to form eleven clusters of GO terms. Significance ($P_{FDR} < 0.01$) of overrepresentation of each GO term is shown in the heat map to the left. The heat map on the right shows similarity of individual GO terms, wherein GO terms are clustered together by similarity and divided by horizontal and vertical lines. The most commonly occurring descriptive terms for each cluster is shown to the right, where more frequent terms are shown in larger text.